Artificial Selection and the Skin Microbiome Independently Predict Parasite Resistance

Rachael D. Kramp[1*], Mary J. Janecka[1,5], Nadine Tardent[2,3], Jukka Jokela[2,3], Kevin D. Kohl[1], Jessica F. Stephenson[1,2,3,4]

1. Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA 15260, USA

2. Department of Aquatic Ecology, EAWAG, Swiss Federal Institute of Aquatic Sci & Tech, Dübendorf 8600, Switzerland.

3. Institute for Integrative Biology, ETH Zürich, Zürich 8092, Switzerland.

4. Center for Adaptation to a Changing Environment, ETH Zürich, Zürich 8092, Switzerland.

5. Department of Biological Sciences, University of Texas at El Paso, El Paso, TX 79968, USA

*Author for correspondence: rdkramp@gmail.com

1. It is well established that host genetics determine much of hosts response to parasites, but recent research has highlighted that the host microbiome plays a role in defense against parasites. Interactions between host resistance to parasites and the microbiome are widely acknowledged; for example, the microbiome can modulate the host immune response, and *vice versa*. However, it remains unclear how the host immune system and the microbiome may together influence the host's overall response to parasites.

2. In many species males and females differ in both their genetically-based parasite resistance and the composition and structure of their associated microbiomes. We may therefore expect interactions between host resistance and the microbiome to differ between males and females, potentially explaining the often-observed variation in their response to parasites.

3. To investigate how host genetics and the microbiome interact to shape host response to parasites, we imposed truncation selection on Trinidadian guppies (*Poecilia reticulata*) for their response to the specialist ectoparasite *Gyrodactylus turnbulli*. The 30% of guppies with the lowest number of parasites, "parasite load", during experimental infection founded the resistant line, and the 30% with the highest parasite load founded the susceptible line, while randomly chosen uninfected fish founded the control line. After 3-6 generations of breeding in the absence of parasites, we sampled the skin-associated microbiome of fish from these lines and then infected them with *G. turnbulli*. We used Dirichlet multinomial modeling (DMM) machine learning to identify the bacterial community types across lines and evaluated the importance of selection line and bacterial community type in explaining variation between fish in parasite load over time 'infection severity".

4. Among females, the resistant line developed significantly lower infection severity, and the susceptible line developed significantly higher infection severity, than any other treatment. Among males, however, while the susceptible line developed significantly higher infection severity than the other lines, there was no difference between the resistant and control lines. We also found that female fish in the resistant line had significantly lower tolerance compared to the control or susceptible line females. This apparent tradeoff between resistance and tolerance was also supported by analysis of the individual-level data.

5. We found that host-associated microbiomes explain as much of the variation in infection severity as host genetics, and that these two factors appear independent and additive in their effects on infection severity. Furthermore, a post-hoc test showed that a model including both artificial selection line and bacterial community type is significantly better at explaining variation in infection severity than models with either variable alone. Our results therefore suggest that the microbiome plays an active role in the interaction between host and parasite and is not a passive reflection of host genetics.

Keywords: Trinidadian guppies, ectoparasite, sex-specific differences, host-parasite interactions, artificial selection, parasite resistance, skin microbiome, pre-infection microbiome, bacterial community, parasite-tolerance

Introduction

Hosts use multiple strategies to defend themselves against parasites and pathogens (parasites hereafter). Host genetic background is commonly credited to determine much of the defense against parasites through immune system, behavior, or other genetic/protein alterations, e.g., thalassemia (Hawley et al., 2011; Janecka et al., 2021; Luzzi et al., 1991; Råberg et al., 2007). However, a more recently highlighted defense, the host-associated microbial communities or microbiome, may also influence parasite defense indirectly through the host immune system or directly via niche occupation or secreting anti-parasitic compounds (Nelson & May, 2020; Speer, 2022). Indeed, the host-associated microbiome can dramatically impact host survival and defense to parasites (Hopkins et al., 2017). For example, chytrid fungus, which is currently decimating amphibian biodiversity, can be suppressed by the addition of a natural skin microbe, *Janthinobacterium lividum* (Becker et al., 2009; Brucker et al., 2008). Similarly, the disruption of the gut microbiome alerts the host immune system to parasitic invasion of *Toxoplasma gondii*, and thus improves disease outcomes (Benson et al., 2009). Moreover, the "pre-infection" microbiome can be a powerful tool to predict individuals that are most susceptible to parasites within a population (Nasal: (Lee et al., 2019)& Skin:(Bates et al., 2022)). However, the question of whether the microbiome is playing an active role in host-parasite infection or is merely a reflection of the host's health/immune status is commonly posed (de Vos et al., 2022; Dheilly et al., 2019; Hanage, 2014; Reverter et al., 2020; Trevelline et al., 2020). While there is abundant evidence that the microbial community can have a direct or indirect role in host-parasite interactions (Abt et al., 2012; Becker et al., 2009; Ichinohe et al., 2011; Le Sage et al., 2021; Nelson & May, 2020; Ost & Round, 2017), there is also evidence that the microbiome is merely a bystander in some diseases and a reflection of the host's genetics, diet, and general health (Ni et al., 2017; Shanahan, 2012).

To fully understand the interplay between host genetics and microbiome, it is essential to consider how hosts influence microbial communities and their colonization. Hosts exert control over their symbionts via immune responses, barrier function, physiological regulation, and other mechanisms, enabling them to shape both the ecology and evolution of their microbiota (Wilde et al., 2024). This host-driven control creates natural selection for microbial traits that benefit the host, which ultimately could affect traits like parasite resistance. The host's genetic makeup influences microbiome composition, leading to selection for specific microbial communities that can be

inherited across generations (Bordenstein & Theis, 2015; Kohl, 2020; Lim & Bordenstein, 2020; Mazel et al., 2018).

For example, siblings display more similar microbiomes than cohoused non-siblings, highlighting the heritable

aspect of these communities (Garrigues et al., 2022; Hand et al., 2013; Robertson et al., 2019). This dynamic

interaction between host genetics and microbiome suggests that hosts are not merely passive environments but are

actively shaping their microbial inhabitants through selective pressures. However, disentangling host genetic effects

from microbial influence presents challenges. These ecological communities are subject to dispersal, priority

effects, ecological drift, and other elements of stochasticity (J. Z. Chen et al., 2024; Debray et al., 2022; Jones et

al., 2022). Further, host immune systems rely on early microbial colonization for proper immune development

(Amenyogbe et al., 2017; Mazel et al., 2018), creating a feedback loop that complicates the ability to fully isolate

genetic versus microbial contributions. Despite the heritability of microbiomes, exact mechanisms governing these

processes remain under investigation, and the influence of external factors such as environment and temporal

changes must be carefully considered (Kohl, 2020). Direct manipulation of the host genetics in traits like parasite

resistance, where the microbiome plays a known role—will aid in disentangling genetic effects from microbial ones.

Adding further complexity, the microbiome and the host immune system are both commonly cited to differ

between the host sexes (Kramp et al., 2022; Sylvia et al., 2017). The microbiome can be altered by physiological

factors that commonly differ between sexes. For example, high levels of testosterone in the blood are known to

repress the immune system and significantly alter the microbial communities present within the gut (Santos-Marcos

et al., 2023). Similar changes in both immune system and microbial community are observed during pregnancy and

menstruation (Chen et al., 2023; Gorczyca et al., 2022). Investigating the immune system and the microbiome in a

sex-specific manner thus provides a more holistic understanding of host-parasite interactions in natural systems,

and it can also be a powerful tool to examine how genetic mechanisms of parasite resistance may co-evolve with

microbiomes to meet differing fitness demands between the sexes.

We artificially selected Trinidadian guppies, *Poecilia reticulata*, for resistance or susceptibility to their

specialist parasite *Gyrodactylus turnbulli*, with resistant fish having lower worm burdens (indication of parasite

suppression) and susceptible fish having higher worm burdens. Additionally, while we did not specifically select

for tissue-specific tolerance (tolerance hereafter), or the ability to minimize the fitness losses caused by a given

parasite load (Stephenson & Adelman, 2022), we did examine the levels of tolerance within our lines. Previous

evidence indicates that guppy resistance or susceptibility to *Gyrodactylus turnbulli* is genetically linked and

heritable, with potential sex differences (Cable & Van Oosterhout, 2007; Dargent et al., 2016; Fraser & Neff, 2009;

Madhavi & Anderson, 1985). The major histocompatibility complex (MHC) appears to be part of genetic-based

resistance to *G. turnbulli*, which functions by presenting peptide fragments from parasites to host immune cells, and

subsequent immune recognition and response (Fraser et al., 2010; Fraser & Neff, 2009; Phillips et al., 2018).

However, both innate and acquired immune responses likely contribute to *G. turnbulli* resistance (Cable & Van

Oosterhout, 2007). Interestingly, skin bacterial alpha diversity has been shown to be positively correlated with

expected MHC heterozygosity and negatively with MHC nucleotide diversity in frogs (Cortazar-Chinarro et al.,

2024). If the same is true for fish, this covariation sets up a clear avenue for the skin microbiome and host genetics

to interactively shape parasite resistance phenotypes within our system.

We found that both host genetics and the microbiome significantly predicted the severity of parasite

infection, measured as the area under the curve of infection load throughout an individual's infection. However, the

microbiome did not significantly differ across our three selection lines. Interestingly, parasite resistance and

tolerance showed different patterns between the artificially selected male and females. These results suggest that

the microbiome plays an active role in host-parasite interactions, rather than simply reflecting the host's genetics.

Overall, our findings highlight the complex interplay between host genetics, the microbiome, and sex-specific

responses to parasite infection.

Materials and methods

Experimental design and sample collection

Host origin and maintenance

We used laboratory-bred descendants of wild guppies from a high-predation population located at an undeveloped,

rural site on the Caura River, Trinidad (UTM: 20 P 679527.7 m E, 1180376.4 m N based on WGS84 Datum;

elevation 112m). Wild guppies (n ~ 600) were transported to Cardiff University in June 2012 (Cefas APB

authorization number CW054-D-187A), where they were prophylactically treated for infection using Binox®
(Nitrofurazone; Jungle Laboratories Corporation®, Cibolo, Texas). This site was found *Gyrodactylus* spp.-free in
a previous survey (Stephenson, van Oosterhout, Mohammed, et al., 2015), and we did not find any *Gyrodactylus*
spp. during the examination of a subset of 80 of the collected wild fish. Fin clips were taken from the caudal fin of
66 wild-caught, parental generation fish for microsatellite genotyping. Fish were housed in 70 L mixed sex stock
tanks on a 12h light: 12h dark lighting regime (overhead fluorescent lighting) at 24±1°C and fed daily with
Aquarian® tropical fish flakes supplemented with *Artemia* and bloodworm. Fry were removed from the breeding
tanks soon after birth, confirmed free of ectoparasitic infection under a microscope, and reared in one of four
juvenile tanks until sex determination was possible (~6-8 weeks; males identified by the initiation of the
modification of the anal fin, females by dark pigmentation at the base of the anal fin), and then reared in sex-specific
tanks for at least a further month before use. All fish used to initiate the selected lines were, therefore, sexually
mature and unmated.

Selection experiment

We experimentally infected 100 male and 100 female F1 fish, scored their resistance, and either assigned them to a
breeding population ('resistant' or 'susceptible'), or euthanized them. There was only one round of artificial
selection on our lines. To initiate experimental infections, a heavily infected fish "culture fish" from a laboratory-
maintained culture of *Gyrodactylus turnbulli* was killed with an overdose of MS222 (0.02% tricaine
methanesulfonate; PHARMAQ Ltd., Fordingbridge, UK) followed by pithing. This culture fish was then placed
next to an anaesthetized experimental fish until two parasites had transmitted, as observed under a dissecting
microscope and fiber optic illumination. The experimental fish were revived and housed individually in 1 L tanks,
and the number of parasites infecting each was counted under anesthetic every other day. All tanks were maintained
under the same conditions as the breeding tanks and received 100% water exchanges every other day. After nine
days of infection, all fish were treated with Levamisole and screened three times with at least four days between
screenings to confirm they were free of infection. Fin clips were taken from the caudal fin of all fish and stored in
100% ethanol for microsatellite genotyping.

The 30 males and 30 females who maintained the highest mean number of worms throughout their nine-

day infection were assigned to the 'susceptible' line and allowed to breed freely in a single 120 L tank. The 30

males and 30 females who maintained the lowest mean number of worms over the nine-day infection were assigned

to the 'resistant' line and allowed to breed freely in a second 120 L tank. As a control line, we randomly selected

30 males and 30 females from the uninfected F1 sexually mature virgin fish and allowed them to breed in a third

120 L tank. All three tanks were checked at least twice weekly for fry, and these were moved to F2 rearing tanks.

In the F2 generation, each line was split into two (CA, CB, RA, RB, SA, SB), again with a population size of 60

fish per tank (30 males, 30 females). This population size was kept consistent across the six subpopulations, with

new tanks set up for each subsequent generation as above.

Experimental infection

In August 2015, all fish in each of the three lines were transported from Cardiff, UK, to Zürich,

Switzerland. In Zürich, guppies were housed at densities of 1-2 fish per L in 4.5 L tanks in a recirculating system

at 24±1°C, on a 12 h light: 12 h dark lighting schedule (overhead fluorescent lighting) and fed daily on commercial

flake food supplemented with *Artemia*. The fish were maintained and bred within the six subpopulations under

these conditions until the start of the experimental infections, for which 130 fish were infected in five experimental

batches between August and November 2017. All fish used for the experimental infections were born in Zürich and

represented generations 3-6 after the imposed selection. Allowing fish to breed in the absence of parasites for

multiple generations may eliminate potential parental and grand-parental effects, which recent research indicates

are important in this system (TGIP MANUSCRIPT CITE). *G. turnbulli* were collected from Qualipet Glatt Zentrum

in June 2017, and an isogenic line was initiated by infecting a parasite-naïve ornamental guppy with a single parasite

individual. This line was subsequently maintained on parasite-naïve ornamental guppies. Experimental infections

were initiated following the protocol described above. The experimental fish were revived and housed individually

in 1 L tanks, and the number of parasites infecting each was counted under anesthetic every other day. All tanks

were maintained under the same conditions as the breeding tanks and received 100% water exchanges every other

day. After 17 days of infection, all fish were fin-clipped (stored in 100% ethanol), treated and screened clear of

infection as above, and placed in a breeding stock tank.

Microsatellite genotyping

We evaluated the genetic diversity and variance in inbreeding within the wild-caught populations, F1, F3-F6

generations to determine if the observed difference in response to infection between the resistant and susceptible

lines could be attributed to differences in inbreeding between the groups. DNA was isolated from fin clips with the

HotSHOT method using 50 µl of alkaline lysis reagent and neutralizing solution followed by incubation at 95 °C

for 30 minutes (Truett et al., 2000). Eight previously designed microsatellite markers were amplified in two

multiplex assays (Becher et al., 2002; Nater et al., 2008; Paterson et al., 2005; van Oosterhout et al., 2006; Watanabe

et al., 2005). Microsatellites samples were generated in collaboration with the Genetic Diversity Centre (GDC),

ETH Zurich. Samples were genotyped on an ABI Genetic Analyzer 3730 (Applied Biosystems, USA). Samples

were scored using GeneMarker (Version 2.6.4; Softgenetics) and binned using the package MSatAllele (Alberto,

2013) in R (R Core Team, 2020). For more detailed microsatellite PCR prep see supplemental materials.

Microbial inventory

Before infection began, we collected samples from a total of 51 fish split between the infection and sham-infection

treatment groups: Control (n=10, sham n=5), Resistant (n=12, sham n=6), Susceptible (n=9, sham n=8). The

collection and inventory of the microbial communities on the fish skin we used the methods found in (Kramp et al.,

2022).

Data analyses

Infection of 3^{rd}-6^{th} generation offspring

We quantified infection resistance/susceptibility as the area under the curve (AUC) of worm number over day of

infection or "infection severity". To test the outcome of our artificial selection, we used the infection severity AUC

as the response variable in a generalized linear mixed models (glmm) in the glmmTMB package, v. 1.1.2.3 (Brooks

et al., 2017) (Gamma error family and logit link function within R v.4.3.2 ((R Core Team, 2020); see supplementary

material). As fixed main effects, we included artificial selection (line; resistant n=19, control n=15, susceptible

n=16), fish sex, length (On day 0 and controlling for sex differences, see supplementary material), body condition

(scaled mass index (Peig & Green, 2009)), generation (gen), number of parasite starting infection (dose), and trial

date (Julien date). We also included a two-way interaction between sex and line based on lower AIC. We used

ggplot2 (Wickham, 2016) for plotting all figures. For all models, we assessed collinearity among predictor variables

using the pairs function (Zuur et al., 2009). Model validation was performed with the 'DHARMa' package (Hartig,

2022) and AIC criteria. To test for significance across models, we applied Type II Wald chi-square tests using the

Anova function from the 'car' package (Fox & Weisberg, 2019).

Tissue-specific tolerance of 3^{rd}-6^{th} generation offspring

We quantified tissue-specific tolerance "ppmass" as the relative per-parasite percentage mass change of the host

during infection by dividing the percentage mass change between day 0 and day 20 by the maximum number of

worms observed on the fish during its infection (Jog et al., 2022). We used "relative" per-parasite percentage mass

change because we do not expect zero percent body mass change throughout the experiment, e.g., a perfectly

tolerant individual would likely gain weight during the experiment. Therefore, we scale all percentage body mass

change relative to the highest percent body mass change observed in the experiment. To test the effect of artificial

selection on tolerance of infection we used tolerance as a response variable in a glmm in the glmmTMB package,

v. 1.1.2.3 (Brooks et al., 2017) (Gaussian error family and logit link function within R v.4.3.2 ((R Core Team,

2020); see supplementary material). As fixed main effects, we included artificial selection (line), fish sex, length

(On day 0 and controlling for sex differences, see supplementary material), generation (gen), number of parasite

starting infection (dose), and trial date (Julien date). We excluded body condition (scaled mass index) as a fixed

effect due to the response variable incorporating mass changes in its calculations. Furthermore, within the

microbiome we tested if the bacterial community types predicted tolerance by adding it as a fixed effect.

Resistance-tolerance Tradeoff

To investigate the relationship between tissue-specific tolerance and resistance we ran a polynomial regression

model of the second degree (quadratic) to account for the non-linear effects. The model was fit using the lm()

function in R v.4.3.2 ((R Core Team, 2020); see supplementary material). We used Tolerance as the response

variable and log transformed AUC (our resistance metric), fish length, and generation as fixed effects. We did not

include AUC within our main tolerance model, described above, due to high collinearity with other fixed effects

within the model, and fit issues due the nonlinear relationship between AUC and ppmass. This model included fixed

effects without collinearity with AUC.

Calculating Heritability

We calculated narrow-sense heritability (assuming that the response to selection is primarily due to additive genetic

variance) of our artificial selection for mean worm and our calculated tolerance metric for susceptible and resistant

lines following the breeder's equation (Lush, 1937). For mean worm, we used the average number of parasites

between 1-9 days of infection, rescaled to a mean of zero to account for the differences in worm growth between

the infections of the parental and offspring generations. For tolerance, we rescaled using the 'rescale' package

(Gelman, 2008) which takes the variable to center (subtract mean) and scale (center and then divides by standard

deviation).

P_b: Average starting phenotype (mean phenotype of the population before selection)

P_s: Average selected phenotype (mean phenotype of the selected parents)

P_r: Average response phenotype (mean phenotype of the offspring generation)

h^2: Narrow-sense heritability

$h^2 = \frac{P_r - P_b}{P_s - P_b}$

Host genetic analysis

We tested allelic richness (A_R), gene diversity (H_S), $F_{IS,}$ and identity disequilibrium (g2) among our artificial

selection lines to evaluate whether the differences in phenotype were attributable to variation in inbreeding

(Smallbone et al., 2016). Allelic richness is the first population genetic metric to reflect bottleneck effects, which

decrease the effective population size and facilitate reductions in heterozygosity due to inbreeding (Nei et al., 1975).

Significant differences in allelic richness among populations are indicative of variation in increases in the

probability that inbreeding may occur. Gene diversity measures the mean heterozygosity per individual per

population, under inbreeding conditions homozygosity increases but gene diversity is not immediately lost

(Charlesworth, 2003). Positive values in F_{IS} can be generated by inbreeding but may also be artificially elevated by

technical artifacts such as large allele drop out or null alleles, which can artificially inflate a deficiency of

heterozygotes (De Meeûs, 2018; Nei et al., 1975). We therefore also estimated identity disequilibrium (g_2), the

correlation of heterozygosity across loci resulting from inbreeding, which is robust in the presence of null alleles

(David et al., 2007). Concurrence among population genetic metrics consistent with patterns inbreeding are needed

to determine if phenotypic differences among populations are attributable to inbreeding.　Allelic richness (A_R)

(rarefied to the lowest population sample size), gene diversity (H_S) and F_{IS}, was calculated in FSTAT 2.39 (Goudet,

01 2001).　Significance levels for F_{IS} were assessed in GENEPOP (web version 4.2;(Raymond & Rousset, 1995)

with 5000 iterations followed by a Bonferroni correction for multiple comparisons. Tests for statistically significant

differences allelic richness (A_R), gene diversity (H_S) and F_{IS} among groups were calculated in FSTAT with 2000

permutations. Identity disequilibrium (g^2) is the difference between joint identity by descent and the product of the

separate probabilities of identity by decent for two loci. g^2 is estimated with microsatellite markers as the excess of

double heterozygous genotypes at all pairs of loci relative to a random distribution of heterozygous genotypes

among individuals at each locus (Cockerham & Weir, 1973; David et al., 2007).　g_2 values significantly greater than

0 are indicative of identity disequilibrium (ID). ID was calculated in inbreedR (Stoffel et al., 2016).

Microbial alpha and beta diversity

Raw sequence data were processed using the bioinformatics program QIIME2 version 2019.7 (Bolyen et

al., 2019). The resulting ASV table was rarefied to 3,654 sequences per sample, which retained all but 1 sample and

excluded kit controls. Within QIIME2, the DADA2 pipeline was used to calculate the unweighted and weighted

UniFrac distances (Lozupone & Knight, 2005) in a pairwise fashion between all pairs of samples to create distance

matrices. Significant differences in microbial community composition/membership across line, sex, length, and

body condition were tested using a PERMANOVA for the adonis package in R (Anderson, 2001). We used Principal

Coordinate Analysis (PCoA) plots with generated in R version 3.4.3. (R Core Team, 2020) to visualize all factors

tested. We also calculated the following alpha diversity indices using QIIME2: the number observed number of

ASVs, Shannon diversity, and Faith's phylogenetic diversity. We compared mean values of these metrics using

Pairwise Kruskal-Wallis preformed on QIIME2 for line and sex.

Bacterial community typing

We used Dirichlet multinomial mixture models (DMM) (Holmes et al., 2012) in R v4.0.0 and the Dirichlet

Multinomial v1.16.0 package (Morgan, 2023) to assign all samples (n=51) to three different bacterial community

types based on the relative frequency of taxa. We determined the number of community types by comparing the

Laplace approximation of the negative log models and identifying the point at Dirichlet components that resulted

in the best model fit (see supplemental materials). All formal statistical inferences are based on the models relating

community types to our outcomes of interest, with any findings being statistically supported by the data.

Infection analysis with bacterial community types

To test if the bacterial community types significantly predicted infection severity, we included bacterial community

types as a fixed effect in the same glmm model used for artificial selection lines analysis, subsetting to only swabbed

fish. We used AUC as the response variable in a glmm in the glmmTMB package, v. 1.1.2.3 (Brooks et al., 2017)

(Gamma error family and logit link function within R v.4.3.2 ((R Core Team, 2020); see supplementary material).

As fixed main effects, we included bacterial community type, artificial selection (line), fish sex, length (On day 0

and controlling for sex differences, see supplementary material), body condition (scaled mass index (Peig & Green,

2009)), generation (gen), number of parasites starting infection (dose), and trial date (Julien date). We also included

a two-way interaction between sex and line based on lower AIC.

Comparing the Predictive Power in our GLMM Models

To compare how well our two GLMM models—one with bacterial community types and the other with the artificial

selection lines only—explained variation in the response variable (AUC), we conducted an ANOVA-like

comparison using the anova function. All three models shared the same fixed effects, differing only in whether

"line" or "bacterial community type" or both was included as a fixed effect (see supplementary material). All models

used the same subset of swabbed fish (n=31) to allow for a fair comparison.

Results

Infection of 3^{rd}-6^{th} generation offspring

The artificial selection lines significantly predicted infection severity (***Figure 1***; χ^2=37.74, P =6.4e-09), and had a

significant interaction with sex (χ^2=13.51, P =0.0012). Additionally, sex and dose were both important to infection

severity in this overall model (sex; χ^2= 24.29, P =8.3e-07 & dose; χ^2= 5.11, P =0.024), but length, scaled mass

index, JulianDate, and generation were not. As a post hoc test to evaluate whether the effect of line was significant

among both sexes, we subset the data by sex and re-ran the model. We found that artificial selection line was still

significant in female and male guppies (female; χ^2=24.33, P =5.2e-06 & male; χ^2=17.57, P =0.00015). A post hoc

Tukey test revealed that the female guppies from susceptible and resistant artificial selection lines differed

significantly from control line, with susceptible female having high infection severities and resistant females lower

compared to control (Susceptible; z=3.49, P<0.001; Resistant; z =-2.74, P=0.016). The male guppies from the

susceptible artificial selection line, but not the resistant line, significantly differed compared to the control line

(Susceptible: z=4.19, P<0.001; Resistant: z =1.86, P=0.15).

Tissue-specific tolerance of 3^{rd}-6^{th} generation offspring

The artificial selection lines significantly differed in relative tolerance (***Figure 2***; χ^2=14.56, P =0.00069), with a

significant interaction with sex (χ^2=10.28, P =0.0059). Additionally, sex and length were both important to tolerance

(length; χ^2=5.94, P =0.015). As a post hoc test to evaluate whether this correlation was significantly different from

0 among either sex, we subset the data by sex and re-ran the model. We found that artificial selection line was only

significant in female guppies (female; χ^2=23.81, P =6.7e-06) and smaller females were less tolerant (length, female;

χ^2=6.96, P =0.0083). However, no fixed effect was significant in the male subset. JulianDate was not significant in

explaining variation within our models. A post hoc Tukey test revealed that resistant line females differed

significantly from control and susceptible line females, with resistant females having lower tolerance and compared

to the other lines (control; z=-3.92, P=0.0003; susceptible; z =4.05, P=0.00016). We did not find that the bacterial

community types significantly predicted tolerance within our study.

Resistance-tolerance Tradeoff

The polynomial regression model revealed a significant relationship between tissue-specific tolerance and infection

severity (lower infection severity indicates higher resistance). AUC explained a substantial proportion of the

variation in tissue-specific tolerance (logAUC: F = 61.95, p < 4e-16, see supplement). No other fixed effects

included in our model were significant.

Host genetic analysis

We tested gene diversity (H_s), allelic richness (A_R), inbreeding coefficient ($F_{IS)}$, , and identity disequilibrium (g^2)

among our artificial selection lines to evaluate whether the difference in phenotype were attributable to variation in

inbreeding. In this case the reduction in population size post-capture can be seen clearly in the decrease in allelic

richness among all of the experimental infection groups. The reduction in allelic richness among all experimental

infection groups relative to the wild caught founders likely reflects this post-capture reduction (*Supplement*

materials; Table 1). We found no significant difference in A_R, (p= 0.589), H_s (p=0.73) or F_{IS} (p=0.42) among the

three lines. We found no significant identity disequilibrium in any of the three lines (*Supplemental 1, Table 2*).

Narrow sense heritability of resistance and tolerance

We found that the heritability of 'meanworm' (the metric used for parental selection, i.e., average number of

parasites throughout infection) was $h^2=0.79$ in the susceptible line, and for the resistant lines $h^2=0.41$. We found

that heritability of tolerance was $h^2=0.88$ in the susceptible line, and $h^2=0.77$ for the resistant line.

Bacterial community types

All skin swab samples were assigned to a community type with the greatest posterior probability. All samples had

a posterior probability of 98% or higher. Each bacterial community type contained between 10.0–66.8% of all

samples (n = 5-33), and no samples were undefined. For top contributing bacterial genera see ***Figure S2*** in

supplemental material.

Skin Microbiome Variation

The artificial selection lines did not significantly differ any skin microbial diversity metrics tested (see

supplemental materials). Interestingly, the sex of the fish was significant in explaining Shannon's index and

observed ASVs, with females having significantly higher alpha diversities. (Kruskal–Wallis pairwise test, SI; H =

4.06, q = 0.04 & ASVs; H = 4.88, q = 0.03). Consistent with Kramp *et al*. 2022, female and male fish had

significantly different Unweighted Unifrac beta diversity (PERMANOVA; F=1.56, Pr=0.006).

Infection outcomes with bacterial community types

DMM modeling found that the lowest Laplace approximation model fit was three bacterial clusters (See

supplemental materials). To confirm that the three bacterial community types (named A, B, and C) within our fish

skin samples were distinct community types, we ran a PERMANOVA (PERMANOVA: F = 7.85, p = 0.001, ***Figure***

3a). Male and female fish numbers were balanced among the bacterial community types A and B, and bacterial

community types clustered evenly across the artificial selection lines (***Figure 3a***; Fisher's exact test, p = 0.8).

Additionally, we found no interaction between line and bacterial community types in our model ($\chi2 = 7.4$ p=0.116).

We found that bacterial community type was a significant in explaining AUC ($\chi2 = 19.83$, p=4.9e-5, ***Figure 3b).***

We found that the bacterial community type A of the skin microbiome on Trinidadian guppies had significantly

reduced infection severity compared to bacterial community type B & C(B: z=3.23, p=0.0034 & C: z=4.92, p=1e-
04). A post-hoc test showed that a model including both artificial selection line and bacterial community type is
significantly better at explaining variation in infection severity than models with either variable alone (*Figure 4*;
Chisq=13.4, Pr=0.0012), see supplement materials.

Discussion

Our artificial selection lines demonstrated the striking heritability of parasite infection resistance and
tolerance. We did not find that the microbiome is correlated or differed significantly between our three selection
lines. Surprisingly, we did find that the microbiome community types predicted infection severity independently
from host lines through bacterial community types found by DMM machine learning, though the microbiome did
not predict metrics of host tolerance. We also observed notable differences between males and females, both in the
response to artificial selection and in the skin microbiome. Below, we explore each of these findings in turn. We
first analyze the implications of the guppies' responses to artificial selection, followed by a discussion on the
independence of the microbiome in predicting resistance. Finally, we delve into the observed differences between
males and females and their potential impact on parasite resistance and microbiome interactions.

Our study is the first, to our knowledge, to use artificial selection to select for guppies that are susceptible
or resistant to their specialist parasite, *G. turnbulli*. Previous studies have examined parasite resistance evolution
through natural population variation (Dargent et al., 2013; van Oosterhout et al., 2003). It appears that selection for
both resistance and susceptibility traits can be realized within a single generation. Our study expands past
investigations by considering susceptibility selection, and not resistance alone, we revealed that both susceptible
male and female guppies were significantly different to control line fish. Selection for susceptible fish could have
multiple genetic mechanisms, one being tolerance and other being loss of resistance, and evidence for evolved
tolerance has been shown in wild guppy populations (Stephenson, van Oosterhout, & Cable, 2015). Our
experimental design could have imposed selection for tolerance, particularly among the susceptible line, as only
those individuals that are able to survive and successfully reproduce after experiencing severe infection loads
contributed to subsequent generations (Medzhitov et al., 2012; Råberg et al., 2007). Tolerance and resistance likely

have different genetic basis, and tradeoffs between the traits have been demonstrated (Mikaberidze & McDonald, 2020; Råberg et al., 2007). Consistent with such a tradeoff, and that our experimental design imposed selection for tolerance in the susceptible line, we found that resistant line fish had lower tolerance compared to control and susceptible line fish. Also, we found that this resistance-tolerance tradeoff exists among all fish sampled: the more resistant an individual is the lower tolerance the individual appears to have. However, it is important to note that research has shown that these tradeoffs do not always occur and both tolerance and resistance traits can coexist within a single individual host, and within a population (Hardy et al., 2024; Pagán & García-Arenal, 2018). Additionally, tolerance can have multiple mechanisms, which we do not touch on, but have profound effects on parasite virulence (Miller et al., 2006).

Our estimates of narrow-sense heritability for 'meanworm' and tolerance were high, particularly for our susceptible lines, though our values are still comparable to heritability estimates to our previous studies (Weiler et al., 2024), and to male guppy ornaments (Morris et al., 2020; van der Bijl et al., 2023), which are correlated with *G. turnbulli* resistance (Stephenson et al., 2020). The greater heritability of susceptibility, and the indication of their higher tolerance may be important for understanding how the behavioral and immunological defenses of Trinidadian guppies interact with the spread and evolution of parasites (Walsman et al., 2022, 2024). In some guppy populations, the prevalence of infection is very high (>90%; (Clark et al., 2023; Stephenson, van Oosterhout, Mohammed, et al., 2015)), selecting for higher tolerance of infection and lower resistance to infection via the 'resistance is futile' effect which can resemble the evolution of tolerance (Walsman et al., 2023). It is important to note that we did not find any notable differences in the three selections line in our genetic metrics: F_{IS} (inbreeding coefficient) or g^2 (identity disequilibrium). These results taken together indicate that the differences we found within our selection lines were not due to inbreeding and appear to be heritable.

Host characteristics such as immune system, metabolism, and growth rate have been shown to interact with the microbiome (Hosseini et al., 2016; Linehan et al., 2018; Speer, 2022; Zheng et al., 2020). More recent evidence has shown that life history and size selection experiments have an indirect effect on guppy resistance to *G. turnbulli* (Bartuseviciute et al., 2022), suggesting more unknown complexities of parasite resistance interacting with

metabolism and growth rates of the host. However, we did not find that the microbiome composition, alpha

diversity, or bacterial community types differed significantly between the resistance, susceptibility, or control lines,

nor did the bacterial communities predict tolerance within our fish. Therefore, our findings suggest that selection

line host genetics and host-associated skin microbiome significantly predict parasite infection severity, but not

tolerance, and that they could do so independently and additively within the host. Several studies have found that

fish skin microbiomes are highly variable and may not reflect the host's genetics but drift due to stochasticity (Jones

et al., 2022; Sadeghi et al., 2023; Wang et al., 2023), in agreement with our results. Nevertheless, while host genetics

within our artificial selection lines do not appear to influence the skin microbiome, that many studies including ours

have shown a strong correlation between bacterial communities and infection severity nevertheless points to

important interactions between the skin microbiome and parasite (Le Sage et al., 2021; Scheifler et al., 2022; Speer,

2022; Xavier et al., 2019). While within our study the skin microbiome was not altered by artificial selection, a

recent study that artificially selected for body size in fish significantly altered the host-associated microbiome and

found that the microbiome in the fast-growing line had the ability to resist parasite colonization (Piazzon et al.,

2020). Piazzon's study indicates that selection on other traits, i.e., size or color could have important effects on the

microbiome that could alter their susceptibility to parasites. Indeed, Bartuseviciute *et al.* that found life history and

size selection experiments have been shown to have an indirect effect on guppies resistance to *G. turnbulli*

(Bartuseviciute et al., 2022), potentially mediated through the microbiome.

The microbiome's role in determining resistance has been seen in other systems, where pre-infection

microbiome composition, alpha diversity, or microbiome function is a powerful tool to predict individuals that are

most susceptible to parasites within a population (Bates et al., 2022; Lee et al., 2019). Our results further emphasize

the potential of the microbiome in predicting infectious disease dynamics and lend further credence to the

microbiome having an active role on host-parasite interactions and not passively reflecting host's genetics. Further

investigation of our bacterial community findings were found to be similar to a recent study of skin microbial

metabolism function in amphibian chytridiomycosis (Bates et al., 2022): we found five of the eight ASVs reported

to be indicator taxa of epizootic dynamics; *Burkholderiales, Pseudomonadales, Sphingomonadales, Rhizobiales,*

Flavobacteriales were also important in our selected bacterial community types (see ***Figure S2*** for all contributing species). These results taken together indicate a complex and dynamic relationship between skin microbiome, ectoparasite infection, and host genetics. (Casadei et al., 2023; Douglas et al., 2021).

Our results show intricate patterns of evolution between the sexes in the multiple defenses against their specialist parasite *G. turnbulli*. We found significant differences between the sexes in the extent to which artificial selection line predicted infection severity This finding is similar to those found in Dargent et al., where male guppies did not show significant evolution of resistance to *Gyrodactylus*, compared to females (Dargent et al., 2016). A potential explanation for this finding is that male guppies in the ancestral population already had greater resistance than did females due to sexual selection, and thus perhaps the evolution of further resistance among males is constrained. Indeed, females in many populations prefer males with larger orange ornaments (Endler et al., 2001; Houde, 1988, 1997), with implications for disease dynamics (Rovenolt et al., 2024). Larger orange ornaments indicate male resistance to this parasite (Stephenson et al., 2020). We also see that males tend to develop lower infection loads in general in the lab (Johnson et al., 2011), are less likely, and less heavily infected in surveys of natural populations (Clark et al., 2023; Gotanda et al., 2013; Stephenson, van Oosterhout, Mohammed, et al., 2015). Furthermore, our experimental design allowed female preference for such males to continue in our experiment, potentially resulting in the overall high and invariable resistance we observe in males.

We saw differences in the evolution of susceptibility between males and females in our lines. Natural selection on female guppies for larger body sizes and longevity has likely resulted in the evolution of increased infection tolerance compared to males (Stephenson et al., 2016).That we saw the tolerance-resistant tradeoff only within our female guppies could have fundamental links to how male and female guppies respond to parasite selection pressure. However, that we saw males within the susceptible lines were significantly different to control lines in infection severity, but not tolerance, might suggest that the sexes face different tradeoffs. This possibility will hopefully inspire exciting and novel exploration of the evolutionary dynamics of resistance and tolerance between the sexes.

We did not find an interaction between bacterial community types and sex within our main models, and

males and females were even spread across the two largest community types A and B. Though, we did find that sex

was a significant indicator of both alpha and beta diversity metrics. Sex has been seen to have influence on skin

microbiome in other studies (Jung et al., 2023; Rodríguez-Barreto et al., 2023; Sadeghi et al., 2023). The observed

sex-specific differences in both artificial selection and microbiome suggest potential co-evolutionary patterns

between host sex and microbiome in parasite resistance that need to be disentangled with methodical experimental

designs.

This study is the first to artificially select guppies for resistance and susceptibility to *G. turnbulli*, revealing

novel insights into the evolution of host-parasite dynamics. By incorporating both resistant and susceptible lines,

we uncovered evidence for a tolerance-resistance tradeoff, particularly in females, where resistant fish showed

lower tolerance to infection. Surprisingly, while the microbiome did not differ across selection lines, bacterial

community types were significantly associated with infection outcomes, suggesting that the microbiome plays an

active role in shaping disease dynamics. This study is the first to highlight the role of the microbiome in the well-

known system between Trinidadian guppies and *G. turnbulli* parasites. These findings have broader implications

for understanding how host genetics and microbiome interact to influence disease susceptibility and tolerance, with

potential applications in disease management, conservation, and evolutionary biology. The study also highlights

sex-specific evolutionary trajectories in response to parasite pressure, encouraging further exploration into how sex

differences shape resistance and tolerance in host populations.

References

Bakke, T. A., Cable, J., & Harris, P. D. (2007). The biology of gyrodactylid monogeneans: the "Russian-doll

killers." *Advances in Parasitology*, *64*, 161–376.

Bates, K. A., Sommer, U., Hopkins, K. P., Shelton, J. M. G., Wierzbicki, C., Sergeant, C., Tapley, B., Michaels, C.

J., Schmeller, D. S., Loyau, A., Bosch, J., Viant, M. R., Harrison, X. A., Garner, T. W. J., & Fisher, M. C.

(2022). Microbiome function predicts amphibian chytridiomycosis disease dynamics. *Microbiome*, *10*(1),

44.

Benson, A., Pifer, R., Behrendt, C. L., Hooper, L. V., & Yarovinsky, F. (2009). Gut commensal bacteria direct a protective immune response against Toxoplasma gondii. *Cell Host & Microbe*, *6*(2), 187–196.

Bernardo-Cravo, A. P., Schmeller, D. S., Chatzinotas, A., Vredenburg, V. T., & Loyau, A. (2020). Environmental Factors and Host Microbiomes Shape Host-Pathogen Dynamics. *Trends in Parasitology*, *36*(7), 616–633.

Bordenstein, S. R., & Theis, K. R. (2015). Host Biology in Light of the Microbiome: Ten Principles of Holobionts and Hologenomes. *PLoS Biology*, *13*(8), e1002226.

Brucker, R. M., Harris, R. N., Schwantes, C. R., Gallaher, T. N., Flaherty, D. C., Lam, B. A., & Minbiole, K. P. C. (2008). Amphibian chemical defense: antifungal metabolites of the microsymbiont Janthinobacterium lividum on the salamander Plethodon cinereus. *Journal of Chemical Ecology*, *34*(11), 1422–1429.

de Vos, W. M., Tilg, H., Van Hul, M., & Cani, P. D. (2022). Gut microbiome and health: mechanistic insights. *Gut*, *71*(5), 1020–1032.

Haehling, M. B., Toscano, J. H. B., Giraldelo, L. A., Santos, I. B., Esteves, S. N., Benavides, M. V., Tholon, P., & Chagas, A. C. S. (2021). Is selection for resistance and resilience to Haemonchus contortus possible in Morada Nova sheep? *Small Ruminant Research: The Journal of the International Goat Association*, *201*, 106447.

Hanage, W. P. (2014, August 20). *Microbiology: Microbiome science needs a healthy dose of scepticism*. Nature Publishing Group UK. https://doi.org/10.1038/512247a

Holmes, I., Harris, K., & Quince, C. (2012). Dirichlet multinomial mixtures: generative models for microbial metagenomics. *PloS One*, *7*(2), e30126.

Hopkins, S. R., Wojdak, J. M., & Belden, L. K. (2017). Defensive Symbionts Mediate Host-Parasite Interactions at Multiple Scales. *Trends in Parasitology*, *33*(1), 53–64.

Le Sage, E. H., LaBumbard, B. C., Reinert, L. K., Miller, B. T., Richards-Zawacki, C. L., Woodhams, D. C., & Rollins-Smith, L. A. (2021). Preparatory immunity: Seasonality of mucosal skin defences and Batrachochytrium infections in Southern leopard frogs. *The Journal of Animal Ecology*, *90*(2), 542–554.

Lee, K. H., Gordon, A., Shedden, K., Kuan, G., Ng, S., Balmaseda, A., & Foxman, B. (2019). The respiratory microbiome and susceptibility to influenza virus infection. *PloS One*, *14*(1), e0207898.

Li, F., Li, C., Chen, Y., Liu, J., Zhang, C., Irving, B., Fitzsimmons, C., Plastow, G., & Guan, L. L. (2019). Host

genetics influence the rumen microbiota and heritable rumen microbial features associate with feed

efficiency in cattle. *Microbiome*, *7*(1), 92.

Libertucci, J., & Young, V. B. (2019). The role of the microbiota in infectious diseases. *Nature Microbiology*, *4*(1),

35–45.

Lorgen-Ritchie, M., Clarkson, M., Chalmers, L., Taylor, J. F., Migaud, H., & Martin, S. A. M. (2022). Temporal

changes in skin and gill microbiomes of Atlantic salmon in a recirculating aquaculture system – Why do

they matter? *Aquaculture* , *558*, 738352.

Mazel, F., Davis, K. M., Loudon, A., Kwong, W. K., Groussin, M., & Parfrey, L. W. (2018). Is Host Filtering the

Main   Driver   of   Phylosymbiosis   across   the   Tree   of   Life?   *MSystems*,   *3*(5).

https://doi.org/10.1128/mSystems.00097-18

Minniti, G., Rød Sandve, S., Padra, J. T., Heldal Hagen, L., Lindén, S., Pope, P. B., Ø Arntzen, M., & Vaaje-

Kolstad, G. (2019). The Farmed Atlantic Salmon (Salmo salar) Skin-Mucus Proteome and Its Nutrient

Potential for the Resident Bacterial Community. *Genes*, *10*(7). https://doi.org/10.3390/genes10070515

Mueller, U. G., Juenger, T. E., Kardish, M. R., Carlson, A. L., Burns, K. M., Edwards, J. A., Smith, C. C., Fang,

C.-C., & Des Marais, D. L. (2021). Artificial Selection on Microbiomes To Breed Microbiomes That Confer

Salt Tolerance to Plants. *MSystems*, *6*(6), e0112521.

Nelson, P., & May, G. (2020). Defensive Symbiosis and the Evolution of Virulence. *The American Naturalist*,

196(3), 333–343.

Parks, D. H., & Beiko, R. G. (2013). Measures of phylogenetic differentiation provide robust and complementary

insights into microbial communities. *The ISME Journal*, *7*(1), 173–183.

Piazzon, M. C., Naya-Català, F., Perera, E., Palenzuela, O., Sitjà-Bobadilla, A., & Pérez-Sánchez, J. (2020). Genetic

selection for growth drives differences in intestinal microbiota composition and parasite disease resistance

in gilthead sea bream. *Microbiome*, *8*(1), 168.

Reverter, M., Sasal, P., Suzuki, M. T., Raviglione, D., Inguimbert, N., Pare, A., Banaigs, B., Voisin, S. N., Bulet,

P., & Tapissier-Bontemps, N. (2020). Insights into the Natural Defenses of a Coral Reef Fish Against Gill

557   Ectoparasites: Integrated Metabolome and Microbiome Approach. *Metabolites*, *10*(6).

558   https://doi.org/10.3390/metabo10060227

559 Shanahan, F. (2012). The microbiota in inflammatory bowel disease: friend, bystander, and sometime-villain.

560   *Nutrition Reviews*, *70 Suppl 1*, S31-7.

561 Smallbone, W., Ellison, A., Poulton, S., van Oosterhout, C., & Cable, J. (2021). Depletion of MHC supertype during

562   domestication can compromise immunocompetence. *Molecular Ecology*, *30*(3), 736–746.

563 Smith, M. I., Yatsunenko, T., Manary, M. J., Trehan, I., Mkakosya, R., Cheng, J., Kau, A. L., Rich, S. S.,

564   Concannon, P., Mychaleckyj, J. C., Liu, J., Houpt, E., Li, J. V., Holmes, E., Nicholson, J., Knights, D.,

565   Ursell, L. K., Knight, R., & Gordon, J. I. (2013). Gut microbiomes of Malawian twin pairs discordant for

566   kwashiorkor. *Science*, *339*(6119), 548–554

567 Abt, M. C., Osborne, L. C., Monticelli, L. A., Doering, T. A., Alenghat, T., Sonnenberg, G. F., Paley, M. A.,

568   Antenus, M., Williams, K. L., Erikson, J., Wherry, E. J., & Artis, D. (2012). Commensal bacteria calibrate

569   the activation threshold of innate antiviral immunity. *Immunity*, *37*(1), 158–170.

570 Alberto, F. (2013). MsatAllele: Visualizes the scoring and binning of microsatellite fragment sizes. *R Package*

571   *Version*.

572 Amenyogbe, N., Kollmann, T. R., & Ben-Othman, R. (2017). Early-Life Host-Microbiome Interphase: The Key

573   Frontier for Immune Development. *Frontiers in Pediatrics*, *5*, 111.

574 Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance. *Austral Ecology*, *26*(1),

575   32–46.

576 Bartuseviciute, V., Diaz Pauli, B., Salvanes, A. G. V., & Heino, M. (2022). Size-selective harvesting affects the

577   immunocompetence of guppies exposed to the parasite Gyrodactylus. *Proceedings. Biological Sciences /*

578   *The Royal Society*, *289*(1981), 20220534.

579 Bates, K. A., Sommer, U., Hopkins, K. P., Shelton, J. M. G., Wierzbicki, C., Sergeant, C., Tapley, B., Michaels, C.

580   J., Schmeller, D. S., Loyau, A., Bosch, J., Viant, M. R., Harrison, X. A., Garner, T. W. J., & Fisher, M. C.

581   (2022). Microbiome function predicts amphibian chytridiomycosis disease dynamics. *Microbiome*, *10*(1),

582   44.

583 Becher, S. A., Russell, S. T., & Magurran, A. E. (2002). Isolation and characterization of polymorphic
584    microsatellites in the Trinidadian guppy (Poecilia reticulata). *Molecular Ecology Notes*, *2*(4), 456–458.

585 Becker, M. H., Brucker, R. M., Schwantes, C. R., Harris, R. N., & Minbiole, K. P. C. (2009). The bacterially
586    produced metabolite violacein is associated with survival of amphibians infected with a lethal fungus.
587    *Applied and Environmental Microbiology*, *75*(21), 6635–6638.

588 Benson, A., Pifer, R., Behrendt, C. L., Hooper, L. V., & Yarovinsky, F. (2009). Gut commensal bacteria direct a
589    protective immune response against Toxoplasma gondii. *Cell Host & Microbe*, *6*(2), 187–196.

590 Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., Alexander, H., Alm, E.
591    J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K., Brejnrod, A., Brislawn, C. J., Brown,
592    C. T., Callahan, B. J., Caraballo-Rodríguez, A. M., Chase, J., … Caporaso, J. G. (2019). Reproducible,
593    interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology*, *37*(8),
594    852–857.

595 Bordenstein, S. R., & Theis, K. R. (2015). Host Biology in Light of the Microbiome: Ten Principles of Holobionts
596    and Hologenomes. *PLoS Biology*, *13*(8), e1002226.

597 Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Maechler,
598    M., & Bolker, B. M. (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated
599    Generalized Linear Mixed Modeling. In *The R Journal* (Vol. 9, Issue 2, pp. 378–400).
600    https://doi.org/10.32614/RJ-2017-066

601 Brucker, R. M., Harris, R. N., Schwantes, C. R., Gallaher, T. N., Flaherty, D. C., Lam, B. A., & Minbiole, K. P. C.
602    (2008). Amphibian chemical defense: antifungal metabolites of the microsymbiont Janthinobacterium
603    lividum on the salamander Plethodon cinereus. *Journal of Chemical Ecology*, *34*(11), 1422–1429.

604 Cable, J., & Van Oosterhout, C. (2007). The role of innate and acquired resistance in two natural populations of
605    guppies (Poecilia reticulata) infected with the ectoparasite Gyrodactylus turnbulli. *Biological Journal of*
606    *the Linnean Society. Linnean Society of London*, *90*(4), 647–655.

Casadei, E., Mani, A., Cisco, M., Vågnes, Ø., Salinas, I., & Patel, S. (2023). Sex-dependent effects of mechanical
delousing on the skin microbiome of broodstock Atlantic salmon (Salmo salar L.). *Scientific Reports*, *13*(1),
10824.

Charlesworth, D. (2003). Effects of inbreeding on the genetic diversity of populations. *Philosophical Transactions*
of the Royal Society of London. Series B, Biological Sciences, *358*(1434), 1051–1070.

Chen, J. Z., Kwong, Z., Gerardo, N. M., & Vega, N. M. (2024). Ecological drift during colonization drives within-
host and between-host heterogeneity in an animal-associated symbiont. *PLoS Biology*, *22*(4), e3002304.

Chen, X., Wu, R., Li, L., Zeng, Y., Chen, J., Wei, M., Feng, Y., Chen, G., Wang, Y., Lin, L., Luo, H., Chen, A.,
Zeng, Z., He, F., Bai, Y., Zhang, S., Han, Y., Wang, Z., Zhao, X., … Gong, S. (2023). Pregnancy-induced
changes to the gut microbiota drive macrophage pyroptosis and exacerbate septic inflammation. *Immunity*,
56(2), 336-352.e9.

Clark, D. R., Young, K. A., Kitzes, J., Moore, P. J., Evans, A. J., & Stephenson, J. F. (2023). Aggregation of
symbionts on hosts depends on interaction type and host traits. *Ecography*.
https://doi.org/10.1111/ecog.06858

Cockerham, C. C., & Weir, B. S. (1973). Descent measures for two loci with some applications. *Theoretical*
Population Biology, *4*(3), 300–330.

Cortazar-Chinarro, M., Richter-Boix, A., Rödin-Mörch, P., Halvarsson, P., Logue, J. B., Laurila, A., & Höglund, J.
(2024). Association between the skin microbiome and MHC class II diversity in an amphibian. *Molecular*
Ecology, *33*(1), e17198.

Dargent, F., Rolshausen, G., Hendry, A. P., Scott, M. E., & Fussmann, G. F. (2016). Parting ways: parasite release
in nature leads to sex-specific evolution of defence. *Journal of Evolutionary Biology*, *29*(1), 23–34.

Dargent, F., Scott, M. E., Hendry, A. P., & Fussmann, G. F. (2013). Experimental elimination of parasites in nature
leads to the evolution of increased resistance in hosts. *Proceedings. Biological Sciences / The Royal Society*,
280(1773), 20132371.

David, P., Pujol, B., Viard, F., Castella, V., & Goudet, J. (2007). Reliable selfing rate estimates from imperfect
population genetic data. *Molecular Ecology*, *16*(12), 2474–2487.

De Meeûs, T. (2018). Revisiting FIS, FST, Wahlund Effects, and Null Alleles. *The Journal of Heredity*, *109*(4),

634    446–456.

635 de Vos, W. M., Tilg, H., Van Hul, M., & Cani, P. D. (2022). Gut microbiome and health: mechanistic insights. *Gut*,

636    *71*(5), 1020–1032.

637 Debray, R., Herbert, R. A., Jaffe, A. L., Crits-Christoph, A., Power, M. E., & Koskella, B. (2022). Priority effects

638    in microbiome assembly. *Nature Reviews. Microbiology*, *20*(2), 109–121.

639 Dheilly, N. M., Martínez Martínez, J., Rosario, K., Brindley, P. J., Fichorova, R. N., Kaye, J. Z., Kohl, K. D., Knoll,

640    L. J., Lukeš, J., Perkins, S. L., Poulin, R., Schriml, L., & Thompson, L. R. (2019). Parasite microbiome

641    project: Grand challenges. *PLoS Pathogens*, *15*(10), e1008028.

642 Douglas, A. J., Hug, L. A., & Katzenback, B. A. (2021). Composition of the North American Wood Frog (Rana

643    sylvatica) Bacterial Skin Microbiome and Seasonal Variation in Community Structure. *Microbial Ecology*,

644    *81*(1), 78–92.

645 Endler, J. A., Basolo, A., Glowacki, S., & Zerr, J. (2001). Variation in response to artificial selection for light

646    sensitivity in guppies (Poecilia reticulata). *The American Naturalist*, *158*(1), 36–48.

647 Fox, J., & Weisberg, S. (2019). Companion to Applied Regression [R package car version 3.1-2]. *Sage*, *3rd*.

648    https://cran.r-project.org/web/packages/car/index.html

649 Fraser, B. A., & Neff, B. D. (2009). MHC class IIB additive and non-additive effects on fitness measures in the

650    guppy *Poecilia reticulata*. *Journal of Fish Biology*, *75*(9), 2299–2312.

651 Fraser, B. A., Ramnarine, I. W., & Neff, B. D. (2010). TEMPORAL VARIATION AT THE MHC CLASS IIB IN

652    WILD POPULATIONS OF THE GUPPY (*POECILIA RETICULATA*). *Evolution; International Journal*

653    *of Organic Evolution*, *64*(7), 2086–2096.

654 Garrigues, Q., Apper, E., Chastant, S., & Mila, H. (2022). Gut microbiota development in the growing dog: A

655    dynamic process influenced by maternal, environmental and host factors. *Frontiers in Veterinary Science*,

656    *9*, 964649.

657 Gelman, A. (2008). Scaling regression inputs by dividing by two standard deviations. *Statistics in Medicine*, *27*(15),

658    2865–2873.

Gorczyca, K., Obuchowska, A., Kimber-Trojnar, Ż., Wierzchowska-Opoka, M., & Leszczyńska-Gorzelak, B. (2022). Changes in the Gut Microbiome and Pathologies in Pregnancy. *International Journal of Environmental Research and Public Health*, *19*(16). https://doi.org/10.3390/ijerph19169961

Gotanda, K. M., Delaire, L. C., Raeymaekers, J. A. M., Pérez-Jvostov, F., Dargent, F., Bentzen, P., Scott, M. E., Fussmann, G. F., & Hendry, A. P. (2013). Adding parasites to the guppy-predation story: insights from field surveys. *Oecologia*, *172*(1), 155–166.

Goudet, J. (01 2001). FSTAT, Version. *1. 2: A Computer Program to Calculate F-Statistics. Heredity*, *86*, 485–486.

Hanage, W. P. (2014, August 20). *Microbiology: Microbiome science needs a healthy dose of scepticism*. Nature Publishing Group UK. https://doi.org/10.1038/512247a

Hand, D., Wallis, C., Colyer, A., & Penn, C. W. (2013). Pyrosequencing the canine faecal microbiota: breadth and depth of biodiversity. *PloS One*, *8*(1), e53115.

Hardy, B. M., Muths, E., Funk, W. C., & Bailey, L. L. (2024). Quantifying intraspecific variation in host resistance and tolerance to a lethal pathogen. *The Journal of Animal Ecology*. https://doi.org/10.1111/1365-2656.14106

Hartig, F. (2022). *DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression models. R package version 0.4. 6.*

Hawley, D. M., Etienne, R. S., Ezenwa, V. O., & Jolles, A. E. (2011). Does animal behavior underlie covariation between hosts' exposure to infectious agents and susceptibility to infection? Implications for disease dynamics. *Integrative and Comparative Biology*, *51*(4), 528–539.

Holmes, I., Harris, K., & Quince, C. (2012). Dirichlet multinomial mixtures: generative models for microbial metagenomics. *PloS One*, *7*(2), e30126.

Hopkins, S. R., Wojdak, J. M., & Belden, L. K. (2017). Defensive Symbionts Mediate Host-Parasite Interactions at Multiple Scales. *Trends in Parasitology*, *33*(1), 53–64.

Hosseini, M., Kolangi Miandare, H., Shabani, A., Hoseinifar, S. H., & Yarahmadi, P. (2016). Dietary Lactobacillus

acidophilus modulated skin mucus protein profile, immune and appetite genes expression in gold fish

(Carassius auratus gibelio). *Fish & Shellfish Immunology*, *59*, 149–154.

Houde, A. E. (1988). Genetic difference in female choice between two guppy populations. *Animal Behaviour*, *36*(2),

510–516.

Houde, A. E. (1997). Effect of artificial selection on male colour patterns on mating preference of female guppies.

Proceedings of the Royal Society of London. Series B: Biological Sciences, *256*(1346), 125–130.

Ichinohe, T., Pang, I. K., Kumamoto, Y., Peaper, D. R., Ho, J. H., Murray, T. S., & Iwasaki, A. (2011). Microbiota

regulates immune defense against respiratory tract influenza A virus infection. *Proceedings of the National

Academy of Sciences of the United States of America*, *108*(13), 5354–5359.

Janecka, M. J., Rovenolt, F., & Stephenson, J. F. (2021). How does host social behavior drive parasite non-selective

evolution from the within-host to the landscape-scale? *Behavioral Ecology and Sociobiology*, *75*(11), 150.

Jog, M. G., Sackett, M. E., Kisty, S. D., Hansen, J. A., & Stephenson, J. F. (2022). The behaviour of infected guppies

depends on social context, parasite tolerance and host sex. *Animal Behaviour*, *187*, 97–104.

Johnson, M. B., Lafferty, K. D., van Oosterhout, C., & Cable, J. (2011). Parasite transmission in social interacting

hosts: monogenean epidemics in guppies. *PloS One*, *6*(8), e22634.

Jones, E. W., Carlson, J. M., Sivak, D. A., & Ludington, W. B. (2022). Stochastic microbiome assembly depends

on context. *Proceedings of the National Academy of Sciences of the United States of America*, *119*(7),

e2115877119.

Jung, Y., Cui, H. S., Joo, S. Y., Lee, E. K., Seo, C. H., & Cho, Y. S. (2023). Sex differences in the skin microbiome

of burn scars. *Wound Repair and Regeneration: Official Publication of the Wound Healing Society [and]

the European Tissue Repair Society*, *31*(4), 547–558.

Kohl, K. D. (2020). Ecological and evolutionary mechanisms underlying patterns of phylosymbiosis in host-

associated microbial communities. *Philosophical Transactions of the Royal Society of London. Series B,

Biological Sciences*, *375*(1798), 20190251.

Kramp, R. D., Kohl, K. D., & Stephenson, J. F. (2022). Skin bacterial microbiome diversity predicts lower activity
levels in female, but not male, guppies, Poecilia reticulata. *Biology Letters*, *18*(8), 20220167.

Le Sage, E. H., LaBumbard, B. C., Reinert, L. K., Miller, B. T., Richards-Zawacki, C. L., Woodhams, D. C., &
Rollins-Smith, L. A. (2021). Preparatory immunity: Seasonality of mucosal skin defences and
Batrachochytrium infections in Southern leopard frogs. *The Journal of Animal Ecology*, *90*(2), 542–554.

Lee, K. H., Gordon, A., Shedden, K., Kuan, G., Ng, S., Balmaseda, A., & Foxman, B. (2019). The respiratory
microbiome and susceptibility to influenza virus infection. *PloS One*, *14*(1), e0207898.

Lim, S. J., & Bordenstein, S. R. (2020). An introduction to phylosymbiosis. *Proceedings. Biological Sciences / The*
Royal Society, *287*(1922), 20192900.

Linehan, J. L., Harrison, O. J., Han, S.-J., Byrd, A. L., Vujkovic-Cvijin, I., Villarino, A. V., Sen, S. K., Shaik, J.,
Smelkinson, M., Tamoutounour, S., Collins, N., Bouladoux, N., Dzutsev, A., Rosshart, S. P., Arbuckle, J.
H., Wang, C.-R., Kristie, T. M., Rehermann, B., Trinchieri, G., … Belkaid, Y. (2018). Non-classical
immunity controls Microbiota impact on skin immunity and tissue repair. *Cell*, *172*(4), 784-796.e18.

Lozupone, C., & Knight, R. (2005). UniFrac: a new phylogenetic method for comparing microbial communities.
Applied and Environmental Microbiology, *71*(12), 8228–8235.

Lush, J. L. (1937). *Animal Breeding Plans - Scholar's Choice Edition*. Bibliolife DBA of Bibilio Bazaar II LLC.

Luzzi, G. A., Merry, A. H., Newbold, C. I., Marsh, K., & Pasvol, G. (1991). Protection by alpha-thalassaemia
against Plasmodium falciparum malaria: modified surface antigen expression rather than impaired growth
or cytoadherence. *Immunology Letters*, *30*(2), 233–240.

Madhavi, R., & Anderson, R. M. (1985). Variability in the susceptibility of the fish host, Poecilia reticulata, to
infection with Gyrodactylus bullatarudis (Monogenea). *Parasitology*, *91*(3), 531–544.

Mazel, F., Davis, K. M., Loudon, A., Kwong, W. K., Groussin, M., & Parfrey, L. W. (2018). Is Host Filtering the
Main Driver of Phylosymbiosis across the Tree of Life? *MSystems*, *3*(5).
https://doi.org/10.1128/mSystems.00097-18

Medzhitov, R., Schneider, D. S., & Soares, M. P. (2012). Disease tolerance as a defense strategy. *Science*,
335(6071), 936–941.

Mikaberidze, A., & McDonald, B. A. (2020). A tradeoff between tolerance and resistance to a major fungal
pathogen in elite wheat cultivars. *The New Phytologist*, *226*(3), 879–890.

Miller, M. R., White, A., & Boots, M. (2006). The evolution of parasites in response to tolerance in their hosts: the
good, the bad, and apparent commensalism. *Evolution; International Journal of Organic Evolution*, *60*(5),
945–956.

Morgan, M. (2023). *DirichletMultinomial: Dirichlet-Multinomial Mixture Model Machine Learning for*
Microbiome Data. https://doi.org/10.18129/B9.bioc.DirichletMultinomial

Morris, J., Darolti, I., van der Bijl, W., & Mank, J. E. (2020). High-resolution characterization of male
ornamentation and re-evaluation of sex linkage in guppies. *Proceedings. Biological Sciences / The Royal*
Society, *287*(1937), 20201677.

Nater, A., Krützen, M., & Lindholm, A. K. (2008). Development of polymorphic microsatellite markers for the
livebearing fish Poecilia parae. *Molecular Ecology Resources*, *8*(4), 857–860.

Nei, M., Maruyama, T., & Chakraborty, R. (1975). The Bottleneck Effect and Genetic Variability in Populations.
Evolution; International Journal of Organic Evolution, *29*(1), 1–10.

Nelson, P., & May, G. (2020). Defensive Symbiosis and the Evolution of Virulence. *The American Naturalist*,
196(3), 333–343.

Ni, J., Wu, G. D., Albenberg, L., & Tomov, V. T. (2017). Gut microbiota and IBD: causation or correlation? *Nature*
Reviews. Gastroenterology & Hepatology, *14*(10), 573–584.

Ost, K. S., & Round, J. L. (2017). [Review of *A Few Good Commensals: Gut Microbes Use IFN-γ to Fight*
Salmonella]. *Immunity*, *46*(6), 977–979.

Pagán, I., & García-Arenal, F. (2018). Tolerance to plant pathogens: Theory and experimental evidence.
International Journal of Molecular Sciences, *19*(3). https://doi.org/10.3390/ijms19030810

Paterson, I. G., Crispo, E., Kinnison, M. T., Hendry, A. P., & Bentzen, P. (2005). Characterization of tetranucleotide
microsatellite markers in guppy (Poecilia reticulata). *Molecular Ecology Notes*, *5*(2), 269–271.

Peig, J., & Green, A. J. (2009). New perspectives for estimating body condition from mass/length data: the scaled
mass index as an alternative method. *Oikos* , *118*(12), 1883–1891.

Phillips, K. P., Cable, J., Mohammed, R. S., Herdegen-Radwan, M., Raubic, J., Przesmycka, K. J., van Oosterhout,
C., & Radwan, J. (2018). Immunogenetic novelty confers a selective advantage in host-pathogen
coevolution. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(7),
1552–1557.

Piazzon, M. C., Naya-Català, F., Perera, E., Palenzuela, O., Sitjà-Bobadilla, A., & Pérez-Sánchez, J. (2020). Genetic
selection for growth drives differences in intestinal microbiota composition and parasite disease resistance
in gilthead sea bream. *Microbiome*, *8*(1), 168.

R Core Team. (2020). *R: A language and environment for statistical computing* (Version v.4.1.0) [R Foundation
for Statistical Computing, Vienna, Austria.].

Råberg, L., Sim, D., & Read, A. F. (2007). Disentangling genetic variation for resistance and tolerance to infectious
diseases in animals. *Science*, *318*(5851), 812–814.

Raymond, M., & Rousset, F. (1995). GENEPOP (Version 1.2): Population Genetics Software for Exact Tests and
Ecumenicism. *The Journal of Heredity*, *86*(3), 248–249.

Reverter, M., Sasal, P., Suzuki, M. T., Raviglione, D., Inguimbert, N., Pare, A., Banaigs, B., Voisin, S. N., Bulet,
P., & Tapissier-Bontemps, N. (2020). Insights into the Natural Defenses of a Coral Reef Fish Against Gill
Ectoparasites: Integrated Metabolome and Microbiome Approach. *Metabolites*, *10*(6).
https://doi.org/10.3390/metabo10060227

Robertson, S. J., Lemire, P., Maughan, H., Goethel, A., Turpin, W., Bedrani, L., Guttman, D. S., Croitoru, K.,
Girardin, S. E., & Philpott, D. J. (2019). Comparison of Co-housing and Littermate Methods for Microbiota
Standardization in Mouse Models. *Cell Reports*, *27*(6), 1910-1919.e2.

Rodríguez-Barreto, D., Sanz-González, J. C., Martín, M. V., Arrieta, J. M., & Almansa, E. (2023). Sex-specific
bacterial microbiome variation in octopus vulgaris skin. *Frontiers in Microbiology*, *14*, 1233661.

Rovenolt, F., Walsman, J., Martin, C., Gotanda, K., & Stephenson, J. (2024). Parasite prevalence depends on female
preference: Integrating parasite-mediated sexual selection and infectious disease dynamics. In *Authorea*
Inc. https://doi.org/10.22541/au.172447480.06236699/v1

Sadeghi, J., Chaganti, S. R., Johnson, T. B., & Heath, D. D. (2023). Host species and habitat shape fish-associated

bacterial communities: phylosymbiosis between fish and their microbiome. *Microbiome*, *11*(1), 258.

Santos-Marcos, J. A., Mora-Ortiz, M., Tena-Sempere, M., Lopez-Miranda, J., & Camargo, A. (2023). Interaction

between gut microbiota and sex hormones and their relation to sexual dimorphism in metabolic diseases.

Biology of Sex Differences, *14*(1), 4.

Scheifler, M., Sanchez-Brosseau, S., Magnanou, E., & Desdevises, Y. (2022). Diversity and structure of sparids

external microbiota (Teleostei) and its link with monogenean ectoparasites. *Animal Microbiome*, *4*(1), 27.

Shanahan, F. (2012). The microbiota in inflammatory bowel disease: friend, bystander, and sometime-villain.

Nutrition Reviews, *70 Suppl 1*, S31-7.

Smallbone, W., Van Oosterhout, C., & Cable, J. (2016). The effects of inbreeding on disease susceptibility:

Gyrodactylus turnbulli infection of guppies, *Poecilia reticulata*. *Experimental Parasitology*, *167*, 32–37.

Speer, K. A. (2022). [Review of *Microbiomes mediate host-parasite interactions*]. *Molecular Ecology*, *31*(7), 1925–

1927.

Stephenson, J. F., & Adelman, J. S. (2022). The behavior of infected hosts: Behavioral tolerance, behavioral

resilience, and their implications for behavioral competence. *Animal Behavior and Parasitism*.

https://books.google.ca/books?hl=en&lr=&id=xSZ_EAAAQBAJ&oi=fnd&pg=PA287&ots=xW36pEEkt

3&sig=asg6-91ifwE5LAwCypVYASxqwO4

Stephenson, J. F., Kinsella, C., & Cable, J. (2016). A further cost for the sicker sex? Evidence for male-biased

parasite-induced vulnerability to predation. *Ecology*.

https://onlinelibrary.wiley.com/doi/abs/10.1002/ece3.2049

Stephenson, J. F., Stevens, M., Troscianko, J., & Jokela, J. (2020). The size, symmetry, and color saturation of a

male guppy's ornaments forecast his resistance to parasites. *The American Naturalist*, *196*(5), 597–608.

Stephenson, J. F., van Oosterhout, C., & Cable, J. (2015). Pace of life, predators and parasites: predator-induced

life-history evolution in Trinidadian guppies predicts decrease in parasite tolerance. *Biology Letters*, *11*(11),

20150806.

Stephenson, J. F., van Oosterhout, C., Mohammed, R. S., & Cable, J. (2015). Parasites of Trinidadian guppies:

evidence for sex- and age-specific trait-mediated indirect effects of predators. *Ecology*, *96*(2), 489–498.

Stoffel, M. A., Esser, M., Kardos, M., Humble, E., Nichols, H., David, P., & Hoffman, J. I. (2016). inbreedR: an R

package for the analysis of inbreeding based on genetic markers. *Methods in Ecology and Evolution*, *7*(11),

1331–1339.

Sylvia, K. E., Jewell, C. P., Rendon, N. M., St John, E. A., & Demas, G. E. (2017). Sex-specific modulation of the

gut microbiome and behavior in Siberian hamsters. *Brain, Behavior, and Immunity*, *60*, 51–62.

Trevelline, B. K., Stephenson, J. F., & Kohl, K. D. (2020). [Review of *Two's company, three's a crowd: Exploring*

how host-parasite-microbiota interactions may influence disease susceptibility and conservation of

wildlife]. *Molecular Ecology*, *29*(8), 1402–1405.

Truett, G. E., Heeger, P., Mynatt, R. L., Truett, A. A., Walker, J. A., & Warman, M. L. (2000). Preparation of PCR-

quality mouse genomic DNA with hot sodium hydroxide and tris (HotSHOT). *BioTechniques*, *29*(1), 52,

54.

van der Bijl, W., Shu, J., Goberdhan, V. S., Sherin, L., Cortazar-Chinarro, M., Corral-Lopez, A., & Mank, J. E.

(2023). Deep learning reveals the role of copy number variation in the genetic architecture of a highly

polymorphic sexual trait. In *bioRxiv* (p. 2023.09.29.560175). https://doi.org/10.1101/2023.09.29.560175

van Oosterhout, C., Harris, P. D., & Cable, J. (2003). Marked variation in parasite resistance between two wild

populations of the Trinidadian guppy, Poecilia reticulata (Pisces: Poeciliidae). *Biological Journal of the*

Linnean Society. Linnean Society of London, *79*(4), 645–651.

van Oosterhout, C., Joyce, D. A., & Cummings, S. M. (2006). Evolution of MHC class IIB in the genome of wild

and ornamental guppies, Poecilia reticulata. *Heredity*, *97*(2), 111–118.

Walsman, J. C., Duffy, M. A., Cáceres, C. E., & Hall, S. R. (2023). "Resistance Is Futile": Weaker Selection for

Resistance by Abundant Parasites Increases Prevalence and Depresses Host Density. *The American*

Naturalist, *201*(6), 864–879.

Walsman, J. C., Janecka, M. J., Clark, D. R., Kramp, R. D., Rovenolt, F., Patrick, R., Mohammed, R. S., Konczal,
M., Cressler, C. E., & Stephenson, J. F. (2022). Shoaling guppies evade predation but have deadlier
parasites. *Nature Ecology & Evolution*, *6*(7), 945–954.

Walsman, J. C., Lambe, M., & Stephenson, J. F. (2024). Associating with kin selects for disease resistance and
against tolerance. *Proceedings. Biological Sciences*, *291*(2023), 20240356.

Wang, L.-C., Chen, L.-H., Chiu, Y.-C., Liou, C.-Y., Chen, H.-C., Lu, C.-Y., & Chen, J.-L. (2023). Teleost skin
microbiome: An intimate interplay between the environment and the host immunity. *Fish & Shellfish*
Immunology, *139*, 108869.

Watanabe, T., Yoshida, M., Nakajima, M., & Taniguchi, N. (2005). Linkage mapping of AFLP and microsatellite
DNA markers with the body color- and sex-determining loci in the guppy (Poecilia reticulata). *Zoological*
Science, *22*(8), 883–889.

Weiler, I. L. G., Kramp, R. D., Rovenolt, F., & Stephenson, J. F. (2024). Sex-dependent effects of infection on
guppy reproductive fitness and offspring parasite resistance. In *bioRxiv* (p. 2023.10.29.564622).
https://doi.org/10.1101/2023.10.29.564622

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer.

Wilde, J., Slack, E., & Foster, K. R. (2024). Host control of the microbiome: Mechanisms, evolution, and disease.
Science (New York, N.Y.), *385*(6706), eadi3338.

Xavier, R., Mazzei, R., Pérez-Losada, M., Rosado, D., Santos, J. L., Veríssimo, A., & Soares, M. C. (2019). A
Risky Business? Habitat and Social Behavior Impact Skin and Gut Microbiomes in Caribbean Cleaning
Gobies. *Frontiers in Microbiology*, *10*, 716.

Zheng, D., Liwinski, T., & Elinav, E. (2020). Interaction between microbiota and immunity in health and disease.
Cell Research, *30*(6), 492–506.

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed Effects Models and Extensions*
in Ecology with R. Springer New York.

Figure 1. Trinidadian guppy artificial selection lines significantly predicted G. turnbulli infection severity (χ^2=37.74, P =6.4e-09). *Infection severity (worm days)* is the area under the curve of infection load over the course of an individual's infection. *Female guppies (n=44) susceptible and resistant artificial selection lines differed significantly from the control line (Susceptible-Control, z=3.49, P=0.001; Resistant-Control, z =-2.74, P=0.016). Male guppies (n=43) susceptible, but not resistant, artificial selection line differed significantly from control line (Susceptible-control, z=4.19, P<0.001; Resistant-Control, z =1.86, P=0.15). Box plots give the median (dark line), interquartile range (box) and values within 1.5× the interquartile range (whiskers) of the infection severity.*

Figure 2. Trinidadian guppy artificial selection lines significantly predicted Relative Tissue-Specific Tolerance (χ^2=14.56, P =0.00069). We quantified relative tissue-specific tolerance as the per-parasite percentage mass change across infection minus max tolerance value, greater values indicate greater tolerance. *Female guppies (n=44) resistant artificial selection lines differed significantly from the control and susceptible lines (Resistant-Control, z=-3.92, P=0.0003; Resistant-Susceptible, z =4.05, P=0.0001). Male guppies (n=43) artificial selection lines did not predict relative tissue specific tolerance (χ^2=2.08, P =0.35). Box plots give the median (dark line), interquartile range (box) and values within 1.5× the interquartile range (whiskers) of the infection severity.*

Figure 3. Uniform distribution of skin bacterial community types across guppy artificial selection lines suggests no association with selection mechanism. A. *Mosaic plot presenting proportion of Bacterial community types (A, B, C) across, Control (c), Resistant (r), and Susceptible lines (s). Bacterial communities do not show a significantly uneven distribution across lines (Fisher's exact test, p = 0.8).* **B**. *A Principal Coordinate (PCoA) analysis plot based on Weighted UniFrac similarity of fish skin microbial communities (n=52) across sex, bacterial community type, and artificial selection line. Percentages represent the proportion of variation explained by each axis. Bacterial community type was significant in explaining variation (PERMANOVA: F=7.37, P=0.0003). No significant pattern or clustering is was detected for artificial selection lines (PERMANOVA: F=0.68, P=0.89).*

Figure 4. Trinidadian guppy artificial selection lines and bacterial community types predict G. turnbulli infection severity. **A.** The artificial selection lines still significantly predicted infection severity (Anova; $\chi^2=9.39$, *P=0.009)* within the swabbed subset of fish (n=31). **B.** Bacterial community types differed in their infection severity (n=31). Bacterial community type A had significantly lower infection severity compared to bacterial community type B & C (A-B, *z=2.59, P=0.0034; A-C, z =4.92, P<0.001*). No interactions between bacterial community and artificial selection line (Anova; $\chi^2=7.4$, *P=0.116).* Post-hoc tests showed that a model including both line and bacterial community types were significantly better at explaining infection severity than either effect alone ($\chi^2=13.40$, *P=0.001).*