Artificial Selection and the Skin Microbiome Independently Predict Parasite Resistance

Rachael D. Kramp[1*], Mary J. Janecka[1,5], Nadine Tardent[2,3], Jukka Jokela[2,3], Kevin D. Kohl[1], Jessica F. Stephenson[1,2,3,4]

1. Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA 15260, USA

2. Department of Aquatic Ecology, EAWAG, Swiss Federal Institute of Aquatic Sci & Tech, Dübendorf 8600, Switzerland.

3. Institute for Integrative Biology, ETH Zürich, Zürich 8092, Switzerland.

4. Center for Adaptation to a Changing Environment, ETH Zürich, Zürich 8092, Switzerland.

5. Department of Biological Sciences, University of Texas at El Paso, El Paso, TX 79968, USA

*Author for correspondence: rdkramp@gmail.com

Abstract

1. Host responses to parasite infection involve several interacting systems. Host genetics determine much of the response, but it is increasingly clear that the host-associated microbiome also plays a role. Genetically determined systems and the microbiome can also interact; for example, the microbiome can modulate the host immune response, and *vice versa*. However, it remains unclear how such interactions between the host immune system and the microbiome may influence the host's overall response to parasites.

2. Host sex is an important source of variation in both immune response and the microbiome across systems, so interactions between them may also therefore differ between males and females, potentially explaining sex-specificity in response to parasites.

3. To investigate how host genetics and the microbiome interact to shape responses to parasites, we imposed truncation selection on the response of Trinidadian guppies (*Poecilia reticulata*) to the specialist ectoparasite *Gyrodactylus turnbulli*. Following experimental infection, we used the 30% of guppies with the lowest parasite load to found the resistant

line, the 30% with the highest load to found the susceptible line, and randomly chosen

uninfected fish to found the control line. After 3-6 generations of breeding without

parasites, we sampled the skin-associated microbiome and then infected fish from each

line. We applied machine-learning to identify bacterial community types across lines and

evaluated how selection line and community type explained variations in infection severity.

4. Our findings showed that among females, the resistant line had significantly lower

infection severity, while the susceptible line had higher infection severity than other lines.

Interestingly, females in the resistant line lost more mass per parasite, i.e., exhibited lower

parasite tolerance, than in the control or susceptible lines. Among males, the susceptible

line experienced higher infection severity than the other lines, but there was no difference

between the resistant and control lines.

5. We found that the microbiome and host genetics independently explain similar variation in

infection severity, highlighting the microbiome's active role in host-parasite interactions.

Our results also indicate sex specificity in the evolution of host defenses.

1. Introduction

Hosts use multiple strategies to defend themselves against parasites and pathogens (parasites

hereafter). Host genetic background is commonly credited with much of the defense against

parasites through the immune system, behavior, or other genetic/protein alterations, e.g.,

thalassemia (Labro, 2022). A recently highlighted alternative defense, that conferred by the host-

associated microbial communities or microbiome, may act indirectly through the host immune

system or directly via niche occupation or secreting anti-parasitic compounds (Bernardo-Cravo et

al., 2020), and may also exhibit heritable components. However, it remains unclear whether the

microbiome plays an active role in host-parasite infection or merely reflects the host's

health/immune status (Trevelline et al., 2020). While there is abundant evidence that the microbial

community can have a direct or indirect role in host-parasite interactions (Becker et al., 2009;

Nelson & May, 2020), there is also evidence that the microbiome is merely a bystander in some

diseases and a reflection of the host's genetics, diet, and general health (Ni et al., 2017).

Host genetics and the microbiome interact throughout host development, with complex feedbacks

on traits like parasite defense. On one hand, the host immune system relies on early microbial

colonization for proper development and the microbiome ultimately shapes immune cell

composition (Amenyogbe et al., 2017). On the other hand, hosts exert control over their symbionts,

shaping their associated microbiomes, via immune responses, barrier function, and physiological

regulation (Wilde et al., 2024). The host's genetic makeup therefore influences microbiome

composition, and specific microbial communities may therefore become heritable (Bordenstein &

Theis, 2015). However, distinguishing host genetic effects from factors such as priority effects,

drift, and other stochastic elements, is an ongoing challenge (Debray et al., 2022). Direct

manipulation of host genetics in traits like parasite resistance, where the microbiome plays a

known role, thus provides a useful tool in disentangling host genetic effects from microbial ones.

Both the host-associated microbiome and the immune system commonly differ between host sexes

(Markle & Fish, 2014). These differences are likely due to other correlated differences, for

example, high testosterone levels in the blood both suppress the immune system and significantly

alter gut microbial communities (Santos-Marcos et al., 2023), and both the immune system and

microbiome change during female reproductive cycles and pregnancy (Chen et al., 2023; Gorczyca

et al., 2022). Given that host sexes across taxa also differ in their response to parasites (Morrow, 2015), it is essential to test for sex-specific interactions between genetic resistance factors, the microbiome, and parasites.

Trinidadian guppies (*Poecilia reticulata*) and their specialist parasite *Gyrodactylus turnbulli* provide a rich context to study host-parasite dynamics. *Gyrodactylus* spp. feed and reproduce (both sexually and asexually, generation time 24hrs) on host skin. Guppies show a large variation in parasite resistance across populations and ecological contexts (Dargent et al., 2016; Phillips et al., 2021; Stephenson, et al., 2015; Van Oosterhout et al., 2003). Guppy resistance is at least partly genetically determined and heritable, with sex differences (Dargent et al., 2016; Van Oosterhout et al., 2003; Weiler et al., 2024). Innate and acquired immunity are important for *G. turnbulli* resistance, although the underlying mechanisms remain unclear (Cable & Van Oosterhout, 2007; Mohammed et al., 2016). Research has largely focused on the major histocompatibility complex (MHC) for genetic-based resistance to *G. turnbulli* (Fraser et al., 2010; Fraser & Neff, 2009; Phillips et al., 2018). Notably, skin bacterial alpha diversity may correlate with MHC heterozygosity in frogs (Cortazar-Chinarro et al., 2024). If also true for fish, this covariation sets up a clear avenue for the skin microbiome and host genetics to interactively shape parasite resistance in our system.

Here, we tested how host genetics and host-associated microbiomes affect the interaction between guppies and *G. turnbulli*. We imposed truncation selection for host parasite resistance and susceptibility, quantified as the mean number of parasites over infection. Additionally, while we did not specifically select for tissue-specific tolerance, or the ability to minimize the fitness losses

caused by a given parasite load ('tolerance' hereafter; (Stephenson & Adelman, 2022)), we

compared tolerance between our lines. Previous literature has demonstrated that tolerance is an

essential factor to consider in guppy response to *Gyrodactylus* parasites (Jog et al., 2022;

Stephenson, van Oosterhout, & Cable, 2015; Tadiri et al., 2021), and the heritability of tolerance

requires additional empirical test (Hayward et al., 2014; Råberg, 2014). Our results highlight the

complex interplay among host genetics, the microbiome, and sex-specific responses to infections.

2. Materials and methods

2.1 Does guppy resistance to Gyrodactylus turnbulli respond to selection?

2.1.1 Selection experiment

We used laboratory-bred descendants of wild guppies from a high-predation population on the

Caura River, Trinidad (UTM: 20 P 679527.7 m E, 1180376.4 m N; elevation 112 m). Wild guppies

(n ~600) were transported to Cardiff University in June 2012 (Cefas APB authorization number

CW054-D-187A) and treated prophylactically with Binox® (Nitrofurazone). The site was

previously confirmed *Gyrodactylus* spp.-free by previous surveys, and no *Gyrodactylus* spp. were

found in a subset of 80 fish examined. We collected fin clips from 66 parental-generation fish for

microsatellite genotyping. Fish were housed in 70 L mixed-sex tanks under a 12h light:12h dark

cycle at 24±1°C and fed daily with Aquarian® tropical fish flakes, supplemented with Artemia and

bloodworm. Fry were transferred to juvenile tanks soon after birth, checked for ectoparasitic

infection, and reared until sex determination (6-8 weeks). They were then kept in sex-specific tanks

for at least another month before use, ensuring all fish were sexually mature and unmated.

We infected 100 male and 100 female F1 guppies to evaluate their resistance and select founders

for our 'resistant' and 'susceptible' lines. To initiate infections, a heavily infected fish from our

parasite culture was euthanized with MS222 (0.02% tricaine methanesulfonate) and placed next to

an anesthetized experimental fish until two *G. turnbulli* parasites transmitted, as observed under a

dissecting scope. Experimental fish were housed individually in 1 L tanks, and parasites were

counted every other day under anaesthetic. After nine days, all fish were treated with Levamisole

(0.002g per L) and screened under anesthetic several times to confirm parasite clearance. Fin clips

were collected for genotyping. The 30 males and 30 females with the highest (lowest) mean

number of worms founded the 'susceptible' ('resistant') line, and randomly selected uninfected F1

fish (30 of each sex) founded the control line. Each line was allowed to breed freely in separate

120 L tanks. Tanks were monitored for fry, which were moved to F2 rearing tanks. The F2

generation was split into two subpopulations, maintaining 60 fish per tank. New tanks were set up

for each subsequent generation. Population size was kept consistent across the six subpopulations,

with new tanks set up for each subsequent generation as above.

To evaluate response to selection, 88 guppies 3-7 generations post selection were experimentally

infected with *G. turnbulli* in five batches between August and November 2017, following the same

procedure as for the F1 generation, described above. 14 guppies from each line were 'sham'

infected and were used in microbiome analysis.

2.1.2 Data analysis

We used R v.4.3.2 (R Core Team, 2020) and provide our code, output, and validation in the

Supporting Information (SI). We used ggplot2 (Wickham, 2016) to plot all the figures. For all

models, we assessed collinearity among predictor variables using the pairs function (Zuur et al.,

2009), validated model fits using the DHARMa package (Hartig, 2022), and used Type II Wald

chi-square tests to test for significance using the *Anova()* function from the 'car' package (Fox &

Weisberg, 2019).

We quantified infection resistance/susceptibility as the area under the curve of parasite number

over time, or "infection severity." To test response to selection, we used infection severity as the

response variable in a generalized linear mixed model (GLMM) in the glmmTMB package, v.

1.1.2.3 (Brooks et al., 2017) (Gamma error family and logit link function). As fixed effects, we

included selection line (resistant n=30, control n=28, susceptible n=29), fish sex, length (pre-

infection, using residuals from a regression on sex to control for sex differences), body condition

(scaled mass index; (Peig & Green, 2009)), generation, number of parasites establishing the

infection (dose), and trial date (Julian date). Length and body condition were calculated including

sham infected fish (resistant n=14, control n=14, susceptible n=14). We also included the

interaction between sex and line.

We quantified tolerance as the host's relative per-parasite percentage mass change during infection,

calculated by dividing the percentage mass change from day 0 to day 20 by the maximum number

of worms on the fish, relative to the highest observed percentage change. To assess the impact of

selection on tolerance, we used tolerance as a response variable in a GLMM, including the fixed

effects of line, sex, length, dose, and Julian date. With interaction between line and sex. We

excluded scaled mass index due to its overlap with mass change. What about interaction?

We additionally tested for a tradeoff between resistance and tolerance using a quadratic polynomial

regression model to account for non-linear effects. We used tolerance as the response variable and

log-transformed infection severity (our resistance metric), length, and generation as fixed effects.

We did not include infection severity due to high collinearity with other fixed effects.

2.1.4 Calculating heritability

We calculated the narrow-sense heritability of resistance and tolerance following the breeder's

equation (Lush, 1937). As our resistance metric, we used the average number of parasites between

1-9 days of infection, which was the metric we originally used in our truncation selection. We used

the same tolerance metric described above. Both metrics were centered to a mean of zero to

account for the differences in worm growth between the infections of the F1 and subsequent

generations.

P_b: Average starting phenotype (mean phenotype of the population before selection)

P_s: Average selected phenotype (mean phenotype of the selected parents)

P_r: Average response phenotype (mean phenotype of the offspring generation)

h^2: Narrow-sense heritability

$h^2 = P_r - P_b P_s - P_b$

2.1.5 Do differences in inbreeding contribute to line differences?

We evaluated the genetic diversity and variance in inbreeding within the wild-caught populations,

F1, and F3-F6 generations, to test if the observed difference between the resistant and susceptible

lines was due to differences in inbreeding (Smallbone et al., 2016). We extracted DNA from fin

clips with the HotSHOT method: using 50 µl of alkaline lysis reagent and neutralizing solution

followed by incubation at 95 °C for 30 minutes (Truett et al., 2000). We used two multiplex assays

to amplify eight previously designed microsatellite markers (Becher et al., 2002; Nater et al., 2008;

Paterson et al., 2005; van Oosterhout et al., 2006; Watanabe et al., 2005) at the Genetic Diversity

Centre (GDC), ETH Zurich. Samples were genotyped on an ABI Genetic Analyzer 3730 (Applied

Biosystems, USA), scored using GeneMarker (Version 2.6.4; Softgenetics), and binned using the

package MSatAllele (Alberto, 2013) in R ((R Core Team, 2020); see SI).

We tested allelic richness (A_R), gene diversity (H_S), F_{IS}, and identity disequilibrium (g2).

Allelic richness is the first population genetic metric to reflect bottleneck effects, which decrease

the effective population size and facilitate reductions in heterozygosity due to inbreeding (Nei et

al., 1975). Significant differences in allelic richness among populations are indicative of variation

in the probability that inbreeding may occur. Gene diversity measures the mean heterozygosity per

individual per population. Under inbreeding conditions homozygosity increases but gene diversity

is not immediately lost (Charlesworth, 2003). Positive values in F_{IS} can be generated by inbreeding

but may also be artificially elevated by large allele drop out or null alleles (De Meeûs, 2018; Nei

et al., 1975). We therefore also estimated identity disequilibrium (g_2), the correlation of

heterozygosity across loci resulting from inbreeding, which is robust in the presence of null alleles

(David et al., 2007). Patterns across several of these population genetic metrics consistent with

inbreeding would indicate that phenotypic differences between lines are attributable to inbreeding.

Allelic richness (A_R) (rarefied to the lowest population sample size), gene diversity (H_S) and F_{IS},

were calculated in FSTAT 2.39 (Goudet, 01 2001). Significance levels for F_{IS} were assessed in

GENEPOP (web version 4.2;(Raymond & Rousset, 1995) with 5000 iterations followed by a

Bonferroni correction for multiple comparisons. Tests for statistically significant differences

allelic richness (A_R), gene diversity (H_S) and F_{IS} among groups were calculated in FSTAT with

2000 permutations. Identity disequilibrium (g^2) is the difference between joint identity by descent

and the product of the separate probabilities of identity by decent for two loci. g^2 is estimated with

microsatellite markers as the excess of double heterozygous genotypes at all pairs of loci relative

to a random distribution of heterozygous genotypes among individuals at each locus (Cockerham

& Weir, 1973; David et al., 2007). g_2 values significantly greater than 0 are indicative of identity

disequilibrium (ID). ID was calculated in inbreedR (Stoffel et al., 2016).

2.2 Does the host-associated microbiome contribute to guppy resistance to Gyrodactylus

infection?

2.2.1 Microbial inventory

Before the F3-F7 infections began, we collected samples from 51 fish split between the infection

and sham-infection treatment groups: control (infected n=10, sham n=5), Resistant (infected n=12,

sham n=6), and Susceptible (infected n=9, sham n=8). We used the methods previously published

to collect and inventory the microbial communities on the fish skin.

Table 1: Replication Statement-Counts of Artificial selection lines in sham, infection, and microbiome

experiment

Scale of inference	Scale at which the factor of interest is applied	Number of replicates at the appropriate scale
Uninfected fish	Three selection lines (Resistant, Susceptible, Control)	Resistant: 14, Control: 14, Susceptible: 14
Infected fish	Three selection lines (Resistant, Susceptible, Control)	Resistant: 30, Control: 28, Susceptible: 29
Microbial community types	Skin swabs across lines	Control: 19, Resistant: 15, Susceptible: 16

2.2.2 Microbial alpha and beta diversity

Raw sequence data were processed using QIIME2 version 2019.7 (Bolyen et al., 2019). We

rarefied the resulting ASV table to 3,654 sequences per sample which resulted in excluding one

sample and kit controls, and used the DADA2 pipeline to calculate the unweighted and weighted

UniFrac distances (Lozupone & Knight, 2005) pairwise between all samples to create distance

matrices. We used a PERMANOVA from the adonis package in R (Anderson, 2001) to test for

significant differences in microbial community across fish line, sex, length, and body condition.

We used QIIME2 to calculate the number observed number of ASVs, Shannon's diversity index,

and Faith's phylogenetic diversity, and compare the mean values of these metrics across line and

sex using Pairwise Kruskal-Wallis tests.

2.2.3 Bacterial community typing

We used Dirichlet multinomial mixture models (DMM) (Holmes et al., 2012) in the Dirichlet

Multinomial v1.16.0 package (Morgan, 2023) to assign all samples (n=50) to three different

bacterial community types based on the relative frequency of taxa. We compared the Laplace

approximation of the negative log models to identify the number of Dirichlet components that

resulted in the best model fit (see SI).

2.2.4 Does bacterial community type predict the severity of Gyrodactylus *infection?*

To test if the bacterial community type significantly predicted infection severity, we used bacterial

community type as a fixed effect in the GLMM model used for the selection line resistance analysis

described in section 2.1.2, with the data subset to just swabbed fish.

 2.3 What is the relative importance of host genetics and the microbiome to guppy resistance to

Gyrodactylus infection?

To compare how well our GLMM models with a) bacterial community type only, b) artificial

selection line only, and c) both factors explained variation in infection severity, we used the

anova() function. All three models shared the same fixed effects, differing only in whether line or

bacterial community type or both were included as a fixed effect, and were run on the reduced

dataset using only the infected and swabbed fish (n=31).

3. Results

3.1 Guppy resistance to, and tolerance of, Gyrodactylus infection responded to selection

3.1.1 Resistance responded to selection

Artificial selection line was a significant predictor of infection severity ($\chi^2(2)$ =37.74, p=6.4e-09),

but the effect differed between sexes (Figure 1; line:sex interaction: $\chi^2(2)$ =13.51, p=0.0012).

Females had significantly more severe infections than males ($\chi^2(1)$ =24.29, p=8.3e-07) and high

starting worm 'dose' resulted in higher infection severity ($\chi^2(1)$ =5.11, p=0.024). Post hoc tests to

evaluate whether selection line significantly explained infection severity among each sex found

that artificial selection line was still significant in both sexes (female; $\chi^2(2)$ =43.69, p=5.2e-06 &

male; $\chi^2(2)$ =16.10, p=0.00032). Post hoc Tukey tests also revealed that the female guppies from

the susceptible line experienced significantly higher (Cohen's d=3.49, p<0.001), and from the

resistant line experienced significantly lower infection severity (Cohen's d=2.74, p=0.016) than

the control. Male guppies from the susceptible line experienced significantly higher infection

severity than control line males (Cohen's d=4.19, p<0.001), but resistant line males did not differ

significantly from control line males (Cohen's d=1.86, p=0.15). The average number of parasites

throughout infection was highly heritable (susceptible line h^2=0.79; resistant line h^2=0.41).

3.1.2 Tolerance responded to selection and trades off with resistance

The artificial selection lines significantly differed in tolerance ($\chi^2(2)$=14.56, p=0.00069) again

depending on host sex (Figure 2; $\chi^2(2)$=10.28, p=0.00586). Larger fish were more tolerant

($\chi^2(1)$=5.94, p=0.014). Post hoc testing found artificial selection line was only significant in

females ($\chi^2(2)$=23.81, p=6.7e-06), and smaller females were less tolerant ($\chi^2(2)$ =6.96, p=0.0083).

Post hoc Tukey tests revealed that resistant line females were significantly less tolerant than control

and susceptible line females (control; Cohen's d=7.04, p=0.0003; susceptible; Cohen's d=8.54,

p<0.00016). No fixed effect explained variation in tolerance among males. Bacterial community

types did not predict tolerance. Tolerance was heritable (susceptible line: h^2=0.52; resistant line:

h^2=0.24).

The polynomial regression model revealed a tradeoff between tolerance and resistance: higher

tolerance was correlated with lower resistance (Figure S1; F= 62.64, p<2e-16). No other fixed

effects explained significant variation in tolerance.

3.1.3 Selection lines did not differ in the extent of inbreeding

We saw a significant reduction in allelic richness among all experimental infection groups relative

to the wild-caught founders, likely reflecting a post-capture reduction in population size (Table

S1). However, we found no significant difference in AR (p= 0.589), Hs (p=0.73), FIS (p=0.42), or

g2 (Table S2) among the three lines.

3.2 The host-associated microbiome and host genetics independently underly guppy resistance to Gyrodactylus infection

DMM modeling found that the lowest Laplace approximation model fit was three bacterial clusters (*Figure S2*). All samples had a posterior probability of 98% or higher. Each bacterial community type contained 10.0–66.8% of all samples (n=5-33), and no samples were undefined. The three bacterial community types (A, B, and C) in our fish skin samples were distinct (PERMANOVA: F = 7.37, p = 0.0003, Figures 3a & S3-4). Males and females were balanced among the bacterial community types A and B, and bacterial community types clustered evenly across the artificial selection lines (Figure 3a; Fisher's exact test, p=0.8). Additionally, we found no interaction between line and bacterial community types in our model ($\chi2(4)$=3.07, p=0.546). All skin swab samples were assigned to a community type with the greatest posterior probability. For top contributing bacterial genera, see *Figure S3*.

The artificial selection lines did not significantly differ in any skin microbial diversity metrics tested. The microbiomes associated with females had significantly higher alpha diversity than those associated with males (Kruskal–Wallis pairwise test, Shannon's index: H=4.06, q=0.04; observed ASVs: H=4.88, q=0.03). Consistent with Kramp et al. (2022), the microbiomes associated with the two sexes also had significantly different beta diversity (unweighted unifrac PERMANOVA; F=1.56, p=0.006).

The skin bacterial community type A had significantly reduced infection severity compared to bacterial community type B & C ($\chi^2(2)$ = 27.54, p=1e-6, Figure 3b). A post-hoc test showed that a

model including both artificial selection line and bacterial community type is significantly better

at explaining variation in infection severity than models with either variable alone (Figure 4;

$\chi^2(14)=15.00$, p=0.00055).

4. Discussion

Our selection lines demonstrated striking heritability of parasite resistance and tolerance. Bacterial

community type predicted infection severity independently from selection line. We also observed

notable differences between males and females in skin microbiome, and in response to selection.

We discuss each of these results in turn.

We found that guppy resistance and susceptibility to *G. turnbulli* responded readily to a single

generation of artificial selection. We found no notable differences in the three selection lines in our

genetic metrics, indicating that the differences were not due to differences in inbreeding among

the lines. Our narrow-sense heritability estimates for resistance and tolerance were high,

particularly for our susceptible lines. However, our values are still comparable to heritability

estimates in previous studies (Weiler et al. 2024) and to male guppy ornaments (Morris et al. 2020),

which are correlated with *G. turnbulli* resistance (Stephenson et al. 2020).

Our results suggest that susceptibility to parasite infection responds more readily to selection than

does resistance. This may be because response to selection could involve an increase in tolerance,

or a loss of resistance. Our experimental design favored tolerance in the susceptible line, as only

individuals surviving severe infections contributed to future generations, and indeed, susceptible

line fish trended towards higher tolerance. Tolerance and resistance likely have different genetic

bases, and hosts appear to invest in one strategy at the expense of the other (Råberg, Sim, and Read

2007). This theory is supported by the fact that the resistant line was less tolerant than control and

susceptible lines, and that we found a resistance-tolerance tradeoff at the individual level.

However, these tradeoffs do not always occur, and both tolerance and resistance traits can coexist

within an individual host or population (Hardy et al. 2024; Pagán and García-Arenal 2018).

Understanding how resistance, susceptibility, and tolerance of *Gyrodactylus* spp. infection respond

to selection is crucial for understanding how Trinidadian guppy behavioral and immunological

defenses interact with parasite spread and evolution (Walsman, Lambe, and Stephenson 2024;

Walsman et al. 2022). Our heritability estimates suggest an important genetic contribution to traits

influencing infection outcomes, which may allow natural selection to act rapidly under high

parasite pressures. In some populations, infection prevalence is very high (>90%; (Clark et al.,

2023; Stephenson, van Oosterhout, Mohammed, et al., 2015)), potentially selecting for increased

infection tolerance and decreased resistance, as 'resistance is futile' (Walsman et al. 2023). In

concert, transgenerational immune priming may reduce the expression of parasite susceptibility

under natural conditions in this system Weiler et al bred fish for several generations in the absence

of parasites to remove such effects. Future work should take into account the full complexity of

eco-evolutionary feedbacks, as well as the heritability estimates we present here (Weiler et al.,

2025).

Despite differences in their infection phenotypes, we did not find that microbiome composition,

alpha diversity, or bacterial community type differed significantly between lines, nor did the

bacterial communities predict tolerance. Nevertheless, we found a strong correlation between

bacterial community and infection severity, pointing to important interactions between the skin

microbiome and parasites, supporting previous studies (Scheifler et al. 2022; Speer 2022). Our

findings therefore suggest that host genetics and host-associated skin microbiome significantly

predict parasite infection severity, but not tolerance, and that they do so independently and

additively to host genetics. This result appears contrary to previous findings that host immune

system, metabolism, and growth rate correlate with the microbiome (Zheng, Liwinski, and Elinav

2020; Speer 2022). Indeed, artificial selection for body size in guppies significantly altered the

host-associated microbiome, and the fast-growing line had greater parasite colonization resistance

(Piazzon et al. 2020), and life history and size selection experiments indirectly affect guppy

resistance to *G. turnbulli* (Bartuseviciute et al. 2022), potentially mediated through the

microbiome. This difference may be due to the fact that we used skin microbiome, and fish skin

microbiomes are highly variable and may not reflect the host's genetics but drift due to stochasticity

(Jones et al. 2022; Wang et al. 2023).

Our results add to the growing body of work demonstrating that the microbiome affects parasite

resistance in several taxa: pre-infection microbiome composition, alpha diversity, or microbiome

function is a powerful tool to predict individuals that are most susceptible to parasites within a

population (Bates et al. 2022; Lee et al. 2019). Further investigation of our bacterial community

findings revealed similarities to a recent study of skin microbial metabolism function in amphibian

chytridiomycosis (Bates et al. 2022): we found five of the eight ASVs reported to be indicator taxa

of epizootic dynamics; *Burkholderiales*, *Pseudomonadales*, *Sphingomonadales*, *Rhizobiales*,

Flavobacteriales were also key contributing taxa in our bacterial community types (see Figure S2).

In alignment with those of Dargent et al. (2016), our results indicate that parasite resistance

evolved more slowly in males than females. This may be due to a pre-existing higher level of, and

less genetic variation in, resistance in males due to sexual selection. This lack of variation in male

resistance may explain why we found evidence for a trade-off between resistance and tolerance

among females, but not among males. Indeed, males in our experiment had less severe infections,

and males are less likely, and less heavily infected in surveys of natural populations (Clark et al.,

2023; Gotanda et al., 2013; Stephenson, van Oosterhout, Mohammed, et al., 2015). Females in

many populations prefer males with larger orange ornaments (Endler et al., 2001; Houde, 1988,

1997), with implications for disease dynamics (Rovenolt et al., 2024). Larger orange ornaments

indicate resistance to this parasite (Stephenson et al., 2020). Our experimental design allowed

female preference to affect male reproductive success, potentially resulting in the overall high and

invariable resistance we observe in males.

Importantly, the apparently sex-specific evolutionary trajectory of parasite resistance we observe

highlights the role of something other than the MHC, which has been a focus in this system to date

(Fraser et al., 2010; Fraser & Neff, 2009; Phillips et al., 2018): MHC resistance is not sex based

but a mixture of both parents' MHC genes (Cruz-Tapias et al., 2013). Our results suggest that the

microbiome may contribute somewhat to the sex-specificity we observe: the sexes differed in both

alpha and beta diversity metrics of bacterial communities, similar to the skin microbiome in other

species (Rodríguez-Barreto et al. 2023). More broadly, our findings highlight potential co-

evolutionary interactions between host sexes and their associated microbiomes in relation to

parasite resistance. These possibilities will hopefully inspire exciting and novel exploration of the

evolutionary dynamics of resistance and tolerance between the sexes.

References

Alberto, F. (2013). MsatAllele: Visualizes the scoring and binning of microsatellite fragment sizes.

R Package Version.

Amenyogbe, N., Kollmann, T. R., & Ben-Othman, R. (2017). Early-Life Host-Microbiome

Interphase: The Key Frontier for Immune Development. *Frontiers in Pediatrics*, *5*, 111.

Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance.

Austral Ecology, *26*(1), 32–46.

Bartuseviciute, V., Diaz Pauli, B., Salvanes, A. G. V., & Heino, M. (2022). Size-selective

harvesting affects the immunocompetence of guppies exposed to the parasite Gyrodactylus.

Proceedings. Biological Sciences / The Royal Society, *289*(1981), 20220534.

Bates, K. A., Sommer, U., Hopkins, K. P., Shelton, J. M. G., Wierzbicki, C., Sergeant, C., Tapley,

B., Michaels, C. J., Schmeller, D. S., Loyau, A., Bosch, J., Viant, M. R., Harrison, X. A.,

Garner, T. W. J., & Fisher, M. C. (2022). Microbiome function predicts amphibian

chytridiomycosis disease dynamics. *Microbiome*, *10*(1), 44.

Becher, S. A., Russell, S. T., & Magurran, A. E. (2002). Isolation and characterization of

polymorphic microsatellites in the Trinidadian guppy (Poecilia reticulata). *Molecular*

Ecology Notes, *2*(4), 456–458.

Becker, M. H., Brucker, R. M., Schwantes, C. R., Harris, R. N., & Minbiole, K. P. C. (2009). The

bacterially produced metabolite violacein is associated with survival of amphibians

infected with a lethal fungus. *Applied and Environmental Microbiology*, *75*(21), 6635–

6638.

Bernardo-Cravo, A. P., Schmeller, D. S., Chatzinotas, A., Vredenburg, V. T., & Loyau, A. (2020).

Environmental Factors and Host Microbiomes Shape Host-Pathogen Dynamics. *Trends in*

Parasitology, *36*(7), 616–633.

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A.,

Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K.,

Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M.,

Chase, J., … Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible

microbiome data science using QIIME 2. *Nature Biotechnology*, *37*(8), 852–857.

Bordenstein, S. R., & Theis, K. R. (2015). Host Biology in Light of the Microbiome: Ten Principles

of Holobionts and Hologenomes. *PLoS Biology*, *13*(8), e1002226.

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A.,

Skaug, H. J., Maechler, M., & Bolker, B. M. (2017). glmmTMB Balances Speed and

Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. In *The*

R Journal (Vol. 9, Issue 2, pp. 378–400). https://doi.org/10.32614/RJ-2017-066

Cable, J., & Van Oosterhout, C. (2007). The role of innate and acquired resistance in two natural

populations of guppies (Poecilia reticulata) infected with the ectoparasite Gyrodactylus

turnbulli. *Biological Journal of the Linnean Society. Linnean Society of London*, *90*(4),

647–655.

Charlesworth, D. (2003). Effects of inbreeding on the genetic diversity of populations.

Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences,

358(1434), 1051–1070.

Chen, X., Wu, R., Li, L., Zeng, Y., Chen, J., Wei, M., Feng, Y., Chen, G., Wang, Y., Lin, L., Luo,

H., Chen, A., Zeng, Z., He, F., Bai, Y., Zhang, S., Han, Y., Wang, Z., Zhao, X., … Gong, S.

(2023). Pregnancy-induced changes to the gut microbiota drive macrophage pyroptosis and

exacerbate septic inflammation. *Immunity*, *56*(2), 336-352.e9.

Clark, D. R., Young, K. A., Kitzes, J., Moore, P. J., Evans, A. J., & Stephenson, J. F. (2023).

Aggregation of symbionts on hosts depends on interaction type and host traits. *Ecography*.

https://doi.org/10.1111/ecog.06858

Cockerham, C. C., & Weir, B. S. (1973). Descent measures for two loci with some applications.

Theoretical Population Biology, *4*(3), 300–330.

Cortazar-Chinarro, M., Richter-Boix, A., Rödin-Mörch, P., Halvarsson, P., Logue, J. B., Laurila,

A., & Höglund, J. (2024). Association between the skin microbiome and MHC class II

diversity in an amphibian. *Molecular Ecology*, *33*(1), e17198.

Cruz-Tapias, P., Castiblanco, J., & Anaya, J.-M. (2013). Major histocompatibility complex:

Antigen processing and presentation. In *Autoimmunity: From Bench to Bedside [Internet]*.

El Rosario University Press.

Dargent, F., Rolshausen, G., Hendry, A. P., Scott, M. E., & Fussmann, G. F. (2016). Parting ways:

parasite release in nature leads to sex-specific evolution of defence. *Journal of*

Evolutionary Biology, *29*(1), 23–34.

David, P., Pujol, B., Viard, F., Castella, V., & Goudet, J. (2007). Reliable selfing rate estimates

from imperfect population genetic data. *Molecular Ecology*, *16*(12), 2474–2487.

De Meeûs, T. (2018). Revisiting FIS, FST, Wahlund Effects, and Null Alleles. *The Journal of*

Heredity, *109*(4), 446–456.

Debray, R., Herbert, R. A., Jaffe, A. L., Crits-Christoph, A., Power, M. E., & Koskella, B. (2022).

Priority effects in microbiome assembly. *Nature Reviews. Microbiology*, *20*(2), 109–121.

Endler, J. A., Basolo, A., Glowacki, S., & Zerr, J. (2001). Variation in response to artificial selection for light sensitivity in guppies (Poecilia reticulata). *The American Naturalist*, *158*(1), 36–48.

Fox, J., & Weisberg, S. (2019). Companion to Applied Regression [R package car version 3.1-2]. *Sage*, *3rd*. https://cran.r-project.org/web/packages/car/index.html

Fraser, B. A., & Neff, B. D. (2009). MHC class IIB additive and non-additive effects on fitness measures in the guppy *Poecilia reticulata*. *Journal of Fish Biology*, *75*(9), 2299–2312.

Fraser, B. A., Ramnarine, I. W., & Neff, B. D. (2010). TEMPORAL VARIATION AT THE MHC CLASS IIB IN WILD POPULATIONS OF THE GUPPY (*POECILIA RETICULATA*). *Evolution; International Journal of Organic Evolution*, *64*(7), 2086–2096.

Gorczyca, K., Obuchowska, A., Kimber-Trojnar, Ż., Wierzchowska-Opoka, M., & Leszczyńska-Gorzelak, B. (2022). Changes in the Gut Microbiome and Pathologies in Pregnancy. *International Journal of Environmental Research and Public Health*, *19*(16). https://doi.org/10.3390/ijerph19169961

Gotanda, K. M., Delaire, L. C., Raeymaekers, J. A. M., Pérez-Jvostov, F., Dargent, F., Bentzen, P., Scott, M. E., Fussmann, G. F., & Hendry, A. P. (2013). Adding parasites to the guppy-predation story: insights from field surveys. *Oecologia*, *172*(1), 155–166.

Goudet, J. (01 2001). FSTAT, Version. *1. 2: A Computer Program to Calculate F-Statistics. Heredity*, *86*, 485–486.

Hardy, B. M., Muths, E., Funk, W. C., & Bailey, L. L. (2024). Quantifying intraspecific variation in host resistance and tolerance to a lethal pathogen. *The Journal of Animal Ecology*. https://doi.org/10.1111/1365-2656.14106

Hartig, F. (2022). *DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression*
models. R package version 0.4. 6.

Hayward, A. D., Nussey, D. H., Wilson, A. J., Berenos, C., Pilkington, J. G., Watt, K. A.,
Pemberton, J. M., & Graham, A. L. (2014). Natural selection on individual variation in
tolerance of gastrointestinal nematode infection. *PLoS Biology*, *12*(7), e1001917.

Holmes, I., Harris, K., & Quince, C. (2012). Dirichlet multinomial mixtures: generative models
for microbial metagenomics. *PloS One*, *7*(2), e30126.

Houde, A. E. (1988). Genetic difference in female choice between two guppy populations. *Animal*
Behaviour, *36*(2), 510–516.

Houde, A. E. (1997). Effect of artificial selection on male colour patterns on mating preference of
female guppies. *Proceedings of the Royal Society of London. Series B: Biological Sciences*,
256(1346), 125–130.

Jog, M. G., Sackett, M. E., Kisty, S. D., Hansen, J. A., & Stephenson, J. F. (2022). The behaviour
of infected guppies depends on social context, parasite tolerance and host sex. *Animal*
Behaviour, *187*, 97–104.

Jones, E. W., Carlson, J. M., Sivak, D. A., & Ludington, W. B. (2022). Stochastic microbiome
assembly depends on context. *Proceedings of the National Academy of Sciences of the*
United States of America, *119*(7), e2115877119.

Labro, M. T. (2022). *Host defense and infection*. CRC Press.

Lee, K. H., Gordon, A., Shedden, K., Kuan, G., Ng, S., Balmaseda, A., & Foxman, B. (2019). The
respiratory microbiome and susceptibility to influenza virus infection. *PloS One*, *14*(1),
e0207898.

Lozupone, C., & Knight, R. (2005). UniFrac: a new phylogenetic method for comparing microbial

communities. *Applied and Environmental Microbiology*, *71*(12), 8228–8235.

Lush, J. L. (1937). *Animal Breeding Plans - Scholar's Choice Edition*. Bibliolife DBA of Bibilio

Bazaar II LLC.

Markle, J. G., & Fish, E. N. (2014). SeXX matters in immunity. *Trends in Immunology*, *35*(3), 97–

104.

Mohammed, R. S., Reynolds, M., James, J., Williams, C., Mohammed, A., Ramsubhag, A., van

Oosterhout, C., & Cable, J. (2016). Getting into hot water: sick guppies frequent warmer

thermal conditions. *Oecologia*, *181*(3), 911–917.

Morgan, M. (2023). *DirichletMultinomial: Dirichlet-Multinomial Mixture Model Machine

Learning for Microbiome Data*. https://doi.org/10.18129/B9.bioc.DirichletMultinomial

Morris, J., Darolti, I., van der Bijl, W., & Mank, J. E. (2020). High-resolution characterization of

male ornamentation and re-evaluation of sex linkage in guppies. *Proceedings. Biological

Sciences / The Royal Society*, *287*(1937), 20201677.

Morrow, E. H. (2015). The evolution of sex differences in disease. *Biology of Sex Differences*,

6(1), 5.

Nater, A., Krützen, M., & Lindholm, A. K. (2008). Development of polymorphic microsatellite

markers for the livebearing fish Poecilia parae. *Molecular Ecology Resources*, *8*(4), 857–

860.

Nei, M., Maruyama, T., & Chakraborty, R. (1975). The Bottleneck Effect and Genetic Variability

in Populations. *Evolution; International Journal of Organic Evolution*, *29*(1), 1–10.

Nelson, P., & May, G. (2020). Defensive Symbiosis and the Evolution of Virulence. *The American

Naturalist*, *196*(3), 333–343.

Ni, J., Wu, G. D., Albenberg, L., & Tomov, V. T. (2017). Gut microbiota and IBD: causation or

correlation? *Nature Reviews. Gastroenterology & Hepatology*, *14*(10), 573–584.

Pagán, I., & García-Arenal, F. (2018). Tolerance to plant pathogens: Theory and experimental

evidence. *International Journal of Molecular Sciences*, *19*(3).

https://doi.org/10.3390/ijms19030810

Paterson, I. G., Crispo, E., Kinnison, M. T., Hendry, A. P., & Bentzen, P. (2005). Characterization

of tetranucleotide microsatellite markers in guppy (Poecilia reticulata). *Molecular Ecology*

Notes, *5*(2), 269–271.

Peig, J., & Green, A. J. (2009). New perspectives for estimating body condition from mass/length

data: the scaled mass index as an alternative method. *Oikos* , *118*(12), 1883–1891.

Phillips, K. P., Cable, J., Mohammed, R. S., Chmielewski, S., Przesmycka, K. J., van Oosterhout,

C., & Radwan, J. (2021). Functional immunogenetic variation, rather than local adaptation,

predicts ectoparasite infection intensity in a model fish species. *Molecular Ecology*, *30*(21),

5588–5604.

Phillips, K. P., Cable, J., Mohammed, R. S., Herdegen-Radwan, M., Raubic, J., Przesmycka, K. J.,

van Oosterhout, C., & Radwan, J. (2018). Immunogenetic novelty confers a selective

advantage in host-pathogen coevolution. *Proceedings of the National Academy of Sciences*

of the United States of America, *115*(7), 1552–1557.

Piazzon, M. C., Naya-Català, F., Perera, E., Palenzuela, O., Sitjà-Bobadilla, A., & Pérez-Sánchez,

J. (2020). Genetic selection for growth drives differences in intestinal microbiota

composition and parasite disease resistance in gilthead sea bream. *Microbiome*, *8*(1), 168.

R Core Team. (2020). *R: A language and environment for statistical computing* (Version v.4.1.0)

[R Foundation for Statistical Computing, Vienna, Austria.].

Råberg, L. (2014). How to live with the enemy: understanding tolerance to parasites. *PLoS*

Biology, *12*(11), e1001989.

Råberg, L., Sim, D., & Read, A. F. (2007). Disentangling genetic variation for resistance and

tolerance to infectious diseases in animals. *Science*, *318*(5851), 812–814.

Raymond, M., & Rousset, F. (1995). GENEPOP (Version 1.2): Population Genetics Software for

Exact Tests and Ecumenicism. *The Journal of Heredity*, *86*(3), 248–249.

Rodríguez-Barreto, D., Sanz-González, J. C., Martín, M. V., Arrieta, J. M., & Almansa, E. (2023).

Sex-specific bacterial microbiome variation in octopus vulgaris skin. *Frontiers in*

Microbiology, *14*, 1233661.

Rovenolt, F., Walsman, J., Martin, C., Gotanda, K., & Stephenson, J. (2024). Parasite prevalence

depends on female preference: Integrating parasite-mediated sexual selection and

infectious disease dynamics. In *Authorea Inc.*

https://doi.org/10.22541/au.172447480.06236699/v1

Santos-Marcos, J. A., Mora-Ortiz, M., Tena-Sempere, M., Lopez-Miranda, J., & Camargo, A.

(2023). Interaction between gut microbiota and sex hormones and their relation to sexual

dimorphism in metabolic diseases. *Biology of Sex Differences*, *14*(1), 4.

Scheifler, M., Sanchez-Brosseau, S., Magnanou, E., & Desdevises, Y. (2022). Diversity and

structure of sparids external microbiota (Teleostei) and its link with monogenean

ectoparasites. *Animal Microbiome*, *4*(1), 27.

Smallbone, W., Van Oosterhout, C., & Cable, J. (2016). The effects of inbreeding on disease

susceptibility: *Gyrodactylus turnbulli* infection of guppies, *Poecilia reticulata*.

Experimental Parasitology, *167*, 32–37.

Speer, K. A. (2022). [Review of *Microbiomes mediate host-parasite interactions*]. *Molecular*

Ecology, *31*(7), 1925–1927.

Stephenson, J. F., & Adelman, J. S. (2022). The behavior of infected hosts: Behavioral tolerance,

behavioral resilience, and their implications for behavioral competence. *Animal Behavior*

and   *Parasitism*.

https://books.google.ca/books?hl=en&lr=&id=xSZ_EAAAQBAJ&oi=fnd&pg=PA287&o

ts=xW36pEEkt3&sig=asg6-91ifwE5LAwCypVYASxqwO4

Stephenson, J. F., Stevens, M., Troscianko, J., & Jokela, J. (2020a). The size, symmetry, and color

saturation of a male guppy's ornaments forecast his resistance to parasites. *The American*

Naturalist, *196*(5), 597–608.

Stephenson, J. F., Stevens, M., Troscianko, J., & Jokela, J. (2020b). The size, symmetry, and color

saturation of a male guppy's ornaments forecast his resistance to parasites. *The American*

Naturalist, *196*(5), 597–608.

Stephenson, J. F., van Oosterhout, C., & Cable, J. (2015). Pace of life, predators and parasites:

predator-induced life-history evolution in Trinidadian guppies predicts decrease in parasite

tolerance. *Biology Letters*, *11*(11), 20150806.

Stephenson, J. F., van Oosterhout, C., Mohammed, R. S., & Cable, J. (2015). Parasites of

Trinidadian guppies: evidence for sex- and age-specific trait-mediated indirect effects of

predators. *Ecology*, *96*(2), 489–498.

Stoffel, M. A., Esser, M., Kardos, M., Humble, E., Nichols, H., David, P., & Hoffman, J. I. (2016).

inbreedR: an R package for the analysis of inbreeding based on genetic markers. *Methods*

in Ecology and Evolution, *7*(11), 1331–1339.

Tadiri, C. P., Fussmann, G. F., & Scott, M. E. (2021). Parasite spread in experimental

metapopulations: resistance, tolerance and host competence. *Oikos (Copenhagen,*

Denmark), *130*(9), 1562–1571.

Trevelline, B. K., Stephenson, J. F., & Kohl, K. D. (2020). [Review of *Two's company, three's a*

crowd: Exploring how host-parasite-microbiota interactions may influence disease

susceptibility and conservation of wildlife]. *Molecular Ecology*, *29*(8), 1402–1405.

Van Oosterhout, C., Harris, P. D., & Cable, J. (2003). Marked variation in parasite resistance

between two wild populations of the Trinidadian guppy, *Poecilia reticulata* (Pisces:

Poeciliidae). *Biological Journal of the Linnean Society. Linnean Society of London*, *79*(4),

645–651.

van Oosterhout, C., Joyce, D. A., Cummings, S. M., Blais, J., Barson, N. J., Ramnarine, I. W.,

Mohammed, R. S., Persad, N., & Cable, J. (2006). Balancing selection, random genetic

drift, and genetic variation at the major histocompatibility complex in two wild populations

of guppies (*Poecilia reticulata*). *Evolution; International Journal of Organic Evolution*,

60(12), 2562–2574.

Walsman, J. C., Duffy, M. A., Cáceres, C. E., & Hall, S. R. (2023). "Resistance Is Futile": Weaker

Selection for Resistance by Abundant Parasites Increases Prevalence and Depresses Host

Density. *The American Naturalist*, *201*(6), 864–879.

Walsman, J. C., Janecka, M. J., Clark, D. R., Kramp, R. D., Rovenolt, F., Patrick, R., Mohammed,

R. S., Konczal, M., Cressler, C. E., & Stephenson, J. F. (2022). Shoaling guppies evade

predation but have deadlier parasites. *Nature Ecology & Evolution*, *6*(7), 945–954.

Walsman, J. C., Lambe, M., & Stephenson, J. F. (2024). Associating with kin selects for disease

resistance and against tolerance. *Proceedings. Biological Sciences*, *291*(2023), 20240356.

Wang, L.-C., Chen, L.-H., Chiu, Y.-C., Liou, C.-Y., Chen, H.-C., Lu, C.-Y., & Chen, J.-L. (2023).

Teleost skin microbiome: An intimate interplay between the environment and the host

immunity. *Fish & Shellfish Immunology*, *139*, 108869.

Watanabe, T., Yoshida, M., Nakajima, M., & Taniguchi, N. (2005). Linkage mapping of AFLP and

microsatellite DNA markers with the body color- and sex-determining loci in the guppy

(Poecilia reticulata). *Zoological Science*, *22*(8), 883–889.

Weiler, I. L. G., Kramp, R. D., Rovenolt, F., & Stephenson, J. F. (2024). Sex-dependent effects of

infection on guppy reproductive fitness and offspring parasite resistance. In *bioRxiv* (p.

2023.10.29.564622). https://doi.org/10.1101/2023.10.29.564622

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer.

Wilde, J., Slack, E., & Foster, K. R. (2024). Host control of the microbiome: Mechanisms,

evolution, and disease. *Science (New York, N.Y.)*, *385*(6706), eadi3338.

Zheng, D., Liwinski, T., & Elinav, E. (2020). Interaction between microbiota and immunity in

health and disease. *Cell Research*, *30*(6), 492–506.

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed Effects Models*

and Extensions in Ecology with R. Springer New York.

Figures

Figure 1. *Artificial selection lines significantly predicted G. turnbulli infection severity in*

Trinidadian guppies. Infection severity (worm days) reflects the area under the infection load

curve. Female guppies (n=44) in susceptible and resistant lines differed from the control line.

Male guppies (n=43) only showed differences in the susceptible line. Box plots illustrate median,

interquartile range, and values within 1.5× the interquartile range of infection severity.

667   *Figure 2. Artificial selection lines also predicted relative tissue-specific tolerance in*

668   *Trinidadian Guppies*. Tolerance measured as per-parasite mass change across infection

669   minus max tolerance. Female guppies (n=44) in resistant lines significantly differed from control

670   and susceptible lines. Male guppies (n=43) did not show significant differences. Box plots

671   illustrate the same metrics as in Figure 1.

Figure 3. The distribution of skin bacterial community types across guppy selection lines

showed no significant association. A. mosaic plot presents bacterial community types (A, B,

C) across control (c), resistant (r), and susceptible (s) lines, with no significant distribution

differences (Fisher's exact test, p=0.8). B. PCoA analysis (n=52) indicated bacterial community

type was significant in variation (PERMANOVA: F=7.37, P=0.0003), but no significant clustering

was detected for selection lines (PERMANOVA: F=0.68, P=0.89).

 Figure 4. Selection lines and bacterial community types predicted G. turnbulli infection

689 *severity.* A. Lines significantly predicted severity in swabbed fish (n=31; ANOVA: χ^2=9.39,

690 P=0.009). B. Community type A had lower infection severity than types B and C (A-B, z=2.59,

691 P=0.0034; A-C, z=4.92, P<0.001). No interactions between community type and selection line

692 were detected (ANOVA: χ^2=7.4, P=0.116). A combined model of line and bacterial types

693 explained infection severity better than either factor alone (χ^2=13.40, P=0.001).

694