Membrane-Dependent Dynamics and Dual Translocation Mechanisms of ABCB4: Insights from Molecular Dynamics Simulations

Veronica Crespi,[1] Ágota Tóth,[1,2] Angelika Janaszkiewicz,[1,3] Thomas Falguières,[4] and Florent Di Meo[1,5*]

[1] *Inserm U1248 Pharmacology & Transplantation, Univ. Limoges, Limoges, France*

[2] *InSilibio, Limoges, France*

[3] *School of Science, RMIT University, Melbourne, Vic, 3001, Australia*

[4] *Inserm U1193 Physiopathogenesis and Treatment of Liver Diseases - Hepatinov, Univ. Paris Saclay, France.*

[5] *Inserm US042/CNRS UAR 2015 Integrative Biology Health Chemistry & Environment, Univ. Limoges, Limoges, France*

Corresponding Author :

Dr. Florent Di Meo

Inserm U1248 Pharmacology & Transplantation, Univ. Limoges,

2 rue du Prof. Descottes,

87000 F-Limoges, France

florent.di-meo@inserm.fr

Tel: +33(0)5 19 56 42 76

Abstract

ABCB4 is an ATP-binding cassette transporter expressed at the canalicular membrane of hepatocytes and responsible for translocating phosphatidylcholine into bile. Despite the recent cryo-EM structures of ABCB4, knowledge about the molecular mechanism of phosphatidylcholine transport remains fragmented. In this study, we used all-atom molecular dynamics simulations to investigate ABCB4 dynamics during its transport cycle, leveraging both symmetric and asymmetric membrane models. Our results demonstrate that membrane composition influences the local conformational dynamics of ABCB4, revealing distinct lipid-binding patterns across different conformers, particularly for cholesterol. We explored the two potential mechanisms for phosphatidylcholine translocation: the canonical ATP-driven alternating access model and the "credit-card swipe" model. Critical residues were identified for phosphatidylcholine binding and transport pathway modulation, supporting the canonical mechanism while also indicating a possible additional pathway. The conformer-specific roles of kinking in transmembrane helices (TMH4 and TMH10) were highlighted as key events in substrate translocation. Overall, ABCB4 may utilize a cooperative transport mechanism, integrating elements of both models to facilitate efficient phosphatidylcholine motion across the membrane. This study provides new insights into the relationship between membrane environment and ABCB4 function, contributing to our understanding of its role in bile physiology and susceptibility to genetic and xenobiotic influences.

Keywords: ABC transporters – Molecular dynamics – Liver – Phosphatidylcholine translocation

1. Introduction

ABCB4 is an ATP-binding cassette (ABC) transporter exclusively expressed at the canalicular membrane of hepatocytes. It is responsible for the translocation of phosphatidylcholine lipids (PC) into the bile canaliculus; being PC one of the main components of bile's mixed micelles, along with bile salts and cholesterol[1]. Variations in *ABCB4* gene are associated with biliary diseases such as progressive familial intrahepatic cholestasis type 3 (PFIC3), cholelithiasis syndrome, intrahepatic cholestasis of pregnancy, and liver cancer[2]. Xenobiotics can inhibit ABCB4 leading to iatrogenic imbalance of bile composition which is associated with an increment in bile toxicity owing to bile salt detergent action on cell membranes [3–5].

Recently, ABCB4 structure was resolved through cryogenic electron microscopy (cryo-EM) adopting the so-called type IV fold[1,3]. ABCB4 structure assembles two consecutive similar subdomains connected by a linker [6]. Each subdomain consists of a transmembrane domain (TMD) made of six transmembrane helices (TMHs) and one so-called nucleotide binding domain (NBD). ABCB4 was resolved under different conformations, namely inward facing ([IF]ABCB4) [3] and close-cleft ([CC]ABCB4) [1]. Different bound states were resolved, namely apo, substrate- and inhibitor-bound states for [IF]ABCB4 [3], and nucleotide-bound state for [CC]ABCB4[1]. Following the commonly accepted two-side alternating access model, both the NBD dimerize upon the binding of two ATP molecules sandwiched in two nucleotide binding sites (NBS)[7]. It is still not fully understood how a PC substrate is translocated during the transport cycle. Likewise, knowledge remains fragmented regarding whether ABCB4-mediated PC lipid is directly released to canalicular micelles or if ABCB4 transport mechanism leads to outer leaflet PC enrichment prior to PC lipid extrusion in bile, even though the latter seems more likely[8].

Two complementary different molecular mechanisms have been suggested regarding PC translocation across lipid bilayer. Nosol et al. proposed the canonical ATP-driven alternating access for PC translocation relying on the resolution of substrate-bound ABCB4 adopting the IF conformation[3]. Similar to type IV ABC-driven substrate translocation[3], [IF]ABCB4 might initially bind PC lipid substrate from the inner leaflet leading to the ligand-bound conformation ([IF]ABCB4-PC). PC substrate is thus expected to enter into ABCB4 binding pocket either by the front gate composed of TMH3/TMH4/TMH6 or the rear (or back) gate formed by TMH9/10/12, the latter being suggested to be less likely[9]. The central role of Trp234 and Phe345 for PC substrate binding was proposed through site-directed mutagenesis experiments and the cryo-EM structure. Upon ATP binding, [IF]ABCB4-PC-(ATP)$_2$ state might undergo large-scale conformational changes leading to PC release out of the protein binding pocket through a proposed transient open outward-facing (OF) conformation.

Interestingly, site-directed mutagenesis of Trp234, Ala231, Phe345, or His989 was associated with a significant decrease of PC translocation ranging from ca. 20 to 50% as compared to WT-ABCB4; but not no total inhibition of ABCB4 function was observed[3]. This may suggest

the existence of a non-canonical PC translocation mechanism. In this context, a joint computational and experimental study proposed an alternative ABCB4-driven PC translocation adopting the so-called credit-card (c/c) swipe mechanism[9]. Such a mechanism relies on the exposure of TMH1 polar amino acids shading PC polar head while maintaining lipid tails in a hydrophobic environment. The c/c swipe mechanism starts with the outward-open or ^{CC}ABCB4 conformation, in which TMH1 polar amino acids bind PC polar head. As for other type IV ABC transporters [6,10] ATP hydrolysis is required for CC-to-IF conformational transition, achieving PC translocation. Subsequently, the binding of ATP in NBS favors conformational changes from ^{IF}ABCB4-(ATP)$_2$ back to ^{CC}ABCB4-(ATP)$_2$ in presence or not of PC substrate.

Taking advantage of the available cryo-EM structures, we investigated the dynamics of the main conformational milestones by means of all-atom molecular dynamics simulations. Being ABCB4 a membrane transporter, special attention is also paid to the impact of the lipid composition onto the structural dynamics of ABCB4 accounting for the known interplay between membrane environment and ABC proteins[11–16]. Two different membrane models were employed, namely symmetric (POPC:POPE:Chol (2:1:1)) and asymmetric distributions consisting in multiple lipids as proposed by Ingólfsson et al.[17]. The present study aims to provide computational perspectives on membrane-dependent ABCB4 structural dynamics as well as the complementarity between the canonical substrate translocation and the credit-card swipe mechanism.

2. Results

2.1. Membrane-dependence of ABCB4 structural dynamics

2.1.1. Overview of the ABC conformational space along transport cycle.

Sampling of the ABCB4 structural dynamics was examined accounting for the different conformations (namely IF and CC) and bound states (PC-, ATP- and PC/ATP-bound). The so-called ABC structural parameters[18,19] *i.e.*, intracellular (IC) angle, NBD distance, and NBD twist were monitored throughout the MD simulations (Figure 1A). These structural parameters were used to investigate structural variabilities of IC opening as well as NBD dimerization and twist along ABCB4 alternating access[18]. Distributions of ABC structural parameters were projected onto the so-called ABC conformational space (see Figure 1B&C). It was defined by ABC structural parameters measured from a large dataset of cryo-EM structures of type IV ABC transporters (including the ABCB4 structures) and adopting different conformations and bound states (Table S1). The three main conformations (namely IF, OF, and the so-called Unlocked-returned – UR) are neatly arranged and recognizable, even though IF structures exhibit large variability regarding IC angle and NBD distance. Ligand- and/or nucleotide-bound states are less clearly distinguishable.

Interestingly, for most of the IF ABC conformations, substrate-bound structures reported larger IC angles than apo states in line with our MD simulations (Figure 1B). Inhibitor-bound ABC IF conformations exhibited an even larger IC opening. This is the case for ^{IF}ABCB4 structures (PDB IDs: 7NIU, 7NIV, and 7NIW being respectively apo, substrate- and inhibitor-bound states, see Figure 1B). While comparing structures of ABCB4 and those obtained from MD simulations, similar behaviors were observed regardless of membrane composition. ABC structural parameters measured from ABCB4 structures exhibit only slight differences with MD simulations, in contrast to similar investigations conducted with the multidrug resistance-associated protein 1 (ABCC1/MRP1)[11]. Cryo-EM IF structures of ABCB4 exhibit slightly larger IF opening as pictured by IC angle and NBD distances as compared to MD simulations. In contrast, structures relaxed during MD simulations, leading to decrease in NBD distance and IC angle for IF conformations. NBD distance slightly increased during the MD simulations for ^{CC}ABCB4-(ATP)$_2$ while NBD twist remained within the same range as cryo-EM structure. As expected, owing to the absence of substrate or nucleotide, apo states exhibit much larger structural variability of IC opening as pictured by NBD distance and IC angle distributions, in agreement with observations made on e.g., MRP1[11] or P-gp[10].

Regardless of membrane composition, MD simulations conducted with IF conformations show that ATP binding is not sufficient to reach the so-called proper NBD dimerization described by the ^{CC}ABCB4-(ATP)$_2$ conformation (Figure 1B&C). This can be explained by the relatively short μs-scale from all-atom MD simulations as compared to the expected alternating access timescale, in line with observations made on other ABC transporters [10,11]. This also suggests that IF-to-OF transition requires slow large-scale motions driven from TMHs to trigger proper NBD dimerization. Indeed, IF conformations showed NBD distance values greater than those of the CC conformations (30.8 ± 0.4 and 30.7 ± 0.3 Å for ^{CC}ABCB4-(ATP)$_2$ respectively in symmetric and asymmetric membrane). Likewise, MD simulations revealed significantly lower variability of NBD twist for ^{CC}ABCB4-(ATP)$_2$ conformation (-145.4° ± 2.0 and -149.1° ± 3.3 in symmetric and asymmetric membrane, respectively).

Binding of PC substrate in the canonical binding pocket is associated with an increase in the IC opening as compared with apo ^{IF}ABCB4 state likely owing to the steric hindrance assigned to PC substrate size. In contrast, NBD distances for ^{IF}ABCB4-PC tend to be smaller than apo ^{IF}ABCB4 state, suggesting an allosteric communication from TMD to NBD regions upon substrate binding. In other words, TMH and NBD motions are not moving altogether as a single block but their motions are coupled, which is assigned to the pivotal role of the TMH-NBD coupling helices as described for type IV ABC transporters [6,7]. Therefore, PC binding to ^{IF}ABCB4 conformation leads to IC opening, which in turn might be associated with the triggering of NBD dimerization.

2.1.2. Impact of the lipid composition onto the overall structural dynamics of ABCB4.

Overall structural dynamics in symmetric or asymmetric membranes exhibit similar patterns according to ABC conformational space, even though a larger variability was observed for asymmetric membrane (Figure 1B&C). Similar behaviors were reported for MRP1 MD simulations conducted in different membrane simple models, for which the lipid bilayer was suggested not to significantly affect local minima, but likely kinetics between milestone steps along transport cycle[11]. Taking advantage of our extensive unbiased MD simulations, conformational subspaces were featured in terms of free energy using the InfleCS clustering method[20] (Figure 2A). It is important to note that in the present study, μs-scale sampling is not sufficient to accurately investigate free energy transitions between conformational states. Likewise, the known TMH kinking upon substrate binding for ABCB4[3] and ABCB1/P-gp[21] was not considered here for conformational transition between different states. We first evaluated different populations per conformation and per bound state (Figures S1-S5) according to the aforementioned ABC structural parameters. MD simulations performed in asymmetric lipid bilayer exhibit a larger structural variability of ABCB4 conformations leading to more conformational sub-states (Table S2 and Figures S1-S5), except for the ^{CC}ABCB4-(ATP)$_2$ conformation (Figure S5); which is restrained in the same subspace regardless of the membrane composition.

Structural overlap between IF states is more favored in an asymmetric membrane than in symmetric lipid bilayer (Figure 1). Therefore, we compiled ABC structural parameters of each conformation and bound state as a single dataset in order to assess potential structural overlaps along alternating access (Figure 2A&B). This is particularly true for ^{IF}ABCB4-PC-(ATP)$_2$ which strongly overlaps with conformational subspace of ^{IF}ABCB4-PC and ^{IF}ABCB4-(ATP)$_2$ states, even though the main representative snapshot looks to be off. Main representative snapshots obtained from InfleCS clustering in symmetric and asymmetric membranes were superimposed to assess the structural deviations (Figure 2C). While ^{IF}ABCB4-PC, ^{IF}ABCB4-(ATP)$_2$, and ^{CC}ABCB4-(ATP)$_2$ states exhibit RMSD differences ranging from 2.6 to 3.7 Å, the largest deviations were obtained for apo ^{IF}ABCB4 and ^{IF}ABCB4-PC-(ATP)$_2$. This can be easily explained for the former since apo ^{IF}ABCB4 exhibits the largest structural dynamics in both membranes as compared with other states, leading to more clusters. For ^{IF}ABCB4-PC-(ATP)$_2$ state, we observed that NBD2 is interestingly shifted as compared to NBD1 in both membranes (Figure 2C). Similar NBD arrangement was also observed with MRP1 state bound to substrate and nucleotide. Such a shift is more important in our asymmetric membrane model suggesting that the TMD conformational transition toward the OF conformation might trigger PC extrusion but also proper NBD dimerization. This

suggests that surrounding lipids may have a more important role in driving conformational transitions rather than modulating ABCB4 conformations.

2.2. On the interplay between surrounding lipids and ABCB4 states.

2.2.1. Conformation-dependent membrane properties of lipid bilayers.

To assess membrane fluidity, we first calculated lipid order parameters (S_{CD}) with particular focus on common lipid tails, namely sn_1-palmitate and sn_2-oleate tails (Figure 3A). Surprisingly, higher lipid order parameters were obtained in an asymmetric membrane than in the symmetric case (Figure 3A), suggesting that the former asymmetric membrane is less fluid than the symmetric one. We distinguished lipid order parameters per leaflet to assess the potential protein-dependent asymmetric fluidity. Interestingly, MD simulations suggest that the aforementioned increase in lipid order is slightly more pronounced within the inner leaflet than in the outer leaflet (Figure 3A) for all ABCB4 conformations and states. Furthermore, we observed stronger impact from ABCB4 conformation onto lipid order in symmetric model. Indeed, MD simulations performed with ^{CC}ABCB4-$(ATP)_2$ exhibited slightly higher fluidity than ^{IF}ABCB4 conformations in a symmetric membrane while in an asymmetric membrane, there is no significant difference between ABCB4 conformations and states. We also measured membrane thickness from phosphate-phosphate distances (Figure S6) showing that the asymmetric membrane is systematically thicker than the symmetric membrane. Interestingly, apo ^{IF}ABCB4 state exhibits slighter thinner bilayers for both asymmetric and symmetric models. This might be assigned to the larger structural variability of apo ^{IF}ABCB4 state leading to membrane adaptation. However, this effect is more pronounced in symmetric model than in asymmetric membrane.

2.2.2. Conformation-dependent cholesterol binding spots

We first investigated potential cholesterol binding sites to compare with former investigations conducted on ABCB1/P-gp since it shares 76% identity and 86% similarity with the ABCB4 transporter[12]. Likewise, key cholesterol binding sites were proposed for their cousins MRP1[11] and ABCB11[13] suggesting an active role for allosteric communication along the protein. Therefore, we probed membrane- and conformation-dependent cholesterol binding sites of ABCB4 from MD simulations.

As observed for P-gp and MRP1, focusing solely on cholesterol recognition consensus motifs (namely CRAC and CARC) is not sufficient to properly describe cholesterol binding. In ABCB4, 4 CRAC and 10 CARC motifs were identified in the transmembrane region, 2 CRAC and 6 CARC motifs for which overlap was observed as for P-gp and MRP1 (Table S3). Cholesterol binding sites were probed by calculating (i) atomistic contacts between cholesterol at the membrane interface along MD simulations and (ii) protein interface for which the likelihood of finding cholesterol molecules is higher than 60% (Figure 3B-3E and Figures S8-S9). Overall, TMH3, TMH4, TMH5, and TMH6 seem to be involved in cholesterol binding spots for both IF

and CC conformations. We also observed that a cholesterol hotspot is found between TMH4 and TMH3 for apo ^{IF}ABCB4 and ^{IF}ABCB4-(ATP)$_2$ systems (Figure S7). MD simulations suggested that bound cholesterol may preclude the entry of PC substrate through the TMH4-TMH6 front gate, even though contact analysis showed a smaller contact fraction in the asymmetric membrane (Figure S8). Interestingly, we observed conformation dependence for potential cholesterol binding sites in which two patterns can be identified for apo ^{IF}ABCB4/^{IF}ABCB4-(ATP)$_2$ and ^{IF}ABCB4-PC/ ^{IF}ABCB4-PC-(ATP)$_2$/^{CC}ABCB4-(ATP)$_2$ (see Figure 3B-3E and Figures S8-S9). The main difference relies on the appearance of two potential binding sites involving TMH10 and TMH11 for PC-bound ^{IF}ABCB4 states as well as ^{cc}ABCB4. Our results confirm that CRAC and CARC motifs are not sufficient to probe cholesterol binding for ABC transporters in agreement with observations made for MRP1[11] and P-gp[12]. Indeed, large scale conformational changes along transport cycle substantially modify residue exposition to lipid bilayer membrane, and therefore protein-lipid interactions.

2.2.3. Non-covalent protein-phospholipid interactions

We here mostly focused on other phospholipids than PC lipids since out of their prominent presence in lipid bilayers, they are also potential substrates of ABCB4, especially considering the credit-card swipe mechanism (see Section 3.3). Given the asymmetric repartition of phospholipids, lipid hotspots were mostly observed at the protein-membrane interface in the inner leaflet (Figure 4A). Interestingly, we observed strong lipid hotspots (*i.e.*, for which polar heads are present for more than 60% of MD simulations and for the three different replicas) for phosphatidylserine (PS) and phosphatidic acid (PA) polar heads in close contact with regions rich in cationic residues at the membrane-water interface (Figure 4A). This suggests the importance of electrostatic contributions in maintaining anionic polar head lipids in close proximity to ABCB4. We also investigated the importance of H-bond networks to rationalize protein-lipid interplay. Interestingly, H-bond protein-lipid network is globally maintained for both asymmetric and asymmetric membranes. However, apo ^{IF}ABCB4 systematically exhibited a weaker H-bond network than all other conformations (Figure 4B). This might be explained by its larger structural variability, for which the larger apo ^{IF}ABCB4 structural dynamics might lead to more frequent H-bond formation/breaking events.

While comparing symmetric and asymmetric membranes, average protein-lipid H-bond counts are similar for all conformations and states but ^{CC}ABCB4-(ATP)$_2$ for which the average H-bond number is slightly smaller in asymmetric membrane than in symmetric one. In asymmetric membrane, the lower abundance of PC lipids in the inner leaflet is compensated by the presence of anionic PS phospholipids to maintain protein-lipid H-bond network. PA phospholipid poorly participated to H-bond network, suggesting that PA hotspots are mostly driven by electrostatic interactions.

Protein-wise, the overall pattern of protein-lipid H-bond network is slightly affected by membrane composition. While focusing on H-bond only present for 35% along MD simulations, 65% of residues are shared between symmetric and asymmetric membranes, with similar scores (Table S4). Overall, the dynamic protein-lipid H-bond network involved mostly intracellular regions of the so-called bundle A, *i.e.*, TMH1, TMH2, TMH11, and TMH12 (see Figure 4C). As expected with the partial atomic charge of polar heads, most of ABCB4 residues involved in protein-lipid H-bond network are polar residues such as glutamate, lysine, and arginine. It is worth mentioning that Glu52 which has been shown to be involved in credit-card swipe mechanism is also strongly involved in protein-lipid H-bond network as its almost neighbor Lys54. Interestingly, these two residues which are respectively anionic and cationic might match with zwitterionic state of PC lipids. MD simulations also revealed the involvement of pre-TMH1 and pre-TMH7 elbow helix residues as key for protein-lipid network (namely, Thr44, Ser696, and to a lesser extend Lys705, see Figure 4C and Table S4). H-bond analyses suggested an almost permanent H-bond with Arg361 located in TMH6, regardless polar head type.

2.3. PC lipid as substrate of ABCB4

2.3.1. Structural impact of PC substrate canonical binding onto ABCB4 structure

PC binding into the canonical substrate binding pocket led to the opening IC angle associated with the triggering of NBD dimerization when nucleotides are present (Figures 1&2). It is worth mentioning that even if the main population of ^{IF}ABCB4-PC-(ATP)$_2$ observed from MD simulations exhibited larger NBD distance (See Figure 1), it represents approximately only 43% (populations C1 and C7, see Figure S4). Other subpopulations (C2 to C6 and C8-C9, see Figure S4) with NBD distances in similar range as ^{IF}ABCB4-(ATP)$_2$ state correspond to 57% of MD simulations. This might be explained by the cryo-EM structure of PC-bound ABCB4 which is likely more open than *in situ*, and the MD sampling was not sufficient to systematically reach the proper NBD distance. To document on PC substrate entry, we conducted structural analyses of front and back (or rear) entry gates that have been described in the literature for ABCB1 [21] and ABCB4 [9]. These gates consist of transmembrane regions of TMH3/TMH4/TMH6 or TMH9/TMH10/TMH12 for front and back gates, respectively (Figure 5A). The front gate exhibited a significant structural difference between PC-bound IF conformations and "apo" states (*i.e.*, apo ^{IF}ABCB4 and ^{IF}ABCB4-(ATP)$_2$) as pictured by RMSD (Figure 5B).

The presence of PC substrate in the canonical binding pocket led to the motion of TMH6 which might be associated with a steric hindrance precluding the entry of other substrates (Figure 5B) while non PC-bound ^{IF}ABCB4 exhibited more open conformation. However, we did not observe any spontaneous PC or other phospholipid entry into the binding pocket. This might be explained by the relatively short sampling of our MD simulations (2μs) while such an event

might require longer timescale. To a lesser extent, back/rear gate is more similar between different states but for one replica (Figure 5B&C). MD simulations suggest a TMH10 motion closing the gate entry (Figure 5C&D). ^{CC}ABCB4-(ATP)$_2$ front and back/rear gates exhibited slightly different conformations as compared to PC-bound states; even though they both remained closed along MD simulations. These observations were made with both symmetric and asymmetric membranes, suggesting that lipid composition does not have a relevant impact.

We performed PCA analysis considering each conformation and each state to better decipher the impact of PC substrate binding onto the TMH arrangement of ABCB4 (See Figure S9). Both TMH4 and TMH10 were identified as main source of structural variability along the alternating access. In presence of PC substrate, TMH4 and TMH10 exhibited kinking event leading to an important discontinuity in TMH helicities (Figure 5D and Table S5). In other words, the presence of PC substrate was associated with TMH4 and TMH10 kinking and adopting a Y-shaped cavity[3] likely to restrict substrate in the binding pocket, precluding its exit. Our findings are in agreement with observations made with ABCB1/P-gp [14]. This was also confirmed by measuring TMH helicity along MD simulations (Table S5), in which *e.g.*, TMH4 and TMH10 exhibit helicities of 0.94 ± 0.21 and 0.85 ± 0.25 in apo ^{IF}ABCB4 and 0.78 ± 0.36 and 0.69 ± 0.35 in ^{IF}ABCB4-PC-(ATP)$_2$ in asymmetric membrane. Interestingly, kinking of TMH10 was also observed in at least one replica of ^{IF}ABCB4-(ATP)$_2$. Noteworthy, both TMH4 and TMH10 are directly involved in the interactions between NBDs and TMDs through the "ball-and-socket" arrangement between NBDs and TMH through the intracellular coupling helices including those connecting TMH4-TMH5 and TMH10-TMH11. Therefore, proper ATP binding might propagate to TMH domains as proposed for other ABC transporters[10,22]. One can thus assume a synergistic effect of ATP and PC substrate binding to favor such an event, even though the latter might be sufficient.

2.3.2. Structural impact of PC substrate canonical binding onto ABCB4 structure

Taking advantage of the cryo-EM structure of ^{IF}ABCB4-PC state, we investigated key amino acid involved in PC binding (Figure 6A). In line with cryo-EM structure, the central role of Trp234, Phe345, His989[3] (located in TMH4, TMH6, and TMH12, respectively) were also observed in our MD simulations. For instance, calculated atomic contact fractions in asymmetric model are respectively 9.23, 30.64 and 25.63 for ^{IF}ABCB4-PC and 29.42, 26.52 and 19.63 for ^{IF}ABCB4-PC-(ATP)$_2$ (see Table S6). MD simulations also suggested Phe305 (TMH5) and Gln725 (TMH7) as involved in PC substrate binding (Figure 6A), contact fractions being respectively 17.14 and 22.38 for ^{IF}ABCB4-PC and 4.61 and 22.15 for ^{IF}ABCB4-PC-(ATP)$_2$. The presence of three aromatic residues suggests a network of cation-π dispersive interactions maintaining PC substrate into a so-called cation-π cage (Figure 6A).

Noteworthy, Trp234 was also proposed to play a central role in (i) substrate entry and then (ii) maintaining PC substrate in the canonical hydrophobic binding pocket by means of cation-π interactions[3]. His989 and Gln725 play also an important role through H-bond network between phosphate moiety with residue side chains (Table S7). For instance, our MD simulations showed that PC substrate and His989 side chain were involved in H-bond for 35% (26%) and 32% (6%) respectively for ^{IF}ABCB4-PC-(ATP)$_2$ and ^{IF}ABCB4-PC in asymmetric (symmetric membrane). Likewise, Gln725 and PC substrate interact by H-bond for 15% (27%) and 7% (33%) for ^{IF}ABCB4-PC-(ATP)$_2$ and ^{IF}ABCB4-PC in asymmetric (symmetric membrane), respectively. We can observe that H-bond exchange might occur between His989 to Gln725 (Figure 6A), the former being deeper in membrane than the latter.

Taking advantage of our 1 µs MD simulations, we also estimated the free energy surface of PC substrate according to the distance of PC polar head with membrane center and the orientation of PC polar head as compared to membrane normal axis (Figure 6B&C). We considered ^{IF}ABCB4-PC-(ATP)$_2$ and ^{IF}ABCB4-PC to assess the coupling between nucleotide binding to NBS with PC substrate (Figure 6A). Even though the sampling of our MD simulations was not sufficient to observe expected PC flip-flop in the binding pocket, the presence of ATP bound to both NBS led to a local free energy minimum in the conformational space of PC polar head crossing the membrane center and properly oriented towards outer leaflet (Figure 6C). Noteworthy, in absence of ATP, PC polar head never reoriented itself toward outer leaflet, even if most of local minima have crossed membrane center. One can thus assume that proper PC binding is sufficient to help PC polar head to cross membrane center, but ATP binding and subsequent TMH arrangement are required to favor PC polar head flip-flop.

2.3.3. Structural insights into the potential credit-card swipe mechanism from µs-scale MD simulations.

Credit-card swipe PC translocation was described from non-conserved polar amino acids (Gln52, Ser58 and Ser69) between ABCB1 and ABCB4, the former being a "drug transporter" for which PC translocation has not been formerly confirmed yet [14,23,24]. These amino acid side chains are oriented toward the membrane in ^{CC}ABCB4-(ATP)$_2$ conformation.

Along our µs-scaled MD simulations, we confirmed previous observations made from shorter simulations in which side chain orientation toward lipid bilayer membrane was maintained, even for the deep Ser69 which is not located at the interface between high-density polar head and high-density lipid tail regions (Figure 6D&E). Unfortunately, we were not able to observe PC flip-flop events along our MD simulations nor specific binding to those amino acids, which is in contrast to the previous work of Prescher et al.[9]. This could be explained by the (i)

insufficient time scale from our µs-MD simulations and (ii) the estimated free energy barrier obtained by umbrella sampling simulations ranging from 10 and 20 kcal.mol^{-1}.

Credit-card swipe relying on a TMH-assisted translocation event, we monitored amino acids that might favor proper flip-flop of PC substrate (*i.e.*, lipid tail reorientation, see Figure 6D&E). We identified polar residues in the outer leaflet that may participate in shielding zwitterionic PC polar head through electrostatic and H-bond interactions, namely Arg212 (TMH3), Glu80 (TMH1), and Asp83 (TMH1). In this context, we measured average distance between Arg212 and Ser69 which exhibit similar values as between Ser58 and Ser69 (ca. 16Å). Likewise, calculated densities (Figure 6E) exhibited Arg212-Ser69 distance ranges from 14.4 and 18.2 Å in asymmetric membrane (Figure 6D). Noteworthy, PC polar head cross sectional diameter was shown to be ca 10 Å[25]. This suggests that while bound to Ser69, PC is within the range of short-distance electrostatic interactions, and translocation may happen from Ser69 to Arg212. In other words, even though we could not confirm PC translocation from all atom MD simulations owing to too short time scale, our results are in line with former investigations suggesting that such a mechanism remains a potential event with ABCB4.

3. Discussion

In the present study, all-atom MD simulations were carried out to investigate the overall conformational space and the lipid-protein interplay of ABCB4. Several conformations and bound states were considered taking advantage of cryo-EM structures resolved in nanodiscs mimicking a native environment of the canalicular membrane. By conducting µs-long unbiased MD simulations, ABC structural patterns of ABCB4 conformations and bound states were investigated considering two lipid bilayer membrane models, namely symmetric (made of POPC:POPE:Chol (2:1:1)) and asymmetric. Protein-lipid interplay was investigated by assessing (i) the impact of membrane composition on the structural dynamics of ABCB4 and (ii) conformation-dependent protein-lipid contact patterns driven by non-covalent interactions. Special attention was also made on substrate-ABCB4 interactions considering either a canonical alternating access[3] as well as a credit-card swipe mechanism[9]. In line with recent observations made for ABCB4 cousin, namely P-gp/ABCB1[14], TMH4 and TMH10 kinking was also investigated given the different TMH conformations observed from cryo-EM resolved structures. Large-scale conformational transitions were not investigated in the present study given the challenge to define the proper driving force responsible for either IF-to-OF or OF-to-UR conformational transitions.

3.1. Membrane-dependent structural dynamics of ABCB4.

While ABCB1/P-gp has been extensively studied over the past decades, recent studies highlighted that knowledge of ABCB family still requires further investigations. This is particularly true for ABCB4 which is not a drug transporter even though studies showed that

few xenobiotics may be translocated by ABCB4[9,26,27]. It is important to note that the present study is limited by forcefield inaccuracies as well as limited conformational sampling. However, our set of MD simulations appear relevant to decipher local conformational space of each conformer and bound state including potential overlaps.

In contrast to what has been observed from MD simulations of ABCC1/MRP1[11], ABC structural parameters (*i.e.*, IC angle, NBD distance, and NBD twist) measured along the ABCB4 simulations remain close to cryo-EM experimental structures. This can be easily explained by MRP1 cryo-EM resolution in detergent leading to wide open IF structure. One can also hypothesize that the degenerated NBS in C-family ABC transporter can be associated with permanent single ATP-bound state requiring the closing of IC gate and NBD dimerization[11]. Notably, ATP hydrolysis for non-degenerated ABC transporter as ABCB4 was however proposed to alternate single ATP hydrolysis from NBS to another along several transport cycles [28].

Our MD simulations are in line with previous experimental and theoretical investigations for which ABC IC angle and NBD distance motions are not necessarily coupled**.** This is particularly true for ABCB4 while considering the canonical alternating access since PC substrate binding is associated with increase of IC angle owing to steric hindrance in the binding pocket. On the other hand, NBD distance shortening requires nucleotide binding since ATP is known to be sandwiched by both NBDs[6]. However, none of our MD simulations conducted with IF ATP-bound state (*i.e.*, ^{IF}ABCB4-(ATP)$_2$ and ^{IF}ABCB4-PC-(ATP)$_2$) reached proper NBD dimerization as pictured by NBD twist distribution. This is due to the insufficient sampling at the μs-scale. One can infer that proper ATP hydrolysis competent NBD dimerization, as observed in ^{CC}ABCB4-(ATP)$_2$, requires TMD large scale conformational transition which then propagate to NBD conformation. We observed translation of NBDs along the membrane when bound to both PC substrate and ATP (*i.e.*, ^{IF}ABCB4-PC-(ATP)$_2$). Interestingly, similar shifted NBD arrangement was also observed for MRP1 bound to nucleotides and substrate [11]. One can hypothesize from these MD simulations conducted with two mammalian ABC transporters that such NBD conformation is a transient state prior to TMD large scale conformational transition. Unfortunately, there is not a single resolved cryo-EM structure of mammalian fold IV ABC transporter to validate such hypothesis.

The MD simulations did not reveal substantial impact of membrane composition onto conformational local minima. Overall, the same conformational space was sampled along MD simulations regardless of membrane composition. Interestingly, asymmetric membrane simulations showed more structural variability and flexibility as compared to simulations conducted in symmetric membrane. The larger structural variability in asymmetric membrane and proposed lower energy barriers between different ABC conformations suggest that membrane composition might mostly affect kinetics of ABCB4 transport cycle. However, free

energy surfaces proposed in the present manuscript must be deemed very carefully in the framework of alternating access. Indeed, used structural parameters used do not reflect the actual driving forces of large-scale conformational transitions for type IV ABC transporters. Recent experimental investigations conducted on MRP4 have proposed that lipid environment might modulate, at least ATPase activity[15], for which native environment was associated with higher activity than in POPC-reconstituted nanodiscs. In other words, lipid composition is likely important for ABC transporter kinetics.

3.2. Protein-lipid interplay and annular lipids of ABCB4

Present MD simulations were also used to investigate protein-lipid interactions and individuate potential conformation-dependent lipid fingerprints. First, by monitoring membrane properties, we observed that sn_1-palmitate and sn_2-oleate lipid order parameters were systematically more ordered in asymmetric membrane than in symmetric model. Such a finding is counter-intuitive with the larger flexibility observed for asymmetric conformational sampling in asymmetric membrane. Noteworthy, lipid order parameters were calculated only for sn_1-palmitate and sn_2-oleate lipids for sake of comparison with symmetric membrane. Therefore, ~20% of lipid tails were not included to properly assess membrane fluidity. Besides, membrane thicknesses were calculated for which asymmetric membrane is ~2 Å thicker than symmetric one, leading to larger protein-lipid interface. Altogether, these findings including those regarding ABCB4 conformational flexibility raise the question of the iterative interplay between protein and membrane lipid bilayer, *i.e.,* how do lipids modulate protein flexibility as well as in turn how protein affect membrane properties.

In this context, our results highlighted conformational-dependent pattern of lipid protein contact and non-covalent interactions. While it is known that the use of CRAC and CARC motifs, at least in ABC transporters [11,12], is not sufficient to predict and investigate cholesterol binding, MD simulations clearly suggest that cholesterol binding as allosteric partners differs from one conformation to another (*e.g.,* ^{IF}ABCB4 and ^{CC}ABCB4). Interestingly, as observed in MRP1[11,29], we observed cholesterol hotspot in the pre-TMH elbow helix suggesting a specific role of cholesterol in this region. Given that annular lipids, including cholesterol, were shown to actively participate to distant communication between *e.g.,* NBS and substrate binding site, one can hypothesize that cholesterol binding might play different roles in IF-to-OF transition and OF-to-UR-to-IF transition. As it has been very recently suggested for P-gp[14], protein-lipid interplay in the inner leaflet remains more important than the outer leaflet. Indeed, anionic lipids in our asymmetric model are exclusively located in the inner leaflet. Strong electrostatic interactions between anionic polar head and cationic ABCB4 residues are suspected to maintain strong non-covalent network which also differs from one conformation to another. This may be explained by different patterns in terms of residue exposure to high density polar head region according to TMH arrangement. Even though the allosteric role of annular lipid

was not within the scope of the present study, one can infer from other studies[11–16] that membrane itself is not only a physical container for ABCB4 but annular lipids are active player for its function. We can thus expect that lipid composition plays a role in modulating ABCB4 function given that cell membrane structure is highly dynamic and for which, *e.g.*, nanodomains and/or lipid rafts might also affect protein structure and dynamics.

However, our present results regarding the protein-lipid contact should be considered very carefully since only all-atom MD simulations were conducted. In such an approach, the sampling and force field inaccuracy leads to a poor description of lipid lateral diffusion. On the contrary, in the recent study regarding P-gp[14], coarse-grained model was used with Martini 3 force field which was shown to provide robust insights regarding surrounding lipid fingerprint. Besides, it is worth mentioning that all atom MD simulations using asymmetric membranes remain challenging and should be considered carefully. Indeed, asymmetric membrane is associated with differential lateral compressibility between inner and outer leaflets. Nevertheless, the present study only paves the way for further investigations.

3.3. TMD-NBD coupling improve canonical alternating access substrate translocation

Particular attention was paid to substrate entry and interactions in the canonical binding pocket. In contrast to MRP1, P-gp and ABCB4 were shown to bind substrate directly from lipid bilayer, through either the front or rear gate made of TMH3/TMH4/TMH6 or TMH9/TMH10/TMH12, respectively [21,30,31]. No substrate entry was observed in our simulations with either apo [IF]ABCB4 nor [IF]ABCB4-(ATP)$_2$. However, our simulations clearly demonstrate two distinct supramolecular arrangements of the front gate. On one hand, apo [IF]ABCB4 and [IF]ABCB4-(ATP)$_2$ exhibit an open front gate with space between TMH3 and TMH4. One the other hand, [CC]ABCB4-(ATP)$_2$ conformation and PC-bound [IF]ABCB4 states showed TMH6 shift between TMH3 and TMH4. We can here hypothesize that PC substrate binding leads to TMH6 motions closing the front gate in order to preclude (i) substrate exit back to the inner leaflet and (ii) other PC lipid entry. Cryo-EM structures of [IF]ABCB4 highlighted the central role of TMH4 Trp234 residue as PC substrate carrier through cation-π interactions between cationic choline moiety and aromatic tryptophane benzimidazole group. It is important to note that current force fields tend to underestimate such dispersive interactions[32], which can explain why we were not able to observe such events. Despite, MD simulations strengthen the proposed central role of cation-π interactions in the substrate binding pocket. Our results confirmed the central role of Trp234 (TMH4), Phe345 (TMH6) and His989 (TMH12) in maintaining PC substrate as shown by mutagenesis experiments. Furthermore, structural dynamics of ABCB4 also identified two other residues, namely Phe305 (TMH5) and Gln725 (TMH7) which might be involved in PC substrate binding and dynamics. The former is involved

in maintaining the cation-π cage between aromatic ring and choline cationic moiety while the latter is proposed to favor PC polar head translocation across membrane center given its position along membrane normal. While comparing substrate position with MD simulations, PC polar head spontaneously crossed the membrane center. MD simulations highlighted the already known distant coupling between nucleotide binding and substrate binding pocket since PC polar head tends to flop only in presence of ATP bound to NBS. This might suggest that PC substrate may reorient itself along transport cycle. However, the full flopping event was not observed owing to the present insufficient sampling, while being within the current standard for unbiased all-atom MD simulations. Interestingly, PC flopping was not initiated in symmetric membrane. It could be explained by the insufficient sampling especially for such a POPC:POPE:Chol (2:1:1) symmetric model. Indeed, this is in line with the aforementioned lower ATPase activity of MRP4 observed in POPC as compared to native nanodiscs[15]. Therefore, this suggests that asymmetry might also be an important driving force for substrate translocation by ABC transporters.

In contrast to C-family ABC transporters, ABCB1 and ABCB4 were resolved with kinked TMH4 and TMH10 helices in presence of substrate. TMH kinking is expected to at least maintain ABCB4 substrate in substrate binding pocket. The role of membrane and annular lipids in modulating the conformational space and flexibility of P-gp TMHs was thoroughly investigated. Our MD simulations also suggest that TMH4 and TMH10 flexibility may participate in substrate translocation by carrying PC lipid to the outer lipid-water interface. Our MD simulations carried out with ABCB4 are in agreement with P-gp MD simulations[14], exhibiting TMH4 and TMH10 kinking events in presence of substrate. Noteworthy, this can be associated with high identity (similarity) scores between ABCB4 and P-gp TMH4 and TMH10, being 85% (93%) and 76% (93%), respectively. However, we did not observe spontaneous kinking events in apo [IF]ABCB4 systems in spite of longer MD simulations. Spontaneous kinking of TMH10 in absence of PC substrate was only observed when ATP is bound to NBS highlighting the coupling existing between NBDs and TMDs. This can be easily explained by the "ball-and-socket" arrangement between coupling helices and NBD[6]. This is particularly true since TMH4 and TMH10 are directly connected to two of four coupling helices. We can thus assume that proper ATP binding may favor TMH4 and TMH10 kinking event. It seems however that TMH4 and TMH10 kinking remains a specific event for at least few members of B-family ABC transporters. This can also be correlated to the existence of two canonical ATP-hydrolysis competent NBS as compared to *e.g.*, C-family ABC transporters for which no TMH kinking event was reported so far, even in presence of substrate and in native environment. Furthermore, [CC]ABCB4-(ATP)$_2$ exhibit straight TMHs clearly suggesting that substrate translocation might be associated to spontaneous unkinking event which does not require ATP hydrolysis.

3.4. Credit-card swipe mechanism as complementary to potentiate ABCB4 PC floppase activity

Another translocation mechanism has also been proposed regarding PC flopping mediated by ABCB4, supported by joint experimental and computational investigations [9]. The so-called "credit-card swipe mechanism" relies on TMH-mediated translocation events in which a PC lipid located in the inner leaflet PC binds to polar residues exposed at the protein- membrane interface, namely Gln52, Ser58, and Ser69. Importantly, biased MD simulations were carried out strengthening this hypothesis supported by site-directed mutagenesis experiments demonstrating that mutations of these residues are associated with lower ATPase activity[9]. In the present work, spontaneous PC flopping was not observed owing to the length of our MD simulations. However, they clearly showed strong H-bond interactions involving Gln52 and Lys54 with surrounding lipids, including PC lipids in spite of their lower abundance in the inner leaflet. Interestingly, by monitoring the different polar residues which may be involved in carrying PC lipid translocation from inner and outer leaflet, we identified a network of polar residues at the outer lipid-water interface, namely Arg212, Glu80 and Asp83. These residues tend to form a dynamic salt-bridge network in the high density polar head region of membrane outer leaflet. Furthermore, z-projected distance between Ser69 and Arg212 is within the same range as between Ser58 and Ser69 (~16 Å). This distance is sufficient to favor the electrostatic interactions that may shield the polar head in the apolar inner membrane. Therefore, our results still require to be confirmed by further investigations, *e.g.*, by (i) considering coarse grained MD simulations to reach sufficient timescale of observing TMH1-assisted PC translocation and (ii) site-directed mutagenesis experiments by mutating Arg212, Asp83, and Glu80.

Overall, by considering both the canonical alternating access and credit-card swipe mechanisms, we can hypothesize that ABCB4 might be more efficient to favor PC translocation from the inner leaflet to the outer leaflet. Indeed, by mutating key PC residues of canonical alternating access, PC translocation was shown to be less efficient but still possible (from ~ 20 to 50% of wild-type ABCB4 activity when mutating Trp234, Phe345, and His989). This may suggest that the credit-card swipe mechanism can be complementary translocation event leading up to two PC translocations per ABCB4 transport cycle, as pictured in Figure 7. Obviously, such an hypothesis requires further experimental evidence to be carried out.

4. Conclusion

Altogether, the present investigation relying on µs-scaled MD simulations provides robust hints regarding the ABCB4 function by conciliating both canonical alternating access and credit-card mechanisms as crucial for ABCB4-mediated PC transport. ABCB4 is not referred to as drug membrane transporter but as an essential physiological transporter to maintain bile

composition and homeostasis. Therefore, the selectivity and specificity of ABCB4 dysfunction cannot be overcome by increasing functions of other transporters as shown for membrane drug transporters (*e.g.*, MRP2/MRP4 pair). This is even more important for ABCB4 which acts only in the liver canalicular membrane while drug transporter compensation can exist remotely according to the remote sensing signaling theory. We also strengthen the importance of surrounding lipids and membrane for modulating structural features that have been proposed for ABCB4 cousin, namely ABCB1, such as TMH4 and TMH10 kinking. These results require further investigations to be also considered for other mammalians B-family ABC transporters, in contrast to C-family ABC transporters for which the transport cycle might differ.

5. Computational procedures

5.1. Structure

ABCB4 five main states and conformations along canonical transport cycle were considered, namely: apo ^{IF}ABCB4, ^{IF}ABCB4-PC, ^{IF}ABCB4-(ATP)$_2$, ^{IF}ABCB4-PC-(ATP)$_2$, and ^{CC}ABCB4-(ATP)$_2$. ^{IF}ABCB4-based models were built using either apo or PC-bound cryo-EM structures (PDB ID: 7NIU, 7NIV) while ^{CC}ABCB4 was constructed using the resolved cryo-EM structure of ABCB4 adopting CC conformation (PDB ID: 6S7P). The so-called linker 1 connecting the NBD1 to TMH7 was not included in the present molecular model given its expected disordered structure to preclude false non-covalent interactions with resolved domains. Few residues of the first extracellular loop connecting TMH1 and TMH2 being unresolved, missing residues (*i.e.*, from Phe92 to Gly106 for ^{IF}ABCB4 structures - *i.e.*, 7NIU and 7NIV – and Lys84 to Pro105 ^{IF}ABCB4 – *i.e.*, 6S7P) were included using the Modeller v9.23 [33] taking advantage of resolved P-gp structures (PDB ID: 6QEX) [16]. The cryo-EM resolved 1,2-dilinoleoyl-sn-glycero-3-phosphatidylcholine (DLPC) lipid was used for substrate-bound states of ^{IF}ABCB4 conformations (*i.e.*, ^{IF}ABCB4-PC and ^{IF}ABCB4-PC-(ATP)$_2$. Since resolved ^{IF}ABCB4 structures do not include ATP and Mg^{2+} molecules, ATP-bound states were built by independently superimposing ^{IF}ABCB4 NBDs onto those obtained from ^{CC}ABCB4 conformation in which ATP and Mg^{2+} were expected to be properly docked to at least NBD A-loop and signature motifs. Protonation states of titratable amino acids were predicted using the propKa3.0 software[34,35] Histidine residues were protonated by visual inspection of their potential involvement in H-bond network with surrounding amino acids. For sake of comparison, the same protonation states were used for each model.

Molecular models were briefly minimized in vacuum to remove potential steric clashes using the Amber 20 package[36,37] prior to embedding into lipid bilayer membranes using the CHARMM-GUI web server[38–40]. Models were first carefully aligned with ^{IF}ABCB4 and ^{CC}ABCB4 coordinates in the OPM (Orientations of proteins in membranes) database[41] to mimic proper protein alignment in membrane models. Two models of lipid bilayers were here considered to

investigate the protein-lipid interplay and the dependence on membrane composition. A classical but simplistic symmetric lipid bilayer made of 1-palmitoyl-2-oleyl-sn-glycero-3-phosphatidylcholine (POPC), 1-palmitoyl-2-oleyl-sn-glycero-3-phosphatidylethanolamine (POPE) and cholesterol (Chol) with a molecular ratio of POPC:POPE:Chol (2:1:1). Besides, a more complex asymmetric lipid bilayer composition was also considered made of different lipids. PC, PE, Chol but also phosphatidic acid (PA) and phosphatidylserine (PS); lipid polar heads were built as well as different lipid tails such palmitoyl (PA, 16:0), oleoyl (OL, 18:1), arachidonyl (AR, 20:4), docosahexaenoyl (DHA, 22:6). Lipid compositions in inner and outer leaflet for each model are reported in Table S8 and S9. Overall, the number of lipids in all the membranes is slightly larger than 400 leading to system size of ca. 120x120x180 $Å^3$ including water and NaCl counterions set up at the extracellular physiological concentration (*i.e.*, 0.15 M, system details are reported in Table S10). The scripts included in AmberTools package (namely, charmmlipid2amber.py and pdb4amber) were employed to convert the CHARMM-GUI inputs in an Amber-compatible file format. For bound states, PC substrate, ATP molecules, and Mg^{2+} ions were then added by superimposing models built initially without lipid bilayer. For each system, the corresponding amount of counterions was removed to maintain the neutrality of the system.

5.2. Molecular dynamics simulations

Amino acids, lipids, water, and ATP molecules were respectively modelled using the FF14SB[42], lipid21[43], TIP3P[44–46], and DNA.OL15-based forcefields[47] for bound models. Apo ^{IF}ABCB4 systems were modelled using the latest FF19SB[48] with OPC[49] model for water molecules. Such a difference of forcefield models between apo and bound states was due to unexpected forcefield incompatibilities. Using FF19SB and OPC forcefield with substrates surprisingly leads to unreliable restraints onto amino acid backbone and sidechains. Mg^{2+} ions and NaCl counterions were modeled using the mono and divalent parameters suited either to TIP3P[50] or OPC [49] water force fields, with FF14SB or FF19SB, respetively. ABCB4 PC substrate (*i.e.*, DLPC) was modelled using the lipid21 forcefield. The cutoff for non-bonded interaction was set at 10 Å for both Coulomb and van der Waals potentials. Long-range electrostatic interactions were computed using the particle mesh Ewald method [51].

Simulations were conducted using the Amber20 and Amber22 packages[52,53] using CPU and GPU versions of the PMEMD code. First, minimizations using energy steepest gradient were conducted as follows: (i) minimization of water O-atoms for 20,000 steps, (ii) all bonds involving H-atoms for 20,000 steps, (iii) water molecules and counterions for 50,000 steps, and (iv) the whole system for 50,000 steps. For each model, system thermalizations were performed using Langevin thermostat in two steps. First, water molecules were thermalized from 0 to 100 K for 50 ps in the NVT ensemble with 0.5 fs as time integration. Then, the whole system was heated up from 100 K to 310 K for 500 ps in the NPT ensemble using a time

integration of 2 fs, and restraining bonds involving H-atoms with the SHAKE algorithm. Pressure and box equilibrations were then achieved by performing short MD simulations for 5ns with a integration time set at 2 fs in the semi-isotropic NPT ensemble for which Berendsen barostat [54] was employed. Finally, for each model, MD productions of 2 μs were performed using 2 fs as integration time step, in the NPT ensemble condition with semi-isotropic scaling. Langevin dynamics thermostat [55] with 1.0 ps^{-1} collision frequency was used to maintain the temperature. The overall pressure was maintained at 1 bar with semi-isotropic pressure scaling using Berendsen barostat.

Noteworthy, ATP-bound models were carried out using a restraint MD approach as used for ABCC1 transporter [11] and initially proposed by Wen et al. in 2013 [56] to ensure the different induced fit for ATP-Mg^{2+}-NBD interactions. Briefly, for equilibration steps, distance between each ATP purine moiety and NBD1&2 A-loop tyrosines (namely Tyr 403 and Tyr 1043) were restrained as well as H-bond between phosphate moieties, Mg^{2+} atoms with surrounding Walker A serine residues (Ser436 and Ser1076) and Q-loop glutamine residues (Gln477 and Gln1206) to maintain proper ATP docking in NBS. For each restraint, a harmonic potential was applied to restrain distances. The distance restraints were applied during the thermalization, equilibration, but also the first 10 ns of the production. They were then smoothly removed over 10 ns. The remaining part of the production run was performed without restraints.

For each model, three replicas were conducted from the thermalization stage to consider a minimum of conformational variability. The aggregated total MD simulation time in the present work is ~60 μs (see Table S11).

5.3. Analysis and visualization

For each system and given time-dependent root-mean square deviations (RMSD) evolution over MD simulations (see Figure S10), analyses were performed over the last 1 μs of each production run. Trajectories were saved every 100 ps. Analyses were conducted with the CPPTRAJ[57] and MDAnalysis[58,59] softwares as well as in-house python scripts. Plots were created using the matplotlib v3.1.1 python package [60]. Trajectories visualization and image rendering were made using the VMD software (versions 1.9.3 and alpha-v1.9.4)[61].

The so-called ABC structural parameters, namely, intracellular (IC) and extracellular (EC) angles, NBD distance, and NBD twist were used to assess the conformational difference of the different models. They were defined according to previous studies by Moradi et al.[18], Hofmann et al.[19], and Tóth et al.[11] (see Figure 1A). Briefly, IC was defined as the angle between vectors made of the EC TMH region center-of-mass towards the centers-of-mass of IC TMH regions bound to either NBD1 (*i.e.*, TMH1, TMH2, TMH3, TMH10, TMH11, and TMH6) or NBD2 (*i.e.*, TMH4, TMH5, TMH7, TMH8, TMH9, and TMH12). Likewise, EC angle was defined by the angles originating from the center-of-mass of intracellular regions of TMHs towards the extracellular regions of either TMH1, TMH2, TMH11, TMH12 or TMH5, TMH6,

TMH7. The NBD distance was defined by the distance between the centers-of-mass of each NBD; and finally, the NBD twist measures the dihedral angle between the two NBDs, defined by four centers-of-mass made of alpha-helical subdomains and beta-sheet subdomain of each NBD. The TMH and NBD topology used for ABCB4 protein is reported in Table S12.

Structural principal component analyses (PCA) were also performed with the CPPTRAJ package[57], focusing on the ABC core *i.e.*, backbone atoms of TMH1 to TMH12 and NBD1 and NBD2. PCA was, as well, performed focusing only on the core part of the helices. In the first case, the system variability was measured by considering each replica per system independently aligned to an average structure of the system. In the second case, PCA was carried out by aligning all the systems to an average structure of the CC conformation to measure the overall variability along the transport cycle. For PC-bound states, contacts between PC substrate and ABCB4 were investigated using the CPPTRAJ package with the keyword *nativecontacts*.

Lipid order was calculated with the CPPTRAJ package using the keyword *lipidscd*. Polar-head oriented lipid distributions and reside[54,55]ncies were measured using the CPPTRAJ package (*grid* keyword with a resolution of 1 Å). For each polar head or entire cholesterol, the threshold of lipid occupancies was set up to highlight regions for which the likelihood of finding a given polar head was higher than 60% over the analysis. Polar head and cholesterol contacts have been investigated by CPPTRAJ package.

6. Data Availability

Initial and final structures for all simulations, MD input examples and ATP force field parameters used are available upon reasonable request as well as MD trajectories.

7. Acknowledgments

The authors gratefully acknowledge support from Xavier Montagutelli (University de Limoges, France) for computational and technical support, from the regional supercomputers CALI ("CAlcul en Limousin") and "Baba Yaga", as well as from the "Jean-Zay" national[56] supercomputer from IDRIS HPC resources under the allocations 2020-A0080711487, 2021-A0100711487 and 2022-A0120711487 made by GENCI. We also thank Prof. Lutz Schmitt for fruitful scientific discussion and suggestions about the present manuscript.

8. Author Contribution – CrediT

V. Crespi: Conceptualization, Investigation, Methodology, Data curation, Formal analysis, Writing – Original draft, Writing – review & editing

A. Tóth: Methodology, Formal analysis, Writing – review & editing

A. Janaszkiewicz: Methodology, Formal analysis, Writing – review & editing

T. Falguières: Investigation, Supervision, Project administration, Writing - review and editing

F. Di Meo: Conceptualization, Investigation, Methodology, Data curation, Formal analysis, Funding acquisition, Project administration, Supervision, Writing – original draft, Writing - review and editing

9. Funding

This work was supported by the [57]French Government's "Plan de relance" (DIGPHAT 22-PESN-0017) and by grants from the French Research Agency (ANR-21-CE18-0030-01, ANR-19-CE17-0008-01) and from Inserm and Région Nouvelle Aquitaine (AAP-NA-ESR 2019 VICTOR and 2023 MUSYPHA).

10. References

(1) Olsen, J. A.; Alam, A.; Kowal, J.; Stieger, B.; Locher, K. P. Structure of the Human Lipid Exporter ABCB4 in a Lipid Environment. *Nat. Struct. Mol. Biol.* **2020**, *27* (1), 62–70. https://doi.org/10.1038/s41594-019-0354-3.

(2) Falguières, T.; Aït-Slimane, T.; Housset, C.; Maurice, M. ABCB4: Insights from Pathobiology into Therapy. *Clin. Res. Hepatol. Gastroenterol.* **2014**, *38* (5), 557–563. https://doi.org/10.1016/j.clinre.2014.03.001.

(3) Nosol, K.; Bang-Sørensen, R.; Irobalieva, R. N.; Erramilli, S. K.; Stieger, B.; Kossiakoff, A. A.; Locher, K. P. Structures of ABCB4 Provide Insight into Phosphatidylcholine Translocation. *Proc. Natl. Acad. Sci.* **2021**, *118* (33), e2106702118. https://doi.org/10.1073/pnas.2106702118.

(4) Mahdi, Z. M.; Synal-Hermanns, U.; Yoker, A.; Locher, K. P.; Stieger, B. Role of Multidrug Resistance Protein 3 in Antifungal-Induced Cholestasis. *Mol. Pharmacol.* **2016**, *90* (1), 23–34. https://doi.org/10.1124/mol.116.103390.

(5) Frampton, J. E.; Scott, L. J. Posaconazole: A Review of Its Use in the Prophylaxis of Invasive Fungal Infections. *Drugs* **2008**, *68* (7), 993–1016. https://doi.org/10.2165/00003495-200868070-00008.

(6) Thomas, C.; Tampé, R. Structural and Mechanistic Principles of ABC Transporters. *Annu. Rev. Biochem.* **2020**, *89* (1), 605–636. https://doi.org/10.1146/annurev-biochem-011520-105201.

(7) Kroll, T.; Prescher, M.; Smits, S. H. J.; Schmitt, L. Structure and Function of Hepatobiliary ATP Binding Cassette Transporters. *Chem. Rev.* **2021**, *121* (9), 5240–5288. https://doi.org/10.1021/acs.chemrev.0c00659.

(8) Prescher, M.; Smits, S. H. J.; Schmitt, L. Stimulation of ABCB4/MDR3 ATPase Activity Requires an Intact Phosphatidylcholine Lipid. *J. Lipid Res.* **2020**, *61* (12), 1605–1616. https://doi.org/10.1194/jlr.RA120000889.

(9) Prescher, M.; Bonus, M.; Stindt, J.; Keitel-Anselmino, V.; Smits, S. H. J.; Gohlke, H.; Schmitt, L. Evidence for a Credit-Card-Swipe Mechanism in the Human PC Floppase ABCB4. *Structure* **2021**, *29* (10), 1144-1155.e5. https://doi.org/10.1016/j.str.2021.05.013.

(10) Verhalen, B.; Dastvan, R.; Thangapandian, S.; Peskova, Y.; Koteiche, H. A.; Nakamoto, R. K.; Tajkhorshid, E.; Mchaourab, H. S. Energy Transduction and Alternating Access of the Mammalian ABC Transporter P-Glycoprotein. *Nature* **2017**, *543* (7647), 738–741. https://doi.org/10.1038/nature21414.

(11) Tóth, Á.; Janaszkiewicz, A.; Crespi, V.; Di Meo, F. On the Interplay between Lipids and Asymmetric Dynamics of an NBS Degenerate ABC Transporter. *Commun. Biol.* **2023**, *6* (1), 149. https://doi.org/10.1038/s42003-023-04537-3.

(12) Thangapandian, S.; Kapoor, K.; Tajkhorshid, E. Probing Cholesterol Binding and Translocation in P-Glycoprotein. *Biochim. Biophys. Acta BBA - Biomembr.* **2020**, *1862* (1), 183090. https://doi.org/10.1016/j.bbamem.2019.183090.

(13) Liu, H.; Irobalieva, R. N.; Kowal, J.; Ni, D.; Nosol, K.; Bang-Sørensen, R.; Lancien, L.; Stahlberg, H.; Stieger, B.; Locher, K. P. Structural Basis of Bile Salt Extrusion and Small-Molecule Inhibition in Human BSEP. *Nat. Commun.* **2023**, *14* (1), 7296. https://doi.org/10.1038/s41467-023-43109-1.

(14) De Vecchis, D.; Schäfer, L. V. Coupling the Role of Lipids to the Conformational Dynamics of the ABC Transporter P-Glycoprotein. *Biophys. J.* **2024**, S000634952400417X. https://doi.org/10.1016/j.bpj.2024.06.020.

(15) Bloch, M.; Raj, I.; Pape, T.; Taylor, N. M. I. Structural and Mechanistic Basis of Substrate Transport by the Multidrug Transporter MRP4. *Structure* **2023**, *31* (11), 1407-1418.e6. https://doi.org/10.1016/j.str.2023.08.014.

(16) Alam, A.; Kowal, J.; Broude, E.; Roninson, I.; Locher, K. P. Structural Insight into Substrate and Inhibitor Discrimination by Human P-Glycoprotein. *Science* **2019**, *363* (6428), 753–756. https://doi.org/10.1126/science.aav7102.

(17) Ingólfsson, H. I.; Arnarez, C.; Periole, X.; Marrink, S. J. Computational 'Microscopy' of Cellular Membranes. *J. Cell Sci.* **2016**, jcs.176040. https://doi.org/10.1242/jcs.176040.

(18) Moradi, M.; Tajkhorshid, E. Mechanistic Picture for Conformational Transition of a Membrane Transporter at Atomic Resolution. *Proc. Natl. Acad. Sci.* **2013**, *110* (47), 18916–18921. https://doi.org/10.1073/pnas.1313202110.

(19) Hofmann, S.; Januliene, D.; Mehdipour, A. R.; Thomas, C.; Stefan, E.; Brüchert, S.; Kuhn, B. T.; Geertsma, E. R.; Hummer, G.; Tampé, R.; Moeller, A. Conformation Space of a Heterodimeric ABC Exporter under Turnover Conditions. *Nature* **2019**, *571* (7766), 580–583. https://doi.org/10.1038/s41586-019-1391-0.

(20) Westerlund, A. M.; Delemotte, L. InfleCS: Clustering Free Energy Landscapes with Gaussian Mixtures. *J. Chem. Theory Comput.* **2019**, *15* (12), 6752–6759. https://doi.org/10.1021/acs.jctc.9b00454.

(21) Nosol, K.; Romane, K.; Irobalieva, R. N.; Alam, A.; Kowal, J.; Fujita, N.; Locher, K. P. Cryo-EM Structures Reveal Distinct Mechanisms of Inhibition of the Human Multidrug Transporter ABCB1. *Proc. Natl. Acad. Sci.* **2020**, *117* (42), 26245–26253. https://doi.org/10.1073/pnas.2010264117.

(22) Weng, J.-W.; Fan, K.-N.; Wang, W.-N. The Conformational Transition Pathway of ATP Binding Cassette Transporter MsbA Revealed by Atomistic Simulations. *J. Biol. Chem.* **2010**, *285* (5), 3053–3063. https://doi.org/10.1074/jbc.M109.056432.

(23) Romsicki, Y.; Sharom, F. J. Phospholipid Flippase Activity of the Reconstituted P-Glycoprotein Multidrug Transporter. *Biochemistry* **2001**, *40* (23), 6937–6947. https://doi.org/10.1021/bi0024456.

(24) Van Helvoort, A.; Smith, A. J.; Sprong, H.; Fritzsche, I.; Schinkel, A. H.; Borst, P.; Van Meer, G. MDR1 P-Glycoprotein Is a Lipid Translocase of Broad Specificity, While MDR3 P-Glycoprotein Specifically Translocates Phosphatidylcholine. *Cell* **1996**, *87* (3), 507–517. https://doi.org/10.1016/S0092-8674(00)81370-7.

(25) Bronstein, L.-G.; Tóth, Á.; Cressey, P.; Rosilio, V.; Di Meo, F.; Makky, A. Phospholipid–Porphyrin Conjugates: Deciphering the Driving Forces behind Their Supramolecular Assemblies. *Nanoscale* **2022**, *14* (19), 7387–7407. https://doi.org/10.1039/D2NR01158A.

(26) Smith, A. J.; Van Helvoort, A.; Van Meer, G.; Szabó, K.; Welker, E.; Szakács, G.; Váradi, A.; Sarkadi, B.; Borst, P. MDR3 P-Glycoprotein, a Phosphatidylcholine Translocase, Transports Several Cytotoxic Drugs and Directly Interacts with Drugs as Judged by Interference with Nucleotide Trapping. *J. Biol. Chem.* **2000**, *275* (31), 23530–23539. https://doi.org/10.1074/jbc.M909002199.

(27) Duan, Z.; Brakora, K. A.; Seiden, M. V. Inhibition of ABCB1 (MDR1) and ABCB4 (MDR3) Expression by Small Interfering RNA and Reversal of Paclitaxel Resistance in Human Ovarian Cancer Cells. *Mol. Cancer Ther.* **2004**, *3* (7), 833–838.

(28) Stockner, T.; Gradisch, R.; Schmitt, L. The Role of the Degenerate Nucleotide Binding Site in Type I ABC Exporters. *FEBS Lett.* **2020**, *594* (23), 3815–3838. https://doi.org/10.1002/1873-3468.13997.

(29) Tóth, Á.; Crespi, V.; Janaszkiewicz, A.; Di Meo, F. Computational and Structural Insights into the Pre- and Post-hydrolysis States of Bovine Multidrug Resistance-associated Protein 1. *Basic Clin. Pharmacol. Toxicol.* **2023**, *133* (5), 508–525. https://doi.org/10.1111/bcpt.13871.

(30) Johnson, Z. L.; Chen, J. Structural Basis of Substrate Recognition by the Multidrug Resistance Protein MRP1. *Cell* **2017**, *168* (6), 1075-1085.e9. https://doi.org/10.1016/j.cell.2017.01.041.

(31) Kroll, T.; Smits, S. H. J.; Schmitt, L. Monomeric Bile Acids Modulate the ATPase Activity of Detergent-Solubilized ABCB4/MDR3. *J. Lipid Res.* **2021**, *62*, 100087. https://doi.org/10.1016/j.jlr.2021.100087.

(32) Mahadevi, A. S.; Sastry, G. N. Cation−π Interaction: Its Role and Relevance in Chemistry, Biology, and Material Science. *Chem. Rev.* **2013**, *113* (3), 2100–2138. https://doi.org/10.1021/cr300222d.

(33) Webb, B.; Sali, A. Comparative Protein Structure Modeling Using MODELLER. *Curr. Protoc. Bioinforma.* **2016**, *54*, 5.6.1-5.6.37. https://doi.org/10.1002/cpbi.3.

(34) Søndergaard, C. R.; Olsson, M. H. M.; Rostkowski, M.; Jensen, J. H. Improved Treatment of Ligands and Coupling Effects in Empirical Calculation and Rationalization of pKa Values. *J. Chem. Theory Comput.* **2011**, *7* (7), 2284–2295. https://doi.org/10.1021/ct200133y.

(35) Olsson, M. H. M.; Søndergaard, C. R.; Rostkowski, M.; Jensen, J. H. PROPKA3: Consistent Treatment of Internal and Surface Residues in Empirical pKa Predictions. *J. Chem. Theory Comput.* **2011**, *7* (2), 525–537. https://doi.org/10.1021/ct100578z.

(36) Salomon-Ferrer, R.; Case, D. A.; Walker, R. C. An Overview of the Amber Biomolecular Simulation Package. *WIREs Comput. Mol. Sci.* **2013**, *3* (2), 198–210. https://doi.org/10.1002/wcms.1121.

(37) Case, D. A.; Aktulga, H. M.; Belfon, K.; Cerutti, D. S.; Cisneros, G. A.; Cruzeiro, V. W. D.; Forouzesh, N.; Giese, T. J.; Götz, A. W.; Gohlke, H.; Izadi, S.; Kasavajhala, K.; Kaymak, M. C.; King, E.; Kurtzman, T.; Lee, T.-S.; Li, P.; Liu, J.; Luchko, T.; Luo, R.; Manathunga, M.; Machado, M. R.; Nguyen, H. M.; O'Hearn, K. A.; Onufriev, A. V.; Pan, F.; Pantano, S.; Qi, R.; Rahnamoun, A.; Risheh, A.; Schott-Verdugo, S.; Shajan, A.; Swails, J.; Wang, J.; Wei, H.; Wu, X.; Wu, Y.; Zhang, S.; Zhao, S.; Zhu, Q.; Cheatham, T. E.; Roe, D. R.; Roitberg, A.; Simmerling, C.; York, D. M.; Nagan, M. C.; Merz, K. M. AmberTools. *J. Chem. Inf. Model.* **2023**, *63* (20), 6183–6191. https://doi.org/10.1021/acs.jcim.3c01153.

(38) Brooks, B. R.; Brooks, C. L.; Mackerell, A. D.; Nilsson, L.; Petrella, R. J.; Roux, B.; Won, Y.; Archontis, G.; Bartels, C.; Boresch, S.; Caflisch, A.; Caves, L.; Cui, Q.; Dinner, A. R.; Feig, M.; Fischer, S.; Gao, J.; Hodoscek, M.; Im, W.; Kuczera, K.; Lazaridis, T.; Ma, J.; Ovchinnikov, V.; Paci, E.; Pastor, R. W.; Post, C. B.; Pu, J. Z.; Schaefer, M.; Tidor, B.; Venable, R. M.; Woodcock, H. L.; Wu, X.; Yang, W.; York, D. M.; Karplus, M. CHARMM: The Biomolecular Simulation Program. *J. Comput. Chem.* **2009**, *30* (10), 1545–1614. https://doi.org/10.1002/jcc.21287.

(39) Jo, S.; Kim, T.; Iyer, V. G.; Im, W. CHARMM-GUI: A Web-based Graphical User Interface for CHARMM. *J. Comput. Chem.* **2008**, *29* (11), 1859–1865. https://doi.org/10.1002/jcc.20945.

(40) Lee, J.; Cheng, X.; Swails, J. M.; Yeom, M. S.; Eastman, P. K.; Lemkul, J. A.; Wei, S.; Buckner, J.; Jeong, J. C.; Qi, Y.; Jo, S.; Pande, V. S.; Case, D. A.; Brooks, C. L.; MacKerell, A. D.; Klauda, J. B.; Im, W. CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J. Chem. Theory Comput.* **2016**, *12* (1), 405–413. https://doi.org/10.1021/acs.jctc.5b00935.

(41) Lomize, M. A.; Pogozheva, I. D.; Joo, H.; Mosberg, H. I.; Lomize, A. L. OPM Database and PPM Web Server: Resources for Positioning of Proteins in Membranes. *Nucleic Acids Res.* **2012**, *40* (D1), D370–D376. https://doi.org/10.1093/nar/gkr703.

(42) Maier, J. A.; Martinez, C.; Kasavajhala, K.; Wickstrom, L.; Hauser, K. E.; Simmerling, C. ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *J. Chem. Theory Comput.* **2015**, *11* (8), 3696–3713. https://doi.org/10.1021/acs.jctc.5b00255.

(43) Dickson, C. J.; Walker, R. C.; Gould, I. R. Lipid21: Complex Lipid Membrane Simulations with AMBER. *J. Chem. Theory Comput.* **2022**, *18* (3), 1726–1736. https://doi.org/10.1021/acs.jctc.1c01217.

(44) Price, D. J.; Brooks, C. L. A Modified TIP3P Water Potential for Simulation with Ewald Summation. *J. Chem. Phys.* **2004**, *121* (20), 10096–10103. https://doi.org/10.1063/1.1808117.

(45) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of Simple Potential Functions for Simulating Liquid Water. *J. Chem. Phys.* **1983**, *79* (2), 926–935. https://doi.org/10.1063/1.445869.

(46) MacKerell, A. D.; Bashford, D.; Bellott, M.; Dunbrack, R. L.; Evanseck, J. D.; Field, M. J.; Fischer, S.; Gao, J.; Guo, H.; Ha, S.; Joseph-McCarthy, D.; Kuchnir, L.; Kuczera, K.; Lau, F. T. K.; Mattos, C.; Michnick, S.; Ngo, T.; Nguyen, D. T.; Prodhom, B.; Reiher, W. E.; Roux, B.; Schlenkrich, M.; Smith, J. C.; Stote, R.; Straub, J.; Watanabe, M.; Wiórkiewicz-Kuczera, J.; Yin, D.; Karplus, M. All-Atom Empirical Potential for Molecular Modeling and Dynamics Studies of Proteins. *J. Phys. Chem. B* **1998**, *102* (18), 3586–3616. https://doi.org/10.1021/jp973084f.

(47) Galindo-Murillo, R.; Robertson, J. C.; Zgarbová, M.; Šponer, J.; Otyepka, M.; Jurečka, P.; Cheatham, T. E. Assessing the Current State of Amber Force Field Modifications for DNA. *J. Chem. Theory Comput.* **2016**, *12* (8), 4114–4127. https://doi.org/10.1021/acs.jctc.6b00186.

(48) Tian, C.; Kasavajhala, K.; Belfon, K. A. A.; Raguette, L.; Huang, H.; Migues, A. N.; Bickel, J.; Wang, Y.; Pincay, J.; Wu, Q.; Simmerling, C. ff19SB: Amino-Acid-Specific Protein Backbone Parameters Trained against Quantum Mechanics Energy Surfaces in Solution. *J. Chem. Theory Comput.* **2020**, *16* (1), 528–552. https://doi.org/10.1021/acs.jctc.9b00591.

(49) Izadi, S.; Anandakrishnan, R.; Onufriev, A. V. Building Water Models: A Different Approach. *J. Phys. Chem. Lett.* **2014**, *5* (21), 3863–3871. https://doi.org/10.1021/jz501780a.

(50) Joung, I. S.; Cheatham, T. E. Determination of Alkali and Halide Monovalent Ion Parameters for Use in Explicitly Solvated Biomolecular Simulations. *J. Phys. Chem. B* **2008**, *112* (30), 9020–9041. https://doi.org/10.1021/jp8001614.

(51) Darden, T.; York, D.; Pedersen, L. Particle Mesh Ewald: An $N \cdot \log(N)$ Method for Ewald Sums in Large Systems. *J. Chem. Phys.* **1993**, *98* (12), 10089–10092. https://doi.org/10.1063/1.464397.

(52) *Amber 2020 Reference Manual*.

(53) *Amber 2022 Reference Manual*.

(54) Berendsen, H. J. C.; Postma, J. P. M.; Van Gunsteren, W. F.; DiNola, A.; Haak, J. R. Molecular Dynamics with Coupling to an External Bath. *J. Chem. Phys.* **1984**, *81* (8), 3684–3690. https://doi.org/10.1063/1.448118.

(55) Davidchack, R. L.; Handel, R.; Tretyakov, M. V. Langevin Thermostat for Rigid Body Dynamics. *J. Chem. Phys.* **2009**, *130* (23), 234101. https://doi.org/10.1063/1.3149788.

(56) Wen, P.-C.; 溫博超; Verhalen, B.; Wilkens, S.; Mchaourab, H. S.; Tajkhorshid, E. On the Origin of Large Flexibility of P-Glycoprotein in the Inward-Facing State. *J. Biol. Chem.* **2013**, *288* (26), 19211–19220. https://doi.org/10.1074/jbc.M113.450114.

(57) Roe, D. R.; Cheatham, T. E. PTRAJ and CPPTRAJ: Software for Processing and Analysis of Molecular Dynamics Trajectory Data. *J. Chem. Theory Comput.* **2013**, *9* (7), 3084–3095. https://doi.org/10.1021/ct400341p.

(58) Gowers, R.; Linke, M.; Barnoud, J.; Reddy, T.; Melo, M.; Seyler, S.; Domański, J.; Dotson, D.; Buchoux, S.; Kenney, I.; Beckstein, O. MDAnalysis: A Python Package for the Rapid Analysis of Molecular Dynamics Simulations; Austin, Texas, 2016; pp 98–105. https://doi.org/10.25080/Majora-629e541a-00e.

(59) Michaud-Agrawal, N.; Denning, E. J.; Woolf, T. B.; Beckstein, O. MDAnalysis: A Toolkit for the Analysis of Molecular Dynamics Simulations. *J. Comput. Chem.* **2011**, *32* (10), 2319–2327. https://doi.org/10.1002/jcc.21787.

(60) Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Comput. Sci. Eng.* **2007**, *9* (3), 90–95. https://doi.org/10.1109/MCSE.2007.55.

(61) Humphrey, W.; Dalke, A.; Schulten, K. VMD: Visual Molecular Dynamics. *J. Mol. Graph.* **1996**, *14* (1), 33–38. https://doi.org/10.1016/0263-7855(96)00018-5.

Figure Captions

Figure 1. Projection of ABCB4 MD simulations onto the ABC conformational space. (A) Definitions of the ABC structural parameters, TMH1-6 and NBD1 are depicted in green and TMH7-12 and NBD2 are depicted in blue. Conformation- and bound state-dependent distributions of (B) IC angle *versus* NBD distance and (C) NBD twist *versus* NBD distance from ABCB4 MD simulations on the ABC conformational space defined by resolved cryo-EM structures in symmetric and asymmetric models.

Figure 2. Local conformational sampling of ABCB4 MD simulations. InfleCS free energy-based conformational clustering of ABCB4 under different conformations and bound states in symmetric and asymmetric membrane models according to (A) IC angle versus NBD distance and (B) NBD twist versus NBD distance. (C) Superimposition of representative snapshots for ABCB4 main populations in symmetric (blue) and asymmetric (red) lipid models. NBD1 and NBD2 are respectively depicted in blue and cyan for symmetric lipid bilayer model and red and pink for asymmetric lipid bilayer model.

Figure 3. Lipid bilayer properties and cholesterol-protein interplay. (A) Conformation- and bound state-dependent order parameters of palmitate and oleate lipid tails of outer and inner leaflet in symmetric (solid lines) and asymmetric models (dashed line). Per-residue atomic contact fractions between cholesterol and (B) apo ^{IF}ABCB4 or (D) ^{CC}ABCB4-(ATP)$_2$ and calculated cholesterol hotspots for (C) apo ^{IF}ABCB4 or (E) ^{CC}ABCB4-(ATP)$_2$.

Figure 4. Lipid-protein non-covalent interactions from ABCB4 MD simulations. Conformation and bound state-dependent (A) polar head hotspots and (B) average lipid-protein H-bond counts from MD simulations. (C) Per-domain calculated H-bond fractions over MD simulations for apo ^{IF}ABCB4, ^{IF}ABCB4-PC-(ATP)$_2$ and ^{CC}ABCB4-(ATP)$_2$.

Figure 5. Comparison of TMH arrangement and helicity of ABCB4 conformations and bound states in symmetric and asymmetric membrane. (A) Time-dependent root-mean-squared deviations of ABCB4 front and back gates for all ABCB4 conformations and substrate

bound states, using initial frame of apo ^{IF}ABCB4 as reference. Superimposition of representative snapshots of (B) front and (C) back gates of ABCB4 conformations and bound states in asymmetric model. (C) Conformation- and bound state-dependent TMH4 and TMH10 kinking over ATP and PC binding event along canonical alternating access.

Figure 6. PC substrate binding in canonical and credit-card swipe alternating access. (A) Representative snapshots of ^{IF}ABCB4-PC and ^{IF}ABCB4-PC-(ATP)$_2$ in asymmetric model highlighting identified residues involved in cation-π cage. (B) Definition of geometrical parameters, namely PC polar head orientation and distance to membrane center used to assess (C) the local free energy surface of substrate binding event of canonical alternating access. (D) Mapping of potential polar amino acid involved in credit-card swipe mechanism including key average distances and (E) residue position along membrane normal from MD simulations.

Figure 7. Proposed overall dual mechanism for PC translocation

A

^{IF}**ABCB4-PC**

Gln725
Phe305
His989
Phe345
Trp234

^{IF}**ABCB4-PC-(ATP)$_2$**

Gln725
Phe305
Phe345
His989
Trp234

B

z
PC Substrate
P-atom
PC substrate
N-atom

C

^{IF}**ABCB4-PC**

C1(44.2%)
C2(36.1%)
C3(13.5%)
C4(6.2%)
Cryo EM (7NIV)

PC orientation (deg)
PC distance to membrane center (Å)
kcal/mol

^{IF}**ABCB4-PC-(ATP)$_2$**

C1(31.0%)
C2(29.7%)
C3(16.9%)
C4(15.0%)
C5(5.0%)
C6(2.5%)
Cryo EM (7NIV)

PC orientation (deg)
PC distance to membrane center (Å)
kcal/mol

D

TMH1
TMH3
Asp83
Glu80
Arg212
Ser69
Ser58
Lys54
Gln52

16.4 ± 0.5 Å
16.5 ± 0.3 Å
10.8 ± 0.4 Å

E

Distance from membrane center (Å)
Density (A.U.)

Gln52
Lys54
Ser58
Ser69
Arg212
Glu80
Asp83