Intrinsic Diversity in Primary Cilia Revealed Through Spatial Proteomics

Jan N. Hansen [1,2], Huangqingbo Sun [2,*], Konstantin Kahnert [2,*], Eini Westenius [3,4], Alexandra Johannesson [1], Kalliopi Tzavlaki [1], Casper Winsnes [1], Emmie Pohjanen [1], Jenny Fall [1], Frederic Ballllosera Navarro [2], Anna Bäckström [1,2], Cecilia Lindskog [5], Fredric Johansson [1], Kalle von Feilitzen [1], Angelica Delado Vega [3,4] Mathias Uhlén [1,6], Ana Martinez Casals [2], Diana Mahdessian [1], Anna Lindstrand [3,4], Ulrika Axelsson [1], Emma Lundberg [†,1,2,7,8]

[1] Science for Life Laboratory, School of Engineering Sciences in Chemistry, Biotechnology and Health, KTH - Royal Institute of Technology, Stockholm, Sweden.
[2] Department of Bioengineering, Stanford University, Stanford, CA, USA.
[3] Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden
[4] Department of Clinical Genetics and Genomics, Karolinska University Hospital, Stockholm, Sweden
[5] Department of Immunology, Genetics and Pathology, Cancer Precision Medicine Research Unit, Uppsala University, SE-75185 Uppsala, Sweden
[6] Department of Neuroscience, Karolinska Institutet, Stockholm, Sweden.
[7] Department of Pathology, Stanford University, Stanford, CA, USA.
[8] Chan Zuckerberg Biohub, San Francisco, San Francisco, CA, USA.
* Contributed equally
†Correspondence: emmalu@stanford.edu

Abstract

Primary cilia are solitary filamentous protrusions on most human cells. Our understanding of their function is informed by the study of ciliopathies, hereditary diseases linked to cilia dysfunction. These disorders present a wide array of symptoms, highlighting the tissue-specific roles of cilia. However, cell-state and cell-type specific functions remain poorly understood due to limited tools for analyzing individual ciliary protein composition. Here, we utilize antibody-based spatial proteomics to expand the Human Protein Atlas with a dedicated section for primary cilia. We identify the subciliary locations of 654 proteins across three cell lines, analyzing 218,412 individual cilia. Our findings reveal that 67% of the ciliary proteins are cell-type specific and that 78% of ciliary proteins show heterogeneous localization patterns within cell populations, suggesting that the functional tuning of individual cilia is substantially driven by protein trafficking rather than gene expression. Our atlas portrays the cilium as a dynamic sensory organelle, suggesting that individual cells can fine-tune their ciliary structure - both in terms of receptors and signaling components - to sense and integrate diverse signals, computing a response to relay to the cell body. We identify 71 novel cilia proteins, establish CFAP300 as a specific and broad marker for cilia, and suggest multiple potential ciliopathy genes including CREB3, where we show that a coding genetic variant can be linked to blindness and skeletal abnormalities. In summary, our spatially resolved cilia atlas paves new avenues for basic and clinical research on cilia and ciliopathies.

Introduction

Cilia are small, hair-like cellular appendages that exist on almost any vertebrate cell. They are classified into motile cilia and immotile, primary cilia. While motile cilia move entire cells or fluid flow along epithelial surfaces, primary cilia fulfill an antenna function [1]. Most of what we know about the function of primary cilia stems from hereditary diseases that are caused by primary cilia dysfunction, so-called ciliopathies [2]. Ciliopathy patients present with a broad spectrum of phenotypes, including polydactyly, cognitive impairments, obesity, kidney defects, or blindness, underlining that primary cilia fulfill pleiotropic and tissue-specific functions [2]. However, the molecular mechanisms underlying primary cilia function are largely unknown.

Functionally, primary cilia represent an organelle optimized for signal transduction: they sense extracellular cues via receptors, locally transduce the signal, and relay it to the cell body [3]. The primary cilium reportedly participates in several signaling pathways: individual components of Hedgehog signaling, G-protein-coupled receptor (GPCR) signaling, receptor-tyrosine-kinase (RTK) signaling as well as transforming growth factor β (TGF-β) signaling, bone morphogenetic protein (BMP) signaling, and Wnt signaling have been observed to localize to the primary cilium [1]. However, the picture of ciliary signaling components is still elusive. While the molecular sequence of events has been mostly understood only for Hedgehog signaling [1], knowledge about how the other components engage mechanistically in signaling pathways and whether and how the cilium can distinguish between signals of different receptors targeting the same signaling mechanism (e.g., GPCRs) remains elusive [4].

The architecture of the cilium and the process of its construction, called ciliogenesis, has been intensively studied. Two ciliogenesis pathways have been described, the internal and the external pathway [5]. They are initiated by the mother centriole docking to the plasma membrane or to a vesicular membrane before the cilium is shaped by the axoneme extending from the mother centriole as a template. The axoneme is the central skeleton structure of the cilium consisting of nine microtubule doublets nucleated at the centriole. Beyond providing the cilium with a skeleton and thereby determining its length, the axoneme serves as a template for protein trafficking in the cilium, which is driven by the intraflagellar transport (IFT) machinery [1]. Early during ciliogenesis, several protein structures are built around the mother centriole, which let it become the "basal body", and at the interface of cilium and cell body, which is called the "transition zone" [5]. For example, transition fibers are formed to mount the basal body to the plasma membrane [6–9]. The transition zone limits exchange of lipids and proteins between cilium and the cell body, which macro-structurally appear as a continuum [10]. The transition zone consists of several protein structures, such as a Septin-based diffusion barrier at the ciliary membrane [11], Y-shaped protein complexes connecting plasma membrane and ciliary axoneme [12], and the distal appendage matrix sealing the space between transition fibers [13]. Yet, the composition and mechanism of the gate between cilium and the cell body remains poorly understood.

Nonetheless, it has been shown that the protein and lipid repertoire of the primary cilium is distinct from the cell body, which is achieved by controlled entry and exit of proteins [14,15]. This tightly controlled shuttling of proteins into and out of the primary cilium fulfills multiple functions:

(1) Enabling signal transduction, e.g., during activation of Hedgehog signaling a competitive spatial exchange of proteins triggers the signaling cascade [16]. (2) Relaying information to the cell body to regulate cellular responses, e.g., in Hedgehog signaling, transcription factors are modified in the primary cilium, shuttle to the nucleus, and instruct a specific gene expression program [16]. (3) Tuning the sensitivity of the ciliary antenna function, e.g., activated GPCRs in the primary cilium are removed from the ciliary membrane and transported out of the cilium [17,18], and the ciliary length determines its sensitivity in mechanosensation [19]. (4) Assembling and disassembling the primary cilium [5]. In conclusion, understanding the ciliary protein repertoire and the underlying spatiotemporal dynamics appears as a key to understanding the antenna function and signaling pathways employed by the cilium, and how it's influenced by and shaping cell states [1].

A major limitation for primary cilia research is imposed by the size of the primary cilium. Its volume can be approximated to be in the range of 0.3 fl, which is more than 1,000-times smaller than the cell body, thus requiring highly sensitive measurement technologies. In addition, extracting and purifying intact cilia is difficult, if not impossible, because the ciliary membrane is a continuum of the plasma membrane [5]. Thus, the ciliary proteome has mainly been characterized using proximity-labeling-based spatial proteomics techniques, such as APEX-, BioID-, or Turbo-ID-based proximity labeling [20–24]. However, the use of proximity-labeling MS features several disadvantages: (1) it does not provide single cell resolution and results represent a mix of cilia in multiple signaling states; (2) technologically, it can only reveal ciliary protein enrichment compared to the cell body; (3) it is not functional in all cell types due to endogenous enzymatic activity; (4) it relies on fusing an enzyme, such as APEX or TurboID, to a ciliary GPCR or ciliary protein fragment, which may bias signaling in the primary cilium; and (5) it is not applicable to study human tissues.

Here, we overcome this challenge and shed light on the physiological function of individual primary cilia on human cells by applying large-scale antibody- and imaging-based spatial proteomics. We establish a new resource for the cilia field: a section in the public Human Protein Atlas (HPA) that provides high-resolution confocal 3D images and subciliary annotations for 654 cilium-related proteins in three ciliated cells of diverse origin. We reveal that the primary cilium shows the most heterogeneous proteome compared to all other organelles, across cell types and within the same cell population, and that the diversity relates to cell-individual customization of the cilium to conduct cell-specific sensory signaling. Furthermore, we demonstrate that our cilia atlas paves new avenues for cilia and ciliopathy research, by leveraging its content for revealing a new cilia marker and identifying a potential novel ciliopathy gene.

Results

Mapping the subcellular and subciliary location for human proteins

To examine the proteomic composition of primary cilia on the single-cilium level, we developed an antibody-based spatial proteomics workflow, employing the antibody library of the Human Protein Atlas (HPA) [25,26] in a co-staining approach with markers for nuclei, cilia, and basal bodies (Fig. 1A) and high-throughput, high-resolution 3D confocal microscopy (Fig. 1B). With this workflow we comparatively analyzed three immortalized primary human cell lines of different tissue origin and sex: (1) starved hTERT-RPE1 cells from the embryonic retina [27], to represent the best studied *in vitro* model for human cilia, (2) RPTEC/TERT1 cells [28], derived from the epithelial kidney proximal tubules, representing a cell type affected in the most common ciliopathy - polycystic kidney disease (PKD), and (3) the mesenchymal, multipotent, stromal ASC52telo cells [29], derived from adipose tissue, to represent a multipotent cell type. The three cell lines are morphologically diverse and likely rely on different signaling and ciliogenesis programs, allowing coverage of a broad band of cellular functionality and ciliary diversity (Fig 1C). The starved hTERT-RPE1 cells represent a synchronized culture, with a high ciliation rate (Fig 1D). While the RPTEC/TERT1 cell culture cycles slowly and provides cilia with a broad length range, the ASC52telo cells cycle fast, presenting short cilia (FIG 1D). hTERT-RPE1 cells form their cilia mostly at the cell-glass-interface *in vitro* and reportedly, predominantly employ the internal ciliogenesis pathway [5]. RPTEC-TERT1 cells are highly polarized and form cilia that can become very long and that mostly stick out from the top of the cells into the fluid (FIG 1C,D). We observed ASC52telo cells to occasionally show dots or small lines of the ciliary marker on one centrosome during mitosis, hinting to the retainment of a ciliary vesicle or cilium during mitosis, similar to reported observations in HEK293T cells [30] (FIG 1E). This was not observed in the other cell lines. However, in RPTEC TERT1 cells, we occasionally observed cells that formed two adjacent cilia (Fig. 1F). Double ciliation was also seen in RPE1 cells, albeit much rarer (Fig. 1G). Interestingly, we also once observed a ciliary bridge in RPTEC/TERT1 cells, where cilia from neighboring cells appeared to be either connected or touching (Fig. 1H).

For the targeted antibody-based spatial proteomics approach, we assembled a list of candidate proteins based on existing curated databases including SYSCILIA gold standard version 2 [31] and the ChlamyFP.org database [15], the tissue resource of the HPA [25,32], MS data sets on primary cilia based on proximity-labeling approaches or purification attempts [20–24,33], predictive studies including the CiliaCarta [34–36], comparative transcriptomics studies [37], phylogenetic studies on flagellated species [38,39], CRISPR screens [40,41], MS studies of flagella or motile cilia [42–46], Cryo-ET studies of sperm flagella [47–49], and studies of the centrosomal proteome [50–55]. For a strong focus on signaling pathways, we extended the screen with key components of diverse signaling pathways, such as AKT kinases, MAP kinases, mTOR, a selection of G-protein coupled receptors (GPCRs), phosphodiesterases (PDEs), or adenylyl cyclases, even without any prior evidence of localizing to cilia. Interestingly, our assembled cilia candidate list highlighted that nearly ten thousand protein-coding genes have been reported or suggested to potentially produce proteins localizing to primary cilia or engaging in their regulation (Fig. 1I, Supplementary Table 1). The overlap of proteins identified in cilia with different methods or in different cell types or species was

particularly small: 4122, 1905, and 1011 genes were attributed to cilia in one, two, or three data sets only, respectively (Fig. 1I). This points to an incredible heterogeneity in the ciliary proteome across species and cell types and potentially many false positives due to technical limitations. Finally, we prioritized the candidate list and selected 1,935 candidate proteins for assessment of ciliary localization in human cells (see Materials and Methods) (Fig. 1J, Supplementary Table 2).

A spatial map of the human cilia proteome

Applying our spatial proteomics data generation pipeline for the 1,935 proteins produced a massive high-resolution 3D confocal image data set of cilia and their protein composition with 17,092 3D image stacks, revealing 463,368 cells, 218,412 of which displayed a cilium. To annotate the 17,092 3D image stacks, we applied the established HPA subcellular workflow, which involves investigation and review of every staining by multiple skilled cell biologists.[26] Beyond the classical 35 subcellular organelles and structures annotated in the HPA, we also annotated protein localization to the primary cilium, basal body, transition zone, and ciliary tip (Fig. 2A,B). We could identify the proteins encoded by 654 of our 1,935 candidate genes at cilia, of which 352 proteins were detected in primary cilia, 59 at the ciliary tip, 95 at the transition zone region, and 403 at the basal bodies (Fig. 2C). Furthermore, we observed additional subpatterns of ciliary localization, such as partial staining of the proximal cilium (e.g., for EFCAB7, EVI2A, and TRIM46, Fig. 2D), which might correspond to a domain distal from the transition zone and close to the "inversin" compartment [56] or a strong staining of a ciliary pocket; spotty staining patterns along the cilium (e.g., for TRIO, Fig. 2D); or enrichment towards the ciliary tip (e.g., for C2CD4C, Fig. 2D); or combinations of spotty patterns and enrichment towards the tip for the same protein (e.g., for TMEM237, Fig. 2D), highlighting the subciliary resolution of our screen.

Among the studied proteins are many established ciliary proteins, such as the intraflagellar transport protein IFT22, SMAD7 localizing to the basal body, the subdistal appendage protein CEP164, the MKS-complex protein TCTN2 at the transition zone, the small GTPase RAB7A in the transition zone, the ciliary trafficking protein ARL6 enriched towards the ciliary tip, the kinesin KIF7 at the ciliary tip, the septin SEPTIN2, the adenylyl cyclase ADCY3, or the small GTPases RAB23, which is known to participate in regulating GLI transcription factors and ciliary Hedgehog signaling (Fig. 2E). [57,58] This confirms the validity of our approach for diverse types of ciliary proteins and provides the cilia field with subcellular resolution pictures of these and many more known proteins in unperturbed human cells. In total, we identified as much as 71 proteins that had never been previously associated with any type of cilium across different species (Fig. 2F, Supplementary Table 3). Additionally, there were 193 proteins that had never been identified to localize to human cilia by MS- or Cryo-ET based approaches (Fig. 2G), and 96 proteins that had never been validated to localize to cilia (Fig. 2H). Notably, for those proteins that were identified only by MS-based approaches, the subciliary localization was not known, whereas we provide subciliary localization resolution.

An interesting new discovery was that we detected three proteasomal subunits in cilia: PSMD10, PSME3, and PSMA6 (Fig. 2I). PSMD10 is among the 71 proteins never related to cilia before. PSME3 has previously been detected in one mouse sperm flagella data set [47] and was bioinformatically predicted to be a cilia protein [34]. PSMA6 was identified in cilia by many

approaches and was listed in databases [15], an analysis of murine kidney cilia with cilia-APEX [22], a CRISPR-based screen [41], a comparative transcriptomic analysis [37], and in motile cilia and sperm flagella studies [42,45,47–49]. Other examples for our data confirming proximity-labeling-MS-approaches in the cilia of model organisms are ZNF330 or the senescence-regulator RSL1D1 (Fig. 2J), which both had previously been identified by cilia-APEX in murine kidney cells [21,22]. Also, we revealed previous predictions in cilia, such as ARL15 (Fig. 2J) [36]. Some of the studies in our candidate list analysis were from motile cilia. Surprisingly, we identified proteins believed to be specific for motile cilia in primary cilia of RPTEC/TERT1 cells, such as the central-pair protein HYDIN, IQCA1, which is part of the nexin-dynein-regulatory complex in motile cilia, or the CATSPER2 channel subunit (Fig. 2K). This is particularly interesting in light of recent descriptions of motile cilia proteins and reportedly active motility of primary cilia in *ex vivo* pancreatic islet cells [59]. Yet, we see that these proteins accumulate towards the tip rather than show staining along the whole cilium, which might indicate that, for HYDIN and IQCA1, they are not in fact implemented into the axoneme but might just transit the cilium. The same applies to CATSPER2, which is expressed in RPTEC/TERT1 cells (according to the HPA cell line sequencing data). Based on our observation we do not know whether CATSPER2 in fact implemented into the plasma membrane or transiting the cilium. Further studies will need to delineate why and to which purpose motile cilia proteins transit RPTEC/TERT1 cilia and whether this observation also holds true *in vivo* in cilia of the human kidney's proximal tubules.

The primary cilium is a hub for diverse sensory signaling pathways

To explore the functions of the human cilia proteins, we performed an enrichment analysis for Gene Ontology (GO) Biological Process terms (Supplementary Fig. 1, Supplementary Table 4). Beyond many terms relating to maintenance processes of the cilium or basal body (e.g. intraciliary transport, adj. P=1.1906e-27 and cilium organization, adj. P=4.7218e-94), we found significant enrichment of terms relating to signaling, such as calcium signaling (calcium-mediated signaling, adj. P=0.0008), Wnt signaling (Wnt signaling pathway, adj. P=0.0006), cAMP signaling (cAMP-mediated signaling, adj. P=0.0028), and morphogenesis and developmental signaling pathways (e.g. regulation of developmental growth adj. P=0.0054) (Fig. 3A, Supplementary Table 4). In fact, we detected signaling proteins of most major signaling pathways in the primary cilium (Table 1) of which many had hitherto not been confirmed in cilia. For example, we revealed in cilia the kinases AKT1 and AKT3, the Serine/threonine-protein kinase TAO1 (TAOK1), a catalytic and regulatory subunit of PI3K (PIK3CA and PIK3R4), the focal adhesion kinase PTK2, the calcium/calmodulin dependent protein kinase CAMK1G, the protein phosphatase 2 (PPP2R3C), the E3 ubiquitin-protein ligase MIB1, the vascular endothelial growth factor receptor 3 FLT4, the GPCR OPRL1, the src-related receptor PDGFRA, the transforming growth factor beta receptor 1 TGFBR1, and the cyclic nucleotide gated ion channel CNGA2 (Fig. 3B). Furthermore, the Triple functional domain protein TRIO mentioned above (Fig. 2D) is a Guanine nucleotide exchange factor (GEF) for RHOA and RAC1 GTPases [60], In conclusion, these results demonstrate that the cilium is a versatile signaling organelle with broad signaling capacities, filled with most receptor types and diverse downstream signaling machinery. Moreover, the fact that we do not only find receptors but also downstream signaling components of all large signaling pathways and central signaling hubs such as AKT kinases emphasizes that the cilium is not just an antenna but

processes signals from receptors and computes a response that it ultimately relays to the cell body.

Table 1 Signaling proteins revealed in cilia or basal bodies

Gene names part of the selected GO terms are shown if they were revealed at the Primary Cilium Tip, Primary Cilium, Primary Cilium transition zone (TZ), or the Basal Body.

Signaling pathway (Gene ontology = GO term)	Primary Cilium Tip	Primary Cilium	Primary Cilium TZ	Basal Body
G-protein coupled receptor signaling pathway (GO:0007186)	CAMK2A, ADCY2, OPRL1	ATP2B4, CAMK2A, PRKACA, ACTN2, TULP3, ADCY2, PDE4D, GPR83, OPRL1, ADCY3, LGR5, AKT1, PRKACB, GPR161, HTR7, GPR180, ADCY9, ADORA1, FZD6, ADRB2, PTAFR, ROBO1, GPRC5C, INSR, LPAR3, MRGPRF, ADCY5, FZD4, SPHK1, FZD8, CHRM2, PDE4B, SSTR3	CAMK2A, ACTN2, TULP3, ADCY3, LGR5, GPR161, GPRC5C, INSR, MRGPRF, SPHK1	ATP2B4, GNAI3, PRKACA, TULP3, RAPGEF4, RPGRIP1L, NHERF1, GNAI2, TACR1, PRKD3, GPR83, GNAI1, PLCB2, ADCY3, LGR5, AKT1, PRKACB, ADCY9, ADGRV1, FZD6, TUB, ADRB2, TAPT1, INSR, FZD2, CHRM2, FFAR4, ADRB3, GABBR1, LGR4, GNAT3
cAMP-mediated signaling (GO:0019933)	ADCY2, OPRL1	ADCY2, PDE4D, OPRL1, PDE11A, PDE4B		PDE4C, CREB1, GNAI1, PDE11A, PDE6B, PDE5A, FFAR4, PDE4C
Calcium-mediated signaling (GO:0019722)	OPRL1, PDPK1, CAMK2D, GSTM2	ATP2B4, PRKACA, FKBP1B, OPRL1, PRKAA1, PDGFRA, PDPK1, CAMK2D, CACNA1C, RCAN2, ATP2A2, SPHK1, PPP3R2, GSTM2, PPP3R1, ACTN3	CAMK2D, SPHK1, GSTM2, ACTN3	ATP2B4, PRKACA, FKBP1B, PLCG1, PRKAA1, CAMTA1, LRRK2
Receptor tyrosine kinase signaling pathway (GO:0007169)	WDR54, NR1H4, CHRNA3, MAPK1, MAPK3, PDPK1, PTK2, PIK3CD	WDR54, NR1H4, FLT4, RAB7A, FGFR1, CHRNA3, MAPK1, MAPK3, IFT20, WNT5A, PIK3CA, DNAI1, BMP2, PRKAA1, PDGFRA, AHI1, MAP2K5, IGF1R, PDPK1, AKT1, EGFR, BRAF, CTNNB1, ADRB2, PTK2, ROBO1, INSR, PIK3CD, FZD4, GRB2	RAB7A, CHRNA3, BMP2, INSR	PRKCZ, FGFR1, CHRNA3, MAPK1, MAPK3, TGFB1, NHERF1, IFT20, GNAI2, CCDC88A, TEK, PLCG1, PRKAA1, MAP2K5, PRICKLE1, AKT1, RASA1, EGFR, BRAF, MAPKAPK2, CRK, CTNNB1, MAP2K1, ADRB2, INSR, HRAS
Transforming growth factor beta receptor signaling pathway (GO:0007179)	TGFBR1, PDPK1, SMAD4, PTK2	FKBP1A, AXIN1, TGFBR1, STK11, BMP2, SMAD6, PDPK1, SMAD4, SMAD3, PTK2, SMAD2, PPARA	BMP2, SMAD3, SMAD2	FKBP1A, SMAD7, TGFB1, SMAD6, SMAD4, SMAD3, PPARA
Vascular endothelial growth factor receptor signaling pathway (GO:0048010)	PTK2	FLT4, PTK2, ROBO1, FZD4		MAPKAPK2

MAPK cascade (GO:0000165)	WDR54, MAPK1, PSMD10, MAPK3, SMAD4	WDR54, MAP4K5, FLT4, MAPK6, FGFR1, CYLD, IGBP1, MAPK1, MAP1LC3A, PSMD10, MAPK3, AXIN1, WNT5A, BMP2, PDGFRA, MAP2K5, IGF1R, SMAD4, EGFR, BRAF, TAOK1, ADORA1, SMAD3, CDC42EP5, CTNNB1, ULK4, ADRB2, ROBO1, PKHD1, INSR, LPAR3, SPHK1, GRB2, AIDA, PJA2	CYLD, BMP2, SMAD3, ULK4, INSR, SPHK1, PJA2	PRKCZ, DLG1, FGFR1, TP73, CYLD, IGBP1, MAPK1, MAPK3, TGFB1, NHERF1, GNAI2, TEK, MAP2K5, PELI2, SMAD4, EGFR, BRAF, DVL3, ATF3, MAPKAPK2, SMAD3, CRK, CTNNB1, MAP2K1, ADRB2, PKHD1, INSR, HRAS, AIDA, FFAR4, ADRB3, LRRK2, PJA2
TOR signaling pathway (GO:0031929)	MAPK3, VHL	PRKACA, MAPK3, AKT3, STK11, MAPKAP1, PIK3CA, PRKAA1, VHL, AKT1, PRKACB, TSC1, PKHD1	TSC1	PRKACA, MAPK3, USP4, CCDC88A, AKT3, PRKAA1, AKT1, PRKACB, PRKAA2, TSC1, PKHD1
BMP signaling pathway (GO:0030509)	MAPK3, SMAD4	MAPK3, WNT5A, BMP2, SMAD6, SMAD4, SMAD2	BMP2, SMAD2	SMAD7, MAPK3, SMAD6, SMAD4
Wnt signaling pathway (GO:0016055)	TTC21B, SMAD4, VCP	CYLD, AXIN1, IFT20, ARL6, NPHP3, WNT5A, STK11, INVS, TTC21B, BMP2, PRKAA1, LGR5, CSNK1D, SMAD4, EGFR, TMEM237, GRHL3, FZD6, VCP, SMAD3, DAPK3, CTNNB1, FZD4, FZD8	CYLD, BMP2, LGR5, TMEM237, VCP, SMAD3	MKS1, CYLD, TGFB1, NHERF1, IFT20, GLI1, INVS, TTC21B, MACF1, PRKAA1, PRICKLE1, LGR5, SMAD4, EGFR, DVL3, PRKAA2, FZD6, VCP, SMAD3, CTNNB1, RUVBL1, FZD2, RUVBL2, LRRK2, CTNND1, LGR4
Notch signaling pathway (GO:0007219)	NR1H4	NR1H4, MIB1, CDKN1B, BMP2, AKT1, PRKCI, ROBO1	CDKN1B, BMP2	IFT74, MIB1, CDKN1B, AKT1
Rho protein signal transduction (GO:0007266)		RHOU, CDC42EP5, ROBO1, ARHGAP1, SEMA4D	SEMA4D	TEK, SHTN1, SEMA4D, LRRK2

Subcellular protein multilocalization and the implications for ciliary function

Protein localization is in general closely associated with protein function. We and others have previously shown that more than half of the human proteome are localized to multiple subcellular compartments, i.e. are multilocalizing [26,61,62]. Here we show that cilium-localizing proteins (including tip and transition zone) are significantly more multilocalizing than other proteins (P=3.7932e-202) - as much as 99% of cilium-localizing proteins are also found in other cellular compartments (Supplementary Fig. 2). Looking at individual stainings, we observed for 90% of the stainings of a protein localizing to the cilium that the protein was multilocalizing to another subcellular compartment in the same cell culture, the most common other compartments being basal bodies, cytosol, nucleoplasm, and microtubules (Fig. 3C). In light of the basal body being the access point to the cilium [10], it makes sense that we observed many proteins also at the basal bodies, since these are transiting the basal body during transport to cilia. Multilocalization with microtubules may indicate that those proteins are part of the ciliary axoneme, a microtubule-based structure.

In contrast, proteins localizing at cilia and in the cytosol and nucleoplasm may rather be proteins involved in relaying information from the cilium to the cell body or potentially engaging in similar or different functions or signaling pathways in different places in the cell (a process called "moonlighting" [63]. In fact we identified kinases that are central in many signaling pathways in cilia, but also in cell body compartments, i.e., the cytosol, such as AKT1 and AKT3, PIK3CA and PIK3R4, or PTK2 (Fig. 3B). Their ability to localize to multiple locations may be attributed to their capacity to transduce distinct signals in various cellular compartments, influenced by the diverse environments with varying stimulators and substrates. We also observed proteins with different ciliary and cell body functions in multiple locations in the cell. For example, the protein TULP3 is well known for a role as a cargo-adaptor in IFT-A mediated trafficking of GPCRs and ARL13B to the cilium [64–66], but may also act as a transcription factor residing at the plasma membrane and relocalizing to the nucleus upon activation [67]. In agreement, we detected TULP3 in cilia, at the plasma membrane, and in the nucleus (Fig. 3B). Many multilocalization observations of cilia proteins may also relate to the cilium relaying information to the cell body. For example, for Gli transcription factors in Hedgehog signaling it is established that post-translational modification of the transcription factors occurs in the cilium, upon which they are transported into the cytosol, where they are - depending on the modification - either degraded or sent to the nucleus to adjust gene transcription [1]. We in fact found additional transcription factors at cilia, such as CRX in cilia (Fig. 3D), or ATF3 and CREB1 at the basal body (Fig. 3D), which co-localized to the nucleoplasm, supporting the hypothesis that the cilium controls gene expression by local modification of transcription factors beyond Sonic Hedgehog signaling.

Another class of proteins that often multilocalize and "moonlight" are enzymes, as we recently demonstrated [68]. Little is known about metabolic and enzymatic activities in the cilium. Notably, not only the proteomic but also the lipid composition of the cilium uniquely makes up the ciliary compartment [69]. While the lipid composition of cilia has been shown to be unique compared to the plasma membrane, the processes involved in its formation - whether through transport mechanisms or local production - remain underexplored. While for phosphoinositides it has been demonstrated that the membrane composition is locally regulated by enzymes, for other lipid species it is not understood how they end up in the ciliary membrane, such as for cholesterol, which is described to essentially regulate Hedgehog signaling [69,70]. Interestingly, we discovered several enzymes in cilia, which also localize to other positions in the cell. For example, an enzyme that is part of sterol synthesis, HSD3B1/HSD3B2 (Fig. 3E). Other examples are the Glutathione S-transferase LANCL1, the Inositol polyphosphate 5-phosphatase OCRL, the carboxypeptidase AGBL5 involved in tubulin de-glutamylation and has been indicated as a negative regulator of ciliogenesis [71], the Sphingosine kinase SPHK1, which regulates ceramide levels and is highly engaged in signaling [72–74], or the UDP-glucose glycoprotein glucosyltransferase 1 (UGGT1), a protein that reportedly acts as a gate-keeper for folding of glycosylated proteins at the endoplasmic reticulum (ER) (Fig. 3E) [75,76].

Our data clearly demonstrates that the cilia mainly contain proteins that are also present elsewhere in the cell, and that the cilium has in fact a local metabolic machinery, supporting the hypothesis that the ciliary lipid composition is shaped and adjusted locally through enzymatic

activity. Notably, finding UGGT1 in cilia also supports that the cilium might even be able to locally repair proteins or refold proteins, which might be required after import with chaperons into the cilium. The regulation of such enzymes in cilia may likely be different from the cell body to create the special ciliary environment, and this environment is likely locally fine-tuned, since the ciliary lipid composition is reportedly tightly intertwined with ciliary signaling [69].

Cilia proteomes reveal cell type specific functional differences

Most of the investigated proteins (67%) localized to cilia specifically only in one or two of the studied cell lines (Fig. 4A). In the cell lines where we did not observe ciliary localization, the protein was either not detectable or localized to other compartments. For example, we identified the calmodulin- and kinase-related signaling protein STRN3 at the plasma membrane / actin cytoskeleton in all three cell lines, but only in RPTEC/TERT1 cells in the cilium and we identified the common kinase PI3K (PIK3CA) in cilia in RPTEC/TERT1 and RPE1 cells but not in ASC52telo cells (Fig 4B). Other examples are the activin receptor ACVR1B, which localizes to cilia in RPE1 cells but not in RPTEC/TERT1 or ASC52telo cells, where it resides in vesicles, the SEPTIN9, which localized to cilia in RPE1 and RPTEC/TERT1 cells, but not in ASC52telo cells, or the RHO GTPase activating protein ARHGAP1, which we detected in cilia in ASC52telo cells and weakly in RPTEC/TERT1 cells, and not in cilia of RPE1 cells, but in the cytosol of all three cell lines (Fig. 4B). This cell type specificity is significantly higher than for other organelles and highlights that the ciliary proteome is much more regulated than other organelles and compartments. In a comparative GO Biological Process term enrichment analysis of the identified cilia proteins between cell lines (Fig. 4C, Supplementary Fig. 3, Supplementary Table 5) we confirmed that indeed each cell line has some unique ciliary specializations. While all cell lines shared GO terms relating to general cilia processes such as (1) ciliogenesis, microtubules, and anchoring of the cilium, (2) transport processes, (3) smoothened signaling, (4) development, and (5) cell polarity (Fig. 4C, Supplementary Fig. 3), we identified cell type specific processes relating to signaling and developmental processes. RPTEC/TERT1 cilia showed enriched functions related to cAMP and kinase signaling, as well as actin and Wnt signaling, while RPE1 cilia showed enriched functions relating to development, and ASC52telo cilia showed functions related to metabolic activity. Furthermore, looking into individual GO terms showed the specific enrichment for the term "establishment of spindle orientation" (adj. P=0.0050) in ASC52telo, supporting our observation that ASC52telo - in contrast to RPTEC/TERT1 - often retain a primary cilium or ciliary vesicle in mitotic cells (Fig. 1E), indicating an engagement of cilia in asymmetric cell division processes of pluripotent cells. Since this study is limited to three cell lines, we expect this heterogeneity to be even higher when extending to more cell types. Notably, the cell type specific ciliary functions align well with the cellular functions of the individual cell lines. RPTEC/TERT1 cells are kidney epithelial cells with their cilia floating in the urine where they may sense hormonal and urine-compositional changes via GPCRs and RTKs, to instruct the cell to furnish the cell surface with adequate transporters for fine-tuned water and ion homeostasis. RPE1 cells show enrichment for processes related to developmental pathways, which might be a consequence of the serum starvation driving them into cell fate changes or senescence. ASC52telo cells have multi-potent features and may require cilium-assisted asymmetrical cell divisions to retain cells with stemness.

Of note, we detected ciliary-motility related proteins, like Hydin, IQCA1, or CATSPER2 only in RPTEC cells and their cilia (Fig. 2K) but not in RPE1 or ASC52telo cells, highlighting that - if some primary cilia have ciliary motility - it may be a tool employed not by all but only some cell types.

Extensive spatiotemporal dynamics of cilia proteins

Thanks to the single cell resolution of our imaging-based approach, we were able to assess variations in protein localization and abundance within primary cilia at the individual cell level. Beyond heterogeneity across cilia in different cell types, we also observed an incredible heterogeneity on the single cell level. 345 of 443 (78%) of the primary cilium, primary cilium tip, or primary cilium transition zone stainings were classified as varying across individual cells. While our images reflect a static snapshot, this heterogeneity within cell lines indicates dynamic spatiotemporal variations of the cilia state in each cell line. Using a comparative GO term enrichment analysis, we explored which processes are particularly stable and which are particularly diverse across individual cilia. As expected, proteins engaging in ciliogenesis, microtubules, and ciliary transport-associated processes appeared in both groups, stable and dynamic (Fig. 4D, Supplementary Fig. 4-5, Supplementary Table 6), while the most dynamic localization was observed for proteins involved in signaling, developmental, and morphogenesis processes.

As expected, we could observe single-cell heterogeneity related to ciliogenesis: For example, RAB8A and RAB34 localized only to individual cilia and are both known to shape the cilium during ciliogenesis by mediating transport of proteins to the growing cilium (Fig. 4E). Interestingly, GPCRs showed varying abundance across cilia even in the serum-starved, synchronized RPE1 cells, such as LPAR3, SSTR3, and 5HT7R (Fig. 4F). Yet, we also detected GPCRs without intensity variation in RPE1 cilia, such as the beta adrenoreceptor ADRB2 (Fig. 4F). In conclusion, we demonstrate that ciliary GPCR signaling even in a population of synchronized cells, i.e., serum-starved hTERT-RPE1 cells, appears dynamic.

Furthermore, many enzymes in cilia showed spatiotemporal variation, such as for HSD3B1/HSD3B2, SPHK1, and AGBL5 (Fig. 4F). Besides locally contributing to shaping a specific lipid environment, such enzymes may also engage in the process of lipid-dependent regulation of signaling, such as for Hedgehog signaling, which has been shown to be regulated by cholesterol [70,77]. Single-cell variation was also seen for Septins, such as for SEPTIN3, which we newly identified in cilia and which heterogeneously localized to RPE1 cilia but not to cilia of the other cell lines (Fig. 4F). Septins are known to be cell type specific and attach to the membrane in a PIP-dependent manner [78]. Observing Septin heterogeneity in single cilia may highlight even more the spatiotemporal dynamics of ciliary lipid-protein-interactions. In conclusion, variation in ciliary localization of lipid-metabolic or -interacting enzymes may relate to the signaling activity of cilia or to fine-tuning of the local ciliary lipid environment.

For some proteins, we observed the variation to correlate with localization of the protein to other cell compartments, or cell cycle stage, such as for CKAP2, where localization to the cilium and part of the microtubule cytoskeleton were seen in some cells, and localization to vesicles in other cells (Fig. 4G). This localization heterogeneity likely relates to the cell cycle, since postmitotic

cells showed high intensity for the vesicle staining. In addition, a transitional stage where the cilium showed staining for CKAP2 only at the shaft could be seen. Further confirming that this heterogeneity may relate to the cell cycle: A similar staining variety among ASC52telo cells was observed, whereas, in RPE1 cells, which were starved and thus, synchronized, the variation was not visible, instead only the stage with microtubules and cilia staining was observed (Fig. 4G).

In summary, compared to other organelles and cellular structures, the primary cilium displays remarkable heterogeneity in multiple regards, both in terms of cell type specificity and in terms of cell-to-cell heterogeneity and subciliary distribution. Altogether this supports the importance of the cilium in being the cellular antenna and computing unit for sensing the environment and computing cellular responses and suggests that it is tightly controlled by extensive spatiotemporal proteome regulation.

Ciliary subcompartments have unique functions in the cell

Spatiotemporal regulation is occuring not only in the cilium as a whole versus other cellular locations but likely also within the cilium. Thanks to the high resolution in our images, we could observe subciliary protein distribution. We distinguished spatial domains in cilia by manual annotation and cataloged proteins by their enrichment at the tip, in the whole cilium, or at the transition zone. Yet, this annotation scheme collected many types of ciliary staining under the term primary cilium (Fig. 2D). An analysis of ciliary intensity profiles with CiliaQ [79] for those proteins localizing to only one of the four annotated cilium-related localizations confirmed the manual annotations scheme (Supplementary Fig. 6A). Yet, due to multilocalization and intensity variation ciliary intensity profiles are extremely diverse, resulting in a high variance in each annotation class, when accepting multilocalizing proteins (Supplementary Fig. 6A).

A comparative GO biological process enrichment analysis of proteins annotated to the different spatial domains in cilia – the tip, cilium, transition zone and basal body - contrasting against cell body regions, revealed certain aspects attributable to subciliary regions compared to the cell body (Supplementary Fig. 6B-C, Supplementary Fig. 7). Beyond the annotated ciliary localizations, we observed a broad variety of localization patterns of enzymes to cilia, e.g., UGGT1 at the transition zone, HSD3B1/HSD3B2 in the proximal cilium region, OCRL more towards the tip, and AGBL5 showed a spotty localization throughout the cilium (Fig. 3E). A similar variety in patterns was observed for signaling proteins (Fig. 3B). We conclude that the subciliary organization is extremely complex and microdomains may exist, but based on the breadth of pattern observations and the continuity between patterns, it is likely that ciliary microdomains may be transient or dynamic, so not only spatially but spatiotemporally regulated, such as what we observed for CKAP2 (Fig. 4G).

A stable protein marker for cilia

Given the multi-localizing and heterogenous aspects of ciliary proteins it is particularly difficult to find suitable marker proteins for cilia imaging assays. A cilia marker protein that is stably and exclusively localizing to primary cilia, could greatly aid in identifying cilia in tissues and cells in research. The common cilia markers such as ARL13B, ADCY3, and acetylated Tubulin all come

with the disadvantage of also localizing to other subcellular structures, which makes it hard to apply them in some cell types, where the non-ciliary aspect of these proteins is dominating or where the proteins are lowly expressed. ADCY3 antibodies appear to not label consistently cilia on all cell types [80,81]. ARL13B also has cytosolic roles [82,83], and, in particular, posttranslational tubulin modifications of the axonemal cytoskeleton, i.e., acetylation, are known to be heterogeneous across cell types and also label cytoskeletal structure in some cells, i.e., neurons [80,84]. In this study, only one single protein - besides the established marker ARL13B - was stably present in the cilia across all three cell lines and for which we did not detect non-cilia expression: CFAP300 (Fig. 5A). Mutations in *CFAP300* have been recently reported in patients with motile ciliopathy phenotypes and abnormal motile cilia structure albeit the function of CFAP300 is not fully understood [85–91]. We here demonstrate that CFAP300 is a suitable marker for primary cilia across multiple human tissues (Fig. 5B). For example, the antibody targeting CFAP300 shows a specific staining in immunohistochemical (IHC) DAB stainings in the fallopian tube (motile cilia), the kidney (primary cilia) and the thyroid gland (primary cilia) (Fig. 5B). Immunofluorescence (IF) co-staining with ARL13B and PCNT show that CFAP300 clearly marks primary cilia in the kidney, the thyroid gland, the cerebellum, and the cortex (Fig. 5C). We thus suggest CFAP300 to be a new reliable cilia marker, complementary to the existing commonly applied marker panel.

Revealing diseases with ciliary defects

To evaluate the functional significance of the newly identified ciliary proteins and to uncover potentially clinically relevant ciliopathy genes, we compared the 654 proteins detected in primary cilium (including tip and transition zone) or basal bodies with phenotype annotations in the Online Mendelian Inheritance in Man (OMIM) database [92]. This analysis resulted in three distinct groups: (1) Genes found in OMIM with a known phenotype (n=300), (2) Genes found in OMIM without an associated phenotype (n=307) and (3) Genes not listed in OMIM (n=47) (Supplementary Table 7, Fig. 6A). Most genes (197 of 300 in group 1, 303 of 307 in group 2, and 46 of 47 in group 3) were not labeled as ciliopathy-causing genes in OMIM (Fig. 6A).

We convolved the clinical analysis of the protein list with the newly discovered proteins that had not been related to cilia before, resulting in the identification of 13 highly interesting proteins with potential disease associations but without described cilia- or ciliopathy-relationships, for which we now provide evidence of localization to cilia (Fig. 6A). Importantly, most of these showed a high spatiotemporal variation across cilia. Surprisingly, the majority of these proteins are enriched in the brain tissue or fulfills key roles in neurons. These include: the cell adhesion molecule NRCAM (Fig. 6B), which is involved in cell-cell contacts and axon outgrowth in the peripheral- and central nerve system [93]; the aromatic-L-amino-acid decarboxylase DDC (Fig. 6B) involved in neurotransmitter metabolism; the Neuronal tyrosine-phosphorylated phosphoinositide-3-kinase adapter 2 NYAP2 (Fig. 6B), which belongs to a family of proteins that regulates neuronal morphogenesis [94]; The calcium/calmodulin-dependent kinase CAMK1G (Fig. 3B), which can act in a calcium/calmodulin dependent signaling cascade important for axon formation [95]; the calcium/calmodulin dependent protein kinase II inhibitor 1 CAMK2N1 (Fig. 6B), which inhibits CAMK2, a key kinase in many neuronal functions [96]. Furthermore, we identified the E3 ubiquitin ligase ZNRF1 at the transition zone (Fig. 6B) - ZNRF1 is enriched in the nervous system and targets various receptors [97], but also interacts with tubulin [98]. The little studied actin binding

transcription modulator ABITRAM was observed in cilia of ASC52telo and hTERT-RPE1, also with single cilium heterogeneity (Fig. 6B). We found the GPCR LPAR3 in individual cilia, as described above (Fig. 4F), highlighting a single-cell-specific ciliary receptor repertoire. Interestingly, LPA signaling through LPA receptors has recently gained more attention in ciliogenesis regulation and glioblastoma, yet focusing on LPA detection through LPAR1 but lacking clarity on where endogenous LPARs localize [99–101]. Furthermore, in a study using zebrafish as a model organism, LPAR3 had been recently implicated to be crucial for establishing Left-Right asymmetry [102], where cilia are known to play important roles, and brain functions [103]. Exogenous FLAG-tagged LPAR1 and LPAR3, introduced by overexpression, have been shown to localize to cilia of immortalized astrocytes [100]. We now reveal endogeneous LPAR3 in human cilia, supporting the relevance for LPA signaling in cilia and placing importance on that LPA signaling may occur through the cilium. An additional protein that we revealed in cilia is AXIN1 (Fig. 6B), reported to be part of the beta-catenin destruction complex and to regulate beta-catenin levels and modulate Wnt signaling [104–106]. Just recently, Wnt signaling has been highlighted as an additional signaling pathway in primary cilia [107]. We add AXIN1 to the picture of the Wnt signaling pathway in cilia. Interestingly, those cilia showing AXIN1 staining in RPE1 cells were off focus compared to the cell nucleus, sticking into the liquid space of the culture well (Fig. 6B). Furthermore, we revealed in cilia the kinesin KIF26B (Fig. 6B), the calcium-dependent regulator of potassium channels ANO3 (Fig. 6B), and the uncharacterized proteins RUNDC3B (Fig. 6B), and ANKRD65 (Fig. 6B).

Lastly, we investigated genes from groups two and three (Fig. 6A) in clinical samples analyzed at the Karolinska University Hospital by whole genome sequencing (WGS) [108,109]. Clinical WGS data from three patients groups were analyzed: 1) patients with a suspected ciliopathy without a causative variant found (n=157), 2) fetuses with an ultrasound-detected malformation without a causative variant found (n= 470) and 3) patients who had undergone a clinical patient-parental trio WGS analysis (n= over 1,500). We searched for potentially clinically relevant single nucleotide variants in the 354 genes in groups two and three (Fig. 6A, Supplementary Table 7) in the WGS data from these three patient groups by one gene at a time, filtering for frameshift variants, nonsense variants and predicted splicing variants in homozygous, compound heterozygous or *de novo* states. Interestingly, we identified one individual in the trio cohort for whom a *de novo* two nucleotide deletion was detected in *CREB3* (NM_006368) (c.810_811del/p.Ser271*) causing a premature stop codon. The individual was a young boy with blindness due to optical nerve hypoplasia. His other symptoms include retinal changes, nystagmus, low bone density, short stature and hypoglycemia. We observed CREB3 in the cilia of RPTEC/TERT1 cells with an enrichment at the ciliary tip, at the transition zone region in ASC52telo cells, and at basal bodies or the transition zone, or at the tip in starved hTERT-RPE1 cells (Fig. 6C). This pattern of localization to base and tip of the cilium might indicate that CREB3 is shuttled through primary cilia. As ciliopathies are rare disorders, patients are often diagnosed after months to years only, or the disease ontogeny remains unresolved since the disease-causing genetic variants are not located in known ciliopathy genes. As we demonstrate here, our new dataset provides new avenues for clinics to reveal potential new ciliopathies and better understand ciliopathies by showing where cilium related proteins localize.

Discussion

Our study provides the first insight into single-cilia proteomic heterogeneity and a unique resource for the cilia community to develop new hypotheses about the cellular functions governed by the primary cilium and their underlying mechanisms. We resolve domains of heterogeneity, i.e., within and across cell populations, highlight the specificity of ciliary signaling pathways, identify a new ciliary marker, and employ our atlas to unlock the mystery behind unresolved ciliopathy-like cases.

While the primary cilium has been studied since the 1960s, we still know very little about it compared to other organelles. Only over the past two decades, research efforts on the cilium took pace, and in manifold studies on ciliary signaling the common hypothesis was established that the cilium likely fulfills an antenna role for the cell [1,3,4]. Still the knowledge on the content, subciliary architecture, mechanisms, and functions of the cilium is very limited, in particular for human cilia. This is mainly due to the challenge of the primary cilium being a tiny cellular appendage and in turn, the lack of methods to study cilia systemically with feasible throughput.

The composition of primary cilia has been studied systematically by proximity-labeling based MS-proteomics approaches, which have provided lists of proteins to localize to cilia of a few murine and human cell types [20–23]. Yet, such approaches require pooling millions of cilia into one measurement and contrasting the measurements of ciliary and other cell compartments to reach subcellular specificity, which in turn renders the detection of multilocalizing proteins that are less abundant in the cilium than in the cytoplasm impossible. Consequently, we lack a systemic understanding of diversity of the ciliary proteome among individual cilia, and of proteins that are multilocalizing to cilia and other cell compartments. Furthermore, comparative studies on the ciliary proteome so far reside on phylogenetic studies or comparison across species [24,39]. Comparing multiple human cell types has not been realized, which might be due to the limitation of proximity-labeling-approaches not being functional in many cell types. In turn, the studies on the cell-specificity of ciliary proteins have focused on a few proteins. For example, many years of mechanistic research, using genetic tools, cellular models, and model organisms have established that GPCRs localize to cilia with cell-type specificity [110]. With our method we greatly advance the understanding of cilia as we show that a broad variety of ciliary proteins, not only receptors but also kinases like PI3K, AKT kinases, or calmodulin-dependent kinases, as well as Cytoskeleton-related proteins, i.e., Septins, and enzymes are in fact cell-type- and even single-cell-specifically localized to the cilium. Importantly, we observe that proteins - when not localizing to cilia - are still expressed and localized to other cell compartments. We show that in 90% of the investigated images a ciliary protein is present in another cellular structure at the same time. We thereby highlight that ciliary specificity across cell types is substantially driven by protein trafficking rather than gene expression. Ciliary import and export has been identified as a signal transduction mechanism, such as in Hedgehog signaling [1]. We demonstrate that many more signaling pathways are underlying spatiotemporal ciliary dynamics and propose that cells rapidly fine-tune their antenna to individually sense their environment depending on their specific task in a cellular consortium. The fact that we also observe downstream signaling machinery in cilia, such as kinases like AKTs, further stresses that the cilium is not just a cellular antenna but also a computer that integrates signals to compute a response decision, to be relayed to the cell body. This

concept has been well established for sperm cells before, where local signaling machinery in the flagellum modifies the flagellar beating for steering the cell [111] - our data indicates that this concept holds true for cilia as well.

Structurally, we go beyond the resolution of previous studies and provide the localization of proteins to cilia with a resolution of confocal microscopy. This is the first proteomic study that distinguishes the protein composition of tip, cilium, transition zone, and basal bodies in the same cells. We thereby demonstrate that the different ciliary regions may be specifically functionalized for different aspects of signaling, which may be modes of shaping the computational architecture of the cilium. Of note, our results hint at spatiotemporal organization beyond the compartments commonly known and that we annotated (Cilium, Tip, Transition zone, Basal Body). We conclude that the ciliary protein architecture is spatially highly organized into microdomains, which are likely dynamic. Our data set can serve as a starting point for systematic super-resolved cilia studies, for example using expansion microscopy or cryo-CLEM.

Importantly, the ciliary compartment is not just organized on the level of proteins. Beyond signaling molecules, we also reveal enzymes in cilia that may close gaps in our understanding of how the ciliary compartment is organized. For example, we reveal an enzyme involved in sterol synthesis supporting the hypothesis that cilia locally metabolize their unique lipid environment for customizing or conducting signaling pathways [69]. The presence of the other enzymes indicates that the cilium has its own metabolism, which may be crucial to further study to understand its contribution to ciliary signaling pathways and functions. The heterogeneity in these factors highlights the importance for single-cell spatially-resolved methods for protein, lipid, and metabolite measurements to further delineate the complex architecture of the ciliary compartment.

Beyond revealing the cilium as a signaling device, our data also highlights new avenues for cilia research. For example, we surprisingly detect the least number of cell-type-unique proteins in ASC52telo cells, which show only 16 or 12 cell-type unique primary cilia or basal body protein localizations - much less than RPE1 (68 or 73, respectively) or RPTEC/TERT1 cells (69 or 62, respectively). We cannot exclude that a technical limitation hampered discovery in ASC52telo cells: they have shorter cilia compared to the other two cell types (Fig. 1D), which typically lie flat on the cells, and they show a significantly lower ciliation rate. However, a biological reason for finding only a few proteins specific to ASC52telo cilia may be that we developed the candidate list of proteins based on existing literature and data sets, which majorly come from epithelial cells, whereas ASC52telo cells are mesenchymal. In addition, the fraction of proteins not previously described for cilia in our candidate list was collected with a strong focus on signaling cascades. By contrast, based on our gene ontology analysis of ASC52telo cells and the fact that they have cilia only for a very short time due to their fast cell cycle, we hypothesize that the cilium of ASC52telo cells is less tuned to receptor-based signaling but rather crucial for driving stemness and asymmetric cell division in a stem-cell population. Such asymmetry may be produced by the ASC52telo cells retaining a ciliary vesicle or whole cilium during cell division. Thus, our data encourage us to next direct our antibody-based spatial proteomics methods to stem cells with a redeveloped candidate list of ciliary genes tailored more to stem cell functions. Furthermore, we surprisingly identified many neuron- and brain-related proteins in cilia, which emphasizes that our

focus should also direct towards neuronal cilia. Of note, some of these proteins relate to synapses or axons, supporting the recent observations of the cilium being involved in synaptic transmission [112] and that the cilium is not just an organelle that is a free-floating antenna but that can also establish contacts with other structures, i.e., synapses, for even more local signal reception.

Antibody-based profiling provides many advantages as discussed above, but also comes with the need for careful assessment of antibody specificity. While the HPA antibodies are highly validated, we cannot exclude that some of the revealed 654 proteins may be false positives due to unspecific binding or cross-reactivity of the antibody, the likelihood for this is particularly low in light of the significant diversity we observe in the ciliary proteins. Given the limitations with antibodies and the difficulty studying a tiny structure like primary cilia, we do not conclude that the proteins not detected in cilia in any cell lines are in fact not in cilia. Biological reasons for not detecting them may be that they localize to cilia only in other cell types or under other culturing conditions, such as upon stimulation with a receptor ligands, or that they are of extremely low abundance. To shed further light on spatiotemporal dynamics in response to stimuli, new avenues for spatiotemporal proteomics will need to be explored.

Our study opens up many new avenues for ciliary research. First of all, we have tested and validated antibodies for 654 proteins in primary cilia in human cells. Most of these antibodies are commercially available and allow the field to probe ciliary signaling and structure in detail, for example, when investigating patient-derived cells to explore disease mechanisms. Second, we also show that our screen has the potential to be translated to deep studies of human tissues, which we pursued for revealing CFAP300 as a new ciliary marker. Notably, many of the other antibodies that we tested are also suitable for tissues, as they are already validated in IHC workflows in the Human Protein Atlas [25,32]. Recently, we also showed that the antibody for ATF3, which we identified at the Basal Body, is also applicable to study the role of cilia in the developing human heart [113]. Third, we generate the basis for revealing additional ciliopathy cases in the clinics. We demonstrate that our new atlas can link a clinical phenotype of a CREB3 variant to the cilium.

In summary, we highlight the primary cilium as a cellular compartment with a diverse proteome involved in signal transduction mechanisms. The ciliary compartment is tightly regulated to bring together certain signaling molecules in time and space. We hypothesize that cells employ their cilia to customize sensation and signal transduction to individual cellular roles. Thereby, the primary cilium might be a space in which the cell can both tailor their environmental sensing abilities and customize the internal computation to allow precise cell type and state-tuned signaling. Our data indicates that there are multiple domains of ciliary heterogeneity potentially relating to cell states, such as cell cycle, ciliogenesis, signaling, and metabolism. Further deep analysis of our data set followed up by functional assays and live measurements of the proteins undergoing heterogeneity, will allow the cilia community to further define these axes of heterogeneity. Hence, this study serves as a starting point for cilia systems biology, providing a template for modeling the cilium as an information processing unit. With all our data being publicly explorable at https://www.proteinatlas.org we pave new avenues for research on decoding the fundamental biology of human primary cilia and clinical implications such as in ciliopathies.

Figures

Figure 1: Spatially profiling the primary cilia proteome

A-B) New staining concept. **A)** Ciliated cells (i.e., starved hTERT-RPE1) cells are co-stained with markers for nuclei (DAPI), cilia (ARL13B antibody), basal bodies (PCNT antibody) and for a protein of interest (i.e., with a polyclonal antibody against IFT43 from the Human Protein Atlas). **B)** Confocal z-stacks are generated, which allow 3D analysis of the whole cell including the cilia, even when they stick out of the cell to the top or bottom (Blue: Nuclei, Red: Cilia, Yellow: Basal Body, Green: Protein of Interest). **C)** Three cell lines are investigated: Starved hTERT-RPE1 cells from the retina, RPTEC/TERT1 cells from the kidney, and ASC52telo cells from the adipose tissue. Example images from all three cell lines are shown with magnified insets, from a co-staining of DAPI (labeling nuclei), ARL13B (labeling cilia), and the protein of interest IFT20. Gender of the cell line indicated. **D)** CiliaQ analysis of all images underlying this study to determine the ciliation rate and cilia length distribution of the studied cell line cultures. In total 463,454 cells with 218,410 cilia were included. Top: Each dot represents all images from one individual experiment day. Bars and error bars represent median and quantiles, respectively. Bottom: Normalized histograms of the cilia lengths for all cilia analyzed, distinguished by cell line. **E-H)** Example images of special cilia observations in the three different cell lines. Only the Nucleus (DAPI), cilia (ARL13B), and Basal Body (PCNT) channels are displayed. **E)** Mitotic ASC52telo cells retaining a cilium. **F)** RPTEC/TERT1 cells with two adjacent cilia. **G)** Starved hTERT-RPE1 cell with two adjacent cilia. **H)** Two adjacent RPTEC/TERT1 cells whose cilia are connected and form a "bridge". **I)** Sunburst plot of all data sets accumulated to form the cilia candidate list of 9634 human genes. Different data sets and studies, as well as additional educated guesses were convolved (see Materials & Methods). Each study is colored by species. **J)** Candidate list of 1,935 genes for which a staining with Human Protein Atlas antibodies was conducted, including a substantial portion of genes that had not been associated with cilia before ("new", purple). Scale bars indicated.

Figure 2: Spatial map of the primary cilia proteome

A) Schematic drawing of a primary cilium and its substructures. **B)** Protein localization to four ciliary substructures was annotated, for which example images are shown: Primary cilium, if the protein was overlapping with the ciliary marker ARL13B; Primary cilium Tip (Tip), if the protein of interest channel showed a signal at the distal end of the cilium; Basal Body, if the protein of interest channel showed signals overlapping or accumulating close to the basal body marker PCNT; Primary cilium transition zone (Transition zone) if the protein of interest channel showed an enrichment between basal body and cilium. **C)** Venn diagram showing the number of genes observed at one or multiple of the four locations specified in B. **D)** Example images (single confocal slice images) of additional subpatterns observed in primary cilia. Gray interrupted line distinguishes two individual images placed adjacently for showing two cilia. **E)** Example images (single confocal slice images) of stainings for proteins that are established as localizing to cilia - gene names are shown. **F)** Comparing the list of genes encoding proteins that were annotated as localizing to at least one of the four cilium locations (B) to existing datasets. 71 proteins were not found in other datasets. **G-H)** Pie charts showing the number of occurrences in other datasets for each gene that encoded a protein found in one of the four annotated cilia compartments. **G)** Comparing the list of revealed proteins to datasets derived from Human samples only. **H)** Comparing the list of revealed proteins to datasets from any species. **I)** Example images of proteasomal subunits observed in cilia. **J)** Example images (single confocal slice images) for proteins observed in cilia that were previously only detected in individual proximity-labeling-based MS approaches. **K)** Motile cilia proteins detected in the cilia of RPTEC/TERT1 cells. Images were derived from stainings of different cell lines: Starved hTERT-RPE1 cells: DNAL4, ZNRF1, EFCAB7, EVI2A, TRIM46, SMAD7, RAB7A, SEPTIN2, RAB23, ARL15; RPTEC/TERT1 cells: CFAP300, TRIO, C2CD4C, TMEM237, IFT22, ARL6, PSMD10, PSME3, PSMA6, HYDIN, IQCA1, CATSPER2; ASC52telo cells: KIF7, CEP164, TCTN2, KIF7, ADCY3, ZNF330, RSL1D1.

Figure 3: The primary cilium is a signaling hub

A) Comparative Gene Ontology enrichment analysis of Biological Process terms between the two major cilia locations primary cilium (including tip) and basal body (including transition zone). The Venn diagram shows the number of terms found to be significantly enriched (adj. P value < 0.01 and q value < 0.01) for only primary cilium (n=125) or basal body (n=129), as well as for both locations (n=198). Significant terms were clustered by semantic similarity, and for each resulting GO term cluster, a word cloud is shown summarizing the biological functions of the terms in the cluster based on keywords enrichment of term names. The font size in the word clouds correlates with -log(P value) of the key word enrichment, i.e. the larger, the more enriched a keyword is compared against background GO vocabulary. The color in the heat maps represents the semantic similarity. **B)** Protein localization of example signaling proteins to the primary cilium. All images show single confocal slice images except for PDGFRA, where a maximum intensity projection of 5 confocal slices is shown to show the whole cilium. The inset for PDGFRA shows the same position with displaying only the PDGFRA channel. **C)** Multilocalization of cilia-related proteins with the other 35 annotated subcellular compartments. Abbreviations: Centriolar Sat. = Centriolar Satellites; Cyt. bridge = cytokinetic bridge; Int. Fil. = Intermediate Filaments; Mitotic chr. = Mitotic chromosome; Nuc. bodies = Nuclear bodies; Nucleoli fib. cent. = Nucleoli fibrillar center; Nuc. Membrane = Nuclear membrane; Nuc. speckles = Nuclear speckles. **D)** Example transcription factors revealed in cilium and nucleus. **E)** Example enzymes revealed in cilia. Images were derived from stainings of different cell lines: Starved hTERT-RPE1 cells: PTK2, PP2R3C, CNGA2, TULP3, CRX, ATF3, CREB1, HSD3B1/HSD3B2, LANCL1, OCRL, SPK1; RPTEC/TERT1 cells: AKT1, AKT3, TAOK1, PIK3CA, PIK3R4, CAMK1G, MIB1, OPRL1, PDGFRA, AGBL5, UGGT1; ASC52telo cells: FLT4, TGFBR1.

Figure 4: Cellular diversity and spatiotemporal dynamics of the primary cilia proteome

A) Venn diagram of the genes for the proteins identified in primary cilia (including tip and transition zone), distinguishing by cell line. Only genes included for which an approved staining was obtained from all three cell lines. (Abbreviations: RPTEC = RPTEC/TERT1, RPE1 = starved hTERT-RPE1) **B)** Example images (single confocal slice images) for stainings across the three cell lines, showing heterogeneity in ciliary localization across cell types. **C)** Comparative Gene Ontology enrichment analysis of Biological Process terms between the three included cell lines to investigate cell-line differences and commonalities. The Venn diagram shows the overlap of significantly enriched terms (adj. P value < 0.01 and q value < 0.01) for each cell line. Significant terms of each section of the Venn diagram were clustered by semantic similarity, and for each resulting GO term cluster, a word cloud is shown summarizing the biological functions of the terms in the cluster based on keywords enrichment of term names. The font size in the word clouds correlates with -log(P value) of the key word enrichment, i.e. the larger, the more enriched a keyword is compared against background GO vocabulary. **D)** Comparative Gene Ontology enrichment analysis of Biological Process terms between proteins annotated as displaying intensity variation in one of the four ciliary locations, and annotated as stable as described in C). **E)** Example images (single confocal slice images) for stainings of the two ciliogenesis-related proteins RAB8A and RAB34. **F)** Example images (single confocal slice images) of stainings revealing proteins with heterogeneous localization to primary cilia. **G)** Example images (maximum intensity projections of whole stack) of CKAP2 staining in the three cell lines.Cell lines for images (if not indicated): Starved hTERT-RPE1 cells: LPAR3, SSTR3, 5HTR7, ADRB2, HSD3B1 HSD3B2, SPHK1, RAB8A, SEPTIN3; RPTEC/TERT1 cells: AGBL5; ASC52telo cells RAB34.

Figure 5: CFAP300 - a new ciliary marker protein

A) Example images for stainings of the CFAP300 in the three cell lines. **B)** Example images from IHC stainings on human tissues (sex and age indicated in brackets) with the CFAP300 antibody. **C)** Confocal microscopy images (Maximum intensity projections of z-stacks) from different human tissues stained for CFAP300, cilia (ARL13B antibody), and nuclei (DAPI).

Figure 6: Clinical relevance of the newly revealed ciliary proteome

A) Flow chart for splitting the 654 genes into three groups based on their OMIM annotations. Additionally the genes were classified by being established ciliopathy proteins. Example genes for each group are written on the right, and shown in B. **B)** Example images for staining the

proteins for genes pointed out in A. Cell lines for images: Starved hTERT-RPE1 cells: NRCAM, DDC, NYAP2, CAMK2N1, ZNRF1, AXIN1, KIF26B, ANO3; ASC52telo cells: ABITRAM. **C)** Example images from stainings of CREB3 in RPE1 cells (serum-starved), ASC52telo cells, or RPTEC/TERT1 cells. The white interrupted line distinguishes two images that were placed adjacently to show two cilia. Abbreviations: TZ = transition zone. BB = Basal Body. The basal body PCNT marker was not shown to better visualize the localizations of CREB3 - The PCNT marker was however used to correctly mark BB and TZ positions with arrows.

Materials and methods

Developing a list of cilium-associated candidate proteins

Lists of genes or proteins detected in primary cilia, motile cilia, flagellum in humans and across other species were extracted from databases in the format of supplementary tables, PDF-files, or exportable csv files. The data from ChlamyFP.org (Human) were downloaded on July 10, 2024. The files were next imported into R and merged into a master list (Supplementary Table 1) by conversion of the protein or gene identifiers into Ensembl gene IDs using the R package biomaRt (version 2.56.1) [114]. Conversions failing in bioMart were converted - if possible - by automated queries to the Ensembl REST API (https://rest.ensembl.org). The data overlap was plotted using the library Plotrix (version 3.8.4). To reduce the list to the candidate list of 1,935 genes, we selected targets based on (1) whether the gene is expressed in the cell lines we aimed to study (using HPA internal cell line RNAsequencing data), (2) whether they represent gold standard cilia proteins, (3) whether the HPA antibody library provided a good, validated antibody for the protein of interest, and (4) the discovery potential of the protein of interest, with a strong focus on revealing signaling proteins in cilia.

Staining prioritization and order of experiments

We started by staining the one or two cell lines with highest RNA expression (HPA internal cell line RNA expression database) and did not pursue further staining in the other cell line(s) if the staining / antibody was deemed unspecific or unreliable or did not show cilium or basal body localization of the protein and in the other cell lines the protein was not indicated as expressed on RNA level. This resulted in not all proteins being stained in all three cell lines (Supplementary Fig. 2a). For all conclusions drawn on inter-cell line diversity, only proteins that have been stained in all three cell lines were included.

Cell culture

hTERT-RPE1 cells (Evercyte GmbH, Vienna, Austria) were acquired from ATCC and cultivated in DMEM/F12 (Thermo Fisher Scientific Inc, GibcoTM, Cat. no. 11320033) + 10% FBS (Merck KGaA, Sigma-Aldrich, Cat. no. F7524) at 37°C and 5% CO_2. ASC52telo and RPTEC/TERT1 cells were acquired from Evercyte GmbH (Vienna, Austria) and cultivated according to manufacturer's instructions. ASC52telo cells were cultivated in Endothelial Cell Growth Medium-2 (Lonza, Cat. no. CC-3162) supplemented with 4% FBS (Merck KGaA, Sigma-Aldrich, Cat. no. F7524) and 200 ug/ml G418 (G-418 Solution, Merck KGaA, Cat. no. 04727878001). RPTEC/TERT1 cells were cultivated in ProxUp2 medium (Evercyte, MHT-003-2).

Immunocytochemistry

Cells were seeded in 80 µl / well of the respective growth medium on 96-well glass-bottom plates (Greiner Sensoplate Plus, Greiner Bio-One, Germany) coated with 50 µl of 12.5 µg/ml human fibronectin (Sigma Aldrich, Darmstadt, Germany) and cultivated at 37 °C and 5% CO_2. For hTERT-RPE1 and ASC52telo cells, approximately 10,000 cells/well were seeded. For

RPTEC/TERT1 cells, approximately 30,000 cells/well were seeded. hTRET-RPE1 cells were starved 24 hours after seeding by medium exchange to DMEM/F12 + 0.2% FBS, and fixed 48 hours after start of starvation. RPTEC/TERT1 cells were fixed 24 hours after seeding. ASC52telo cells were fixed 48 hours after seeding. For fixation, the cells were washed with phosphate-buffered saline (PBS, pH 7) and fixed with 40 µl 4% ice-cold paraformaldehyde (PFA: Sigma Aldrich, Darmstadt, Germany) dissolved in PBS for 15 min. Cells were washed with PBS and permeabilized with 40 µl 0.1% Triton X-100 (Sigma Aldrich) in PBS for 3 × 5 min. Rabbit polyclonal HPA antibodies targeting the proteins of interest were dissolved to 2–4 µg/ml in blocking buffer (PBS + 4% FBS) containing the cilia marker antibodies against ARL13B (Abcam, ab136648, mouse IgG2a, 1:2000) and PCNT (Abcam, ab28144, mouse IgG1, 1:2000). After washing with PBS, the diluted primary antibodies were added (40 µl per well), and the plates were incubated overnight at 4 °C. After overnight incubation, wells were washed with PBS for 3 × 10 min. Secondary antibodies, goat-anti-mouse-IgG1k-Cross-Adsorbed-Alexa555 (ThermoFisher, A-21127, 1:2000), goat-anti-mouse-IgG2a-Cross-Adsorbed-Alexa647 (ThermoFisher, A-21241, 1:1000), and goat-anti-rabbit-IgG (H+L)-Highly-Cross-Adsorbed-Alexa488 (Invitrogen, A11034, 1:800) diluted in blocking buffer were added and the plates were incubated for 90 min at room temperature. Next, cells were incubated with 572 nM = 200 ng/ml of DAPI (4', 6-diamidino-2-phenylindole) in PBS for 10 min, and washed 3 x 5 min with PBS.

Antibodies

Except for the marker antibodies ARL13B (Abcam, ab136648) and PCNT (Abcam, ab28144), the antibodies used in this study were generated within the HPA project. All HPA antibodies are rabbit polyclonal antibodies generated using recombinant protein epitope signature tags (PrEST) as antigens for immunization [115]. The PrESTs are designed to enable the antibody to bind specifically to as many isoforms of the target protein as possible. The resulting antibodies were affinity purified using the PrEST antigen as an affinity ligand. Commercial antibodies were provided by the suppliers and used according to the supplier's recommendations. As previously described, the antibodies were initially quality controlled for sensitivity and lack of cross-reactivity to other proteins using western blotting and protein arrays [115]. Antibodies that passed the initial quality assessment, with no literature contradicting the results obtained in immunofluorescence (IF), were labeled 'approved'. If there was independent data in UniProt supporting the results in IF, the antibody was labeled 'supported'. For enhanced application-specific validation, we used the strategies outlined by the International Working Group for Antibody Validation (IWGAV) [116], including i) validation by gene silencing/knockout, ii) validation by co-staining of fluorescent protein-tagged protein expressed at endogenous levels, iii) validation by independent antibody targeting a different epitope, and iv) capture validation by mass spectrometry [116,117]. If the antibody had been validated according to the IWGAV guidelines, it was labeled 'enhanced'.

Fluorescence image acquisition of cells

All fluorescent images of cells were acquired with an inverted Leica SP8 confocal microscope equipped with a motorized and piezo z stage, an HC PL APO CS2 63×/1.20 water-immersion objective (Leica Microsystems, Mannheim, Germany), Laser Diodes 405, 488, 552, and 638, and HyD detectors. The settings for each image were Pinhole 1 Airy unit, 16-bit acquisition, 1.28x

zoom, 2048x2048 pixels field of view, 700 Hz scan speed, resulting in a pixel size of 0.07 μm. All channels were detected on HyD detectors in multiple individual series to reduce bleed through: (1) DAPI and Cilia (Alexa 647) were detected, (2) the protein of interest was detected (Alexa 488), (3) the basal body channel was detected (Alexa 555). The detector gain and the laser intensity for the protein of interest (Alexa 488) channel were constant for all images acquired. On every plate a negative control (no primary rabbit antibody) was included and recorded to confirm that no nonspecific signals were visible in the Alexa488 channel. Detector gains and laser intensities for the other channels were minimally adjusted between different plates and cell lines, if needed (the different cell lines showed different strength of ARL13B PCNT and DAPI signals and thus, customization was required). Acquisition of images was conducted fully automatically, at least 4 images at random positions per well were acquired. If these images did not contain cilia or were deemed insufficient for deciding on a subcellular location (in the IF image annotation workflow described below), additional images at random positions were acquired. For ASC52telo cells, in this additional acquisition step, the field of views were placed manually in ciliated regions with many cells (without looking at the protein of interest channel).

Image processing

Images were exported to OME-TIFF format through Leica LASX's inbuilt 3D viewer software. The OME-TIFF images were next reprocessed by a custom-written ImageJ plugin that complements and corrects metadata in the OME-XML strings in the tiff files based on the xml-based metadata files co-saved from Leica´s LAS-X software. Next all images were imported into the Laboratory Internal Management System (LIMS) of the Human Protein Atlas. Here, the intensity range for all channels was adjusted by an automatic script lowering the maximum displayed intensity value to improve visibility of stainings. The maximum display intensity was, however, never allowed to fall below 10,000 a.u., which corresponds to a value where in a negative control staining no signals were visible (Negative control staining was obtained by keeping all markers and secondary antibodies in the procedure but leaving out an antibody for the protein of interest). During manual annotation (see below), the maximum displayed intensity value was - if needed - further adjusted to better visualize ciliary signals but also never allowed to be set below 10,000 a.u. For display on the https://www.proteinatlas.org webpage, all images were converted from TIFF to JPEG file format based on the set maximum display intensity range.

IF image annotation

The subcellular location of each protein was manually determined based on the signal pattern and relation to the markers for nucleus (DAPI), cilia (ARL13B), and basal bodies (PCNT). 35 different subcellular locations (Intermediate filaments, Actin cytoskeleton, Microtubules, Microtubule ends, Centriolar satellites, Mitochondria, Aggresome, Cytosol, Cytoplasmic bodies, Rods & Rings, ER, Golgi, Vesicles, Peroxisomes, Endosomes, Lysosomes, Lipid droplets, Plasma membrane, Cell Junctions, Focal Adhesions, Nucleoplasm, Nuclear membrane, Nucleoli, Nucleoli Fibrillar center, Nucleoli rim, Nuclear speckles, Nuclear bodies, Micronucleus, Kinetochore, Mitotic chromosome, Cytokinetic bridge, Midbody, Midbody ring, Cleavage furrow, Mitotic spindle), the standard for the subcellular section of the human protein atlas[26], and four additional cilia-related locations were annotated for all images: primary cilium, basal body, primary

cilium tip, primary cilium transition zone. If more than one location was detected, they were defined as the main or additional location depending on the relative signal strength between the location and the most common location when including multiple cell lines, including also non-ciliated cell lines in the HPA. Variation between single cells were annotated as variation in the intensity or spatial distribution based on a visual inspection. The staining was not annotated if considered negative or unspecific.

Of note, it is likely that in many cases - given the resolution of the method - it was not possible to distinguish a transition zone and primary cilium staining from an only primary cilium staining unless the transition zone staining is significantly brighter than the whole cilium staining. The proteins annotated to stain the transition zone showed a clear only transition zone staining in at least one cell line. The same applied to ciliary tip proteins.

Image analysis

OME-TIFF files for individual channels and slices were fetched from the internal image database of the Human Protein Atlas and composed into z-stack tif-files. To segment individual cilia and basal body from the 3-dimensional image stack, we firstly blurred the image using a Gaussian kernel (standard deviation of 1) and subtracted the background which is obtained by blurring the original image using a larger Gaussian kernel (standard deviation of 10). Next, we set threshold values to segment the cilia and basal body objects. We empirically found that Otsu thresholding and 10 times of triangle thresholding values work best for basal body and cilia, respectively. To refine the segmented structures, we performed image opening using a ball kernel of radius 1, then removed small unconnected objects from the segmentation (objects smaller than 75 and 200 pixels for basal body and cilia, respectively). We also removed the falsely segmented cilia that is not attached to any segmented basal bodies.

To analyze how many cells and cilia were in the data set, as well as the ciliation rate, we used a new version of CiliaQ that tracks basal bodies / centrosomes and cilia (CiliaQ Version v0.2.1) [79].

Considering the different sizes of intensity profiles associated with each cilia, we projected each path onto the same 1-dimensional grid from 0 to 1 with 100 intervals (the starting and ending points are mapped to 0 and 1 respectively) and interpolated the profile according to the grid. Specifically, for cilia that are longer than 1.16 micrometers, we partitioned it into two pieces (0 to 1.16 micrometers and 1.16 micrometers to its end) and projected the first part onto the grid of 0 - 20 and the second part onto the grid of 21 - 100. For the cilia shorter than 1.16 micrometers, we projected it onto the grid points 0 to 20 and filled the vector by 0s after that. The rationale here is to prevent stretching protein intensity at the bottom of short cilia too much and it helps us to better characterize the non-deformed basal body / transition zone. It results in protein intensity profiles as vectors of length 100. In addition, we normalized profiles by the maximal values among all profiles belonging to the same staining and smoothened each profile using a 1-dimensional Gaussian kernel (standard deviation of 8). To visualize the differences between profiles associated with proteins at different subcellular structures, for example, proteins located at basal body, primary cilium, primary cilium tip and transition zone, we plotted the median, 25 and 75 percentile among intensity profiles at each grid point along the normalized cilia path for proteins localized in each of those structures.

Multilocalization analysis

To assess whether ciliary proteins localize to significantly more subcellular compartments than the overall proteome, we first counted the number of subcellular compartments annotated for each protein in the Human Protein Atlas. We then compared the distribution of the number of subcellular localization annotations for all proteins in the HPA (with any subcellular location annotation) to that of ciliary proteins—specifically, those annotated with "primary cilium tip," "primary cilium," and "primary cilium transition zone." A two-sided Mann-Whitney U test was used to determine statistical significance, with a P-value threshold set at $P < 0.01$.

To study the locations, where multilocalizing proteins were observed, the annotations on the staining level (data split by protein and cell line) from all cell lines were converted into edges with the python package networkx (version 3.2) and next plotted using the python package d3blocks (version 1.4.6).

Functional enrichment analyses

Functional enrichment analyses were conducted in R v4.4.1 using clusterProfiler v4.12.6 [118], enrichplot v1.24.4, and simplifyEnrichment v1.14.1 [119]. Overrepresentation of Gene Ontology Biological Process terms compared to all human genes was computed using the compareCluster function of clusterProfiler with org.Hs.eg.db as annotation database. The results were controlled for multiple hypothesis testing by calculating adjusted P values using the Benjamini-Hochberg Procedure and q values and restricting the results to adjusted P values < 0.01 and q values < 0.01. Results were visualized using the dotplot function from enrichplot. Additionally, the resulting list of significantly enriched GO terms for each variable in each comparison was simplified by calculating and clustering the corresponding semantic similarity matrix of the significant GO terms. Relational similarity was used as the similarity measure and binary_cut as clustering method. For each of the resulting clusters, a word cloud was drawn summarizing the biological functions with keywords in every GO term cluster. For this, the enrichment of keywords compared to the background GO vocabulary is determined using Fisher's Exact test and mapped to the fontsize of the keywords in the word clouds. Overrepresentation of Reactome pathways was determined equivalently, but were not further simplified.

To compare ciliary locations to cellular locations the 35 HPA locations were simplified into four groups: cytoplasm, membrane, nucleus and mitotic. Cytoplasm collected annotations for Intermediate filaments, Actin cytoskeleton, Microtubules, Microtubule ends, Centriolar satellites, Mitochondria, Aggresome, Cytosol, Cytoplasmic bodies, Rods & Rings. Membrane contained the endo-membrane system, collecting annotations for ER, Golgi, Vesicles, Peroxisomes, Endosomes, Lysosomes, Lipid droplets, Plasma membrane, Cell Junctions, Focal Adhesions. Nucleus collected annotations for Nucleoplasm, Nuclear membrane, Nucleoli, Nucleoli Fibrillar center, Nucleoli rim, Nuclear speckles, Nuclear bodies, Micronucleus. Mitotic collected annotations for Kinetochore, Mitotic chromosome, Cytokinetic bridge, Midbody, Midbody ring, Cleavage furrow, Mitotic spindle.

Human tissue sample preparation

The tissue sample preparation protocol used was the same as that described earlier by Danielsson et al. [120]. The tissue samples were processed following Swedish laws and regulation, obtained from the Department of Pathology, Uppsala University Hospital, Sweden, as included in the sample collection governed by the Uppsala Biobank (http://www.uppsalabiobank.uu.se/en/). All human tissue samples used were anonymized in accordance with an approval and advisory report from the Uppsala Ethical Review Board (Dnr Ups 02-577). Tissue MicroArrays (TMA) were constructed with strategies used in the Human Protein Atlas previously described [121,122], in Formalin-fixed paraffin-embedded (FFPE) format. In summary, hematoxylin-eosin stained tissue sections from FFPE donor blocks were examined to confirm histology and identify representative regions for sampling into the TMA. Normal tissue was characterized as microscopically normal, predominantly sourced from proximity areas to surgically excised tumors. Two TMA blocks were created, each containing triplicate 1 mm cores from a total of 45 different types of normal tissue (tissues listed in Supplementary Table 10).

Immunofluorescence staining of tissues

We used the same indirect immunofluorescence staining protocol described by Martinez Casals et al. [123]. with some adjustments. In brief: single section from each of the human TMA FFPE blocks was sectioned at 4 µm thickness. Each section was dewaxed and underwent heat-induced epitope retrieval (HIER) treatment using pH 9 EDTA buffer (Akoya Biosciences, AR900250ML) in a pressure cooker (Bio SB TintoRetriever, BSB-7087) at 114–121 °C during 20 minutes. After two ddH2O washes, the material was exposed to a photobleaching procedure to reduce tissue autofluorescence; following a blocking step using TNB buffer containing 0.1 M Tris-HCl /0.15 M NaCl/0.5% blocking reagent pH 7.5 (Akoya, SKU FP1020) for 30 min. Primary antibodies targeting ARL13B (RRID:AB_3073658 - Abcam, ab136648, mouse IgG2a, 1:300), PCNT (RRID:AB_2160664 - Abcam, ab28144, mouse IgG1, 1:500), and CFAP300 (HPA, HPA038585, rabbit polyclonal, 1:100) were diluted in antibody diluent containing 0.3 % Triton (Sigma Aldrich, T8787) 1x PBS pH 7.4 and added on the TMA section to be incubated overnight at 4°C. The next day, the material was washed with TBS-T and blocked using TNB for 30 min. Secondary antibodies, goat-anti-mouse-IgG2a-Cross-Adsorbed-Alexa647 (ThermoFisher, A-21241, 1:600), goat-anti-mouse-IgG1-Cross-Adsorbed-Alexa555 (ThermoFisher, A-21127, 1:600), goat-anti-rabbit-IgG (H+L)-Highly-Cross-Adsorbed-Alexa488 (Invitrogen, A11034, 1:600) were diluted in TNB, containing Hoescht (ThermoFihser, H3570), and added on the section for 90 min incubation at room temperature. Lastly, the stained material was mounted with Fluoromount-G (Thermo Fisher Scientific, 00-4958-02) to proceed with image acquisition.

Fluorescence image acquisition of tissue cores from TMA sections

Regions of interest in selected cores of the TMA were image acquired using Leica STELLARIS 8 confocal microscope equipped with a 63 HC PL APO 63x/1.20 water-immersion CS2 objective (Leica Microsystems. Mannheim, Germany). The settings for each image were Pinhole 1 Airy unit, 16-bit acquisition, 1x zoom, 2048x2048 pixels field of view, Z-stack size 9.1 µm, number of steps 14, resulting in a pixel size of 0.09 µm and a z-step size of 0.7 µm. Detectors used: HyD S1 was

used for Hoechst channel, HyD H2 for A488 channel, Hyd S3 regarding the A555 channel, and Hyd X4 for Alexa 647 channel. The "Lightning" deconvolution software (Leica Microsystems, Mannheim, Germany), which was integrated into the microscope software, was used to deconvolve the images after recording.

Immunohistochemistry images

The immunohistochemical stainings were performed as part of the regular Human Protein Atlas workflow, as described elsewhere. [25,32]

Exploration of clinical databases

Clinical samples analyzed at the Karolinska University Hospital by whole genome sequencing (WGS) were searched for interesting variants in genes who were classified in OMIM [92] as without a phenotype or not annotated in OMIM at all (Fig. 6A, Supplementary Table 9). The overall detailed analytical process of WGS has previously been described in Stranneheim et al. [108] and in Ek et al. [109] Clinical WGS data from three groups of patients were queried: 1) patients with a suspected ciliopathy disorder without a causative variant found in a clinical WGS (n=157), 2) fetuses with an ultrasound-detected malformation without a causative variant found in a clinical WGS (n=470) and 3) patients who had undergone a clinical patient-parental trio WGS analysis (n>1,500). We searched for potentially clinically relevant single nucleotide variants in the WGS data from these three patient groups by one gene at a time, filtering for the found variants so that only frameshift variants, nonsense variants and predicted splicing variants in a homozygous, compound heterozygous or *de novo* state were retained.

Data availability

The data presented in this study, including staining annotation, is available in the Human Protein Atlas (v24) at https://www.proteinatlas.org.

Acknowledgement

This research was funded by grants from Göran Gustafsson Foundation, Erling Persson Foundation, Chan-Zuckerberg Biohub, and Knut and Alice Wallenberg Foundation to EL (KAW 2021.0346) as well as to MU (HPA). JNH was supported by a Postdoctoral Fellowship from the Wenner-Gren Foundations, and by an EMBO Postdoctoral Fellowship (ALTF 556-2022).

Author contribution

JNH, UA, and EL conceived the project and designed the study. JNH, UA, AJ, KT, AMC, EP, JF, and AB conducted experiments. UA, JNH, KT, JF, EL, and DM annotated images. JNH, UA, EL, HS, KK, EW, CW, and DM analyzed and interpreted data. MU and CLB contributed to the discussion and critical reagents. JNH, UA, and EL wrote the paper. ADV and AL supervised the clinical analysis. EL supervised the project. KVF, FJ, and UA built the cilia atlas in HPA. All authors edited and commented on the manuscript.

References

1.   Mill, P., Christensen, S.T., and Pedersen, L.B. (2023). Primary cilia as dynamic and diverse signalling hubs in development and disease. Nat Rev Genet. https://doi.org/10.1038/s41576-023-00587-9.

2.   Reiter, J.F., and Leroux, M.R. (2017). Genes and molecular pathways underpinning ciliopathies. Nature reviews. Molecular cell biology *18*, 533–547. https://doi.org/10.1038/nrm.2017.60.

3.   Pazour, G.J., and Witman, G.B. (2003). The vertebrate primary cilium is a sensory organelle. Current opinion in cell biology *15*, 105–110.

4.   Gopalakrishnan, J., Feistel, K., Friedrich, B.M., Grapin-Botton, A., Jurisch-Yaksi, N., Mass, E., Mick, D.U., Müller, R., May-Simera, H., Schermer, B., et al. (2023). Emerging principles of primary cilia dynamics in controlling tissue organization and function. The EMBO Journal *42*, e113891. https://doi.org/10.15252/embj.2023113891.

5.   Breslow, D.K., and Holland, A.J. (2019). Mechanism and Regulation of Centriole and Cilium Biogenesis. Annu. Rev. Biochem. *88*, 691–724. https://doi.org/10.1146/annurev-biochem-013118-111153.

6.   Anderson, R.G. (1972). The three-dimensional structure of the basal body from the rhesus monkey oviduct. J. Cell Biol. *54*, 246–265. https://doi.org/10.1083/jcb.54.2.246.

7.   Gibbons, I.R., and Grimstone, A.V. (1960). On flagellar structure in certain flagellates. J. Biophys. Biochem. Cytol. *7*, 697–716. https://doi.org/10.1083/jcb.7.4.697.

8.   O'Toole, E.T., Giddings, T.H., and Dutcher, S.K. (2007). Understanding microtubule organizing centers by comparing mutant and wild-type structures with electron tomography. Methods Cell Biol. *79*, 125–143. https://doi.org/10.1016/S0091-679X(06)79005-7.

9.   Ringo, D.L. (1967). Flagellar motion and fine structure of the flagellar apparatus in Chlamydomonas. J. Cell Biol. *33*, 543–571. https://doi.org/10.1083/jcb.33.3.543.

10.   Moran, A.L., Louzao-Martinez, L., Norris, D.P., Peters, D.J.M., and Blacque, O.E. (2023). Transport and barrier mechanisms that regulate ciliary compartmentalization and ciliopathies. Nat Rev Nephrol. https://doi.org/10.1038/s41581-023-00773-2.

11.   Hu, Q., Milenkovic, L., Jin, H., Scott, M.P., Nachury, M.V., Spiliotis, E.T., and Nelson, W.J. (2010). A septin diffusion barrier at the base of the primary cilium maintains ciliary membrane protein distribution. Science *329*, 436–439. https://doi.org/10.1126/science.1191054.

12.   Gilula, N.B., and Satir, P. (1972). The ciliary necklace. A ciliary membrane specialization. The Journal of cell biology *53*, 494–509. https://doi.org/10.1083/jcb.53.2.494.

13.   Yang, T.T., Chong, W.M., Wang, W.J., Mazo, G., Tanos, B., Chen, Z., Tran, T.M.N., Chen, Y.D., Weng, R.R., Huang, C.E., et al. (2018). Super-resolution architecture of mammalian centriole distal appendages reveals distinct blade and matrix functional components. Nature communication *9*, 2023. https://doi.org/10.1038/s41467-018-04469-1.

14.   Garcia, G., Raleigh, D.R., and Reiter, J.F. (2018). How the Ciliary Membrane Is Organized Inside-Out to Communicate Outside-In. Curr. Biol. *28*, R421–R434. https://doi.org/10.1016/j.cub.2018.03.010.

15.   Pazour, G.J., Agrin, N., Leszyk, J., and Witman, G.B. (2005). Proteomic analysis of a eukaryotic cilium. J. Cell Biol. *170*, 103–113. https://doi.org/10.1083/jcb.200504008.

16.   Gigante, E.D., and Caspary, T. (2020). Signaling in the primary cilium through the lens of the Hedgehog pathway. WIREs Dev. Biol. *9*, e377. https://doi.org/10.1002/wdev.377.

17.   Shinde, S.R., Nager, A.R., and Nachury, M.V. (2020). Ubiquitin chains earmark GPCRs for BBSome-mediated removal from cilia. J. Cell Biol. *219*. https://doi.org/10.1083/jcb.202003020.

18.   Ye, F., Nager, A.R., and Nachury, M.V. (2018). BBSome trains remove activated GPCRs from cilia by enabling passage through the transition zone. J. Cell Biol. *217*, 1847–1868. https://doi.org/10.1083/jcb.201709041.

19.   Spasic, M., and Jacobs, C.R. (2017). Lengthening primary cilia enhances cellular mechanosensitivity. Eur. Cell. Mater. *33*, 158–168. https://doi.org/10.22203/eCM.v033a12.

20.   Mick, D.U., Rodrigues, R.B., Leib, R.D., Adams, C.M., Chien, A.S., Gygi, S.P., and Nachury, M.V. (2015). Proteomics of Primary Cilia by Proximity Labeling. Dev. Cell *35*, 497–512. https://doi.org/10.1016/j.devcel.2015.10.015.

21.   Kohli, P., Hohne, M., Jungst, C., Bertsch, S., Ebert, L.K., Schauss, A.C., Benzing, T., Rinschen, M.M., and Schermer, B. (2017). The ciliary membrane-associated proteome reveals actin-binding proteins as key components of cilia. EMBO rep. *18*, 1521–1535. https://doi.org/10.15252/embr.201643846.

22.   May, E.A., Kalocsay, M., D'Auriac, I.G., Schuster, P.S., Gygi, S.P., Nachury, M.V., and Mick, D.U. (2021). Time-resolved proteomics profiling of the ciliary Hedgehog response. J. Cell Biol. *220*. https://doi.org/10.1083/jcb.202007207.

23.   Gupta, G.D., Coyaud, E., Goncalves, J., Mojarad, B.A., Liu, Y., Wu, Q., Gheiratmand, L., Comartin, D., Tkach, J.M., Cheung, S.W., et al. (2015). A Dynamic Protein Interaction

Landscape of the Human Centrosome-Cilium Interface. Cell *163*, 1484–1499. https://doi.org/10.1016/j.cell.2015.10.065.

24. Aslanyan, M.G., Doornbos, C., Diwan, G.D., Anvarian, Z., Beyer, T., Junger, K., Van Beersum, S.E.C., Russell, R.B., Ueffing, M., Ludwig, A., et al. (2023). A targeted multi-proteomics approach generates a blueprint of the ciliary ubiquitinome. Front. Cell Dev. Biol. *11*, 1113656. https://doi.org/10.3389/fcell.2023.1113656.

25. Uhlén, M., Fagerberg, L., Hallström, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A., Sivertsson, Å., Kampf, C., Sjöstedt, E., Asplund, A., et al. (2015). Tissue-based map of the human proteome. Science *347*, 1260419. https://doi.org/10.1126/science.1260419.

26. Thul, P.J., Åkesson, L., Wiking, M., Mahdessian, D., Geladaki, A., Ait Blal, H., Alm, T., Asplund, A., Björk, L., Breckels, L.M., et al. (2017). A subcellular map of the human proteome. Science *356*, eaal3321. https://doi.org/10.1126/science.aal3321.

27. Rambhatla, L., Chiu, C.-P., Glickman, R.D., and Rowe-Rendleman, C. (2002). In vitro differentiation capacity of telomerase immortalized human RPE cells. Invest Ophthalmol Vis Sci *43*, 1622–1630.

28. Wieser, M., Stadler, G., Jennings, P., Streubel, B., Pfaller, W., Ambros, P., Riedl, C., Katinger, H., Grillari, J., and Grillari-Voglauer, R. (2008). hTERT alone immortalizes epithelial cells of renal proximal tubules without changing their functional characteristics. American Journal of Physiology-Renal Physiology *295*, F1365–F1375. https://doi.org/10.1152/ajprenal.90405.2008.

29. Wolbank, S., Stadler, G., Peterbauer, A., Gillich, A., Karbiener, M., Streubel, B., Wieser, M., Katinger, H., Van Griensven, M., Redl, H., et al. (2009). *Telomerase* Immortalized Human Amnion- and Adipose-Derived Mesenchymal Stem Cells: Maintenance of Differentiation and Immunomodulatory Characteristics. Tissue Engineering Part A *15*, 1843–1854. https://doi.org/10.1089/ten.tea.2008.0205.

30. Paridaen, J.T., Wilsch-Brauninger, M., and Huttner, W.B. (2013). Asymmetric inheritance of centrosome-associated primary cilium membrane directs ciliogenesis after cell division. Cell *155*, 333–344. https://doi.org/10.1016/j.cell.2013.08.060.

31. Vasquez, S.S.V., van Dam, J., and Wheway, G. (2021). An updated SYSCILIA gold standard (SCGSv2) of known ciliary genes, revealing the vast progress that has been made in the cilia research field. Mol. Biol. Cell *32*, br13. https://doi.org/10.1091/mbc.E21-05-0226.

32. Lindskog, C., Norradin, F.H., Digre, A., Méar, L., Schutten, R., Lundberg, E., Hansen, J., and Uhlén, M. (2024). A high-resolution spatial map of cilia-associated proteins based on characterization of the human fallopian tube-specific proteome. Preprint, https://doi.org/10.21203/rs.3.rs-3914234/v1 https://doi.org/10.21203/rs.3.rs-3914234/v1.

33. Ishikawa, H., Thompson, J., Yates, J.R., and Marshall, W.F. (2012). Proteomic analysis of mammalian primary cilia. Curr. Biol. *22*, 414–419. https://doi.org/10.1016/j.cub.2012.01.031.

34. Karunakaran, K.B., Chaparala, S., Lo, C.W., and Ganapathiraju, M.K. (2020). Cilia interactome with predicted protein-protein interactions reveals connections to Alzheimer's disease, aging and other neuropsychiatric processes. Sci. Rep. *10*, 15629. https://doi.org/10.1038/s41598-020-72024-4.

35.   van Dam, T.J.P., Kennedy, J., van der Lee, R., de Vrieze, E., Wunderlich, K.A., Rix, S., Dougherty, G.W., Lambacher, N.J., Li, C., Jensen, V.L., et al. (2019). CiliaCarta: An integrated and validated compendium of ciliary genes. PloS one *14*, e0216705. https://doi.org/10.1371/journal.pone.0216705.

36.   Pir, M.S., Begar, E., Yenisert, F., Demirci, H.C., Korkmaz, M.E., Karaman, A., Tsiropoulou, S., Firat-Karalar, E.N., Blacque, O.E., Oner, S.S., et al. (2024). CilioGenics: an integrated method and database for predicting novel ciliary genes. Nucleic Acids Research, gkae554. https://doi.org/10.1093/nar/gkae554.

37.   Elliott, K.H., Balchand, S.K., Bonatto Paese, C.L., Chang, C.-F., Yang, Y., Brown, K.M., Rasicci, D.T., He, H., Thorner, K., Chaturvedi, P., et al. (2023). Identification of a heterogeneous and dynamic ciliome during embryonic development and cell differentiation. Development *150*, dev201237. https://doi.org/10.1242/dev.201237.

38.   Nevers, Y., Prasad, M.K., Poidevin, L., Chennen, K., Allot, A., Kress, A., Ripp, R., Thompson, J.D., Dollfus, H., Poch, O., et al. (2017). Insights into Ciliary Genes and Evolution from Multi-Level Phylogenetic Profiling. Mol. Biol. Evol. *34*, 2016–2034. https://doi.org/10.1093/molbev/msx146.

39.   Sigg, M.A., Menchen, T., Lee, C., Johnson, J., Jungnickel, M.K., Choksi, S.P., Garcia, G., Busengdal, H., Dougherty, G.W., Pennekamp, P., et al. (2017). Evolutionary Proteomics Uncovers Ancient Associations of Cilia with Signaling Pathways. Dev. Cell *43*, 744-762 e11. https://doi.org/10.1016/j.devcel.2017.11.014.

40.   Breslow, D.K., Hoogendoorn, S., Kopp, A.R., Morgens, D.W., Vu, B.K., Kennedy, M.C., Han, K., Li, A., Hess, G.T., Bassik, M.C., et al. (2018). A CRISPR-based screen for Hedgehog signaling provides insights into ciliary function and ciliopathies. Nat. Genet. https://doi.org/10.1038/s41588-018-0054-7.

41.   Pusapati, G.V., Kong, J.H., Patel, B.B., Krishnan, A., Sagner, A., Kinnebrew, M., Briscoe, J., Aravind, L., and Rohatgi, R. (2018). CRISPR Screens Uncover Genes that Regulate Target Cell Sensitivity to the Morphogen Sonic Hedgehog. Dev. Cell *44*, 113-129 e8. https://doi.org/10.1016/j.devcel.2017.12.003.

42.   Blackburn, K., Bustamante-Marin, X., Yin, W., Goshe, M.B., and Ostrowski, L.E. (2017). Quantitative Proteomic Analysis of Human Airway Cilia Identifies Previously Uncharacterized Proteins of High Abundance. J. Proteome Res. *16*, 1579–1592. https://doi.org/10.1021/acs.jproteome.6b00972.

43.   Sim, H.J., Yun, S., Kim, H.E., Kwon, K.Y., Kim, G.-H., Yun, S., Kim, B.G., Myung, K., Park, T.J., and Kwon, T. (2020). Simple Method To Characterize the Ciliary Proteome of Multiciliated Cells. J. Proteome Res. *19*, 391–400. https://doi.org/10.1021/acs.jproteome.9b00589.

44.   Baker, M.A., Naumovski, N., Hetherington, L., Weinberg, A., Velkov, T., and Aitken, R.J. (2013). Head and flagella subcompartmental proteomic analysis of human spermatozoa. Proteomics *13*, 61–74. https://doi.org/10.1002/pmic.201200350.

45.   Amaral, A., Castillo, J., Estanyol, J.M., Ballescà, J.L., Ramalho-Santos, J., and Oliva, R. (2013). Human Sperm Tail Proteome Suggests New Endogenous Metabolic Pathways. Molecular & Cellular Proteomics *12*, 330–342. https://doi.org/10.1074/mcp.M112.020552.

46.   Firat-Karalar, E.N., Sante, J., Elliott, S., and Stearns, T. (2014). Proteomic analysis of mammalian sperm cells identifies new components of the centrosome. Journal of Cell Science, jcs.157008. https://doi.org/10.1242/jcs.157008.

47.   Tai, L., Yin, G., Huang, X., Sun, F., and Zhu, Y. (2023). In-cell structural insight into the stability of sperm microtubule doublet. Cell Discov *9*, 116. https://doi.org/10.1038/s41421-023-00606-3.

48.   Zhou, L., Liu, H., Liu, S., Yang, X., Dong, Y., Pan, Y., Xiao, Z., Zheng, B., Sun, Y., Huang, P., et al. (2023). Structures of sperm flagellar doublet microtubules expand the genetic spectrum of male infertility. Cell *186*, 2897-2910.e19. https://doi.org/10.1016/j.cell.2023.05.009.

49.   Gui, M., Croft, J.T., Zabeo, D., Acharya, V., Kollman, J.M., Burgoyne, T., Höög, J.L., and Brown, A. (2022). SPACA9 is a lumenal protein of human ciliary singlet and doublet microtubules. Proc. Natl. Acad. Sci. U.S.A. *119*, e2207605119. https://doi.org/10.1073/pnas.2207605119.

50.   Firat-Karalar, E.N., Rauniyar, N., Yates, J.R., and Stearns, T. (2014). Proximity interactions among centrosome components identify regulators of centriole duplication. Curr. Biol. *24*, 664–670. https://doi.org/10.1016/j.cub.2014.01.067.

51.   Carden, S., Vitiello, E., Rosa E Silva, I., Holder, J., Quarantotti, V., Kishore, K., Roamio Franklin, V.N., D'Santos, C., Ochi, T., Van Breugel, M., et al. (2023). Proteomic profiling of centrosomes across multiple mammalian cell and tissue types by an affinity capture method. Developmental Cell *58*, 2393-2410.e9. https://doi.org/10.1016/j.devcel.2023.09.008.

52.   Andersen, J.S., Wilkinson, C.J., Mayor, T., Mortensen, P., Nigg, E.A., and Mann, M. (2003). Proteomic characterization of the human centrosome by protein correlation profiling. Nature *426*, 570–574. https://doi.org/10.1038/nature02166.

53.   Jakobsen, L., Vanselow, K., Skogs, M., Toyoda, Y., Lundberg, E., Poser, I., Falkenby, L.G., Bennetzen, M., Westendorf, J., Nigg, E.A., et al. (2011). Novel asymmetrically localizing components of human centrosomes identified by complementary proteomics methods: Centrosome proteomics. The EMBO Journal *30*, 1520–1535. https://doi.org/10.1038/emboj.2011.63.

54.   Gheiratmand, L., Coyaud, E., Gupta, G.D., Laurent, E.M., Hasegan, M., Prosser, S.L., Gonçalves, J., Raught, B., and Pelletier, L. (2019). Spatial and proteomic profiling reveals centrosome-independent features of centriolar satellites. The EMBO Journal *38*, e101109. https://doi.org/10.15252/embj.2018101109.

55.   Conkar, D., Culfa, E., Odabasi, E., Rauniyar, N., Yates, J.R., and Firat-Karalar, E.N. (2017). The centriolar satellite protein CCDC66 interacts with CEP290 and functions in cilium formation and trafficking. J. Cell Sci. *130*, 1450–1462. https://doi.org/10.1242/jcs.196832.

56.   Bennett, H.W., Gustavsson, A.K., Bayas, C.A., Petrov, P.N., Mooney, N., Moerner, W.E., and Jackson, P.K. (2020). Novel fibrillar structure in the inversin compartment of primary cilia revealed by 3D single-molecule superresolution microscopy. Molecular biology of the cell *31*, 619–639. https://doi.org/10.1091/mbc.E19-09-0499.

57.   Boehlke, C., Bashkurov, M., Buescher, A., Krick, T., John, A.-K., Nitschke, R., Walz, G., and Kuehn, E.W. (2010). Differential role of Rab proteins in ciliary trafficking: Rab23

regulates smoothened levels. J Cell Sci *123*, 1460–1467. https://doi.org/10.1242/jcs.058883.

58. Chi, S., Xie, G., Liu, H., Chen, K., Zhang, X., Li, C., and Xie, J. (2012). Rab23 negatively regulates Gli1 transcriptional factor in a Su(Fu)-dependent manner. Cell Signal *24*, 1222–1228. https://doi.org/10.1016/j.cellsig.2012.02.004.

59. Cho, J.H., Li, Z.A., Zhu, L., Muegge, B.D., Roseman, H.F., Lee, E.Y., Utterback, T., Woodhams, L.G., Bayly, P.V., and Hughes, J.W. (2022). Islet primary cilia motility controls insulin secretion. Sci. Adv. *8*, eabq8486. https://doi.org/10.1126/sciadv.abq8486.

60. van Rijssel, J., and van Buul, J.D. (2012). The many faces of the guanine-nucleotide exchange factor trio. Cell Adh Migr *6*, 482–487. https://doi.org/10.4161/cam.21418.

61. Ota, M., Gonja, H., Koike, R., and Fukuchi, S. (2016). Multiple-Localization and Hub Proteins. PLoS ONE *11*, e0156455. https://doi.org/10.1371/journal.pone.0156455.

62. Mulvey, C.M., Breckels, L.M., Geladaki, A., Britovšek, N.K., Nightingale, D.J.H., Christoforou, A., Elzek, M., Deery, M.J., Gatto, L., and Lilley, K.S. (2017). Using hyperLOPIT to perform high-resolution mapping of the spatial proteome. Nat Protoc *12*, 1110–1135. https://doi.org/10.1038/nprot.2017.026.

63. Singh, N., and Bhalla, N. (2020). Moonlighting Proteins. Annu. Rev. Genet. *54*, 265–285. https://doi.org/10.1146/annurev-genet-030620-102906.

64. Mukhopadhyay, S., Wen, X., Chih, B., Nelson, C.D., Lane, W.S., Scales, S.J., and Jackson, P.K. (2010). TULP3 bridges the IFT-A complex and membrane phosphoinositides to promote trafficking of G protein-coupled receptors into primary cilia. Genes Dev. *24*, 2180–2193. https://doi.org/10.1101/gad.1966210.

65. Han, S., Miyoshi, K., Shikada, S., Amano, G., Wang, Y., Yoshimura, T., and Katayama, T. (2019). TULP3 is required for localization of membrane-associated proteins ARL13B and INPP5E to primary cilia. Biochemical and Biophysical Research Communications *509*, 227–234. https://doi.org/10.1016/j.bbrc.2018.12.109.

66. Fagan, R.R., Lee, D.F., Geron, M., Scherrer, G., Von Zastrow, M., and Ehrlich, A.T. (2024). Selective targeting of mu opioid receptors to primary cilia. Cell Reports *43*, 114164. https://doi.org/10.1016/j.celrep.2024.114164.

67. Santagata, S., Boggon, T.J., Baird, C.L., Gomez, C.A., Zhao, J., Shan, W.S., Myszka, D.G., and Shapiro, L. (2001). G-Protein Signaling Through Tubby Proteins. Science *292*, 2041–2050. https://doi.org/10.1126/science.1061233.

68. Gnann, C., Cesnik, A., Sigaeva, A., Le, T., Sariyar, S., Mahdessian, D., Schutten, R., Raghavan, P., Leonetti, M.D., Lindskog, C., et al. (2024). Dissecting autonomous enzyme variability in single cells. Preprint, https://doi.org/10.1101/2024.10.03.616530 https://doi.org/10.1101/2024.10.03.616530.

69. Nguyen, T.D., Truong, M.E., and Reiter, J.F. (2022). The Intimate Connection Between Lipids and Hedgehog Signaling. Front. Cell Dev. Biol. *10*, 876815. https://doi.org/10.3389/fcell.2022.876815.

70. Kinnebrew, M., Iverson, E.J., Patel, B.B., Pusapati, G.V., Kong, J.H., Johnson, K.A., Luchetti, G., Eckert, K.M., McDonald, J.G., Covey, D.F., et al. (2019). Cholesterol accessibility at the ciliary membrane controls hedgehog signaling. eLife *8*, e50051. https://doi.org/10.7554/eLife.50051.

71. Wang, Y., Zhang, Y., Guo, X., Zheng, Y., Zhang, X., Feng, S., and Wu, H.-Y. (2023). CCP5 and CCP6 retain CP110 and negatively regulate ciliogenesis. BMC Biol *21*, 124. https://doi.org/10.1186/s12915-023-01622-1.

72. Adada, M.M., Orr-Gandy, K.A., Snider, A.J., Canals, D., Hannun, Y.A., Obeid, L.M., and Clarke, C.J. (2013). Sphingosine kinase 1 regulates tumor necrosis factor-mediated RANTES induction through p38 mitogen-activated protein kinase but independently of nuclear factor κB activation. J Biol Chem *288*, 27667–27679. https://doi.org/10.1074/jbc.M113.489443.

73. Maceyka, M., Sankala, H., Hait, N.C., Le Stunff, H., Liu, H., Toman, R., Collier, C., Zhang, M., Satin, L.S., Merrill, A.H., et al. (2005). SphK1 and SphK2, sphingosine kinase isoenzymes with opposing functions in sphingolipid metabolism. J Biol Chem *280*, 37118–37129. https://doi.org/10.1074/jbc.M502207200.

74. Alvarez, S.E., Harikumar, K.B., Hait, N.C., Allegood, J., Strub, G.M., Kim, E.Y., Maceyka, M., Jiang, H., Luo, C., Kordula, T., et al. (2010). Sphingosine-1-phosphate is a missing cofactor for the E3 ubiquitin ligase TRAF2. Nature *465*, 1084–1088. https://doi.org/10.1038/nature09128.

75. Helenius, A., and Ritter, C. (2000). Recognition of local glycoprotein misfolding by the ER folding sensor UDP-glucose:glycoprotein glucosyltransferase. Nat. Struct Biol. *7*, 278–280. https://doi.org/10.1038/74035.

76. Wang, N., Seko, A., Takeda, Y., and Ito, Y. (2020). Glycan dependent refolding activity of ER glucosyltransferase (UGGT). Biochimica et Biophysica Acta (BBA) - General Subjects *1864*, 129709. https://doi.org/10.1016/j.bbagen.2020.129709.

77. Kinnebrew, M., Luchetti, G., Sircar, R., Frigui, S., Viti, L.V., Naito, T., Beckert, F., Saheki, Y., Siebold, C., Radhakrishnan, A., et al. (2021). Patched 1 reduces the accessibility of cholesterol in the outer leaflet of membranes. eLife *10*, e70504. https://doi.org/10.7554/eLife.70504.

78. Palander, O., El-Zeiry, M., and Trimble, W.S. (2017). Uncovering the Roles of Septins in Cilia. Front. Cell Dev. Biol. *5*. https://doi.org/10.3389/fcell.2017.00036.

79. Hansen, J.N., Rassmann, S., Stuven, B., Jurisch-Yaksi, N., and Wachten, D. (2021). CiliaQ: a simple, open-source software for automated quantification of ciliary morphology and fluorescence in 2D, 3D, and 4D images. Eur. Phys. J. E Soft Matter *44*, 18. https://doi.org/10.1140/epje/s10189-021-00031-y.

80. Bishop, G.A., Berbari, N.F., Lewis, J., and Mykytyn, K. (2007). Type III adenylyl cyclase localizes to primary cilia throughout the adult mouse brain. J. Comp. Neurol. *505*, 562–571. https://doi.org/10.1002/cne.21510.

81. Hua, K., and Ferland, R.J. (2017). Fixation methods can differentially affect ciliary protein immunolabeling. Cilia *6*, 5. https://doi.org/10.1186/s13630-017-0045-9.

82. Gigante, E.D., Taylor, M.R., Ivanova, A.A., Kahn, R.A., and Caspary, T. (2020). ARL13B regulates Sonic hedgehog signaling from outside primary cilia. eLife *9*, e50434. https://doi.org/10.7554/eLife.50434.

83. Ferent, J., Constable, S., Gigante, E.D., Yam, P.T., Mariani, L.E., Legué, E., Liem, K.F., Caspary, T., and Charron, F. (2019). The Ciliary Protein Arl13b Functions Outside of the Primary Cilium in Shh-Mediated Axon Guidance. Cell Reports *29*, 3356-3366.e3. https://doi.org/10.1016/j.celrep.2019.11.015.

84. Janke, C., and Montagnac, G. (2017). Causes and Consequences of Microtubule Acetylation. Curr. Biol. *27*, R1287–R1292. https://doi.org/10.1016/j.cub.2017.10.044.

85. Zlotina, A., Barashkova, S., Zhuk, S., Skitchenko, R., Usoltsev, D., Sokolnikova, P., Artomov, M., Alekseenko, S., Simanova, T., Goloborodko, M., et al. (2024). Characterization of pathogenic genetic variants in Russian patients with primary ciliary dyskinesia using gene panel sequencing and transcript analysis. Orphanet J Rare Dis *19*, 310. https://doi.org/10.1186/s13023-024-03318-3.

86. Jat, K.R., Faruq, M., Jindal, S., Bari, S., Soni, A., Sharma, P., Mathews, S., Shamim, U., Ahuja, V., Uppilli, B., et al. (2024). Genetics of 67 patients of suspected primary ciliary dyskinesia from India. Clinical Genetics, cge.14590. https://doi.org/10.1111/cge.14590.

87. Schultz, R., Elenius, V., Fassad, M.R., Freke, G., Rogers, A., Shoemark, A., Koistinen, T., Mohamed, M.A., Lim, J.S.Y., Mitchison, H.M., et al. (2022). CFAP300 mutation causing primary ciliary dyskinesia in Finland. Front. Genet. *13*, 985227. https://doi.org/10.3389/fgene.2022.985227.

88. Yiallouros, P.K., Kouis, P., Kyriacou, K., Evriviadou, A., Anagnostopoulou, P., Matthaiou, A., Tsiolakis, I., Pirpa, P., Michailidou, K., Potamiti, L., et al. (2021). Implementation of multigene panel NGS diagnosis in the national primary ciliary dyskinesia cohort of Cyprus: An island with a high disease prevalence. Human Mutation *42*. https://doi.org/10.1002/humu.24196.

89. Aprea, I., Raidt, J., Höben, I.M., Loges, N.T., Nöthe-Menchen, T., Pennekamp, P., Olbrich, H., Kaiser, T., Biebach, L., Tüttelmann, F., et al. (2021). Defects in the cytoplasmic assembly of axonemal dynein arms cause morphological abnormalities and dysmotility in sperm cells leading to male infertility. PLoS Genet *17*, e1009306. https://doi.org/10.1371/journal.pgen.1009306.

90. Zietkiewicz, E., Bukowy-Bieryllo, Z., Rabiasz, A., Daca-Roszak, P., Wojda, A., Voelkel, K., Rutkiewicz, E., Pogorzelski, A., Rasteiro, M., and Witt, M. (2019). *CFAP300*: Mutations in Slavic Patients with Primary Ciliary Dyskinesia and a Role in Ciliary Dynein Arms Trafficking. Am J Respir Cell Mol Biol *61*, 440–449. https://doi.org/10.1165/rcmb.2018-0260OC.

91. Fassad, M.R., Shoemark, A., Le Borgne, P., Koll, F., Patel, M., Dixon, M., Hayward, J., Richardson, C., Frost, E., Jenkins, L., et al. (2018). C11orf70 Mutations Disrupting the Intraflagellar Transport-Dependent Assembly of Multiple Axonemal Dyneins Cause Primary Ciliary Dyskinesia. The American Journal of Human Genetics *102*, 956–972. https://doi.org/10.1016/j.ajhg.2018.03.024.

92. McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University (Baltimore, MD) Online Mendelian Inheritance in Man, OMIM®. https://omim.org/.

93. Sakurai, T. (2012). The role of NrCAM in neural development and disorders--beyond a simple glue in the brain. Mol Cell Neurosci *49*, 351–363. https://doi.org/10.1016/j.mcn.2011.12.002.

94. Yokoyama, K., Tezuka, T., Kotani, M., Nakazawa, T., Hoshina, N., Shimoda, Y., Kakuta, S., Sudo, K., Watanabe, K., Iwakura, Y., et al. (2011). NYAP: a phosphoprotein family that links PI3K to WAVE1 signalling in neurons: NYAPs link PI3K to WAVE1 in neurons. The EMBO Journal *30*, 4739–4754. https://doi.org/10.1038/emboj.2011.348.

95.   Davare, M.A., Fortin, D.A., Saneyoshi, T., Nygaard, S., Kaech, S., Banker, G., Soderling, T.R., and Wayman, G.A. (2009). Transient Receptor Potential Canonical 5 Channels Activate Ca $^{2+}$/Calmodulin Kinase Iγ to Promote Axon Formation in Hippocampal Neurons. J. Neurosci. *29*, 9794–9808. https://doi.org/10.1523/JNEUROSCI.1544-09.2009.

96.   Lucchesi, W., Mizuno, K., and Giese, K.P. (2011). Novel insights into CaMKII function and regulation during memory formation. Brain Research Bulletin *85*, 2–8. https://doi.org/10.1016/j.brainresbull.2010.10.009.

97.   Araki, T., and Wakatsuki, S. (2019). Regulation of neuronal/axonal degeneration by ZNRF1 ubiquitin ligase. Neuroscience Research *139*, 21–25. https://doi.org/10.1016/j.neures.2018.07.008.

98.   Yoshida, K., Watanabe, M., and Hatakeyama, S. (2009). ZNRF1 interacts with tubulin and regulates cell morphogenesis. Biochemical and Biophysical Research Communications *389*, 506–511. https://doi.org/10.1016/j.bbrc.2009.09.011.

99.   Hu, H.-B., Song, Z.-Q., Song, G.-P., Li, S., Tu, H.-Q., Wu, M., Zhang, Y.-C., Yuan, J.-F., Li, T.-T., Li, P.-Y., et al. (2021). LPA signaling acts as a cell-extrinsic mechanism to initiate cilia disassembly and promote neurogenesis. Nat Commun *12*, 662. https://doi.org/10.1038/s41467-021-20986-y.

100.   Loskutov, Y.V., Griffin, C.L., Marinak, K.M., Bobko, A., Margaryan, N.V., Geldenhuys, W.J., Sarkaria, J.N., and Pugacheva, E.N. (2018). LPA signaling is regulated through the primary cilium: a novel target in glioblastoma. Oncogene *37*, 1457–1471. https://doi.org/10.1038/s41388-017-0049-3.

101.   Walia, V., Cuenca, A., Vetter, M., Insinna, C., Perera, S., Lu, Q., Ritt, D.A., Semler, E., Specht, S., Stauffer, J., et al. (2019). Akt Regulates a Rab11-Effector Switch Required for Ciliogenesis. Developmental Cell *50*, 229-246.e7. https://doi.org/10.1016/j.devcel.2019.05.022.

102.   Lai, S.-L., Yao, W.-L., Tsao, K.-C., Houben, A.J.S., Albers, H.M.H.G., Ovaa, H., Moolenaar, W.H., and Lee, S.-J. (2012). Autotaxin/Lpar3 signaling regulates Kupffer's vesicle formation and left-right asymmetry in zebrafish. Development *139*, 4439–4448. https://doi.org/10.1242/dev.081745.

103.   Lin, Y.-N., Audira, G., Malhotra, N., Ngoc Anh, N.T., Siregar, P., Lu, J.-H., Lee, H., and Hsiao, C.-D. (2020). A Novel Function of the Lysophosphatidic Acid Receptor 3 (LPAR3) Gene in Zebrafish on Modulating Anxiety, Circadian Rhythm Locomotor Activity, and Short-Term Memory. IJMS *21*, 2837. https://doi.org/10.3390/ijms21082837.

104.   Kusano, S., and Raab-Traub, N. (2002). I-mfa domain proteins interact with Axin and affect its regulation of the Wnt and c-Jun N-terminal kinase signaling pathways. Mol Cell Biol *22*, 6393–6405. https://doi.org/10.1128/MCB.22.18.6393-6405.2002.

105.   Goto, T., Matsuzawa, J., Iemura, S.-I., Natsume, T., and Shibuya, H. (2016). WDR26 is a new partner of Axin1 in the canonical Wnt signaling pathway. FEBS Lett *590*, 1291–1303. https://doi.org/10.1002/1873-3468.12180.

106.   Liu, W., Rui, H., Wang, J., Lin, S., He, Y., Chen, M., Li, Q., Ye, Z., Zhang, S., Chan, S.C., et al. (2006). Axin is a scaffold protein in TGF-beta signaling that promotes degradation of Smad7 by Arkadia. EMBO J *25*, 1646–1658. https://doi.org/10.1038/sj.emboj.7601057.

107. Zhang, K., Da Silva, F., Seidl, C., Wilsch-Bräuninger, M., Herbst, J., Huttner, W.B., and Niehrs, C. (2023). Primary cilia are WNT-transducing organelles whose biogenesis is controlled by a WNT-PP1 axis. Developmental Cell *58*, 139-154.e8. https://doi.org/10.1016/j.devcel.2022.12.006.

108. Stranneheim, H., Lagerstedt-Robinson, K., Magnusson, M., Kvarnung, M., Nilsson, D., Lesko, N., Engvall, M., Anderlid, B.-M., Arnell, H., Johansson, C.B., et al. (2021). Integration of whole genome sequencing into a healthcare setting: high diagnostic rates across multiple clinical entities in 3219 rare disease patients. Genome Med *13*, 40. https://doi.org/10.1186/s13073-021-00855-5.

109. Ek, M., Nilsson, D., Engvall, M., Malmgren, H., Thonberg, H., Pettersson, M., Anderlid, B.-M., Hammarsjö, A., Helgadottir, H.T., Arnardottir, S., et al. (2023). Genome sequencing with comprehensive variant calling identifies structural variants and repeat expansions in a large fraction of individuals with ataxia and/or neuromuscular disorders. Front. Neurol. *14*, 1170005. https://doi.org/10.3389/fneur.2023.1170005.

110. Wachten, D., and Mick, D.U. (2021). Signal transduction in primary cilia - analyzing and manipulating GPCR and second messenger signaling. Pharmacol. Ther. *224*, 107836. https://doi.org/10.1016/j.pharmthera.2021.107836.

111. Alvarez, L., Friedrich, B.M., Gompper, G., and Kaupp, U.B. (2014). The computational sperm cell. Trends in cell biology *24*, 198–207. https://doi.org/10.1016/j.tcb.2013.10.004.

112. Sheu, S.-H., Upadhyayula, S., Dupuy, V., Pang, S., Deng, F., Wan, J., Walpita, D., Pasolli, H.A., Houser, J., Sanchez-Martinez, S., et al. (2022). A serotonergic axon-cilium synapse drives nuclear signaling to alter chromatin accessibility. Cell *185*, 3390-3407.e18. https://doi.org/10.1016/j.cell.2022.07.026.

113. Lázár, E., Mauron, R., Andrusivová, Ž., Foyer, J., He, M., Larsson, L., Shakari, N., Salas, S.M., Avenel, C., Sariyar, S., et al. (2024). Spatial Dynamics of the Developing Human Heart. Preprint, https://doi.org/10.1101/2024.03.12.584577 https://doi.org/10.1101/2024.03.12.584577.

114. Durinck, S., Spellman, P.T., Birney, E., and Huber, W. (2009). Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt. Nat Protoc *4*, 1184–1191. https://doi.org/10.1038/nprot.2009.97.

115. Nilsson, P., Paavilainen, L., Larsson, K., Ödling, J., Sundberg, M., Andersson, A.-C., Kampf, C., Persson, A., Szigyarto, C.A.-K., Ottosson, J., et al. (2005). Towards a human proteome atlas: High-throughput generation of mono-specific antibodies for tissue profiling. Proteomics *5*, 4327–4337. https://doi.org/10.1002/pmic.200500072.

116. Uhlen, M., Bandrowski, A., Carr, S., Edwards, A., Ellenberg, J., Lundberg, E., Rimm, D.L., Rodriguez, H., Hiltke, T., Snyder, M., et al. (2016). A proposal for validation of antibodies. Nat Methods *13*, 823–827. https://doi.org/10.1038/nmeth.3995.

117. Edfors, F., Hober, A., Linderbäck, K., Maddalo, G., Azimi, A., Sivertsson, Å., Tegel, H., Hober, S., Szigyarto, C.A.-K., Fagerberg, L., et al. (2018). Enhanced validation of antibodies for research applications. Nat Commun *9*, 4130. https://doi.org/10.1038/s41467-018-06642-y.

118. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., et al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. The Innovation *2*, 100141. https://doi.org/10.1016/j.xinn.2021.100141.

119. Gu, Z., and Hübschmann, D. (2023). *SimplifyEnrichment* : A Bioconductor Package for Clustering and Visualizing Functional Enrichment Results. Genomics, Proteomics & Bioinformatics *21*, 190–202. https://doi.org/10.1016/j.gpb.2022.04.008.

120. Danielsson, A., Pontén, F., Fagerberg, L., Hallström, B.M., Schwenk, J.M., Uhlén, M., Korsgren, O., and Lindskog, C. (2014). The Human Pancreas Proteome Defined by Transcriptomics and Antibody-Based Profiling. PLoS ONE *9*, e115421. https://doi.org/10.1371/journal.pone.0115421.

121. Uhlen, M., Oksvold, P., Fagerberg, L., Lundberg, E., Jonasson, K., Forsberg, M., Zwahlen, M., Kampf, C., Wester, K., Hober, S., et al. (2010). Towards a knowledge-based Human Protein Atlas. Nat Biotechnol *28*, 1248–1250. https://doi.org/10.1038/nbt1210-1248.

122. Kampf, C., Olsson, I., Ryberg, U., Sjöstedt, E., and Pontén, F. (2012). Production of Tissue Microarrays, Immunohistochemistry Staining and Digitalization Within the Human Protein Atlas. JoVE, 3620. https://doi.org/10.3791/3620.

123. Martinez Casals, A. (2023). Indirect immunofluorescence - tissue staining in TMA and whole tissue FFPE sections v1. Preprint, https://doi.org/10.17504/protocols.io.6qpvr4w6ogmk/v1.