1 **The Zoonotic Potential of Poultry-Associated Nitrofurantoin-Resistant**

2 ***Escherichia coli***

5 Jordan E Sealey[1], Beth Astley[1] & Matthew B Avison[1*]

7 [1]School of Cellular & Molecular Medicine, University of Bristol. United Kingdom

8 *Correspondence to Matthew B. Avison, bimba@bristol.ac.uk

Abstract

Escherichia coli is the most common cause of urinary tract infection (UTI) in humans. Nitrofurantoin is the most frequent UTI therapy in the United Kingdom. Here we identify poultry as a potential zoonotic reservoir of nitrofurantoin resistant (NFT-R) and pre-resistant (one mutation away from NFT-R) *E. coli*. We found NFT-R *E. coli* excreted by dogs fed a raw meat diet, as well as NFT-R and pre-resistant *E. coli* contaminating chicken meat sold for human consumption and chicken-based raw dog food. By whole genome sequence analysis, with reference to additional data from public repositories, we identified NFT-R or pre-resistant *E. coli* clones. These were ST919/8874, ST665, ST6805, ST69, ST752 and ST7529, which were dominated by isolates from poultry sources in Europe, though the ST752 clone is globally disseminated. Five of these clones included isolates from human clinical samples, of which several were confirmed urinary isolates. Our findings have important implications for public health and domestic hygiene, particularly for people being treated for UTI. They also increase concerns about the roles of raw-fed dogs as vectors for the entry of resistant *E. coli* into the home. It is likely that nitrofuran use in poultry production was the initial driver for the emergence of NFT-R on poultry farms. Therefore, whilst NFT-R *E. coli* sample-level positivity in poultry meat was far lower than resistance to more recently used fluoroquinolones, NFT-R and pre-resistant *E. coli* appear to have persisted among poultry flocks despite 30 years of zero legal use in Europe.

Introduction

Nitrofurantoin is the recommended first line treatment for uncomplicated urinary tract infection (UTI) in adults in the United Kingdom, most of which are caused by *Escherichia coli*.[1] Despite widespread use, nitrofurantoin resistance is rare.[2]

The ability of nitrofurantoin to kill relies on two oxygen-insensitive nitroreductases produced in the bacterium, NfsA and NfsB, which reduce nitrofurantoin to liberate highly reactive intermediates.[3] Nitrofurantoin resistance occurs predominately by loss of function mutations, deletions or insertional inactivation of both *nfsA* and *nfsB*, and in rare cases, by mutation of *ribE* or carriage of the plasmid-mediated *oqxAB* efflux pump genes.[2,4-6] We have recently identified the presence of nitrofurantoin "pre-resistant" urinary *E. coli*, where one gene (more commonly *nfsA*) is disrupted, but the other remains intact.[2]

The requirement for mutations in two genes – *nfsA* and *nfsB* – may form a barrier to nitrofurantoin resistance emerging among *E. coli* in an infection[7] and there is currently no evidence for nitrofurantoin-resistant (NFT-R) *E. coli* clones circulating in the human population.[2] If such clones emerge, this could spark a rise in NFT-R UTIs. The primary aim of this work, therefore, was to consider the possibility of animal reservoirs of NFT-R or pre-resistant *E. coli* with the potential to transfer into the human population.

Nitrofurans were widely used in food producing animals, particularly poultry, until the 1990s when the European Union banned this due to the risks associated with toxic chemical residues in meat. Similar bans are in place in the USA, Australia, the Philippines, Thailand and Brazil.[8] In some countries, nitrofurantoin is used to treat ornamental birds and fish, and as an off-label therapy for recurrent UTI in dogs, cats

and horses[9], however, in the United Kingdom, nitrofurantoin is not licensed for use in

any animal.[10]

Surveys of NFT-R bacteria from animals are scarce, and in the few studies that have

considered NFT-R *E. coli*, resistance was very rare or non-existent.[11-14], but

emergence of NFT-R *E. coli* in dogs during treatment of UTI has been reported.[15]

We have established a 50 x 50 km study region in the South West of England in

which we have sequenced resistant *E. coli* from human infections[2,16-18], and from the

faeces of farmed[17,19], companion[20,21], and zoo animals.[22] In the work reported here,

we investigated whether NFT-R *E. coli* are excreted by dogs in our study region or

are found on meat being sold in stores in the area, and used whole genome

sequence (WGS) data to test whether these NFT-R *E. coli* are shared across One

Health compartments, including human infections. In so doing we have identified that

there is a zoonotic reservoir of NFT-R and pre-resistant *E. coli* in poultry and in dogs

fed raw meat.

Results

NFT-R E. coli excreted by dogs being fed raw (uncooked) meat.

We tested faecal samples from 600 healthy dogs living in our 50 x 50 km study

area.[20] Two dogs living in the city of Bristol excreted NFT-R *E. coli* at detectable

levels. Both NFT-R *E. coli* had loss of function mutations in *nfsA* and *nfsB*; one was

ST6805 and carried horizontally-acquired genes conferring resistance to multiple

antibiotics, the other was ST919 and did not carry any acquired resistance genes

(**Table 1**).

We collected owner-completed survey data about the lives of the 600 dogs.[20,21]

Since only two dogs excreted NFT-R *E. coli* it is not possible to be confident about

any statistical assessment of associations between excretion and any lifestyle

practice, as determined from the owner-completed surveys. Nonetheless, we noted

(**Table S1**) that both dogs were fed raw meat at the time of faecal sampling, which

was a practice observed for only 31/297 dogs living in the city of Bristol (Fisher Exact

p=0.011).

NFT-R E. coli in uncooked chicken meat and chicken-based commercial raw dog

food.

In the United Kingdom, raw meat fed to dogs is sourced as commercial raw dog food

(RDF) or meat sold primarily for human consumption (once cooked).[23] Accordingly,

to investigate whether meat potentially being fed to dogs in the city of Bristol was

contaminated with NFT-R *E. coli* we tested 73 samples of meat from across the city:

15 packets each of beef, chicken and lamb, and 13 of pork purchased from large

chain grocery stores, and 15 packets of chicken-based commercial RDF of various

brands.[24]

Two out of 58 (3%) samples of meat from large chain grocery stores tested positive

for NFT-R *E. coli*; both were chicken meat. WGS showed that the NFT-R *E. coli* had

loss of function mutations in *nfsA* and *nfsB* and were ST665 (carrying multiple

antibiotic resistance genes) and ST8874 (carrying none) (**Table 1**). One out of fifteen

(7%) packets of chicken-based commercial RDF was positive for NFT-R *E. coli.*

WGS showed this to be ST8874, the same ST found in one packet of chicken meat.

Despite their different origins, the two isolates carried the same mutations in *nfsA*

and *nfsB* and both lacked mobile resistance genes (**Table 1**).

Poultry-associated NFT-R and NFT pre-resistant E. coli clones found across multiple

countries and One Health compartments, including human clinical specimens

The NFT-R ST8874 (n=3) and ST919 (n=4) isolates found on meat or excreted by

one dog in Bristol had the same mutations in *nfsA* and in *nfsB* and none of them

carried acquired resistance genes (**Table 1**). We hypothesized that these STs are

closely related and using multi-locus sequence typing found all except one of the

alleles (*adk*38 in ST8874 versus *adk*861 in ST919) were identical across the two

STs.

There was WGS data for only one ST8874 isolate on Enterobase (search date, 2[nd]

July 2024) and 34 of ST919. The ST8874 isolate (Accession SAMN12241725) came

from a chicken yoke sack infection in Czechia[25] and has the same *nfsA* and *nfsB*

mutations found in the Bristol ST919/ST8874 isolates (**Table 1**). Of the 34

Enterobase ST919 isolates, 56% were from poultry or wildfowl sources and three

had *nfsA* and *nfsB* mutations identical to the Bristol ST919/ST8874 isolates (**Table

1**). Two were of poultry origin, one from the United Kingdom (SAMN11444779) and

one from Canada (SAMN32247434). The third (SAMEA7025515) is a published

NFT-R *E. coli* from a human UTI from London, United Kingdom,[6] which is

approximately 200 km from Bristol.

The Bristol ST8874 isolates and the Czech chicken isolate were very closely related

to each other (<65 SNPs across a core genome alignment), and to the ST919 United

Kingdom chicken isolate on Enterobase (<120 SNPs) which was only 57 SNPs

different from the NFT-R human urinary isolate from London. The remaining ST919

isolates were variable in their relatedness with the most diverse being from a Bristol

dog (**Fig. 1**).

Among 94 ST665 sequences on Enterobase there was a source bias of 41% poultry

and 10% human, including two isolates from the United Kingdom. One of these

(SAMN21845675) had the same *nfsA* and *nfsB* mutations as the NFT-R ST665

isolate from uncooked chicken meat in Bristol, and the same seven mobile

resistance genes (**Table 1**), with an additional *dfrA1*. The two isolates were 2394

SNPs, apart.

Of the seven other human ST665 isolates on Enterobase, multiple isolates from

across Europe and into the Middle East had loss of function mutations in *nfsA* but

wild-type *nfsB*, and so are considered nitrofurantoin pre-resistant.[2] In poultry ST665

isolates, *nfsA* loss but with a wild type *nfsB* was also seen in multiple countries, with,

in some cases, identical *nfsA* mutations to those seen in human isolates (**Table S2**).

Among four ST6805 isolates on Enterobase, one Danish poultry isolate

(SAMEA112328101) shared the same *nfsA* mutation as a Bristol dog isolate (**Table**

1), but it had a wild type *nfsB*, hence is also considered pre-resistant. These two

isolates are 2281 SNPs apart.

Based on the above findings, we suspected that there might be additional

nitrofurantoin pre-resistant clones present on chicken meat being sold in Bristol. To

test this, we sequenced 92 nitrofurantoin susceptible (NFT-S) *E. coli* from RDF and

76 from uncooked chicken meat, cultured from the same samples as the NFT-R

isolates (above) and found 17 isolates from 12/15 samples of chicken meat and 10

isolates from 4/15 samples of chicken-based RDF, had obvious loss of function in

nfsA but wild-type *nfsB*, and so are defined as pre-resistant. These isolates,

accounting for 16.1% of all NFT-S isolates from chicken meat or RDF sequenced,

spanned 11 STs (**Table S3**). They included ST665 isolates from two chicken and one

RDF samples with identical *nfsA* mutations, and pairwise SNP distances ranging

from 84 to 880, reminiscent of the NFT-R clones described above. Furthermore, pre-

resistant ST69 isolates, with identical *nfsA* mutations from six chicken meat and one

RDF samples were identified with SNP distances of 10 to 41.

Evidence of a more longstanding nitrofuoranotin pre-resistant clone having *nfsA* loss

of function was also seen for ST752, where isolates from different samples of

chicken meat in Bristol were separated by in some cases as little as 88 SNPs (**Table**

S4). Of ST752 isolates on Enterobase (n=503), the majority were from poultry

(33.0%) or human (28.2%) sources. Of these, 91.9% of human isolates and 84.9% of

poultry isolates carried the same *nfsA* mutation as also seen in the Bristol chicken

meat isolates suggesting they are nitrofurantoin pre-resistant. We selected ST752

nfsA mutant Enterobase isolates from poultry and humans (n=10 for each) for

comparison with the six isolates from chicken meat in Bristol. Two human isolates,

one from the United Kingdom and the other from the Netherlands were 43 and 61

SNPs different, respectively, from one of the Bristol chicken meat isolates (**Table**

S4). Furthermore, there were four incidences where ST752 isolates from humans

were <100 SNPs from Enterobase poultry isolates, with the closest pair being 48

SNPs apart (**Table S4**).

Finally, for ST7529, a human clinical Enterobase isolate from the United Kingdom

was found to be 103 SNPs different from the two Bristol RDF isolates, which were

themselves identical. All carried the same *nfsA* loss of function mutation (**Table S3**).

Discussion

Because dogs in the United Kingdom are almost never treated with nitrofurantoin, we were surprised to find NFT-R *E. coli* in faeces excreted by two dogs out of a cohort of 600 based on testing one faecal sample per dog with a limit of detection of approximately 1000 viable bacteria per gramme of faeces. The NFT-R *E. coli* are clonally related to those found in poultry meat and poultry farm environment samples in the United Kingdom, Canada and Czechia. Given that we also found NFT-R *E. coli* in uncooked chicken meat, and chicken-based RDF purchased in the same city, the most likely explanation for these dogs excreting poultry-associated NFT-R *E. coli* is the fact – reported by their owners – that they were being raw fed at the time of sampling. Whilst we have shown compelling evidence that feeding raw meat to dogs is associated with them excreting fluoroquinolone resistant *E. coli*,[21] the evidence for NFT-R *E. coli* presented here is much weaker. This may be related to the relative degrees of contamination of meat with *E. coli* resistant to these different agents. For example, in parallel, we identified here 10% (3/30) of chicken meat or chicken-based RDF samples were positive for NFT-R *E. coli*, but 47% (14/30) of these same samples were positive for fluoroquinolone-resistant *E. coli*.[24]

A number of studies have reported sharing of resistant *E. coli* between dogs and their owners[26-29], so it is possible that dogs excreting NFT-R *E. coli* may pass these on to their owners. This is in addition to any associated risks of accidental ingestion when handling and preparing RDF or other uncooked poultry, which we have shown here can be contaminated with NFT-R or pre-resistant *E. coli*. Accordingly, it would seem prudent for those at risk of UTI, and particularly those who are taking

nitrofurantoin, to be cautious when handling uncooked meat or interacting with pets

fed uncooked meat, and practice enhanced hygiene regimes. This is clearly sound

advice for people taking other antibiotics, and particularly ciprofloxacin, but since the

number of nitrofurantoin courses dispensed in the community in some countries is

very high, and since resistance rates among urinary *E. coli* from humans are very

low, the additional zoonotic burden exposed here is potentially important. Perhaps

the most significant example of zoonotic antibiotic resistance transmission to be

uncovered since the identification of porcine-associated *mcr*-positive *E. coli* with

reduced colistin susceptibility in China.[30]

Whilst nitrofuran use in poultry production in many countries globally has been

banned for decades due to toxicity concerns if found in meat,[11] it is likely that NFT-R

and pre-resistant poultry-associated *E. coli* identified here evolved a long time ago.

This is apparent due to the range of SNP distances between isolates from the

various clones reported here.

It still remains unclear how prevalent NFT-R or NFT pre-resistant *E. coli* are in meat

sold for human consumption around the world, and whether other poultry-associated

clones exist, but the findings of this study should stimulate interest in answering that

question through widespread surveillance in a number of One Health compartments.

Methods

Canine faecal sample collection and processing

Ethical approval was obtained from the Faculty of Health Research Ethics

Committee, University of Bristol (ref: 89282) alongside Ethical Approval of an

Investigation Involving Animals (ref: UB/19/057). Details of canine faecal sample

collection are published[20,21], but briefly, 303 rural dogs and 297 urban dogs were

recruited at local dog-walking sites between September 2019 and September 2020.

All owners gave informed consent to participate. Faecal samples were collected and

a standardized questionnaire was provided to collect demographic data and

variables chosen as being potentially associated with carriage of ABR *E. coli* in dogs.

Samples were refrigerated and processed within 48 h. Faecal samples were weighed

(0.1–0.5 g) and PBS was added at 10 mL/g before vortexing and mixing with an

equal volume of 50% sterile glycerol. Samples were stored at -75°C.

Meat sample collection and processing

Details of meat sample collection and processing are published[24], but briefly, we

sampled beef, chicken, lamb and pork meat collected from large chain grocery

stores in the city of Bristol between October and November 2022, totalling 15

samples per type of meat, except for pork where due to availability only 13 were

collected. Samples were either processed on the day or kept at 4°C and processed

within the use by date. For raw chicken sold as RDF, one of each brand sold across

five RDF stores in Bristol was purchased. A total of 15 packets were collected in

September 2023, covering 11 brands of RDF.

Under sterile conditions, all meat and RDF samples were weighed to 200g and

placed into a stomacher bag with 200 mL of EC broth (Oxoid LTD). Bags were

sealed and incubated for 5 hours at 37°C, 180 rpm. The liquid culture was aliquot

into Eppendorf tubes with 25% (v/v) glycerol and stored at -75°C.

Microbiology

Twenty microlitres of each sample was spread onto Tryptone Bile X-Glucuronide

agar plates (Sigma) containing Nitrofurantoin (64 mg/L) based on EUCAST

resistance breakpoints for human infections[31] and incubated overnight at 37°C.

Putative NFT-R *E. coli* were re-streaked to confirm resistance. Isolates were stored

on Cryobeads at -75°C. Resistance was confirmed by MIC using broth dilution

according to EUCAST methodology.

WGS and phylogenetic analysis

WGS was performed by MicrobesNG to achieve a minimum 30-fold coverage.

Preparation of isolates was carried out following MicrobesNG strain submission

procedures as described in detail elsewhere.[32] Genomes were sequenced on an

Illumina NovaSeq 6000 (Illumina, San Diego, USA) using a 250 bp paired end

protocol. Sequencing data was analysed using ResFinder 4.4.2[33], Hound[34] and

MLST 2.0.[35]

Sequence alignment and phylogenetic analysis was carried out using the

computational facilities of the Advanced Computing Research Centre, University of

Bristol (http://www.bristol.ac.uk/acrc/) and the Bioconda environment management

system.[36] Alignment of sequences and reference sequence (**Table S5**) were

performed using Snippy and Snippy-Core software

(https://github.com/tseemann/snippy) and SNP distances were determined using

SNP-dists software (https://github.com/tseemann/snp-dists). Maximum likelihood

trees were produced using RAxML and the GTRGAMMA model of rate of

heterogeneity.[37] Phylogenetic trees were illustrated using Microreact.[38]

Data Availability

WGS data are deposited with ENA under accessions PRJEB74489 (Isolates from meat) and PRJEB74488 (isolates from canine faeces).

Funding

This work was funded by grant NE/N01961X/1 to M.B.A. from the Antimicrobial Resistance Cross Council Initiative supported by the seven United Kingdom research councils. It was also funded by grant BB/X012670/1 from the Biotechnology and Biological Sciences Research Council. J.E.S. was supported by a scholarship from the Medical Research Foundation National PhD Training Programme in Antimicrobial Resistance Research (MRF-145-0004-TPG-AVISO).

Transparency declaration

M.B.A. is married to the owner of a veterinary practice that sells various mass-manufactured dog foods amounting to a value less than 5% of total turnover. Otherwise, the authors declare no competing interests.

Author Contributions

Conceived the Study: J.E.S., M.B.A.

Collection of Data: J.E.S, B.A, K.R.

Cleaning and Analysis of Data: J.E.S.

Initial Drafting of Manuscript: J.E.S., M.B.A.

290   Corrected   and   Approved   Manuscript: All   authors

Table 1 NFT-R *E. coli* isolates excreted by dogs, and from raw poultry in Bristol, United Kingdom.

Isolates were from raw chicken meat sold for human consumption (RCM), chicken raw dog food (RDF), and canine faeces (DOG), sequence type (ST) and missense and nonsense mutations, deletions and insertions affecting NfsA and NfsB sequences are noted. Green represents those variants predicted to be functionally wild- type.[2] Red represents those conferring loss of function.

| Isolate | ST | Phylo-group | Effect of non-synonymous mutations relative to *E. coli* | | Other resistance genes |
			NfsA	NfsB	
RCM1	ST8874	B2	I117T, K141E, G187D, 212Stop	46Stop	None
RCM2	ST665	A	138FS	M75I, V93A, 101Stop	*aadA1*, *aph(3”)-Ib*, *aph(6)-Id*, *sul1*, *sul2*, *bla*TEM-1B, *tet*(A)
RDF15	ST8874	B2	I117T, K141E, G187D, 212Stop	46Stop	None
DOG1030	ST919	B2	I117T, K141E, G187D, 212Stop	46Stop	None
DOG1110	ST6805	A	Insertional inactivation after codon 225	M75I, 79Stop	*aadA5*, *dfrA17*, *sul2*, *tet*(A)

Figure 1 Maximum Likelihood phylogenetic tree based on a core genome alignment illustrating the evolutionary relationship between NFT-R ST919 and ST8874 isolates. Core genomes from a human UTI isolate from Enterobase (blue), dog faecal isolate (green), chicken isolates from Enterobase (yellow) and raw chicken meat (pink) were aligned using Snippy and Phylogenetic trees were constructed based on RaXML analysis. Reported SNP distances between pairs of isolates were determined using SNPdist. RCM, Raw Chicken Meat.

References

1. Howard B, Furman B. 2018. Nitrofurantoin. Reference Module in Biomedical Sciences: Elsevier; 72.

2. Dulyayangkul P, Sealey JE, Lee WWY, Satapoomin N, Reding C, Heesom KJ, Williams PB, Avison MB. 2024. Improving Nitrofurantoin Resistance Prediction in Escherichia coli from Whole Genome Sequence by Integrating NfsA/B Enzyme Assays. Antimicrob Agents Chemother. 68:e0024224.

3. McOsker CC, Fitzpatrick PM. 1994. Nitrofurantoin: mechanism of action and implications for resistance development in common uropathogens. J Antimicrob Chemother. 33 Suppl A:23-30.

4. Whiteway J, Koziarz P, Veall J, Sandhu N, Kumar P, Hoecher B, Lambert IB. 1998. Oxygen-insensitive nitroreductases: Analysis of the roles of nfsA and nfsB in development of resistance to 5-nitrofuran derivatives in Escherichia coli. J Bacteriol. 180:5529–5539.

5. Vervoort J, Xavier BB, Stewardson A, Coenen S, Godycki-Cwirko M, Adriaenssens N, Kowalczyk A, Lammens C, Harbarth S, Goossens H, Malhotra-Kumar S. 2014. An in vitro deletion in ribE encoding lumazine synthase contributes to nitrofurantoin resistance in Escherichia coli. Antimicrob Agents Chemother. 58:7225-33.

6. Wan Y, Mills E, Leung RCY, Vieira A, Zhi X, Croucher NJ, Woodford N, Jauneikaite E, Ellington MJ, Sriskandan S. 2021. Alterations in chromosomal genes nfsA, nfsB, and ribE are associated with nitrofurantoin resistance in Escherichia coli from the United Kingdom. Microb Genom. 7:000702.

7. Vallée M, Harding C, Hall J, Aldridge PD, Tan A. 2023. Exploring the in situ evolution of nitrofurantoin resistance in clinically derived uropathogenic Escherichia coli isolates. J Antimicrob Chemother. 78:373-379.

8. EFSA Panel on Contaminants in the Food Chain. 2015. Scientific Opinion on nitrofurans and their metabolites in food. EFSA J 13:4140, 217 pp.

9. Plumb DC. 2011. Veterinary Drug Handbook. 7th ed. Wiley—Blackwell; Hoboken, NJ, USA

10. https://www.vmd.defra.gov.uk/productinformationdatabase; searched 26[th] July 2024.

11. Sannes MR, Kuskowski MA, Johnson JR. 2004. Antimicrobial resistance of Escherichia coli strains isolated from urine of women with cystitis or pyelonephritis and feces of dogs and healthy humans. J Am Vet Med Assoc. 225:368-73.

12. Sayah RS, Kaneene JB, Johnson Y, Miller R. 2005. Patterns of antimicrobial resistance observed in Escherichia coli isolates obtained from domestic- and wild-animal fecal samples, human septage, and surface water. Appl Environ Microbiol. 71:1394-404.

13. Maaland M, Guardabassi L. 2011. In vitro antimicrobial activity of nitrofurantoin against Escherichia coli and Staphylococcus pseudintermedius isolated from dogs and cats. Vet Microbiol. 151:396-9.

14. Suay-García B, Galán F, Rodríguez-Iglesias MA, Pérez-Gracia MT. 2019. Detection and Characterization of Extended-Spectrum Beta-Lactamases-Producing Escherichia coli in Animals. Vector Borne Zoonotic Dis. 19:115-120.

15. Leuin AS, Hartmann F, Viviano K. 2021. Administration of nitrofurantoin in dogs with lower urinary tract infections: 14 cases (2013-2019). J Small Anim Pract. 62:42-48

16. Findlay J, Gould VC, North P, Bowker KE, Williams MO, MacGowan AP, Avison MB. 2020. Characterization of cefotaxime-resistant urinary Escherichia coli from primary care in South-West England 2017-18. J Antimicrob Chemother. 75:65-71.

17. Mounsey O, Schubert H, Findlay J, Morley K, Puddy EF, Gould VC, North P, Bowker KE, Williams OM, Williams PB, Barrett DC, Cogan TA, Turner KM, MacGowan AP, Reyher KK, Avison MB. 2021. Limited phylogenetic overlap between fluoroquinolone-resistant Escherichia coli isolated on dairy farms and those causing bacteriuria in humans living in the same geographical region. J Antimicrob Chemother. 76:3144-3150.

18. Dulyayangkul P, Beavis T, Lee WWY, Ardagh R, Edwards F, Hamilton F, Head I, Heesom KJ, Mounsey O, Murarik M, Reding C, Satapoomin N, Shaw JM, Takebayashi Y, Tooke CL, Spencer J, Williams PB, Avison MB. 2024. Harvesting and amplifying gene cassettes confers cross-resistance to critically important antibiotics. PLoS Pathog. 20:e1012235

19. Findlay J, Mounsey O, Lee WWY, Newbold N, Morley K, Schubert H, Gould VC, Cogan TA, Reyher KK, Avison MB. 2020. Molecular Epidemiology of Escherichia coli Producing CTX-M and pAmpC β-Lactamases from Dairy Farms Identifies a Dominant Plasmid Encoding CTX-M-32 but No Evidence for Transmission to Humans in the Same Geographical Region. Appl Environ Microbiol. 87:e01842-20.

20. Sealey JE, Hammond A, Mounsey O, Gould VC, Reyher KK, Avison MB. 2022. Molecular ecology and risk factors for third-generation cephalosporin-resistant

Escherichia coli carriage by dogs living in urban and nearby rural settings. J Antimicrob Chemother. 77:2399-2405.

21. Sealey JE, Hammond A, Reyher KK, Avison MB. 2023. One health transmission of fluoroquinolone-resistant Escherichia coli and risk factors for their excretion by dogs living in urban and nearby rural settings. One Health. 17:100640.

22. Sealey JE, Saunders R, Horspool T, Barrows MG, Avison MB. 2023. Molecular ecology of highest priority critically important antibiotic resistant Escherichia coli from mammals housed at an urban zoo. J Antimicrob Chemother. 78:1667-1671.

23. Morgan G, Pinchbeck G, Taymaz E, Chattaway MA, Schmidt V, Williams N. 2024. An investigation of the presence and antimicrobial susceptibility of Enterobacteriaceae in raw and cooked kibble diets for dogs in the United Kingdom. Front Microbiol. 14:1301841.

24. Sealey JE, Astley B, Rollings K, Avison MB. 2024. Antibiotic Resistant Escherichia coli in Uncooked Meat Purchased from Large Chain Grocery Stores and in Raw Dog Food Purchased From Pet Stores in the Same City. BioRxiv doi: https://doi.org/10.1101/2024.03.03.583175.

25. Papouskova A, Masarikova M, Valcek A, Senk D, Cejkova D, Jahodarova E, Cizek A. 2020. Genomic analysis of Escherichia coli strains isolated from diseased chicken in the Czech Republic. BMC Vet Res. 16:189.

26. Toombs-Ruane LJ, Benschop J, French NP, Biggs PJ, Midwinter AC, Marshall JC, Chan M, Drinković D, Fayaz A, Baker MG, Douwes J, Roberts MG, Burgess SA. 2020. Carriage of extended-spectrum beta-lactamase-and AmpC beta-lactamase-producing Escherichia coli from humans and pets in the same households. Appl. Environ. Microbiol. 86:e01613-20.

27. Kidsley AK, White RT, Beatson SA, Saputra S, Schembri MA, Gordon D, Johnson JR, O'Dea M, Mollinger JL, Abraham S, Trott DJ. 2020. Companion animals are spillover hosts of the multidrug-resistant human extraintestinal Escherichia coli pandemic clones ST131 and ST1193. Front. Microbiol. 11:1968.

28. Hong JS, Song W, Park HM, Oh JY, Chae JC, Jeong S, Jeong SH. 2020. Molecular characterization of fecal extended-spectrum β-lactamase- and AmpC β-lactamase-producing Escherichia coli from healthy companion animals and cohabiting humans in South Korea. Front. Microbiol. 11:674.

29. Hong JS, Song W, Park HM, Oh JY, Chae JC, Shin S, Jeong SH. 2019. Clonal spread of extended-spectrum cephalosporin-resistant Enterobacteriaceae between companion animals and humans in South Korea. Front. Microbiol. 10:1371.

30. Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X, Yu LF, Gu D, Ren H, Chen X, Lv L, He D, Zhou H, Liang Z, Liu JH, Shen J. 2016. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. Lancet Infect Dis. 16:161-8.

31. EUCAST. Breakpoint Tables for Interpretation of MICs and Zone Diameters, Version 13.10. 2023.

32. Mounsey O, Marchetti L, Parada J, Alarcón LV, Aliverti F, Avison MB, Ayala CS, Ballesteros C, Best CM, Bettridge J, Buchamer A, Buldain D, Carranza A, Corti Isgro M, Demeritt D, Escobar MP, Gortari Castillo L, Jaureguiberry M, Lucas MF, Madoz LV, Marconi MJ, Moiso N, Nievas HD, Ramirez Montes De Oca MA, Reding C, Reyher KK, Vass L, Williams S, Giraudo J, De La Sota RL, Mestorino N, Moredo FA, Pellegrino M. 2024. Genomic epidemiology of third-generation

cephalosporin-resistant Escherichia coli from Argentinian pig and dairy farms reveals animal-specific patterns of co-resistance and resistance mechanisms. Appl Environ Microbiol. 90:e0179123.

33. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, Philippon A, Allesoe RL, Rebelo AR, Florensa AF, Fagelhauer L, Chakraborty T, Neumann B, Werner G, Bender JK, Stingl K, Nguyen M, Coppens J, Xavier BB, Malhotra-Kumar S, Westh H, Pinholt M, Anjum MF, Duggett NA, Kempf I, Nykäsenoja S, Olkkola S, Wieczorek K, Amaro A, Clemente L, Mossong J, Losch S, Ragimbeau C, Lund O, Aarestrup FM. 2020. ResFinder 4.0 for predictions of phenotypes from genotypes. J Antimicrob Chemother. 75:3491–500.

34. Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Pontén T, Ussery DW, Aarestrup FM, Lund O. 2012. Multilocus Sequence Typing of Total Genome Sequenced Bacteria. J Clin Microbiol. 50:1355-1361.

35. Reding C, Satapoomin N, Avison MB. 2024. Hound: a novel tool for automated mapping of genotype to phenotype in bacterial genomes assembled de novo. Brief Bioinform. 25:bbae057.

36. Grüning, Björn, Ryan Dale, Andreas Sjödin, Brad A. Chapman, Jillian Rowe, Christopher H. Tomkins-Tinch, Renan Valieris, the Bioconda Team, and Johannes Köster. 2018. Bioconda: Sustainable and Comprehensive Software Distribution for the Life Sciences. Nat Methods. 15:475-476.

37. Stamatakis, A. 2014. RAxML Version 8: A tool for Phylogenetic Analysis and Post-Analysis of Large Phylogenies. Bioinformatics 30: 1312-1313

38. Argimón S, Abudahab K, Goater RJE, Fedosejev A, Bhai J, Glasner C, Feil EJ, Holden MTG, Yeats CA, Grundmann H, Spratt BG, Aanensen DM. 2016.

Microreact: visualizing and sharing data for genomic epidemiology and phylogeography. Microb Genom 2:e000093.