The protease interpain A of *Prevotella intermedia* promotes human oral squamous cell carcinoma cells proliferation and migration

Szu-Min Chang[1][*], I-Hsiu Huang[2][*], Yun-Jhen Ciou[3], Wen-Rong Wang[3], Ni-Hung Wu[3], Chih-Pin Chuu[4], Jenn-Ren Hsiao[5], Jeffrey S Chang[6], Jang-Yang Chang [7,8], Jenn-Wei Chen[1, 3, 9]

[1] Institute of Basic Medical Sciences, College of Medicine, National Cheng Kung University, Tainan 70101, Taiwan
[2] Department of Biochemistry and Microbiology, Oklahoma State University Center for Health Sciences, Tulsa, OK 74107, USA.
[3] Department of Microbiology and Immunology, College of Medicine, National Cheng Kung University, Tainan 70101, Taiwan
[4] Institute of Cellular and System Medicine, National Health Research Institutes, Miaoli 35053, Taiwan
[5] Department of Otolaryngology, National Cheng Kung University Hospital, Tainan 70101, Taiwan
[6] National Institute of Cancer Research, National Health Research Institutes, Tainan 70101, Taiwan
[7] Institute of Biotechnology and Pharmaceutical Research, National Health Research Institutes, Miaoli 35053, Taiwan
[8] TMU Research Center of Cancer Translational Medicine, Taipei Medical University, Taipei 110301, Taiwan
[9] Center of Infectious Disease and Signaling Research, National Cheng Kung University, Tainan 70101, Taiwan

[*] These authors contributed equally to this article.

Running title: Interpain A promotes OSCC cells proliferation and migration

Key words: *Prevotella intermedia*; OSCC, interpain A; PAR-2

Corresponding Author: Jenn-Wei Chen, Department of Microbiology and Immunology, College of Medicine, National Cheng Kung University, Tainan 70101, Taiwan. E-mail: jc923@mail.ncku.edu.tw

Competing Interest Statement: The authors declare no potential conflicts of interest.

Abstract

Prevotella intermedia has been increasingly recognized as a potential contributor to oral squamous cell carcinoma (OSCC), yet the underlying mechanisms remain poorly defined. In this study, we identified interpain A (InpA), a cysteine protease secreted by *P. intermedia*, as a key virulence factor that promotes oral epithelial cell proliferation and OSCC cell migration. Conditioned medium (CM) derived from *P. intermedia* strain ATCC 25611 enhanced proliferation in both normal oral keratinocytes (SG cells) and OSCC cell lines (SCC-15, SAS). The pro-proliferative effect was abolished upon heat inactivation of the CM and inhibited by E64, a cysteine protease inhibitor, and FSLLRY-NH$_2$, a PAR-2 antagonist, suggesting the involvement of protease-activated receptor-2 (PAR-2) signaling. InpA was highly secreted by strain ATCC 25611 but not by clinical isolates lacking proliferative effects, and RT-qPCR confirmed higher *inpA* expression in OSCC-derived strains compared to non-cancer controls. Recombinant InpA (rInpA) mimicked the effect of CM, inducing proliferation and migration, which were blocked by PAR-2 inhibition. Transcriptomic and protein-level screening in SG cells revealed activation of EGFR and downstream RAS–RAF–MEK–ERK signaling. Furthermore, in a colorectal cancer (CRC) mouse model, oral administration of *P. intermedia* led to increased tumor formation, suggesting a broader oncogenic potential. These findings highlight InpA as a PAR-2–activating protease that may contribute to OSCC and potentially other cancers associated with oral microbiota dysbiosis.

Introduction

Oral squamous cell carcinoma (OSCC) is a common malignancy worldwide and ranks as the fourth most prevalent cancer among men in Taiwan (1). It represents the most frequent form of head and neck cancer and can develop in various parts of the oral cavity, including the lips, tongue, cheeks, palate, and throat. The majority of oral cancers are squamous cell carcinomas, which are associated with poor prognosis and high mortality, with a 5-year survival rate of approximately 50%. Numerous factors contribute to OSCC development, such as gender, human papillomavirus (HPV) infection, alcohol consumption, tobacco use, betel nut chewing, poor oral hygiene, and alterations in the oral microbiota (2, 3). Notably, chronic periodontitis, a gum disease characterized by gingival inflammation and tooth loss, has been strongly linked to oral cancer (4). Poor oral hygiene facilitates the formation of biofilms, known as dental plaques, which contribute to oral microbiota dysbiosis. Anaerobic bacteria residing within these plaques produce acids and other virulence factors that contribute to periodontal inflammation and disease (5). Anaerobic bacteria residing within these plaques produce acids and other virulence factors that contribute to periodontal inflammation and disease.

Several bacterial genera, including *Porphyromonas*, *Fusobacterium*, and *Prevotella*, are found in higher abundance in patients with periodontitis compared to healthy individuals. (6, 7). *Porphyromonas gingivalis*, a Gram-negative anaerobe, is associated with periodontal disease and systemic conditions such as Alzheimer's disease (8). Studies have also linked *P. gingivalis* to various cancers, including lung, pancreatic, prostate, colorectal cancer (CRC), and OSCC (9, 10). This bacterium produces multiple virulence factors, such as lipopolysaccharides (LPS), outer membrane vesicles (OMVs), capsules, and gingipains—cysteine proteases implicated

in promoting tumor invasion and migration (11). Another oral bacterium,

Fusobacterium nucleatum, another Gram-negative anaerobic bacterium, has been

associated with CRC, breast, pancreatic, and oral cancers (12, 13). It secretes virulence

factors including OMVs, LPS, and adhesins. The adhesin FadA interacts with E-

cadherin to activate WNT/β-catenin signaling and promote cell proliferation (14).

While Fap2 binds tumor-associated oligosaccharides and the immune checkpoint

receptor TIGIT, potentially facilitating immune evasion (15). While much attention has

focused on *P. gingivalis* and *F. nucleatum* in OSCC, the role of *Prevotella intermedia*

remains poorly understood. *P. intermedia* is a Gram-negative, obligate anaerobic rod-

shaped bacterium that requires vitamin K and hemin for growth and is associated with

periodontal disease and acute necrotizing ulcerative gingivitis. Recent studies have

reported elevated levels of *P. intermedia* in the oral microbiota of OSCC patients

compared to healthy controls. Although this correlation suggests a potential role in oral

carcinogenesis, the molecular contributions of this association remain unclear.

In this study, we demonstrate that *P. intermedia* strain ATCC 25611 promotes

proliferation of normal oral epithelial cells (SG cell line) at a multiplicity of infection

(MOI) of 100. Its conditioned medium (CM) also enhanced proliferation in SG, SCC-

15, and SAS cell lines. Proteomic analysis using capillary electrophoresis/liquid

chromatography-mass spectrometry (CE/LC-MS) revealed that a cysteine protease,

interpain A (InpA), secreted by strain ATCC 25611, may be the critical factor driving

this effect. In addition to stimulating proliferation, InpA also enhanced migration of

OSCC cells. Furthermore, in a murine colorectal cancer model, colonization with *P.*

intermedia increased tumor burden. These findings support a role for *P. intermedia*,

and specifically InpA, in promoting OSCC development.

Materials and methods

Bacterial strains and cultures

P. intermedia strain ATCC 25611 was obtained from the Bioresource Collection and Research Center (BCRC, Taiwan), and culture conditions were adapted from a previously published protocol (16). Bacteria were grown anaerobically at 37 °C in supplemented Tryptic Soy Broth (ATCC 2722), consisting of tryptic soy broth (#211825, BD) enriched with 0.5% yeast extract (#212750, BD), 0.05% L-cysteine (#A3694, Applichem Panreac), 0.1% hemin (#H9039, Sigma), and 0.02% menadione (#M5625, Sigma). Clinical isolates of *P. intermedia* were obtained from the saliva of patients at National Cheng Kung University Hospital (NCKUH), under Institutional Review Board approval (IRB no. B-ER-106-234). Saliva samples were diluted in PBS and plated on Kanamycin-Vancomycin Laked Blood (KVLB) agar (TPM, Yilan, Taiwan). After 2–3 days of anaerobic incubation (5% CO_2, 5% H_2, 90% N_2), black-pigmented colonies were selected. Colony identity was confirmed by Gram staining and PCR using *P. intermedia*-specific primers Pi-192 and Pi-486 (Supplementary Table 1). The PCR conditions were as follows: initial denaturation at 95 °C for 5 min; 35 cycles of 95 °C for 30 s, 55 °C for 30 s, and 72 °C for 30 s; and a final extension at 72 °C for 7 min.

Cell lines and cultures

The immortalized human gingival keratinocyte, Smulow–Glickman (SG) cell (17), was kindly provided by Dr. Chuan-Fa Chang (NCKU) and cultured with Dulbecco's Modified Eagle Medium (DMEM) (#11965-092, Gibco/Life Technology, Carisbad, CA, USA). Human tongue carcinoma cell lines, SAS cells (18), and SCC-15 cells (19) were obtained from Dr. Jhen-Wei Ruan (NCKU) and maintained in DMEM/F12

(D/F12) medium (#11330-032, Gibco). All media were supplemented with 10% FBS

(#10437-028, Gibco) and 1% 100x Penicillin-Streptomycin-Amphotericin B solution

(#CC501-0100, Simply Biotechnology, San Diego, CA, USA). Cells were incubated

at 37 °C in a humidified atmosphere containing 5% CO_2.

Conditioned medium (CM) preparation

The preparation of *P. intermedia* condtioned medium was adapted from a previously

published protocol (20). *P. intermedia* strains were anaerobically cultured in ATCC

2722 broth at 37 °C and were harvested when the OD600 value reached 1.0. The

bacterial culture was centrifuged at 3220 xg for 3 min and discarded supernatant. The

pellet was resuspended with a double volume of cell culture medium and anaerobically

incubated at 37 °C for an additional 8 hr. Then the supernatant was collected after being

centrifuged at 3220 xg for 10 min, filtered with a 0.22 µm filter (Merck Millipore,

Burlington, MA, USA), and stored at 4 °C before use.

Cell proliferation assay

The cell proliferation assay was performed as previously described (21). SG, SAS, or

SCC-15 cells were dissociated by 0.05% trypsin-EDTA solution (#25300-054, Gibco)

and resuspended in DMEM or D/F12 medium. A total of 1×10^4 cells were seeded into

each well of 24-well tissue culture plates (Corning, NY, USA) and treated with bacteria

or CM treatment. CM was fractionated into four components using centrifugal filter

devices with molecular weight cutoffs of 3 kDa, 10 kDa, and 30 kDa (Merck Millipore).

Treatments included 10% CM; the PAR-2 antagonist FSLLRY-NH2 at 40 µM for SG

cells or 20 µM for SCC-15 and SAS cells (#4751, TOCRIS/Bio-Techne, Minneapolis,

MN, USA); and the cysteine protease inhibitor E64 at 5 µM for SG cells or 0.5 µM for

SCC-15 and SAS cells (#324890, Merck Millipore). All treatments were administered

in DMEM or D/F12 containing 10% FBS to a final volume of 1 ml. After incubation

for 24, 48, or 72 hr at 37 °C in a 5% CO_2 atmosphere, cells were washed once with 1×

PBS and then dissociated using 0.05% trypsin-EDTA. The resulting cell suspension

was mixed with an equal volume of 0.4% trypan blue (#15250-061, Gibco), and viable

cells were counted using a Luna cell counter (Logos Biosystems, Gyeonggi-do, South

Korea).

Capillary electrophoresis/Liquid Chromatography-Mass spectrometry (CE/LC

MASS) analysis

A total of 100 µg of protein in 60 µl rehydration buffer (7 M urea, 2 M thiourea, and

2% CHAPS) was mixed with 9.3 µl of 7.5% SDS (#75746, Sigma, Burlington, MA,

USA) and 0.7 µl of 1 M dithiothreitol (#UR-DTT, UniRegion Bio-Tech, Taichung,

Taiwan) prepared in 7 M urea and 0.01 M sodium acetate. The mixture was heated at

95 °C for 5 min, then placed on ice for 3 min. Next, 4 µl of 500 mM IAM

(iodoacetamide in 50 mM ammonium bicarbonate) was added and incubated at room

temperature for 30 min. Afterward, 52 µl of 50% trichloroacetic acid (TCA) (#100807,

Merck Millipore) was added, and the solution was placed on ice for 15 min. The

precipitate was collected by centrifugation at 15,871 ×g for 15 min at 4 °C, resuspended

in 150 µl of 10% TCA, and centrifuged again at 15,871 ×g for 5 min at 4 °C. This

washing step was repeated three times. The final pellet was resuspended in 100 µl of

100 mM ammonium bicarbonate and digested with 20 µl of trypsin solution (#V5111,

Promega, Madison, WI, USA) at 37 °C for 16 hr. To inactivate trypsin, samples were

stored overnight at −80 °C. Prior to analysis, samples were desalted using Ziptips

(Merck Millipore). Protein identification was performed using CE/LC-MS on a Q

Exactive Orbitrap Mass Spectrometer (Thermo Fisher Scientific).

Interpain A gene expression quantitative analysis (RT-qPCR)

RNA extraction and RT-qPCR were adapted from previously described methods (22).

A 2 ml culture of *P. intermedia* (OD_{600} = 0.8) was mixed with 4 ml of RNAprotect

Bacteria Reagent (QIAGEN, Venlo, Netherlands). After centrifugation at 2,200 ×g for

5 min, the supernatant was discarded, and the pellet was stored at −80 °C until RNA

extraction. RNA was purified using the RNeasy Mini Kit (QIAGEN), and residual

DNA was removed using DNase (Promega). The resulting RNA was concentrated with

the Clean & Concentrator Kit (ZYMO GENETICS, Seattle, WA, USA). cDNA was

synthesized from the RNA using the SuperScript III First-Strand System with random

hexamers (Invitrogen, Waltham, MA, USA). RT-qPCR was performed using GoTaq

qPCR Master Mix (Promega) on a MyGo Pro real-time PCR instrument (IT-IS Life

Science Ltd/DKSH, St Leonards, Australia) with primers inpA F, inpA R, Pi 16S rRNA

F, and Pi 16S rRNA R (Supplementary Table 1). The thermal cycling conditions were

as follows: initial denaturation at 95 °C for 10 min, followed by 40 cycles of

denaturation at 95 °C for 15 s and annealing/extension at 60 °C for 1 min. A melt curve

analysis was performed from 60 to 97 °C at a ramp rate of 0.1 °C/s. All procedures

were conducted in accordance with the manufacturers' instructions.

InpA$_{N381}$ expression vector construction and recombinant InpA$_{N381}$ purification

Genomic DNA of *P. intermedia* ATCC 25611 was extracted using the MasterPure^{TM}

complete DNA and RNA purification kit (Lucigen Corp, Middleton, WI, USA)

according to the manufacturer's instructions. The gene region encoding the signal

peptide, pro-domain, and catalytic domain (inpA$_{N381}$) was amplified using a mixture of

GoTaq G2 Flexi DNA Polymerase (Promega) and Pfu polymerase (Promega) with

primers $inpA_{N381}$ F and $inpA_{N381}$ R (Supplementary Table1). The PCR product was

digested with BamHI and XhoI and ligated into the pET21b(+) expression vector pre-

digested with the same enzymes. The resulting plasmid, pET21b(+)::$inpA_{N381}$, was

verified by sequencing and chemically transformed into *Escherichia coli* C41 (DE3)

cells. A single colony of *E. coli* C41 (DE3) harboring pET21b(+)::$inpA_{N381}$ was

inoculated into 20 ml LB broth supplemented with ampicillin (100 µg/ml) and cultured

at 37 ℃ for 14-16 hr with shaking at 170 rpm. The bacterial culture was then refreshed

with 2 liters LB at the ratio of 1:100 and grew at 37 ℃ with shaking at 170 rpm until

OD600 reached 0.4. Protein expression was induced by adding 0.2 mM isopropyl-β-

D-thiogalactoside (IPTG; #101-367-93-1, Cyrusbioscience, Inc., Seattle, WA, USA),

followed by incubation at 25 ℃ for an additional 16 hr with shaking (23). After

induction, the bacterial culture was centrifuged at 3,300 ×g for 40 min at 4 ℃. The

resulting pellet was resuspended in 100 ml of 1× PBS and lysed by ultrasonication on

ice (20 s pulse on, 100 s pulse off) until complete cell disruption. The lysate was then

centrifuged at 16,000 ×g for 40 min at 4 ℃. $InpA_{N381}$ protein was purified from the

supernatant using a HisTrap HP His-tag affinity column (Cytiva, Washington D.C.,

USA) according to the manufacturer's instructions. The purified $InpA_{N381}$ protein was

washed with 100 volumes of 1× PBS and stored at −20 ℃ until use.

Transwell assay

The Transwell assay was conducted as previously described (24). SAS or SCC-15 cells

were resuspended in D/F12 medium supplemented with 10% FBS. Cell concentration

was adjusted to 1×10^5 cells/ml. A total of 100 µl of cell suspension was added to the

upper chamber of Transwell inserts (8 µm pore size, Falcon/Corning, NY, USA) placed

in a 24-well plate and incubated at 37 °C in 5% CO_2. The lower chamber was filled

with 10% CM, 20 μM FSLLRY-NH$_2$, or 0.5 μM E64, in D/F12 containing 10% FBS,

to a final volume of 600 μl. After 24 hr, cells in the upper chamber were removed with

cotton swabs, and migrated cells on the lower side of the membrane were fixed with

70% ethanol for 10 min and stained with 0.2% crystal violet for 5 min. Five random

fields per membrane were selected for cell counting under an inverted microscope

(Leica Microsystems, Wetzlar, Germany).

Wound healing assay

The wound healing assay was performed according to a previously reported method

(24). SAS or SCC-15 cells were seeded in both compartments of each block insert (SPL

Life Sciences Co., Gyeonggi-do, Korea), placed in 24-well plates, and incubated at

37 °C with 5% CO_2 for 24 hr. The inserts were then removed, and the wells were

washed once with 1× PBS. Cells were treated with 10% CM, 20 μM FSLLRY-NH$_2$, or

0.5 μM E64, with the total volume adjusted to 1 ml using D/F12 containing 10% FBS.

Images were captured every 2 hr using a microscope (OLYMPUS, Tokyo, Japan) to

monitor gap closure. Gap widths were measured using ImageJ, and wound closure rates

were calculated by dividing the change in width over time.

Colorectal cancer (CRC) mouse model

The mouse model was modified from previously published protocols (25). Seven-

week-old specific pathogen-free (SPF) male C57BL/6 mice were purchased from the

Laboratory Animal Center of NCKU. Mice received a single intraperitoneal injection

of azoxymethane (AOM; Sigma-Aldrich) at 12.5 mg/kg, followed by administration of

2% dextran sodium sulfate (DSS; MP Biomedicals, Irvine, CA, USA) in drinking water

to induce colorectal cancer. Twenty mice were randomly assigned to four groups:

MOCK, AOM/DSS, AOM/DSS + *F. nucleatum*, and AOM/DSS + *P. intermedia*. Mice

underwent three cycles of DSS and bacterial treatment. Each cycle lasted 3 weeks: 5

days of 2% DSS in the first week, followed by two weeks of oral gavage with either

1× PBS or 1×10^9 CFU of *F. nucleatum* (ATCC 23726) or *P. intermedia* (ATCC 25611)

once per week. Mice were sacrificed at week 10, and colons were harvested,

longitudinally opened, and tumor numbers were counted.

Statistical analysis

Statistical analyses were conducted using GraphPad Prism 8 software. All data were

obtained from experiments performed in triplicate. Comparisons between two groups

were assessed using the Student's t-test, while multiple group comparisons were

analyzed by one-way ANOVA. Statistical significance was indicated as follows: $p <$

0.05 (*), $p < 0.01$ (**), $p < 0.001$ (***), and $p < 0.0001$ (****).

Results

***P. intermedia* ATCC 25611 enhances human oral cell proliferation**

A previous study at National Cheng Kung University Hospital (NCKUH)

reported a significant increase in *P. intermedia* abundance in patients with oral

squamous cell carcinoma (OSCC) compared to healthy controls (6). To investigate the

potential contribution of *P. intermedia* to OSCC development, we assessed its effect

on oral epithelial cell proliferation. The reference strain ATCC 25611 was tested at

multiple multiplicities of infection (MOIs) using the human normal oral epithelial cell

line SG. After 24 hours of co-culture, *P. intermedia* ATCC 25611 significantly

increased SG cell numbers at an MOI of 100 compared to the uninfected control group

(Fig. 1A, 1B). In addition to ATCC 25611, we also examined six clinical isolates of

P. intermedia obtained from patient saliva samples at NCKUH. Among these, strains

H26219-3, H29383, and H29703 were isolated from OSCC patients, while strains

H31082, H29823, and H31226 were obtained from individuals with non-cancerous oral

conditions. SG cells were co-cultured with each clinical isolate at MOI 100. Unlike

ATCC 25611, these strains did not induce a proliferation increase in SG cells; notably,

strain H29383 even reduced SG cell numbers after 24 hours (Fig. 1B; Fig. S1A). We

further found that only live *P. intermedia* was capable of promoting SG cell

proliferation (Fig. 1C), likely due to its limited viability under aerobic conditions,

which may confine its activity to the early co-culture period. Similarly, conditioned

medium (CM) from strain ATCC 25611 increased SG cell proliferation, whereas CM

from the clinical strains failed to elicit the same response (Fig. 1D; Fig. S1B). These

findings were corroborated using EdU incorporation assays (Fig. S1C), which supports

the role of a secreted factor in enhancing proliferation. To identify this active factor,

we subjected the CM to heat inactivation, which abolished its proliferative effect (Fig.

1D), indicating the involvement of a heat-sensitive, likely proteinaceous molecule.

Further size-fractionation of CM revealed that only the >30 kDa fraction retained

stimulatory activity (Fig. 1E). Additionally, CM from ATCC 25611 significantly

enhanced proliferation in OSCC cell lines SCC-15 and SAS after 48 and 72 hours,

respectively (Fig. 1F, 1G), suggesting that *P. intermedia* promotes oral epithelial cell

proliferation through the action of one or more secreted proteins.

Interpain A plays a central role in *P. intermedia* ATCC 25611 promoting oral cell

proliferation

Following our observation that strain ATCC 25611 and several clinical isolates

exhibited divergent effects on SG cell proliferation, we hypothesized that these

differences could be attributed to the expression of specific secreted protein(s). To

investigate this, we conducted proteomic analysis of conditioned medium (CM) from

strain ATCC 25611 and the non-proliferative strain H31226 using capillary

electrophoresis/liquid chromatography-mass spectrometry (CE/LC-MS). Several

proteins were found to be more abundant in the CM of ATCC 25611 compared to that

of H31226 (Supplementary Table 2). Among these, interpain A (InpA), a secreted

cysteine protease, was particularly enriched—present at levels thirteen times higher in

ATCC 25611 than in H31226 (Fig. 2A). InpA is a 90 kDa cysteine protease previously

reported to facilitate heme acquisition for bacterial growth and to degrade complement

factor C3, aiding immune evasion (26-28). Notably, *P. gingivalis*, another oral

anaerobe strongly associated with chronic periodontitis and cancer, produces similar

cysteine proteases called gingipains, which have been shown to activate protease-

activated receptors (PARs) on oral epithelial cells (11). Gingipains, in particular, have

been implicated in OSCC progression through activation of PAR-2 signaling,

enhancing cell migration and invasiveness (29, 30). Given these functional similarities,

we hypothesized that InpA may enhance oral epithelial cell proliferation via activation

of the PAR-2 signaling pathway. To test this hypothesis, SG cells were treated with

CM from ATCC 25611 in the presence of either E64, a cysteine protease inhibitor, or

FSLLRY-NH$_2$, a PAR-2 antagonist. After 24 hours, the proliferation-promoting effects

of CM were significantly attenuated by both inhibitors (Fig. 2B). A similar inhibition

pattern was observed under direct bacterial co-culture conditions. These results were

consistent across SCC-15 and SAS oral cancer cell lines, where the proliferative effects

observed after 48 hours were also blocked by E64 or FSLLRY-NH$_2$ treatment (Fig. 2C,

2D). To further confirm InpA's direct effect, we expressed and purified the

recombinant catalytic domain of InpA (InpA$_{N381}$) in *E. coli* strain C41. InpA$_{N381}$

stimulated SCC-15 proliferation in a manner consistent with that of native CM, and

again, this effect was neutralized by either E64 or FSLLRY-NH$_2$ treatment (Fig. 2E).

Importantly, treatment with E64 or FSLLRY-NH$_2$ alone had no effect on the

proliferation of SG, SCC-15, or SAS cells (Fig. S2), confirming that the inhibitors'

effects were specific to the activity of InpA or the associated pathway. We next

investigated whether differences in InpA production could explain the observed strain-

dependent variability in proliferation. RT-qPCR analysis revealed significantly higher

inpA expression in ATCC 25611 compared to all clinical isolates tested (Fig. 2F).

Furthermore, among the clinical strains, those isolated from OSCC patients (H26219-

3, H29383, and H29703) exhibited higher *inpA* expression than strains from patients

with non-cancerous oral diseases (H31082, H29823, and H31226). These findings

collectively indicate that *P. intermedia* promotes oral epithelial cell proliferation via

InpA and the PAR-2 signaling axis and that this activity is strain-dependent, correlating

with InpA production levels.

InpA-Mediated Promotion of OSCC Cell Migration

During cancer progression, tumor cells initially undergo uncontrolled growth

at the primary site, followed by invasion and dissemination to distant organs—a

process known as metastasis. Metastasis involves the degradation of the extracellular

matrix, intravasation into the vasculature, and colonization of secondary tissues, and is

a major contributor to cancer-related mortality (31). Given our findings that *P.*

intermedia promotes oral epithelial cell proliferation, we next investigated whether it

also influences OSCC cell migration. To address this, we employed two

complementary assays—wound healing and transwell migration assays—using SCC-

15 and SAS OSCC cell lines. In the wound healing assay, treatment with conditioned

medium (CM) from *P. intermedia* strain ATCC 25611 significantly enhanced the

migration rate of both cell lines at 2 and 4 hours post-treatment compared to untreated

controls (Fig. 3A, 3B). To determine whether this migration-promoting effect involved

InpA and PAR-2 signaling, we co-treated cells with either E64 or FSLLRY-NH$_2$. Both

inhibitors effectively abrogated the CM-induced enhancement of cell migration.

Similarly, in the transwell migration assay, CM from ATCC 25611 significantly

increased the number of migrating SCC-15 and SAS cells after 24 hours (Fig. 3C, 3D).

Treatment with E64 or FSLLRY-NH$_2$ had no significant effect on SCC-15 and SAS

cell migration in either wound healing or transwell assays (Fig. S3). Co-treatment with

E64 or FSLLRY-NH$_2$ suppressed this enhanced migratory response. Furthermore,

recombinant InpA$_{N381}$ purified from *E. coli* elevated SCC-15 cell migration (Fig. 3E),

and this effect was also blocked by both E64 and FSLLRY-NH$_2$. Together, these

findings suggest that *P. intermedia* facilitates OSCC cell migration through an InpA-

dependent mechanism involving PAR-2 signaling, paralleling its role in cell

proliferation.

***P. intermedia* increases tumor burden in a colorectal cancer mouse model**

An increasing body of evidence suggests that oral microbiota not only

contributes to oral diseases but also plays a role in systemic conditions such as cystic

fibrosis, cardiovascular disease, rheumatoid arthritis, and gastrointestinal disorders

(32). Furthermore, several oral bacteria have been linked to cancers in distal organs,

including the pancreas, breast, stomach, esophagus, and colon (32). The concept of the

oral-gut axis proposes that oral bacteria can translocate to the gastrointestinal tract,

where they may disrupt the gut microbial balance and contribute to diseases such as

inflammatory bowel disease and colorectal cancer (CRC) (33, 34). Notably,

Fusobacterium nucleatum, an oral anaerobe, has been identified as an important

contributor to CRC progression (35). Given the pro-tumorigenic effects of *P.*

intermedia observed in OSCC, we hypothesized that it may also influence CRC

development. To test this, we used an azoxymethane (AOM) and 2% dextran sulfate

sodium (DSS)-induced CRC model in C57BL/6 mice, followed by oral gavage with

either *F. nucleatum* or *P. intermedia*. Compared to the AOM/DSS-only group, both

bacterial treatments significantly increased tumor formation and elevated the number

of colonic tumors (Fig. 4B, 4C). These results suggest that *P. intermedia*, like *F.*

nucleatum, may contribute to CRC pathogenesis and should be considered a relevant

microbial factor in colorectal tumorigenesis.

Discussion

The present study demonstrated that *P. intermedia* may play a significant role in

oral cancer development. Interpain A (InpA), a cysteine protease secreted by *P.*

intermedia, functioned as an effector that enhanced both oral epithelial cell

proliferation (Fig. 2B) and OSCC cell migration (Fig. 3A–D). Importantly, these

effects were linked to the protease-activated receptor-2 (PAR-2) signaling pathway

activation. Prior studies examining salivary samples from OSCC patients reported

increased expression of various oral bacterial virulence genes, including *fadA* and *inpA*

(36). Notably, PAR-2 has been implicated in promoting cell proliferation, metastasis,

and invasion in multiple cancers, including OSCC, esophageal carcinoma, and

pancreatic cancer (37). Previous studies have also shown that gingipains, secreted

cysteine proteases from *P. gingivalis*, can activate the PAR-2/NF-κB signaling axis and

induce proMMP9 expression, thereby facilitating OSCC invasion (11). These findings are consistent with our observations implicating InpA and PAR-2 signaling in OSCC progression.

Our results showed that *P. intermedia*-CM treatment promoted proliferation in both normal oral keratinocytes (SG cells) and OSCC cell lines (SCC-15 and SAS) (Fig. 1D–G). Interestingly, the proliferative effect manifested at different time points between cell types: SG cells responded within 24 hours, whereas SCC-15 and SAS cells showed continued stimulation at 48 and 72 hours. Moreover, SG cell numbers decreased after 48 hours, suggesting that normal oral cells may retain mechanisms to counterbalance proliferative signals, while cancer cells have lost this regulatory capacity. A previous study of *P. gingivalis* reported G_1 cell cycle arrest and apoptosis in trophoblasts, as well as in osteoblastic/stromal cells following gingipain exposure (38, 39). Gingipains have also been shown to induce apoptosis in oral epithelial cells (40). It is possible that InpA exerts similar effects in SG cells following initial proliferation, potentially through the regulation of cell cycle or apoptotic pathways. This observation warrants further investigation. Our findings also highlight the need to clarify whether InpA differentially activates downstream PAR-2 signaling in normal versus malignant oral cells.

The concept of an oral-gut microbiome axis is increasingly supported by evidence showing that oral dysbiosis can influence both local and systemic health (41); A growing number of studies have linked oral microbiota imbalances with gastrointestinal diseases, including colorectal cancer (CRC) (42). Recent findings suggest that *P. intermedia* enhances CRC cell proliferation and migration through secreted protein effectors, similar to those identified in our OSCC model (43). The promoting effect was also from *P. intermedia*-secreted protein(s). Those results were

similar to our findings (Fig. 2C-E). In agreement with these findings, our data showed

that oral administration of *P. intermedia* increased tumor burden in an AOM/DSS-

induced CRC mouse model (Fig. 4). These results reinforce the idea that *P. intermedia*

may act as a tumor-promoting microbe across multiple tissue contexts. Beyond cancer,

P. intermedia and other oral pathogens have been implicated in the pathogenesis of

various systemic diseases (44). Oral bacteria are capable of releasing inflammatory

mediators that may affect the central nervous system. Elevated serum antibody levels

against *P. gingivalis*, *F. nucleatum,* and *P. intermedia* have been observed in patients

with Alzheimer's disease (45). Gingipains from *P. gingivalis* have been detected in the

brains of Alzheimer's patients and are known to increase blood–brain barrier

permeability (8, 46). Moreover, gingipains have been associated with other

neurodegenerative diseases, such as Parkinson's disease (47). Periodontal bacteria

have also been identified in atheromatous plaques and are believed to contribute to

cardiovascular disease pathogenesis (48, 49). These findings position *P. intermedia* as

a potentially important contributor to systemic inflammatory and degenerative diseases.

Our study also supports the notion that the pro-proliferative effects of *P.*

intermedia are strain-dependent. In SG cells, only strain ATCC 25611 induced

proliferation at MOI 100 (Fig. 1B, S1A), and only its conditioned medium promoted

proliferation (Fig. 1D, S1B). Proteomic analysis confirmed that this strain secreted

significantly higher levels of InpA than clinical isolates such as H31226 (Fig. 2A), and

RT-qPCR revealed higher *inpA* expression in strains isolated from OSCC patients

compared to non-cancer controls (Fig. 2F). These observations raise the possibility of

using salivary InpA levels as a biomarker for OSCC risk or diagnosis. Although there

was no statistically significant difference between clinical strain H26219-3 or H29383

and the untreated control group at the 24-hour time point (Fig. S1B), SG cell

proliferation showed a mild upward trend in response to both strains isolated from

OSCC patients. This observation suggests that these strains may exert weak but

measurable biological effects. Given that such bacteria can persistently colonize the

oral cavity, it is plausible that their cumulative impact over time could become

biologically relevant. However, short-term cell-based assays are unlikely to capture

such long-term effects.

To explore the mechanism underlying InpA-dependent proliferation, we

performed a Micro-Western Array focusing on signaling pathways related to cell

growth in SG cells (Fig. S4). We found that the EGFR signaling pathway was

upregulated, with RAF expression increasing at 8 hours and decreasing at 24 hours.

This aligns with prior studies showing that the EGFR–RAS–RAF–MEK–ERK axis is

frequently upregulated in OSCC (50). While these findings provide a preliminary

mechanistic link between InpA and cancer signaling pathways, further validation is

required.

In conclusion, our study demonstrates that InpA, a cysteine protease secreted by

P. intermedia, plays a pivotal role in promoting oral epithelial cell proliferation and

OSCC cell migration. These effects appear to be mediated via the PAR-2 signaling

pathway and are strain-dependent. The results suggest a potential microbial mechanism

contributing to OSCC development and lay the groundwork for future investigations

into therapeutic strategies targeting microbial virulence factors.

Figure legends

Figure 1. *P. intermedia* ATCC 25611 enhances proliferation of normal and

cancerous oral epithelial cells. (A) Human normal oral epithelial SG cells were

infected with live *P. intermedia* strain ATCC 25611 at a multiplicity of infection (MOI)

of 100 or 1000 for 24 or 48 hours. (B) SG cells were co-cultured with strain ATCC

25611 or clinical isolates H29383 and H31226 at MOI 100. (C) Comparison of live

versus UV-inactivated *P. intermedia* strain ATCC 25611 on SG cell proliferation. (D)

Proliferative effect of conditioned medium (CM) from ATCC 25611 versus heat-

inactivated CM on SG cells. (E) CM of strain ATCC 25611 was fractionated by

molecular weight using centrifugal filters: >30 kDa, 10–30 kDa, 3–10 kDa, and <3 kDa.

(F–G) CM from ATCC 25611 was applied to oral squamous carcinoma cell lines SCC-

15 and SAS for 24, 48, and 72 hours. All results are representative of three independent

experiments. Data are shown as mean $\pm$ SD. Statistical analysis was performed by one-

way ANOVA (A–C) or t-test (D). ns: not significant; *$p < 0.05$; **$p < 0.01$; ***$p <$

0.001; ****$p < 0.0001$.

Figure 2. InpA promotes oral cell proliferation via PAR-2 activation. (A) CE/LC-

MS identified elevated levels of InpA in strain ATCC 25611 compared to clinical strain

H31226, which did not stimulate SG cell proliferation. (B) SG cell proliferation in

response to CM or live *P. intermedia* (MOI 100), with or without 5 µM E64 (protease

inhibitor) or 40 µM FSLLRY-NH₂ (PAR-2 antagonist). (C-D) SCC-15 and SAS cell

proliferation following treatment with CM from ATCC 25611, with or without 0.5 µM

E64 or 20 µM FSLLRY-NH₂. (E) Five micrograms of recombinant InpA (rInpA) were

applied to SCC-15 cells, with or without E64 or FSLLRY-NH₂. PBS served as a vehicle

control; CM from ATCC 25611 served as a positive control. (F) Quantitative RT-PCR

analysis of *inpA* gene expression in ATCC 25611 and six clinical isolates. Strains

H26219-3, H29383, and H29703 were from oral cancer patients (CA), while H31082,

H29823, and H31226 were from control patients (CO). Expression levels were

calculated using the $2^{-\Delta\Delta Ct}$ method.

All results are representative of three independent experiments. Data are expressed as

mean ± SD and analyzed by one-way ANOVA. ns: not significant; $*p < 0.05$; $****p <$

0.0001.

Figure 3. InpA enhances OSCC cell migration through PAR-2 signaling. (A–B)

Wound healing assays in SCC-15 (A) and SAS (B) cells treated with standard medium

(D/F12), CM from ATCC 25611, or CM with 0.5 μM E64 or 20 μM FSLLRY-NH$_2$.

Cell migration was assessed at 2 and 4 hours and calculated as migration rate (μm/hr).

(C–D) Transwell migration assays in SCC-15 (C) and SAS (D) cells treated as above

for 24 hours. Migrated cells were quantified under a phase-contrast microscope. (E)

Transwell migration of SCC-15 cells treated with 5 μg/mL recombinant InpA (rInpA)

with or without E64 or FSLLRY-NH$_2$. All data represent triplicate experiments and are

shown as mean ± SD. One-way ANOVA was used for statistical analysis. ns: not

significant; $**p < 0.01$; $***p < 0.001$; $****p < 0.0001$.

Figure 4. Oral administration of *P. intermedia* increases colorectal tumor burden

in mice. (A) Schematic diagram of the azoxymethane (AOM)/dextran sulfate sodium

(DSS)-induced colorectal cancer (CRC) mouse model. Mice received a single

intraperitoneal injection of AOM (12.5 mg/kg), followed by three cycles of 2% DSS

in drinking water (5 days per cycle, separated by 16 days of regular water). During the

DSS cycles, 1×10^9 CFU of *P. intermedia* ATCC 25611 or *F. nucleatum* ATCC 23726

was administered via oral gavage. Mice were euthanized at week 10. (B)

Representative images of distal colon morphology from AOM/DSS mice with or

without bacterial infection. (C) Quantification of tumor number in the distal colon.

Each group included five mice. ns: not significant; $*p < 0.05$.

Acknowledgements. We thank the Laboratory Animal Center, College of Medicine at National Cheng Kung University (NCKU), Taiwan, accredited by AAALAC International and Taiwan Animal Consortium, for their support in caring for mice. We appreciate the participation of all patients in this study. We are grateful to all the unmentioned clinicians and recruiters. We also thank Dr. Chuan-Fa Chang and Dr. Jhen-Wei Ruan provided the cell lines.

Financial support: This work was supported by the National Science and Technology Council (Grants: 113-2314-B-006-043-MY3) awarded to J. W. C. and OSU-CHS Startup awarded to I.H.H.. The funding agencies played no role in study design, data collection or analysis, manuscript preparation, or decision to publish.

Authors' contributions: S.-M. C., Y.-J. C., W.-R. W. and N.-H. W. performed the experiments. S.-M. C. wrote this manuscript. I-H. H. assisted with the experimental design and reviewed this manuscript. C.-P. C. performed the microwestern array experiments and reviewed this manuscript. J.-R. H., J. S C., and J.-Y. C. provided clinical patients' saliva. J.-W. C. contributed to design, provide the concept, and supervise this manuscript. All authors have read and approved the final version of this manuscript.

References:

1.   Jhuang JR, Su SY, Chiang CJ, Yang YW, Lin LJ, Hsu TH, et al. Forecast of peak attainment and imminent decline after 2017 of oral cancer incidence in men in Taiwan. Sci Rep. 2022;12(1):5726.
2.   Tuominen H, Rautava J. Oral Microbiota and Cancer Development. Pathobiology. 2021;88(2):116-26.
3.   Zhang L, Liu Y, Zheng HJ, Zhang CP. The Oral Microbiota May Have Influence on Oral Cancer. Front Cell Infect Microbiol. 2019;9:476.
4.   Karmakar S, Kar A, Thakur S, Rao VUS. Periodontitis and oral Cancer-A striking link. Oral Oncology. 2020;106:104630.
5.   Kirst ME, Li EC, Alfant B, Chi Y-Y, Walker C, Magnusson I, et al. Dysbiosis

and alterations in predicted functions of the subgingival microbiome in chronic
periodontitis. Appl Environ Microbiol. 2015;81(2):783-93.
6.   Hsiao JR, Chang CC, Lee WT, Huang CC, Ou CY, Tsai ST, et al. The
interplay between oral microbiome, lifestyle factors and genetic polymorphisms in
the risk of oral squamous cell carcinoma. Carcinogenesis. 2018;39(6):778-87.
7.   Chattopadhyay I, Verma M, Panda M. Role of Oral Microbiome Signatures in
Diagnosis and Prognosis of Oral Cancer. Technol Cancer Res Treat.
2019;18:1533033819867354.
8.   Dominy SS, Lynch C, Ermini F, Benedyk M, Marczyk A, Konradi A, et al.
Porphyromonas gingivalis in Alzheimer's disease brains: Evidence for disease
causation and treatment with small-molecule inhibitors. Sci Adv.
2019;5(1):eaau3333.
9.   Wen L, Mu W, Lu H, Wang X, Fang J, Jia Y, et al. Porphyromonas gingivalis
Promotes Oral Squamous Cell Carcinoma Progression in an Immune
Microenvironment. Journal of Dental Research. 2020;99(6):666-75.
10.   Liu Y, Yuan X, Chen K, Zhou F, Yang H, Yang H, et al. Clinical significance
and prognostic value of Porphyromonas gingivalis infection in lung cancer. Transl
Oncol. 2021;14(1):100972.
11.   Inaba H, Sugita H, Kuboniwa M, Iwai S, Hamada M, Noda T, et al.
Porphyromonas gingivalis promotes invasion of oral squamous cell carcinoma
through induction of proMMP9 and its activation. Cell Microbiol. 2014;16(1):131-
45.
12.   Geng F, Zhang Y, Lu Z, Zhang S, Pan Y. Fusobacterium nucleatum Caused
DNA Damage and Promoted Cell Proliferation by the Ku70/p53 Pathway in Oral
Cancer Cells. DNA Cell Biol. 2020;39(1):144-51.
13.   Parhi L, Alon-Maimon T, Sol A, Nejman D, Shhadeh A, Fainsod-Levi T, et
al. Breast cancer colonization by Fusobacterium nucleatum accelerates tumor growth
and metastatic progression. Nature Communications. 2020;11(1):3259.
14.   Rubinstein MR, Wang X, Liu W, Hao Y, Cai G, Han YW. Fusobacterium
nucleatum promotes colorectal carcinogenesis by modulating E-cadherin/β-catenin
signaling via its FadA adhesin. Cell Host Microbe. 2013;14(2):195-206.
15.   Gur C, Ibrahim Y, Isaacson B, Yamin R, Abed J, Gamliel M, et al. Binding of
the Fap2 protein of Fusobacterium nucleatum to human inhibitory receptor TIGIT
protects tumors from immune cell attack. Immunity. 2015;42(2):344-55.
16.   Tompkins GR, Wood DP, Birchmeier KR. Detection and comparison of
specific hemin binding by Porphyromonas gingivalis and Prevotella intermedia. J
Bacteriol. 1997;179(3):620-6.
17.   Babich H, Tipton DA. In vitro response of human gingival epithelioid S-G
cells to minocycline. Toxicol In Vitro. 2002;16(1):11-21.
18.   TAKAHASHI K, KANAZAWA H, AKIYAMA Y, TAZAKI S,
TAKAHARA M, MUTO T, et al. Establishment and characterization of a cell
line(SAS) from poorly differentiated human squamous cell carcinoma of the tongue.
Journal of The Japanese Stomatological Society. 1989;38(1):20-8.
19.   Rheinwald JG, Beckett MA. Tumorigenic keratinocyte lines requiring
anchorage and fibroblast support cultured from human squamous cell carcinomas.
Cancer research. 1981;41 5:1657-63.
20.   Scheres N, de Vries TJ, Brunner J, Crielaard W, Laine ML, Everts V. Diverse
effects of Porphyromonas gingivalis on human osteoclast formation. Microb Pathog.
2011;51(3):149-55.
21.   Yang Y, Weng W, Peng J, Hong L, Yang L, Toiyama Y, et al. Fusobacterium

nucleatum Increases Proliferation of Colorectal Cancer Cells and Tumor Development in Mice by Activating Toll-Like Receptor 4 Signaling to Nuclear Factor-κB, and Up-regulating Expression of MicroRNA-21. Gastroenterology. 2017;152(4):851-66.e24.

22. Chen KY, Rathod J, Chiu YC, Chen JW, Tsai PJ, Huang IH. The Transcriptional Regulator Lrp Contributes to Toxin Expression, Sporulation, and Swimming Motility in Clostridium difficile. Front Cell Infect Microbiol. 2019;9:356.

23. Wang ST, Kuo CJ, Huang CW, Lee TM, Chen JW, Chen CS. OmpR coordinates the expression of virulence factors of Enterohemorrhagic Escherichia coli in the alimentary tract of Caenorhabditis elegans. Mol Microbiol. 2021;116(1):168-83.

24. Justus CR, Marie MA, Sanderlin EJ, Yang LV. Transwell In Vitro Cell Migration and Invasion Assays. Methods Mol Biol. 2023;2644:349-59.

25. Yu MR, Kim HJ, Park HR. Fusobacterium nucleatum Accelerates the Progression of Colitis-Associated Colorectal Cancer by Promoting EMT. Cancers (Basel). 2020;12(10).

26. Mallorquí-Fernández N, Manandhar SP, Mallorquí-Fernández G, Usón I, Wawrzonek K, Kantyka T, et al. A new autocatalytic activation mechanism for cysteine proteases revealed by Prevotella intermedia interpain A. J Biol Chem. 2008;283(5):2871-82.

27. Potempa M, Potempa J, Kantyka T, Nguyen KA, Wawrzonek K, Manandhar SP, et al. Interpain A, a cysteine proteinase from Prevotella intermedia, inhibits complement by degrading complement factor C3. PLoS Pathog. 2009;5(2):e1000316.

28. Byrne DP, Wawrzonek K, Jaworska A, Birss AJ, Potempa J, Smalley JW. Role of the cysteine protease interpain A of Prevotella intermedia in breakdown and release of haem from haemoglobin. Biochem J. 2009;425(1):257-64.

29. Liu Y, Wu Z, Nakanishi Y, Ni J, Hayashi Y, Takayama F, et al. Infection of microglia with Porphyromonas gingivalis promotes cell migration and an inflammatory response through the gingipain-mediated activation of protease-activated receptor-2 in mice. Sci Rep. 2017;7(1):11759.

30. Nonaka S, Nakanishi H. Secreted gingipains from Porphyromonas gingivalis induce microglia migration through endosomal signaling by protease-activated receptor 2. Neurochem Int. 2020;140:104840.

31. Chaffer CL, Weinberg RA. A perspective on cancer cell metastasis. Science. 2011;331(6024):1559-64.

32. Peng X, Cheng L, You Y, Tang C, Ren B, Li Y, et al. Oral microbiota in human systematic diseases. International Journal of Oral Science. 2022;14(1):14.

33. Read E, Curtis MA, Neves JF. The role of oral bacteria in inflammatory bowel disease. Nature Reviews Gastroenterology & Hepatology. 2021;18(10):731-42.

34. Kudra A, Muszyński D, Sobocki BK, Atzeni A, Carbone L, Kaźmierczak-Siedlecka K, et al. Insights into oral microbiome and colorectal cancer - on the way of searching new perspectives. Front Cell Infect Microbiol. 2023;13:1159822.

35. Wu N, Feng YQ, Lyu N, Wang D, Yu WD, Hu YF. Fusobacterium nucleatum promotes colon cancer progression by changing the mucosal microbiota and colon transcriptome in a mouse model. World J Gastroenterol. 2022;28(18):1981-95.

36. Moghimi M, Bakhtiari R, Mehrabadi JF, Jamshidi N, Jamshidi N, Siyadatpanah A, et al. Interaction of human oral cancer and the expression of virulence genes of dental pathogenic bacteria. Microb Pathog. 2020;149:104464.

37. Sheng J, Deng X, Zhang Q, Liu H, Wang N, Liu Z, et al. PAR-2 promotes

invasion and migration of esophageal cancer cells by activating MEK/ERK and PI3K/Akt signaling pathway. Int J Clin Exp Pathol. 2019;12(3):787-97.

38.   Inaba H, Kuboniwa M, Sugita H, Lamont RJ, Amano A. Identification of signaling pathways mediating cell cycle arrest and apoptosis induced by Porphyromonas gingivalis in human trophoblasts. Infect Immun. 2012;80(8):2847-57.

39.   Kato T, Tsuda T, Inaba H, Kawai S, Okahashi N, Shibata Y, et al. Porphyromonas gingivalis gingipains cause G(1) arrest in osteoblastic/stromal cells. Oral Microbiol Immunol. 2008;23(2):158-64.

40.   Stathopoulou PG, Galicia JC, Benakanakere MR, Garcia CA, Potempa J, Kinane DF. Porphyromonas gingivalis induce apoptosis in human gingival epithelial cells through a gingipain-dependent mechanism. BMC Microbiol. 2009;9:107.

41.   Park SY, Hwang BO, Lim M, Ok SH, Lee SK, Chun KS, et al. Oral-Gut Microbiome Axis in Gastrointestinal Disease and Cancer. Cancers (Basel). 2021;13(9).

42.   Zhang S, Kong C, Yang Y, Cai S, Li X, Cai G, et al. Human oral microbiome dysbiosis as a novel non-invasive biomarker in detection of colorectal cancer. Theranostics. 2020;10(25):11595-606.

43.   Lo CH, Wu DC, Jao SW, Wu CC, Lin CY, Chuang CH, et al. Enrichment of Prevotella intermedia in human colorectal cancer and its additive effects with Fusobacterium nucleatum on the malignant transformation of colorectal adenomas. J Biomed Sci. 2022;29(1):88.

44.   Bui FQ, Almeida-da-Silva CLC, Huynh B, Trinh A, Liu J, Woodward J, et al. Association between periodontal pathogens and systemic disease. Biomed J. 2019;42(1):27-35.

45.   Sparks Stein P, Steffen MJ, Smith C, Jicha G, Ebersole JL, Abner E, et al. Serum antibodies to periodontal pathogens are a risk factor for Alzheimer's disease. Alzheimers Dement. 2012;8(3):196-203.

46.   Li F, Ma C, Lei S, Pan Y, Lin L, Pan C, et al. Gingipains may be one of the key virulence factors of Porphyromonas gingivalis to impair cognition and enhance blood-brain barrier permeability: An animal study. J Clin Periodontol. 2024;51(7):818-39.

47.   Adams B, Nunes JM, Page MJ, Roberts T, Carr J, Nell TA, et al. Parkinson's Disease: A Systemic Inflammatory Disease Accompanied by Bacterial Inflammagens. Front Aging Neurosci. 2019;11:210.

48.   Romano F, Barbui A, Aimetti M. Periodontal pathogens in periodontal pockets and in carotid atheromatous plaques. Minerva Stomatol. 2007;56(4):169-79.

49.   Figuero E, Sánchez-Beltrán M, Cuesta-Frechoso S, Tejerina JM, del Castro JA, Gutiérrez JM, et al. Detection of periodontal bacteria in atheromatous plaque by nested polymerase chain reaction. J Periodontol. 2011;82(10):1469-77.

Fig. 1

Fig. 1

E

alive cell numbers(*10^4)

*
**
ns
ns
ns

time (hrs)

■ Untreated (SG cells)
■ ATCC 25611 (CM)
▨ ATCC 25611 (CM >30 kDa)
▨ ATCC 25611 (CM 10~30 kDa)
▥ ATCC 25611 (CM 3~10 kDa)
▦ ATCC 25611 (CM <3 kDa)

F

alive cell numbers(*10^4)

***

**

ns

0   24   48   72

time (hrs)

▼ Untreated (SCC-15 cells)
■ ATCC 25611 (CM)

G

alive cell numbers(*10^4)

***

*

ns

0   24   48   72

time (hrs)

▲ Untreated (SAS cells)
■ ATCC 25611 (CM)

Fig. 2

A

Strain	Protein	Prot_acc	Prot_hit_num	
ATCC 25611	Interpain A	A0A0D9NCF8 A9J7N5	273 273	13-fold
H31226		A0A0D9NCF8 A9J7N5 A0A0S3UMK9	21 21 21	

Fig. 2

Fig. 3

Fig. 4

A

AOM (12.5 mg/kg)

Fn/ Pi (1X10⁹ CFU)

Fn/ Pi (1X10⁹ CFU)

Fn/ Pi (1X10⁹ CFU)

Sacrifice

5 Days 2% DSS

5 Days 2% DSS

5 Days 2% DSS

0 1 2 3 4 5 6 7 8 9 10 Weeks

B

MOCK AOM/DSS

AOM/DSS + Fn AOM/DSS + Pi

C

Colon tumor number