Title: Medicinal chemistry of benzo[d]isoxazole-4,7-dione analogues identifies the

ribosomal small subunit as a target for specific mycobacterial translational fidelity

Authors: Jie Wu[1], Swarnava Chaudhuri[2], Sarah Caldwell Feid[3], Miaomiao Pan[2],

Qingwen Kawaji[2], Gang Liu[1*] and Babak Javid[2, 3*]

JW, SC and SCF contributed equally to this work

GL and BJ contributed equally to this work

[*]To whom correspondence should be addressed: babak.javid@ucsf.edu (BJ) or

gangliu27@tsinghua.edu.cn (GL)

Running title: Benzo[d]isoxazole-4,7-dione analogues decrease mycobacterial

mistranslation

Abstract

Mycobacteria have high rates of error in the translation of glutamine and asparagine

codons due to a two-step indirect tRNA aminoacylation pathway involving the essential

amidotransferase GatCAB. This specific mycobacterial mistranslation is both necessary

and sufficient for tolerance to the antibiotic rifampicin. Furthermore, we identified the

natural product kasugamycin as a small molecule that could decrease this type of

translational error, and potentiate rifampicin-mediated killing of mycobacteria, but

whether other small molecules can decrease this specific type of mistranslation is

unknown. We screened a library of synthetic small molecules using a pathway-specific

whole-cell screen in a reporter strain of *Mycobacterium smegmatis* for their ability to

decrease specific mycobacterial mistranslation. We synthesized 110 analogs of a parent

hit compound and analyzed their bioactivity. To identify the cellular target of the

compounds, we biotinylated active and inactive analogs and performed pull-down and

mass-spectrometry of bound cellular proteins. We identified benzo[d]isoxazole-4,7-

diones as a class of small molecules able to decrease mycobacterial mistranslation. Of the

110 analogs synthesized, we identified 10 with superior potency to the parent compound

and analyzed the compounds to infer structure-activity relationships. Pull-down and mass

spectrometry of bound cellular proteins identified RpS5 as a potential target of active

compounds. Our study identified synthetic small molecules able to decrease

mycobacterial mistranslation and confirmed that the small ribosomal subunit can be

targeted to decrease mistranslation rates arising from physiologically mischarged tRNAs.

Targeting mistranslation may be a potentially attractive means of targeting antibiotic-

tolerant bacteria.

Keywords: mistranslation; translational fidelity; small molecule; antibiotic tolerance;

medicinal chemistry; mycobacteria

INTRODUCTION

Genetically susceptible bacteria may still withstand killing by antibiotics, a phenomenon known as antibiotic tolerance.[1, 2] There are several potential mechanisms for tolerance, and these may reflect different physiological states of the bacteria.[3-5] Non-replicating persisters are usually multi-drug tolerant and have a relatively, but not entirely, quiescent physiological state.[5-8] We recently described a form of tolerance in mycobacteria to the first-line antibiotic rifampicin, which we termed RNA polymerase-specific phenotypic resistance (RSPR).[9-11] By contrast with non-replicating persisters, the antibiotic-tolerant bacteria in RSPR are able to grow in bulk-lethal concentrations of antibiotic that can kill the majority of bacilli in the culture.[9] Importantly, RSPR was apparent in *Mycobacterium tuberculosis* freshly isolated from sputum from a small number of patients, suggesting that it may be a clinically relevant mode of antibiotic tolerance.[10, 12]

We had identified at least two mechanisms that contribute to mycobacterial RSPR.[10, 11] One of these mechanisms involved modulation of translational fidelity in the indirect tRNA aminoacylation pathway – or specific mycobacterial mistranslation.[10, 13] Unlike cytosolic eukaryotic gene translation, the vast majority of bacteria, excepting *Escherichia coli* and a few related proteobacteria, lack the specific tRNA aminoacyl synthetases for either glutamine or asparagine or both.[14-18] All mycobacteria lack both glutaminyl- and asparaginyl-synthetases. Instead, they undergo a two-step pathway to specifically aminoacylate glutamine and asparagine tRNAs. First, a non-discriminatory glutamyl- (or aspartyl-) synthetase physiologically misacylates the glutamine or asparagine tRNA respectively to Glu-tRNA^{Gln} and Asp-tRNA^{Asn}. This misacylated complex is then

specifically recognized by the essential glutamine amidotransferase GatCAB.[10] Using

cytosolic glutamine as an amide donor, GatCAB amidates the misacylated tRNAs to their

cognate Gln-tRNA^{Gln} and Asn-tRNA^{Asn} respectively, thereby allowing preservation of

the genetic code. Therefore, mutations that alter GatCAB structure and stability can

increase mistranslation in mycobacteria.[10, 13] Moreover, we had identified that naturally

occurring mutations in *gatA* in clinical isolates of *Mycobacterium tuberculosis* both

increased rates of specific mistranslation and tolerance to rifampicin.[10] Furthermore,

GatCAB could be limiting in wild-type mycobacteria, and sorting cells by decreased

expression of wild-type *gatCAB* in an isogenic population identified subpopulations with

both increased rates of mistranslation and antibiotic tolerance.[10] We also showed that

mistranslation of a specific, critical, residue in the rifampicin binding region of the

cellular target of rifampicin, RNA polymerase (RNAP), contributed substantially to

RSPR in wild-type mycobacteria.[10] Together, these data compellingly suggested that

specific mycobacterial mistranslation was an important contributor to RSPR.

Since mycobacterial mistranslation contributed to RSPR even in isogenic wild-type

bacteria, we hypothesized that this pathway may be amenable to pharmacological

intervention. We identified the natural product kasugamycin as a small molecule that

could both decrease specific mycobacterial mistranslation and potentiate rifampicin both

in vitro and in a murine model of tuberculosis.[19] However, other than the naturally

occurring peptide Edeine, which showed some activity in our assays,[19] no other

molecules capable of decreasing this specific form of bacterial mistranslation, which is

mechanistically distinct from ribosomal decoding errors,[10] have been identified. We

decided to perform a pathway-specific whole-cell screen[20] that could identify synthetic

small molecules that could target mycobacterial mistranslation. In this study, we utilized

our mistranslation inhibition assay, which had earlier been used to identify

kasugamycin's ability to increase mycobacterial translational fidelity,[19, 21] in a high-

throughput format. Here, we identify analogs of benzo[d]isoxazole-4,7-diones that can

decrease mistranslation rates and potentiate rifampicin *in vitro*. Furthermore, we identify

the potential target of the most potent analog as the small ribosomal subunit protein RpS5.

MATERIALS & METHODS

Bacterial strains, culture, antibiotics and test compounds

Wildtype *M. smegmatis* mc^2-155 and its mistranslation reporter derivative strain were

grown in Middlebrook 7H9 broth supplemented with 0.2% glycerol, 0.05% Tween-80, 10%

Albumin-dextrose-salt (ADS), and on Luria-Bertani agar (LB agar) for plate assays.

Rifampicin and kasugamycin (Ksg) were purchased from Sigma. Rifampicin was

dissolved in dimethyl sulfoxide (DMSO); Ksg was dissolved in water, and filter sterilized.

The test compounds for screening were all dissolved in DMSO.

Screening of compounds that decrease mycobacterial mistranslation

Screening of the chemical library to identify compounds that decrease mycobacterial

mistranslation was done using the previously developed N-luc/GFP mistranslation

assay.[11] Briefly, the mistranslation reporter *M. smegmatis* strain was grown in 7H9 broth

containing hygromycin (50µg/ml) and kanamycin (20µg/ml) at 37˚C to late log phase

(OD$_{600}$ = 2.0). N-luc expression was induced by adding anhydrotetracycline (ATc, 50

ng/ml), and the bacterial culture was immediately aliquoted into a 96 well plate (100 µl,

i.e. approximately 6×10^7 cells per well). 50 µM of each of the test compound or 50 µg/ml

of Ksg as positive control were added, and the cultures incubated for 16 hours at 37°C

with shaking. Then the cultures were transferred to a 96-well black plate and GFP

fluorescence was measured by Fluoroskan Ascent FL Fluorometer and Luminometer

(Thermo Scientific). The plate was then centrifuged (4000 rpm for 10 minutes), and

supernatants transferred to a 96-well white luminescence plate. The nano-luciferase assay

was performed using Nano-Glo luciferase assay kit (Promega), and luminescence

measured by the same machine. For confirmation of hits in *Mycobacterium bovis*-BCG

(BCG), the same protocol as above was followed except compounds were co-incubated

with bacteria for 24 hours prior to analysis.

Rifampicin specific phenotypic resistance (RSPR) assay

RSPR assay was done as previously described.[19] Briefly, *M. smegmatis* mc^2-155 was

grown to stationary phase, after which the cultures were serially diluted and spread on

LB-agar plates containing rifampicin (50µg/ml) only, or rifampicin and 50µM of each

test compound. LB agar containing rifampicin and Ksg (100µg/ml) was used as positive

control. The plates were incubated at 37°C for 5 days after which the number of colony

forming units (cfu) were enumerated. The number of bacterial cells in the inoculum was

calculated by counting colonies after spreading serial dilutions of the culture on LB-agar

plates without any antibiotic.

Target identification of 9788

To identify the intracellular target(s) of compounds that decrease mycobacterial

mistranslation,   9788, a biotinylated version of compound 9787 that decreased

mycobacterial mistranslation, and 9791, a biotinylated version of compound 9789 that cannot decrease mycobacterial mistranslation were used. A protocol modified from an earlier study to identify mycobacterial drug targets was used.[22] To prepare mycobacterial cell lysate, *M. smegmatis* mc^2-155 was grown in 7H9 broth up to OD_{600} 1.0, and then centrifuged at 4000 rpm for 10 minutes. The pellet was resuspended in phosphate-buffered saline (PBS) containing a tablet of protease inhibitor (Roche), and the cells were lysed by beads beating twice for 45 seconds each. Then the mixture was centrifuged at 15000 rpm for 15 minutes, the supernatant was collected and added to EZ view^{TM} Red streptavidin affinity gel (Sigma Aldrich), pre-equilibrated with the lysis buffer (PBS plus protease inhibitor cocktail). The mixture was incubated at 4˚C for an hour in an orbital shaker, and then centrifuged at 15000 rpm for an hour to collect the supernatant. The supernatant was then distributed to five tubes, and the protein concentration of each tube was adjusted to 1 mg/ml. The following compounds were then added one to each tube: 1) 1% DMSO, 2) 100 μM 9788, 3) 100 μM 9788 + 500 μM 9787, 4) 100 μM 9791, 5) 100 μM 9791 + 500 μM 9789. After 1 hour incubation at room temperature, the mixture from each tube was added to pre-equilibrated EZ view^{TM} Red streptavidin affinity gel, and incubated at 4˚C for an hour in an orbital shaker. Then, the mixtures were centrifuged at 15000 rpm for 20 mins. at 4˚C. The pellets were then resuspended in Laemmli buffer, ran in SDS PAGE, and sent for mass-spectrometry at the Tsinghua University Mass-Spectrometry facility.

RESULTS

We developed a reporter system in *M. smegmatis* to sensitively measure relative

mistranslation of asparagine to aspartate – one of the two form of mistranslation

implicated in the indirect tRNA aminoacylation pathway (Figure 1A).[19, 21] We then used

this strain to screen 9000 synthetic heterocyclic small molecules in 96 well-plate format

(Fig. 1B). By using a reporter strain, a common pitfall of *in vitro* screens is avoided: such

screens identify many hits that are ultimately inactive against whole cells due to the

complex and thick mycobacterial cell wall and envelope.[20] However, unlike a traditional

whole-cell screen in which all hits that could kill bacteria are identified, leading to 're-

discovery' of common antimicrobials or identification of non-specifically toxic

molecules, the system retains the advantage of being "pathway specific". Only hits that

decrease specific mycobacterial mistranslation are identified.

The basis of the screen was as follows: Nluc luciferase was mutated at a critical aspartate

residue (D140) to asparagine, which resulted in >99% loss of activity.[19, 21] The

'physiological' mycobacterial mistranslation rate of asparagine->aspartate resulted in a

baseline detectable Nluc activity of ~1% of the wild-type enzyme (corrected for cell

density and protein expression by normalizing to GFP fluorescence). If a small molecule

decreased mistranslation rates, the corrected Nluc/GFP would be reduced below the

baseline physiological rates (**Fig 1A**). Hits were classified as those molecules that

decreased mistranslation rates by > 1 standard deviation. Due to the inherent noisy signal

in measuring translational error, screen performance was variable (**Fig 1B**). 30% of the

plates had a z' <0.34 and were not considered in the analysis of hit compounds.

Upon completion of the screen, we identified 45 hits from well-performing plates. Of

note, 3 were from the same chemical scaffold, benzo[d]isoxazole-4,7-diones (**Fig. 1C**).

These 3 compounds, **7006**, **7013** and **7020**, were retested for their ability to reduce

mycobacterial mistranslation and RSPR, confirming them as valid hits (**Fig, 2A and 2B**).

We proceeded to analyze this scaffold further by medicinal chemistry, with the goal of

developing tool compounds to test the effects of modulating mycobacterial mistranslation

on antibiotic tolerance. Therefore, we prioritized compounds without major direct

antimicrobial activity by themselves in order to deconvolute antimicrobial activity from

anti-mistranslation activity. We synthesized 110 analogs of **7020**, chosen because it

modulates mistranslation without antimicrobial activity and is amenable to medicinal

chemistry analysis, of which 10 compounds (**Fig. 3**), showed superior potency to the

parent compound, **7020**, both in terms of decreasing mistranslation, and potentiating

rifampicin in an assay used to measure RSPR (**Table 1**). The compounds also had

minimal direct antimicrobial activity by themselves (**Table 1**), making them suitable for

dissecting the contribution of targeting mycobacterial mistranslation on antibiotic

tolerance without confounding issues due to inhibitory activity.

Our preliminary investigation of structure-activity relationships (SAR) is summarized in

Fig. 4 and **Fig. 5**. Our SAR analysis showed that the benzoquinone part of the compound

is essential for its ability to reduce mistranslation; all analogs of 7020 synthesized

without the benzoquinone structure are poor inhibitors of mistranslation (**Fig. 4**). We

further observed that the methyl groups at positions 5 and 6 of the benzo[d]isoxazole-4,7-

diones scaffold are not necessary for their bioactivities in terms of inhibition of

mistranslation and RSPR (**Fig. 5**). Replacement of these methyl groups with other

functional groups such as chlorine, as in **9744**, or with an aromatic ring, as in **9775**, **9782**,

9783 and **9787**, improved bioactivities of the compounds. (**Table. 1** and **Fig. 5**).

We chose **9787** as the most promising candidate for attempting to determine the cellular

target(s) of the hits. **9787** was more potent than the initial hit **7020** (**Fig. 6A**). Moreover,

9787 was able to decrease mistranslation in another closely related mycobacterial species,

M. bovis BCG (**Fig. 6B**).

We wished to identify the cellular target by which **9787** mediated a decrease in specific

mistranslation rates. Since the compounds lacked significant antimicrobial activity by

themselves, it was not possible to raise mutants resistant to the compound and sequence

for mutations in the genome to identify potential targets.[23] We, therefore, decided to

identify the target by a modified pull-down and mass-spectrometry approach.[23-26] For this

chemical proteomic procedure, **9787** was biotinylated to yield **9788**, which retained

activity (**Fig. 7A and B**). An inactive analog, **9789** from the initial medicinal chemistry

approach was also biotinylated to make inactive **9791** (**Fig. 7A and C**). We incubated

9788 or **9791** with mycobacterial cell lysate. To increase the specificity of the identified

target protein(s), we also included conditions in which 5-fold molar excess of non-

biotinylated **9787** or **9789** were added to the tubes in addition to the biotinylated

compound (**Fig. 7D** and see methods). Following incubation, the biotinylated compounds

and putative cellular binding partners were pulled down using a streptavidin column,

washed, and then the eluates were resolved on an SDS-PAGE gel and bands cut out and

sent for analysis by mass-spectrometry (Supplementary Data 1). Our rationale for

identifying the target for **9787** was as follows: When the active biotinylated compound

(**9788**) was used, the quality score of the target protein should be significantly higher than

that obtained with the inactive biotinylated compound (**9791**). Moreover, in the presence

of competing non-biotinylated compounds, less target protein(s) are expected to bind to

the streptavidin column. Hence the quality score of the target protein obtained with the

active **9788** alone should be significantly higher than that obtained with a mixture of

9788 and excess **9787**. Furthermore, no such phenomenon should be observed in case of

the inactive compound (**9789**) and its biotinylated form (**9791**). We plotted the ratio of

protein quality scores obtained with the active and inactive biotinylated compounds

(**9788/9791**) against the ratio of protein quality scores obtained with **9788** alone, and

9788 and excess **9787** (9788/[9788 + excess 9787]). Out of 1925 proteins, only the 30S

ribosomal protein S5 (Rps5) derived from *rpsE* gene, could fulfill all the criteria and

significantly stood out against the others. (**Fig. 7E**). These data strongly suggest that

RpS5 is the target of active **9787**.

DISCUSSION

Phenotypic screening has been extensively used in developing novel drug candidates

against *Mycobacterium tuberculosis*.[27-30] Phenotypic screening is, in general considered a

better approach than biochemical screening, which often yields 'hits' that cannot actually

penetrate the bacterial cell. This is particularly true for mycobacteria that have a complex

capsule. On the other hand, hits obtained from whole-cell phenotypic screening exhibit

their effect after penetrating the bacterial cell membrane. Hence, they are usually better

candidates for drug development than those obtained from biochemical screening.

In this study, we have successfully expanded one of the luciferase-based mycobacterial

mistranslation assays[19] developed in our laboratory into a larger-scale screening platform.

Throughout the screening procedure, kasugamycin was used as a positive control to

enhance the validity of the screen. Although the performance of the screening platform

was variable, it nonetheless identified several hits that could be further validated and

characterized. Most of the hits also significantly decreased mycobacterial RSPR. This

further confirms the connection between mistranslation and RSPR in mycobacteria.

Our chemical proteomic approach was effective in identifying the 30S ribosomal protein

S5 (RpS5) as the molecular target of our test compound **9787**. During mycobacterial

protein translation, RpS5 plays a crucial role in the binding of mRNA to the ribosome.

Moreover, it is necessary for proper ribosomal decoding of the mRNA.[31] Our results

show that binding of a small molecule to RpS5 can further also increase fidelity of

ribosomal discrimination of mischarged aminoacyl-tRNAs. The mechanisms by which

the small ribosomal subunit, concerned with ribosomal decoding, contributes to

discrimination of aminoacyl identity of tRNAs at the ribosome are unknown and the

subject of ongoing investigation. Interestingly, kasugamycin, our previously identified

inhibitor of mycobacterial mistranslation, also targets the 30S ribosomal subunit. In

Escherichia coli, it binds near the residues A794 and G926 of the 16S rRNA.[32, 33] Due to

the universally conserved nature of these two residues, kasugamycin likely binds to a

similar location in mycobacteria. Hence, the 30S ribosomal subunit appears to be a

common target for compounds that decrease mistranslation in mycobacteria. The

ribosome itself has been shown to have proof-reading activity of codon•anti-codon

interactions during translation to ensure fidelity of protein synthesis.[34] However, the

ability of the ribosome to detect mischarged tRNAs is less well understood. It will be

interesting to study ribosomal conformational changes after binding of molecules such as

9787 to RpS5 in the future.

Although maintaining translational fidelity is believed to be essential for all living

systems, bacteria have been shown to tolerate up to 10% mistranslation of a specific

codon without major deleterious effects.[35] Moreover, mistranslation has been proposed to

be a mechanism by which bacteria such as *Mycobacterium* adapt to environmental

stresses such as low pH and the presence of rifampicin. Under such conditions,

mistranslation results in increased phenotypic heterogeneity and improves bacterial

survival.[36, 37]

Rifampicin is a crucial antibiotic in anti-TB chemotherapy. Its introduction in TB therapy

has drastically reduced the treatment time. However, due to drug-tolerance of the bacteria,

the standard treatment time for drug-susceptible TB is still 6 months, and thus has room

for further improvement. Empirical attempts have been made to improve the efficacy of

rifampicin by changing the therapeutic concentrations, as well as by using new

combinations with other drugs.[38-40] We, for the first time, had earlier proposed a

mechanism of enhancing the efficacy of rifampicin in TB treatment by targeting

mycobacterial mistranslation.[19] In this work we have further expanded our search for

compounds that can decrease mycobacterial mistranslation. Importantly, this gives us

confidence in using our platform to identify and develop future drug candidates.

Moreover, the 30S ribosomal subunit can be used as an attractive drug target for reducing

the burden of mistranslation-mediated antibiotic-tolerant bacteria.

ASSOCIATED CONTENT

Supporting information available: The experimental procedures of chemistry and

spectroscopic data of all compounds.

WORKS CITED

1.   Gold B, Nathan C. Targeting Phenotypically Tolerant Mycobacterium tuberculosis.
Microbiology spectrum. 2017;5(1). doi: 10.1128/microbiolspec.TBTB2-0031-2016. PubMed
PMID: 28233509; PMCID: PMC5367488.
2.   Keren I, Kaldalu N, Spoering A, Wang Y, Lewis K. Persister cells and tolerance to
antimicrobials. FEMS Microbiol Lett. 2004;230(1):13-8. doi: 10.1016/S0378-1097(03)00856-5.
PubMed PMID: 14734160.
3.   Aldridge BB, Keren I, Fortune SM. The Spectrum of Drug Susceptibility in Mycobacteria.
Microbiology spectrum. 2014;2(5). doi: 10.1128/microbiolspec.MGM2-0031-2013. PubMed
PMID: 26104360.
4.   Balaban NQ, Gerdes K, Lewis K, McKinney JD. A problem of persistence: still more
questions than answers? Nat Rev Microbiol. 2013;11(8):587-91. PubMed PMID: 24020075.
5.   Kaldalu N, Hauryliuk V, Tenson T. Persisters-as elusive as ever. Appl Microbiol Biotechnol.
2016;100(15):6545-53. Epub 20160604. doi: 10.1007/s00253-016-7648-8. PubMed PMID:
27262568; PMCID: PMC4939303.
6.   Maisonneuve E, Gerdes K. Molecular mechanisms underlying bacterial persisters. Cell.
2014;157(3):539-48. doi: 10.1016/j.cell.2014.02.050. PubMed PMID: 24766804.
7.   Orman MA, Brynildsen MP. Dormancy is not necessary or sufficient for bacterial
persistence. Antimicrob Agents Chemother. 2013;57(7):3230-9. doi: 10.1128/AAC.00243-13.
PubMed PMID: 23629720; PMCID: 3697331.
8.   Wakamoto Y, Dhar N, Chait R, Schneider K, Signorino-Gelo F, Leibler S, McKinney JD.
Dynamic persistence of antibiotic-stressed mycobacteria. Science. 2013;339(6115):91-5. Epub
2013/01/05. doi: 10.1126/science.1229858

339/6115/91 [pii]. PubMed PMID: 23288538.
9.   Javid B, Sorrentino F, Toosky M, Zheng W, Pinkham JT, Jain N, Pan M, Deighan P, Rubin
EJ. Mycobacterial mistranslation is necessary and sufficient for rifampicin phenotypic resistance.
Proc Natl Acad Sci U S A. 2014;111(3):1132-7. doi: 10.1073/pnas.1317580111. PubMed PMID:
24395793; PMCID: PMC3903211.
10.   Su HW, Zhu JH, Li H, Cai RJ, Ealand C, Wang X, Chen YX, Kayani MU, Zhu TF,
Moradigaravand D, Huang H, Kana BD, Javid B. The essential mycobacterial amidotransferase
GatCAB is a modulator of specific translational fidelity. Nat Microbiol. 2016;1(11):16147. doi:
10.1038/nmicrobiol.2016.147. PubMed PMID: 27564922.
11.   Zhu JH, Wang BW, Pan M, Zeng YN, Rego H, Javid B. Rifampicin can induce antibiotic
tolerance in mycobacteria via paradoxical changes in rpoB transcription. Nature
communications. 2018;9(1):4218. doi: 10.1038/s41467-018-06667-3. PubMed PMID: 30310059;
PMCID: PMC6181997.
12.   Li YY, Cai RJ, Yang JY, Hendrickson TL, Xiang Y, Javid B. Clinically Relevant Mutations of
Mycobacterial GatCAB Inform Regulation of Translational Fidelity. mBio. 2021;12(4):e0110021.
Epub 20210706. doi: 10.1128/mBio.01100-21. PubMed PMID: 34225495; PMCID: PMC8406222.
13.   Cai RJ, Su HW, Li YY, Javid B. Forward Genetics Reveals a gatC-gatA Fusion Polypeptide
Causes Mistranslation and Rifampicin Tolerance in Mycobacterium smegmatis. Front Microbiol.
2020;11:577756. Epub 2020/10/20. doi: 10.3389/fmicb.2020.577756. PubMed PMID: 33072044;
PMCID: PMC7541841.
14.   Bailly M, Blaise M, Lorber B, Becker HD, Kern D. The transamidosome: a dynamic
ribonucleoprotein particle dedicated to prokaryotic tRNA-dependent asparagine biosynthesis.
Mol Cell. 2007;28(2):228-39. doi: 10.1016/j.molcel.2007.08.017. PubMed PMID: 17964262.

15.   Curnow AW, Hong K, Yuan R, Kim S, Martins O, Winkler W, Henkin TM, Soll D. Glu-
tRNAGln amidotransferase: a novel heterotrimeric enzyme required for correct decoding of
glutamine codons during translation. Proc Natl Acad Sci U S A. 1997;94(22):11819-26. doi:
10.1073/pnas.94.22.11819. PubMed PMID: 9342321; PMCID: PMC23611.
16.   Huot JL, Fischer F, Corbeil J, Madore E, Lorber B, Diss G, Hendrickson TL, Kern D,
Lapointe J. Gln-tRNAGln synthesis in a dynamic transamidosome from Helicobacter pylori,
where GluRS2 hydrolyzes excess Glu-tRNAGln. Nucleic Acids Res. 2011;39(21):9306-15. doi:
10.1093/nar/gkr619. PubMed PMID: 21813455; PMCID: 3241645.
17.   Rathnayake UM, Wood WN, Hendrickson TL. Indirect tRNA aminoacylation during
accurate translation and phenotypic mistranslation. Curr Opin Chem Biol. 2017;41:114-22. Epub
2017/11/21. doi: 10.1016/j.cbpa.2017.10.009. PubMed PMID: 29156229.
18.   Sheppard K, Yuan J, Hohn MJ, Jester B, Devine KM, Soll D. From one amino acid to
another: tRNA-dependent amino acid biosynthesis. Nucleic Acids Res. 2008;36(6):1813-25. Epub
20080205. doi: 10.1093/nar/gkn015. PubMed PMID: 18252769; PMCID: PMC2330236.
19.   Chaudhuri S, Li L, Zimmerman M, Chen Y, Chen YX, Toosky MN, Gardner M, Pan M, Li YY,
Kawaji Q, Zhu JH, Su HW, Martinot AJ, Rubin EJ, Dartois VA, Javid B. Kasugamycin potentiates
rifampicin and limits emergence of resistance in Mycobacterium tuberculosis by specifically
decreasing mycobacterial mistranslation. Elife. 2018;7. doi: 10.7554/eLife.36782. PubMed PMID:
30152756; PMCID: PMC6160228.
20.   Moreira W, Ngan GJ, Low JL, Poulsen A, Chia BC, Ang MJ, Yap A, Fulwood J, Lakshmanan
U, Lim J, Khoo AY, Flotow H, Hill J, Raju RM, Rubin EJ, Dick T. Target mechanism-based whole-cell
screening identifies bortezomib as an inhibitor of caseinolytic protease in mycobacteria. mBio.
2015;6(3):e00253-15. Epub 20150505. doi: 10.1128/mBio.00253-15. PubMed PMID: 25944857;
PMCID: PMC4436076.
21.   Chen YX, Pan M, Chen YM, Javid B. Measurement of Specific Mycobacterial
Mistranslation Rates with Gain-of-function Reporter Systems. J Vis Exp. 2019(146). doi:
10.3791/59453. PubMed PMID: 31081825.
22.   Cox JA, Abrahams KA, Alemparte C, Ghidelli-Disse S, Rullas J, Angulo-Barturen I, Singh A,
Gurcha SS, Nataraj V, Bethell S, Remuinan MJ, Encinas L, Jervis PJ, Cammack NC, Bhatt A, Kruse
U, Bantscheff M, Futterer K, Barros D, Ballell L, Drewes G, Besra GS. THPP target assignment
reveals EchA6 as an essential fatty acid shuttle in mycobacteria. Nat Microbiol. 2016;1:15006.
Epub 20160118. doi: 10.1038/nmicrobiol.2015.6. PubMed PMID: 27571973.
23.   La Rosa V, Poce G, Canseco JO, Buroni S, Pasca MR, Biava M, Raju RM, Porretta GC,
Alfonso S, Battilocchio C, Javid B, Sorrentino F, Ioerger TR, Sacchettini JC, Manetti F, Botta M, De
Logu A, Rubin EJ, De Rossi E. MmpL3 is the cellular target of the antitubercular pyrrole derivative
BM212.   Antimicrob   Agents   Chemother.   2012;56(1):324-31.   Epub   20111024.   doi:
10.1128/AAC.05270-11. PubMed PMID: 22024828; PMCID: PMC3256021.
24.   Abrahams KA, Chung CW, Ghidelli-Disse S, Rullas J, Rebollo-Lopez MJ, Gurcha SS, Cox JA,
Mendoza A, Jimenez-Navarro E, Martinez-Martinez MS, Neu M, Shillings A, Homes P, Argyrou A,
Casanueva R, Loman NJ, Moynihan PJ, Lelievre J, Selenski C, Axtman M, Kremer L, Bantscheff M,
Angulo-Barturen I, Izquierdo MC, Cammack NC, Drewes G, Ballell L, Barros D, Besra GS, Bates RH.
Identification of KasA as the cellular target of an anti-tubercular scaffold. Nat Commun.
2016;7:12581. Epub 20160901. doi: 10.1038/ncomms12581. PubMed PMID: 27581223; PMCID:
PMC5025758.
25.   Abrahams KA, Cox JA, Spivey VL, Loman NJ, Pallen MJ, Constantinidou C, Fernandez R,
Alemparte C, Remuinan MJ, Barros D, Ballell L, Besra GS. Identification of novel imidazo[1,2-
a]pyridine inhibitors targeting M. tuberculosis QcrB. PLoS One. 2012;7(12):e52951. Epub
20121231. doi: 10.1371/journal.pone.0052951. PubMed PMID: 23300833; PMCID: PMC3534098.

26.   Bantscheff M, Hopf C, Savitski MM, Dittmann A, Grandi P, Michon AM, Schlegl J,
Abraham Y, Becher I, Bergamini G, Boesche M, Delling M, Dumpelfeld B, Eberhard D,
Huthmacher C, Mathieson T, Poeckel D, Reader V, Strunk K, Sweetman G, Kruse U, Neubauer G,
Ramsden NG, Drewes G. Chemoproteomics profiling of HDAC inhibitors reveals selective
targeting of HDAC complexes. Nat Biotechnol. 2011;29(3):255-65. Epub 20110123. doi:
10.1038/nbt.1759. PubMed PMID: 21258344.
27.   Alumasa JN, Manzanillo PS, Peterson ND, Lundrigan T, Baughn AD, Cox JS, Keiler KC.
Ribosome Rescue Inhibitors Kill Actively Growing and Nonreplicating Persister Mycobacterium
tuberculosis Cells. ACS Infect Dis. 2017;3(9):634-44. doi: 10.1021/acsinfecdis.7b00028. PubMed
PMID: 28762275; PMCID: PMC5594445.
28.   Burke C, Jankute M, Moynihan P, Gonzalez Del Rio R, Li X, Esquivias J, Lelievre J, Cox JAG,
Sacchettini J, Besra GS. Development of a novel secondary phenotypic screen to identify hits
within the mycobacterial protein synthesis pipeline. FASEB Bioadv. 2020;2(10):600-12. Epub
20200820. doi: 10.1096/fba.2020-00022. PubMed PMID: 33089076; PMCID: PMC7566049.
29.   Sukheja P, Kumar P, Mittal N, Li SG, Singleton E, Russo R, Perryman AL, Shrestha R,
Awasthi D, Husain S, Soteropoulos P, Brukh R, Connell N, Freundlich JS, Alland D. A Novel Small-
Molecule   Inhibitor   of   the   Mycobacterium   tuberculosis   Demethylmenaquinone
Methyltransferase MenG Is Bactericidal to Both Growing and Nutritionally Deprived Persister
Cells. MBio. 2017;8(1). doi: 10.1128/mBio.02022-16. PubMed PMID: 28196957; PMCID:
PMC5312080.
30.   Zheng P, Somersan-Karakaya S, Lu S, Roberts J, Pingle M, Warrier T, Little D, Guo X,
Brickner SJ, Nathan CF, Gold B, Liu G. Synthetic calanolides with bactericidal activity against
replicating and nonreplicating Mycobacterium tuberculosis. J Med Chem. 2014;57(9):3755-72.
doi: 10.1021/jm4019228. PubMed PMID: 24694175.
31.   Hentschel J, Burnside C, Mignot I, Leibundgut M, Boehringer D, Ban N. The Complete
Structure of the Mycobacterium smegmatis 70S Ribosome. Cell reports. 2017;20(1):149-60. doi:
10.1016/j.celrep.2017.06.029. PubMed PMID: 28683309.
32.   Schuwirth BS, Day JM, Hau CW, Janssen GR, Dahlberg AE, Cate JH, Vila-Sanjurjo A.
Structural   analysis   of   kasugamycin   inhibition   of   translation.   Nat   Struct   Mol   Biol.
2006;13(10):879-86. doi: 10.1038/nsmb1150. PubMed PMID: 16998486; PMCID: PMC2636691.
33.   Wilson DN. Ribosome-targeting antibiotics and mechanisms of bacterial resistance. Nat
Rev Microbiol. 2014;12(1):35-48. doi: 10.1038/nrmicro3155. PubMed PMID: 24336183.
34.   Zaher HS, Green R. Quality control by the ribosome following peptide bond formation.
Nature. 2009;457(7226):161-6. Epub 2008/12/19. doi: nature07582 [pii]

10.1038/nature07582. PubMed PMID: 19092806.
35.   Ruan B, Palioura S, Sabina J, Marvin-Guy L, Kochhar S, Larossa RA, Soll D. Quality control
despite mistranslation caused by an ambiguous genetic code. Proc Natl Acad Sci U S A.
2008;105(43):16502-7. Epub 2008/10/24. doi: 0809179105 [pii]

10.1073/pnas.0809179105. PubMed PMID: 18946032.
36.   Mohler K, Ibba M. Translational fidelity and mistranslation in the cellular response to
stress. Nat Microbiol. 2017;2:17117. Epub 2017/08/25. doi: 10.1038/nmicrobiol.2017.117.
PubMed PMID: 28836574; PMCID: PMC5697424.
37.   Ribas de Pouplana L, Santos MA, Zhu JH, Farabaugh PJ, Javid B. Protein mistranslation:
friend   or   foe?   Trends   in   biochemical   sciences.   2014;39(8):355-62.   doi:
10.1016/j.tibs.2014.06.002. PubMed PMID: 25023410.
38.   Boeree MJ, Heinrich N, Aarnoutse R, Diacon AH, Dawson R, Rehal S, Kibiki GS,
Churchyard G, Sanne I, Ntinginya NE, Minja LT, Hunt RD, Charalambous S, Hanekom M, Semvua

HH, Mpagama SG, Manyama C, Mtafya B, Reither K, Wallis RS, Venter A, Narunsky K, Mekota A,
Henne S, Colbers A, van Balen GP, Gillespie SH, Phillips PPJ, Hoelscher M, Pan Ac. High-dose
rifampicin, moxifloxacin, and SQ109 for treating tuberculosis: a multi-arm, multi-stage
randomised controlled trial. Lancet Infect Dis. 2017;17(1):39-49. Epub 20161026. doi:
10.1016/S1473-3099(16)30274-2. PubMed PMID: 28100438; PMCID: PMC5159618.

39.   de Steenwinkel JE, Aarnoutse RE, de Knegt GJ, ten Kate MT, Teulen M, Verbrugh HA,
Boeree MJ, van Soolingen D, Bakker-Woudenberg IA. Optimization of the rifampin dosage to
improve the therapeutic efficacy in tuberculosis treatment using a murine model. Am J Respir
Crit Care Med. 2013;187(10):1127-34. doi: 10.1164/rccm.201207-1210OC. PubMed PMID:
23525933.

40.   Seijger C, Hoefsloot W, Bergsma-de Guchteneire I, Te Brake L, van Ingen J, Kuipers S, van
Crevel R, Aarnoutse R, Boeree M, Magis-Escurra C. High-dose rifampicin in tuberculosis:
Experiences from a Dutch tuberculosis centre. PLoS One. 2019;14(3):e0213718. Epub 20190314.
doi: 10.1371/journal.pone.0213718. PubMed PMID: 30870476; PMCID: PMC6417786.

     Table 1. Bioactivity of analogs of 7020 with superior potency.

Compound	Mistranslation Inhibition (%)	RSPR Inhibition (%)	Bacterial Growth Restriction in the Absence of Rifampicin (%)
7020	53.24	57.69	0
9719	83.21	61.23	0
9741	74.38	78.34	0
9744	86.95	87.32	1
9759	59.64	86.02	0
9767	64.77	81.43	1.2
9775	86.89	71.35	0
9782	83.73	86.78	0
9783	66.99	88.93	0
9787	75.51	98.14	1.15

Figure 1. A phenotypic whole-cell screen identified benzo[d]isoxazole-4,7-dione analogs as modulators of mycobacterial mistranslation (A) Schematic of the N-luc luciferase/GFP reporter system for screening of chemicals that decrease mistranslation in *Mycobacterium smegmatis*. (B) Screening of chemical library using the N-luc luciferase/GFP reporter for inhibitors of mycobacterial mistranslation. (Materials & Methods for details) The quality of the assay was assessed by the Z-factor, which was calculated by the formula $\text{Z-factor} = 1 - \frac{3\sigma_p + 3\sigma_n}{|\mu_n - \mu_p|}$, where σ_p, σ_n, μ_p and μ_n are standard deviations (σ) and means (μ) of the positive (p) and negative (n) controls respectively. Throughout the screening process, kasugamycin (Ksg) and dimethyl sulfoxide (DMSO) have been used as the positive and negative controls respectively.   (C) Structure of benzo[d]isoxazole-4,7-dione analogs identified in the preliminary screen (7006, 7013, 7020).

Figure 2. **The benzo[d]isoxazole-4,7-dione analogs 7006, 7013 and 7020 decrease mycobacterial mistranslation and RSPR.** (A) The compounds 7006, 7013 and 7020 could significantly decrease mistranslation in *M.smegmatis* expressing the N-luc/GFP reporter system. (B) *M. smegmatis* was plated on rifampicin containing media in presence 7006/ 7013/ 7020. All of them could significantly enhance rifampicin mediated mycobacterial killing. DMSO and Kasugamycin (Ksg) have been used as negative and positive controls in both the experiments. ***$p < 0.001$ by Student's t-test.

Figure 3. Structural analogs of 7020 which decrease mistranslation and RSPR in *M. smegmatis.* Several analogs of 7020 were synthesized, of which 10 compounds were found to be better than 7020 in inhibiting mycobacterial mistranslation and RSPR (Table 1). The structures of the analog are shown here along with the parent compound 7020.

Figure 4. **Analogs of 7020 lacking the benzoquione structure are poor inhibitors of mistranslation.** The structure of analogs of 7020 synthesized without the benzoquinone structure (boxed in green on 7020) and their percent inhibition of mistranslation are shown alongside 7020.

Figure 5. Replacement of 7020 methyl groups improves bioactivity of analog compounds. The structures and percent inhibition of mistranslation and RSPR inhibition of 7020 analogs with improved bioactivity that have replacements of the methyl groups at positions 5 and 6 (boxed in blue on 7020) are shown alongside 7020. In 9744, the methyl

groups have been replaced by chlorine. In 9775, 9782, 9783 and 9787, the methyl groups

have been replaced by an aromatic ring.

Figure 6. Compound 9787 decreases mistranslation more effectively than compound

7020 and decreases mistranslation in *M. bovis* BCG. (A) The N-luc/GFP reporter was

used to show the ability of 9787 to inhibit mycobacterial mistranslation. Different doses

of 9787 and the parent compound 7020 were used to check the improved potency of 9787.

Dimethyl sulfoxide (DMSO) and kasugamycin (Ksg) have been used as the negative and

the positive controls respectively. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$ by Student's t-test

(B) Other than *M. smegmatis*, the N-luc/GFP mistranslation reporter could be expressed

in *M.bovis* BCG strain. 9787 was able to decrease mistranslation in this strain as well.

$**p < 0.01$, $***p < 0.001$ by Student's t-test.

Figure 7. 30S ribosomal protein S5 (RpS5) is the molecular target of compound 9787.

(A) Structures of 9787, active biotinylated analog 9787, inactive analog 9789, and

inactive biotinylated analog 9791. (B) The N-luc/GFP reporter was used to show the

ability of 9787 and biotinylated 9788 to inhibit mycobacterial mistranslation. . $*p < 0.05$,

$**p < 0.01$, $***p < 0.001$ by Student's t-test (C) 9789, unable to reduce mycobacterial

mistranslation, was biotinylated to produce compound 9791. Neither of the compounds

could inhibit mycobacterial mistranslation. (D) Schematic of the experiment for target

identification of 9787. The compound of interest (C) is tagged with biotin (B) and

incubated with mycobacterial cell lysate. After it binds with the target (T), the whole

complex is pulled down using streptavidin (S) beads (Materials & Methods for details).

The pulled-down proteins are then analyzed by mass-spectrometry. (E) The ratio of the

quality scores of the proteins pulled down by the active biotinylated compound (9788)

and the inactive biotinylated compound (9791) has been plotted against the ratio of the

quality scores of proteins pulled down by the active alone (9788), and the mixture of

9788 and excess non-biotinylated active compound (9787). The 30S ribosomal protein S5

(Rps5) is represented by the red dot.

Supplementary Data 1. Pull-down mass spectrometry to identify the binding target

of 9787. Putative binding proteins identified by mass spectrometry after mycobacterial

cell lysates were incubated with (**A**) DMSO, (**B**) 9791, inactive biotinylated compound,

(**C**) 9791 + 9789, inactive biotinylated compound with excess inactive non-biotinylated

compound, (**D**) 9788, active biotinylated compound, and (**E**) 9788 + 9787, active

biotinylated compound with excess active non-biotinylated compound.

Fig. 1

A

D140N

N-luc

physiological
mistranslation

+ Compound

GFP

+ Compound

B

C

7006   7013   7020

Figure 1. A phenotypic whole-cell screen identified benzo[d]isoxazole-4,7-dione analogs as modulators of mycobacterial mistranslation (A) Schematic of the N-luc luciferase/GFP reporter system for screening of chemicals that decrease mistranslation in *Mycobacterium smegmatis*. (B) Screening of chemical library using the N-luc luciferase/GFP reporter for inhibitors of mycobacterial mistranslation. (Materials & Methods for details) The quality of the assay was assessed by the Z-factor, which was calculated by the formula $\text{Z-factor} = 1 - \frac{3\sigma_p + 3\sigma_n}{|\mu_n - \mu_p|}$, where σ_p, σ_n, μ_p and μ_n are standard deviations (σ) and means (μ) of the positive (p) and negative (n) controls respectively. Throughout the screening process, kasugamycin (Ksg) and dimethyl sulfoxide (DMSO) have been used as the positive and negative controls respectively. (C) Structure of benzo[d]isoxazole-4,7-dione analogs identified in the preliminary screen (7006, 7013, 7020).

Fig. 2

Figure 2. **The benzo[d]isoxazole-4,7-dione analogs 7006, 7013 and 7020 decrease mycobacterial mistranslation and RSPR.** (A) The compounds 7006, 7013 and 7020 could significantly decrease mistranslation in *M.smegmatis* expressing the N-luc/GFP reporter system. (B) *M. smegmatis* was plated on rifampicin containing media in presence 7006/ 7013/ 7020. All of them could significantly enhance rifampicin mediated mycobacterial killing. DMSO and Kasugamycin (Ksg) have been used as negative and positive controls in both the experiments. ***$p < 0.001$ by Student's t-test.

Fig. 3

Figure 3. Structural analogs of 7020 which decrease mistranslation and RSPR in *M. smegmatis.*
Several analogs of 7020 were synthesized, of which 10 compounds were found to be better than 7020
in inhibiting mycobacterial mistranslation and RSPR (Table 1). The structures of the analog are shown
here along with the parent compound 7020.

Fig. 4

Figure 4. **Analogs of 7020 lacking the benzoquione structure are poor inhibitors of mistranslation.** The structure of analogs of 7020 synthesized without the benzoquinone structure (boxed in green on 7020) and their percent inhibition of mistranslation are shown alongside 7020.

Fig. 5

7020	9744	9775
MI: 53.24%	MI: 86.95%	MI: 86.89%
RSPRI: 57.69%	RSPRI: 87.32%	RSPRI: 71.35%

9782	9783	9787
MI: 83.73%	MI: 66.99%	MI: 75.51%
RSPRI: 86.78%	RSPRI: 88.93%	RSPRI: 98.14%

Figure 5. Replacement of 7020 methyl groups improves bioactivity of analog compounds.
The structures and percent inhibition of mistranslation and RSPR inhibition of 7020 analogs
with improved bioactivity that have replacements of the methyl groups at positions 5 and 6
(boxed in blue on 7020) are shown alongside 7020. In 9744, the methyl groups have been
replaced by chlorine. In 9775, 9782, 9783 and 9787, the methyl groups have been replaced by
an aromatic ring.

Fig. 6

Figure 6. Compound 9787 decreases mistranslation more effectively than compound 7020 and decreases mistranslation in *M. bovis* BCG. (A) The N-luc/GFP reporter was used to show the ability of 9787 to inhibit mycobacterial mistranslation. Different doses of 9787 and the parent compound 7020 were used to check the improved potency of 9787. Dimethyl sulfoxide (DMSO) and kasugamycin (Ksg) have been used as the negative and the positive controls respectively. *p < 0.05, **p < 0.01, ***p < 0.001 by Student's t-test (B) Other than *M. smegmatis*, the N-luc/GFP mistranslation reporter could be expressed in *M.bovis* BCG strain. 9787 was able to decrease mistranslation in this strain as well. **p < 0.01, ***p < 0.001 by Student's t-test.

Fig. 7

E

Figure 7. 30S ribosomal protein S5 (RpS5) is the molecular target of compound 9787. (A) Structures of 9787, active biotinylated analog 9787, inactive analog 9789, and inactive biotinylated analog 9791. (B) The N-luc/GFP reporter was used to show the ability of 9787 and biotinylated 9788 to inhibit mycobacterial mistranslation. . *p < 0.05, **p < 0.01, ***p < 0.001 by Student's t-test (C) 9789, unable to reduce mycobacterial mistranslation, was biotinylated to produce compound 9791. Neither of the compounds could inhibit mycobacterial mistranslation. (D) Schematic of the experiment for target identification of 9787. The compound of interest (C) is tagged with biotin (B) and incubated with mycobacterial cell lysate. After it binds with the target (T), the whole complex is pulled down using streptavidin (S) beads (Materials & Methods for details). The pulled-down proteins are then analyzed by mass-spectrometry. (E) The ratio of the quality scores of the proteins pulled down by the active biotinylated compound (9788) and the inactive biotinylated compound (9791) has been plotted against the ratio of the quality scores of proteins pulled down by the active alone (9788), and the mixture of 9788 and excess non-biotinylated active compound (9787). The 30S ribosomal protein S5 (Rps5) is represented by the red dot.