Investigating the curvature sensing ability of Huntingtin's N17 domain

Neha Nanajkar,[†] Abhilash Sahoo,[‡,∥] Shelli L. Frey,[¶] and Silvina Matysiak[*,§]

[†]Department of Biology, University of Maryland, College Park

[‡]Biophysics Program, Institute of Physical Science and Technology, University of Maryland, College Park, MD, USA

[¶]Department of Chemistry, Gettysburg College, Gettysburg, Pennsylvania 17325, USA

[§]Fischell Department of Bioengineering, University of Maryland, College Park, MD, USA

[∥]Current address: Flatiron Institute, NY

E-mail: matysiak@umd.edu

Phone: +1(301) 405-0313

Abstract

N17, the N-terminal domain of the huntingtin protein (htt), has garnered significant attention for its role in htt's membrane binding and aggregation processes. Previous studies have identified a nuclear export sequence within the N17 domain and demonstrated its localization at various cellular organelles. Recent evidence suggests that, like other amphipathic helices, N17 can sense and preferentially bind to curved membranes. Gaining deeper insight into this behavior is essential to fully understand the function of this domain.

In this study, we combine coarse-grained molecular dynamics simulations with circular dichroism (CD) spectroscopy to investigate the mechanism behind N17's curvature sensing. We generated a unique hemispherical-planar membrane model, where 36% of

the upper leaflet surface is curved, allowing us to evaluate N17's binding preferences. Our findings show that N17 exhibits a strong preference for curved regions, with approximately 78 ± 7 % of peptides binding to these areas. This interaction is primarily mediated by the terminal F residues, indicating that N17's curvature sensing is driven by its ability to detect lipid packing defects. Furthermore, N17 not only senses these defects but also amplifies them by coalescing smaller pockets. Mutating the F residues to methionine, a smaller hydrophobic residue, significantly reduces N17's curvature sensitivity, resulting in equal binding to both curved and planar regions. CD spectroscopy corroborates these results, showing that N17 binds more strongly to highly curved small unilamellar vesicles (SUVs) compared to larger, less curved large unilamellar vesicles (LUVs).

Introduction

Huntingtin (htt) is an intracellular protein implicated in the development of Huntington's disease (HD). In HD, htt has an aberrant expansion of the polyglutamine region, resulting in the formation of toxic aggregates, subsequently leading to neurodegeneration.[1] The flanking domains of polyglutamine, N17 (MATLEKLMKAFESLKSF) and proline rich domains are known to promote and impede aggregation respectively.[2] N17, referring to the first 17 residues in the N-terminus, is particularly important with respect to membrane interaction. Composed of hydrophobic and polar residues, this fragment is disordered in solution but forms an amphipathic α-helix when in contact with membranes or in aggregated form.[3,4] N17's ability to self-associate results in higher rates of aggregation and fibril formation through an increase in local density of the polyglutamine stretch.[2,5] It also plays a pivotal role in cellular functioning. Studies indicate that N17 acts as the membrane anchoring domain of the first exon of htt, with reports of subcellular localization at the mitochondria, endoplasmic reticulum and the Golgi apparatus.[6–8] A leucine rich nuclear export motif present within the fragment is responsible for the localization of the htt protein.[5] Specific

mutations or complete deletion can result in nuclear accumulation of htt and a subsequent rise in cell toxicity. N17 is also the target of post-translational modifications that alter its aggregation and localization propensities.[9,10] Phosphomimetic mutations to S13/S16 of the N17 domain resulted in reduced cellular toxicity even at mutant htt fragments.[11] Targeted modifications of the N17 domain are being considered a possible avenue for the treatment of HD.[12] These and other works underscore the significance of the N17 domain in Huntington's Disease pathology.

Recent reports have suggested that N17 has a preference towards curved membranes. Studies using EPR spectroscopy demonstrate that the percentage of N17 binding to vesicles reduces as vesicle diameter increases.[13] In addition, curvature affinity of the synthetic N-terminal htt peptide (N17Q35P10KK) was observed through atomic force microscopy.[14] Silica nanoparticles were placed under a bilayer to create regions of artificial curvature of approximately 50 nm in diameter. Upon incubating the bilayer with N17Q35P10KK, the peptides accumulated on the curved rather than planar regions of the bilayer, indicative of preferential binding.

Curvature sensing has been observed in other peptides associated with neurodegeneration, such as α-synuclein and amyloid-β,[15,16] suggesting a relationship between neurodegeneration and membrane curvature. Prior work on curvature sensing has attributed sensing to electrostatic interactions or the hydrophobic effect. α-synuclein and ALPS, two commonly studied curvature sensors, were found to have differing modes of curvature sensing.[17] α-synuclein was found to have a poorly developed hydrophobic face, resulting in its sensitivity to anionic curvatures. In contrast, ALPS motifs have a very well developed hydrophobic face and few charged residues, leading to a hydrophobic effect driven sensing mechanism. It is unclear which of these categories N17 falls into, necessitating further examination.

Experimental and computational studies performed so far have unveiled a great deal about the interactions of N17 with lipids. All atom molecular dynamics simulations revealed characteristic features of N17 binding to zwitterionic lipid membranes,[18,19] including

tilt angle, insertion depth, and the effects of binding on the bilayer itself. Experimental studies using solid state NMR and circular dichroism examined the structural features of N17 in solution and while bound to vesicles and micelles.[20] From these studies, changes in membrane thickness, lipid order parameter and the structure of the N-terminal in varying environments were observed. There is no doubt these experimental works were instrumental in understanding N17's interaction with membranes. However, molecular level insights into the mechanisms responsible for curvature sensing are still missing. Traditional all atom MD is capable of modeling systems in high resolution but, it is computational challenging to capture large-scale biological phenomena such as membrane binding and peptide aggregation. Coarse-grained molecular dynamics (CG-MD) simulations, however, are uniquely poised to address these shortcomings. In CG-MD, similar atoms are grouped together to form beads. By "coarse-graining" the system, larger systems can be simulated, and for longer periods of time.

In this work, CG-MD is applied to simulate the binding of N17 to a membrane architecture containing both curved and planar regions, as depicted in Figure 1. As expected, we find that N17 does in fact sense curvature, with the bulk of binding occur at regions with lipid packing defects. The terminal F residue drives the interaction process, with peptide adsorption only occurring after partitioning of the two F residues (F11 and F17). Mutation of the F residues to M, a smaller hydrophobic amino acid, suppresses the curvature sensing ability. Circular dichroism experiments performed on N17 and its mutant are in agreement with these findings. Finally, we show that N17 seeks out regions with higher area per lipid, such that the bulky sidechains of F can partition to the membrane, at regions of lipid packing defects. Since mutated version of N17 has smaller sidechain residues it is capable of partitioning to regions without packing defects.

Figure 1: **Hemispherical-planar membrane architecture used in this study, consisting of both planar and curved regions.** The curved region of the upper leaflet accounts for 36% of the upper membrane surface area.

Results and Discussion

N17 prefers interacting with curved regions of the membrane

To assess whether N17 can sense curvature, we ran simulations of N17 with a hemispherical-planar membrane architecture, pictured in Fig. 1. The curved region of the membrane (M_c) accounts for around 36 % of the total surface area, while the planar region (M_p) makes up the remaining 64%. Thus, to characterize N17 as a curvature sensor, an excess of 36% of the peptides in the system must bind to the curved region. Fig. 2A presents a timeseries analysis of percentage of N17 binding in the system- either interacting with M_c, M_p. The percentage of peptides bound to M_c increases to around 85 % (blue) by the end of the simulation, far outweighing the 36% required to qualify N17 as a curvature sensor. Other replicates for N17 follow a similar trend (Figure S1), clearly positioning N17 as a curvature sensor. Note, at the studied timescales, we do not see complete binding of N17 to the membrane surface. Instead, N17 forms aggregates in solution that are, on average, 8-10 peptides in size (see Figure S2A).

Experimental verification of these findings was carried out by collecting circular dichro-

Figure 2: **N17 preferentially binds to curved regions of the membrane.** A) Binding percentages for one replicate of N17 to the curved and planar regions are denoted blue and purple respectively. The dotted lines represent the minimum percent binding required to qualify as having preference for curved and planar regions. The solid green line represents the percentage of membrane bound peptides in the system, B) Circular dichroism spectra for N17 in varying solutions. Circular dichroism of N17 in varying solutions. 10 mM pH 7.3 PBS (blue); 10 mM pH 7.3 PBS with 2 mM sodium dodecyl sulfate (SDS) to induce helical folding (orange); 3:1 POPC:POPS SUVs in 10 mM pH 7.3 PBS (green); and 3:1 POPC:POPS LUVs in 10 mM pH 7.3 PBS (red). SUVs (small unilamellar vesicles) are around 30nm in diameter and LUVs (large unilamellar vesicles) are around 120 nm in diameter, C) Area per lipid (APL) plot of the control simulation (no peptides), D) APL plot for one replicate of N17. Black dots indicate the center of mass of peptides that have bound to the membrane.

ism (CD) spectra on N17 in the presence of small and large unilamellar vesicles. Small Unilamellar Vesicles (SUVs) have a diameter of approximately 30 nm and the Large Unilamellar Vesicles (LUVs) have a diameter of approximately 120 nm. As vesicle size increases, its curvature reduces. Thus, a curvature sensing peptide should then prefer binding to an SUV over an LUV. To enhance the signal in CD binding experiments, the addition of anionic lipids is required, because the membrane partitioning constant of N17 with SUVs is 2 orders of magnitude larger in the 3:1 POPC:POPS when compared to 100% POPC.[19] Of note, NMR and atomistic simulations have shown N17 binding and helix formation in 100% POPC bilayers,[18–20] as observed in our simulations. We observed a tilt angle of 99.3 $\pm$ 7.9, in good agreement with prior works.[20] Figure 2B shows the spectra for N17. In PBS, N17 adopts a random coil structure, while in 2 mM SDS N17 is completely helical, with minimas at 208 and 222 nm. When bound to the vesicles, N17 adopts a combination of random coil and helical structures, evident through the minimas at 205 and 222 nm. However, in the presence of SUVs (green), N17 exhibits a significantly more intense spectra, indicating more alpha helical signal and more binding overall.

Figure 2 C, D present overhead views of area per lipid (APL) for the hemispherical-planar membrane architecture in control and with peptides simulations respectively. In the planar region L_d is predictably lower, as lipids tend to be more ordered and closely packed. There is a marked increase in APL at the curved region because of the formation of lipid packing defects. Figure 2D presents the APL at the last frame of one 500 ns replicate. Here, the black dots represent the center of mass of the peptides interacting with the membrane. For this replica, the majority of N17 binding indeed occurs at M_c, with only one peptide binding to M_p. The APL plots for the 2 other replicates reflect this trend, with most binding occurring at M_c (see Figure S3) Thus, we confirm that N17 is sensitive to and prefers binding to curved membranes.

Bulky hydrophobic residues are necessary for sensing curvature

We calculated the probability of first interaction across all residues of N17 to better understand how N17 interacts with the membrane. Fig. 3A reveals that about 30 % of the initial peptide-lipid interaction occurs through the terminal F residue, highlighting the importance of large hydrophobic groups in the curvature sensing process. In addition, there is a 20% chance the peptide interacts with the membrane via residues located on the N-terminus (M-L). N17 forms small aggregates in solution prior to interacting with the membrane, and the peptides orient their hydrophobic sides facing inward, forming helical bundles, resulting in residues at either termini being more likely to participate in the first interaction. However, F is significantly more hydrophobic than M, thus it is expected that the first interaction would occur primarily through the terminal F residue.

Figure 3: **N17-membrane interaction greatly depends on the F residues.** A) The bulk of N17-membrane interactions occur through the C-terminal F residue, B) Snapshots of N17 interacting with the upper leaflet of the membrane and C) N17 partitions to the upper leaflet upon insertion of the F residues. Protein backbones and F sidechains are denoted in red and orange respectively.

Figure S2A presents the average aggregate size for N17 interacting with M_c, M_p and remaining in solution. The average size of aggregates on M_p and M_c are 3 and 5 respectively. Figs. 3B and C present snapshots of the peptide interacting with the membrane. N17 first approaches the membrane, followed by interaction of the terminal F residue with the membrane surface. Only when both F11 and F17 residue insert to the surface does N17 partition completely to the membrane. In Fig. 3C, we note multiple peptides adsorbed to the curved region of the membrane, with deep insertion of the F residues (orange triangles).

Removal of F residues leads to a loss in curvature sensitivity

Insertions of hydrophobic residues into regions with lipid packing defects is one method by which amphipathic helices can sense, and in some cases generate curvature. In particular, the ALPS motif senses deep lipid packing defects through the insertion of its large hydrophobic residues.[21] Having established the role of the F residues in membrane interaction, we tested a mutated version of N17, henceforth called N17$^{F11M/F17M}$, where the bulky F residues in positions 11 and 17 were mutated to methionine. Methionine is also a hydrophobic amino acid but is smaller than phenylalanine, so the amphipathic nature of N17 is not disrupted. Furthermore, methionine is a residue commonly found in helical structures, thus the helical nature of N17 should not be perturbed.[22] This mutation allows us test the contribution of the bulky hydrophobic groups in the curvature sensing process.

Although Figure 3A indicates that F17 is the responsible for the bulk of first interactions with the membrane, we elected to mutate both F11 and F17 to methionine. This was done to eradicate the possibility of bundling between peptides. Prior work by our group has demonstrated that N17 domains of htt have a propensity to bundle together, arranging so the hydrophobic residues point inwards forming a hydrophobic core.[23] The large hydrophobic residues on N17 give way to two distinct and competing behaviors: either N17 bundles together, or it partitions to the membrane by inserting the F residues. Figure S4A presents a boxplot of the number of hydrophobic interactions within aggregates (in solution) of N17 and N17$^{F11M/F17M}$ respectively. Here, higher values indicate more interactions between the side chain beads of hydrophobic residues, ie. a larger hydrophobic core. Thus, it is clear the aggregates of N17 in solution experience more hydrophobic interactions. As a result of this, we see differences in the average aggregate size. Where N17 can form aggregates in the range of 8-10 peptides, aggregates of N17$^{F11M/F17M}$ are smaller at 4-6 peptides on average, as seen in Figure S2B. Although the formation of large oligomers of N17 is atypical in experimental studies, simulations usually require concentrations higher than physiological concentrations to ensure sufficient sampling and to guarantee concentration dependent

Figure 4: **A mutated version of N17, N17$^{F11M/F17M}$ has no specific binding preferences, binding to both the curved and planar regions.** A) Binding percentages for one replicate of N17$^{F11M/F17M}$ to the curved and planar regions are denoted blue and purple respectively. The dotted lines represent the minimum percent binding required to qualify as having preference for curved and planar regions. The solid green line represents the percentage of membrane bound peptides in the system, B) Circular dichroism of N17$^{F11M/F17M}$ in varying solutions. 10 mM pH 7.3 PBS (blue); 10 mM pH 7.3 PBS with 2 mM sodium dodecyl sulfate (SDS) to induce helical folding (orange); 3:1 POPC:POPS SUVs in 10 mM pH 7.3 PBS (green); and 3:1 POPC:POPS LUVs in 10 mM pH 7.3 PBS (red). SUVs (small unilamellar vesicles) are around 30nm in diameter and LUVs (large unilamellar vesicles) are around 120 nm in diameter, C) First interaction probability for residues of N17$^{F11M/F17M}$ with the lipid headgroups, D) APL plot for one replicate of N17$^{F11M/F17M}$. Black dots indicate the center of mass of bound peptides.

phenomena, like aggregation, can be effectively studied. The concentration of N17 in CD experiments amounts to 76 μM, while the peptide concentration in simulation is closer to 10mM. At these elevated concentrations, N17 aggregation is not an unexpected phenomena. Figure S4B presents a binary classification of the competing interactions. Most peptides that are involved in interpeptide hydrophobic interactions do not interact with the membrane. Thus, by mutating both F to M, we eradicate the possibility of bundling and achieve comparable binding fractions for both the wild-type N17 or its mutated counterpart (Figure S4C).

Fig. 4A presents the binding percentage plot of one replicate of N17$^{F11M/F17M}$, where a loss in curvature specificity can be noted for the mutated peptide. By the end of the simulation, we see that the majority of peptide is bound to the planar region, though the percentage fluctuates around 64 %. The circular dichroism spectra of N17$^{F11M/F17M}$, shown in Fig. 4B, is markedly different than that of wild-type N17. N17 demonstrates a clear preference towards SUVs (more curvature) over LUVs, demarcated by the significant increase in spectral intensity (Fig. 2B). On the other hand, N17$^{F11M/F17M}$ only indicates a slight preference for SUVs over LUVs. This indicates that removal of the bulky hydrophobic amino acids from the amphipathic N17 segment, greatly affects its binding preferences and its tendency to localize to membranes of curvature.

Similar to N17, N17$^{F11M/F17M}$ also displays a high interaction probability at either termini, as seen in Fig. 4C. Since the mutation from F to M does not affect the amphipathic nature of N17$^{F11M/F17M}$, the mutated peptides also orient with their hydrophobic face pointing inwards, resulting in first interactions that can occur from either the N or C terminal residues. Fig. 4D presents the APL for one replicate of the N17$^{F11M/F17M}$ system. Here, N17$^{F11M/F17M}$ binds to both M_c and M_p, displaying no preference for either region. Replicates of these simulations, shown in Figure S5, substantiate these trends.

Mechanisms of curvature sensing

To better understand the mechanism of curvature sensing, we analyzed the effect of N17 binding on the membrane. Figure 5A illustrates the distribution of area per lipid (APL) at M_c for both the control simulation and the simulation with peptides. In the control, the APL at M_c peaks around 1.5 nm^2, with a median value of 0.8 nm^2. In a planar POPC bilayer, our lipid model has APL values of 0.62 nm^2,[24] falling within the range of accepted values for POPC.[25,26] The extension to 1.5 nm^2 highlights the existence of packing defects, formed due to the curved region. However, these defects are enhanced in the presence of N17, where the APL distribution is vastly extended, ranging from with values reaching up to 4 nm^2, in sharp contrast to the condensed APL distribution of the control simulation . Figures 2 C,D tell the same story, with the curved region having higher APL values than the surrounding planar regions. In the control simulation, the curved region has APL values ranging from 0.625 nm^2 to 1.562nm^2, but upon binding of the peptide to the curved region, a jump in APL can be noted, with some regions having an APL of 2.5 nm^2. Thus, binding of N17 further enhances lipid packing defects, corresponding to the extended tail seen in Figure 5A. The inset in Figure 5A presents a schematic of the process. Peptides may partition to different regions of the membrane surface either as a monomer, or as a small aggregate (such as a dimer/trimer/tetramer). Over time, the peptides bound to M_c are able to diffuse towards each other, causing the increase in defects in the curved region of the membrane.

We have confirmed that N17 is sensitive to membrane curvature through its affinity for lipid packing defects, and lack of the bulky hydrophobic groups in N17$^{F11M/F17M}$ results in loss of curvature specificity. To distinguish the binding of N17 and N17$^{F11M/F17M}$, we considered APL at the site of initial peptide binding, as seen in Fig. 5B. Due to its sensitivity for curved regions, N17 displays distinct partitioning preferences when interacting with M_c and M_p respectively. It is entropically more favorable for the large F residues of N17 to find regions of higher APL (and thus more space between headgroups) to bind. These regions primarily occur in the curved region of the membrane. In the case of N17$^{F11M/F17M}$,

Figure 5: **N17 detects regions of curvature by sensing lipid packing defects.** A) Violin plots plot of area per lipid (APL) of the curved region of the membrane in the control simulation (no peptide) and simulation with N17. The inset contains a schematic describing the mechanism by which N17 can enhance membrane defects, ie. N17 binds to regions with packing defects and these defects consolidate into larger defects leading to the extended range of APL values, B) Area per lipid (APL) at the site of first peptide interaction for both N17 and N17$^{F11M/F17M}$, C) and D) present diffusion maps for N17 and N17$^{F11M/F17M}$ respectively. Positions denote the center of mass of the peptide and each peptide is marked in a different color.

however, the distribution of APL at the initial peptide binding event are virtually identical. The smaller size of methionine allows N17 to bind to regions where the defects are not nearly as large, as evidenced by the indiscriminate binding to both planar and curved regions. An interesting feature observed in the simulations of N17$^{F11M/F17M}$, is the diffusion of the peptide across the bilayer, something not observed in the simulations with N17. In one replicate, we observed N17$^{F11M/F17M}$ initially binding to the planar region of the membrane, then diffuse to the curved region such that the binding percentage at the curved region is higher (see Figure S6B).

Figures 5C and D present diffusion maps for one replicate of N17 and N17$^{F11M/F17M}$, respectively. Here, the center of mass of peptides in direct contact with the membrane is tracked over time, and each peptide is marked in a different color. N17 and N17$^{F11M/F17M}$ demonstrate vastly different behaviour while interacting with the lipid. N17 binds to the curved region of the bilayer (center of simulation box), and displays very little diffusion. On the other hand, N17$^{F11M/F17M}$, diffuses over the entire membrane surface. We can connect back to Figure 5B, where it is more entropically favorable for N17 to bind regions of high APL to accommodate its bulky hydrophobic groups, which then restrict diffusion along the membrane surface. The smaller sidechains of N17$^{F11M/F17M}$ allow it to diffuse along the membrane, even from the curved region to the planar.

Conclusions

In this work, we apply coarse-grain molecular dynamics simulations to investigate the interactions of N17, the N-terminal domain of htt protein, with curved membranes. A plethora of studies have detailed the importance of N17, not just in Huntington's Disease pathology, but also in healthy cellular function. As the membrane binding domain of htt, it is crucial to understand how N17 interacts with the complex cellular environment, specifically, how N17 associates with curved membranes, a common intracellular feature.

The curvature sensing ability of N17 has been noted in prior experiments via AFM microscopy and EPR spectroscopy.[13,14] We aimed to uncover a molecular mechanism that can explain this behavior. To do so, we use a membrane architecture containing both planar and curved regions, as depicted in Figure 1. The curved region(M_c) accounts for 36% of the upper leaflet surface. Thus if more than 36 % of the peptides are bound to M_c, the peptide was classified as a curvature sensor.

Nearly 79 % N17-M_c binding was observed, firmly establishing N17 as a curvature sensor. We found that membrane binding is largely dependent on the C-terminal F residue, with over 30% of interactions occurring through this residue. N17 domains arrange as bundles in solution due to their amphipathic nature,[23] thus interactions can occur through either termini. Since the C-terminus is more hydrophobic, we observe most interaction from this terminus. Mutation of the F residues to M, in $N17^{F11M/F17M}$, revealed the pivotal role that bulky hydrophobic residues play in curvature sensing. $N17^{F11M/F17M}$ displays no binding preferences, interacting with both M_c and M_p.

These findings were found to be in agreement with circular dichroism spectra. N17 could differentiate between SUVs (more curvature) and LUVs (less curvature), as seen by the variance in signal intensity in Figure 2B. By contrast, $N17^{F11M/F17M}$ does not distinguish between LUVs and SUVs to the same extent. In the case of N17, the bulky hydrophobic groups gravitate to regions with higher APL, where it is entropically more favorable to bind. This occurs primarily in the curved region of the membrane, where lipid packing defects are abundant. $N17^{F11M/F17M}$ does not discriminate between curved and planar to the same extent, as shown by the similar APL distributions at the sight of first lipid interaction. Here, the smaller M residues are able bind to regions of the membrane, even if the area per lipid is low.

Two main strategies have been proposed to explain the curvature sensing phenomena.[27,28] Certain peptides, such as the BAR domain proteins, possess a slightly curved surface populated with charged residues, resulting in electrostatic attraction to curved membranes. On

the other hand, ALPS motifs bind preferentially to curvature through sensing the lipid packing defects in curved membranes. They often possess large hydrophobic residues on one face and small polar groups (serine and threonine) on the polar face, leading to membrane curvature being a necessity for binding. α-synuclein is one such curvature sensor, with a poorly developed hydrophobic face, but a comparatively stronger polar face, granting it sensitivity to curved anionic lipid bilayers. Interestingly, mutating key polar residues to W, results in α-synuclein developing a sensitivity to zwitterionic lipid curvatures as well.[17] Unlike α-synuclein and ALPS motifs, N17 has a well developed hydrophobic face, with 2 aromatic F residues and a sizable polar face of 5 charged residues. Thus, the mechanism by which N17 senses curvature is distinct. While N17 is able to detect membrane curvature, it can also bind to planar membranes, placing it in a unique category of curvature sensor We suspect these distinctions may have larger effects in downstream cellular signaling.

The insights obtained from this study will be crucial to better develop our understanding of neuro-degenerative diseases, with Huntington's Disease as a model system. A plethora of work has highlighted the link between curvature sensing and curvature generation, with some peptides sensing curvature at low concentrations and generating curvature at high concentrations.[29,30] Although our simulations employ high concentrations, we cannot assess the potential curvature generation due to the restraining potentials applied to retain the hemispherical-planar architecture. Our future works will feature updated lipid systems to examine possible membrane curvature generation upon N17 binding.

Methods

Simulations were performed on GROMACS 2019.4 simulation engine.[31] VMD 1.9.4[32] was used for visualization. Analysis was performed on MDAnalysis 1.1.1[33] using in-house scripts.

Simulation parameters

Forcefield. In this work, we use the previously developed ProMPT forcefield[34] to set up all simulations. ProMPT is a coarse-grained protein force-field that employs free-energy based parameterization and a 4:1 bead mapping. To account for the orientation of the atoms underlying the coarse grained beads, flexible dipoles consisting of opposing charges are incorporated into the coarse-grained beads. The electrostatic interactions induced by these dipoles introduce structural polarization to the model, allowing us to capture secondary structure changes in response to fluctuations in the local environment. ProMPT has been validated on a number of small proteins[35] and more recently was employed to study aggregation of N17 and the following polyQ domain.[23] Figure S8 presents the ProMPT representation of the protein. A helical dihedral potential with competing minima was applied to the N17 backbone beads, utilizing a force constant of 10 kcal/mol (see Figure S9). The lipid model, titled WEPMEM, is similarly designed, in that it features a 4:1 bead mapping with the addition of explicit dipole to polar coarse grained beads.[24] The protein and lipid forcefields are used in conjunction with the polarizable Martini water.[36]

Curved membrane setup. Experimentally, curvature is typically studied either using vesicles or supported bilayers. To reproduce curvature in silico, membrane architecture containing both curved and planar regions was created using the BUMPy package.[37] The hemispherical-planar membrane architecture pictured in Figure 1 features POPC, a phospholipid routinely encountered in the cellular environment and has a diameter of approximately 15 nm. To retain the shape of the membrane through the course of the simulation, a grid of dummy particles was restrained close to the phosphate group of the lipid, as detailed in the BUMPy paper.[37] These dummy particles have a repulsive interaction with the acyl tail of the lipids (C_{12}= 0.02581 kJ mol^{-1} nm^2), ensuring that the membrane shape is maintained through the course of the simulation. The dummy particles do not engage in interactions with any other molecules in the system. To promote binding of N17 to the positive curvature (upper leaflet), we instituted a repulsive interaction (C_{12}= 0.02581 kJ mol^{-1} nm^2) between

the headgroup of the lower leaflet with the protein backbone bead.

To stabilize the curved setup, an energy minimization of 5000 steos was performed, followed by an NVT equilibration for 10 nanoseconds with soft restraint on phosphate group. Next, an NPT simulation was run for 50 nanoseconds without any restraints on the phosphate. The final diameter of the membrane architecture is around 15 nm.

Simulation Setup. The control simulation (no peptide) was run for 250 ns using the NPT ensemble. For the simulations with N17 and $N17^{F11M/F17M}$, peptides were run added to the water box for a peptide concentration of approximately 20mM, amounting to 36 peptides and approximately 1400 POPC molecules, or peptide:lipid ratio of 1:40. The simulation box had x,y,z dimensions of 26nm, 26nm and 13 nm respectively. Following this, a short energy minimization of 5000 steps and equilibration of 5000 steps was performed. Counterions and a salt concentration amounting to 125mM were then added to the simulation box. 3 consecutive equilibrations of 5000 steps were performed with timesteps of 1fs, 5fs and 10fs respectively. Production was run for 500 ns with the NPT ensemble.

The Nose-Hoover thermostat, with a time constant of 1 picosecond, was utilized to maintain a fixed temperature of 300K and the Parrinello-Rahman barostat kept the pressure constant at 1 bar. A relative dielectric constant of 2.5 was applied, and electrostatic calculations were performed using Particle Mesh Ewald. Constraints were evaluated using the LINCS algorithm, and the neighbor list was updated every 10 steps.

Peptide preparation

N17, the first 17 amino acids of htt (MATLEKLMKAFESLKSF), and $N17^{F11M/F17M}$ (N17 with an F11M/F17M substitution), both with C-terminal amidation, were obtained via custom synthesis (GenScript, Piscataway, New Jersey). Disaggregation and solubilization of N17 and $N17^{F11M/F17M}$ peptides was achieved using established protocols.[38] Briefly, peptide was dissolved overnight at a concentration of 0.5 mg/mL in a 1:1 mixture of trifluoroacetic acid (TFA, Sigma) and hexafluoroisopropanol (HFIP, Sigma). After vortexing, a N_2 stream

was used to evaporate solvent. To remove residual solvent, samples were placed in a vacufuge concentrator overnight, resulting in thin peptide films which were then stored at $-20°C$. For each experiment, peptide films were resuspended in ultrapure water that had been adjusted to pH 3 with TFA and then diluted in phosphate-buffered saline (PBS) (10 mM phosphate, 138 mM NaCl, 2.7 mM KCl, pH 7.3) (Dulbecco) to a suitable final concentration, typically 100-200 µM.

Vesicle preparation

Large unilamellar vesicles (LUVs) and small unilamellar vesicles (SUVs) composed of 3:1 1-palmitoyl-2-oleoyl-glycero-3-phosphocholine (POPC): 1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine (POPS) (Avanti Polar Lipids, Alabaster, AL) were prepared using extrusion and sonication methods. Briefly, a thin lipid film of lipid, dried under nitrogen from a chloroform stock, was hydrated at 2.5 mg/ml in PBS (10 mM phosphate, 138 mM NaCl, 2.7 mM KCl, pH 7.3) and vortexed for 10 min. For LUVs, this solution was treated with 10 freeze-thaw cycles using ethanol/dry ice and warm water baths and then extruded 11 times through a 100 nm pore polycarbonate filter (Avanti Polar Lipids). For SUVs, this solution was bath sonicated for 30-60 min. Dynamic light scattering measurements with a Zetasizer ZS90(Malvern, Worcestershire, UK) were used to measure vesicle diameter (LUVs 130 nm; SUVs 30 nm) and ensure sample homogeneity.

Circular dichroism

CD spectra were collected on a JASCO J-1500 spectropolarimeter using a 1 mm pathlength quartzcuvette. Samples contained 0.15 mg/mL peptide (76 µM) in PBS at pH 7.3 and spectra were normalized to protein concentration measured by UV-visible spectroscopy at 205 nm (ϵ_{N17}= 65340 M^{-1} cm^{-1}; $\epsilon_{N17F11M/F17M}$ = 58570 M^{-1} cm^{-1}).[39] Lipid interfaces were introduced via 1.25 mg/mL lipid (1.60 mM) SUVs or LUVs. Spectra are baseline corrected and averaged over three scans, measured from 260 nm to 190 nm with a 0.2 nm data interval,

and 4 s data averaging.

Analysis of simulations

To conduct the analysis of the simulations, we first established specific criteria. A distance cutoff of 7 Å was used to evaluate contacts across all analyses. Peptides were considered interacting or bound to another peptide, or adsorbed to the membrane, if they had five or more heavy atom contacts with the interacting partner (either another peptide or the lipid phosphate group). We further categorized peptide interactions with the lipid as either direct or indirect. Direct interactions were defined as a peptide having five or more heavy atom contacts with the phosphate group of the lipid. Indirect interactions were defined as instances where peptide X did not directly interact with the lipid bilayer but instead interacted with another peptide Y, which was in contact with the bilayer. In this scenario, peptide X is considered to be bound to the lipid indirectly through its interaction with peptide Y.

Analysis performed on the proteins considered all heavy atoms (backbone and sidechain beads) unless explicitly mentioned. For the lipids, only the phosphate bead was considered.

To calculate the percentage of peptides bound to the curved and planar regions shown in Figure 2A, we counted the number of peptides interacting with M_c and M_p using the above mentioned criteria, and normalised by the total number of bound peptides in the system at each time point.

Area per lipid plots shown in Figures 2 and 4 were calculated from the FATSLiM package,[40] which provides the APL for each lipid in the system. Here, the black dots depict the center of mass of peptides bound to the curved region.

To classify the initial interaction between the peptide and lipid (Figures 3A and 4C), we defined the first interaction as lasting for at least 5 ns. This approach helped eliminate any transient interactions. The probabilities shown in the first interaction plots are averaged across all replicates, representing the frequency of peptide-lipid interaction per residue.

Figure 5 was created using the violinplot function from the seaborn package. Diffusion

plots were created by plotting the center of mass of the peptides in direct contact with the membrane. Here, peptides are differentiated by color.

Acknowledgement

This research was supported by the National Science Foundation under grants CHE-1454948 and CHE-2202281. The authors acknowledge the University of Maryland supercomputing resources (http://hpcc.umd.edu) made available for conducting the research reported in this paper.

Supporting Information Available

Analysis performed on replicates of N17 and N17$^{F11M/F17M}$ including, binding percentages, area per lipid, and diffusion plots. Details on peptide model also included.

References

(1) Schulte, J.; Littleton, J. T. The biological function of the Huntingtin protein and its relevance to Huntington's Disease pathology. *Current trends in neurology* **2011**, *5*, 65.

(2) Crick, S. L.; Ruff, K. M.; Garai, K.; Frieden, C.; Pappu, R. V. Unmasking the roles of N-and C-terminal flanking sequences from exon 1 of huntingtin as modulators of polyglutamine aggregation. *Proceedings of the National Academy of Sciences* **2013**, *110*, 20075–20080.

(3) Thakur, A. K.; Jayaraman, M.; Mishra, R.; Thakur, M.; Chellgren, V. M.; L Byeon, I.-J.; Anjum, D. H.; Kodali, R.; Creamer, T. P.; Conway, J. F., et al. Polyglutamine disruption of the huntingtin exon 1 N terminus triggers a complex aggregation mechanism. *Nature structural & molecular biology* **2009**, *16*, 380–389.

(4) Jayaraman, M.; Kodali, R.; Sahoo, B.; Thakur, A. K.; Mayasundari, A.; Mishra, R.; Peterson, C. B.; Wetzel, R. Slow amyloid nucleation via α-helix-rich oligomeric intermediates in short polyglutamine-containing huntingtin fragments. *Journal of molecular biology* **2012**, *415*, 881–899.

(5) Zheng, Z.; Li, A.; Holmes, B. B.; Marasa, J. C.; Diamond, M. I. An N-terminal nuclear export signal regulates trafficking and aggregation of Huntingtin (Htt) protein exon 1. *Journal of Biological Chemistry* **2013**, *288*, 6063–6071.

(6) Orr, A. L.; Li, S.; Wang, C.-E.; Li, H.; Wang, J.; Rong, J.; Xu, X.; Mastrobernardino, P. G.; Greenamyre, J. T.; Li, X.-J. N-terminal mutant huntingtin associates with mitochondria and impairs mitochondrial trafficking. *Journal of Neuroscience* **2008**, *28*, 2783–2792.

(7) Rockabrand, E.; Slepko, N.; Pantalone, A.; Nukala, V. N.; Kazantsev, A.; Marsh, J. L.; Sullivan, P. G.; Steffan, J. S.; Sensi, S. L.; Thompson, L. M. The first 17 amino acids of Huntingtin modulate its sub-cellular localization, aggregation and effects on calcium homeostasis. *Human molecular genetics* **2007**, *16*, 61–77.

(8) Atwal, R. S.; Xia, J.; Pinchev, D.; Taylor, J.; Epand, R. M.; Truant, R. Huntingtin has a membrane association signal that can modulate huntingtin aggregation, nuclear entry and toxicity. *Human molecular genetics* **2007**, *16*, 2600–2615.

(9) DiGiovanni, L. F.; Mocle, A. J.; Xia, J.; Truant, R. Huntingtin N17 domain is a reactive oxygen species sensor regulating huntingtin phosphorylation and localization. *Human molecular genetics* **2016**, *25*, 3937–3945.

(10) Yalinca, H.; Gehin, C. J. C.; Oleinikovas, V.; Lashuel, H. A.; Gervasio, F. L.; Pastore, A. The role of post-translational modifications on the energy landscape of Huntingtin N-terminus. *Frontiers in molecular biosciences* **2019**, *6*, 95.

(11) Thompson, L. M.; Aiken, C. T.; Kaltenbach, L. S.; Agrawal, N.; Illes, K.; Khoshnan, A.; Martinez-Vincente, M.; Arrasate, M.; O'Rourke, J. G.; Khashwji, H., et al. IKK phosphorylates Huntingtin and targets it for degradation by the proteasome and lysosome. *Journal of Cell Biology* **2009**, *187*, 1083–1099.

(12) Kumar, V.; Singh, A. Targeting N17 domain as a potential therapeutic target for the treatment of Huntington disease: An opinion. *EXCLI journal* **2021**, *20*, 1086.

(13) Tao, M.; Pandey, N. K.; Barnes, R.; Han, S.; Langen, R. Structure of membrane-bound huntingtin exon 1 reveals membrane interaction and aggregation mechanisms. *Structure* **2019**, *27*, 1570–1580.

(14) Chaibva, M.; Burke, K. A.; Legleiter, J. Curvature enhances binding and aggregation of huntingtin at lipid membranes. *Biochemistry* **2014**, *53*, 2355–2365.

(15) Jensen, M. B.; Bhatia, V. K.; Jao, C. C.; Rasmussen, J. E.; Pedersen, S. L.; Jensen, K. J.; Langen, R.; Stamou, D. Membrane curvature sensing by amphipathic helices: a single liposome study using α-synuclein and annexin B12. *Journal of Biological Chemistry* **2011**, *286*, 42603–42614.

(16) Kegulian, N. C.; Sankhagowit, S.; Apostolidou, M.; Jayasinghe, S. A.; Malmstadt, N.; Butler, P. C.; Langen, R. Membrane Curvature-sensing and Curvature-inducing Activity of Islet Amyloid Polypeptide and Its Implications for Membrane Disruption*. *Journal of Biological Chemistry* **2015**, *290*, 25782–25793.

(17) Pranke, I. M.; Morello, V.; Bigay, J.; Gibson, K.; Verbavatz, J.-M.; Antonny, B.; Jackson, C. L. α-Synuclein and ALPS motifs are membrane curvature sensors whose contrasting chemistry mediates selective vesicle binding. *Journal of Cell Biology* **2011**, *194*, 89–103.

(18) Côté, S.; Binette, V.; Salnikov, E. S.; Bechinger, B.; Mousseau, N. Probing the Hunt-

ingtin 1-17 membrane anchor on a phospholipid bilayer by using all-atom simulations. *Biophysical Journal* **2015**, *108*, 1187–1198.

(19) Michalek, M.; Salnikov, E. S.; Werten, S.; Bechinger, B. Membrane interactions of the amphipathic amino terminus of huntingtin. *Biochemistry* **2013**, *52*, 847–858.

(20) Michalek, M.; Salnikov, E. S.; Bechinger, B. Structure and topology of the huntingtin 1–17 membrane anchor by a combined solution and solid-state NMR approach. *Biophysical journal* **2013**, *105*, 699–710.

(21) Mouret, L.; Vamparys, L.; von Filseck, J. M.; Fuchs, P.; Bondon, A.; Drin, G. Structural basis of membrane curvature recognition by the ALPS motifs. *Biophysical Journal* **2015**, *108*, 2a.

(22) Okamoto, Y. Helix-forming tendencies of nonpolar amino acids predicted by Monte Carlo simulated annealing. *Proteins: Structure, Function, and Bioinformatics* **1994**, *19*, 14–23.

(23) Nanajkar, N.; Sahoo, A.; Matysiak, S. Unraveling the Molecular Complexity of N-Terminus Huntingtin Oligomers: Insights into Polymorphic Structures. *The Journal of Physical Chemistry B* **2024**,

(24) Ganesan, S. J.; Xu, H.; Matysiak, S. Effect of lipid head group interactions on membrane properties and membrane-induced cationic β-hairpin folding. *Physical Chemistry Chemical Physics* **2016**, *18*, 17836–17850.

(25) Leekumjorn, S.; Sum, A. K. Molecular characterization of gel and liquid-crystalline structures of fully hydrated POPC and POPE bilayers. *The Journal of Physical Chemistry B* **2007**, *111*, 6026–6033.

(26) Janosi, L.; Gorfe, A. A. Simulating POPC and POPC/POPG bilayers: conserved pack-

ing and altered surface reactivity. *Journal of chemical theory and computation* **2010**, *6*, 3267–3273.

(27) Antonny, B. Mechanisms of membrane curvature sensing. *Annual review of biochemistry* **2011**, *80*, 101–123.

(28) Drin, G.; Casella, J.-F.; Gautier, R.; Boehmer, T.; Schwartz, T. U.; Antonny, B. A general amphipathic α-helical motif for sensing membrane curvature. *Nature structural & molecular biology* **2007**, *14*, 138–146.

(29) Suetsugu, S.; Gautreau, A. Synergistic BAR–NPF interactions in actin-driven membrane remodeling. *Trends in cell biology* **2012**, *22*, 141–150.

(30) Simunovic, M.; Bassereau, P.; Voth, G. A. Organizing membrane-curving proteins: the emerging dynamical picture. *Current opinion in structural biology* **2018**, *51*, 99–105.

(31) Abraham, M. J.; Murtola, T.; Schulz, R.; Páll, S.; Smith, J. C.; Hess, B.; Lindahl, E. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* **2015**, *1*, 19–25.

(32) Humphrey, W.; Dalke, A.; Schulten, K. VMD: visual molecular dynamics. *Journal of molecular graphics* **1996**, *14*, 33–38.

(33) Gowers, R. J.; Linke, M.; Barnoud, J.; Reddy, T. J.; Melo, M. N.; Seyler, S. L.; Domanski, J.; Dotson, D. L.; Buchoux, S.; Kenney, I. M., et al. MDAnalysis: a Python package for the rapid analysis of molecular dynamics simulations. Proceedings of the 15th python in science conference. 2016; p 105.

(34) Sahoo, A.; Lee, P.-Y.; Matysiak, S. Transferable and Polarizable Coarse Grained Model for Proteins- ProMPT. *Journal of Chemical Theory and Computation* **2022**, *18*, 5046–5055.

(35) Lee, P.-Y.; Sahoo, A.; Matysiak, S. Folding and modulation of the helical conformation of Glycophorin A by point mutations. *Physical Chemistry Chemical Physics* **2023**, *25*, 10885–10893.

(36) Yesylevskyy, S. O.; Schäfer, L. V.; Sengupta, D.; Marrink, S. J. Polarizable water model for the coarse-grained MARTINI force field. *PLoS computational biology* **2010**, *6*, e1000810.

(37) Boyd, K. J.; May, E. R. BUMPy: a model-independent tool for constructing lipid bilayers of varying curvature and composition. *Journal of chemical theory and computation* **2018**, *14*, 6642–6652.

(38) Chen, S.; Berthelier, V.; Yang, W.; Wetzel, R. Polyglutamine aggregation behavior in vitro supports a recruitment mechanism of cytotoxicity. *Journal of molecular biology* **2001**, *311*, 173–182.

(39) Anthis, N. J.; Clore, G. M. Sequence-specific determination of protein and peptide concentrations by absorbance at 205 nm. *Protein Science* **2013**, *22*, 851–858.

(40) Buchoux, S. FATSLiM: a fast and robust software to analyze MD simulations of membranes. *Bioinformatics* **2017**, *33*, 133–134.

TOC Graphic

Some journals require a graphical entry for the Table of Contents. This should be laid out "print ready" so that the sizing of the text is correct.

Inside the `tocentry` environment, the font used is Helvetica 8 pt, as required by *Journal of the American Chemical Society*.

The surrounding frame is 9 cm by 3.5 cm, which is the maximum permitted for *Journal of the American Chemical Society* graphical table of content entries. The box will not resize if the content is too big: instead it will overflow the edge of the box.

This box and the associated title will always be printed on a separate page at the end of the document.