Natural variation in chalcone isomerase defines a major locus controlling growth variation between *Populus nigra* populations

Duruflé Harold*[1], Déjardin Annabelle[1], Jorge Véronique, Pégard Marie[2], Pilate Gilles[1], Rogier Odile1, Sanchez Leopoldo[1], Segura Vincent*[1,3]

[1] INRAE, ONF, BioForA, UMR 0588, 45075 Orléans, France
[2] INRAE, URP3F, UR 0004, 86600 Lusignan, France
[2] UMR AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, F-34398 Montpellier, France

*Corresponding authors
Correspondence: harold.durufle@inrae.fr; vincent.segura@inrae.fr

ABSTRACT

Poplar is a promising resource for wood production and the development of lignocellulosic biomass, but currently available varieties have not been optimized for these purposes. Therefore, it is critical to investigate the genetic variability and mechanisms underlying traits that affect biomass yield. Previous studies have shown that target traits in different poplar species are complex, with a small number of genetic factors having relatively low effects compared to medium to high heritability. In this study, a systems biology approach was implemented, combining genomic, transcriptomic, and phenotypic information from a large collection of individuals from natural populations of black poplar from Western Europe. Such an approach identified a QTL and a gene, chalcone isomerase (CHI), as a candidate for controlling radial growth. Additionally, analysis of the structure and diversity of traits as well as CHI gene expression revealed a high allelic fixation index, linked to the geographical origin of the natural populations under study. These findings provide insights into how adaptive traits arise, are selected, and maintained in the populations. Overall, this study contributes to enhancing the use of poplar as a valuable resource for sustainable biomass production.

Keywords: Poplar, GWAS, transcriptomics, systems biology, adaptation

Introduction

Trees play a crucial role in mitigating climate change by sequestering carbon from the atmosphere through photosynthesis, and forest ecosystems are considered the largest terrestrial carbon sinks on Earth (Pan et al., 2011; Harris et al., 2021). The future evolution of carbon sequestration in forests relies heavily on how the growth rate and lifespan of trees respond to the changing climate (Brienen et al., 2020; Zhou, 2022). Trees keep accumulating carbon in their trunks, branches, and roots as they grow, which enables them to capture and store atmospheric carbon for several decades or possibly centuries (Green & Keenan, 2022). The major part of the tree trunk is created by the cambium, and the developing xylem constitutes a complex and dynamic system that generates wood in accordance with the seasonal cycle (Rathgeber et al., 2016). However, we still lack an integrative theory to understand growth patterns because wood formation requires the coordination of many metabolic pathways (Bryant et al., 2023).

Knowing and understanding the links between phenotypes and genetic mutations is a major challenge. Such studies have emerged for poplar, a model for tree biology, genomics, evolutionary and ecological genetics (Jansson & Douglas, 2007; Douglas, 2017). Furthermore, cultivated poplars have commercial value for peeling and veneer, lumber, paper pulp and are also used as bioenergy feedstock due to their high biomass production and favourable cell wall chemistry (Porth & El-Kassaby, 2015; Taylor et al., 2016; An et al., 2021; Abreu et al., 2022). *Populus nigra* is a deciduous tree species native to Europe, Asia and North Africa that occupies riparian ecosystems with diverse climate ranges (De Rigo et al., 2016). The genetic structure of this species in its natural distribution area is not extensively known. Yet, some studies have shown high genetic diversity within populations and low but significant genetic differentiation between river basins, suggesting high levels of gene flow in Western parts of the distribution (Smulders et al., 2008; Dewoody et al., 2015; Wójkiewicz et al., 2021). Seven ancestral genetic clusters were found in the first genome-wide genotyping study of 838 native individuals from 12 Western European river basins (Faivre-Rampant et al., 2016). However, another study of seven species showed that black poplar is highly structured with low diversity within populations (Milesi et al., 2024). These results may be due to the fact that the ecology of the species is strongly influenced by a very dynamic environment, the alluvial banks where it breeds, resulting in a complex structure (Gurnell & Petts, 2006; Alimpić et al., 2022).

Black poplar also shows a wide phenotypic diversity which can be observed on latitudinal clines such as that observed for leaf functional traits in response to drought (Viger et al., 2016) or on leaf morphology and structure (Guet et al., 2015b). Among the observable phenotypes, growth traits and wood production are considered fundamental for the adaptation and productivity of planted forests (Grattapaglia et al., 2009). For example, the biosynthetic pathway of lignin, an essential component of wood, is known to affect abiotic tolerance and growth in *Populus* (Xie et al., 2018). However, few genetic studies have been carried out on traits related to growth, and even fewer at the genomic level using the natural intraspecific diversity of trees. Genetic differentiation between natural populations of *P. trichocarpa* was found for growth and phenology, which was higher than the rather weak differentiation observed at the genome level (Evans et al., 2014; Oubida et al., 2015). This suggests that local adaptation explains patterns of variation in these traits better than genetic drift alone. The adaptive traits of poplar populations show variations depending on the local climate at their geographic origin. Using genome-wide association studies (GWAS) on provenances of *P. trichocarpa*, candidate loci underlying bud phenology and biomass have already been identified (Evans et al., 2014; Zhang et al., 2019). Based on 113 natural *P. tremula* genotypes from Sweden, a study showed significant natural variation in growth and wood-related traits and allowed the identification of genetic markers associated with these traits (Escamez et al., 2023). In this context, the OGDH enzyme (2-oxoglutarate dehydrogenase) was found to be associated with variation in tree volume and constitutes an interesting potential candidate for improving stem volume. Within the same collection of Aspen trees, a major and unique locus was also discovered. It determines the timing of bud formation and facilitates adaptation to different growing seasons and colder climates

(Wang et al., 2018). A systems genetics approach in a subset of the same collection linked
natural variation in lignin content and composition to responses to mechanical stimuli and
nutrient availability (Luomaranta et al., 2024). Furthermore, QTLs were identified for stem and
biomass traits in several mapping populations involving as parental species those typically used
to generate cultivated hybrids (*P. deltoides*, *P. nigra* and *P. trichocarpa*). Of note, these studies
reported several QTL hotspots for biomass accumulation in different environments (Rae et al.,
2008, 2009; Dillen et al., 2009; Monclus et al., 2012).
Although QTL mapping studies in segregating progenies have reported QTL hotspots that
explain a large part of genetic variation for growth, the QTL resolution was too limited to identify
the underlying candidate genes (Rae et al., 2009). On the other hand, GWAS can make use of
the rapid decay of linkage disequilibrium in forest trees (Neale & Kremer, 2011), but most studies
carried out so far for growth traits have reported a limited number of loci that individually do not
explain a large proportion of the genetic variance of this heritable trait (Mckown et al., 2014;
Allwright et al., 2016). Many studies suggest that complex traits are controlled by multiple loci,
each with rather small effects (Bradshaw & Stettler, 1995; Grattapaglia et al., 1996; Rae et al.,
2007; Wade et al., 2022). To go further in understanding phenotypes and adaptation, the
genomics toolbox and statistical methods as systems biology approach made available for
research are constantly evolving (Pazhamala et al., 2021). The revolution comes in particular
from the applications that "omics" technologies have made possible for plants such as forest
trees (Plomion et al., 2016; Borthakur et al., 2022). Thus, with the progression of methodologies
and the reduction in the costs of these approaches, a certain number of studies have examined
at large scale of endophenotypes like transcriptomic (Chateigner et al., 2020), proteomic
(Plomion et al., 2006; Castillejo et al., 2023; Teyssier et al., 2023) or even metabolomic
(Rodrigues et al., 2021) in trees. Another study has advocated the use of RNAseq to jointly
identify polymorphisms and quantify the transcriptomic variability across natural populations (De
Wit et al., 2015). Such an approach could contribute to filling the gap between the genome and
phenotypic variation for complex traits and further contribute to the explanation of their missing
heritability (Maher, 2008; Chandler et al., 2014).
Here, we report a GWAS for growth using phenotypic data from natural populations of *P.*
nigra evaluated in two common garden experiments together with SNP data from RNAseq
(Chateigner et al., 2020; Rogier et al., 2023). We further make use of the transcriptomic data to
dissect a major QTL for growth and pinpoint a candidate gene from the flavonoid pathway.
Finally, we studied the genomic and transcriptomic diversity of the candidate gene and the
phenotypic diversity across the natural populations and could show that the polymorphism is
involved in growth differentiation, suggesting an implication in local adaptation.

Material and methods

Plant material and field experiments.

The complete plant material and field management was previously described (Guet et al.,
2015a; Gebreselassie et al., 2017). Briefly, an initial experimental design based on a total of
1,160 genotypes of *P. nigra*, representative of the species range in Western Europe, was
established in two contrasting common gardens located at Orléans (France, ORL) and
Savigliano (Italy, SAV) in 2008. In both sites, the genotypes were replicated 6 times in a
randomized complete block design. A previous study, using a 12k Infinium array (Faivre-
Rampant et al., 2016) was used to characterize the genetic diversity within this collection. A
subset of 241 genotypes representative of the natural diversity and originating from 10 river
basins was selected.

Climate data.

Climatic variations across the locations of origin of the populations were analysed by applying
principal component analysis on 19 annual bioclimatic variables obtained from the WorldClim

dataset (Hijmans et al., 2005). The values used are the 30-year average (1960 to 1990) with a
resolution of 1 km² per grid cell obtained from the GPS location of the original natural
populations. The first two principal components (PC1 and PC2) corresponded to the weighted
precipitation and temperature variables, respectively.

Phenotyping.

We have described in detail the phenotyping of 21 traits in previous works (Chateigner et al.,
2020; Wade et al., 2022). Only the circumference and the basic density of the wood (Infraden)
were used in this study. Briefly, trees were pruned at the base after one (SAV) or two years of
growth (ORL), to remove a potential cutting effect. Circumferences at 1-m above the ground
were measured on 2-year-old trees in winter 2010-2011 at SAV and in winter 2011-2012 at ORL.
Basic density was determined as previously reported as described in (Chateigner et al., 2020).
Briefly, it was measured on a piece of wood from the stem section harvested for RNA
sequencing (see hereafter) following the Technical Association of Pulp and Paper Industry
(TAPPI) standard test method T 258 "Basic density and moisture content of pulpwood". For each
site, the phenotypic data were analyzed with a linear mixed model to compute genotypic means
adjusted for micro-environmental effects as described in (Gebreselassie et al., 2017). Before the
adjustment of the model, a square root transformation was made to ensure the normality and
homoscedasticity of the residuals.

Transcriptomic data.

RNA sequencing was carried out on young differentiating xylem and cambium tissues
collected in 2015 from two replicates of the 241 genotypes located in two blocks of the Orleans
common garden, as described in (Chateigner et al., 2020). Sequencing reads were obtained to
provide both transcriptomic and genomic data. Briefly, frozen milled tissue was used to isolate
total RNA with RNeasy Plant kit (Qiagen, France), according to manufacturer's recommendations
and a treatment with DNase I (Qiagen, France) was made. Samples of young differentiating
xylem and cambium tissues of the same tree were pooled in an equimolar extract before sending
it for the sequencing at the POPS platform with Illumina Hiseq2000. Reads were mapped to the
P. trichocarpa v3.0 primary transcripts using bowtie2 v2.4.1 (Langmead & Salzberg, 2012) and
only transcripts with at least 1 count in 10% of the samples were kept, yielding 34,229 features.
The raw count data were normalized by Trimmed Mean of M-values using the R package edgeR
v3.26.4, calculated in counts per millions (CPM) and computed in $\log_2(n+1)$. At the end, the
CPM were fitted with a linear mixed model including batch and genetic effects to extract their
genotypic Best Linear Unbiased Predictors (BLUPs). These genotypic BLUPs of transcripts were
used for the rest of our analysis.

Genotypic data.

The full details of genotypic analysis have been described in (Rogier et al., 2023), including
software used, data filtering criteria and final SNP selection. Briefly, genotyping data were
obtained, using BWA-MEM v0.7.12 to map the reads into the *P. trichocarpa* v3.0 reference
genome and the SNPs were called using 3 callers to generate a high-confidence SNP set. Only
the SNPs identified by at least 2 of the 3 callers and with less than 50% of missing values were
selected. Missing values were imputed using the Fimpute v.2.2 program (Sargolzaei et al., 2014)
and complementary genotyping data previously obtained with a 12 k Illumina Infinium Bead-Chip
array (Faivre-Rampant et al., 2016). At the end, we obtained 878,957 SNPs and from these,
440,292 SNPs were retained for this study after filtering for a minimum allele frequency of 0.05.

Genetic analyses

185   Unless otherwise stated, all analyses have been carried out with R v4.4.1 (R Core Team,
186 2021) under the RStudio environment (RStudio Team, 2020).

 Partition of variance
188   The following bivariate mixed model was fitted to partition the variance in circumference
189   across the two sites into between- and within-population genetic variation and their interaction
190   with site:

191   (1)
$$y = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = X\beta + Z_b b + Z_w w + \epsilon$$

Where y is a vector of genotypic adjusted means for circumference in ORL and SAV, X, Z_b
and Z_w are design matrices relating observations to fixed and random effects, β is the fixed
effect of site and b and w are between and within random genetic effects. b and w follow a

multivariate normal distribution with mean 0 and variances: $\begin{bmatrix} \sigma^2_{b_1} & \sigma_{b_{12}} \\ \sigma_{b_{21}} & \sigma^2_{b_2} \end{bmatrix} \otimes K_b$ and

$\begin{bmatrix} \sigma^2_{w_1} & \sigma_{w_{12}} \\ \sigma_{w_{21}} & \sigma^2_{w_2} \end{bmatrix} \otimes K_w$. K_b and K_w are genomic relationship matrices between and within

populations. They were estimated from the full genomic relationship matrix computed with ldak
software v5 (Speed et al., 2012), by averaging the kinships per population for K_b and setting the
kinships at zero across populations for K_w.

Population genetics
F_{ST} was estimated using Weir and Cockerham method (Weir & Cockerham, 1984) and
implemented in plink (v1.90b6.3). Q_{ST} was estimated using variance parameters from the

previously described mixed-model as: $Q_{ST} = \dfrac{\sigma^2_b}{\left(\sigma^2_b + 2\sigma^2_w\right)}$.

GWAS
GWAS was performed for circumference in each site with genotypic adjusted means and
SNPs, using a linear mixed model as originally proposed by (Yu et al., 2005) and implemented in
the R package MM4LMM (Laporte et al., 2022). This model included a random polygenic effect
with a covariance structure defined by a genomic relationship matrix computed with the software
ldak to account for linkage disequilibrium between SNPs (Speed et al., 2012). We also performed
multi-locus GWAS using the multi-locus mixed-model (MLMM) approach implemented in the R
package MLMM v0.1.1 (Segura et al., 2012), as well as multi-environment GWAS carried out
with the MTMM approach implemented in R (Korte et al., 2012). Linkage disequilibrium between
significantly associated SNPs was estimated in R as the squared allelic coefficient.
GWAS were also carried out using transcriptomic data (eQTL analysis) but focusing only on 2
genes of particular interest in this work. The analyses were done using both single- and multi-
locus approaches, as presented for circumference.
We also looked at associations between our candidate SNP, latitude of origin and climatic
data at the population level using a Pearson correlation test.
Further tests were carried with data previously published by (Pégard et al., 2020) on a multi-
parental population of *P. nigra* (factorial mating design). This dataset consisted of 629 individuals
with genotypic and circumference data. We retrieved 46 SNPs within the interval
[chr10:20105000, chr10:20125000] corresponding to the region of interest in the present study,
and carried out association tests between these SNPs and the phenotype using a simple linear
model.

Results

A QTL controlling radial growth is highlighted by a genome-wide association study.

We performed a GWAS for circumference using 428,836 SNPs and detected a significant signal for this trait phenotyped in Savigliano (**Fig. 1a**), with a total of 18 significant SNPs, including 11 on chromosome 10 in strong linkage disequilibrium. Closer examination of this region showed that the signal is distributed over two gene models: Potri.010G212900, annotated as a Beta-Hexosaminidase 1 (Hexo1) and Potri.010G213000, annotated as a chalcone isomerase family protein (CHI) (**Fig. 1b**). In the MLMM approach, the whole signal vanishes out after conditioning on the top SNP, suggesting that a single allele is associated with the trait in the region (**Fig. S1**). This top SNP explains more than 50% of the phenotypic variation (without accounting for population structure, **Fig. 1c**). While non-significant at the genome-wide level when considering circumference at Orleans (**Fig. S2**), this top SNP still explains more than 20% of the phenotypic variation in this common garden and its effect is in the same direction as found in Savigliano (**Fig. 1c**). Consequently, a multi-trait GWAS combining phenotypes from the two common gardens confirmed this signal but detected only a total of 7 significant SNPs (**Fig. S3**), mainly for the global effect (i.e., common to the two sites). These 7 SNPs identified in the multi-trait GWAS, are included in the 11 detected in single-trait GWAS at Savigliano and constitute our core set of candidate SNPs (**Tab. S1**). Among them, 6 are exonic (4 non-synonymous and 2 synonymous) and 1 is in 5'UTR, unsurprisingly as they come from RNAseq reads. In addition, they are all located on the CHI gene except one. It is worth mentioning that the top SNP is located in an exon of CHI gene and is predicted to be non-synonymous.

Figure 1 - GWAS of the circumference phenotype. a) Genome-wide Manhattan plot highlighting QTL on chromosome 10 performed using a single locus mixed model and 428,836 SNPs markers from natural *P. nigra* diversity phenotype at Savigliano; b) Manhattan plots focused on SNPs with lowest p-values obtained and concerning 2 gene regions, with corresponding mean coverage of RNAseq reads across individuals; c) Box plot of the circumference in both experimental sites (transformed with a square root, sqrt), depending on the allele count of the candidate SNP with the lowest p-value.

256 **Gene expression sustains CHI as a candidate gene.**

257   We used the RNAseq data, generated from the xylem and cambium tissues of poplars grown
258 at Orléans as an endophenotype to test whether the expression of our candidate genes
259 correlated with the phenotypes and could be linked to the effect of one of them. Negative
260 correlations were found between gene expressions and phenotypes, and their magnitude was
261 higher for CHI than for Hexo1 (**Fig. 2a**, **Fig. 2b**), with R^2 of 0.53 and 0.35 for circumference
262 evaluated in Savigliano and Orleans, respectively. When using the expression of both genes to
263 jointly explain phenotypes, the correlation between CHI gene expression and circumference was
264 maintained (R^2 = 0.49 at Savigliano and R^2 = 0.25 at Orleans) while it drastically dropped for
265 Hexo1 (R^2 = 0.004 at Savigliano and R^2 = 0.05 at Orleans). We also made use of transcriptomic
266 data for CHI and Hexo1 to perform an eQTL analysis, which highlighted a strong cis control for
267 the 2 genes (**Fig. 2c**). The fact that these two genes are close from each other and in opposite
268 directions on the genome, together with the existence of strong LD in the region (**Fig. 1b**),
269 generates a positive correlation between their expressions (R^2 = 0.19). But, when focusing on the
270 region of interest, we observed different patterns of eQTL signal between the 2 genes (**Fig. 2d**).
271 Interestingly, the pattern of eQTL for CHI gene was similar to the one observed for circumference
272 (**Fig. 2d**, **Fig. 1b**). Altogether, these results supported CHI as a candidate gene for the control of
273 circumference variability.

274
275

277   **Figure 2** - eQTL analysis sustains CHI (Potri.01G213000) as a candidate gene for
278   the control of circumference variation. a) correlation between the circumference
279   and the expression level of Hexo1 primary transcript (Potri.010G212900.1) ; b)
280   correlation between the circumference and the expression level of CHI primary
281   transcript (Potri.010G213000). c) Manhattan plot and d) focus on the candidate
282   region of the eQTL analysis using the variations in the expression level of the 2
283   primary transcripts previously highlighted as phenotypes. Circumference was
284   transformed with a square root (sqrt). The expression level of transcripted have
285   been standardized with a genetic analysis.

Structure of the diversity of the CHI gene highlighted by population-scale analyses.

To further characterize the effect of the top SNP on the phenotypic variability, we partitioned the variance of circumference across locations into between-population and within-population genetic effects, their interaction with location, and a residual term (**Fig. 3**). This analysis showed that a large part of the phenotypic variation (35%) was due to genetic differences between populations, followed by interaction variance between genetics within populations and location (25%), genetic variance across populations (20%), and interaction variance between genetics across populations and location (17%). Interestingly, when the top SNP was included as a fixed effect in this variance partitioning model, it explained up to 24% of the total phenotypic variance, and this part of variability was mainly from the between population genetic component (**Fig. 3**, model 2). This analysis suggests that the QTL, previously identified by GWAS, is driven by differences in radial growth at the population level.

Figure 3 - Partition of phenotypic variance for circumference across two locations using two models: Model 1 (mod.1) refers to the model of variance partition without the top SNP (Chr10:20120195), while model 2 (mod. 2) is the model that includes the top SNP as a cofactor. Btw-pop and With-pop refer to between and within population variances, while G and GE refer to genetic and genetic by environment variance, respectively.

To confirm this observation, we computed the fixation index (F_{ST}) of the 428,836 SNPs and looked at the value of the top SNP detected by the GWAS. This SNP displayed a high F_{ST} value (0.69) well above the 99[th] percentile (0.28) of the genome-wide F_{ST} distribution (**Fig. 4a**). Such a high fixation index is due to a fixation of the reference allele in several populations mainly from the north-east of the studied area (NL, Kuhkopf, Rhin, Ticino), a fixation of the alternative allele in some population from central (Loire, Val d'Allier) and southern (Ramières) France and southern Italy (Basento), and a balanced situation in intermediate populations between these extremes (Dranse and Paglia) as well as in the population of south-western France (Adour) (**Fig. 4b**). Interestingly, such a genetic differentiation is also observed at the phenotypic level as well as at the transcriptomic level for CHI gene, as highlighted by high Q_{ST} values (**Fig. 4a**) and population differences (**Fig. 4c**). Consequently, associations between SNP and traits (**Fig. 5a**) or gene expression (**Fig. 5b**), as well as correlations between traits and gene expression (**Fig. 5c**), were high and significant when estimated at the population level, except for the trait evaluated at Orleans, which is consistent with the results obtained at the individual level (**Fig. S4**).

Figure 4 - Structure of the diversity. a) Distribution of genome-wide Fst together with specific values indicated by vertical lines: 99th percentile of the distribution, top SNP (Chr10:20120195) Fst, CHI (Potri.010G213000.1) expression Qst, circumference at Orléans and Savigliano Qsts. b) Geographical origin of populations together with the distribution of alleles within each population for the top SNP (Chr10:20120195), the size of the pie is proportional to the size of the population. c) Distribution of circumferences at Orléans and Savigliano, as well as CHI (Potri.010G213000.1) expression across populations.

Figure 5 - Associations at the population scale. a) Correlation between circumference at Savigliano and allele frequencies for the top SNP (Chr10:20120195); b) correlation between CHI (Potri.010G213000.1) expression and allele frequencies for the top SNP (Chr10:20120195); c) correlation between circumference at Savigliano and CHI (Potri.010G213000.1) expression.

Validations.

To support our findings, we complemented our study by several analyses. First, we looked at
co-localizations between the QTL detected in the present study and QTLs previously reported in
the literature. Of particular interest, we found in the same genomic region a QTL previously
reported by Rae et al. (2009) for several traits related to biomass production in an interspecific
poplar progeny, and named poplar biomass locus 3 (PBL3). PBL3 included several QTLs for
height and diameter found across multiple years, and it was delimited by two SSRs (ORPM149
and PMGC2786). We retrieved the coordinates of these markers on the *P. trichocarpa* reference
genome by blasting their priming sequences. The resulting interval in bp was [17566502,
21189318] (**Fig. S5**). It thus fully includes the QTL reported here which spans the interval
[20105000, 20125000] (**Fig. 1b**). Second, we retrieved data from intraspecific crosses of *P. nigra*
carried out within the French breeding program and previously used and reported by (Pégard et
al., 2020) for genomic prediction. From the SNP set in this previous study, we identified 46 SNPs
that fell within the interval and tested associations between each of these SNPs and the
phenotype circumference in a panel of 629 individuals resulting from those crosses. The most
significant association (p = 2.43e-05) was found for a SNP located at 20 119 788 bp (407 bp
from the top SNP) (**Fig. S6**), which was also found significant in the present study with an effect
in the same direction (alternative allele associated with an increase in circumference).

Discussion

We made use of growth data collected in two common garden experiments together with
transcriptome-wide SNP data to search for genetic associations between genotype and
phenotype in *P. nigra*. Such analysis pinpointed a small genomic region located at the distal end
of chromosome 10 which encompassed 2 gene models, of which one was annotated as a
chalcone isomerase (CHI). Transcriptomic data within one of the two common gardens further
supported an implication of CHI in the phenotypic variation. Because the black poplar collection
was structured into subpopulations corresponding to the geographic origins of the accessions,
we further focused on differences between subpopulations and found that CHI diversity is a main
driver of growth differences at the subpopulation scale. Such findings suggest an implication of
this gene in local adaptation. Finally, we seek to validate our results with data from previous
works and found that our significant loci match a previously reported QTL hotspot for biomass
accumulation in an interspecific poplar family (Rae et al., 2009). We further validated the effect of
the QTL in an independent panel with a *P. nigra* pedigree from the French breeding program
(Pégard et al., 2020).
The strongest effect in the GWAS was found for the phenotypic data collected in the common
garden (SAV) where the genetic variability for growth was the largest. This site enabled a better
expression of the phenotypic variability for growth. Unfortunately, transcriptomic evaluation was
carried out in the other common garden (ORL). Consequently, it is hard to conclude on the
interplay between SNP variation and gene expression to explain the variation in growth. Indeed,
if we run a mediation analysis, as proposed by Sasaki et al. (2018), using phenotypic data from
SAV, we cannot conclude that the expression of CHI mediates the genetic association (data not
shown). While if we repeat such analysis with phenotypic data from ORL we find that the
association is mediated by CHI expression, although the association with growth at ORL is not
significant genome-wide. Yet, the fact that gene expression data were collected from a different
site than the one in which the GWAS is significant, underlines the robustness of the results.
Another complication with the loci detected originates from the confounding effect of
population structure. Indeed, the phenotype, gene expression as well as polymorphisms display
a significant variability across populations which drives the correlations and associations
between them. Such a situation is not ideal for association mapping, and the confounding effect
attributed to population structure has to be specifically handled by the statistical model applied,
usually a linear mixed model with a random polygenic effect having as covariance the genomic
relationship between individuals (Yu et al., 2005). In addition, we used the program ldak to

account for LD between polymorphisms in the estimation of the relationships (Speed et al.,
2012), required to be independent, because our SNPs come from RNAseq and are thus
clustered by genes with potentially some strong LD between neighbouring SNPs. The correction
applied within the linear mixed model with such a matrix appeared to be milder than the one
achieved with a regular GRM such as the one estimated following (VanRaden, 2008), resulting in
a significant signal. Such a complication underlines the need to validate the association, which
was achieved through two main approaches. First, we found that the detected locus falls within a
QTL hotspot for biomass previously reported in several mapping populations (Rae et al., 2008,
2009; Dillen et al., 2009; Monclus et al., 2012). Second, we have shown that one of the
significant SNP affects also the growth in a large collection of *P. nigra* from a breeding pedigree
previously used for testing genomic prediction in black poplar (Pégard et al., 2020). While
statistically significant, the effect of the SNP is lower than in the natural populations. This could
potentially be explained by GxE interaction since we already found that the SNP effect is different
between the two common garden experiments and the *P. nigra* pedigree from Pégard et al.
(2020) was evaluated in a different location within a quite different climatic area (oceanic
climate).
Another way of validating the locus would be to gain insights into the biological mechanism
relating the polymorphisms to the trait through the expression of CHI. Considering the
polymorphisms, we could identify four non-synonymous SNPs significantly associated with the
phenotype, one in the first exon and three in the second exon of the gene, including the top SNP
(**Tab. S1**). We could hypothesize that one or several of these SNPs affect the enzymatic activity
of CHI, for which a cellular response could be an overexpression of the gene as compensation.
This would be consistent with the observed positive relationship between the most significant
polymorphism and the expression of CHI (**Fig. 5b**). Also, the decrease in the enzymatic activity
of CHI for individuals carrying the alternate allele of the top SNP could be consistent with the
decrease in growth observed in these individuals (**Fig. 1c**). Another hypothesis to explain the
negative correlation between growth and CHI expression could be a trade-off between growth
and wood quality (Novaes et al., 2010). To test this hypothesis, we retrieved data on wood
density measured on samples collected at Orléans. The top SNP displayed a significant
association with wood density, with a positive effect of the alternate allele, which was thus
opposite to the effect found for circumference (**Fig. S7a**). Similarly, a significant positive
correlation was found between wood density and CHI gene expression while such correlation
was positive for circumference (**Fig. S7b**). These results provide some evidence for the effect of
CHI on the trade-off between wood growth and density. Interestingly, a preceding study revealed
that silencing hydroxycinnamoyl-CoA shikimate/quinate hydroxycinnamoyl transferase (HCT) in
Arabidopsis thaliana involves an accumulation of flavonoids and a reduction of plant growth
(Besseau et al., 2007). This is consistent with our results since CHI is one of the first key
enzymes of the flavonoid pathway (Dare et al., 2020). HCT is a key enzyme of the lignin
pathway, which is well documented and consists of a metabolic grid that modifies phenylalanine
in multiple steps to ultimately produce the monolignols p-coumaryl, coniferyl, and sinapyl
alcohols. In a previous study, screening a wide diversity of populations and focusing on the lignin
biosynthesis pathway made it possible to identify common and rare functional variants in several
genes (Marroni et al., 2011), including a natural defective allele for HCT (Vanholme et al., 2013).
However, no effect on growth could be detected in this work.
When looking at the loci diversity at the population level we found a strong differentiation, far
above the genome-wide level (**Fig. 4**). Such a differentiation is thus more likely to result from
differential selection than genetic drift. Of particular interest, the differentiation across natural
populations was also found for CHI expression and circumference (**Fig. 4**), and we could show
that the top SNP contributed mainly to the between-population component of genetic variation for
growth (**Fig. 3**). As a result, highly significant correlations were found between allele frequencies,
gene expression, and phenotype at the population level (**Fig. 5**). When looking at the repartition
of alleles on a map representing the geographic origin of the populations, a clear North-East
versus South-West differentiation appears. Such a tendency was confirmed by the significant

correlation found between latitude of origin and allele frequencies (**Fig. S8a**). One could thus
hypothesize that the differentiation could be related to climatic differences across Western
Europe, which was confirmed by the significant correlation detected between allelic frequencies
and a temperature proxy of the climate of origin (**Fig. S8b**). If we go back to the phenotypic data
across populations, it's worth noting that the southern populations with the alternate allele fixed
display a lower growth and higher wood density. These data support the idea that southern
populations are growing slowly as an adaptation to high summer temperatures, which ultimately
underlines the adaptive relevance of the locus reported here.
This work strengthens the interest in combining transcriptomics with genomics data across
large natural populations to unravel locus and genes involved in key adaptive processes such as
the trade-off between growth and wood formation. Such results provide some guidance to breed
future varieties of trees with improved efficiency to store carbon.

Acknowledgements

455 We thank GBFOR (INRAE, Forest Genetics and Biomass Facility),
456 https://doi.org/10.15454/1.5572308287502317E12 for management of the common garden
457 experiment.

Funding

459   This work was done within the SYBIOPOP project (ANR-13-JSV6-0001) funded by the
460 French National Research Agency (ANR). The platform POPS benefits from the support of the
461 LabEx Saclay Plant Sciences-SPS (ANR-10-LABX-0040-SPS).

Conflict of interest disclosure

463   The authors declare that they comply with the PCI rule of having no financial conflicts of
464 interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

Data are available online: https://doi.org/10.57745/DSBTGG.
Information about the RNA-seq project, from which the gene expression data come, is
available in the Gene Expression Omnibus (GEO) from NCBI (accession number: GSE128482).
Raw sequences (FASTQ) are available in the Sequence Read Archive (SRA) from NCBI
(accession number: SRP188754).
Information on the studied genotypes is available in the GnpIS Information System
(Steinbach et al., 2013) via the FAIDARE data portal (https://urgi.versailles.inra.fr/faidare/), using
the keys "Black poplar" and "POPULUS NIGRA RNASEQ PANEL" for the fields "Crops" and
"Germplasm list", respectively.
Scripts and code are available online: https://forgemia.inra.fr/vincent.segura/sybiopop_chi.git.
Supplementary information is available online: https://doi.org/10.5281/zenodo.13950469

References

Abreu M, Silva L, Ribeiro B, Ferreira A, Alves L, Paixão SM, Gouveia L, Moura P, Carvalheiro F,
Duarte LC, et al. 2022. Low Indirect Land Use Change (ILUC) Energy Crops to Bioenergy and
Biofuels—A Review. *Energies*, **15**, 4348. https://doi.org/10.3390/en15124348
Alimpić F, Milovanović J, Pielech R, Hinkov G, Jansson R, Dufour S, Beza M, Bilir N, del Blanco
LS, Božič G, et al. 2022. The status and role of genetic diversity of trees for the conservation
and management of riparian ecosystems: A European experts' perspective. *Journal of*
Applied Ecology, **59**, 2476–2485. https://doi.org/10.1111/1365-2664.14247

Allwright MR, Payne A, Emiliani G, Milner S, Viger M, Rouse F, Keurentjes JJB, Bérard A, Wildhagen H, Faivre-Rampant P, et al. 2016. Biomass traits and candidate genes for bioenergy revealed through association genetics in coppiced European Populus nigra (L.). *Biotechnology for Biofuels and Bioproducts*, **9**, 1–22. https://doi.org/10.1186/s13068-016-0603-1

An Y, Liu Y, Liu Y, Lu M, Kang X, Mansfield SD, Zeng W, Zhang J. 2021. Opportunities and barriers for biofuel and bioenergy production from poplar. *GCB Bioenergy*, **13**, 905–913. https://doi.org/10.1111/gcbb.12829

Besseau S, Hoffmann L, Geoffroy P, Lapierre C, Pollet B, Legrand M. 2007. Flavonoid Accumulation in Arabidopsis Repressed in Lignin Synthesis Affects Auxin Transport and Plant Growth. *The Plant Cell*, **19**, 148–162. https://doi.org/10.1105/tpc.106.044495

Borthakur D, Busov V, Cao XH, Du Q, Gailing O, Isik F, Ko JH, Li C, Li Q, Niu S, et al. 2022. Current status and trends in forest genomics. *Forestry Research*, **2**. https://doi.org/10.48130/FR-2022-0011

Bradshaw HD, Stettler RF. 1995. Molecular genetics of growth and development in Populus. IV. Mapping QTLs with large effects on growth, form, and phenology traits in a forest tree. *Genetics*, **139**, 963–973. https://doi.org/10.1093/genetics/139.2.963

Brienen RJW, Caldwell L, Duchesne L, Voelker S, Barichivich J, Baliva M, Ceccantini G, Di Filippo A, Helama S, Locosselli GM, et al. 2020. Forest carbon sink neutralized by pervasive growth-lifespan trade-offs. *Nature Communications*, **11**, 1–10. https://doi.org/10.1038/s41467-020-17966-z

Bryant N, Zhang J, Feng K, Shu M, Ployet R, Chen J-G, Muchero W, Yoo CG, Tschaplinski TJ, Pu Y, et al. 2023. Novel candidate genes for lignin structure identified through genome-wide association study of naturally varying Populus trichocarpa. *Frontiers in Plant Science*, **14**, 1373. https://doi.org/10.3389/fpls.2023.1153113

Castillejo MA, Pascual J, Jorrín-Novo J V., Balbuena TS. 2023. Proteomics research in forest trees: A 2012-2022 update. *Frontiers in Plant Science*, **14**, 1130665. https://doi.org/10.3389/fpls.2023.1130665

Chandler CH, Chari S, Tack D, Dworkin I. 2014. Causes and consequences of genetic background effects illuminated by integrative genomic analysis. *Genetics*, **196**, 1321–1336. https://doi.org/10.1534/genetics.113.159426

Chateigner A, Lesage-Descauses M-C, Rogier O, Jorge V, Leplé J-C, Brunaud V, Roux CP-L, Soubigou-Taconnat L, Martin-Magniette M-L, Sanchez L, et al. 2020. Gene expression predictions and networks in natural populations supports the omnigenic theory. *BMC Genomics*, **21**. https://doi.org/10.1186/s12864-020-06809-2

Dare AP, Tomes S, McGhie TK, van Klink JW, Sandanayaka M, Hallett IC, Atkinson RG. 2020. Overexpression of chalcone isomerase in apple reduces phloridzin accumulation and increases susceptibility to herbivory by two-spotted mites. *The Plant Journal*, **103**, 293–307. https://doi.org/10.1111/tpj.14729

Dewoody J, Trewin H, Taylor G. 2015. Genetic and morphological differentiation in Populus nigra L.: isolation by colonization or isolation by adaptation? *Molecular Ecology*, **24**, 2641–2655. https://doi.org/10.1111/mec.13192

Dillen SY, Storme V, Marron N, Bastien C, Neyrinck S, Steenackers M, Ceulemans R, Boerjan W. 2009. Genomic regions involved in productivity of two interspecific poplar families in Europe. 1. Stem height, circumference and volume. *Tree Genetics and Genomes*, **5**, 147–164. https://doi.org/10.1007/s11295-009-0199-8

Douglas CJ. 2017. Populus as a Model Tree. In: Comparative and Evolutionary Genomics of Angiosperm Trees. Springer, Cham, 61–84. https://doi.org/10.1007/7397_2016_12

Escamez S, Robinson KM, Luomaranta M, Gandla ML, Mähler N, Yassin Z, Grahn T, Scheepers G, Stener LG, Jansson S, et al. 2023. Genetic markers and tree properties predicting wood biorefining potential in aspen (Populus tremula) bioenergy feedstock. *Biotechnology for Biofuels and Bioproducts*, **16**, 1–16. https://doi.org/10.1186/s13068-023-02315-1

Evans LM, Slavov GT, Rodgers-Melnick E, Martin J, Ranjan P, Muchero W, Brunner AM, Schackwitz W, Gunter L, Chen JG, et al. 2014. Population genomics of Populus trichocarpa identifies signatures of selection and adaptive trait associations. *Nature Genetics*, **46**, 1089–1096. https://doi.org/10.1038/ng.3075

Faivre-Rampant P, Zaina G, Jorge V, Giacomello S, Segura V, Scalabrin S, Guérin V, De Paoli E, Aluome C, Viger M, et al. 2016. New resources for genetic studies in Populus nigra: genome-wide SNP discovery and development of a 12k Infinium array. *Molecular Ecology Resources*, **16**, 1023–1036. https://doi.org/10.1111/1755-0998.12513

Gebreselassie MN, Ader K, Boizot N, Millier F, Charpentier JP, Alves A, Simões R, Rodrigues JC, Bodineau G, Fabbrini F, et al. 2017. Near-infrared spectroscopy enables the genetic analysis of chemical properties in a large set of wood samples from Populus nigra (L.) natural populations. *Industrial Crops and Products*, **107**, 159–171. https://doi.org/10.1016/j.indcrop.2017.05.013

Grattapaglia D, Bertolucci FLG, Penchel R, Sederoff RR. 1996. Genetic Mapping of Quantitative Trait Loci Controlling Growth and Wood Quality Traits in Eucalyptus grandis Using a Maternal Half-sib Family and RAPD Markers. *Genetics*, **144**, 1205–1214. https://doi.org/10.1093/genetics/144.3.1205

Grattapaglia D, Plomion C, Kirst M, Sederoff RR. 2009. Genomics of growth traits in forest trees. *Current Opinion in Plant Biology*, **12**, 148–156. https://doi.org/10.1016/j.pbi.2008.12.008

Green JK, Keenan TF. 2022. The limits of forest carbon sequestration. *Science*, **376**, 692–693. https://doi.org/10.1126/science.abo6547

Guet J, Fabbrini F, Fichot R, Sabatti M, Bastien C, Brignolas F. 2015a. Genetic variation for leaf morphology, leaf structure and leaf carbon isotope discrimination in European populations of black poplar (Populus nigra L.). *Tree Physiology*, **35**, 850–863. https://doi.org/10.1093/treephys/tpv056

Guet J, Fichot R, Lédée C, Laurans F, Cochard H, Delzon S, Bastien C, Brignolas F. 2015b. Stem xylem resistance to cavitation is related to xylem structure but not to growth and water-use efficiency at the within-population level in Populus nigra L. *Journal of Experimental Botany*, **66**, 4643–4652. https://doi.org/10.1093/jxb/erv232

Gurnell A, Petts G. 2006. Trees as riparian engineers: The Tagliamento River, Italy. *Earth Surface Processes and Landforms*, **31**, 1558–1574. https://doi.org/10.1002/esp.1342

Harris NL, Gibbs DA, Baccini A, Birdsey RA, de Bruin S, Farina M, Fatoyinbo L, Hansen MC, Herold M, Houghton RA, et al. 2021. Global maps of twenty-first century forest carbon fluxes. *Nature Climate Change*, **11**, 234–240. https://doi.org/10.1038/s41558-020-00976-6

Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, **25**, 1965–1978. https://doi.org/10.1002/joc.1276

Jansson S, Douglas CJ. 2007. Populus: A Model System for Plant Biology. *Annual Review of Plant Biology*, **58**, 435–458. https://doi.org/10.1146/annurev.arplant.58.032806.103956

Korte A, Vilhjálmsson BJ, Segura V, Platt A, Long Q, Nordborg M. 2012. A mixed-model approach for genome-wide association studies of correlated traits in structured populations. *Nature Genetics*, **44**, 1066–1071. https://doi.org/10.1038/ng.2376

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nature Methods*, **9**, 357–359. https://doi.org/10.1038/nmeth.1923

Laporte F, Charcosset A, Mary-Huard T. 2022. Efficient ReML inference in variance component mixed models using a Min-Max algorithm. *PLOS Computational Biology*, **18**, e1009659. https://doi.org/10.1371/journal.pcbi.1009659

Luomaranta M, Grones C, Choudhary S, Milhinhos A, Kalman TA, Nilsson O, Robinson KM, Street NR, Tuominen H. 2024. Systems genetic analysis of lignin biosynthesis in Populus tremula. *New Phytologist*, **243**, 2157–2174. https://doi.org/10.1111/nph.19993

Maher B. 2008. Personal genomes: The case of the missing heritability. *Nature*, **456**, 18–21. https://doi.org/10.1038/456018a

Marroni F, Pinosio S, Di Centa E, Jurman I, Boerjan W, Felice N, Cattonaro F, Morgante M. 2011. Large-scale detection of rare variants via pooled multiplexed next-generation sequencing: towards next-generation Ecotilling. *The Plant Journal*, **67**, 736–745. https://doi.org/10.1111/j.1365-313X.2011.04627.x

Mckown AD, Guy RD, Klápště J, Geraldes A, Friedmann M, Cronk QCB, El-Kassaby YA, Mansfield SD, Douglas CJ. 2014. Geographical and environmental gradients shape phenotypic trait variation and genetic structure in Populus trichocarpa. *New Phytologist*, **201**, 1263–1276. https://doi.org/10.1111/nph.12601

Milesi P, Kastally C, Dauphin B, Cervantes S, Bagnoli F, Budde KB, Cavers S, Fady B, Faivre-Rampant P, González-Martínez SC, et al. 2024 Resilience of genetic diversity in forest trees over the Quaternary. *Nature Communications*, **15**, 8538. https://doi.org/10.1038/s41467-024-52612-y

Monclus R, Leplé JC, Bastien C, Bert PF, Villar M, Marron N, Brignolas F, Jorge V. 2012. Integrating genome annotation and QTL position to identify candidate genes for productivity, architecture and water-use efficiency in Populus spp. *BMC plant biology*, **12**, 173. https://doi.org/10.1186/1471-2229-12-173

Neale DB, Kremer A. 2011. Forest tree genomics: growing resources and applications. Nature Reviews Genetics 2011 12:2 12: 111–122. https://doi.org/10.1038/nrg2931

Novaes E, Kirst M, Chiang V, Winter-Sederoff H, Sederoff R. 2010. Lignin and Biomass: A Negative Correlation for Wood Formation and Lignin Content in Trees. *Plant Physiology*, **154**, 555–561. https://doi.org/10.1104/pp.110.161281

Oubida RW, Gantulga D, Zhang M, Zhou L, Bawa R, Holliday JA. 2015. Partitioning of multivariate phenotypes using regression trees reveals complex patterns of adaptation to climate across the range of black cottonwood (Populus trichocarpa). *Frontiers in Plant Science*, **6**. https://doi.org/10.3389/fpls.2015.00181

Pan Y, Birdsey RA, Fang J, Houghton R, Kauppi PE, Kurz WA, Phillips OL, Shvidenko A, Lewis SL, Canadell JG, et al. 2011. A large and persistent carbon sink in the world's forests. *Science*, **333**, 988–993. https://doi.org/10.1126/science.1201609

Pazhamala LT, Kudapa H, Weckwerth W, Millar AH, Varshney RK. 2021. Systems biology for crop improvement. *The Plant Genome*, **14**, e20098. https://doi.org/10.1002/tpg2.20098

Pégard M, Segura V, Muñoz F, Bastien C, Jorge V, Sanchez L. 2020. Favorable Conditions for Genomic Evaluation to Outperform Classical Pedigree Evaluation Highlighted by a Proof-of-Concept Study in Poplar. *Frontiers in Plant Science*, **11**, 581954. https://doi.org/10.3389/fpls.2020.581954

Plomion C, Bastien C, Bogeat-Triboulot MB, Bouffier L, Déjardin A, Duplessis S, Fady B, Heuertz M, Le Gac AL, Le Provost G, et al. 2016. Forest tree genomics: 10 achievements from the past 10 years and future prospects. *Annals of Forest Science*, **73**, 77–103. https://doi.org/10.1007/s13595-015-0488-3

Plomion C, Lalanne C, Claverol S, Meddour H, Kohler A, Bogeat-Triboulot MB, Barre A, Le Provost G, Dumazet H, Jacob D, et al. 2006. Mapping the proteome of poplar and application to the discovery of drought-stress responsive proteins. *Proteomics*, **6**, 6509–6527. https://doi.org/10.1002/pmic.200600362

Porth I, El-Kassaby YA. 2015. Using Populus as a lignocellulosic feedstock for bioethanol. *Biotechnology Journal*, **10**, 510–524. https://doi.org/10.1002/biot.201400194

R Core Team. 2021. R: A Language and Environment for Statistical Computing.

Rae AM, Pinel MPC, Bastien C, Sabatti M, Street NR, Tucker J, Dixon C, Marron N, Dillen SY, Taylor G. 2008. QTL for yield in bioenergy Populus: Identifying G×E interactions from growth at three contrasting sites. *Tree Genetics and Genomes*, **4**, 97–112. https://doi.org/10.1007/s11295-007-0091-3

Rae AM, Street NR, Robinson KM, Harris N, Taylor G. 2009. Five QTL hotspots for yield in short rotation coppice bioenergy poplar: The Poplar Biomass Loci. *BMC Plant Biology*, **9**, 1–13. https://doi.org/10.1186/1471-2229-9-23

Rae AM, Tricker PJ, Bunn SM, Taylor G. 2007. Adaptation of tree growth to elevated CO2: quantitative trait loci for biomass in Populus. *New Phytologist*, **175**, 59–69. https://doi.org/10.1111/j.1469-8137.2007.02091.x

Rathgeber CBK, Cuny HE, Fonti P. 2016. Biological basis of tree-ring formation: A crash course. *Frontiers in Plant Science*, **7**, 734. https://doi.org/10.3389/fpls.2016.00734

de Rigo D, Enescu CM, Durrant TH, Caudullo G. 2016. Populus nigra in Europe: distribution, habitat, usage and threats. In: SanMiguel-Ayanz J, de Rigo D, Caudullo G, Houston Durrant T, Mauri A (eds.), European Atlas of Forest Tree Species. Publ. Off. EU, Luxembourg, pp. e0182a4+. https://w3id.org/mtv/FISE-Comm/v01/e01aa69

Rodrigues AM, Miguel C, Chaves I, António C. 2021. Mass spectrometry-based forest tree metabolomics. *Mass Spectrometry Reviews*, **40**, 126–157. https://doi.org/10.1002/mas.21603

Rogier O, Chateigner A, Lesage-Descauses MC, Mandin C, Brunaud V, Caius J, Soubigou-Taconnat L, Almeida-Falcon J, Bastien C, Benoit V, et al. 2023. RNAseq based variant dataset in a black poplar association panel. *BMC research notes*, **16**, 248. https://doi.org/10.1186/s13104-023-06521-w

RStudio Team. 2020. RStudio: Integrated Development Environment for R.

Sargolzaei M, Chesnais JP, Schenkel FS. 2014. A new approach for efficient genotype imputation using information from relatives. *BMC Genomics*, **15**, 1–12. https://doi.org/10.1186/1471-2164-15-478

Sasaki E, Frommlet F, Nordborg M. 2018. GWAS with Heterogeneous Data: Estimating the Fraction of Phenotypic Variation Mediated by Gene Expression Data. *G3 Genes|Genomes|Genetics*, **8**, 3059–3068. https://doi.org/10.1534/g3.118.200571

Segura V, Vilhjálmsson BJ, Platt A, Korte A, Seren Ü, Long Q, Nordborg M. 2012. An efficient multi-locus mixed-model approach for genome-wide association studies in structured populations. *Nature Genetics*, **44**, 825–830. https://doi.org/10.1038/ng.2314

Smulders MJM, Cottrell JE, Lefèvre F, van der Schoot J, Arens P, Vosman B, Tabbener HE, Grassi F, Fossati T, Castiglione S, et al. 2008. Structure of the genetic diversity in black poplar (Populus nigra L.) populations across European river systems: Consequences for conservation and restoration. *Forest Ecology and Management*, **255**, 1388–1399. https://doi.org/10.1016/j.foreco.2007.10.063

Speed D, Hemani G, Johnson MR, Balding DJ. 2012. Improved heritability estimation from genome-wide SNPs. *American Journal of Human Genetics*, **91**, 1011–1021. https://doi.org/10.1016/j.ajhg.2012.10.010

Steinbach D, Alaux M, Amselem J, Choisne N, Durand S, Flores R, Keliet A-O, Kimmel E, Lapalu N, Luyten I, et al. 2013. GnpIS: an information system to integrate genetic and genomic data from plants and fungi. *Database*, **2013**, bat058. https://doi.org/10.1093/database/bat058

Taylor G, Allwright MR, Smith HK, Polle A, Wildhagen H, Hertzberg M, Bhalerao R, Joost Keurentjes JJB, Scalabrin S, Scaglione D, et al. 2016. Bioenergy trees: Genetic and genomic strategies to improve yield. In: Barth S, Murphy-Bokern D, Kalinina O, Taylor G, Jones M (eds) Perennial Biomass Crops for a Resource-Constrained World. Springer, Cham. https://doi.org/10.1007/978-3-319-44530-4_15

Teyssier C, Rogier O, Claverol S, Gautier F, Lelu-Walter M-A, Duruflé H. 2023. Comprehensive Organ-Specific Profiling of Douglas Fir (Pseudotsuga menziesii) Proteome. *Biomolecules*, **13**, 1400. https://doi.org/10.3390/biom13091400

Vanholme B, Cesarino I, Goeminne G, Kim H, Marroni F, Van Acker R, Vanholme R, Morreel K, Ivens B, Pinosio S, et al. 2013. Breeding with rare defective alleles (BRDA): a natural Populus nigra HCT mutant with modified lignin as a case study. *New Phytologist*, **198**, 765–776. https://doi.org/10.1111/nph.12179

VanRaden PM. 2008. Efficient Methods to Compute Genomic Predictions. Journal of Dairy Science 91: 4414–4423. https://doi.org/10.3168/jds.2007-0980

Viger M, Smith HK, Cohen D, Dewoody J, Trewin H, Steenackers M, Bastien C, Taylor G. 2016. Adaptive mechanisms and genomic plasticity for drought tolerance identified in European

black poplar (Populus nigra L.). *Tree Physiology*, **36**, 909–928. https://doi.org/10.1093/treephys/tpw017

Wade AR, Duruflé H, Sanchez L, Segura V. 2022. eQTLs are key players in the integration of genomic and transcriptomic data for phenotype prediction. *BMC Genomics*, **23**, 2021.09.07.459279. https://doi.org/10.1186/s12864-022-08690-7

Wang J, Ding J, Tan B, Robinson KM, Michelson IH, Johansson A, Nystedt B, Scofield DG, Nilsson O, Jansson S, et al. 2018. A major locus controls local adaptation and adaptive life history variation in a perennial plant. *Genome Biology*, **19**, 1–17. https://doi.org/10.1186/s13059-018-1444-y

Weir BS, Cockerham CC. 1984. Estimating F-Statistics for the Analysis of Population Structure. *Evolution*, **38**, 1358. https://doi.org/10.2307/2408641

de Wit P, Pespeni MH, Palumbi SR. 2015. SNP genotyping and population genomics from expressed sequences – current advances and future possibilities. *Molecular Ecology*, **24**, 2310–2323. https://doi.org/10.1111/mec.13165

Wójkiewicz B, Lewandowski A, Żukowska WB, Litkowiec M, Wachowiak W. 2021. Low effective population size and high spatial genetic structure of black poplar populations from the Oder valley in Poland. *Annals of Forest Science*, **78**, 1–24. https://doi.org/10.1007/s13595-021-01055-2

Xie M, Zhang J, Tschaplinski TJ, Tuskan GA, Chen JG, Muchero W. 2018. Regulation of lignin biosynthesis and its role in growth-defense tradeoffs. *Frontiers in Plant Science*, **9**, 407961. https://doi.org/10.3389/fpls.2018.01427

Yu J, Pressoir G, Briggs WH, Bi IV, Yamasaki M, Doebley JF, McMullen MD, Gaut BS, Nielsen DM, Holland JB, et al. 2005. A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. *Nature Genetics*, **38**, 203–208. https://doi.org/10.1038/ng1702

Zhang M, Suren H, Holliday JA. 2019. Phenotypic and Genomic Local Adaptation across Latitude and Altitude in Populus trichocarpa. *Genome Biology and Evolution*, **11**, 2256–2272. https://doi.org/10.1093/gbe/evz151

Zhou Y. 2022. Understanding urban plant phenology for sustainable cities and planet. *Nature Climate Change,* **12,** 302–304. https://doi.org/10.1038/s41558-022-01331-7