Disentangling heritability and plasticity effects on *Populus fremontii* leaf reflectance across a temperature gradient

Seeley, M. M.[1*†]; Thomson, E.[2]*; Allan, G. J[3]; Wiebe, B. C.[4]; Whitham, T. G.[3]; Hultine, K. R.[5]; Cooper, H. F.[3,6]; Asner, G. P.[1]; Doughty, C.E.[4]

1. Center for Global Discovery and Conservation Science, Arizona State University, Hilo, HI 96706
2. School of Geography & the Environment, University of Oxford, Oxford, UK
3. Department of Biological Sciences and Center for Adaptable Western Landscapes, Northern Arizona University, Flagstaff, AZ 86011, USA
4. School of Informatics, Computing, and Cyber Systems, Northern Arizona University, Flagstaff, AZ, USA
5. Department of Research, Conservation and Collections, Desert Botanical Garden, 1201 North Galvin Parkway, Phoenix, AZ 85008, USA
6. Center for Adaptable Western Landscapes, Northern Arizona University, Flagstaff, AZ 86011, USA

*Equal first authors

†Corresponding author; mseeley1@asu.edu

ORCID iDs

Megan M. Seeley https://orcid.org/0000-0003-1945-1162

Eleanor Thomson https://orcid.org/0000-0003-1670-8970

Gerard J. Allan https://orcid.org/0000-0002-8007-4784

Benjamin C. Wiebe https://orcid.org/0000-0002-9325-1540

Thomas G. Whitham https://orcid.org/0000-0002-1262-4342

Kevin Hultine https://orcid.org/0000-0001-9747-6037

Hillary F. Cooper https://orcid.org/0000-0003-2634-1404

Gregory P. Asner https://orcid.org/0000-0001-7893-6421

Christopher E. Doughty https://orcid.org/0000-0003-3985-7960

Keywords

Populus fremontii, cottonwood, spectroscopy, hyperspectral, plasticity, heritability, genotypic

variation, extreme temperatures; forest; tree; heat

Abstract

Globally, vegetation biodiversity is expected to decline as the rate of plant adaptation struggles

to keep pace with rising temperatures. To support conservation efforts through remote sensing,

we disentangled the nested effects of genetic and environmental influences on reflectance

spectra, leveraging spectroscopy to assess plant adaptations to temperature. Specifically, we

quantified the relative effect of plasticity and heritability on *Populus fremontii* (Fremont

cottonwood) leaf reflectance using clonal replicates propagated from 16 populations and grown

across three common gardens spanning a mean annual temperature gradient representing the

thermal range of *P. fremontii*. We used variance partitioning to decompose phenotypic variation

expressed in the leaf spectra into genotypic and environmental components to estimate broad-

sense heritability. Heritability was strongly expressed in the spectral red edge (~680-750nm) and

shortwave infrared (~1400-3000nm), though the heritability peak in the red edge was sensitive to

extreme temperatures. By comparing distances of group centroids in principal component space,

we determined that *P. fremontii* intraspecific spectral variation was shaped by the interaction

between common garden site conditions and source population. Support vector machine models

indicated pronounced environmental influence on spectral variation, as *P. fremontii* source

population and garden location were classified at 71.8% and 92.6% accuracy, respectively. These

findings emphasize the utility of reflectance data in separating genetic and environmental

influences on plant phenotypes, offering a pathway to scale these insights across broader

landscapes and aid in the conservation and management of vulnerable ecosystems in a warming

climate.

Introduction

Globally, forests are threatened by climate change through increasing temperatures and drought, leading to widespread tree mortalities (Allen et al., 2010). Many regions have already experienced declines in biodiversity, and these declines are predicted to accelerate (Pimm et al., 2014). Remote sensing, specifically imaging spectroscopy, has emerged as a powerful tool for addressing these challenges, offering insights into biodiversity, species distributions, and ecosystem health at the landscape scale (Rose et al., 2015; Seeley & Asner, 2021; Turner et al., 2003). For example, forest conservation applications of imaging spectroscopy include accurate, high-resolution species mapping (e.g. Balzotti et al., 2020; Seeley et al., 2023c), large-scale biodiversity estimations to inform and support protected area establishment (Asner et al., 2017; Féret & Asner, 2014), and forest physiological and chemical monitoring (Brodrick & Asner, 2017; Roberts et al., 2003; Weingarten et al., 2021). Imaging spectroscopy can also be used to explore plastic responses to changing environmental conditions and heritable, intraspecific trait variation that enables species to adapt to changing climates and novel conditions (Czyż et al., 2020; Seeley & Asner, 2023; Wang et al., 2022).

Previous research has successfully linked leaf and canopy spectra to both environmental and genetic factors. For example, visible through shortwave-infrared (VSWIR) reflectance data have been used to track environmental responses, such as changes in leaf traits (e.g. leaf mass per area, chlorophyll) due to stress or climatic variation (Czyż et al., 2020; Seeley & Asner, 2023). It has also been used to distinguish individuals within a species based on their genotype (Blonder et al., 2020; Cavender-Bares et al., 2016; Seeley et al., 2023b). However, many of these studies focus on either genetic or environmental influences, with limited attention to how environmental sensitivity of plant traits, and therefore spectra, can affect phylogenetic studies or

how genetic differences in plasticity influence spectral responses to environmental adaptation.

Together, these considerations have the potential to reveal insights into how plants adapt and

respond to changing conditions.

We used a common garden experiment to assess plastic and genotypically constrained

responses of leaf reflectance spectra in *Populus fremontii* (Fremont cottonwood). *P. fremontii*

inhabits riparian zones in the southwest United States and Mexico and is recognized as one of the

most important foundation species in these regions (Ellison et al., 2005; Whitham et al., 2006)

due to its ability to structure communities across multiple trophic levels, drive ecosystem

processes, and influence biodiversity via genotypically based functional trait variation (Ferrier et

al., 2012; Grady et al., 2013). Despite active planting projects, the areal extent of *P. fremontii* has

declined dramatically over the last century due to water extraction, competition with invasive

species, and climate change (Braatne et al., 1996; Shafroth et al., 1995; Stromberg, 1998). To

support management decisions for *P. fremontii* and protect riparian ecosystems vulnerable to

climate change, we attempted to link leaf-level spectral data to temperature adaptations. By

deconstructing VSWIR spectra into heritable and plastic components, we aimed to map and

monitor *P. fremontii* responses and tolerance to changing climate conditions and to derive a

method for quantifying the heritable resilience (intraspecific variation in plant traits that could

result in greater ecosystem resilience; Turner, 2023; Vahsen et al., 2023) of *P. fremontii*

populations to extreme temperatures at large spatial scales.

Heritability is a statistic that estimates the degree of variation in a phenotypic trait that is

due to genetic variation between individuals of a population. Measuring heritability in plant traits

has long been a practice in selective breeding programs searching for heritable traits to maximize

yield and quality (Bradshaw, 2017). Broad-sense heritability is defined as $H^2 = v(G)/v(E)$, where

v(G) is the proportion of genetic variance and v(E) is the proportion of environmental variation,

typically quantified using clonal replicates (Hühn, 1975). Thus, broad-sense heritability can vary

according to the amount of environmental variation experienced by individuals. As spectroscopy

integrates a vast array of leaf traits in its analyses and is a better predictor of plant and ecosystem

function than a subset of foliar traits (Yan et al., 2021), quantifying heritability across the

reflectance spectrum may yield novel insights into genetic versus environmental controls on leaf

spectra, which are linked to functional traits, and reveal parts of the spectrum useful for

retrieving phylogenetic information.

By examining *P. fremontii* leaf reflectance across three common gardens spanning the

thermal range of *P. fremontii*, we sought to disentangle the genetic and environmental factors

shaping its adaptive capacity to varying temperatures. Specifically, our objectives were to: 1)

quantify the relative contributions of genetics (source population) and environmental influences

(common garden site) to intraspecific spectral variation, 2) explore adaptations to environmental

extremes by analyzing reflectance spectra, with a focus on phenotypic plasticity and genetic

differentiation under extreme temperatures, and 3) identify conserved and variable regions across

the VSWIR spectra of *P. fremontii* across the common gardens. Where we do not specifically

assess genetic variation in spectral plasticity, we simplified our general discussion of plasticity

here as the degree to which leaf spectra change across different environments, acknowledging

that plasticity has a genetic component (Cooper et al., 2019; Laitinen & Nikoloski, 2019;

Scheiner, 1993).

Materials and Methods

Study Sites

Three common garden sites were established across the thermal range of *P. fremontii* in

2014. Two were located at the thermal edges of the *P. fremontii* temperature range (Yuma,

Arizona: 22.8°C and Canyonlands, Utah: 10.7°C; Figure S1), and one garden was in the middle

of this range (Agua Fria, Arizona: 17.2°C; Figure 1; Table 1). As Yuma experiences extreme

temperatures where ambient temperatures could exceed the critical temperature at which

Photosystem II is disrupted (t_{crit}) (Moran et al., 2023), we were able to assess the responses of

leaf spectra when trees experienced air temperatures approaching and surpassing their critical

leaf temperature thresholds (Hultine et al., 2020; Moran et al., 2023). Given the large

temperature gradients across the three common garden sites, temperature was expected to be the

primary factor influencing physiological processes (Figure 2; Cooper et al., 2019, 2022).

Each garden hosted cuttings from sixteen geographically separate (minimum 13.5 km) *P.*

fremontii populations across Arizona spanning 12.2ºC Mean Annual Temperature. From each

cutting, clonal replicates were propagated in a greenhouse. 32 clonal replicates of each cutting

were planted in each common garden, across 4 stratified random blocks to minimize the effect of

within-garden variation on a population. Each common garden hosted 16 wild populations of *P.*

fremontii, eight genotypes (cuttings) from each population (each located >20 m away), and 32

clonal replicates from each genotype, resulting in 4096 *P. fremontii* saplings per garden. Trees at

the common gardens were routinely watered to facilitate establishment in the arid southwest. A

detailed description of the common garden establishment can be found in Cooper et al. (2019,

2022).

We note that, while all environmental differences between common garden sites are

included in the variance explained by site, we assume that the primary factor affecting

individuals at each common garden site is temperature following Cooper et al. (2019, 2022).

P. fremontii in the common gardens represented two distinct ecotypes, the Colorado

Plateau and Sonoran Desert ecotypes (Blasini et al., 2021; Bothwell et al., 2023; Ikeda et al.,

2017). *P. fremontii* were collected from sixteen distinct source populations (Table 1), three of

which were located on the Mogollon Rim and are part of the Colorado Plateau ecotype (also

referred to as the Utah High Plateau/Mogollon Rim ecotype; Blasini et al., 2021; Ikeda et al.,

2017). The remaining thirteen source populations were part of the Sonoran Desert ecotype.

Within the Sonoran Desert ecotype, three sub-ecotypes (northern and southern) were represented

(Bothwell et al., 2023). The larger, northern sub-ecotype includes nine source populations and is

referred to as the main Sonoran Desert cluster, while the southeast (SE) Sonoran Desert sub-

ecotype includes three populations in the southeastern portion of Arizona. Lastly, one population

(KWF) did not fall into either Sonoran Desert sub-ecotype.

Leaf Measurements

We measured leaf reflectance spectra from trees at the three common gardens in May

2018, four years after the saplings had been planted. Due to the differential survivorship of the

saplings, the number of trees sampled from each population at each garden varied (0-52; Table

S1). Three mature leaves were collected from each tree, and, where possible, leaves were

harvested from different branches at the same leaf position on the shoot, preferentially just below

the new growth, to standardize for development. Only fully expanded, sunlit adult leaves were

chosen. Leaves were harvested by hand and transported in a cool, dark box to the nearby field

station for processing the same day.

Spectral measurements were collected using an Analytical Spectra Devices (ASD) leaf

clip, plant probe, and field spectrometer from 350 to 2500 nm at 1-nm intervals (Analytical

Spectra Devices Inc., Boulder, CO, USA). Three measurements were taken on the adaxial

surface of each leaf in the center and averaged. The spectrometer was optimized using a

Spectralon white reference taken after every three leaves. To minimize noise, wavelengths with a

low signal-to-noise ratio were removed (<400 nm and >2450 nm). Brightness normalization was

applied to all spectral measurements to minimize variation not related to leaf chemistry and

structure (Kruse et al., 1993; Myneni et al., 1989).

Leaf trait measurements and analyses

Following spectral measurements, leaves were scanned using a digital scanner (Canon

LiDE 110) to determine their area, oven dried at 70 ∘C for three days and weighed for their dry

mass. Leaf mass per area (LMA) was calculated by dividing the leaf's dry mass (g) by its fresh

one-sided area (m^2).

We next calculated solar-weighted reflectance, or albedo, for the visible (400-700nm)

and near-infrared (NIR)-shortwave infrared (SWIR; 700-2500nm). Reflectance spectra (not

brightness normalized) were weighted by the solar irradiance spectrum (ASTM G173-03

Reference Spectra derived from SMARTS version 2.9; Gueymard, 2004) to calculate broadband

albedo (Bartlett et al., 2011) according to Equation 2, where λ represents each wavelength:

Equation 1: $$Albedo \approx \frac{\sum_{i=min}^{max} Reflectance\,(\lambda_i) \cdot Incident\,Irradiance\,(\lambda_i) \cdot \delta\lambda}{\sum_{i=min}^{max} Incident\,Irradiance\,(\lambda_i) \cdot \delta\lambda}$$

Lastly, we estimated chlorophyll content using the chlorophyll index – green (CIG)

according to Equation 2.

$$\text{Equation 2: } CIG = \frac{\rho_{800}}{\rho_{550}} - 1$$

ρ_{800} is the reflectance at 800 nm (in the NIR) and ρ_{500} is that at 550 nm (green). As

chlorophyll absorbs green light, higher CIG values indicate greater leaf chlorophyll content. To

determine whether the common garden sites induced changes in leaf traits, a Tukey-Kramer

multiple comparison test from the scipy python package (Virtanen et al., 2020) was applied to

the albedo, LMA, and CIG.

Phenotypic variation in leaf spectra

To explore phenotypic variation in leaf spectra, we first visualized spectral variation

across the three common gardens. Mean source population spectra at each common garden site

were plotted and colored according to site. Spectral differences between the same populations at

different common garden sites were further visualized using principal component analysis

(PCA). PCA reduces the dimensionality of the spectral data from over 2000 dimensions to a

handful of dimensions that explain the primary spectral variation across multiple channels

(Boardman & Green, 2000; Green & Boardman, 2000; Thompson et al., 2017). We randomly

selected 104 spectra from among the five populations that were represented in all three common

gardens (CLF, KWF, BCE, PSA, and SCT). Using scikit-learn (Pedregosa et al., 2011), we

applied PCA to spectra data, plotting the first two principal components (PC) along the x and y

axes. As higher PC are often important in spectroscopy applications for discriminating

differences in surface chemistry and other traits (Dai et al., 2022; Thompson et al., 2017), we

further visualized spectral differences across ecotypes and common gardens in the first 20 PCs.

As the effect of extreme heat at Yuma was obvious in the first two PCs (Figure 4), this second

analysis included spectral data from only Agua Fria and Canyonlands. Data were grouped by

ecotype (Colorado Plateau, Sonoran Desert Main, and Sonoran Desert Southeast (SE)) and

garden, randomly selecting spectra from 40 trees from each ecotype-garden combination. We

then calculated the centroids (group mean) of each ecotype-garden combination in the first 20

PCs and measured the Euclidean distances between these centroids. Distances were then

projected into 2D space using multidimensional scaling (MDS), following the approach of

Seeley et al. (2023c).

Decomposing genotypic and environmental variation across the spectra and estimating broad-

sense heritability

To quantify the relative effects of environmental versus genotypic variation across the

VSWIR, we used variance partitioning to decompose observed variance at each wavelength into

the relative contribution of site, ecotype, population, genotype, or clone. All sampled data was

used across all common garden sites. Following the method of Asner et al. (2014) and Fyllas et

al. (2009), variance partitioning was calculated using a linear mixed-effects model for each

wavelength across the VSWIR using the lme4 *R* package (Bates et al., 2015). The model was

fitted using Residual Maximum Likelihood (REML). All effects were treated as random.

Common garden site was included as a crossed random effect, while clones were nested within

genotypes, which were nested within populations, which were nested within ecotype. The total

spectral variance about the mean for a given wavelength was, therefore, quantitatively parsed

into the variance explained by site ($\sigma 2_s$), ecotype ($\sigma 2_e$), populations within ecotype ($\sigma 2_p$),

genotypes within populations ($\sigma 2_g$), clones within genotypes ($\sigma 2_c$) and residual variance ($\sigma 2_r$)

according to Equation 3:

242   Equation 3: $\sigma 2_{total} = \sigma 2_s + \sigma 2_e + \sigma 2_p + \sigma 2_g + \sigma 2_c + \sigma 2_r$

We next calculated broad-sense heritability across the VSWIR. Heritability at each

wavelength was calculated using the classic broad sense heritability equation ($H^2 = V_G/V_P$),

where V_G is genotypic variation, and V_P is total variation ($V_{G+}V_E$). Variance explained by

ecotypes, populations, and genotypes is categorized as genotypic variance (V_G), as the genetic

material within these groups differs. Variance explained by site and clones, as well as residual

variance, is categorized as environmental variation (V_E), or spectral plasticity, as the genetic

material within these groups is identical, so any observed differences in the spectra must be due

to environment or measurement error. Broad-sense heritability was plotted across the VSWIR

spectrum. This analysis was applied first to data from all three sites and then to data from

Canyonlands and Agua Fria alone. To obtain confidence intervals for H^2, we employed a

nonparametric bootstrap approach. Bootstrapping, a method commonly used for obtaining

uncertainty estimates for heritability (Schweiger et al., 2016), involves resampling the dataset

with replacement and recalculating the variance with each subsampled dataset. We performed

1000 iterations for each wavelength, sampling 31 spectra from each common garden x ecotype

combination, resulting in n=279 for each iteration. Mean H^2 of the 1000 iterations were

presented along with the 95% confidence interval calculated from the bootstrapped estimates.

Using leaf spectra to predict genotypic and environmental information

To explore whether leaf spectra could be used to extract both genotypic and

environmental information, we used a support vector machine (SVM) to predict *P. fremontii*

source population (genotypic variation) and garden location (environmental variation). SVMs

efficiently handle highly dimensional datasets by creating decision boundaries in n-dimensional

space that maximize the distance between class boundaries and sample points (Melgani &

Bruzzone, 2004). SVMs tend to outperform other machine learning algorithms such as random

forest for species and genotype classifications when applied to spectroscopy data (e.g. Dalponte

et al., 2012). For each radial basis function kernel SVM, we optimized hyperparameter selection

through a cross-validation grid search implemented in python with the scikit-learn package

(Pedregosa et al., 2011). As with the PCA analyses, 104 samples from each garden were

randomly selected from the same five source populations (CLF, KWF, BCE, PSA, and SCT).

For all models, data were divided into training/test datasets using a 70/30 stratified random split.

To assess the generalizability of models in unseen and extreme environments in

predicting genotypic information, we trained three additional SVM models. Each model was

trained to predict the source population using *P. fremontii* spectra from two of the gardens (n =

208) and tested on unseen spectra from the third garden (n = 104). We repeated this experiment

three times, with a different garden withheld each time. This allowed us to compare the

applicability of the model in extreme environments and assess how well the model can

extrapolate outside of its training data.

Results

VSWIR leaf spectra responded to common garden treatment, particularly at Yuma, which

represents the upper end of the cottonwood thermal range. Brightness-normalized reflectance of

all populations at the Yuma site diverged from that at both Canyonlands and Agua Fria (Figures

2a & 3a). Albedo in both the visible and NIR-SWIR differed across the common gardens

(p<0.05) and was highest at Yuma. In the visible, albedo was lowest at Canyonlands, and in the

NIR-SWIR, albedo was lowest at Agua Fria. CIG, indicative of chlorophyll content, was highest

in Canyonlands (mean = 4.8) and Agua Fria (mean = 4.7). Yuma had the lowest CIG, with a

mean of 2.8 (p<0.05).

LMA, an important driver of spectral variation, was highest in the Yuma common garden

and lowest in Canyonlands (p<0.05; Figure 4b). Mean LMA across all populations at

Canyonlands was 72.9 g/m^2 with a standard deviation of 10.2 g/m^2. At Agua Fria, the mean

LMA was 88.1 g/m^2, though it was the most variable with a standard deviation of 19.1 g/m^2.

Yuma exhibited the highest LMA at 108.4 g/m^2 and a standard deviation of 11.2 g/m^2. In

addition to differing across the gardens, LMA also varied among source populations; a two-way

ANOVA revealed significant main effects of both garden environment and population on LMA

(p<0.05 for both factors).

In PC space, spectra differed across the common gardens and among the ecotypes (Figure

4). The first two PC of spectra from all common gardens explained 57.8 and 35.3% of the

variance, respectively. These components primarily separated Yuma spectra from those at

Canyonlands and Agua Fria (Figure 4a). When comparing spectra from different ecotypes at

Canyonlands and Agua Fria, the Colorado Plateau ecotype separated from both Sonoran Desert

ecotypes regardless of common garden in the first 20 PCs. Within the Sonoran Desert ecotypes

(Main and SE), common garden was the primary driver of spectral divergence in PC space.

Variance partitioning and broad-sense heritability

Across all gardens, spectral variance was highest in the visible (~400-680 nm) and red-

edge wavelengths (~680-750 nm). When Yuma was included in the analysis, variance in these

wavelengths was driven largely by the common garden site (Figures 2 and 3), and heritability in

these regions was correspondingly low with a maximum heritability of 0.08 for the visible

(Figure 5). The red-edge heritability peak present when Yuma was excluded collapsed with the

inclusion of Yuma, leaving a smaller heritability peak at the longer wavelengths. Additionally,

site-level variation in the visible (~400-680 nm) and red-edge wavelengths (~680-750 nm) was

minimal without Yuma. Instead, variance at these wavelengths was driven more by genetic

factors (population and genotype), with corresponding high spectral heritability (~0.36 and

~0.66, respectively).

Across all gardens, heritability in *P. fremontii* reflectance was greatest in the SWIR

(1400 – 2450 nm) and the transition from red-edge to NIR (~750 nm), regardless of the inclusion

of Yuma (Figure 5). In the shorter wavelengths of the NIR (~750-1140 nm) and SWIR, site-level

variation explained less of the variance than genotypic factors in general, and the spectral

differences caused by Yuma were relatively minimal, resulting in high heritability (max: 0.51;

min: 0.31). When Yuma was excluded, heritability peaked in the red-edge (740 nm; heritability:

0.67). Heritability had a negative relationship with wavelength in the NIR until ~1320 nm where

heritability increased and remained high in the SWIR, although it varied across this region (max:

~0.56, min: ~0.37). Across the VSWIR, particularly heritable regions of the spectra included the

red-edge/NIR transition (~750 nm) and the SWIR. More environmentally sensitive (plastic)

regions included the visible and red-edge as well as much of the NIR.

SVM analysis and leaf predictions

SVM models were able to discriminate between spectra based on both common garden

site and source population at 92.6% and 71.8% accuracy, respectively (Figure 6). The SVM

trained to classify spectra based on the common garden site (environmental variation)

outperformed the SVM trained to discriminate spectra based on source population (genotypic

variation). The common garden site SVM accurately predicted spectra from the Yuma site 100%

of the time; confusion in this model occurred primarily between Canyonlands and Agua Fria,

with the true positive rate for Canyonlands and Agua Fria being 94% and 84%, respectively. For

the source population SVM, the true positive rate ranged from 61% for the KWF population to

84% for CLF. When predicting source population using one common garden site as a holdout for

testing, model performance declined (Figure S4). Source populations at Canyonlands were

predicted with 38.5% accuracy when trained on data from Agua Fria and Yuma. At Agua Fria

and Yuma, accuracy was 37.5% and 31.7%, respectively, when SVM models were trained on

data from the other common garden sites.

Discussion

High temperatures (>50°C) induced physiological stress in *P. fremontii,* altering their

spectral signatures. The environmental variability introduced by Yuma, the hottest common

garden, resulted in lower heritability in the visible and red edge regions of the reflectance

spectra, discussed further below. We confirmed that trees growing at Yuma experienced thermal

stress, as indicated by the environmental data and lower CIG. Moran et al. (2023) documented

temporal changes in leaf thermal tolerance (t_{crit}) at Yuma, confirming that *P. fremontii* undergoes

physiological adjustments in response to exposure to extreme temperatures. Further, Sankey et

al. (2020) demonstrated GxE interactions among *P. fremontii* at the Agua Fria common garden,

observing differences in canopy temperature between genotypes using thermal imagery from

unmanned aerial vehicles. Plastic responses at the individual level can include increased LMA

(observed at Yuma), increased heat shock proteins (HSP) to help repair heat damage,

concentrations of which have been found to correlate with LMA (Knight & Ackerly, 2003),

trichome development to promote self-shading (Banowetz et al., 2008), an accumulation of

phenolics and isoprenoids for their ability to protect photosynthetic apparatuses and antioxidant

properties (Bartwal et al., 2013; Smirnoff, 2005), and increased production of soluble sugars

(Wahid & Close, 2007) and saturated fatty acids (Ahmad & Prasad, 2011; Tang et al., 2016). In

addition to the physiological changes, Cooper et al. (2019) identified phenological (e.g. bud set,

bud flush) differences between clones located at different gardens; for example, bud flush

occurred earlier at the Yuma garden.

As many of the physical and chemical changes such as increased LMA, plant phenols,

soluble sugars, and fatty acids can be estimated using spectroscopy (Akkaya, 2018; Asner &

Martin, 2009, 2016), it is unsurprising that *P. fremontii* at Yuma differed spectrally from those at

the other sites. For *Panax ginseng*, differences between VSWIR reflectance of heat-stressed and

control plants corresponded to spectral regions linked to fatty acids, sugars, protein, chlorophyll,

and nitrogen (Park et al., 2021). Similarly, other studies have found differences between control

and heat-stressed plants using fluorescence (Faqeerzada et al., 2023) and visible-NIR (Kim et al.,

2022) spectroscopy. While plastic responses to heat stress may decrease spectral heritability,

environmentally sensitive regions of the spectra offer opportunities to understand physiological

responses to extreme temperatures at large scales and may serve as environmental indicators of

stress events. Yuma is near the southern edge of the distribution of *P. fremontii* where it may not

survive future climatic conditions (e.g., Blasini et al., 2021; Moran et al., 2023), so identifying

environmental indicators of extreme environmental conditions may allow us to monitor stress at

the shifting range edge of *P. fremontii.*

Visible and Red Edge (400-750nm)

The visible and red-edge regions demonstrated the greatest sensitivity to extreme temperatures. Without Yuma, heritability estimates in the visible red-edge regions were higher, suggesting that genetic factors played a stronger role in explaining trait variation under non-stress conditions. Heritability in the visible and red edge has been documented across many species (Čepl et al., 2018; Czyż et al., 2020; Feng et al., 2017; McManus et al., 2016). For example, *Metrosideros polymorpha* varieties that organize along environmental gradients diverged in chlorophyll content, which was associated with differences in the visible wavelengths (Seeley et al. 2023b). Feng et al. (2017) found that the red edge was predictive of rice genotypes, and both Čepal et al. (2018) and Czyż et al. (2020) observed a red edge signal in the heritability of *Pinus sylvestris* (Scotts pine) and *Fagus sylvatica* (European beech) leaf reflectance, respectively. In heat-stressed conditions, however, increased physiological plasticity, such as changes in chlorophyll concentration, can reduce the expression of genetic differences (Posch et al., 2024; Seeley et al., 2025b; Stone et al., 2018), thereby lowering heritability.

The inclusion of Yuma, characterized by elevated temperatures, likely introduced greater environmental variance and potentially GxE interactions, altering the proportion of phenotypic variance attributable to genetics. Although we did not directly measure chlorophyll concentration, both the visible and red-edge regions are closely associated with chlorophyll and other photosynthetic pigments (Horler et al., 1983; Vogelmann et al., 1993). Further, the CIG indicated that chlorophyll content at Yuma was lower than that at the other gardens. Heat stress can lead to disruption of the photosystems, characterized by chlorophyll degradation and changes in pigment ratios (Ivanov et al., 2017; Mathur et al., 2014; Tafesse et al., 2022), which can

reduce absorption in the visible range and shift the red-edge position (Tirado et al., 2021; Xie et

al., 2019). Reflectance of trees at Yuma were consistent with these processes, as they exhibited

higher visible reflectance, similar to that reported for heat-stressed rice (Xie et al., 2019) and

maize (Tirado et al., 2021). The change in reflectance suggests physiological changes that both

serve as a mechanism of photosynthetic apparatus protection and as are a side effect of the stress

response.

Near-Infrared (750-1400nm)

Heritability across the NIR declined with increasing wavelength in *P. fremontii* and was

slightly lower with the inclusion of Yuma. NIR reflectance is primarily influenced by the internal

structure of leaves, particularly the spongy mesophyll (Gausman, 1974). LMA, which influences

NIR reflectance (Asner et al., 2011; Gara et al., 2021; Serbin et al., 2019), increased across the

common gardens and varied among source populations, indicating that both genetic and

environmental factors affected LMA. Knight and Ackerly (2003) similarly found elevated LMA

under thermal stress in desert plants, linking it to enhanced photosynthetic recovery and

identifying high LMA as a convergent trait in plants adapted to hot, arid environments, like *P.*

fremontii. The increase in LMA in response to thermal stress at Yuma may have contributed to

the decline in heritability in the NIR, though this effect was less pronounced than in the visible

region. Heritability in the NIR remained relatively high, however, perhaps because other traits

that could further reduce heritability, such as leaf water content (Gausman, 1974; Seeley et al.,

2025b), were controlled across the gardens. The heritability peaks we observed in the NIR are

consistent with those reported in *Pinus sylvestris* and *Fagus sylvatica* (Čepl et al., 2018; Czyż et

al., 2020), though *P. sylvestris* showed an additional region of high heritability between 1130–

1330 nm that corresponded to low heritability in *P. fremontii*. Given that much of the NIR signal

reflects internal leaf structure (Knipling, 1970; Slaton et al., 2001), these interspecific differences

may arise from contrasting leaf morphologies—broadleaf vs. needleleaf.

Shortwave-infrared (1400-2500nm)

In the SWIR, heritability remained relatively high regardless of the inclusion of the hotter

Yuma garden, suggesting that genetic control over traits expressed in the SWIR was robust to

environmental variation. The SWIR region is associated with structural and biochemical leaf

components such as lignin, phenols, and cellulose (Asner & Martin, 2016), which are

influenced by both species-level differences and intra-specific genetic variation (Asner & Martin,

2009; Seeley et al., 2023). These traits are often under stronger genetic regulation and/or less

plastic in response to temperature stress compared to traits captured in other spectral regions

(Novaes et al., 2010), likely explaining the stability of heritability in this SWIR. Previous studies

have likewise identified wavelengths in the SWIR as having high phylogenetic signal (Czyż et

al., 2020; McManus et al. 2016; Meireles et al. 2020).

While reflectance in the SWIR has a strong genetic component, this region is also

important for separating heat-stressed and control plants, as noted by Park et al. (2021). This is

because temperature-dependent leaf traits (e.g. ribulose bisphosphate carboxylation rate) affect

reflectance in the SWIR (Park et al., 2021; Serbin et al., 2012), as was demonstrated by Serbin et

al. (2012) in two *Populus* species, *P. tremuloides* and *P. deltoides* (Serbin et al., 2012). As LMA

is also expressed in the SWIR (Asner et al., 2011; Gara et al., 2021; Serbin et al., 2019), this trait

may further contribute to the sensitivity of the SWIR to heat stress. Further, SWIR-associated

traits can span a broad heritability range; for example, some phenolic compounds exhibit strong

genetic determination, whereas others show considerable plasticity in response to environmental

stressors such as heat (Klaper et al., 2001; Rivero et al., 2001). In *P. fremontii,* site-level

variation between Agua Fria and Canyonlands was primarily observed in the SWIR,

emphasizing the importance of using the full spectrum to distinguish between temperature

treatments.

Albedo

Across the entire VSWIR spectra, albedo increased in response to extreme temperature,

likely lowering canopy temperatures and helping mitigate further heat stress. In the NIR – SWIR,

albedo decreased from Canyonlands to Agua Fria, supporting the results from Doughty et al.

(2018), where albedo declined across a temperature gradient in Peru. Regardless of which region

of the VSWIR was analyzed, albedo at Yuma was higher than at the other common gardens.

While additional evidence is required to confirm whether the increase in leaf albedo is a

universal plant response to extremely high temperatures, this phenomenon aligns with plant

ecophysiological principles; many stress factors, including heat, as was demonstrated here, are

known to decrease chlorophyll concentration (Carter & Knapp, 2001).

Heritability and plasticity of reflectance spectra

Intraspecific variation in *P. fremontii* reflectance spectra was driven by GxE interactions,

with extreme temperatures at the Yuma site playing a dominant role. Ambient temperatures at

Yuma, which approach the upper thermal tolerance limit of *P. fremontii* (Moran et al., 2023),

likely caused a threshold change in spectral expression and heritability, as evidenced by the

heritability estimates, discussed above, and PCA analyses. In the two PCA scenarios, with and

without Yuma, spectra separated according to garden and ecotype, respectively, demonstrating

hierarchical effects of environmental and genetic factors. Similar findings were reported by Li et

al. (2023), who observed that PC1 of *Nicotiana attenuata* spectra was primarily driven by

common garden location, while genotype structured variation along PCs 2–4. Here, the extreme

temperatures at Yuma induced leaf physiological responses (e.g. changes in chlorophyll content,

LMA) that dominated the spectral signal. This dominant effect of environmental on spectra was

further supported by our classification models: SVMs trained to predict garden site achieved

~14% higher accuracy than those trained to distinguish source populations. Cavender-Bares et al.

(2016) similarly found that water treatment was classified more accurately than population origin

for *Quercus oleoides*.

When we trained SVMs on data from two common garden sites and tested on the third,

classification accuracy for predicting *P. fremontii* source population declined by over 34%,

despite controlling for genetic identity. This decline aligns with previous studies demonstrating

that the environmental sensitivity of spectra limits large-scale species classifications (e.g., Seeley

et al., 2024; 2025a). While *in situ*, ecological mechanisms behind species classification

limitations may include distance decay (Rocchini, 2007), intraspecific spectral variation (Czyż et

al., 2020; Seeley & Asner, 2023; 2025a), and interspecific trait convergence (Rodríguez-Alarcón

et al., 2022), this common garden framework allowed us to control for genetic identity and

isolate environmental effects on classification accuracy. The sharp drop in model performance

under novel garden conditions suggested that plasticity in spectral traits, arising from site-level

environmental variation, was a key limiting factor in model transferability.

Below the physiological temperature threshold at Yuma, spectral differences were

predominantly governed by genetic factors, with environmental conditions playing a significant

role in shaping phenotypic expression within genetically similar groups. Excluding Yuma in the

PCA resulted in the source population (genotype) being the primary driver of separability,

observed by the separation of the Sonoran Desert and Colorado Plateau ecotypes in PC space.

However, within finer sub-ecotype categories, differences were primarily driven by the common

garden, reinforcing the interconnected roles of genetics and environment in shaping spectral

traits.

Conclusions

Reflectance spectra of *P. fremontii* contain valuable information about both the

environmental conditions in which they grow and their genetic identities. Intraspecific spectral

variation of *P. fremontii* was shaped by both common garden site and source population, with

environmental conditions playing a dominant role under extreme stress, such as the high

temperatures at the Yuma site. At the upper thermal threshold of *P. fremontii*, the spectral

response was most pronounced in the visible and red-edge regions, corresponding to documented

heat stress responses. In these regions, spectral heritability decreased when Yuma was

considered in the analysis, suggesting that extreme environmental conditions may drive a

threshold shift in spectral expression. However, genetic information was retained, and

heritability was more consistent in parts of the NIR and SWIR.

These results underscore the potential for using VSWIR reflectance data to disentangle

genetic and environmental contributions to plant phenotypes, providing a path forward for

scaling these insights to larger landscapes. As airborne and satellite-based imaging spectroscopy

data continue to become more widely available, they offer a powerful opportunity to assess

phenotypic plasticity and heritable resilience of *P. fremontii* to extreme conditions across broad

spatial scales. Spectroscopy may be a powerful tool for monitoring the adaptive capacity of this

foundation species in the face of rapid environmental change, supporting efforts to preserve and

manage at-risk ecosystems in a warming climate.

Declarations

Funding: The common gardens were initially developed under NSF Macrosystems grant DEB

2017877 awarded to GJA, TGW, and KCG. Funding for this research was provided by the NSF

MacroSystems grant DEB-1340852 awarded to GJA, TGW, and CED as well as the NSF

DEB☐1340856 awarded to KRH.

Author contributions: Original idea: CED, ET, GPA; common garden development common

gardens: GJA, TGW, KCG, KRH, and HFC; data collection: ET; statistical analysis: MMS, ET,

methodology: MMS, ET, CED, BCW; writing the manuscript: MMS, ET. All authors reviewed

several drafts and agreed with the final version.

Availability of data: All data files will be made available on the Figshare database upon

acceptance of the manuscript at DOI: 10.6084/m9.figshare.25719585.

Code availability: Code is available on GitHub and is maintained by Seeley (2025)

https://github.com/MegsSeeley/temperature_cottonwood.

Conflict of interest: The authors have declared that no competing interests exist.

References

Ahmad, P., & Prasad, M. N. V. (2011). *Environmental Adaptations and Stress Tolerance of*

Plants in the Era of Climate Change. Springer Science & Business Media.

Akkaya, M. R. (2018). Prediction of fatty acid composition of sunflower seeds by near-infrared

reflectance spectroscopy. *Journal of Food Science and Technology*, *55*(6), 2318–2325.

https://doi.org/10.1007/s13197-018-3150-x

Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., …

Cobb, N. (2010). A global overview of drought and heat-induced tree mortality reveals

emerging climate change risks for forests. *Forest Ecology and Management*, *259*(4),

660–684. https://doi.org/10.1016/j.foreco.2009.09.001

Asner, G. P., & Martin, R. E. (2009). Airborne spectranomics: Mapping canopy chemical and

taxonomic diversity in tropical forests. *Frontiers in Ecology and the Environment*, *7*(5),

269–276. https://doi.org/10.1890/070152

Asner, G. P., & Martin, R. E. (2016). Spectranomics: Emerging science and conservation

opportunities at the interface of biodiversity and remote sensing. *Global Ecology and*

Conservation, *8*, 212–219. https://doi.org/10.1016/j.gecco.2016.09.010

Asner, G. P., Martin, R. E., Tupayachi, R., Anderson, C. B., Sinca, F., Carranza-Jiménez, L., &

Martinez, P. (2014). Amazonian functional diversity from forest canopy chemical

assembly. *Proceedings of the National Academy of Sciences*, *111*(15), 5604–5609.

https://doi.org/10.1073/pnas.1401181111

Asner, G. P., Martin, R. E., Tupayachi, R., Emerson, R., Martinez, P., Sinca, F., ... & Lugo, A. E.

(2011). Taxonomy and remote sensing of leaf mass per area (LMA) in humid tropical

forests. *Ecological Applications*, *21*(1), 85-98.

Asner, G. P., Martin, R. E., Tupayachi, R., & Llactayo, W. (2017). Conservation assessment of

the Peruvian Andes and Amazon based on mapped forest functional diversity. *Biological*

Conservation, *210*, 80–88. https://doi.org/10.1016/j.biocon.2017.04.008

Balzotti, C. S., Asner, G. P., Adkins, E. D., & Parsons, E. W. (2020). Spatial drivers of

composition and connectivity across endangered tropical dry forests. *Journal of Applied*

Ecology, *57*(8), 1593–1604. https://doi.org/10.1111/1365-2664.13632

Banowetz, G. M., Azevedo, M. D., & Stout, R. (2008). Morphological adaptations of hot springs

panic grass (Dichanthelium lanigunosum var sericeum (Schmoll) to thermal stress.

Journal of Thermal Biology, *33*(2), 106–116.

https://doi.org/10.1016/j.jtherbio.2007.08.006

Bartlett, M. K., Ollinger, S. V., Hollinger, D. Y., Wicklein, H. F., & Richardson, A. D. (2011).

Canopy-scale relationships between foliar nitrogen and albedo are not observed in leaf

reflectance and transmittance within temperate deciduous tree species. *Botany*, *89*(7),

491–497. https://doi.org/10.1139/b11-037

Bartwal, A., Mall, R., Lohani, P., Guru, S. K., & Arora, S. (2013). Role of Secondary

Metabolites and Brassinosteroids in Plant Defense Against Environmental Stresses.

Journal of Plant Growth Regulation, *32*(1), 216–232. https://doi.org/10.1007/s00344-

012-9272-x

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software*, *67*, 1–48.

https://doi.org/10.18637/jss.v067.i01

Blasini, D. E., Koepke, D. F., Grady, K. C., Allan, G. J., Gehring, C. A., Whitham, T. G., …

Hultine, K. R. (2021). Adaptive trait syndromes along multiple economic spectra define

cold and warm adapted ecotypes in a widely distributed foundation tree species. *Journal*

of Ecology, *109*(3), 1298–1318. https://doi.org/10.1111/1365-2745.13557

Blonder, B., Graae, B. J., Greer, B., Haagsma, M., Helsen, K., Kapás, R. E., … Strimbeck, R.

(2020). Remote sensing of ploidy level in quaking aspen (Populus tremuloides Michx.).

Journal of Ecology, *108*(1), 175–188. https://doi.org/10.1111/1365-2745.13296

Boardman, J. W., & Green, R. O. (2000). *Exploring the spectral variability of the Earth as*

measured by AVIRIS in 1999. https://trs.jpl.nasa.gov/handle/2014/16602

Bothwell, H. M., Keith, A. R., Cooper, H. F., Hull, J. B., Andrews, L. V., Wehenkel, C., …

Allan, G. J. (2023). Microevolutionary Processes in a Foundation Tree Inform

Macrosystem Patterns of Community Biodiversity and Structure. *Forests*, *14*(5), Article

5. https://doi.org/10.3390/f14050943

Braatne, J., Rood, S., & Heilman, P. (1996). *Life history, ecology, and conservation of riparian*

cottonwoods in North America (pp. 57–85).

Bradshaw, J. E. (2017). Plant breeding: Past, present and future. *Euphytica*, *213*(3), 60.

https://doi.org/10.1007/s10681-016-1815-y

Brodrick, P. G., & Asner, G. P. (2017). Remotely sensed predictors of conifer tree mortality

during severe drought. *Environmental Research Letters*, *12*(11), 115013.

https://doi.org/10.1088/1748-9326/aa8f55

Buddenbaum, H., Rock, G., Hill, J., & Werner, W. (2015). Measuring Stress Reactions of Beech

Seedlings with PRI, Fluorescence, Temperatures and Emissivity from VNIR and Thermal

Field Imaging Spectroscopy. *European Journal of Remote Sensing*, *48*(1), 263–282.

https://doi.org/10.5721/EuJRS20154815

Carter, G. A., & Knapp, A. K. (2001). Leaf optical properties in higher plants: Linking spectral

characteristics to stress and chlorophyll concentration. *American Journal of Botany*,

88(4), 677–684. https://doi.org/10.2307/2657068

Cavender-Bares, J., Meireles, J. E., Couture, J. J., Kaproth, M. A., Kingdon, C. C., Singh, A., …

Townsend, P. A. (2016). Associations of Leaf Spectra with Genetic and Phylogenetic

Variation in Oaks: Prospects for Remote Detection of Biodiversity. *Remote Sensing*, *8*(3),

Article 3. https://doi.org/10.3390/rs8030221

Čepl, J., Stejskal, J., Lhotáková, Z., Holá, D., Korecký, J., Lstibůrek, M., … Albrechtová, J.

(2018). Heritable variation in needle spectral reflectance of Scots pine (Pinus sylvestris

L.) peaks in red edge. *Remote Sensing of Environment*, *219*, 89–98.

https://doi.org/10.1016/j.rse.2018.10.001

Cooper, H. F., Best, R. J., Andrews, L. V., Corbin, J. P. M., Garthwaite, I., Grady, K. C., …

Allan, G. J. (2022). Evidence of climate-driven selection on tree traits and trait plasticity

across the climatic range of a riparian foundation species. *Molecular Ecology*, *31*(19),

5024–5040. https://doi.org/10.1111/mec.16645

Cooper, H. F., Grady, K. C., Cowan, J. A., Best, R. J., Allan, G. J., & Whitham, T. G. (2019).

Genotypic variation in phenological plasticity: Reciprocal common gardens reveal

adaptive responses to warmer springs but not to fall frost. *Global Change Biology*, *25*(1),

187–200. https://doi.org/10.1111/gcb.14494

Czyż, E. A., Guillén Escribà, C., Wulf, H., Tedder, A., Schuman, M. C., Schneider, F. D., &

Schaepman, M. E. (2020). Intraspecific genetic variation of a Fagus sylvatica population

in a temperate forest derived from airborne imaging spectroscopy time series. *Ecology*

and Evolution, *10*(14), 7419–7430. https://doi.org/10.1002/ece3.6469

Dai, J., Vaughn, N. R., Seeley, M., Heckler, J., Thompson, D. R., & Asner, G. P. (2022).

Spectral dimensionality of imaging spectroscopy data over diverse landscapes and spatial

resolutions. *Journal of Applied Remote Sensing*, *16*(4), 044518.

https://doi.org/10.1117/1.JRS.16.044518

Dalponte, M., Bruzzone, L., & Gianelle, D. (2012). Tree species classification in the Southern

Alps based on the fusion of very high geometrical resolution multispectral/hyperspectral

images and LiDAR data. *Remote Sensing of Environment*, *123*, 258–270.

https://doi.org/10.1016/j.rse.2012.03.013

Doughty, C. E., Santos-Andrade, P. E., Shenkin, A., Goldsmith, G. R., Bentley, L. P., Blonder,

B., … Malhi, Y. (2018). Tropical forest leaves may darken in response to climate change.

Nature Ecology & Evolution, *2*(12), 1918–1924. https://doi.org/10.1038/s41559-018-

0716-y

Ellison, A. M., Bank, M. S., Clinton, B. D., Colburn, E. A., Elliott, K., Ford, C. R., … Webster,

J. R. (2005). Loss of foundation species: Consequences for the structure and dynamics of

forested ecosystems. *Frontiers in Ecology and the Environment*, *3*(9), 479–486.

https://doi.org/10.1890/1540-9295(2005)003[0479:LOFSCF]2.0.CO;2

Faqeerzada, M. A., Park, E., Kim, T., Kim, M. S., Baek, I., Joshi, R., … Cho, B.-K. (2023).

Fluorescence Hyperspectral Imaging for Early Diagnosis of Heat-Stressed Ginseng

Plants. *Applied Sciences*, *13*(1), Article 1. https://doi.org/10.3390/app13010031

Feng, H., Guo, Z., Yang, W., Huang, C., Chen, G., Fang, W., … Liu, Q. (2017). An integrated

hyperspectral imaging and genome-wide association analysis platform provides spectral

and genetic insights into the natural variation in rice. *Scientific Reports*, *7*(1), Article 1.

https://doi.org/10.1038/s41598-017-04668-8

Féret, J.-B., & Asner, G. P. (2014). Mapping tropical forest canopy diversity using high-fidelity

imaging spectroscopy. *Ecological Applications*, *24*(6), 1289–1296.

https://doi.org/10.1890/13-1824.1

Ferrier, S. M., Bangert, R. K., Hersch-Green, E. I., Bailey, J. K., Allan, G. J., & Whitham, T. G.

(2012). Unique arthropod communities on different host-plant genotypes results in

greater arthropod diversity. *Arthropod-Plant Interactions*, *6*(2), 187–195.

https://doi.org/10.1007/s11829-011-9177-9

Fyllas, N. M., Patiño, S., Baker, T. R., Bielefeld Nardoto, G., Martinelli, L. A., Quesada, C. A.,

… Lloyd, J. (2009). Basin-wide variations in foliar properties of Amazonian forest:

Phylogeny, soils and climate. *Biogeosciences*, *6*, 2677–2708. https://doi.org/10.5194/bg-

6-2677-200910.5194/bgd-6-3707-2009

Gara, T. W., Rahimzadeh-Bajgiran, P., & Darvishzadeh, R. (2021). Forest leaf mass per area

(LMA) through the eye of optical remote sensing: a review and future outlook. *Remote

sensing*, *13*(17), 3352.

Gausman, H. W. (1974). Leaf reflectance of near-infrared. *Photogrammetric Engineering*, *40*(2),

183-191.

Grady, K. C., Laughlin, D. C., Ferrier, S. M., Kolb, T. E., Hart, S. C., Allan, G. J., & Whitham,

T. G. (2013). Conservative leaf economic traits correlate with fast growth of genotypes of

a foundation riparian species near the thermal maximum extent of its geographic range.

Functional Ecology, *27*(2), 428–438. https://doi.org/10.1111/1365-2435.12060

Green, R. O., & Boardman, J. W. (2000). Exploration of the relationship between information

content and signal-to-noise ratio and spatial resolution in AVIRIS spectral data.

Spectrum, *7*(8).

Gueymard, C. A. (2004). The sun's total and spectral irradiance for solar energy applications and

solar radiation models. *Solar Energy*, *76*(4), 423–453.

https://doi.org/10.1016/j.solener.2003.08.039

Hernández-Clemente, R., Navarro-Cerrillo, R. M., Suárez, L., Morales, F., & Zarco-Tejada, P. J.

(2011). Assessing structural effects on PRI for stress detection in conifer forests. *Remote*

Sensing of Environment, *115*(9), 2360–2375. https://doi.org/10.1016/j.rse.2011.04.036

Horler, D. N. H., Dockray, M., & Barber, J. (1983). The red edge of plant leaf reflectance.

International Journal of Remote Sensing, *4*(2), 273–288.

https://doi.org/10.1080/01431168308948546

Hühn, M. (1975). Estimation of broad sense heritability in plant populations: An improved

method. *Theoretical and Applied Genetics*, *46*(2), 87–99.

https://doi.org/10.1007/BF00281647

Hultine, K. R., Allan, G. J., Blasini, D., Bothwell, H. M., Cadmus, A., Cooper, H. F., …

Whitham, T. G. (2020). Adaptive capacity in the foundation tree species Populus

fremontii: Implications for resilience to climate change and non-native species invasion

in the American Southwest. *Conservation Physiology*, *8*(1), coaa061.

https://doi.org/10.1093/conphys/coaa061

Ikeda, D. H., Max, T. L., Allan, G. J., Lau, M. K., Shuster, S. M., & Whitham, T. G. (2017).

Genetically informed ecological niche models improve climate change predictions.

Global Change Biology, *23*(1), 164–176. https://doi.org/10.1111/gcb.13470

Ivanov, A. G., Velitchkova, M. Y., Allakhverdiev, S. I., & Huner, N. P. (2017). Heat stress-

induced effects of photosystem I: an overview of structural and functional

responses. *Photosynthesis research*, *133*(1), 17-30.

Kim, G., Lee, H., Wi, S. H., & Cho, B.-K. (2022). Snapshot-Based Visible-Near Infrared

Multispectral Imaging for Early Screening of Heat Injury during Growth of Chinese

Cabbage. *Applied Sciences*, *12*(18), Article 18. https://doi.org/10.3390/app12189340

Klaper, R., Ritland, K., Mousseau, T. A., & Hunter, M. D. (2001). Heritability of phenolics in

Quercus laevis inferred using molecular markers. *Journal of Heredity*, *92*(5), 421-426.

Knight, C. A., & Ackerly, D. D. (2003). Evolution and plasticity of photosynthetic thermal

tolerance, specific leaf area and leaf size: Congeneric species from desert and coastal

environments. *New Phytologist*, *160*(2), 337–347. https://doi.org/10.1046/j.1469-

8137.2003.00880.x

Knipling, E. B. (1970). Physical and physiological basis for the reflectance of visible and near-

infrared radiation from vegetation. *Remote Sensing of Environment*, *1*(3), 155–159.

https://doi.org/10.1016/S0034-4257(70)80021-9

Kruse, F. A., Heidebrecht, K. B., Shapiro, A. T., Barloon, P. J., & Goetz, A. F. H. (1993). The

Spectral Image Processing System (SIPS) Interactive Visualization and Analysis of

Imaging Spectrometer Data. *Remote Sensing of Environment*, *44*, 145–163.

Li, C., Czyż, E. A., Halitschke, R., Baldwin, I. T., Schaepman, M. E., & Schuman, M. C. (2023).

Evaluating potential of leaf reflectance spectra to monitor plant genetic variation. *Plant*

Methods, *19*(1), 108. https://doi.org/10.1186/s13007-023-01089-9

Mathur, S., Agrawal, D., & Jajoo, A. (2014). Photosynthesis: Response to high temperature

stress. *Journal of Photochemistry and Photobiology B: Biology*, *137*, 116–126.

https://doi.org/10.1016/j.jphotobiol.2014.01.010

McManus, K. M., Asner, G. P., Martin, R. E., Dexter, K. G., Kress, W. J., & Field, C. B. (2016).

Phylogenetic Structure of Foliar Spectral Traits in Tropical Forest Canopies. *Remote*

Sensing, *8*(3), Article 3. https://doi.org/10.3390/rs8030196

Meireles, J. E., Cavender-Bares, J., Townsend, P. A., Ustin, S., Gamon, J. A., Schweiger, A. K.,

… O'Meara, B. C. (2020). Leaf reflectance spectra capture the evolutionary history of

seed plants. *New Phytologist*, *228*(2), 485–493. https://doi.org/10.1111/nph.16771

Melgani, F., & Bruzzone, L. (2004). Classification of hyperspectral remote sensing images with

support vector machines. *IEEE Transactions on Geoscience and Remote Sensing*, *42*(8),

1778–1790. IEEE Transactions on Geoscience and Remote Sensing.

https://doi.org/10.1109/TGRS.2004.831865

Moran, M. E., Aparecido, L. M. T., Koepke, D. F., Cooper, H. F., Doughty, C. E., Gehring, C.

A., … Hultine, K. R. (2023). Limits of thermal and hydrological tolerance in a foundation

tree species (Populus fremontii) in the desert southwestern United States. *New*

Phytologist, *240*(6), 2298–2311. https://doi.org/10.1111/nph.19247

Myneni, R. B., Ross, J., & Asrar, G. (1989). A review on the theory of photon transport in leaf

canopies. *Agricultural and Forest Meteorology*, *45*(1–2), 1–153.

https://doi.org/10.1016/0168-1923(89)90002-6

Novaes, E., Kirst, M., Chiang, V., Winter-Sederoff, H., & Sederoff, R. (2010). Lignin and

biomass: a negative correlation for wood formation and lignin content in trees. *Plant*

physiology, *154*(2), 555-561.

Park, E., Kim, Y.-S., Omari, M. K., Suh, H.-K., Faqeerzada, M. A., Kim, M. S., … Cho, B.-K.

(2021). High-Throughput Phenotyping Approach for the Evaluation of Heat Stress in

Korean Ginseng (Panax ginseng Meyer) Using a Hyperspectral Reflectance Image.

Sensors, *21*(16), Article 16. https://doi.org/10.3390/s21165634

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., … Duchesnay,

É. (2011). Scikit-learn: Machine Learning in Python. *Journal of Machine Learning*

Research, *12*(85), 2825–2830.

Pimm, S. L., Jenkins, C. N., Abell, R., Brooks, T. M., Gittleman, J. L., Joppa, L. N., … Sexton,

J. O. (2014). The biodiversity of species and their rates of extinction, distribution, and

protection. *Science*, *344*(6187), 1246752. https://doi.org/10.1126/science.1246752

Rivero, R. M., Ruiz, J. M., Garcıa, P. C., Lopez-Lefebre, L. R., Sánchez, E., & Romero, L.

(2001). Resistance to cold and heat stress: accumulation of phenolic compounds in

tomato and watermelon plants. *Plant science*, *160*(2), 315-321.

Roberts, D. A., Dennison, P. E., Gardner, M. E., Hetzel, Y., Ustin, S. L., & Lee, C. T. (2003).

Evaluation of the potential of Hyperion for fire danger assessment by comparison to the

Airborne Visible/Infrared Imaging Spectrometer. *IEEE Transactions on Geoscience and*

Remote Sensing, *41*(6), 1297–1310. IEEE Transactions on Geoscience and Remote

Sensing. https://doi.org/10.1109/TGRS.2003.812904

Rose, R. A., Byler, D., Eastman, J. R., Fleishman, E., Geller, G., Goetz, S., … Wilson, C. (2015).

Ten ways remote sensing can contribute to conservation. *Conservation Biology*, *29*(2),

350–359. https://doi.org/10.1111/cobi.12397

Sankey, T., Hultine, K., Blasini, D., Koepke, D., Bransky, N., Grady, K., ... & Allan, G. (2021).

UAV thermal image detects genetic trait differences among populations and genotypes of

Fremont cottonwood (Populus fremontii, Salicaceae). *Remote Sensing in Ecology and*

Conservation, *7*(2), 245-258. https://doi.org/10.1002/rse2.185

Schweiger, R., Kaufman, S., Laaksonen, R., Kleber, M. E., März, W., Eskin, E., … Halperin, E.

(2016). Fast and Accurate Construction of Confidence Intervals for Heritability.

American Journal of Human Genetics, *98*(6), 1181–1192.

https://doi.org/10.1016/j.ajhg.2016.04.016

Seeley, M. M., & Asner, G. P. (2021). Imaging Spectroscopy for Conservation Applications.

Remote Sensing, *13*(2), Article 2. https://doi.org/10.3390/rs13020292

Seeley, M. M., & Asner, G. P. (2023). Large-Scale Controls on the Leaf Economic Spectrum of

the Overstory Tree Species Metrosideros polymorpha. *Remote Sensing*, *15*(19), Article

19. https://doi.org/10.3390/rs15194707

Seeley, M. M., Martin, R. E., Vaughn, N. R., Thompson, D. R., Dai, J., & Asner, G. P. (2023a).

Quantifying the Variation in Reflectance Spectra of Metrosideros polymorpha Canopies

across Environmental Gradients. *Remote Sensing*, *15*(6), Article 6.

https://doi.org/10.3390/rs15061614

Seeley, M. M., Stacy, E. A., Martin, R. E., & Asner, G. P. (2023b). Foliar functional and genetic

variation in a keystone Hawaiian tree species estimated through spectroscopy. *Oecologia*.

https://doi.org/10.1007/s00442-023-05374-1

Seeley, M. M., Vaughn, N. R., & Asner, G. A. (2024). Evaluating individual tree species

classification performance across diverse environments. *Environmental Research:*

Ecology, *3*(1), 011001. https://doi.org/10.1088/2752-664X/ad1f49

Seeley, M. M., Vaughn, N. R., Shanks, B. L., Martin, R. E., König, M., & Asner, G. P. (2023c).

Classifying a highly polymorphic tree species across landscapes using airborne imaging

spectroscopy. *Remote Sensing*, *15*(8), 4365. https://doi.org/10.3390/rs15184365

Seeley, M. M.; Wiebe, B. C.; Asner, G. P.; Abraham, A. J.; Cooper, H. F.; Gehring, C. A.;

Hultine, K. R.; Allan, G. J.; Whitham, T. G.; Doughty, C. E. (2025a) Why Do Models Go

Wrong? The Importance of Adaptations and Acclimations in Driving Landscape-Level

Spectral Variation. Available at SSRN http://dx.doi.org/10.2139/ssrn.5379735

Seeley, M. M., Wiebe, B. C., Gehring, C. A., Hultine, K. R., Posch, B. C., Cooper, H. F., ... &

Doughty, C. E. (2025b). Remote sensing reveals inter□and intraspecific variation in

riparian cottonwood (*Populus* spp) response to drought. *Journal of Ecology*.

https://doi.org/10.1111/1365-2745.70061

Serbin, S. P., Dillaway, D. N., Kruger, E. L., & Townsend, P. A. (2012). Leaf optical properties

reflect variation in photosynthetic metabolism and its sensitivity to temperature. *Journal*

of Experimental Botany, *63*(1), 489–502. https://doi.org/10.1093/jxb/err294

Shafroth, P. B., Friedman, J. M., & Ischinger, L. S. (1995). Effects of Salinity on Establishment

of Populus Fremontii (cottonwood) and Tamarix Ramosissima (saltcedar) in

Southwestern United States. *The Great Basin Naturalist*, *55*(1), 58–65.

Slaton, M. R., Raymond Hunt Jr., E., & Smith, W. K. (2001). Estimating near-infrared leaf

reflectance from leaf structural characteristics. *American Journal of Botany*, *88*(2), 278–

284. https://doi.org/10.2307/2657019

Smirnoff, N. (2005). *Antioxidants and Reactive Oxygen Species in Plants*. John Wiley & Sons,

Ltd. https://doi.org/10.1002/9780470988565.fmatter

Stromberg, J. (1998). Dynamics of Fremont cottonwood (*Populus fremontii*) and saltcedar

(*Tamarix chinensis*) populations along the San Pedro River, Arizona. *Journal of Arid*

Environments, *40*(2), 133–155. https://doi.org/10.1006/jare.1998.0438

Tafesse, E. G., Warkentin, T. D., Shirtliffe, S., Noble, S., & Bueckert, R. (2022). Leaf pigments,

surface wax and spectral vegetation indices for heat stress resistance in

pea. *Agronomy*, *12*(3), 739.

Tang, T., Liu, P., Zheng, G., & Li, W. (2016). Two phases of response to long-term moderate

heat: Variation in thermotolerance between Arabidopsis thaliana and its relative Arabis

paniculata. *Phytochemistry*, *122*, 81–90. https://doi.org/10.1016/j.phytochem.2016.01.003

Thompson, D. R., Boardman, J. W., Eastwood, M. L., & Green, R. O. (2017). A large airborne

survey of Earth's visible-infrared spectral dimensionality. *Optics Express*, *25*(8), 9186.

https://doi.org/10.1364/OE.25.009186

Tirado, S. B., Dennis, S. St., Enders, T. A., & Springer, N. M. (2021). Utilizing spatial variability

from hyperspectral imaging to assess variation in maize seedlings. *The Plant Phenome*

Journal, *4*(1), e20013. https://doi.org/10.1002/ppj2.20013

Turner, M. (2023). Heritable trait variation for climate resilience. *Nature Ecology & Evolution*,

7(4), 492–492. https://doi.org/10.1038/s41559-023-02004-0

Turner, W., Spector, S., Gardiner, N., Fladeland, M., Sterling, E., & Steininger, M. (2003).

Remote sensing for biodiversity science and conservation. *Trends in Ecology &*

Evolution, *18*(6), 306–314. https://doi.org/10.1016/S0169-5347(03)00070-3

Vahsen, M. L., Blum, M. J., Megonigal, J. P., Emrich, S. J., Holmquist, J. R., Stiller, B., …

McLachlan, J. S. (2023). Rapid plant trait evolution can alter coastal wetland resilience to

sea level rise. *Science*, *379*(6630), 393–398. https://doi.org/10.1126/science.abq0595

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., …

Vázquez-Baeza, Y. (2020). SciPy 1.0: Fundamental algorithms for scientific computing

in Python. *Nature Methods*, *17*(3), 261–272. https://doi.org/10.1038/s41592-019-0686-2

Vogelmann, J. E., Rock, B. N., & Moss, D. M. (1993). Red edge spectral measurements from

sugar maple leaves. *Remote Sensing*, *14*(8), 1563–1575.

https://doi.org/10.1080/01431169308953986

Wahid, A., & Close, T. J. (2007). Expression of dehydrins under heat stress and their relationship

with water relations of sugarcane leaves. *Biologia Plantarum*, *51*(1), 104–109.

https://doi.org/10.1007/s10535-007-0021-0

Wang, Z., Townsend, P. A., & Kruger, E. L. (2022). Leaf spectroscopy reveals divergent inter-

and intra-species foliar trait covariation and trait–environment relationships across

NEON domains. *New Phytologist*, *235*(3), 923–938. https://doi.org/10.1111/nph.18204

Weingarten, E., Martin, R., Hughes, F., Vaughn, N., Schafron, E., & Asner, G. P. (2021). Early

Detection of a Tree Pathogen using Airborne Remote Sensing. *Ecological Applications.*,

21(2).

Whitham, T. G., Bailey, J. K., Schweitzer, J. A., Shuster, S. M., Bangert, R. K., LeRoy, C. J., …

Wooley, S. C. (2006). A framework for community and ecosystem genetics: From genes

to ecosystems. *Nature Reviews Genetics*, *7*(7), Article 7. https://doi.org/10.1038/nrg1877

Xie, X. j., Zhang, Y. h., Li, R. y., Shen, S. h., & Bao, Y. x. (2019). Prediction model of rice

crude protein content, amylose content and actual yield under high temperature stress

based on hyper-spectral remote sensing. *Quality Assurance and Safety of Crops & Foods*,

11(6), 517–527. https://doi.org/10.3920/QAS2018.1295

Yan, Z., Guo, Z., Serbin, S. P., Song, G., Zhao, Y., Chen, Y., … Wu, J. (2021). Spectroscopy

outperforms leaf trait relationships for predicting photosynthetic capacity across different

forest types. *New Phytologist*, *232*(1), 134–147. https://doi.org/10.1111/nph.17579

Source Population	Elevation	Mean Annual Temperature (°C)	Mean Annual Precipitation (mm)	Ecotype
MRN	1774	10.4	593	Sonoran Desert Main
KKH	1920	10.7	258	Colorado Plateau
JLA	1507	12.3	212	Colorado Plateau
CLF	1299	14.1	176	Colorado Plateau
KWF	1126	15	243	Sonoran Desert KWF
BCE	1109	15.6	432	Sonoran Desert Main
PSA	1234	15.7	471	Sonoran Desert SE
TSZ	1219	16.9	322	Sonoran Desert SE
CAF	988	17.2	440	Sonoran Desert Main
TSE	986	17.5	402	Sonoran Desert SE
WHY	575	19.6	284	Sonoran Desert Main
CCU	696	19.9	349	Sonoran Desert Main
NRV	666	19.9	337	Sonoran Desert Main
SCT	26	22.1	88	Sonoran Desert Main
LBW	143	22.3	137	Sonoran Desert Main
CCR	70	22.6	97	Sonoran Desert Main
Common Garden Site				
Canyonlands	1581	10.7	225	
Agua Fria	998	17.2	453	
Yuma	49	22.8	93	

Table 1: Source population and study site environmental conditions. For source populations, the

ecotype is listed.

Figure Captions

Figure 1: Source population and common garden site locations in Arizona and southern Utah.

Background colors represent Köppen climate types.

Figure 2: Environmental conditions at the common garden sites from 2014 (garden

establishment) to 2018 (data collection). All data represent the mean monthly values.

Photosynthetically active radiation (PAR) data are the daily sum radiation values collected every

three hours by the Moderate Resolution Imaging Spectroradiometer (MODIS). Mean, minimum,

and maximum temperature and vapor pressure deficit (VPD) were obtained from the PRISM

Gridded Climate Data.

Figure 3: a) Mean brightness normalized reflectance of each source population, colored by

common garden site. See Figure S2 for reflectance prior to brightness normalization. Boxplots of

albedo calculated across the b) visible and c) near-infrared-shortwave infrared, d) leaf mass per

area (LMA), and e) chlorophyll index – green (CIG). Within each garden, boxplots are ordered

according to source population mean annual temperature, with cooler populations on the left and

warmer populations on the right. Gaps correspond to populations that were not sampled at a

given garden. Boxplots represent a different number of sampled trees per population per site.

Figure 4: a) First and second principal components (PC) of leaf spectra colored according to the

common garden site. b) Distance between the centroids of source population grouped by ecotype

x common garden in PC space. Colors represent common garden (blue: Canyonlands, grey:

Agua Fria). Shapes correspond to the ecotype (x: Colorado Plateau, circle: Sonoran Desert,

triangle: Sonoran Desert SE).

Figure 5: Heritability of reflectance spectra (solid line) and its 95% bootstrapped confidence

interval (grey fill), mean brightness normalized reflectance spectra (dotted line), and the source

of variability (filled colors) from the visible to shortwave infrared for *P. fremontii* at a) all sites

and b) Canyonlands and Agua Fria.

Figure 6: Support vector machine confusion matrix results for predicting a) common garden site

and b) source population. Source populations are ordered left to right and top to bottom from

lowest to highest mean annual temperature.

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6