Title:

Estimating and correcting index hopping misassignments in single-cell RNA-seq data

Short title:

Index hopping correction in single-cell RNA-seq data

Authors and affiliations:

Lingling Miao[1], Loren Collado[1], Savannah Barkdull[1], Yoshine Saito[1], Jay-Hyun Jo[1],

Jungmin Han[1], Stefania Dell'Orso[2], Michael C. Kelly[3], Sean Conlan[4], Heidi H. Kong[1],

and Isaac Brownell[1]

1 Dermatology Branch, NIAMS, NIH, Bethesda, Maryland, USA

2 Genomic Technology Section, NIAMS, NIH, Bethesda, Maryland, USA

3 Single Cell Analysis Facility, Cancer Research Technology Program, Frederick

National Laboratory, Bethesda, Maryland, USA

4 Microbial Genomics Section, Translational and Functional Genomics Branch, NHGRI,

NIH, Bethesda, Maryland, USA

Corresponding author:

Isaac Brownell, Isaac.brownell@nih.gov

ABSTRACT

Index hopping causes read assignment errors in data from multiplexed sequencing libraries. This issue has become more prevalent with the widespread use of high-capacity sequencers and highly multiplexed single-cell RNA sequencing (scRNA-seq) libraries. We conducted deep, plate-based scRNA-seq on a mixed population of mouse skin cells. Analysis of transcriptomes from 1152 cells identified four distinct cell types. To estimate the error rate in sample assignment due to index hopping, we employed differential expression analysis to identify signature genes that were highly and specifically expressed in each cell type. We quantified the proportion of misassigned reads by examining the detection rates of signature genes in other cell types. Remarkably, regardless of gene expression levels, we estimated that 0.65% of reads per gene were assigned to incorrect cell across our data. To computationally compensate for index hopping, we developed a simple correction method wherein, for each gene, 0.65% of the library's average expression level was subtracted from the expression of each cell. This correction had notable effects on transcriptome analyses, including increased cell-cell clustering distance and alterations in intermediate state assignments of cell differentiation. These findings underscore the potential impact of index hopping on experimental results. In conclusion, we devised a straightforward method to estimate and correct for the index hopping rate by quantifying misassigned genes in distinct cell types within an scRNA-seq library. This approach can be applied to any barcoded, multiplexed scRNA-seq library containing cells with distinct expression profiles, allowing for correction of the expression matrix before conducting biological analysis.

AUTHOR SUMMARY

Index hopping causes errors in multiplexed high-throughput sequencing data whereby inaccurate barcoding leads to the misassignment of sequence reads to an incorrect source. This can result in gene expression assignments to the wrong cell in single-cell RNA sequencing (scRNA-seq). We performed scRNA-seq on sorted mouse skin cells and identified four distinct cell types based on gene expression profiles. Using genes unique to each cell type, we discovered that approximately 0.65% of total reads per gene were incorrectly assigned to another cell due to index hopping, regardless of gene expression levels. To correct for the misassigned reads, we proportionally adjusted each cell's gene expression data. Applying this correction improved analyses allowing for enhanced cell clustering accuracy and refining the identification of intermediate cell states during cell differentiation. Our study emphasizes the significance of index hopping in scRNA-seq experiments and presents a practical correction approach to remove misassigned reads that can be applied to any scRNA-seq study involving cells with distinct gene expression profiles or where non-expressed genes are introduced as a quality control measure during library preparation.

INTRODUCTION

The advent of next-generation sequencing (NGS) has revolutionized genomic research, with contemporary sequencers allowing cost-effective sequencing of large-scale multiplexed libraries [1]. Multiplexing is a common strategy for sequencing multiple samples that involves adding a unique short DNA sequence (index) to each fragment in a sample and pooling samples together for sequencing. However, a notable limitation of

this approach is index misassignment, where sequence reads are erroneously paired

with an index from another sample. Index misassignment can occur at various stages

during NGS including reagent contamination, sample handling, PCR amplification,

patterned flow cell "spread-of-signal", and bioinformatic errors [2, 3]. Among these

mechanisms, a major contributor to index hopping (also known as index swapping or

index switching), is incomplete library cleanup which allows free-floating index primers

to randomly amplify DNA fragments from multiple samples within a pooled library during

sequencing [4]. This phenomenon leads to data contamination through index

misassignment. The impact of index hopping has grown significantly as high capacity

modern sequencing technologies increasingly rely on multiplexing samples. This issue

is particularly evident in single-cell RNA sequencing (scRNA-seq) studies that involve

pooling large numbers of cDNA libraries for sequencing [4-7].

Index hopping poses a challenge in scRNA-seq, impacting both droplet and plate-based

methods [4-7]. Recent investigations analyzing multiplexed droplet-based scRNA-seq

datasets estimated sample index hopping probabilities ranging from 0.3% to 0.9% [4].

Plate-based scRNA-seq studies utilizing the "i5+i7" dual-index multiplexing strategy

reported more variable and higher index hopping rates, ranging from 0.25% to 10% on

Hiseq platforms [5-7]. The index hopping rate varies across different Illumina platforms,

with advanced Hiseq systems, such as Hiseq 3000, which employ patterned flow cells

and exclusion amplification chemistry, exhibiting higher index hopping levels compared

to earlier platforms like Hiseq 2500 [3-5, 8, 9]. Illumina recommends reducing free index

primers or the use of unique dual indexes as experimental approaches to mitigate index

hopping when using dual-index multiplexing strategies [10]. However, unique dual

indexing may not be practical with large samples due to the limitation in barcode

numbers, especially in high-volume scRNA-seq systems. In addition, unique indexing is

not a complete solution as index hopping has been observed in droplet-based scRNA-

seq and multiplexed long-read sequencing which use unique sample barcodes [4, 5,

11]. A few recent studies have explored using computational algorithm to estimate and

correct the contamination effect of index hopping in plated-based and droplet-based

scRNA-seq data [4-6]. However, their approach requires knowing the ground truth of

non-multiplexed sequencing.

Given the limitations of existing strategies, there is a need for an objective method to

estimate and correct for index hopping in scRNA-seq data. In this study, we performed

an in-depth analysis of plate-based scRNA-seq data obtained from mouse skin cells.

We developed a simplified method that leverages distinct expression profiles among

cells to estimate the index hopping rate and implemented a correction approach to

adjust for the contamination caused by index hopping. We further demonstrate the

impact of correcting for index hopping before conducting biological analysis of scRNA-

seq data. Our findings contribute to advancing the understanding of index hopping and

provide a practical solution for its estimation and correction in scRNA-seq data analysis.

RESULTS

Index hopping rates can be estimated by the frequency of non-self genes in

scRNA-seq data

We performed plate-based scRNA-seq on FACS sorted dissociated mouse skin cells.

The skin is a complex tissue that encompasses multiple cell types including, but not

limited to, Merkel cells, dermal papilla fibroblasts, hematopoietic cells, and epithelial

keratinocytes (Fig. S1). The transcription factor SRY-Box Transcription Factor 2 (SOX2)

is expressed in Merkel cells and dermal papilla cells [12], whereas Growth Factor

Independence 1 (GFI1) is expressed in Merkel cells and hematopoietic cells [13, 14].

We sorted single cells expressing GFP from the skin of $Sox2^{GFP/+}$ or $Gfi1^{GFP/+}$ reporter

mice at various developmental stages, along with GFP-negative epidermal cells

expected to be keratinocytes (Fig. 1A). Single cells were sorted into lysis buffer in 96

well plates, and the Smart-seq2 method was used to generate cDNA from these sorted

individual cells. Cleaved cDNA was tagged with dual index adaptors prior to being

pooled into three 384-sample sequencing libraries (Fig. 2A, S2A-C). On average, each

cell yielded ~3 million uniquely mapped reads, allowing for the detection of

approximately 6000 genes per cell. By employing Seurat clustering, we grouped the

individual cells into four clusters based on their transcriptome similarities (Fig. 1B).

Furthermore, by identifying highly differentially expressed genes (marker genes) within

these four clusters, we were able to distinguish them as four distinct cell types: Merkel

cells (MCs), dermal papilla cells (DCs), hematopoietic cells (HCs), and keratinocytes

(KCs) (Fig. 1C). As the pooled libraries included all four cell types, index hopping had

the potential to misassign reads from one cell type to another based on their sharing of

i7 or i5 indexes (Fig. 2A, S2A-B).

Figure 1. scRNA-seq identifies different skin cell types.

(A) Overview of scRNA-seq experimental procedure.

(B) tSNE plot showing clustering of the gene expression profiles of single skin cells.

(C) Heatmap showing highly differentially expressed genes (marker genes) for each cell cluster identified

by Seurat clustering.

Index hopping occurs when cDNA from one cell is indexed erroneously with another

cell's barcodes. After library pooling, unincorporated free index primers associated with

a row or column on the barcoding plate can potentially be added to cDNA from another

row/column creating the risk for index hopping and misassignments of reads. Detection

of such an event is difficult when the two wells contain similar transcriptomes. However,

if the two cells are of different lineages and have distinct gene expression, index

hopping contamination can be measured by detecting non-self genes (transcript not

expressed in that cell type) in a specific cell or from an empty well in plated-based

sequencing libraries. To identify non-self genes, we used the differentially expressed

marker genes for each of the four cell types in our library (Fig. 1C). We were able to

detect marker genes for each of the four cell types labeled with indexes for empty wells.

We then calculated the average library expression levels for all genes by dividing the

total transcripts per million (TPM) for each gene in the library by the number of cells in

the library. Similarly, we calculated the average detection levels for each gene in empty

wells by dividing the total TPM from the empty wells by the number of empty wells.

Comparing the average gene detection levels in the library and in empty wells, we

observed a positive correlation (Fig. 2B, S2D), indicating that misassignments were

likely a result of a stochastic process like index hopping rather than technical errors in

sample preparation.

Figure 2. Index hopping can be detected when occurring between differing cell types.

(A) Overview of the indexing used for 384 single-cell cDNA libraries in the first pooled sequencing library.

Cell types together with empty wells and quality control (QC) failed wells are color-coded. Arrows show for

each individual Merkel cells (MCs), the potential index hopping with dermal cells (DCs, orange arrow) or

hematopoietic cells (HCs, brown arrow). Vertical arrows show i7 index hopping, and horizontal arrows show

i5 index hopping.

(B) Scatter plot showing the correlation of average TPM of cell type marker genes in empty wells and the

entire library, with values transformed using a log2 scale (log2(average TPM +1). Color labels marker genes

for the 4 cell types.

To assess index hopping in our library, we quantified non-self gene detection in the four

cell types. Considering the location of single cell samples within the multi-well plates

and the application of dual index primer sets in our libraries, there was the potential for

dual index hopping (misassignment can happen by either i5 or i7 hopping), single index

hopping (misassignment can happen only by i5 or i7 hopping), and no potential index

hopping between DCs and MCs, while dual index hopping and i5 index hopping were

possible between HC and MC (Fig. 2A, S2A-B). To estimate the rate of read

misassignments, we calculated the following:

Firstly, we computed the average transcripts per million (TPM) of a gene in MCs by

dividing the total TPM for that gene in MCs by the number of MCs (gene A as an

example, $\overline{TPM}_{geneA,MC}$). The data were shown on a log2 scale (log2($\overline{TPM}_{geneA,MC}$ + 1).

$$\overline{TPM}_{geneA,MC} = \frac{1}{n_{MC}}\sum_{i=1}^{n_{MC}} TPM_i$$

$\overline{TPM}_{geneA,MC}$: average TPM for gene A in MC

n_{MC} : number of single MCs

Then, to determine the ratio of reads assigned to individual MCs (detection ratio) for

each gene (gene A as an example, $\varphi_{geneA,cellB:lib}$), we divided the TPM of each gene in

single MC by its average TPM in the library.

$$\overline{TPM}_{geneA,lib} = \frac{1}{n_{lib}} \sum_{i=1}^{n_{lib}} TPM_i$$

$\overline{TPM}_{geneA,lib}$: average TPM for gene A in sequencing library

n_{lib} : number of single cells in library

$$\varphi_{geneA,cellB:lib} = \frac{TPM_{geneA,cellB}}{\overline{TPM}_{geneA,lib}}$$

$\varphi_{geneA,cellB:lib}$: for gene A, the detection ratio between single cell B and average

expression in library

And we were able to calculate the average detection ratio of each gene between MCs

and the entire library (gene A as an example, $\overline{\varphi}_{geneA,MC:lib}$).

$$\overline{\varphi}_{geneA,MC:lib} = \frac{1}{n_{MC}} \sum_{i=1}^{n_{MC}} \varphi_{geneA,cell_i:lib}$$

$\overline{\varphi}_{geneA,MC:lib}$: for gene A, the average detection ratio between MCs and average

expression in library

n_{MC} : number of single MCs

If index hopping played a significant role in misassignment, the detection ratio of non-

self genes in MCs would correspond to the index hopping rate. Notably, the detection

ratio of top DC genes in MCs was higher in MCs with dual index hopping from DCs

compared to those with single index hopping from DCs. Conversely, when no index

hopping occurred between MC and DC, the detection ratio of DC genes in MCs was

nearly zero (Fig. 3A-B, S3A-B). Similarly, the detection ratio of top HC genes in MCs

with dual index hopping from HCs was higher than those with i5 index hopping from

HCs (Fig. 3C, D). Furthermore, the detection ratio of DC and HC genes in MCs

exhibited a positive correlation with the number of index-hopped DCs (Fig. S3C) and

HCs (Fig. S3D), respectively.

Figure 3. Non-self marker genes are detected at different levels in MCs with dual, single and no index hopping to other cell types.

(A) and **(B)** MCs which had dual (A) and no (B) index hopping potential from DCs: Left, scatter plot showing the average TPM for individual genes in single MCs correlated with their average TPM in the library. TPMs are displayed on a log2 scale (log2 (average TPM + 1)). Middle, scatter plot showing the average detection ratios of DC genes in single MCs. Colored labels indicate the top marker genes for different cell types. Box plots show the median and interquartile range ± SD of average detection ratio for color-labeled genes. Ave., average. Right, violin plot showing the detection ratio of the top 6 DC marker genes in individual MCs

(C) and **(D)** MCs which had dual (C) and i5 (D) index hopping potential from HCs: Left, scatter plot showing the average TPM in single MCs correlated with their average TPM in the library. TPMs are displayed on a log2 scale (log2 (average TPM + 1)). Middle, scatter plot showing the average detection ratio of HC genes in single MCs. Colored labels indicate the top marker genes for different cell types. Box plots show the median and interquartile range ± SD of average detection ratio for color-labeled genes. Ave., average. Right, violin plot showing the detection ratio of the top 6 HC marker genes in individual MCs.

Adjusting gene expression levels by the estimated index hopping rate compensates for misassigned reads

Having observed that index hopping was relatively consistent, we sought to correct for its effect. A group of highly-expressed HC and DC marker genes with no expected expression in MCs was detected in MCs at levels consistent with their average TPM in the library (Fig. 4A). We referred to this distinctive group of genes with their misassigned reads in MCs as the "side population". The detection ratios of these genes in MCs ($\overline{\varphi}$) were consistently below 1%, regardless of their varying expression levels in the libraries (Fig. 4B). Notably, there were other highly-expressed HC and DC marker genes that appeared to have native expression in MC as indicated by their detection ratios being greater than 1%. These non-specific marker genes clustered away from the side population genes (Fig. 4A-B). To estimate average index hopping rates in MC, we employed a methodology inspired by the Rank Ordering of Super-Enhancers (ROSE) tool for super-enhancer identification from ChIP-seq data [15]. We first ranked and

plotted all genes increasingly by the reciprocal of their average gene detection ratio ($\overline{\varphi}$)

in MCs. This plot demonstrated a distinct inflection point where the reciprocal of the

detection ratio in MCs began increasing rapidly. To geometrically define this point, we

rescaled the data so that the x and y axes ranged from 0 to 1. Next, we determined

where a tangent to the rank curve had a slope of 1. The gene detection ratio at this

point was calculated to be 0.65%, which we used as the estimated index hopping rate in

our sequencing library. Genes ranked below this threshold were classified as

predominantly MC-expressed genes, whereas genes ranked above this threshold were

considered as genes whose detection in MCs was due, in part, to index hopping

misassignment (Fig. 4C). This classification was validated by identifying the rank order

position of the differentially expressed marker genes for each cell type (Fig. S4A-D).

Notably, genes in the side population of top HC and DC genes were ranked well above

the inflection point, and the non-specific marker genes ranked near the inflection point

(Fig. 4D). Consistent with MC and KC having shared epithelial progenitors and similar

transcriptomes, the detection ratio of KC marker genes in MCs was typically higher than

the index hopping rate reflecting the relative lack of specific KC marker genes (Fig. S5A,

B). Following this pattern, in cases where marker genes appeared to be specific to the

two cell types being compared, a distinct side population was observed when visualizing

average expression levels and detection ratios (Fig. S5C-H).

Figure 4. Index hopping rate can be estimated using non-self genes

(A) Scatter plot showing the average TPM of top HC and DC genes in single MCs correlated to their average TPM in library. Blue line separates highly-expressed HC and DC "side" population genes where the level in MCs is proportional to their average TPM in the library. TPMs are displayed on a log2 scale (log2 (average TPM + 1)). Colors indicate the top MC, HC and DC marker genes.

(B) Scatter plot showing the average gene detection ratios in single MCs. Box plots show median and interquartile range ± SD of average detection ratio in MCs for side population genes. Colors indicate the top HC and DC marker genes.

(C) Rank ordered scatter plot showing distribution of gene detection ratios in MCs across all genes from mm10 genome annotated by Ensembl. Blue box indicates the MC expressed genes, red box indicates genes presumptively detected due to index hopping, grey box indicates genes with no detection in MCs. The cyan tangent line with slope = 1 identifies the inflection point.

(D) Focused rank ordered scatter plot showing genes near the inflection point from plot in (C).

To address the background gene detection resulting from index hopping, we

implemented a simple correction method. We subtracted 0.65% of the average

transcripts per million (TPM) for each gene in our library from the single-cell gene TPM

matrix. Following the correction, the detection ratios of non-self genes in MCs (Fig. 5A-

F) and other cell types (Fig. S5A'-H') was notably reduced. Additionally, we investigated

the impact of the correction on individual gene TPM values in MCs. As expected, highly

expressed HC and DC marker genes showed notably reduced detection in MCs after

correction, whereas the reduction in MC marker genes in MCs was inconsequential

(Fig. 5G-H, S6A-D). Similarly, when applying the correction in HCs, only non-self genes

with high library TPM values were notably impacted, whereas HC marker genes

remained comparable to pre-correction levels in HCs (Fig. S6E-J).

Figure 5. Index hopping correction reduces the detection of non-self marker genes.

(A) and **(B)** Same as Figure 4A and B – precorrection scatter plots for comparison.

(C) and **(D)** Scatter plots showing the average TPM (C) and detection ratios (D) of individual genes in single

MCs after index hopping correction. Colored labels indicate the top marker genes for different cell types.

The blue line in (C) is same as the one in (A) which separates the side population. The cyan dashed line in

(B and D) represents the estimated index hopping rate before correction (0.65%). TPMs are displayed on

a log2 scale (log2 (average TPM + 1)). Box plots depict the median and interquartile range ± SD of average

detection ratio for the side population genes after correction.

(E) and **(F)** Violin plots showing the detection ratios of the top 6 DC (E) and top 6 HC (F) marker genes in

MCs before and after correction.

(G) and **(H)** Scatter plots showing the log2 scaled average TPM of individual genes in single MCs before

and after correction. Colors indicate the marker genes in different cell types (G) and represent levels of

average TPM in library (H).

Correcting for index hopping impacts single-cell analysis and interpretation

To assess the impact of index hopping correction on the bioinformatic analysis of the

scRNA-seq data, we analyzed our data set with and without correction. To evaluate the

heterogeneity among single cells, we employed cell-to-cell distance, which quantifies

transcriptomic similarities between cells. A smaller cell distance indicates a higher level

of similarity between two cells [16]. We compared the cell distances among all cells in

the library before and after correction. As misassigned read contamination would be

expected to artificially make distinct cell types more similar, we anticipated that index

hopping correction would increase distances between cells. The density plot of cell

distances demonstrated that index hopping correction did indeed increase measured

heterogeneity within the DC and HC clusters (Fig. 6A, B). This suggests that index

hopping correction can mitigate artificial cell-cell similarities caused by randomly

misassigned reads across scRNA-seq libraries.

Figure 6. Index hopping correction increases the detection of single cell heterogeneity.
Density plots showing the distribution of cell-to-cell distance among individual DCs (A) and HCs (B) before
and after correction. Kolmogorov-Smirnov test: p < 2.2 x 10^{-16} for cell-to-cell distance before correction vs.
after correction.

Next, we investigated how index hopping correction impacted the analysis of

developmental stage classification. Our scRNA-seq dataset included single cells

spanning a wide range of developmental timepoints, from embryonic to postnatal skin.

Focusing on DCs, we subjected uncorrected and corrected data to Seurat clustering

identifying three cell clusters (Fig. S7A, E). Consistent with the cells' chronological age,

each cluster represented a distinct stage of skin development (Fig. S7B, F). We then

used Monocle trajectory analysis to plot the single DCs along a developmental

pseudotime before (Fig. S7C, D) and after (Fig. S7G, H) index hopping correction.

Comparing the developmental stage classification of each DC before and after

correction, we observed that 79.9% of DC cells retained their stage classification

whereas 20.1% of cells were reclassified after correction (Fig. 7A). Consistent with

stage reclassifications, there was a reordering of the DC pseudotime trajectory after

index hopping correction (Fig. 7B), suggesting that index hopping misassignments had

impacted the results of both cell clustering and state hierarchy analyses.

Figure 7. Index hopping correction impacts the single-cell analysis of cell developmental stages.

(A) and **(E)** Upper: Scatter plots showing the cell developmental stages of individual DCs (A) and HCs (E)

assigned by Seurat clustering before and after index hopping correction. Lower: Histogram showing the

frequency of cell stage assignment changes from before correction to after correction.

(B) and **(F)** Scatter plots showing the pseudotime order of individual DCs (B) and HCs (F) identified by

Monocle trajectory analysis before and after correction. Colors indicate the age of mouse source for

individual cells.

(C) and **(G)** Venn diagrams comparing differentially expressed marker genes for any developmental stage

of DCs (C) and HCs (G) identified by Seurat before and after correction.

(D) and **(H)** Venn diagrams comparing pseudotime-dependent genes identified by Monocle for DCs (D) and

HCs (H) before and after correction.

We performed a similar pre and post correction comparison for two age groups of HCs

(E16.5 and E17.5). Clustering identified two distinct cell clusters before (Fig. S8A, B)

and after (Fig. S8E, F) correction, and trajectory analysis revealed a potential

developmental trajectory before (Fig. S8C, D) and after (Fig. S8G, H) correction. Index

hopping correction also exhibited an analogous, albeit smaller, effect on the stage

classification and pseudotime order of some HCs (Fig. 7E, F).

Analysis of differentially expressed genes is often used to define transcript signatures

for cell clusters whereas identification of genes that are dynamically expressed over a

pseudotime trajectory can suggest biologically relevant genes. Accordingly, we explored

the impact of index hopping correction on these analyses. Using Seurat and Monocle,

we identified differentially expressed developmental stage marker genes as well as

genes that significantly changed over the developmental pseudotime for both DCs and

HCs. Comparing the aggregate set of stage marker genes and pseudotime-dependent

genes before and after index hopping correction, we observed a substantial overlap in

each gene set for DCs (Fig. 7C, D) and HCs (Fig. 7G, H). Nonetheless, index hopping

correction altered the composition of these gene lists in each case, once again

suggesting that index hopping misassignments can impact the gene composition of cell

state signatures. Collectively, these results demonstrate that index hopping measurably

influences the biological interpretation of scRNA-seq data.

Hiseq2500 generates a lower index hopping rate compared to Hiseq3000

As sequencing technology has advanced, the use of patterned flow cells has increased

throughput and efficiency. However, sequencing platforms utilizing a patterned flow cell

with Exclusion Amplification chemistry, such as the HiSeq3000 and NovaSeq, appear to

exhibit higher levels of index hopping compared to platforms without patterned flow cells

[3-5, 8, 9]. Our libraries were sequenced using the HiSeq3000 platform. To test if the

patterned flow cell technology impacted the degree of index hopping, we re-sequenced

one library (Pool #2) on the HiSeq2500, which has a similar sequencing depth but does

not employ a patterned flow cell. We compared the gene detection ratios in MCs

between the HiSeq2500 and HiSeq3000 platforms. As expected, when analyzing MCs,

the detection ratios of genes in the side population, including top HC markers, was

reduced on the HiSeq2500 compared to the HiSeq3000, despite the average library

TPM remaining unchanged between the two platforms (Fig. 8A-D). The detection ratio

of top DC and HC genes in individual MCs was also lower on the HiSeq2500 compared

to the HiSeq3000 (Fig. 8E, F). Interestingly, the detection ratio of non-self genes in MCs

on the HiSeq2500 was comparable to their HiSeq3000 post-correction levels (Fig. 8A-F,

5C-F, S5A'-H'). Similar trends were observed in HCs, where the detection ratios of non-

HC genes were lower on the HiSeq2500 compared to the HiSeq3000 (Fig. S9).

Furthermore, the cell distance among single cells in Pool #2 was greater on the

HiSeq2500 than on the HiSeq3000 (Fig. 8G). These findings confirm that patterned flow

cell technologies are more susceptible to index hopping [3-5, 8, 9], and that our

correction strategy compensates for this vulnerability.

Figure 8. Index hopping is less pronounced in Hiseq2500 platform compared to Hiseq3000.

(A-D) Scatter plots showing the average TPM (A, C) and detection ratio (B, D) without index hopping

correction of individual genes in single MCs in library pool #2 sequenced on Hiseq2500 (A, B) and

Hiseq3000 (C, D) platforms. TPMs are displayed on a log2 scale (log2 (average TPM + 1)). Colors indicate

the top marker genes for different cell types. Box plots depict the median and interquartile range ± SD of

average detection ratio for the color-labeled HC genes.

(E) and **(F)** Violin plots showing the detection ratios in MCs of the top 6 DC (E) and HC (F) marker genes

using sequencing data from Hiseq2500 and Hiseq3000 platforms without index hopping correction.

(G) Density plot showing the distribution of cell-to-cell distance among single cells in library pool #2

sequenced on Hiseq2500 and Hiseq3000. Kolmogorov-Smirnov test: $p < 2.2 \times 10^{-16}$ for cell-to-cell distance

from Hiseq2500 vs. Hiseq3000.

DISCUSSION

In this study, we developed an approach to estimate and correct for index hopping in

multiplexed scRNA-seq data. We performed plate-based single cell sequencing of four

distinct cell types sorted from mouse skin with a focus on MCs as the most abundant

cell type in the libraries. To estimate the rate of read misassignments, we analyzed

differentially expressed genes across each cell type. Top genes expected to be

expressed exclusively in their respective cell types were designated as marker genes,

and the presence of non-self marker genes in a particular cell was interpreted as

contamination from index hopping. We calculated the gene detection ratio for each gene

in MCs and estimated the index hopping rate based on the ratio of non-self genes.

Since the detection ratio of non-self genes in MCs was consistent regardless of their

overall detection levels in the library, we assumed a fixed index hopping rate across all

samples in the libraries. To correct for index hopping, we reduced the TPM for each

gene by the index hopping rate multiplied by the transcript's average expression across

the library. The correction significantly reduced read misassignments, as evidenced by

the decreased detection ratio of non-self genes. Comparing data obtained from

Hiseq3000 and Hiseq2500 sequencing platforms confirmed that index hopping was

more pronounced in platforms employing patterned flow cells (e.g., Hiseq3000/4000/X,

NextSeq 1000/2000, NovaSeq X/X Plus, and NovaSeq 6000). Furthermore, we

evaluated the impact of index hopping correction on biological analysis. Our findings

revealed that correction of index hopping mitigated artificial cell-cell similarities,

impacted the results of cell clustering and state hierarchy analyses, and influenced the

gene composition of cell state signatures. In summary, we propose a simple and

practical methodology for estimating and correcting index hopping that appears to

improve the accuracy of transcriptome data and the cell biology inferred from its

analysis.

By strategically pooling cells with distinct transcriptomes, we were able to effectively

identify and characterize the extent of index hopping. We propose that this method can

be adapted to detect index hopping in multiplex sequencing experiments involving both

heterogeneous and homogeneous libraries. In the case of a homogeneous cell

population, one approach is to introduce single cells from a different cell type as "spike-

ins" during the library preparation process. This deliberate addition of distinct cells will

introduce non-self genes into the otherwise homogenous cell library allowing for the

generation of a side population of non-self genes if index hopping events occur. By

leveraging this spike-in strategy, researchers can effectively assess the presence and

impact of index hopping contamination, thus enhancing the quality and reliability of

scRNA-seq data in various experimental settings.

Our approach to index hopping has certain limitations. An Illumina white paper

demonstrated a correlation between the concentration of free index primers and the rate

of index hopping [10]. Variations in the amounts of cDNA and free index primers can

produce concentration inconsistencies when multiplexing and pooling libraries. To

mitigate this variability, we normalized single-cell cDNA concentrations prior to

tagmentation and indexing. However, multiplexed libraries with varied cDNA

concentrations per sample may result in differing index hopping rate across libraries.

Consequently, estimating a universal index hopping rate for correction may not be

possible. Therefore, we recommend normalizing cDNA concentration prior to indexing

when possible. In addition, our approach was applied to dual indexed multiplexed

libraries. Although unique indexing has been proposed as a strategy to address index

hopping, this issue can still arise, especially due to the amplification of free index

primers on sequencing platforms with patterned flow cells, as evidenced by the droplet-

based scRNA-seq studies [4]. Finally, our comparative analyses of scRNA-seq data

before and after correction revealed substantial effects on biological analyses. It is

crucial to note that the impact of correction on biological analysis may hinge on the

specific index hopping rate within each dataset. Balancing the decision to correct for

index hopping is paramount, as over-correction can also introduce false results,

particularly in analyses that are sensitive to minor changes in gene expression, such as

genes with low expression levels. Each experimental scenario requires a thoughtful

assessment to determine the most appropriate course of action when considering index

hopping correction.

In conclusion, our comprehensive analysis of index hopping in plate-based deep

scRNA-seq data suggests that having cells with distinct transcriptomes, due to intrinsic

diversity or the inclusion of cell "spike-ins" during library preparation, can serve as a

valuable strategy to address index hopping data contamination. To compensate for

index hopping, we developed a simplified method that enables both the estimation of

the index hopping rate and the subsequent correction of read misassignments.

However, it is crucial to consider the specific biological questions being addressed by

the dataset when making the decision to apply index hopping correction. By employing

our approach, researchers can gain a better understanding of the index hopping

phenomenon and its impact on scRNA-seq data. This knowledge can help improve the

accuracy and reliability of downstream analyses, ultimately leading to more robust and

meaningful biological insights.

MATERIALS AND METHODS

Animals

Sox2$^{GFP/+}$ [17] and *Gfi1$^{GFP/+}$* [14] mice were derived from breeders purchased from Jax

(#017592 and #016162, respectively). Mice were housed and bred on an outcrossed

Swiss Webster background in a pathogen-free facility at NIH, Bethesda, MD. Genotyping

of mice was performed by allele-specific PCR on DNA extracted from tail tissue. All

experiments were performed in accordance with institutional guidelines according to

Institutional Animal Care and Use Committee-approved protocols.

Cell dissociation and sorting

Postnatal day 0 (P0) *Sox2GFP/+* and *Gfi1GFP/+* mice were euthanized, and the skin was

harvested, removing the subcutaneous tissue. For embryonic day 18.5 (E18.5) cells, the

pregnant *Sox2GFP/+* female mice were euthanized, and the embryos were harvested and

euthanized. The skin was harvested from the embryos. The harvested E18.5 or P0 skin

was floated in dispase (Corning, #354235) at room temperature for 1 hour to separate the

epidermal sheet. The epidermis was chopped into small pieces, and the epidermal cells

were dissociated in 0.05% Trypsin at 37 °C for 15 minutes, followed by the addition of

DMEM containing 10% FBS. The cells were then sequentially filtered through 70 µm and

40 µm cell strainers. Subsequently, the cells were centrifuged at 300 x g for 5 minutes,

and the supernatant was discarded. The cell pellets were resuspended and processed

for cell sorting using the Beckman MoFlo Astrios or BD influx cell sorter. DAPI was added

immediately before cell sorting, and single DAPI⁻GFP⁺ cells were sorted into 96-well

plates pre-loaded with lysis buffer (lysis buffer prepared following the smart-seq2 method

[18]). Single DAPI⁻GFP⁻ cells from P0 epidermis were sorted as controls.

For embryonic day 16.5 (E16.5) and embryonic day 17.5 (E17.5) cells, pregnant

Sox2GFP/+ (E16.5) and *Gfi1GFP/+* (E16.5 and E17.5) female mice were euthanized, and

the embryos were subsequently harvested and euthanized. The skin from the embryos

was then harvested, chopped into small pieces, and treated with 0.05% Trypsin at 37 °C

for 15 minutes. This was followed by the same cell dissociation procedure used for P0

and E18.5 skin, without separating the epidermis.

Postnatal day 6 (P6) *Sox2GFP/+* and *Gfi1GFP/+* mice were euthanized. The harvested skin

was then incubated in 0.15% trypsin at 37°C for 30 minutes. The epidermis was

obtained by scraping with a scalpel in 0.05% trypsin. Epidermal cells were then

processed using the same method employed for neonatal epidermis.

scRNA-seq

Libraries from isolated single cells were generated following the Smart-seq2 protocol [18]

with the following modifications. Prior to tagmentation, the concentration of single-cell

cDNA libraries was determined using the Quant-iT PicoGreen dsDNA Assay (Thermo

Fisher Scientific). Subsequently, the cDNA libraries were relocated to new plates with

normalized concentration. Libraries from each set of 384 cells with unique barcode

combinations were pooled together, resulting in each pooled library containing cells from

all ages and both reporters. Sequencing was performed on the Hiseq3000 sequencer

(Illumina), resulting in three pooled libraries. Additionally, the second pooled libraries

underwent repeated sequencing on the Hiseq2500 sequencer (Illumina), achieving a

sequencing depth comparable to that of the Hiseq3000. The sequencing generated

paired-end reads of either 125bp or 100bp. Subsequently, reads of 125bp were trimmed

to 100bp for further processing. The sequencing reads were mapped to the mouse mm10

reference genome using STAR 2.5.2b, and gene expression was quantified using RSEM

1.3.0. The gene expression matrix with transcript per million values was used for further

analysis. Log$_2$-scaled TPM values (log$_2$(TPM+1)) were used in all figures where TPM was

shown, unless otherwise specified (log$_2$TPM in Figure 5G, Figure 5H, and Figure S6). For

non-self genes in MCs, their average detection ratios in MCs are considered as the index

hopping rate.

Cell biology analysis

Cell-to-cell distance was analyzed using the R package 'sincell' based on Spearman

correlation [16]. Cell clustering was analyzed using Seurat 2.3 [19]. Cell trajectory was

analyzed using Monocle 2 [20, 21].

Statistical analysis

For comparisons of gene detection in single cells between two different conditions (such

as before vs. after correction, Hiseq2500 vs. Hiseq3000), paired t-test was applied.

DATA AND CODE AVAILABILITY

The scRNA-seq data generated in this study can be accessed in the NCBI Gene

Expression Omnibus (GEO) under the accession number GSE213104. All analyses

were performed in R using established packages. Scripts utilized in this study can be

provided upon request.

ACKNOWLEDGEMENTS

We thank Drs. Livak Ferenc and Teresa Hawley from the CCR Flow Cytometry Core, as

well as Drs. James Simone, Jeff Lay and Kevin Tinsley from the NIAMS Flow Cytometry

Section, for their assistance with cell sorting. We acknowledge Dr. Valery Bliskovsky

from the CCR Genomics Core for his help with single-cell cDNA library preparation, and

Faiza Naz from the NIMAS Genomic Technology Section for her assistance with

sequencing. This work utilized the computational resources of the NIH HPC Biowulf

cluster (https://hpc.nih.gov). This research was supported by the NIAMS intramural

research program (ZIA AR041221, to IB). Its contents are solely the responsibility of the

authors and do not necessarily represent the official views of the NIH.

AUTHOR CONTRIBUTIONS

Conceptualization: IB; Methodology: LM, SD, MCK; Data curation: LM, SD; Resources: IB, SD; Formal analysis: LM, JJ, JH, SC; Investigation: LM, LC, SB, YS; Writing - original draft preparation: LM; Writing - review and editing: all authors; Supervision: IB, HK; Funding acquisition: IB.

REFERENCES

1. Koboldt DC, Steinberg KM, Larson DE, Wilson RK, Mardis ER. The next-generation sequencing revolution and its impact on genomics. Cell. 2013;155(1):27-38. Epub 2013/10/01. doi: 10.1016/j.cell.2013.09.006. PubMed PMID: 24074859; PubMed Central PMCID: PMCPMC3969849.

2. Li Q, Zhao X, Zhang W, Wang L, Wang J, Xu D, et al. Reliable multiplex sequencing with rare index mis-assignment on DNB-based NGS platform. BMC Genomics. 2019;20(1):215. Epub 2019/03/15. doi: 10.1186/s12864-019-5569-5. PubMed PMID: 30866797; PubMed Central PMCID: PMCPMC6416933.

3. MacConaill LE, Burns RT, Nag A, Coleman HA, Slevin MK, Giorda K, et al. Unique, dual-indexed sequencing adapters with UMIs effectively eliminate index cross-talk and significantly improve sensitivity of massively parallel sequencing. BMC Genomics. 2018;19(1):30. Epub 2018/01/10. doi: 10.1186/s12864-017-4428-5. PubMed PMID: 29310587; PubMed Central PMCID: PMCPMC5759201.

4. Farouni R, Djambazian H, Ferri LE, Ragoussis J, Najafabadi HS. Model-based analysis of sample index hopping reveals its widespread artifacts in multiplexed single-cell RNA-sequencing. Nat Commun. 2020;11(1):2704. Epub 2020/06/03. doi: 10.1038/s41467-020-16522-z. PubMed PMID: 32483174; PubMed Central PMCID: PMCPMC7264361.

5. Griffiths JA, Richard AC, Bach K, Lun ATL, Marioni JC. Detection and removal of barcode swapping in single-cell RNA-seq data. Nat Commun. 2018;9(1):2667. Epub 2018/07/12. doi: 10.1038/s41467-018-05083-x. PubMed PMID: 29991676; PubMed Central PMCID: PMCPMC6039488.

6. Larsson AJM, Stanley G, Sinha R, Weissman IL, Sandberg R. Computational correction of index switching in multiplexed sequencing libraries. Nat Methods. 2018;15(5):305-7. Epub 2018/04/28. doi: 10.1038/nmeth.4666. PubMed PMID: 29702636.

7. Sinha R, Stanley G, Gulati GS, Ezran C, Travaglini KJ, Wei E, et al. Index switching causes "spreading-of-signal" among multiplexed samples in Illumina HiSeq 4000 DNA sequencing. BioRxiv. 2017.

8. Costello M, Fleharty M, Abreu J, Farjoun Y, Ferriera S, Holmes L, et al. Characterization and remediation of sample index swaps by non-redundant dual indexing on massively parallel sequencing platforms. BMC Genomics. 2018;19(1):332. Epub 2018/05/10. doi: 10.1186/s12864-018-4703-0. PubMed PMID: 29739332; PubMed Central PMCID: PMCPMC5941783.

9.   Owens GL, Todesco M, Drummond EBM, Yeaman S, Rieseberg LH. A novel post hoc
method for detecting index switching finds no evidence for increased switching on the Illumina
HiSeq X. BioRxiv. 2017.
10.   Illumina. Effects of index misassignment on multiplexing and downstream
analysis. 2017. Available from: https://www.illumina.com/content/dam/illumina-
marketing/documents/products/whitepapers/index-hopping-white-paper-770-2017-004.pdf.
11.   Ezpeleta J, Garcia Labari I, Villanova GV, Bulacio P, Lavista-Llanos S, Posner V, et al.
Robust and scalable barcoding for massively parallel long-read sequencing. Sci Rep.
2022;12(1):7619. Epub 20220510. doi: 10.1038/s41598-022-11656-0. PubMed PMID:
35538127; PubMed Central PMCID: PMCPMC9090787.
12.   Xiao Y, Thoresen DT, Miao L, Williams JS, Wang C, Atit RP, et al. A Cascade of Wnt, Eda,
and Shh Signaling Is Essential for Touch Dome Merkel Cell Development. PLoS Genet.
2016;12(7):e1006150. Epub 2016/07/16. doi: 10.1371/journal.pgen.1006150. PubMed PMID:
27414798; PubMed Central PMCID: PMCPMC4944988.
13.   Vassen L, Okayama T, Moroy T. Gfi1b:green fluorescent protein knock-in mice reveal a
dynamic expression pattern of Gfi1b during hematopoiesis that is largely complementary to
Gfi1. Blood. 2007;109(6):2356-64. Epub 2006/11/11. doi: 10.1182/blood-2006-06-030031.
PubMed PMID: 17095621.
14.   Yucel R, Kosan C, Heyd F, Moroy T. Gfi1:green fluorescent protein knock-in mutant
reveals differential expression and autoregulation of the growth factor independence 1 (Gfi1)
gene during lymphocyte development. J Biol Chem. 2004;279(39):40906-17. Epub 2004/07/15.
doi: 10.1074/jbc.M400808200. PubMed PMID: 15252036.
15.   Whyte WA, Orlando DA, Hnisz D, Abraham BJ, Lin CY, Kagey MH, et al. Master
transcription factors and mediator establish super-enhancers at key cell identity genes. Cell.
2013;153(2):307-19. Epub 2013/04/16. doi: 10.1016/j.cell.2013.03.035. PubMed PMID:
23582322; PubMed Central PMCID: PMCPMC3653129.
16.   Julia M, Telenti A, Rausell A. Sincell: an R/Bioconductor package for statistical
assessment of cell-state hierarchies from single-cell RNA-seq. Bioinformatics. 2015;31(20):3380-
2. Epub 2015/06/24. doi: 10.1093/bioinformatics/btv368. PubMed PMID: 26099264; PubMed
Central PMCID: PMCPMC4595899.
17.   Andoniadou CL, Matsushima D, Mousavy Gharavy SN, Signore M, Mackintosh AI,
Schaeffer M, et al. Sox2(+) stem/progenitor cells in the adult mouse pituitary support organ
homeostasis and have tumor-inducing potential. Cell Stem Cell. 2013;13(4):433-45. Epub
2013/10/08. doi: 10.1016/j.stem.2013.07.004. PubMed PMID: 24094324.
18.   Picelli S, Faridani OR, Bjorklund AK, Winberg G, Sagasser S, Sandberg R. Full-length RNA-
seq from single cells using Smart-seq2. Nat Protoc. 2014;9(1):171-81. Epub 2014/01/05. doi:
10.1038/nprot.2014.006. PubMed PMID: 24385147.
19.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic
data across different conditions, technologies, and species. Nat Biotechnol. 2018;36(5):411-20.
Epub 20180402. doi: 10.1038/nbt.4096. PubMed PMID: 29608179; PubMed Central PMCID:
PMCPMC6700744.
20.   Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics and
regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. Nat

Biotechnol. 2014;32(4):381-6. Epub 20140323. doi: 10.1038/nbt.2859. PubMed PMID:
24658644; PubMed Central PMCID: PMCPMC4122333.
21.   Qiu X, Mao Q, Tang Y, Wang L, Chawla R, Pliner HA, et al. Reversed graph embedding
resolves complex single-cell trajectories. Nat Methods. 2017;14(10):979-82. Epub 20170821.
doi: 10.1038/nmeth.4402. PubMed PMID: 28825705; PubMed Central PMCID:
PMCPMC5764547.
655

SUPPORTING INFORMATION

Figures S1-S9:

Figure S1. Skin schematic showing cells labeled by GFP in Sox2GFP/+ and Gfi1GFP/+

reporter mice.

Figure S2. Index hopping can be detected when occurring among different cell types.

Figure S3. Top DC markers genes are detected in MCs with single index hopping

potential from DCs and the detection ratio is correlated with number of different cell types

with the potential for index hopping.

Figure S4. Rank ordered scatter plot showing distribution of gene detection ratios in MCs

across all genes from mm10 genome annotated by Ensembl.

Figure S5. Index hopping correction reduces the detection of non-self marker genes in

different cell types.

Figure S6. Index hopping correction reduces the detection of non-self marker genes.

Figure S7. Assigned developmental stages of single DCs are altered by index hoping

correction.

Figure S8. Assigned developmental stages of single HCs are altered by index hoping

correction.

Figure S9. Detection ratios of top MC marker genes in HCs show that index hopping is

reduced on Hiseq2500 compared to Hiseq3000 sequencing platforms.