Iridescent structural coloration in a crested Cretaceous enantiornithine

bird from Jehol Biota

Short title: Iridescent structural coloration in a crested enantiornithine bird

Elife digestion: Three-dimensional reconstruction of melanosome organization in an

enantiornithine feather crest reveals convergent iridescent coloration.

Authors

Zhiheng Li[1†*], Jinsheng Hu[2†], Thomas A. Stidham[1,3], Mao Ye[2], Min Wang[1], Yanhong Pan[4],

Tao Zhao[5], Jingshu Li[1], Zhonghe Zhou[1,6], Julia A. Clarke[7]

1 Key Laboratory of Vertebrate Evolution and Human Origins, Institute of Vertebrate

Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing 100044,

China

2 School of Instrumentation and Optoelectronic Engineering, Beihang University, Beijing

100191, China

3 Department of Biology, Austin College, Sherman, Texas 75090, USA

4 State Key Laboratory for Mineral Deposits Research, School of Earth Sciences and

Engineering, Centre for Research and Education on Biological Evolution and Environment

and Frontiers Science Center for Critical Earth Material Cycling, Nanjing

University, Nanjing 210023, China

5 Institute of Palaeontology, Yunnan Key Laboratory of Earth System Science, Yunnan

University, Kunming 650500, China

6 College of Earth and Planetary Sciences, University of Chinese Academy of Sciences,

Beijing 100049, China

7 Department of Earth and Planetary Sciences, Jackson School of Geosciences, The

University of Texas at Austin, Texas 78712, USA

†These authors contributed equally

*Correspondence: lizhiheng@ivpp.ac.cn (Z.L.)

Abstract

A combination of sectioning and microscopy techniques, along with the application of

finite-difference-time-domain modeling on a fossil feather, results in the novel estimation

of the range of iridescent colors from the fossilized melanosome type and organization

preserved in the elongate head crest feathers of a new Cretaceous enantiornithine bird.

The densely packed rod-like melanosomes are estimated to have yielded from red to

deep blue iridescent coloration of the head feathers. The shape and density of these

melanosomes also may have further increased the feather's structural strength. This

occurrence on a likely male individual is highly suggestive of both a signaling function of

the iridescent crest, and a potential behavioral role in adjusting the angle of light incidence

to control the display of this iridescent structural coloration.

Introduction

The recognition and study of melanosomes in fossil feathers of non-avialan dinosaurs and birds has added greater complexity to the pattern of their diversity and macroevolution, demonstrating the intertwined evolution of coloration with feather functions across display, thermoregulation, and other uses (Li, Gao et al. 2012, Li, Clarke et al. 2014, Terrill and Shultz 2023). This linkage among colors, structures, and functions early in feather evolution extends even to convergence in iridescent feathers among non-avialan theropods, and it is highly suggestive of commonalities shared among stem and crown birds in terms of their use of colored pigments for camouflage, structural strength, and display (Hu, Clarke et al. 2018, Foth and Rauhut 2020). In a comparative setting with the melanosome feather coloration of living birds, workers have endeavored to reconstruct the colors of fossil feathers using quantitative means focused predominantly on melanosome geometry and density, and the results point to the fossilized melanosomes having produced patterns of black, brown, rufous, grey, and other colors (Vinther, Briggs et al. 2008, Li, Gao et al. 2010, Zhang, Kearns et al. 2010, Li, Gao et al. 2012, Peteya, Clarke et al. 2017, Benton, Dhouailly et al. 2019). Furthermore, the specific shape and arrangement of some fossilized feather melanosomes derived from both non-avialan theropod dinosaurs and birds are similar to those required for producing structurally-based iridescent feather coloration in living birds, a case of independent evolution (Stoddard and Prum 2011, Hu, Clarke et al. 2018, Pan, Li et al. 2022).

Feathers serve multiple functions in birds from comprising an insulatory epidermal layer to

forming aerodynamic flight surfaces, and as structures for the containment and display of

various pigments. Living birds exhibit diverse feather morphologies and colors, and the

significant diversity of known fossil feathers, along with their preserved pigments, points to

their past use for camouflage, as well as display and signaling (Prum 2006, Foth and

Rauhut 2020, Terrill and Shultz 2023). While the display of most colored feathers are

passive and available to any viewer, birds can behaviorally control their signaling through:

the erection of feathered ornaments such as the head crest in hoopoes (*Upupa*) or

Peacocks (*Pavo*) tails; the exposure of normally hidden brightly colored crown feathers as

in Kinglets (*Regulus*); and even the refined control of the black and iridescent coloration in

birds such as hummingbirds (Trochilidae) and birds of paradise (Paradisaeidae) (Zi, Yu et

al. 2003, Wilts, Michielsen et al. 2014, Stoddard, Eyster et al. 2020). This great variation in

structural or iridescent colors results from differences in pigments, their organization, and

other structural modifications. For example, the variety of mixed structural colors present

in a peacock feather are achieved through changes in the spacing in the keratinous matrix

and melanosome organization (Zi, Yu et al. 2003). In addition to the normal range of

spectral coloration, hummingbirds can see non-spectral colors as well, and their utility is

supported by vision perception experiments (Stoddard, Eyster et al. 2020). The interplay

of bird of paradise feather coloration and their vision similarly indicates coordination of

their mating displays with spectral properties through the avian visual system (Wilts,

Michielsen et al. 2014). Individual birds actively control the signaling functions of their

feathers with feathers or ornaments hidden/displayed or even the color presented (as

dark/black vs. vibrant/iridescent) being consciously restricted or directed to particular

viewers such as potential mates or rivals (Wilts, Michielsen et al. 2014).

Fossilized feathers typically have been evaluated within two-dimensional surface

views, limiting the understanding of their overall ultrastructure and perhaps their true

coloration. As a step forward in our reconstruction of the structure and function (color

display) of early bird feathers, we examined the unique head-crest feathers of a

well-preserved specimen (Institute of Vertebrate Paleontology and Paleoanthropology,

IVPP V26899) of the Early Cretaceous (~120 Ma) enantiornithine *Shangyang* sp. from the

Jiufotang Formation of northeastern China. This fossil preserves a diverse feather

assemblage across the body with dense contour feathers, flight remiges, and two

rachis-dominated tail feathers (Figure 1 and figure S1). Among its diverse feather

morphologies, this specimen also displays a strikingly enlarged feathered crest on the

head that uniquely extends from the caudal end of the skull to a position rostral to the

nasal bone (Figure 1). For our study, we sampled a long isolated mature feather that is

displaced from the caudal part of the crest, as well as three feather fragments extracted

from the *in-situ* position of the head crest.

In this study, we utilized serial ultrathin histological sectioning, scanning/transmission

electron microscopy (S/TEM) imaging, and finite-difference time-domain (FDTD) modeling

to investigate the potential iridescent coloration mechanisms in the elongated crest

feathers from this newly discovered enantiornithine specimen of the Jehol Biota. Differing

from previous statistical reconstructions of fossil color that measured melanosome aspect

ratios, our new approach takes the three-dimensional shape and packing patterns of

melanosomes, as well as the keratinous matrix, into consideration. With these novel

modeling approaches, a more accurate coloration estimation was produced for its crest

feather.

Results

In the new fossil, head crest feathers were selected from the caudal side of the head and

sectioned using ultrathin serial sectioning. The overall shape of the sampled crest feather

is well-preserved and comprised of elongate and relatively wide barbs. The dominant barb

length is proportionally longer than the feather barb examined from the neck and head

region of *Confuciusornis* (Foth 2012). The feather is different from the bilaterally

symmetric plumulaceous feathers of living birds, and we define it as a barb-dominant

feather, since most of the exposed feather tissue is composed of long barbs with a very

short and thick calamus. It is estimated to correlate with evolutionary stage II and IIIa of

(Prum and Brush 2002), similar to the intermediate feather type proposed as found in an

amber fossil (Perrichot, Marion et al. 2008). The new feather described here is distinct

from the amber feather in showing an asymmetric shape, radically diverging toward one

side (caudal side), and is much larger in size (Perrichot, Marion et al. 2008). There are

dense clusters inside the barbs as seen from images of both SEM and TEM.

Ultrathin histological sectioning and application of SEM and S/TEM imaging of the fossil

feather samples (see Supplementary materials) resulted in the identification of partially

interlinked melanosomes with a derived asymmetric packing pattern (Figures 2 and 3).

The long axial crack in Figure 2 may represent the surface of the barbs, resembling a

crack formed in maturation experiments (Zhao, Hu et al. 2020). Those melanosome

clusters are present and concentrated mostly within the barbs rather than in the barbules

of extant bird feathers based on similar size (Figure 2). The long axis of the densely

packed melanosomes is aligned overall parallel with the barb's long axis (Figure 2), which

has been confirmed by detailed histological data as well. The melanosomes are arranged

in an asymmetric hexagonal conformation, clearly visible in the different sections (see

Figure 3).

SEM imaging of the cross section and surface view of the sampled feather indicate the

presence of a few major barbs without barbules (figures S1 and S3-4). The thickness of

the complete feather barb with filled melanosomes (preserved) is around 10 μm, which is

much greater than the typical diameter of known barbules, but similar to barbs in extant

birds (Freyer, Wilts et al. 2021). The three-dimensionally packed melanosomes were

studied in both cross and nearly longitudinal sections within the dark colored barbs (one in

the middle of the barbs and another in the lateral side of the feather block (Figure 3a, b).

Hooks are commonly observed on the oval-shaped melanosomes in cross-sectional

views, with two dominant types identified on the dorsal and ventral sides (Figure 3c-d, red

arrows). These hooks are deflected in opposing directions, linking melanosomes from

different arrays (dorsal-ventral) together. The major axis(y) of the oval-shaped

melanosomes (mean = 283 nm) is oriented toward the left side in cross-section, while the

shorter axis(x) measures approximately 186 nm (Table S2). In oblique or near-longitudinal

sections (Figure 3e-f), the hooked structures' connections to the distal and proximal sides

of neighboring melanosomes are clearly visible (blue arrows, Figure 3f). The mean long

axis (z) of the melanosomes is approximately 1774 nm (Table S2). A similar pattern

occurs in two additional regions of interest within the same feather (figure S5). Although

the smaller proximal hooks in these sections are less distinct, this may reflect

developmental variation during melanosome formation along the feather barb.

Significantly smaller hooks were also observed in cross-sections of *in-situ* feather barbs

from the anterior side of the feather crest (figure S6). Based on these observations, we

propose that the hooked structures—particularly those on the dorsal, ventral, proximal,

and distal sides of the melanosomes—enhance the structural integrity of the barb (figure

S7). However, these features may be teratological and unique to this individual, as no

similar structures have been reported in other species. These hooks may stabilize the

stacked melanosome rods and contribute to increased barb dimensions, such as diameter

and length. The sections exhibit modified (or asymmetric) hexagonally packed

melanosomes with presence of extra hooked linkages (Figure 3c-d and e-f). The long

rod-like melanosomes are different from all other known feather melanosomes from both

extant and extinct taxa in having some extra hooks and an oblique ellipse shape in cross

and longitudinal sections of individual melanosomes (Durrer 1986, Zhang, Kearns et al.

2010). The asymmetric packing of the melanosomes (the major axis leans leftward in

cross section) played a major role in the reduction of fossilized keratinous matrix within

In the computational (FDTD) simulation, when light is incident at a small angle ($\leq 30°$, both

s- and p-polarized) in the X-Y plane, no evident reflection peak was found in the visible

spectrum, as shown in Figure 4c (e.g., when a viewer observes at this angle, the light is

almost completely absorbed, and the crest feather would have a darkened appearance).

As the incident angle varies from 40° to 70°, the wavelength corresponding to the

reflection peak shifts from 687 nm to 475 nm, and the color of the reflection peak gradually

changes from red to deep blue (inset in Figure 4c clearly shows how the resulting color

varies with the angle of incidence). From the perspective of optical modeling, the

generation of iridescent color (reflection peaks in the visible spectrum) is caused by the

constructive interference between scattered light produced by these uniquely shaped

melanosomes that are compactly arranged in the barbs (Vinther, Briggs et al. 2010).

Between blue and red, reflection peaks of green and yellow also are present,

corresponding to the wavelengths of 512 nm and 578 nm respectively. It is worth noting

that in the case of both p- and s-polarized incident light, the positions of the reflection

peaks are nearly identical, with only a slight difference in peak reflectance. Therefore, the

three-dimensional multilayer nanostructure model is somehow equivalent to

one-dimensional multilayer biphotonic crystals (Saranathan, Forster et al. 2012). In

summary, the crest feather of the Cretaceous enantiornithine shows a strikingly

angle-dependent and iridescent coloration pattern. When we consider the shrinkage effect

on melanosomes demonstrated experimentally (McNamara, Briggs et al. 2013) and

various distances between melanosomes resulting from compression effects (Pan et al.

2019; Zhao et al. 2020), our additional FDTD modeling shows a similar reflection

spectrum with a narrower bandwidth of the color peaks (Supplemental Materials).

Discussion

This study of a fossilized feather with a combination of microscopic imaging, ultrathin

sectioning, and three-dimensional modeling has provided a more accurate examination of

ancient structural coloration and uncovered unusual melanosome structures. Using

previous conventional approaches alone, we would have classified this feather crest

simply as black or close to iridescent based on its melanosome aspect ratio (Nordén,

Eliason et al. 2021). The recognition of overall iridescence links to a commonly found

extant bird color range only known previously from the platelet-shaped dinosaurian

melanosomes (Hu, Clarke et al. 2018) and a few fossil feathers (Pan, Li et al. 2022)

(Nordén, Faber et al. 2019, Nordén, Eliason et al. 2021). The independent evolution of

overall iridescence, similar to that found in living birds, was reconstructed here based on a

novel array of melanosomes in a feather barb in an extinct non-crown group bird (Nordén,

Faber et al. 2019, Yoshioka and Akiyama 2021). The occurrence of iridescent structural

coloration in an enlarged head crest on this ancient bird is similar to structures present

across many living birds that employ iridescence for intimidation, signaling, and

particularly for mate attraction or intraspecific territorial defense (Terrill and Shultz 2023).

Given that this individual bird likely was a male based on the occurrence of paired rachis

dominant tail feathers (figure S1), the vibrant coloration on a prominent feathered crest

suggests that sexual selection may have long played an important role in avialan (feather)

macroevolution (Foth 2020, Wang, O'Connor et al. 2021). The vibrant color and its

presence on an enlarged feather crest likely are interlinked, and probably indicate an

evolutionary and selective relationship among color, feather position, and feather

morphology/size.

Our discovery of new melanosome shapes and networks suggests that birds evolved

previously unrecognized patterns of color production outside of the limits present among

crown group birds. Additionally, these pigments and their dense organization may have

made the foundation of the feather darker and allowed for a more intense or vibrant

iridescent coloration, a potential relationship that is understudied among living birds. While

the interrelationship between melanin content in feathers and increases in structural

strength have long been known (Burtt 1981, Burtt Jr and Ichida 2004), perhaps the

increased density of melanosomes in this fossil feather barbs added even more structural

strength to the feathered crest of this bird in compromise for its lack of a dominant rachis.

Additionally, the interlocking hooks among the melanosomes might have aided individual

feather barb in resisting shear forces. The modified closely packed melanosome arrays

reconstructed here also are potentially advantageous in increasing barb structural

integrity and stability in the long barb ramus, and likely affected the visualized spectrum

during the display of iridescent coloration. We cannot completely rule out the increased

melanosome density in the fossil feather is the result of taphonomy. Size reduction in

melanosomes was found previously in a simulation experiment (McNamara, Briggs et al.

2013), but such a reduction is very unlikely to result in the highly organized and consistent

melanosome packing present here (McNamara, Briggs et al. 2013, Zhao, Hu et al. 2020).

To test the potential impact of melanosome shrinkage and potential taphonomic changes

to their spacing, we conducted an additional analysis (See Supplementary Materials), and

the results yield similar iridescent spectral peaks as the original setting.

Since the iridescent coloration of living birds is produced by the accumulation of

melanosomes mostly positioned in barbules, the identified fossil barbs filled with

melanosomes represents a unique phenotype related to an early feather evolutionary and

developmental stage. Given the lack of barbules in most early primitive feather forms (or

filamentary structures) in non-avialan dinosaurs and pterosaurs, the positioning of

melanosomes in barbs might be a plesiomorphic feature for ornithodirans broadly (Xu

2019, Cincotta, Nicolaï et al. 2022). With different destinations of melanosome transport

during feather formation, the pattern recovered here suggests fundamental differences in

feather pigmentation among Cretaceous stem taxa as compared to crown birds. The

melanosome-filled barbs evolved earlier in feather evolution before the later appearance

of innovations such as the presence of a stiff rachis and colorful barbules observed across

living feathers (Yu, Wu et al. 2002).

A focus on the two-dimensional examination of fossil feathers has uncovered a large

number of black or dark melanosomes across many early birds and feathered theropods

(Li, Clarke et al. 2014, Hu, Clarke et al. 2018). However, our three-dimensional study of a

fossilized Jehol feather specimen demonstrates that ultra-histological detail and spatial

relationships can be preserved at the microscopic level. Importantly, our discovery

suggests that reexamination of previously published 'black' pigmented fossil feathers

might uncover a wider occurrence of 'surprising' iridescence and structural coloration

among these early birds and non-avialan taxa. Such discoveries will aid in documenting

both independent and convergent evolutionary pathways in the history of birds. This

research further demonstrates that Jehol fossil feather tissue can be treated as

(microscopically) three dimensional specimens allowing for the collection of valuable

comparative data and application of comparative analytical techniques that facilitate a

better understanding of early feather evolution and function.

Here, we recovered iridescent coloration in fossil bird feathers that was produced by

three-dimensional melanosome packing patterns. This discovery dramatically expands

the range of fossil bird coloration space, and this diversity in feather coloration in an early

bird fits well with its forested environment, as proposed by the coeval fossil plant

community (Wu, Li et al. 2023). Tropical forest environments are accompanied by colorful

flowing plants and green leaves, and the structural color of this feathered crest is

advantageous for bird signaling as well. Forested environments (flowers, leaves, light, and

shadow) can be advantageous for birds and their signaling as more colorful feathers are

found today in tropical regions as demonstrated among extant vibrant coloration of

passeriform birds in particular (Cooney, He et al. 2022).

Materials and Methods

Fossil material

The referred specimen of the enantiornithine bird *Shangyang* sp. (IVPP V 26899) was

recovered near La-Ma-Dong Village in Jian-Chang County, within the Jiufotang Formation

(dated ~120 Ma). It can be referred to *Shangyang* by the presence of derived features in

the premaxillae, tibiotarsus, and sternum (Figure 1 and Supplementary Materials).

The feather fragment (labeled sample Figure 1) extracted from the specimen were mainly

analyzed using SEM (Scanning Electron Microscopy) and S/TEM

(Scanning/Transmission Electron Microscopy). The extracted feather sample is one of the

longest crest feathers, was displaced far from the caudal position of the skull (sample in

Figure 1), and belongs to a fully mature asymmetric barb dominant feather. In addition to

the singular feather, three fragments of other feathers were extracted from the *in-situ*

position of the head crest and sampled to validate the revealed structure (Supplementary

Materials, figure S6). A small fragment of the right femur was extracted to produce a ground-section to evaluate osteo-histological features in relation to its ontogenetic stage (figures S1 and S2).

Scanning electron microscopy (SEM)

Feather samples coated with gold were examined with a FEI ESEM Quanta 450, which is operated at the Tectonic Laboratory in the China Geological Survey in Beijing, China. The SEM was operated with about 10 mm working distance of the sample under a high voltage of 20 KV. Not only the surface of feather sample was examined, but the raw material of the cross-section exposed in the ultrathin sliced sections also were imaged to characterize an overall distribution and thickness of soft tissue in comparison with the adjacent sediment or matrix (Figure 2 and figures S3). Additionally, a FIB-SEM (Heliox 5CX Due Beam system) also was further applied to validate the melanosome packing in an extracted volume, and the imaged block is about 11 x 13 x 9μm (figure S7).

Scanning/Transmission Electron Microscopy

The crest feather fragments were embedded in epoxy resin using the SPI-PON 812 Embedding Kit (MNA, EPOK, DDSA, DMP-30)(Luft 1961) at the Key Laboratory of Vertebrate Evolution and Human Origins of the Chinese Academy of Sciences in Beijing in 2022. After a step-wised polymerization at 37°C for 12 hours, 45°C for 12 hours, and 60°C for 48 hours, the epoxy-embedded block was prepared and cut using an ultrathin slicing machine (Lecia EM UC7), equipped with a diamond knife (ultra 45°, 3 mm) for further TEM observations. The ultrathin slices were placed on copper grids with carbon

film (200 meshes in ~3 mm diameter copper gird). The thickness of a prepared ultrathin

slice is approximately ~70 nm, and the slices were consecutively placed on ~3-5 copper

grids. The removed thickness of the sample for making those ~3-5 copper grids for one

run of sectioning (serial numbered grids were made) took about ~20 μm of the prepared

pyramidal-cubic shaped block (Figures 3 and 5). The direction of slicing was either

perpendicular to the long axis of the barbs, or nearly parallel with the long axis of the barb,

for the two spots selected for cutting.

S/TEM imaging was conducted on a Thermo Scientific (FEI) Talos F200X G2 TEM at the

IVPP, Beijing during 2021-2022. The Talos F200X was operated at 200 KV with an

analytical scanning transmission electron microscope mode (S/TEM). TEM slices were

made from the isolated feather adjacent to the head and rostral portion of the skull. The

two specific micron-sized spots of the feather were targeted for examination of their

microscopic features by ultrathin slicing and examination as shown in Figure 3. For the

long contour feather, one spot close to the middle of the feather was made as a cross

section of the barb, and for the other spot, the slice was made close to the left side of the

fragment with a longitudinal slice (Figure 3). The strategic consideration of the two

directions (and spots) of cutting was to examine the melanosomes and other features in

different views in order to visualize these different features and obtain three-dimensional

data (see Supplementary Materials).

X-ray computed tomography

The fossil slab was scanned using a GE |phenix| micro-scanner with a resolution of 24.3

µm at Key Laboratory of Vertebrate Evolution and Human Origins, IVPP, Chinese

Academy of Sciences, Beijing. For the skeletal anatomy of the bird, µ-CT scans and digital

rendering were applied to aid in visualizing the skeletal features of the new specimen

(Figure 1). The digital rendering of the specimen was completed using Avizo 9.0. (Figure

1). Measurements were made using digital calipers, as well as validated in the CT data

digitally (Table S1). To achieve optimized image resolution, the field of view was adjusted,

resulting in part of the feet being excluded from the CT scan.

Technical protocol for ultrathin slicing

Prior to the work presented here, we obtained numerous practice sections from a few

samples of fossil feathers. In our prior samples, the fragmentary feather tissues were used

to improve the technician's capability and precision in sectioning exercises by comparison

with the desired direction and sectioned direction for given a fossil. There are several key

steps elaborated below for the successful repetition of practices of sectioning in the

correct alignment. First, we examined the fossil feathers with light microscopy and SEM

prior to the embedding process to make sure the barbs were aligned correctly in the epoxy

resin. Meanwhile, the small part of the extracted feather sample was aligned, using the

microscope (Lecia UC 7), into a proper position during the emending and stage trimming

process. The attention to feather direction was performed to assure that the barb major

extension direction was aligned with the long axis of the bullet mold (embedding capsule).

We also checked the position of the feather fragment in the SPI-Pon 812 during

embedding and polymerization. The sliced section should be precise enough to acquire

the desired images and information after planning and optimization of each step described

above, and before the sectioning. Of course, the ultrathin slicing practices can improve the

technician's skill and increase cutting precision which require a gradual learning process.

FDTD modeling of the feather's reflectance

To explore the coloration of the three-dimensionally packed melanosomes, Lumerical

finite-difference-time-domain (FDTD) software (Ansys Canada Ltd.) (Inan and Marshall

2011, Sullivan 2013) was used to simulate the visualization of light propagation on the

crest feather with regular arrangements of melanosomes (Figure 1). The FDTD method

allows explicit simulation of light interaction with particular-shaped melanosomes by

numerically solving Maxwell's equation in the time domain. The construction of a

three-dimensional nanostructure model is based on our characterized data (Figure 4 and

Table S2), and detailed modeling steps are included in Supplementary Materials. The

dispersion of the complex refractive index that was applied in our simulation is described

and shown in figure S9. In order to mimic real visual conditions, several variables were

considered during the simulation, including angle of light incidence in two orthogonal

incident planes (X-Y and Y-Z plane) (Figure 4b) and polarization state (s- and p-polarized).

Usually, p-polarized light is understood to have an electric field direction parallel to the

plane of incidence, and s-polarized light has the electric field oriented perpendicular to the

incident plane. Reflectance spectra are presented in Figure 4c-d.

Caveats about potential taphonomic melanosome changes

The simulated colors here are robust even when taking potential taphonomic changes into

consideration. The chance that the highly organized and anatomically oriented nanoscale

pattern formed here is the result of geological compression or pressure during fossilization

and diagenesis is quite low, and such structures are absent or unseen among known

taphonomic artifacts (McNamara, Briggs et al. 2013, Zhao, Hu et al. 2020). At present, we

cannot exclude the possibility that sheer stress could result in the observed asymmetric

packing of melanosomes. However, that pattern has not been reported in any known

experiment or geological process of fossilization including those related to Jehol bird or

non-avialan dinosaur fossils. Therefore, we suggest the pattern we obtained here

represents original biological structures. Given experimental observations of melanosome

shrinkage at high pressures and temperatures (~10% shrinkage) (McNamara, Briggs et al.

2013, Zhao, Hu et al. 2020), we also completed additional modeling with the fossil

melanosomes enlarged (110%, see Table S3) along with the fossilized keratinous cortex

and matrix. This additional modeling resulted in all of the spectral peaks present in the

original modeling being recovered. Therefore, our results are robust even when

considering known potential taphonomic artifacts, and both approaches produce similar

results with the red-to-blue iridescent coloration having been present in the crown feather

(figure S10).

References

Benton, M. J., D. Dhouailly, B. Jiang and M. McNamara (2019). "The early origin of feathers."

Trends in ecology & evolution **34**(9): 856-869.

Burtt, E. H., Jr. (1981). "The Adaptiveness of Animal Colors." BioScience **31**(10): 723-729.

Burtt Jr, E. H. and J. M. Ichida (2004). "Gloger's rule, feather-degrading bacteria, and color

variation among song sparrows." The Condor **106**(3): 681-686.

Cincotta, A., M. Nicolaï, H. B. N. Campos, M. McNamara, L. D'Alba, M. D. Shawkey, E.-E.

Kischlat, J. Yans, R. Carleer, F. Escuillié and P. Godefroit (2022). "Pterosaur melanosomes

support signalling functions for early feathers." Nature **604**(7907): 684-688.

Cooney, C. R., Y. He, Z. K. Varley, L. O. Nouri, C. J. Moody, M. D. Jardine, A. Liker, T. Székely

and G. H. Thomas (2022). "Latitudinal gradients in avian colourfulness." Nature Ecology &

Evolution **6**(5): 622-629.

Durrer, H. (1986). "Colouration." Biology of the integument: 2 vertebrates: 239-247.

Eliason, C. M. and M. D. Shawkey (2012). "A photonic heterostructure produces diverse

iridescent colours in duck wing patches." Journal of The Royal Society Interface **9**(74):

2279-2289.

Foth, C. (2012). "On the identification of feather structures in stem-line representatives of birds:

evidence from fossils and actuopalaeontology." Paläontologische Zeitschrift **86**(1): 91-102.

Foth, C. (2020). "A morphological review of the enigmatic elongated tail feathers of stem

birds." The evolution of feathers: from their origin to the present: 173-184.

Foth, C. and O. W. Rauhut (2020). "The evolution of feathers." From Their Origin to the

Present. Cham: Springer.

Freyer, P., B. D. Wilts and D. G. Stavenga (2021). "Cortex thickness is key for the colors of

iridescent starling feather barbules with a single, organized melanosome layer." <u>Frontiers in</u>

<u>Ecology and Evolution</u> **9**: 746254.

Hu, D., J. A. Clarke, C. M. Eliason, R. Qiu, Q. Li, M. D. Shawkey, C. Zhao, L. D'Alba, J. Jiang

and X. Xu (2018). "A bony-crested Jurassic dinosaur with evidence of iridescent plumage

highlights complexity in early paravian evolution." <u>Nature Communications</u> **9**(1): 217.

Inan, U. S. and R. A. Marshall (2011). <u>Numerical electromagnetics: the FDTD method</u>,

Cambridge University Press.

Li, Q., J. A. Clarke, K.-Q. Gao, C.-F. Zhou, Q. Meng, D. Li, L. D'Alba and M. D. Shawkey

(2014). "Melanosome evolution indicates a key physiological shift within feathered dinosaurs."

<u>Nature</u> **507**(7492): 350-353.

Li, Q., K.-Q. Gao, Q. Meng, J. A. Clarke, M. D. Shawkey, L. D'Alba, R. Pei, M. Ellison, M. A.

Norell and J. Vinther (2012). "Reconstruction of *Microraptor* and the evolution of iridescent

plumage." <u>Science</u> **335**(6073): 1215-1219.

Li, Q., K.-Q. Gao, J. Vinther, M. D. Shawkey, J. A. Clarke, L. D'Alba, Q. Meng, D. E. Briggs

and R. O. Prum (2010). "Plumage color patterns of an extinct dinosaur." <u>science</u> **327**(5971):

1369-1372.

Luft, J. H. (1961). "Improvements in epoxy resin embedding methods." <u>The Journal of Cell</u>

<u>Biology</u> **9**(2): 409-414.

McNamara, M. E., D. E. Briggs, P. J. Orr, D. J. Field and Z. Wang (2013). "Experimental

maturation of feathers: implications for reconstructions of fossil feather colour." <u>Biology Letters</u>

9(3): 20130184.

Nordén, K. K., C. M. Eliason and M. C. Stoddard (2021). "Evolution of brilliant iridescent

feather nanostructures." <u>eLife</u> **10**: e71179.

Nordén, K. K., J. W. Faber, F. Babarović, T. L. Stubbs, T. Selly, J. D. Schiffbauer, P. Peharec

Štefanić, G. Mayr, F. M. Smithwick and J. Vinther (2019). "Melanosome diversity and

convergence in the evolution of iridescent avian feathers — Implications for paleocolor

reconstruction." <u>Evolution</u> **73**(1): 15-27.

Pan, Y., Z. Li, M. Wang, T. Zhao, X. Wang and X. Zheng (2022). "Unambiguous evidence of

brilliant iridescent feather color from hollow melanosomes in an Early Cretaceous bird."

<u>National Science Review</u> **9**(2): nwab227.

Perrichot, V., L. Marion, D. Néraudeau, R. Vullo and P. Tafforeau (2008). "The early evolution

of feathers: fossil evidence from Cretaceous amber of France." <u>Proceedings of the Royal</u>

<u>Society of London B: Biological Sciences</u> **275**(1639): 1197-1202.

Peteya, J. A., J. A. Clarke, Q. Li, K. Q. Gao and M. D. Shawkey (2017). "The plumage and

colouration of an enantiornithine bird from the Early Cretaceous of China." <u>Palaeontology</u>

60(1): 55-71.

Prum, R. (2006). "Bird Coloration, Vol. 1, Mechanisms and Measurements."

Prum, R. O. and A. H. Brush (2002). "The evolutionary origin and diversification of feathers."

<u>The Quarterly review of biology</u> **77**(3): 261-295.

Saranathan, V., J. D. Forster, H. Noh, S. F. Liew, S. G. Mochrie, H. Cao, E. R. Dufresne and R.

O. Prum (2012). "Structure and optical function of amorphous photonic nanostructures from

avian feather barbs: a comparative small angle X-ray scattering (SAXS) analysis of 230 bird

species." <u>J R Soc Interface</u> **9**(75): 2563-2580.

Stoddard, M. C., H. N. Eyster, B. G. Hogan, D. H. Morris, E. R. Soucy and D. W. Inouye (2020).

"Wild hummingbirds discriminate nonspectral colors." Proceedings of the National Academy of

Sciences **117**(26): 15112-15122.

Stoddard, M. C. and R. O. Prum (2011). "How colorful are birds? Evolution of the avian

plumage color gamut." Behavioral Ecology **22**(5): 1042-1052.

Sullivan, D. M. (2013). Electromagnetic simulation using the FDTD method, John Wiley &

Sons.

Terrill, R. S. and A. J. Shultz (2023). "Feather function and the evolution of birds." Biological

Reviews **98**(2): 540-566.

Vinther, J., D. E. Briggs, J. Clarke, G. Mayr and R. O. Prum (2010). "Structural coloration in a

fossil feather." Biol Lett **6**(1): 128-131.

Vinther, J., D. E. Briggs, R. O. Prum and V. Saranathan (2008). "The colour of fossil feathers."

Biology Letters **4**(5): 522-525.

Wang, M., J. K. O'Connor, T. Zhao, Y. Pan, X. Zheng, X. Wang and Z. Zhou (2021). "An Early

Cretaceous enantiornithine bird with a pintail." Current Biology **31**(21): 4845-4852.e4842.

Wilts, B. D., K. Michielsen, H. De Raedt and D. G. Stavenga (2014). "Sparkling feather

reflections of a bird-of-paradise explained by finite-difference time-domain modeling."

Proceedings of the National Academy of Sciences **111**(12): 4363-4368.

Wu, Y., Z. Li, T. A. Stidham and S. Y. Foong (2023). "Phytolith evidence reveals the origin of

angiosperm leaf eating habits among ancient Cretaceous birds." The Innovation **4**(5).

Xu, X. (2019). "Feather evolution: looking up close and through deep time." Sci Bull (Beijing)

64(9): 563-564.

Yoshioka, S. and T. Akiyama (2021). Mechanisms of Feather Structural Coloration and Pattern

Formation in Birds. Pigments, Pigment Cells and Pigment Patterns. H. Hashimoto, M. Goda, R.

Futahashi, R. Kelsh and T. Akiyama. Singapore, Springer Singapore: 343-366.

Yu, M., P. Wu, R. B. Widelitz and C.-M. Chuong (2002). "The morphogenesis of feathers."

Nature **420**(6913): 308-312.

Zhang, F., S. L. Kearns, P. J. Orr, M. J. Benton, Z. Zhou, D. Johnson, X. Xu and X. Wang

(2010). "Fossilized melanosomes and the colour of Cretaceous dinosaurs and birds." Nature

463(7284): 1075-1078.

Zhao, T., J. Hu, L. Hu and Y. Pan (2020). "Experimental maturation of feathers: implications for

interpretations of fossil feathers." Palaios **35**(2): 67-76.

Zi, J., X. Yu, Y. Li, X. Hu, C. Xu, X. Wang, X. Liu and R. Fu (2003). "Coloration strategies in

peacock feathers." Proceedings of the National Academy of Sciences **100**(22): 12576-12578.

Acknowledgments

We thank: Yang JM, Ma JB, Wen XL, and Li DS for technical assistance in sampling and

ultrathin sectioning and STEM imaging; Hou YM and Yin PF for assistance with micro-CT

scanning of the specimen; Gao W for photography; and Xu Y for the artistic reconstruction.

Two reviewers are grateful in providing valuable suggestions for improving earlier version

of the manuscript.

Funding:

National Natural Science Foundation of China 42288201(ZZ), National Key Research and

Development Project of China (2024YFF0807603) (ZL) and (2024YFF0807602) (YP);

NSFC 42472047 (ZL), 37110101 (MY), and NSFC Fund for Excellent Young Scholars KZ

37110101 (MY).

533   Author contributions:

Conceptualization: ZL and ZZ

Methodology: JH, MY and ZL

Investigation: ZL, TAS, JH, YM, and JAC

Visualization: JH and ZL

Supervision: ZL and ZZ

Writing—original draft: TAS, ZL, YP, MW, JL and TZ

Writing—review & editing: ZL, JH, TAS, MY, MW, YP, TZ, JL, ZZ, and JAC

542   Competing interests: All other authors declare they have no competing interests.

544   Data and materials availability: All data are available in the main text or the supplementary

545   materials

549   Figures Legends

Figure 1 Digital rendering, photography, and line-drawing of the enantiornithine fossil specimen (*Shanyang* sp., IVPP V26899) (a), (c), (d) - dorsal; (b)- ventral view showing the major skeletal elements. The feather extracted from the crown are labeled as feather sample (c). Abbreviations: al, alular metacarpal; co, coracoid; cv, cervical vertebra; fe, femur; fi, fibula; fr frontal; fu, furcula; hac, hallucal ungual; hc, head crest; hu, humerus; hy, hyoid; il, ilium; is, ischium; lt, lateral trabecula; mcf, molted crest feather; mc ll, major metacarpal; mp, midline process; ph II-1, first digit of phalanx II; phIII-1, first digit of

phalanx III; pa, parietal; pu, pubis; py, pygostyle; r, radius; rec, rectrices; rem, remiges; rs,

rostrum; rw, right wing; sc, scapula; sk, skull; st, sternum; tb, tibiotarsus; and tmt,

tarsometatarsus.

Figure 2 Photograph (a) and simplified line-diagram (b) of the sampled isolated long

feather with a focused area from SEM imaging (c-g). As shown in the SEM images, the

long elliptical or oval-shaped melanosomes are tightly aligned with their long axis parallel

to the elongated barbs (**e, g**). The position of enlarged images (**d, e, f, and g**) is indicated

as square in (**c, d**); black arrows in (**e**) and (**g**) indicate the melanosome clusters.

Figure 3. Photograph of the crest feather (a) and prepared blocks of the feather barb (b) for thin sectioning. The sampled spots for the sliced feather are indicated as long red-colored cubes with their respective positions in (b). (c) to (f) are (STEM) images obtained from individual slices as indicated in (b). The shape and arrangement of the melanosome packing (ACHP, asymmetric compact hexagonal packing) is clearly revealed in (d); (c) and (d) are derived from the cross-section; (e) and (f) are derived from the nearly

longitudinal section. As indicated by the red arrows in (d) and (f), the dorsal and ventral

hooks are dominant in the cross-sectional images. On the other hand, hooks connecting

the proximal and distal arrays of melanosomes are better visualized in the longitudinal

section, as labeled with blue arrows in (f). Orientation of the slices were labeled (left, right,

proximal, distal) with their respective petitioning in the fossil before and during sectioning.

Figure 4. FDTD modeling results. (a) Representative multilayer melanosome model; (**b**)

Schematic drawing of the reflectance calculation setup of the feather barb. X-Y incident

plane (blue) is perpendicular, and the Y-Z incident plane (orange) is parallel to the special

melanosome longitudinal axis. θ_1 and θ_2 are incident angles in X-Y and Y-Z incident

planes, respectively. The blue dashed box (R) indicates the area where the reflection

spectra are simulated and calculated. (**c**) Angle-dependent reflectance spectra calculated

for p- and s-polarized light by FDTD modeling with X-Y incident plane. (**d**)

Angle-dependent reflectance spectra calculated for p- and s-polarized light by FDTD

modeling with Y-Z incident plane. Keratin cortex layer was considered in the modeling.

Supplementary Materials

595