Iridescent structural coloration in a crested Cretaceous

enantiornithine bird from Jehol Biota

Short title: Iridescent structural coloration in a crested enantiornithine bird

Elife digestion: Three-dimensional reconstruction of melanosome organization in an
enantiornithine feather crest reveals convergent iridescent coloration.

Authors
Zhiheng Li[1†*], Jinsheng Hu[2†], Thomas A. Stidham[1,3], Mao Ye[2], Min Wang[1], Yanhong Pan[4], Tao
Zhao[5], Jingshu Li[1], Zhonghe Zhou[1,3], Julia A. Clarke[6]

1 Key Laboratory of Vertebrate Evolution and Human Origins, Institute of Vertebrate
Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing 100044, China
2 School of Instrumentation and Optoelectronic Engineering, Beihang University, Beijing
100191, China
3 College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing
100049, China
4 State Key Laboratory for Mineral Deposits Research, School of Earth Sciences and
Engineering, Centre for Research and Education on Biological Evolution and Environment and
Frontiers Science Center for Critical Earth Material Cycling, Nanjing University, Nanjing 210023,
China
5 Institute of Paleontology, Yunnan Key Laboratory of Earth System Science, Yunnan
University, Kunming 650500, China
6 University of Texas at Austin, Department of Earth and Planetary Sciences, Jackson School
of Geosciences, The University of Texas, TX 78712, USA

†These authors contributed equally

*Correspondence: lizhiheng@ivpp.ac.cn (Z.L.)

Abstract
A combination of sectioning and microscopy techniques, along with the application of finite-
difference-time-domain modeling on a fossil feather, novelly results in the estimation of the
range of iridescent colors from fossilized melanosomes preserved in the elongate head crest
feathers of a new Cretaceous enantiornithine bird. The densely packed rod-like melanosomes
yield are estimated to have yielded from red to deep blue iridescent coloration of the head
feathers. The shape and density of these melanosome may have also further increased the
feather's structural strength. This occurrence on a likely male individual is highly suggestive
of both a signaling function of the iridescent crest, and a potential behavioral role in adjusting
the angle of light incidence to control the display of this iridescent structural coloration.

Introduction

The recognition and study of melanosomes in fossil feathers of non-avialan dinosaurs and birds has added greater complexity to the pattern of their diversity and macroevolution, demonstrating the intertwined evolution of coloration with feather functions across display, thermoregulation, and other uses (Li, Gao et al. 2012, Li, Clarke et al. 2014, Terrill and Shultz 2023). This linkage among colors, structures, and functions early in feather evolution extends even to convergence in iridescent feathers among non-avialan theropods, and it is highly suggestive of commonalities shared among stem and crown birds in terms of their use of colored pigments for camouflage, structural strength, and display(Hu, Clarke et al. 2018, Foth and Rauhut 2020). In a comparative setting with the melanosome feather coloration of living birds, workers have endeavored to reconstruct the colors of fossil feathers using quantitative means focused predominantly on melanosome geometry and density, and the results point to the fossilized   melanosomes having produced patterns of black, brown, rufous, grey, and other colors(Vinther, Briggs et al. 2008, Li, Gao et al. 2010, Zhang, Kearns et al. 2010, Li, Gao et al. 2012, Peteya, Clarke et al. 2017, Benton, Dhouailly et al. 2019). Furthermore, the specific shape and arrangement of some fossilized feather melanosomes derived from both non-avialan theropod dinosaurs and birds are similar to those required for producing structurally-based iridescent feather coloration in living birds, a case of independent evolution(Stoddard and Prum 2011, Hu, Clarke et al. 2018, Pan, Li et al. 2022). However, fossilized feathers have been evaluated typically within the two-dimensional surface views, limiting the understanding of their overall ultrastructure and perhaps their true coloration.

As a step forward in our reconstruction of the structure and function (color display) of early bird feathers, we examined the unique head-crest feathers of a well-preserved specimen (Institute of Vertebrate Paleontology and Paleoanthropology, IVPP V26899) of the Early Cretaceous (~120 Ma) enantiornithine *Shangyang* sp. from the Jiufotang Formation of northeastern China. This fossil preserves a diverse feather assemblage across the body with dense contour feathers, flight remiges, and two rachis-dominated tail feathers (Figure 1 and figure S1). Among its diverse feather morphologies, this specimen also displays an enlarged feathered crest on the head that uniquely extends from the caudal end of the skull to a position rostral to the nasal bone (Figure 1). For our study, we sampled an isolated mature feather fell from the caudal part of the crest.

Feathers serve multiple functions in birds from comprising an insulatory epidermal layer to forming aerodynamic flight surfaces, and as structures for the containment and display of various pigments. Living birds exhibit diverse feather morphologies and colors, and the significant diversity of known fossil feathers, along with their preserved pigments, points to their past use for camouflage, as well as display and signaling (Prum 2006, Foth and Rauhut 2020, Terrill and Shultz 2023). While the display of most colored feathers are passive and available to any viewer, birds can behaviorally control their signaling through: the erection of feathered ornaments such as the head crest in hoopoes (*Upupa*) or Peacocks (*Pavo*) tails; the exposure of normally hidden brightly colored crown feathers as in Kinglets (*Regulus*); and

even the refined control of the black and iridescent coloration in birds such as hummingbirds
(Trochilidae) and birds of paradise (Paradiseidae)(Zi, Yu et al. 2003, Wilts, Michielsen et al.
2014, Stoddard, Eyster et al. 2020). This great variation in structural or iridescent colors results
from differences in pigments, their organization, and other structural modifications. For
example, the variety of mixed structural colors present in a peacock feather are achieved
through changing spacing in keratin matrix and melanosome packing (Zi, Yu et al. 2003). In
addition to the normal range of spectral coloration, hummingbirds can see non-spectral
colors as well, and their utility is supported by vision perception experiments (Stoddard, Eyster
et al. 2020). The interplay of bird of paradise feather coloration and their vision similarly
indicates coordination of their mating displays with spectral properties through avian visual
system(Wilts, Michielsen et al. 2014). Individual birds actively control the signaling functions
of their feathers with feathers or ornaments hidden/displayed or even the color presented (as
dark/black vs. vibrant/iridescent) being consciously restricted or directed to particular viewers
such as potential mates or rivals (Wilts, Michielsen et al. 2014). In this study, we utilized
ultrathin histological sectioning, scanning/transmission electron microscopy (S/TEM) imaging,
and finite-difference time-domain (FDTD) modeling to investigate the potential iridescent
coloration mechanisms in the elongated crest feathers from a newly discovered
enantiornithine specimen of the Jehol Biota

Results

In the new fossil, head crest feathers were selected from caudal of the head and sectioned
using ultrathin serial section. The overall shape of the sampled crest feather is comprise of
elongate and relatively thick barbs, which may not be connected by tiny barbules. The feather
is different from the bilaterally symmetric plumulaceous feathers of living birds, and we define
it as a barb-dominant feather, since most of the exposed feather tissue is composed of long
barbs with a very short calamus. It is estimated to correlate with evolutionary stage II and IIIa
of Prum and Brush (2002), similar to the intermediate feather type proposed as found in
amber fossil (Perrichot, Marion et al. 2008). The new feather described here is distinct from
the amber feather in showing an asymmetric shape, in almost parallel diverging toward one
side (caudal side), and is much larger in size (Perrichot, Marion et al. 2008). With internal side
of the fossilized barbs, very dense melanosomes cluster is the major component as seen
images of both SEM and TEM.
115
Ultrathin histological sectioning and application of SEM and S/TEM imaging of the fossil
feather samples (see Supplementary materials) resulted in the identification of partially
interlinked melanosomes with asymmetric packing (Figures 2 and 3). The long axial crack in
Figure 2 may represent the surface of the barbs, resembling a crack formed in maturation
experiments(Zhao, Hu et al. 2020). Those melanosome clusters are present and concentrated
mostly within the barbs rather than in the barbules of extant bird feathers based on similar
size (Figure 2). The long axis of the densely packed melanosomes is aligned overall parallel
with the barb's long axis (Figure 2), which has been confirmed by detailed histological data

as well. The melanosomes are arranged in an asymmetric hexagonal conformation, clearly
visible in the different sections (see Figure 3). SEM imaging of the cross section and surface
view of the sampled feather indicate the presence of a few major barbs without barbules
(figures S2 and S3). The thickness of the complete feather barb with filled melanosomes
(preserved) is around 10 μm, which is much greater than the typical diameter of known
barbules but similar to barbs in extant birds(Freyer, Wilts et al. 2021). The three-dimensionally
packed melanosomes were studied in both cross and nearly longitudinal sections within the
dark colored barbs (one in the middle of the barbs and another in the lateral side of the
feather block (Figure 3a, b). The sections exhibit modified (or asymmetric) hexagonally packed
melanosomes with the occasional presence of extra hooked linkages (Figure 3c-f). The long
rod-like melanosomes are different from all other known feather melanosomes from both
extant and extinct taxa in having some extra hooks and an oblique ellipse shape in cross and
longitudinal sections of individual melanosomes(Durrer 1986, Zhang, Kearns et al. 2010). The
oblique packing of the melanosomes played a major role in the reduction of keratinized space
within the barbs, which may correspond to a novel structural coloration in this extinct bird.
The close packed hexagonal melanosome pattern found in extant avian feathers yield
rounded melanosome outlines in contrast to the oval-shaped melanosomes (see figure S4,
x<y) in perpendicular section here. The modified asymmetric hexagonal packing of the
melanosomes is different from the known pattern of melanosomes formed in the structure
barbules of extant birds(Eliason and Shawkey 2012), which has been seen as regular
hexagonal packing. The packing of the melanosomes in an asymmetric way, on the
microscopic level, might be related to the asymmetrical path of the barb extension direction
observed at the macroscopic level.
147
When light is incident at a small angle (≤30°, both s- and p-polarized) in the X-Y plane, no
evident reflection peak was found in the visible spectrum, as shown in Figure 4c (e.g., when a
viewer observes at this angle, the light is almost completely absorbed, and the crest feather
would have a darkened appearance). As the incident angle varies from 40° to 70°, the
wavelength corresponding to the reflection peak shifts from 687 nm to 475 nm, and the color
of the reflection peak gradually changes from red to deep blue (inset in Figure 4c clearly
shows how the resulting color varies with the angle of incidence). From the perspective of
optical modeling, the generation of iridescent color (reflection peaks in the visible spectrum)
is caused by the constructive interference between scattered light produced by these
particularly-shaped melanosomes that are compactly arranged in the barbs (Vinther, Briggs
et al. 2010). Between blue and red, reflection peaks of green and yellow also are present,
corresponding to the wavelengths of 512 nm and 578 nm respectively. It is worth noting that
in the case of both p- and s-polarized incident light, the positions of the reflection peaks are
nearly identical, with only a slight difference in peak reflectance. Therefore, the three-
dimensional multilayer nanostructure model is somehow equivalent to one-dimensional
multilayer biphotonic crystals(Saranathan, Forster et al. 2012). In summary, the crest feather
of the Cretaceous enantiornithine shows a strikingly angle-dependent and iridescent
coloration pattern. When we consider the shrinkage effect on melanosomes demonstrated
experimentally (McNamara, Briggs et al. 2013), our additional FDTD modeling shows a similar
reflection spectrum with a narrower bandwidth of the color peaks (Supplemental Materials).

Discussion

This study of a fossilized feather with a combination of microscopic imaging, ultrathin sectioning, and three-dimensional modeling has provided a more accurate examination of ancient structural coloration and uncovered unusual melanosome structures. Using previous conventional approaches alone, we would have classified this feather crest simply as black or close to iridescent based on its melanosome aspect ratio(Nordén, Eliason et al. 2021). The recognition of overall iridescence links to a commonly found extant bird color range only known previously from the platelet-shaped dinosaurian melanosomes(Hu, Clarke et al. 2018) and a few fossil feathers(Pan, Li et al. 2022, Nordén, Faber et al. 2019, Nordén, Eliason et al. 2021). The independent evolution of overall iridescence, similar to that found in living birds, was reconstructed here based on a novel array of melanosomes in a feather barb in an extinct non-crown group bird(Nordén, Faber et al. 2019, Yoshioka and Akiyama 2021). The occurrence of iridescent structural coloration in an enlarged head crest on this ancient bird is similar to structures present across many living birds that employ iridescence for intimidation, signaling, and particularly for mate attraction or intraspecific territorial defense(Terrill and Shultz 2023). Given that this individual bird likely was a male based on the occurrence of paired rachis dominant tail feathers (figure S1), the vibrant coloration on a prominent feathered crest suggests that sexual selection may have long played an important role in avialan (feather) macroevolution(Foth 2020, Wang, O'Connor et al. 2021). The vibrant color and its presence on an enlarged feather crest likely are interlinked, and probably indicate an evolutionary and selective relationship among color, feather position, and feather morphology/size.

Our discovery of new melanosome shapes and networks suggests that birds evolved previously unrecognized patterns of color production outside the limits present among crown group birds. Additionally, these pigments and their dense organization may have made the foundation of the feather darker and allowed for a more intense or vibrant iridescent coloration, a potential relationship that is understudied among living birds. While the interrelationship between melanin content in feathers and increases in structural strength have long been known(Burtt 1981, Burtt Jr and Ichida 2004), perhaps the increased density of melanosomes in this fossil feather barbs added even more structural strength to the feathered crest of this bird in compromise the lacking of a dominant rachis. The modified closely packed melanosome arrays reconstructed here also are potentially advantageous in increasing barb stability in the long barb ramus, and likely affected the visualized spectrum during the display of iridescent coloration. We cannot completely rule out the increased melanosome density in the fossil feather is the result of taphonomy. Size reduction in melanosomes was found previously in a simulation experiment(McNamara, Briggs et al. 2013), but such a reduction is highly unlikely to result in the organized and consistent melanosome packing seen here(McNamara, Briggs et al. 2013, Zhao, Hu et al. 2020). To test the potential impact of melanosome shrinkage and potential taphonomic changes to their spacing, we conducted an additional analysis (See Supplementary Materials), and the results yield similar spectral peaks as the original setting.

Since the iridescent coloration of living birds is produced by the accumulation of
melanosomes mostly positioned in barbules, the identified fossil barbs filled with
melanosomes represents a unique phenotype related to an early feather evolutionary and
developmental stage. Given the lack of barbules in most early primitive feather forms (or
filamentary structures) in non-avialan dinosaurs and pterosaurs, the positioning of
melanosomes in barbs might be a plesiomorphic feature for ornithodirans broadly(Xu 2019,
Cincotta, Nicolaï et al. 2022). With different destinations of melanosome transport during
feather formation, the pattern recovered here suggests fundamental differences in feather
pigmentation among Cretaceous stem taxa as compared to crown birds. The melanosome-
filled barbs evolved earlier in feather evolution before the later appearance of innovations
such as the presence of a stiff rachis and colorful barbules observed across living feathers(Yu,
Wu et al. 2002).
223
A focus on the two-dimensional examination of fossil feathers has uncovered a large number
of black or dark melanosomes across many early birds and feathered theropods(Li, Clarke et
al. 2014, Hu, Clarke et al. 2018). However, our three-dimensional study of a fossilized Jehol
feather specimen demonstrates that histological detail and spatial relationships can be
preserved at the microscopic level. Importantly, our discovery suggests that reexamination of
previously published 'black' pigmented fossil feathers might uncover a wider occurrence of
'surprising' iridescence and structural coloration among these early birds and non-avialan
taxa. Such discoveries will aid in documenting both independent and convergent evolutionary
pathways in the history of birds. This research further demonstrates that Jehol fossil feather
tissue can be treated as (microscopically) three dimensional specimens allowing for the
collection of valuable comparative data and application of comparative analytical techniques
that facilitate a better understanding of early feather evolution and function.
236
Here, we recovered iridescent coloration in fossil bird feathers that was produced by three-
dimensional melanosome packing patterns. This discovery dramatically expands the range of
fossil bird coloration space, and this diversity in feather coloration in an early bird fits well
with its forested environment, as proposed by the coeval fossil plant community(Wu, Li et al.
2023). Tropical forest environments are accompanied by colorful flowing plants and green
leaves, and the structural color of this feathered crest is advantageous for bird signaling as
well. Forested environments (flowers, leaves, light, and shadow) can be advantageous for
birds and their signaling as more colorful feathers are found today in tropical regions as
demonstrated among extant vibrant coloration of passeriform birds in particular(Cooney, He
et al. 2022).
247

Materials and Methods

Fossil material

250 The referred specimen of the enantiornithine bird *Shangyang* sp. (IVPP V 26899) was
251 recovered near La-Ma-Dong Village in Jian-Chang County, within the Jiufotang Formation
252 (dated ~120 Ma). It can be referred to Shangyang by the presence of derived features in the

premaxillae, tibiotarsus, and sternum (Figure 1 and Supplementary Materials).

The feather fragment (labeled sample Figure 1) extracted from the specimen were mainly analyzed using SEM (Scanning Electron Microscopy) and S/TEM (Scanning/Transmission Electron Microscopy). The extracted feather sample is one of the longest crest feathers fell fall from the caudal position of the skull (sample in Figure 1), belonging to a fully mature asymmetric barb dominant feather.

Scanning electron microscopy (SEM)

Feather samples coated with gold were examined with a FEI ESEM Quanta 450, which is operated at the Tectonic Laboratory in the China Geological Survey in Beijing, China. The SEM was operated with about 10 mm working distance of the sample under a high voltage of 20 KV. Not only the surface of feather sample was examined, but the raw material of the cross-section exposed in the ultrathin sliced sections also were imaged to characterize an overall distribution and thickness of soft tissue in comparison with the adjacent sediment or matrix (Figure 2 and figures S2 and S3).

Scanning/Transmission Electron Microscopy

The crest feather fragments were embedded in epoxy resin using the SPI-PON 812 Embedding Kit (MNA, EPOK, DDSA, DMP-30)(Luft 1961) at the Key Laboratory of Vertebrate Evolution and Human Origins of the Chinese Academy of Sciences in Beijing in 2022. After a step-wised polymerization at 37°C for 12 hours, 45°C for 12 hours, and 60°C for 48 hours, the epoxy-embedded block was prepared and cut using an ultrathin slicing machine (Lecia EM UC7), equipped with a diamond knife (ultra 45°, 3 mm) for further TEM observations. The ultrathin slices were placed on copper grids with carbon film (200 meshes in ~3 mm diameter copper gird). The thickness of a prepared ultrathin slice is approximately ~70 nm, and the slices were consecutively placed on ~3-5 copper grids. The removed thickness of the sample for making those ~3-5 copper grids for one run of sectioning (serial numbered grids were made) took about ~20 μm of the prepared pyramidal-cubic shaped block (Figures 3 and 5). The direction of slicing was either perpendicular to the long axis of the barbs, or nearly parallel with the long axis of the barb, for the two spots selected for cutting.

S/TEM imaging was conducted on a Thermo Scientific (FEI) Talos F200X G2 TEM at the IVPP, Beijing during 2021-2022. The Talos F200X was operated at 200 KV with an analytical scanning transmission electron microscope mode (S/TEM). TEM slices were made from the isolated feather adjacent to the head and rostral portion of the skull. The two specific micron-sized spots of the feather were targeted for examination of their microscopic features by ultrathin slicing and examination as shown in Figure 3. For the long contour feather, one spot close to the middle of the feather was made as a cross section of the barb, and for the other spot, the slice was made close to the left side of the fragment with a longitudinal slice (Figure 3). The strategic consideration of the two directions (and spots) of cutting was to examine the melanosomes and other features in different views in order to visualize these different features and obtain three-dimensional data (see Supplementary Materials).

X-ray computed tomography

The fossil slab was scanned using a GE |phenix| micro-scanner with a resolution of 24.3 μm at Key Laboratory of Vertebrate Evolution and Human Origins, IVPP, Chinese Academy of Sciences, Beijing. For the skeletal anatomy of the bird, μ-CT scans and digital rendering were applied to aid in visualizing the skeletal features of the new specimen (Figure 1). The digital rendering of the specimen was completed using Avizo 9.0. (Figure 1). Measurements were made using digital calipers, as well as validated in the CT data digitally (Table S1).

Technical protocol for ultrathin slicing

Prior to the work presented here, we obtained numerous practice sections from a few samples of fossil feathers. In our prior samples, the fragmentary feather tissues were used to improve the technician's capability and precision in sectioning excises by comparison with the desired direction and sectioned direction for given a fossil. There are several key steps elaborated below for the successful repetition of practices of sectioning in the correct alignment. First, we examined the fossil feathers with light microscopy and SEM prior to the embedding process to make sure the barbs were aligned correctly in the epoxy resin. Meanwhile, the small part of the extracted feather sample was aligned, using the microscope (Lecia UC 7), into a proper position during the emending and stage trimming process. The attention to feather direction was performed to assure that the barb major extension direction was aligned with the long axis of the bullet mold (embedding capsule). We also checked the position of the feather fragment in the SPI-Pon 812 during embedding and polymerization. The sliced section should be precise enough to acquire the desired images and information after planning and optimization of each step described above, and before the sectioning. Of course, the ultrathin slicing practices can improve the technician's skill and increase cutting precision which require a gradual learning process.

FDTD modeling of the feather's reflectance

To explore the coloration of the three-dimensionally packed melanosomes, Lumerical finite-difference-time-domain (FDTD) software (Ansys Canada Ltd.)(Inan and Marshall 2011, Sullivan 2013) was used to simulate the visualization of light propagation on the crest feather with regular arrangements of melanosomes (Figure 1). The FDTD method allows explicit simulation of light interaction with particular-shaped melanosomes by numerically solving Maxwell's equation in the time domain. The construction of a three-dimensional nanostructure model is based on our characterized data (Figure 4 and Table S2), and detailed modeling steps are included in Supplementary Materials. The dispersion of the complex refractive index that was applied in our simulation is described and shown in figure S5. In order to mimic real visual conditions, several variables were considered during the simulation, including angle of light incidence in two orthogonal incident planes (X-Y and Y-Z plane) (Figure 4b) and polarization state (s- and p-polarized). Usually, p-polarized light is understood to have an electric field direction parallel to the plane of incidence, and s-polarized light has the electric field oriented perpendicular to the incident plane. Reflectance spectra are presented in Figure 4c-d.

To explore the coloration of the three-dimensionally packed melanosomes, Lumerical finite-

difference-time-domain (FDTD) software (Ansys Canada Ltd.) (Inan and Marshall 2011,
Sullivan 2013) was used to simulate the visualization of light propagation on the crest feather
with regular arrangements of melanosomes (Figure 4). The FDTD method allows explicit
simulation of light interaction with particular-shaped melanosomes by numerically solving
Maxwell's equation in the time domain. The construction of a three-dimensional multilayer
nanostructure model is based on our characterized data (figure S4 and Table S2), and detailed
modeling steps are included in supplemental materials. The dispersion of the complex
refractive index that was applied in our simulation is described and shown in figure S4. In
order to mimic real eye-observing conditions, several variables are considered during the
simulation, including angle of light incidence in two orthogonal incident planes (X-Y and Y-
Z plane) (Figure 4b) and polarization state (s- and p-polarized). Usually, p-polarized light is
understood to have an electric field direction parallel to the plane of incidence, and s-
polarized light has an electric field oriented perpendicular to the incident plane. Reflectance
spectra are presented in Figure 4c-d.
352
Caveats about potential taphonomic melanosome changes
The simulated colors here are robust even when taking potential taphonomic changes into
consideration. The chance that the highly organized and anatomically oriented nanoscale
pattern formed here is the result of geological compression or pressure during fossilization
and diagenesis is quite low, and such structures are absent or unseen among known
taphonomic artifacts(McNamara, Briggs et al. 2013, Zhao, Hu et al. 2020). At present, we
cannot exclude the possibility that sheer stress could result in the observed asymmetric
packing of melanosomes. However, that pattern has not been reported in any known
experiment or geological process of fossilization including those related to Jehol bird fossils.
Therefore, we suggest the pattern we obtained here represents original biological structures.
Given experimental observations of melanosome shrinking at high pressure and temperature
(~10% shrinkage)(McNamara, Briggs et al. 2013, Zhao, Hu et al. 2020), we also completed
additional modeling with the fossil melanosomes enlarged (110%, see Table S3) with keratin
space. This additional modeling resulted in all of the spectral peaks present in the original
modeling being recovered. Therefore, our results are robust even when considering known
potential taphonomic artifacts, and both approaches produce similar results with the red-to-
blue iridescent coloration having been present in the crown feather (figure S6).
370
371
References

Benton, M. J., D. Dhouailly, B. Jiang and M. McNamara (2019). "The early origin of feathers." Trends
in ecology & evolution **34**(9): 856-869.
Burtt, E. H., Jr. (1981). "The Adaptiveness of Animal Colors." BioScience **31**(10): 723-729.
Burtt Jr, E. H. and J. M. Ichida (2004). "Gloger's rule, feather-degrading bacteria, and color variation
among song sparrows." The Condor **106**(3): 681-686.
Chuong, C.-M., R. Chodankar, R. B. Widelitz and T.-X. Jiang (2000). "Evo-devo of feathers and
scales: building complex epithelial appendages." Current opinion in genetics & development **10**(4):
449.
Cincotta, A., M. Nicolaï, H. B. N. Campos, M. McNamara, L. D'Alba, M. D. Shawkey, E.-E. Kischlat, J.

Yans, R. Carleer, F. Escuillié and P. Godefroit (2022). "Pterosaur melanosomes support signalling
functions for early feathers." Nature **604**(7907): 684-688.

Cooney, C. R., Y. He, Z. K. Varley, L. O. Nouri, C. J. Moody, M. D. Jardine, A. Liker, T. Székely and G.
H. Thomas (2022). "Latitudinal gradients in avian colourfulness." Nature Ecology & Evolution **6**(5):
622-629.

Durrer, H. (1986). "Colouration." Biology of the integument: 2 vertebrates: 239-247.

Eliason, C. M. and M. D. Shawkey (2012). "A photonic heterostructure produces diverse iridescent
colours in duck wing patches." Journal of The Royal Society Interface **9**(74): 2279-2289.

Foth, C. (2012). "On the identification of feather structures in stem-line representatives of birds:
evidence from fossils and actuopalaeontology." Paläontologische Zeitschrift **86**(1): 91-102.

Foth, C. (2020). "A morphological review of the enigmatic elongated tail feathers of stem birds."
The evolution of feathers: from their origin to the present: 173-184.

Foth, C. and O. W. Rauhut (2020). "The evolution of feathers." From Their Origin to the Present.
Cham: Springer.

Freyer, P., B. D. Wilts and D. G. Stavenga (2021). "Cortex thickness is key for the colors of iridescent
starling feather barbules with a single, organized melanosome layer." Frontiers in Ecology and
Evolution **9**: 746254.

Hu, D., J. A. Clarke, C. M. Eliason, R. Qiu, Q. Li, M. D. Shawkey, C. Zhao, L. D'Alba, J. Jiang and X. Xu
(2018). "A bony-crested Jurassic dinosaur with evidence of iridescent plumage highlights
complexity in early paravian evolution." Nature Communications **9**(1): 217.

Inan, U. S. and R. A. Marshall (2011). Numerical electromagnetics: the FDTD method, Cambridge
University Press.

Li, Q., J. A. Clarke, K.-Q. Gao, C.-F. Zhou, Q. Meng, D. Li, L. D'Alba and M. D. Shawkey (2014).
"Melanosome evolution indicates a key physiological shift within feathered dinosaurs." Nature
507(7492): 350-353.

Li, Q., K.-Q. Gao, Q. Meng, J. A. Clarke, M. D. Shawkey, L. D'Alba, R. Pei, M. Ellison, M. A. Norell
and J. Vinther (2012). "Reconstruction of Microraptor and the evolution of iridescent plumage."
Science **335**(6073): 1215-1219.

Li, Q., K.-Q. Gao, J. Vinther, M. D. Shawkey, J. A. Clarke, L. D'Alba, Q. Meng, D. E. Briggs and R. O.
Prum (2010). "Plumage color patterns of an extinct dinosaur." science **327**(5971): 1369-1372.

Luft, J. H. (1961). "Improvements in epoxy resin embedding methods." The Journal of Cell Biology
9(2): 409-414.

Maia, R., R. H. Macedo and M. D. Shawkey (2012). "Nanostructural self-assembly of iridescent
feather barbules through depletion attraction of melanosomes during keratinization." J R Soc
Interface **9**(69): 734-743.

McNamara, M. E., D. E. Briggs, P. J. Orr, D. J. Field and Z. Wang (2013). "Experimental maturation
of feathers: implications for reconstructions of fossil feather colour." Biology Letters **9**(3): 20130184.

Nordén, K. K., C. M. Eliason and M. C. Stoddard (2021). "Evolution of brilliant iridescent feather
nanostructures." eLife **10**: e71179.

Nordén, K. K., J. W. Faber, F. Babarović, T. L. Stubbs, T. Selly, J. D. Schiffbauer, P. Peharec Štefanić,
G. Mayr, F. M. Smithwick and J. Vinther (2019). "Melanosome diversity and convergence in the
evolution of iridescent avian feathers—Implications for paleocolor reconstruction." Evolution **73**(1):
15-27.

Pan, Y., Z. Li, M. Wang, T. Zhao, X. Wang and X. Zheng (2022). "Unambiguous evidence of brilliant

iridescent feather color from hollow melanosomes in an Early Cretaceous bird." <u>National Science</u>
<u>Review</u> **9**(2): nwab227.

Perrichot, V., L. Marion, D. Néraudeau, R. Vullo and P. Tafforeau (2008). "The early evolution of
feathers: fossil evidence from Cretaceous amber of France." <u>Proceedings of the Royal Society of</u>
<u>London B: Biological Sciences</u> **275**(1639): 1197-1202.

Peteya, J. A., J. A. Clarke, Q. Li, K. Q. Gao and M. D. Shawkey (2017). "The plumage and colouration
of an enantiornithine bird from the Early Cretaceous of China." <u>Palaeontology</u> **60**(1): 55-71.

Prum, R. (2006). "Bird Coloration, Vol. 1, Mechanisms and Measurements."

Prum, R. O. (2005). "Evolution of the morphological innovations of feathers." <u>Journal of</u>
<u>Experimental Zoology Part B: Molecular and Developmental Evolution</u> **304**(6): 570-579.

Prum, R. O. and A. H. Brush (2002). "The evolutionary origin and diversification of feathers." <u>The</u>
<u>Quarterly review of biology</u> **77**(3): 261-295.

Prum, R. O., E. R. Dufresne, T. Quinn and K. Waters (2009). "Development of colour-producing β
-keratin nanostructures in avian feather barbs." <u>Journal of the Royal Society Interface</u> **6**(suppl_2):
S253-S265.

Saitta, E. T., C. S. Rogers, R. A. Brooker and J. Vinther (2017). "Experimental taphonomy of keratin:
a structural analysis of early taphonomic changes." <u>Palaios</u> **32**(10): 647-657.

Saranathan, V. and C. Finet (2021). "Cellular and developmental basis of avian structural
coloration." <u>Current Opinion in Genetics & Development</u> **69**: 56-64.

Saranathan, V., J. D. Forster, H. Noh, S. F. Liew, S. G. Mochrie, H. Cao, E. R. Dufresne and R. O. Prum
(2012). "Structure and optical function of amorphous photonic nanostructures from avian feather
barbs: a comparative small angle X-ray scattering (SAXS) analysis of 230 bird species." <u>J R Soc</u>
<u>Interface</u> **9**(75): 2563-2580.

Stoddard, M. C., H. N. Eyster, B. G. Hogan, D. H. Morris, E. R. Soucy and D. W. Inouye (2020). "Wild
hummingbirds discriminate nonspectral colors." <u>Proceedings of the National Academy of Sciences</u>
117(26): 15112-15122.

Stoddard, M. C. and R. O. Prum (2011). "How colorful are birds? Evolution of the avian plumage
color gamut." <u>Behavioral Ecology</u> **22**(5): 1042-1052.

Sullivan, D. M. (2013). <u>Electromagnetic simulation using the FDTD method</u>, John Wiley & Sons.

Terrill, R. S. and A. J. Shultz (2023). "Feather function and the evolution of birds." <u>Biological Reviews</u>
98(2): 540-566.

Vinther, J., D. E. Briggs, J. Clarke, G. Mayr and R. O. Prum (2010). "Structural coloration in a fossil
feather." <u>Biol Lett</u> **6**(1): 128-131.

Vinther, J., D. E. Briggs, R. O. Prum and V. Saranathan (2008). "The colour of fossil feathers." <u>Biology</u>
<u>Letters</u> **4**(5): 522-525.

Wang, M., J. K. O'Connor, T. Zhao, Y. Pan, X. Zheng, X. Wang and Z. Zhou (2021). "An Early
Cretaceous enantiornithine bird with a pintail." <u>Current Biology</u> **31**(21): 4845-4852.e4842.

Wilts, B. D., K. Michielsen, H. De Raedt and D. G. Stavenga (2014). "Sparkling feather reflections of
a bird-of-paradise explained by finite-difference time-domain modeling." <u>Proceedings of the</u>
<u>National Academy of Sciences</u> **111**(12): 4363-4368.

Wu, Y., Z. Li, T. A. Stidham and S. Y. Foong (2023). "Phytolith evidence reveals the origin of
angiosperm leaf eating habits among ancient Cretaceous birds." <u>The Innovation</u> **4**(5).

Xu, X. (2019). "Feather evolution: looking up close and through deep time." <u>Sci Bull (Beijing)</u> **64**(9):
563-564.

Xu, X. (2020). "Filamentous integuments in nonavialan theropods and their kin: advances and future perspectives for understanding the evolution of feathers." The evolution of feathers: from their origin to the present: 67-78.

Yoshioka, S. and T. Akiyama (2021). Mechanisms of Feather Structural Coloration and Pattern Formation in Birds. Pigments, Pigment Cells and Pigment Patterns. H. Hashimoto, M. Goda, R. Futahashi, R. Kelsh and T. Akiyama. Singapore, Springer Singapore: 343-366.

Yu, M., P. Wu, R. B. Widelitz and C.-M. Chuong (2002). "The morphogenesis of feathers." Nature **420**(6913): 308-312.

Zhang, F., S. L. Kearns, P. J. Orr, M. J. Benton, Z. Zhou, D. Johnson, X. Xu and X. Wang (2010). "Fossilized melanosomes and the colour of Cretaceous dinosaurs and birds." Nature **463**(7284): 1075-1078.

Zhao, T., J. Hu, L. Hu and Y. Pan (2020). "Experimental maturation of feathers: implications for interpretations of fossil feathers." Palaios **35**(2): 67-76.

Zi, J., X. Yu, Y. Li, X. Hu, C. Xu, X. Wang, X. Liu and R. Fu (2003). "Coloration strategies in peacock feathers." Proceedings of the National Academy of Sciences **100**(22): 12576-12578.

Acknowledgments

We thank: Yang JM, Wen XL, and Li DS for technical assistance in sampling and ultrathin sectioning; Hou YM and Yin PF for assistance with micro-CT scanning of the specimen; Gao W for photography; and Xu Y for the artistic reconstruction. Two reviewers are grateful in providing valuable suggestions for improving earlier version of the manuscript.

Funding:

National Natural Science Foundation of China 42288201(ZZ) and 42472047 (ZL);
CAS Strategic Priority Program XDB26000000 CAS (ZZ, ZL)
NSFC Fund for Excellent Young Scholars KZ37110101 (MY)

Author contributions:

Conceptualization: ZL and ZZ
Methodology: JH, MY and ZL
Investigation: ZL, TAS, JH YM and JAC
Visualization: JH and ZL
Supervision: ZL and ZZ
Writing—original draft: TAS, ZL, YP, MW, JL and TZ
Writing—review & editing: ZL, JH, TAS, MY, MW, YP, TZ, JL ZZ and JAC

Competing interests: All other authors declare they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary materials

Figure 1 Digital rendering, photography, and line-drawing of the enantiornithine fossil specimen (*Shanyang* sp., IVPP V26899) (a), (c), (d) - dorsal; (b)- ventral view showing the major skeletal elements. The feather extracted from the crown are labeled as feather sample (**c**). Abbreviations: al, alular metacarpal; co, coracoid; cv, cervical; fe, femur; fi, fibula; fr frontal; fu, furcula; hac, hallucal ungual; hc, head crest; hu, humerus; hy, hyoid; il, ilium; is, ischium; lt, lateral trabecula; mcf, molted crest feather; mc II, major metacarpal; mp, midline process; ph II-1, first digit of phalanx II; phIII-1, first digit of phalanx III; pa, parietal; pu, pubis; py, pygostyle; r, radius; rec, rectrices; rem, remiges; rs, rostrum; rw, right wing; sc, scapula; sk, skull; st, sternum; tb, tibiotarsus; and tmt, tarsometatarsus.

Figure 2 Photograph (a) and simplified line-diagram (b) of the sampled isolated long
feather with a focused area from SEM imaging (c-g). As shown in the SEM images, the
long elliptical or oval-shaped melanosomes are tightly aligned with their long axis parallel to
the elongated barbs **(e, g)**. The position of enlarged images (**d, e, f, and g**) is indicated as
square in (**c, d**); black arrows in (**e**) and (**g**) indicate the melanosome clusters.

Figure 3. Photograph of the crest feather (a) and prepared blocks of the feather barb (b)
for thin sectioning. The sampled spots for the sliced feather are indicated as long red-
colored cubes with their respective positions in (**b**). (**c**) to (**f**) are (STEM) images obtained from
individual slices as indicated in (**b**). The shape and arrangement of the melanosome packing
(ACHP, asymmetric compact hexagonal packing) is clearly revealed and labeled with white
arrows in (**d**); (**c**) and (**d**) are derived from the cross-section; (**e**) and (**f**) are derived from the
nearly longitudinal section. The red triangles in (**d**) and (**f**) indicate the hooked connections
of the melanosomes.

Figure 4. FDTD modeling results. (a) Representative multilayer melanosome model; **(b)** Schematic drawing of the reflectance calculation setup of the feather barb. X-Y incident plane (blue) is perpendicular and the Y-Z incident plane (orange) is parallel to the special melanosome longitudinal axis. θ_1 and θ_2 are incident angles in X-Y and Y-Z incident planes, respectively. The blue dashed box (R) indicates the area where the reflection spectra are simulated and calculated. **(c)** Angle-dependent reflectance spectra calculated for p- and s-polarized light by FDTD modeling with X-Y incident plane. **(d)** Angle-dependent reflectance spectra calculated for p- and s-polarized light by FDTD modeling with Y-Z incident plane.

Supplementary Materials