Title: Scrutinized lipid utilization disrupts Amphotericin-B responsiveness in clinical
isolates of *Leishmania donovani*

Supratim Pradhan[1], Dhruba Dhar[1], Debolina Manna[1], Shubhangi Chakraborty[1], Arkapriya
Bhattacharyya[1], Khushi Chauhan[1], Rimi Mukherjee[2], Abhik Sen[2], Krishna Pandey[2], Soumen Das[1]
and Budhaditya Mukherjee[1]*.

[1] School of Medical Science and Technology, Indian Institute of Technology Kharagpur, India-
721302

[2] ICMR-Rajendra Memorial Research Institute of Medical Sciences, Patna, India-800007

*Corresponding author: Budhaditya Mukherjee, Email: aditya26884@gmail.com,
bmukherjee@smst.iitkgp.ac.in

Running Title: Neutral Lipid Build-Up Disrupts Amp-B responsiveness

14
15

16
17

Abstract
The management of *Leishmania donova*ni (LD), responsible for fatal visceral leishmaniasis (VL),
faces increasing challenges due to rising drug-unresponsiveness, leading to increasing treatment
failures. While hypolipidemia characterizes VL, LD, a cholesterol auxotroph, relies on host lipid
scavenging for its intracellular survival. The aggressive pathology, in terms of increased organ
parasite load, observed in hosts infected with antimony-unresponsive-LD (LD-R) as compared to
their sensitive counterparts (LD-S), highlights LD-R's heightened reliance on host lipids. Here we
report that LD-R-infection promotes fluid-phase endocytosis in the host, selectively accumulating
neutral lipids while excluding oxidized-LDL. LD-R enhances the fusion of endocytosed LDL-
vesicles with its phagolysosomal membrane and inhibits cholesterol mobilization from these
vesicles by suppressing NPC-1. This provides LD-R amastigotes with excess lipids, supporting
their rapid proliferation and membrane synthesis. This excess LDL-influx leads to an eventual
accumulation of neutral lipid droplets around LD-R amastigotes, thereby increasing their
unresponsiveness towards Amphotericin-B, a second-line amphiphilic antileishmanial. Notably, VL
patients showing relapse with Amphotericin-B treatment exhibited significantly lower serum LDL
and cholesterol than cured cases. Treatment with Aspirin, a lipid droplet blocker, reduced lipid
droplets around LD-R amastigotes, restoring Amphotericin-B responsiveness.
35

36 **Keywords:** Clinical Leishmania donovani isolates / drug-unresponsiveness / Fluid phase
37 endocytosis /vesicle fusion/LDL-cholesterol / Lipid droplets

Introduction

Antimony, which was mainstay for treating fatal visceral leishmaniasis (VL), has been withdrawn from Indian Subcontinent for more than a decade now due to emergence of resistance. Despite this, recent clinical LD isolates remain unresponsive to antimony (LD-R), indicating persistent phenotype [1]. Notably, LD-R also exhibits more aggressive pathology in clinical and experimental infection featuring increased organ parasite burden as compared to LD-S-infection [2-5]. Increased metacyclogenesis among LD-R was initially thought to be the primary factor contributing to higher infectivity, resulting in organ-parasite overload [6]. However, later studies revealed that LD-R promastigotes have a greater replicative potential than LD-S, possibly offering a selective intracellular survival advantage [7, 8]. This implies LD-R might be more adept at acquiring and metabolizing host derived nutrients within their parasitophorous vacuole (PV), facilitating their heightened intracellular proliferation. Lipids are crucial carbon sources for energy in proliferating intracellular pathogens. Several intracellular bacteria, protozoans and viruses utilizes cellular lipid droplets and plasma lipoproteins for entry, energy acquisition and biomembrane synthesis, demonstrating intricate interactions between intracellular pathogens and host lipid metabolism [9, 10]. *Leishmania* parasites are cholesterol auxotrophs [11], and need (i) host membrane cholesterol for successful attachment and infection [12]. (ii) Intracellular leishmania amastigotes replicating within membrane-bound PVs, accumulates cholesterol and lipid-droplets [13]. (iii) *Leishmania* amastigotes isolated from infected host have been shown to have increased cholesterol content in them [14], all these clearly suggest replicating *Leishmania* amastigotes might co-opt host derived lipids to facilitate their intracellular proliferation and membrane synthesis. In fact, hypocholesteremia is a hallmark of VL, with intracellular LD depleting host cholesterol and lipoproteins [15, 16]. However, the mechanism by which PV-bound LD amastigotes acquire this lipids, and how variations in lipid type, source, and utilization dynamics affect disease pathology and drug responsiveness, remains unclear.

So far, drug responsiveness in leishmaniasis has mostly been studied through in vitro selection of drug-resistant promastigotes, with limited focus on the drug susceptibility profiles of circulating clinical strains of *Leishmania* [17-19]. However, growing evidences indicates that drug responsiveness and treatment failure in VL are multifactorial, influenced by the host's immune-metabolic status [3, 20]. Importantly, recent reports have also linked emergence of drug unresponsiveness among intracellular pathogens with their ability to modulate host lipid metabolism [21, 22]. For instance, increased lipid accumulation can hinder drug penetration by trapping lipophilic drugs, fostering resistance [23-25]. VL treatment in recent years have seen a rise in treatment failure against Amphotericin-B (Amp-B), a second-line lipophilic drug [2], making it critical to investigate these failures in the context of lipid utilization during LD infection.

Using clinical antimony unresponsive LD-strains isolated from patients this study elucidates how LD-R-infection leads to an increased uptake of low-density lipoprotein (LDL) leading to eventual accumulation of neutral lipids droplets within their PV. In vitro infection of liver-resident Kupffer cells (KCs), primary sites for cholesterol synthesis and LD amastigote residence, demonstrated that LD-R infection facilitates fluid-phase endocytosis, increasing LDL-vesicle uptake in infected-KCs cells. LD-R-PV exhibits a higher propensity to fuse with endocytosed LDL-vesicles, coupled with the inhibition of cholesterol recycling from the vesicles back to the plasma membrane of Infected-MΦs. Our observation thus explains the heightened energy source and cholesterol availability for PV and daughter membrane synthesis to accommodate higher number of LD-R amastigotes observed in clinical infection. We also observed a suppressed expression of MSR-1, a scavenger receptor of ox-LDL [26], in LD-R infected murine liver, suggesting LD-R-infection limits the uptake of ox-LDL,

reducing inflammatory responses. Furthermore, our findings indicate that lipid droplet accumulation
around LD-R amastigotes contributes to Amp-B unresponsiveness, explaining recent reports of
Amp-B cross-resistance in clinical LD-R isolates from India [27, 28]. Significant low lipid profiles in
VL patients experiencing treatment failure against liposomal Amp-B, combined with complete
clearance of LD-R amastigotes using a combination treatment of Aspirin (a lipid droplet formation
blocker) and Amp-B, confirmed critical role of assimilating host lipids in developing Amp-B
unresponsiveness among rapidly replicating LD amastigotes exhibiting primary unresponsiveness
towards antimony. This study identifies sequence of events that enables LD-R amastigotes to
redirect host nutrients for their own aggravated sustenance, and how this preferential lipid utilization
might equip them with unresponsiveness against Amp-B contributing to the gradual spread of
cross-resistance among clinical LD-R isolates.
111
112　**Results**
113
1.　LD-R infection caused severe dyslipidaemia as compared to LD-S infection
Intracellular amastigotes of *Leishmania* parasites exhibit lipid auxotrophy [11], with dyslipidaemia
being a hallmark for leishmaniasis [15], including VL patients. Previously, LD amastigote load in
splenic aspirates of VL patients have been shown to exhibit an inverse correlation with serum
cholesterol level [29]. Since infection with antimony-unresponsive-LD (LD-R) has been previously
reported to exhibit higher amastigote load in spleen and liver of infected host, as compared to
infection with their responsive counterparts (LD-S) [7, 30], we wanted to investigate possible extent
of dyslipidaemia in response to LD-R and LD-S-infection and how it might affect disease pathology.
Mice infected with equal number of sorted metacyclics (**Fig S1A**) of natural clinical LD-isolates with
confirmed LD-R phenotype, MHOM/IN/09/BHU575/0 (LD-R[1]) and MHOM/IN/10/BHU814/1 (LD-
R[2]), showed a significantly lower serum levels of LDL, HDL, and cholesterol as compared to
MHOM/IN/83/AG83 (LD-S[1]) and MHOM/IN/80/DD8 (LD-S[2]), infected mice (**Figure 1A).** Liver
serves as a primary site for lipid synthesis and liver-resident MΦs (Kupffer cells, KC) are one of the
primary sites harboring LD amastigotes [31]. Therefore, we decided to test if KCs can be
established as an *in vitro* model for LD-infection to investigate the possible link of host lipid
utilization with respect to aggressive pathology shown by LD-R. Purified KCs (**Figure S1B**) infected
with equal number of sorted metacyclics for LD-S and LD-R-isolates as before displayed no
significant difference in their initial attachment to host membrane (**Figure 1B** and **Movie 1A, 1B).**
However, although initial number of intracellular LD-S and LD-R-amastigotes 4Hrs post infection
(p.i) showed no difference, a significantly higher number of LD-R amastigotes were observed in
24Hrs infected-KC (**Figure 1Ci, ii**). Interestingly, as compared to a significant and sharp rise in the
number of intracellular amastigotes between 4Hrs and 24Hrs post infected KC in response to LD-
R infection, the number of intracellular amastigotes although increased significantly did not doubled
from 24Hrs to 48Hrs p.i. suggesting exponential LD amastigote replication between 4Hrs and 24Hrs
time frame and slowing down after that (**Figure 1Ci, ii**). Moreover, it was also noticed that at 72Hrs
p.i. a notable number of infected-KC began detaching from the wells with extracellular amastigotes
probably egressing out from the infected-KCs (**Movie 2**). Thus, 24Hrs time point was selected to
conduct all further infection studies involving KCs. Parallel infection performed in peritoneal exudate
macrophages (PEC), which serves as a long-standing model for *in vitro* LD-infection, under
identical experimental conditions also resulted in similar observation with LD-R strains showing
significantly higher amastigote load at 24Hrs p.i than LD-S strains (**Figure S1Ci, ii, iii**), as
previously reported by multiple groups [1, 4, 6]. Moreover, KCs infected with LD-S or LD-R
amastigotes isolated from 28 days infected mouse spleens, also resulted in significantly higher

parasite load for LD-R (**Figure S1D**), clearly indicating that higher parasite load is a conserved LD-
R phenotype which does not depends on initial infectivity or MΦ origin. Notably, infected-KCs had
a slightly higher inherent amastigote load compared to infected-PECs for both LD-S and LD-R-
infections, with no significant difference in parasite load within the two independent LD-S or LD-R
strains (**Figure 1C, Figure S1C, S1D**). Hence, BHU575 was selected as representative LD-R and
AG83 as representative LD-S to conduct all further comparisons with the inclusion of other
representative strains as and when mentioned. Higher intracellular amastigote load in response to
LD-R-infection was further confirmed by imaging LD-S and LD-R-PV at 4Hrs and 24Hrs infected-
KCs (**Figure 1D**). Comparative analysis revealed significantly higher number of LD-R-PVs as
compared to LD-S-PVs at 24Hrs p.i, although this difference was not observed at 4Hrs p.i (**Figure
1Di, Dii**). Interestingly, 3D image reconstruction for LD-R-PV at 24Hrs p.i, presented it as a larger
fused structure as compared to smaller individual segmented appearance of LD-S-PV (**Figure
1Diii**).
160
**2.   Lipid deprivation is detrimental for LD-R while inherent hypercholesterolemia
facilitates its proliferation**
To determine if extracellular lipids are absolutely essential for proliferation of LD-R amastigotes,
LD-S and LD-R-infections were performed in high, low or lipid-free medium. Reduction in
extracellular lipids severely compromised proliferation of LD-R but not LD-S amastigotes in
infected-KCs, confirming LD-R's dependency on external lipid availability (**Figure 2A i, ii**).
Alternatively, infection performed in KCs isolated from ApoE-knockout (KO)-mice(*Apoe*$^{-/-}$) with
inherent diet induced hyperlipidemia [32], resulted in significant increase in amastigote load as
compared to wild type BL/6-infected-KCs, only for LD-R-infection, with LD-S infection resulting in
comparable amastigote load (**Figure 2B i, ii**). Previous report has suggested that LD infection in
hypercholesteremic Apoe$^{-/-}$ mice triggers a heightened inflammatory response at approximately six
weeks' post-infection compared to wild type BL/6 mice, leading to more efficient parasite clearance.
This is owing to unique membrane composition of Apoe$^{-/-}$ which rectifies leishmania mediated
defective antigen presentation at a later stage of LD infection [20]. Additionally, previous studies
have also indicated that LD infection is well-established in mice within 6 to 11 days post-infection
in murine models [33]. Thus to evaluate impact of initial lipid utilization on LD amastigote replication
in vivo, BL/6 and diet-induced hypercholesterolemic Apoe$^{-/-}$ mice were infected with GFP
expressing LD-S or LD-R promastigotes and sacrificed 11 days p.i. Interestingly, however, we
observed *Apoe*$^{-/-}$ mice infected with GFP expressing LD-S or LD-R resulted in increased splenic
parasite burden as compared to wild type BL/6 mice at an earlier time point,11 days p.i (**Figure 2C
i, ii**), This observation possibly indicates a higher initial infection propensity in *Apoe*$^{-/-}$ mice, which
might get resolved later on due to increasing T-cell activity. Our observation also showed that
splenic parasite load for LD-R is significantly higher in *Apoe*$^{-/-}$ than BL/6 (**Figure 2C ii, iii**) again
confirming that higher availability of extracellular lipids might help in increased proliferation of LD-
R amastigotes.
186
**3. Quenching of host membrane cholesterol remains consistent during initial infection with
LD-S and LD-R**
Host lipids scavenging is essential for any intracellular pathogens exhibiting lipid auxotrophy [22].
Thus, to support their aggressive division, LD-R amastigotes should require significantly higher
amount of host lipids, which in turn will allow excess supply of cholesterol for increased PV and
daughter membrane biogenesis as compared to LD-S amastigotes. *Leishmania* amastigotes are
lipid auxotrophs [11, 14], and it has been previously reported that *L. Mexicana* can acquire host

membrane cholesterol during its initial entry into host MΦs [34]. Thus, host membrane Cholesterol
was investigated as a possible source and potential mechanism by which LD-R acquires host-
derived lipids distinctively from their LD-S counterparts. However, confocal image analysis of NBD-
Cholesterol labeled KCs infected with RFP expressing LD-S or LD-R revealed that both LD-S and
LD-R can extract NBD-Cholesterol from the KC membrane during their initial infection (4Hrs p.i)
with comparable efficiency (**Figure S2A i, ii**). This observation was further confirmed by
quantification of NBD-Cholesterol between freshly invaded LD-S and LD-R parasites isolated from
infected-KC by flow cytometry (**Figure S2B i, ii**).
202
4. LD-R-infection mobilizes extracellular LDL within infected-KCs to support its intracellular
proliferation
Since the quenching of host membrane cholesterol during initial entry of LD promastigotes does
not significantly differ in LD-S and LD-R mediated infections (**Figure S2**), we explored other
potential host lipid sources for supporting LD-R's increased intracellular division and membrane
synthesis. LDL serves as the primary source of cellular lipids which helps to maintain lipid
homeostasis, and gets readily metabolized to produce bioavailable cholesterol [35]. Our previous
observation showing a significant drop in LDL levels between LD-S and LD-R-infected mice (**Figure**
1), led us to investigate LDL as a possible source which can differentially contribute in providing
excess lipids, sterols, and fatty acids to rapidly proliferating intracellular LD-R amastigotes. Live
cell imaging comparing uptake of fluorescent red-LDL in EGFP-LD-S or EGFP-LD-R-infected KCs
at 24Hrs p.i revealed a significantly higher red-LDL accumulation within EGFP-positive LD-R-
infected KCs as compared to LD-S-infection (**Figure 3Ai**). Fluorescence plate reader-based
quantification further supported this observation showing significantly higher LDL accumulation in
LD-R-infected-KCs over time, which was not observed by performing infection with killed parasites
(**Figure 3A ii**), indicating that, LDL-influx is mediated by replicating LD amastigotes. Confocal
microscopic images at 24Hrs p.i. of LD-S and LD-R-infected-KCs confirmed significantly higher
LDL co-localizing with LD-R amastigotes compared to LD-S amastigotes (**Figure 3B i, ii**).
221
5.   LD-R-infection results in a distinct metabolic shift in lipid peaks of infected host
Next, we wanted to confirm if higher LDL-uptake could result in accumulation of excess lipids,
particularly cholesterol, in LD-R-PV that would provide LD-R amastigotes with higher replicative
capabilities and associated enhanced membrane synthesis potential. Parasitophorous vacuole
fractions were isolated from LD-S and LD-R-infected KCs at 24Hrs p.i. using a previously
established protocol [36]. Following isolation, PV purity was confirmed through LAMP-1 staining
which showed a significant enrichment around isolated PV in Confocal microscopy (**Figure 3C i**).
Purity of isolated PV fractions was further confirmed by Western blot which showed an enhanced
enrichment of LAMP-1 for LD-R-PV fraction as compared to LD-S-PV fraction, while PV excluded
cellular fraction showed residual LAMP-1 expression confirming the purity of the isolated PV
fractions **(Figure 3C ii, iii).** Following isolation, protein concentration was measured for isolated PV
fractions using the Bradford assay, and PV fractions from both LD-S- and LD-R-infected KCs were
normalized accordingly. Quantitative analysis of total cholesterol using Amplex red kit (**Figure S3A**)
and thin layer chromatography (TLC) (**Figure S3B**) revealed significantly higher cholesterol content
in LD-R-PV as compared to LD-S-PV. Furthermore, GC-MS of LD-S and LD-R-PV under identical
experimental condition identified prominent peaks for cholesterol and fatty acids in LD-R-PV which
were missing for LD-S-PV (**Figure 3D i, ii**). Apart from this biochemical characterization, we also
performed biophysical analysis of lipid related peaks using non-invasive Confocal Raman
spectroscopy to capture a snapshot of lipid distribution in LD-S and LD-R-infected-KCs under native

conditions (**Figure 3E i, ii**). This analysis highlighted a distinct clustering of lipid peaks representing
cholesterol, triglycerides, and fatty acids [37] around LD-R amastigotes, contrasting with their
minimal clustering around LD-S amastigotes which resemble uninfected-KCs more closely (**Figure**
3E i). Raman Shift analysis within the biologically active region (300cm^{-1}-1800cm^{-1}) [38], further
confirmed a significant difference in relative abundance for lipid related peaks' at wavelengths: 540-
560 cm^{-1} (cholesterol), 1080-1090 cm^{-1} (phospholipids), 1270-1280 cm^{-1} (triglycerides), 1300-1340
cm^{-1} (amide-III bond), 1440-1453 cm^{-1} (fatty acids and triglycerides) [37, 39], along with olefinic
stretch of C=C double bonds at 1,655 cm^{-1} between LD-S and LD-R-infected-KC, at 24Hrs p.i.
(Figure 3E ii and S3C i, ii). We also noticed a distinct shift by plotting area under the curve from
(Right to Left) in response to LD-R-infection (**Figure S3 D**), indicative of distinctly different
lipidomes [40].
252
6. LDL-uptake in LD-infected-MΦs occurs through fluid phase endocytosis
As LD-R amastigotes heavy reliance on increased host lipid acquisition for rapid proliferation
became evident, we wanted to investigate molecular basis of this acquisition. Considering that
MΦs primarily uptake LDL via the LDL receptor (LDLr) [41], we checked its status in response to
LD-S and LD-R-infection. However, immunofluorescence analysis showed a decrease in LDLr
expression in infected-KCs harboring LD-amastigotes as compared to adjacent uninfected KCs,
regardless of infection type (**Figure 4A**), along with no significant changes in total LDLr expression
between uninfected, LD-S, and LD-R-infected-KCs, as observed from Western blot (**Figure S4A**).
This indicated that LDL-uptake by LD-infected-KCs might be receptor-independent. Our attempts
to generate siRNA mediated Knock down (KD) in KCs were unsuccessful, hence to convincingly
determine the role of LDLr in LD-infected-MΦs we generated LDLr-KD-PEC (**Figure S4B i, ii**) and
performed infection with LD-S or LD-R. Infecting LDLr-KD-PEC with LD-S or LD-R resulted in no
significant changes in intracellular amastigote numbers for both infection types as compared to
scrambled RNA control (**Figure SC**) confirming that although LD-R relies on increased LDL-influx
for heightened proliferation, this influx is LDLr-independent. Apart from LDLr, MΦs can also acquire
LDL through receptor-independent fluid phase endocytosis [42]. To confirm this possibility, we
checked the status of Cofilin, an actin modulating protein, which serves as a marker for fluid phase
endocytosis. Western blot analysis of infected-KCs revealed an increase in Cofilin levels with a
concurrent decrease in its phosphorylated form as LD-R-infection progressed, indicating active fluid
phase endocytosis (**Figure 4Bi, ii**). In contrast, LD-S-infection did not induce such changes (**Figure**
4B i, ii). Interestingly, this decrease in Cofilin phosphorylation was also observed in LD-R-infected-
PEC (**Figure S4D i, ii**), signifying that enhanced fluid phase endocytosis may be a conserved
mechanism for facilitating LDL-influx in LD-R-infected-MΦs. This enhanced fluid phase endocytosis
in response to LD-R-infection was further confirmed by structural illumination imaging (SIM), which
showed increased Cofilin turnover with dispersed distribution in LD-R-infected-KCs, leading to
enhanced F-Actin rearrangement and protrusion formation [43], which was absent for uninfected
or LD-S-infected-KCs (**Figure 4Ci, ii and Figure 4Di, ii**). Finally, we infected the KCs with GFP
expressing LD-R for 4Hrs, washed and allowed the infection to proceed in presence of fluorescent
red-LDL and Latrunculin-A (5µM), a compound which specifically inhibits fluid phase endocytosis
by inducing actin depolymerization [42]. Real-time fluorescence tracking demonstrated that
Latrunculin-A treatment not only prevented the uptake of fluorescent red-LDL but also severely
impacted intracellular proliferation of LD-R amastigotes (**Movie 3A and 3B and Figure 4E**). In
contrast, treatment with Cytochalasin-D, which alters cellular F-actin organization but does not
affect fluid phase endocytosis [42], had no effect on the intracellular proliferation of LD-R

irrespective of Cytochalasin-D concentrations (2.5µg/ml and 5µg/ml respectively) (**Figure 4F and**
Figure S4E).
289
7.   Endocytosed-LDL traffics and fuses with and LD-R-PV with increased proficiency
Typically, LDL-vesicles fuses with lysosomes upon entry into MΦs, leading to subsequent transfer
of LDL-derived cholesterol into the endoplasmic reticulum and cytoplasm [44]. As we have already
confirmed an increased influx with higher presence of endocytosed-LDL vesicles around LD-R
amastigotes along with significantly higher accumulation of cholesterol and fatty acids in LD-R-PV
as compared to LD-S-PV (**Figure 2**), we hypothesized that the mode of trafficking of endocytosed-
LDL might differ between LD-S and LD-R-infection. LD-PV are modified phagolysosomes [36, 45],
and we observed an increased convergence of LDL-vesicles with LAMP-1 positive LD-R-PV even
at 4Hrs p.i, while such convergence was not noticed in LD-S-infected-KC (**Figure 5A i, ii**).
Structured Illumination Microscopy confirmed a complete convergence of LDL-vesicles with LD-R-
PV at 24Hrs p.i, while LD-S-PV showed only partial convergence with LDL-vesicles even at this
late infection point (**Figure 5B i, ii**). Interestingly, while, Western blot analysis revealed minimal
expression of LAMP-1 in response to LD-S-infection at 4Hrs p.i, which increased considerably as
the infection progressed (**Figure 5C i**), however, a considerable amount of LAMP-1 expression
was observed for LD-R-infected-KCs from 4Hrs p.i, which also increased as infection progresses
(**Figure 5C ii**), indicating that a higher fusion of LDL-vesicles in LD-R-PV may be accompanied by
increased lysosomal biogenesis required for degrading LDL into cholesterol. Filipin staining of
EGFP-LD-S and EGFP-LD-R-infected KCs at 24Hrs p.i. revealed enhanced cholesterol
accumulation surrounding LD-R amastigotes, as indicated by a significantly higher co-localization
coefficient compared to LD-S amastigotes (**Figure 5D i, ii, iii**). Importantly, this cholesterol
enrichment was absent around dextran beads harboring LD-R amastigotes, suggesting that
cholesterol accumulation is a process specifically associated with LD-R infection. Infected-KCs
harboring both LD-R amastigotes and Fluorescent Latex Beads, showed a concentrated staining
of cholesterol around LD-R amastigotes, with no positive Cholesterol staining around internalized
latex beads similar to LD-S amastigotes (**Figure 5E**).
315
8.   LD-R-infection suppress release of metabolized cholesterol from PV
Once LDL is broken down to cholesterol by lysosomal acid hydrolases, metabolized cholesterol is
released back into cytosol from LDL-vesicles by two transmembrane proteins namely NPC-1 and
NPC-2 [44]. We suspected that LD-R must prevent or delay this process as this will limit cholesterol
accessibility to replicating LD-R amastigotes. Analysis of available *RNAseq* (INRP000146) data for
uninfected, LD-S and LD-R-infected-PEC (**Figure 6A**) revealed a suppression of *npc-1* expression
along with its transcription factor *srebp2* [46], in response to LD-R-infection. Western Blot analysis
of LD-S and LD-R-infected-KCs confirmed this low expression of SREBP2 and NPC-1 in LD-R-
infected-KC 24Hrs p.i (**Figure 6B i, ii**). More interestingly, Confocal imaging not only confirmed this
low expression of SREBP2, but even in those cases of LD-R-infected-KCs where we did observe
some residual expression of SREBP2, it was entirely restricted to cytoplasm, unlike LD-S or
uninfected-KCs where SREBP2 did get translocated in the nucleus (**Figure 6C**). Furthermore, as
SREBP2, acts as the master regulator of cholesterol biosynthesis, its downregulation and inhibition
of nuclear translocation further impacted expression of the downstream cholesterol biosynthetic
enzyme HMGCR (**Figure 6D**) specifically in LD-R-infected KCs. This again supports the *RNAseq*
expression data (**Figure 6A**) and further confirms that LD-R-infected KCs has a shutdown of *de*
novo cholesterol synthesis. For NPC-1, immunofluorescence imaging comparing adjacent
uninfected and infected-KCs harboring LD-S or LD-R, clearly confirmed unlike LD-S, LD-R

specifically suppresses NPC-1 to prevent release of LDL-derived cholesterol from fused vesicles making it accessible to amastigotes (**Figure 6E**). We confirmed this observation by performing a Total Internal Reflection Microscopy (TIRF) which clearly showed a low plasma membrane cholesterol for LD-R-infected-KCs as compared to LD-S-infected-KCs (**Figure 6Fi**) and subsequently revalidated this by quantifying the cholesterol content in isolated membrane fraction for individual experimental set as mentioned above (**Figure 6F ii**). Finally, siRNA mediated knock down of NPC-1 in PEC (due to failure of successful KD generation in KCs), resulted in significant increase in amastigote load in response to LD-R-infection, while no such change in amastigote load was observed for LD-S-infected NPC-1-KD-PECs (**Figure 6G i, ii, iii, iv**).

9. LD-R-Infection specifically excludes oxidized-LDL in infected murine liver to limit inflammation

The increased accumulation of LDL within MΦs typically triggers an inflammatory response [47]. This possessed an inherent conceptual dilemma to our observation as increased inflammatory response should be detrimental for LD-R amastigotes [48, 49]. To check this, we measured inflammatory IFN-γ by co-culturing splenic T-cells isolated from LD-S or LD-R-infected mice, with LD-S and LD-R-infected-KCs, respectively (**Figure 7A**). Surprisingly, our ELISA results showed a low IFN-γ levels in the culture supernatant of LD-R-infected-KCs even after 24Hrs of co-culture, while for LD-S-infection a significant increase was noted at this point (**Figure 7Bi**). Similarly, stimulation of LD-infected splenocytes obtained from LD-S and LD-R-infected $Apoe^{-/-}$ mice with specific soluble leishmamia antigens (SLA) of LD-S or LD-R also confirmed this restricted IFN-γ response to LD-R-infection (**Figure 7B ii)**, thus suggesting that although LD-R-infection results in increased accumulation of lipids in host MΦs, it elicits a suppressed inflammatory response as compared to LD-S. Notably, accumulation of oxidized-LDL (ox-LDL), and not its neutral form has been reported to cause inflammatory response [47, 50]. LD-S or LD-R-infected-KCs incubated with ox-LDL showed a significant purple positive staining of ox-LDL only for LD-S-infected-KCs (**Figure 7C i**), with LD-R-infected-KCs seemingly inhibiting uptake of ox-LDL. Infecting KCs with higher number of LD-S or lower number of LD-R did not alter their ox-LDL uptake capabilities, clearly suggesting LD-R-infection inherently suppress ox-LDL uptake, irrespective of the number of intracellular amastigotes (**Figure 7C. ii,**). Although, receptor of ox-LDL, MSR-1 [26], have shown a slightly lower expression in LD-R-infected-PEC in the *RNAseq* analysis (**Figure 6A**), however owing to reported dominant expression of *msr-1* in the liver [50], we checked expression of MSR-1 in the LD-S and LD-R-infected-KCs. Confocal images and Western blot analysis confirmed a significant lower expression of MSR-1 in LD-R-infected-KCs as compared to LD-S-infected-KCs (**Figure 7D i, ii**), thus explaining reduced staining for ox-LDL in response to LD-R infection. Furthermore, increased expression of *atp6vod2* transcripts, implicated in LD-PV acidification and lipid utilization [36], was also noted to be slightly higher for LD-R-infected-PECs in the *RNAseq* (**Figure 6A**). However, western blot analysis showed a slightly increased expression of ATP6V0D2 in LD-R-infected-KCs as compared to LD-S-infected-KCs (**Figure 7E**), suggesting that it might also play a role in maintaining LD-PV acidity, facilitating lipid utilization and limiting ox-LDL uptake, as previously reported for *L. mexicana* infection [36]. Finally, immunostaining of liver sections from 28 days LD-S and LD-R-infected-mice **(Figure 7F i)** confirmed a significant lower expression of MSR-1 in LD-R-infected liver with higher accumulation of neutral lipid droplets as compared to LD-S-infected murine liver **(Figure 7F ii, iii),** thus confirming LD-R specifically accumulates neutral lipids by restricting MSR-1 expression.

10.   Lipid droplets accumulation decreases Amphotericin-B responsiveness in LD-R amastigotes

Increased LDL-uptake has been reported to cause high lipid droplets formation [51, 52], and lipid droplets formation has been reported to play critical role in intracellular proliferation of several pathogens, including LD-amastigotes [53-55]. As we have already observed significant accumulation of lipid droplets in LD-R-infected murine liver (**Figure 7**), we further investigated its role in response to LD-S and LD-R-infection. Lipid droplets staining of LD-S and LD-R-infected-KCs revealed significantly higher lipid droplets accumulation around LD-R amastigotes (**Figure S5A** and **Movie 4).** For better visualization, we expanded [56] EGFP-LD-R-infected-KCs and again performed lipid droplets staining on them which confirmed assimilation of neutral lipid into LD-R amastigotes (**Figure 8A**). This was further confirmed by performing TEM in LD-R-infected-KC which showed Lipid droplets accumulating around LD-R amastigotes **(Figure 8B)**. Lipid accumulation has been previously reported to cause Amphotericin-B unresponsiveness among pathogens by hindering drug penetration [23]. Interestingly, we noted that two LD-R-strains (BHU575 and BHU814) which we have included in this study also shows reduced responsiveness against Amp-B, a second-line-antileishmanial [57], and we validate this by determining their EC50 against Amp-B in LD-infected-KCs, which showed an EC50 of 0.39 μM and 0.31 μM for LD-R[1] and LD-R[2] amastigotes respectively, as compared to EC50 of 0.03 μM for LD-S[1] and 0.032 μM LD-S[2]. We suspected that significant lipid droplets accumulation around LD-R amastigotes might be responsible for their reduced susceptibility towards Amp-B by limiting drug accessibility. To test this possibility, we performed independent infection with two LD-R strains in low lipid and high lipid medium and access their EC50 against Amp-B. Low lipid medium results in significant drop in lipid droplets accumulation within LD-R-infected-KCs (**Figure S5B**) and also resulted in enhanced susceptibility for both the LD-R strains against Amp-B with EC50 reduced to 0.39 μM to 0.16 μM for LD-R[1] and 0.31 μM to 0.06 μM for LD-R[2] as compared to high lipid medium (**Figure 8C i, ii**). Interestingly, infection with lab-generated Amphotericin-B-unresponsive LD promastigotes (LD-S[Amp-B-R]) without primary unresponsiveness towards antimony did not result in any significant lipid droplets accumulation around LD-R amastigotes (**Figure S5A**) and neither showed any significant alteration in EC50 between low (0.09 μM) and high lipid medium (0.11 μM) (**Figure 8C iii**). This observation clearly suggests that this high LDL-influx along with higher lipid droplet accumulation responsible for increased unresponsiveness of LD-R isolates towards Amp-B is a specific attribute of LD isolates with primary unresponsiveness against antimony and may vary significantly from lab generated Amp-B unresponsive strains. Previously, Aspirin (Asp), a lipid droplet blocker has been reported to block lipid droplet formation in *T.cruzi* and LD-infected-MΦs [53, 58]. We also observed a drastic drop in lipid droplets formation in LD-R-infected-KCs treated with Aspirin as compared untreated LD-R-infected-KCs (**Figure S5C).** Interestingly, while treatment with Aspirin alone did not significantly reduce the amastigote load in LD-R-infected-KCs, combining Aspirin with previously determined low dose of Amp-B (0.1 μM), completely restored Amp-B responsiveness against intracellular LD-R amastigotes (**Figure 8E, F**). This suggests that the enhanced uptake of LDL leading to lipid droplet accumulation might be a primarily responsible for increasing Amp-B unresponsiveness among clinical LD-R isolates. Importantly, this observation is consistent with findings from human VL patients from the endemic region of Bihar, India, where those who relapsed after Amp-B treatment had significantly lower serum LDL and cholesterol levels compared to Amp-B treatment responders (**Figure 8G and Table 2**).

Discussion

Our findings revolve around understanding how antimony-resistant *Leishmania donovani* (LD-R) has evolved as a persistent strain by adopting to a strategy of better utilization of host derived lipids to sustain their own aggressive intracellular proliferation. Intracellular pathogens often alter the host's immune-metabolic state for better survival [59, 60]. Our observations pointed out that high organ parasite load linked with LD-R-infection does not rely only on initial infectivity as commonly believed to date [7], rather is indicative of a selective higher intracellular proliferation in host. We observed LD-R amastigotes scrutinize and access specific type of host lipoproteins to support their aggressive proliferation as seen in clinical infection (**Figure 1, S1**) [15]. Previously increased amastigote load in VL patients has been inversely linked with their serum cholesterol level [29], and we also observed that mice infected with LD-R isolates had significantly lower HDL, LDL, and cholesterol compared to LD-S-infected mice (**Figure 1**).

LD amastigotes are lipid auxtrophs [11, 14], and this severe dyslipidemia in response to LD-R-infection can be logically interpreted in terms of LD-R's increase need to support their aggravated intracellular proliferation in the infected-host. In liver-resident macrophages (Kupffer cells), primary sites for LD infection and host cholesterol synthesis, we observed significantly higher number of LD-R amastigotes and parasitophorous vacuoles formation (PV), compared to LD-S-infection, indicating LD-R's better capacity to utilize host cholesterol (**Figure 1, S1**). Also, in hyperlipidemic Apoe$^{-/-}$ mice, LD-R-infection led to organ parasite overload compared to LD-S-infection, confirming LD-R's increased dependence on extracellular lipid sources to sustain its aggressive proliferation (**Figure 2**). Thus, next we focused on investigating LD-R's ability to differentially regulate host's immune-metabolic status. Previous reports suggested that LD with altered drug-responsiveness can modulate their host differentially resulting in different disease outcome [3, 61]. Our results confirmed that while both LD-R and LD-S promastigotes initially quench cholesterol from the host plasma membrane similarly (**Figure S2**), LD-R amastigotes accumulate significantly more cholesterol and fatty acids in their PV by specifically promoting increased endocytosis of LDL as the infection progresses (**Figure 3, S3**). This possibly also explains the similar initial attachment and infectivity of LD-S and LD-R promastigotes with the host MΦs again indicating towards their ability to infect their host equally (**Figure 1, S1, Movie 1**).

Interestingly, this LD-R-induced LDL-endocytosis is independent of the host's LDL receptor and relies on fluid-phase endocytosis, involving significant actin cytoskeleton reorientation in LD-R-infected-MΦs (**Figure 4, S4**). It is worth mentioning that LD-R-infection has been reported to induce high levels of IL-10 [3], which is linked with enhanced fluid-phase endocytosis [42].

Importantly, we also noted an increased convergence of endocytosed LDL-vesicles with lysosomal associated marker protein (LAMP-1), along with higher LAMP-1 expression around LD-R amastigotes as compared to LD-S ones (**Figure 5**). Lysosomal acid hydrolases metabolize LDL into bioavailable cholesterol, and higher fusion of lysosomal vesicles with LD-R-PV explains the significant Filipin-cholesterol staining around LD-R amastigotes (**Figure 5**). These findings not only confirm that endocytosed-LDL can be better utilized by LD-R amastigotes but probably also indicates towards a rapid maturation of LD-R-PV due to increased LDL-vesicle fusion [13, 36].

Although available RNAseq analysis (**Figure 6**) did not support this increased expression of l*amp-1* in the transcript level, it did reflect a notable upregulation of vesicular fusion protein (VSP) *vamp8* and *stx1a* in response to LD-R-infection. How, LD infection can regulate LAMP-1 expression, and the role of VSPs in LDL-vesicle fusion with LD-R-PV is worthy of further investigation. It is possible and has been earlier reported that LD infection can regulate host proteins expression through post transcriptional and post translational modifications [62-64]. It is tempting to speculate that LD-R amastigote might be promoting an increased lysosomal biogenesis through any such mechanism

to increase supply of bioavailable cholesterol through action of lysosomal acid hydrolases on LDL.
Additionally, RNAseq analysis also showed suppressed expression of *npc-1* in LD-R-infected-MΦs.
NPC-1 is crucial for mobilizing vesicular cholesterol to the cytoplasm and plasma membrane [44],
and its suppressed expression correlates with higher cholesterol availability for aggressive LD-R
proliferation (**Figure 6**).
Besides LDL endocytosis, *de novo* cholesterol synthesis could also serve as a lipid source for
intracellular LD amastigotes [65]. Interestingly, LD-R-infected macrophages exhibited suppressed
expression of HMGCR and SREBP2, the key regulators of de novo cholesterol biosynthesis.
Additionally, SREBP2 showed minimal nuclear translocation in LD-R-infected macrophages
compared to LD-S-infected cells, (**Figure 6**), consistent with reports of a shutdown in de novo
cholesterol synthesis in MΦs with high endocytosed LDL accumulation [66]. Notably, SREBP2
regulates NPC-1 expression [46] and has been and has been shown to get activated in response
to LD-S infection [66]. Further RNA sequencing data also revealed a significant downregulation of
hmgcs (3-hydroxy-3-methylglutaryl-CoA synthase) in LD-R infected PECs as compared to LD-S
infection. Downregulation of HMGCS which catalyzes the condensation of acetyl-CoA with
acetoacetyl-CoA to form 3-hydroxy-3-methylglutaryl-CoA (HMG-CoA), which serves as an
intermediate in both cholesterol biosynthesis and ketogenesis further supports our observation that
LD-R-infected PECs preferentially rely on endocytosed low-density lipoprotein (LDL)-derived
cholesterol rather than *de novo* synthesized cholesterol to support their metabolic needs.
Surprisingly, LD-R-Infected-MΦs showed minimal inflammatory response despite high lipid
accumulation, due to their ability to successfully exclude uptake of oxidized LDL (ox-LDL). Low
expression of MSR-1, a scavenger receptor of ox-LDL, in the liver of LD-R-infected mice explained
this observation, leading to the accumulation of neutral lipid droplets around LD-R amastigotes
(**Figure 7, S5, Movie 4**). Expansion Microscopy (ExM) [56] along with TEM confirmed significant
assimilation of these neutral lipids into LD-R amastigotes (**Figure 8**). Several intracellular
pathogens like *Mycobacterium*, *Toxoplasma*, HCV have been reported to transfer stored
triacylglycerols and sterol esters from lipid droplets to support their proliferation [10, 67, 68]. While
the understanding of lipid uptake and metabolizing genes in LD is limited, a recent genome-wide
CRISPR screen in *L. Mexicana* [69], have identified such potential candidates "essential" for
parasite survival. Close proximity lipid droplets with LD-R amastigotes makes it tempting to
speculate existence of lipid droplets contact sites [13], or possible copy number variations in these
lipid metabolizing genes [70], providing survival advantages to LD-R. This speculation is further
supported by the fact that although clinical LD-R isolates with neutral lipid droplet accumulation in
infected MΦs displayed increased unresponsiveness to Amphotericin-B (Amp-B), lab-generated
Amp-B unresponsive promastigotes failed to do so in infected MΦs (**Figure 8**). Fact that LD-R
amastigotes regained their responsiveness towards Amp-B in low lipid medium or in presence of
Aspirin [53], further confirms that eliminating lipid droplets around LD-R amastigotes might be a
weapon of choice to combat treatment failure against Amp-B [27, 28, 71]. Our results thus suggest
that natural clinical LD-R-isolates have developed an inherent adaptability to gradually develop
cross-resistance towards Amp-B, probably as an additional consequence of their ability to
accumulate increased host derived lipids as a rapidly proliferating intracellular amastigote
(represented graphically in **Figure 9**). Our work explains increasing Amp-B treatment failure among
VL patients [27, 28, 71] and correlates with low serum LDL and cholesterol levels in patients
relapsing after Amp-B treatment (**Figure 9, Table 2**).
518
519
520

Methods

Reagents

Laurdan (Merck), photoRED-LDL (Thermo Fisher Scientific), NBD Cholesterol (Sigma), Nile red (Sigma-Aldrich), Oil red O (Thermo Fisher Scientific), Filipin (Sigma-Aldrich), Collagenase (Gibco) were purchased. Tissue culture plasticware were purchased from NUNC (Roskilde, Denmark). Invitrogen Amplex-Red assay kit was purchased from Thermo Fisher Scientific. Primary antibodies required for the experiments were anti-ATP6V0D2 (ABclonal), anti-MSR1 (ABclonal), anti-LAMP1 (Thermo Fisher Scientific), anti-SREBP2 (Thermo Fisher Scientific), anti-NPC1 (Santa Cruz Biotechnology), anti-HMGCR(Santa Cruz Biotechnology), anti-LDLr (Santa Cruz Biotechnology), anti-KMP-11(Invitrogen), goat anti-mouse IgG Alexa Fluor 488 (Thermo Fisher Scientific), goat anti-mouse IgG Alexa Fluor 594 (Thermo Fisher Scientific), goat anti-Rabbit IgG-HRP (Thermo Fisher Scientific), goat anti-Mouse IgG-HRP (Thermo Fisher Scientific), β actin (Cell Signalling Technology), CD11b (Cell Signalling Technology), Coffilin (Cell Signalling Technology), and phospho-Coffilin (Cell Signalling Technology). RhodaminePhalloidin reagent was purchased from Thermo Fisher Scientific and ELISA Kits for IFN-γ were purchased from R&D Systems, Aspirin generic composition, SAG (Albert David)

Animals

BALB/c, C57BL/6 and *Apoe*$^{-/-}$ mice were maintained and bred under pathogen-free conditions with food supplements and filtered water. All experiments were conducted in accordance to Institutional Animal Ethics Committee (IAEC), IIT Kharagpur guidelines India (IAEC/BI/1 55/2021) and approved by National Regulatory Guidelines issued by Committee for the Purpose of Supervision of Experiments on Animals, Ministry of Environment and Forest, Government of India.

Ethical statement

In this study, all procedures involving human participants were conducted in accordance with the ethical standards of the 1964 Helsinki Declaration and approval of the "Institutional Human Ethical Committee" of ICMR-Rajendra Memorial Research Institute of Medical Sciences, Patna, India (Approval No: RMRI/EC/20/2020). Following information about the potential risks, benefits, and the investigational nature of the study consent from all study participants was obtained in the informed consent form.

Parasite Cultures and Maintenance

Leishmania donovani (LD) isolates used in this study are natural clinical isolates obtained from the patients MHOM/IN/09/BHU569/0, MHOM/IN/2009/BHU575/0, MHOM/IN/10/BHU814/1, MHOM/IN/83/AG83, MHOM/IN/09/BHU777/0, MHOM/IN/80/DD8 [2, 3], and were maintained in BALB/c mice as per institute biosafety guidelines (BT/BS/17/27/97-PID). Susceptibility of each of these LD isolates against antimony were checked and confirmed against previously reported EC50 value for antimony before they were included in the study (This can be reported in a common table, Table 1). Lab generated Amp-B resistant strain (DD8-AmpB) strain was kind from Dr. Arun Kumar Haldar (CSIR-CDRI). Amastigotes were isolated by macerating the spleen of 21 days infected mice under sterile conditions, and subsequently transformed into promastigotes by maintaining in M199 (Gibco) supplemented with 5% (v/v) heat-inactivated fetal bovine serum (FBS) (Gibco) and 1% (v/v) Penicillin Streptomycin solution (Gibco) under drug selection.

Cell Culture and Infection

Peritoneal exudate cells (PECs) were harvested from BALB/c mice as mentioned in already published protocol, with slight modifications [72]. After 48Hrs of 4% starch intraperitoneal injection, PECs were isolated and plated on 35-mm tissue culture petri dishes, sterile 22-mm square coverslips in 35-mm petri dishes or 24-well plates at a density of 1×10^6, 1.5×10^5 or 1.5×10^5, respectively, all in RPMI 1640 medium (Gibco) supplemented with 10% (v/v) heat-inactivated FBS, 1% (v/v) Penicillin Streptomycin solution. The cells were allowed to adhere for 48Hrs at 37°C in 5% CO_2 before the infection process.

In vivo **infection**

For *in vivo* infections, healthy female C57BL/6 mice or *Apoe$^{-/-}$* mice (4-6 weeks old and 20-25 g each) were injected via tail vein route with 1×10^7 GFP-expressing *L. donovani* stationary phase metacyclic promastigotes (suspended in 100 µl 1X PBS) as reported previously [73]. After infection mice were kept in atherogenic diet till sacrifice. Each group contained five mice that were sacrificed 11 days' post-infection (p.i). Splenocytes were isolated and GFP positive population was measured through flow cytometry. Each experiment was performed in triplicate.

Isolation of Kupffer cells

Kupffer cells (KCs) were harvested from mice livers following a previously published protocol, with slight modifications [74]. Wild-type BALB/c mice were anesthetized, and warm PBS containing 0.05% collagenase IV was used for liver perfusion. The livers were swiftly excised and minced into small fragments on an ice bath. Tissue digestion was performed using a 0.1% Type IV Collagenase solution in RPMI 1640 for 30 minutes at 37°C with intermittent shaking for every 10 minutes. Following digestion, KCs were separated from hepatocytes and other cellular components through sequential centrifugation steps and plated on well plates, as required, for 4Hrs in RPMI 1640 media supplemented with 10% (v/v) Penicillin Streptomycin and 10% (v/v) FBS. After 24Hrs of resting at 37°C in 5% CO_2, the infection studies commenced. To assess the purity of the isolated KCs by flow cytometry, KCs were plated on a 6 well plate. After 4Hrs, KCs were scraped out, centrifuged at 300g for 5 minutes and stained with monoclonal anti-F4/80 PE conjugated antibody (BD Biosciences). The KCs were then washed and fixed with 2% paraformaldehyde for 5 minutes, analyzed using a flow cytometer (ARIA FACS DIVA), and further processed using FLOW JO software for data analysis. Purified KCs were infected with stationary phase sorted metacyclic LD promastigotes as mentioned previously. PECs and KCs are collectively designated as MΦs (macrophages).

Sorting of Metacyclic LD and infection in MΦs

Metacyclic parasites were sorted through Beckman Coulter CytoFLEX cell sorter, as per manufacturer's protocol. The LD promastigotes were washed and resuspended in PBS before being subjected to light scatter analysis using flow cytometry. Sorted population were checked with forward scatter (FSC) versus side scatter (SSC) and dot plot was generated to represent the acquisition of 10,000 events using a FACS Aria II system. Required number of sorted metacyclic LD were used to perform subsequent MΦ and mice infections. MΦs were infected with stationary-phase Flow sorted metacyclic LD promastigotes at a MOI 1:10 ratio (in some cases 1:5 and 1:20) for 4Hrs which was considered as the 0th point of the infection. MΦs was then washed to remove any extracellular loosely attached parasites and incubated further as per experimental requirement.

Intracellular amastigote load determination

To enumerate intracellular amastigote load, resting stage PECs and KCs on glass coverslips were infected with sorted metacyclic LD promastigotes of various strains at a ratio of 1:10, as described in previous protocols [3, 75]. Following a 4Hrs incubation, the PECs or KCs were washed and subsequently cultured for specific durations. Coverslips were retrieved after specific time points, rinsed with PBS, fixed using methanol, and stained with Giemsa. These prepared coverslips were then affixed onto glass slides and examined utilizing a light microscope (Olympus). Parasite quantification was conducted by counting amastigotes per 100 macrophages. Image processing related to Figure 1A and Figure S1 D was performed using Fiji by separating the spectral overlap between the light and dark stained region to Giemsa stained MΦs. Next, for better visualization, pseudo colours were assigned to represent nucleus (white) and cytoplasm (red) with a black background. Colour separated images were then merged showing LD within PV.

Raman Spectroscopy of LD-infected-KCs

KCs were plated on Raman-grade stainless steel plate and kept for 24Hrs in a humidified environment at 37°C with 5% CO_2, followed by infection with LD as already described. At 24Hrs p.i., the KCs were washed using PBS, fixed with 2% paraformaldehyde and proceeded to perform Raman spectroscopy (Witec, alpha300R). The Raman grade stainless steel plates were positioned on the microscope stage and focused with a 100x air objective and measurements were performed using a 532 nm laser source producing a laser spot of approximately 0.3 μm with 20mW laser power. Spectral measurements were obtained through the Project FIVE interface (Witec, Germany) after setting the grating to 600 gratings/mm. The spectral resolution was set to 2.87 cm^{-1}. The Raman spectra were taken with integration time set to 5 seconds, and ten accumulations were selected after confirming its effectiveness on the KCs being studied. Baseline correction and other spectral denoising procedures were carried out using MATLAB 2017b (MathWorks), and the processed spectra were analysed as described preciously [37, 39]. For confocal Raman spectroscopy, spectral data were acquired from individual cells at 1000× magnification using a 100 × 100 μm scanning area, following previously established specifications. After spectral acquisition, distinct Raman shifts corresponding to specific biomolecular signatures were extracted for further analysis. These included: Cholesterol (535–545 cm^{-1}), Nuclear components (780–790 cm^{-1}), Lipid structures (1262–1272 cm^{-1}), Fatty acids (1436–1446 cm^{-1}) Following spectral extraction, pseudo-color mapping was applied to highlight the spatial distribution of each biomolecular component within the cell. These processed spectral images are presented in Figure 3D1, where the first four panels illustrate the individual biomolecular distributions. A merged composite image was then generated to visualize the co-localizaton of these biomolecules within the cellular microenvironment, with the final panel specifically representing the spatial distribution of key biomolecules.

Quantification of membrane cholesterol with Amplex Red assay kit

The KC membrane preparation was conducted following the method described previously [20], with slight modifications. KCs were ruptured by repeated freezing and thawing cycles followed by probe sonicating for 4 cycles with 30secs burst and 30sec gap. The resulting homogenate was then centrifuged at 900g for 10mins at 4°C. The supernatant was filtered through a nylon mesh (100 μm), while the pellet was discarded. The filtered supernatant was centrifuged at 100,000g for 1hr at 4°C, and resulting pellet, representing native membrane, was suspended in a buffer containing 50 mM Tris (Ph 7.4), 1 mM EDTA, 0.24 mM PMSF, and 10 mM iodoacetamide. The protein content of the membrane was quantified using Bradford reagent (Biorad), following manufacturer's protocol.

The total cholesterol content was determined using an Amplex Red reagent kit [76], with the results expressed in moles of cholesterol per gram of protein.

Parasitophorous vacuole (PV) Isolation and cholesterol measurement
Parasitophorous vacuoles (PV) were isolated using a previously outlined protocol with slight modifications [77]. 10^7 KCs were seeded in a 100 mm plate and allowed to adhere for 24Hrs. Following this infection was performed with Leishmania donovani (LD) for 24Hrs, the infected KCs were then harvested by gentle scraping and lysed through five successive passages through an insulin needle to ensure membrane disruption while preserving organelle integrity. The lysate was centrifuged at 200 × g for 10mins at 4°C to remove intact cells and large debris. The resulting supernatant was carefully collected and subjected to a discontinuous sucrose density gradient (60%, 40%, and 20%). The gradient was centrifuged at 700 × g for 25mins at 4°C to facilitate organelle separation. The interphase between the 40% and 60% sucrose layers, enriched with PVs, was carefully collected and subjected to a final centrifugation step at 12,000 × g for 25mins at 4°C. The supernatant was discarded, and the resulting pellet was enriched for purified parasitophorous vacuoles, suitable for downstream biochemical and molecular analyses. Cholesterol and protein contents in PV were determined by an Amplex Red assay kit and Bradford assay, respectively. Resulting data were represented as micrograms of cholesterol per microgram of protein.

GC-MS analysis of LD-S and LD-R-PV
Following a 24Hrs infection period, KCs were harvested, washed with phosphate-buffered saline (PBS), and pelleted. Subsequent to this, PV isolation was carried out using the previously described protocol [36]. After PV isolation Bradford assay was carried out for normalizing the protein concentration. The resulting equal volume of PV pellet was suspended in 20 ml of dichloromethane: methanol (2:1, vol/vol) and incubated at 4°C for 24hours. After centrifugation (11,000 g, 1 hour, 4°C), the supernatant was checked through thin layer chromatography (TLC) and subsequently evaporated under vacuum. The residue and pellet were saponified with 30% potassium hydroxide (KOH) in methanol at 80°C for 2 hours. Sterols were extracted with n-hexane, evaporated, and dissolved in dichloromethane. A portion of the clear yellow sterol solution was treated with N, O-bis(trimethylsilyl)trifluoroacetamide (BSTFA) and heated at 80°C for 1 hour to form trimethylsilyl (TMS) ethers. Gas chromatography/mass spectrometry (GC/MS) analysis was performed using a Varian model 3400 chromatograph equipped with DB5 columns (methyl-phenylsiloxane ratio, 95/5; dimensions, 30 m by 0.25 mm). Helium was used as the gas carrier (1 ml/min). The column temperature was maintained at 270°C, with the injector and detector set at 300°C. A linear gradient from 150 to 180°C at 10°C/min was used for methyl esters, with MS conditions set at 280°C, 70 eV, and 2.2 kV[78].

Cytokine measurement
Supernatants from KCs infected with stationary-phase LD co-culture with T cells, isolated from the mice infected with LD. For detection of IFN-γ from Apoe-/-, splenocytes isolated from LD infected mice and cultured with SLA. IFN-γ levels in the supernatants were measured using a sandwich ELISA Kit (R&D Systems) as per manufacturer's protocol. Detection limit for these kits were 8pg/ml for IFN-γ. Cytokine levels were determined by measuring the OD at 450 nm using a MultiSkan FC microplate photometer (Thermo Fisher Scientific). Data was demonstrated as mean ± SD of all five individual experiments.

Western Blot Analysis

Lysates of LD-infected PECs and KCs were prepared, and western blotting was performed for different proteins with endogenous control β-actin (Cell Signaling Technology). Blots were probed with specific antibodies. Binding of secondary HRP-labeled goat anti-rabbit or goat anti-mouse Abs was analyzed using SuperSignalR West Pico or West Dura Chemiluminescent substrate (Pierce).

Confocal and Structural Illumination Microscopy (SIM)

PECs and KCs were cultured on 13-mm diameter glass coverslips at a density of $2*10^5$, and infected with LD as mentioned previously. After varying time intervals, infected PECs or KCs were rinsed with PBS, fixed with 2% paraformaldehyde, and permeabilized using 0.2% Triton/PBS for 20mins on an orbital shaker. Non-specific binding was blocked with 2% BSA in 0.2% Triton/PBS for another 20mins on an orbital shaker. The PECs and KCs were then incubated with primary antibodies for 1Hr, followed by washing and staining with IgG Alexa Fluor 594 or IgG Alexa Fluor 488, as required. Mounting was performed using Fluromount-G-DAPI (Thermo Scientific). Images were captured using Olympus confocal microscopy (FV3000) and super-resolution microscopy ZEISS (SIM) with a 63x objective. 3D image visualization processed through Fiji 3D viewer and 3D project plugin. Live cell microscopy was performed using ZEISS phase contrast microscope and ZEISS Apotome. MΦs were plated on confocal dish and incubated with photoRED-LDL for live cell imaging of LDL uptake (**Movie 3)**.

Image processing and analysis

Image processing and analysis were conducted using Fiji (ImageJ). For optimal visualization, Giemsa-stained macrophages (MΦs) were represented in grayscale to enhance contrast and structural clarity. To improve the distinction of different fluorescent signals, pseudo-colors were assigned to fluorescence images, ensuring better differentiation between various cellular components. For colocalization analysis (Figures 3, Figure 5, Figure 6, and Figure S2), we utilized the RGB profile plot plugin in ImageJ, which allows for the precise assessment of signal overlap by generating fluorescence intensity profiles across selected regions of interest. This approach provided quantitative insights into the spatial relationship between labelled molecules within infected cells. Additionally, for analysing the distribution of cofilin in Figure 4, the ImageJ surface plot plugin was employed. This tool enabled three-dimensional visualization of fluorescence intensity variations, facilitating a more detailed examination of cofilin localization and its potential reorganization in response to infection.

Lipid droplet staining

LD-infected-KCs were first fixed using 2% paraformaldehyde (PFA) and subsequently permeabilized with 0.25% Triton X-100. Coverslips were then treated with Nile red (Sigma, diluted 1:100) in PBS for at least 1Hr to stain lipids, followed by DNA staining with DAPI. The coverslips were mounted onto glass slides using Fluoromount-G and visualized with confocal microscope (LEICA STELLARIS 5). 3D stacks were acquired to visualize distribution of lipid droplets. Images were processed and analysed, and stacks were merged to reconstruct 3D images using Fiji.

Ultrastructure expansion microscopy

Ultrastructure Expansion Microscopy (U-ExM) was conducted on LD-infected-MΦs following established experimental protocols [56]. Protein crosslinking was initiated by immersing the coverslip in a solution containing 1.4% formaldehyde, 2% acrylamide, and PBS for 5Hrs at 37°C. Monomer solution (19% sodium acrylate, 10% acrylamide, and 0.1% N,N'-methylenbisacrylamide

in PBS) was mixed with 10% TEMED and 10% APS solution on ice and added to 6-well pates.
Gelation was performed on ice, with cell-plated coverslip placed face down on the gelation solution
in a humid chamber for 5 minutes at 37°C for 1 hour. The gel and coverslip were together
transferred to a 6-well plate filled with denaturation buffer (200 mM SDS, 200 mM NaCl, and 50
mM Tris pH 9.0) and incubated face up for 15mins at room temperature under agitation, followed
by incubation at 95°C for 1.5Hrs. The first expansion phase involved incubating the gel three times
in ddH2O for 30mins each. Previously mentioned IFA steps were adopted for antibody staining.
Images were captured using an Olympus confocal microscope (FV3000) equipped with a 63X 1.4
NA oil objective and Small Volume Computational Clearing mode was used to obtain deconvolved
images. 3D stacks were acquired and analyzed using Fiji software.

Transmission electron microscopy (TEM)
768 For transmission electron microscopy, PECs infected with the LD-R strain were fixed after 24 hours
769 of infection using 2.5% glutaraldehyde, an electron microscopy (EM) grade fixative. The cells were
770 then processed according to previously established protocols[58]. Following this, the samples were
771 examined using a JEM-F200 transmission electron microscope, allowing for lipid droplet imaging
772 and detailed visualization of the cellular structures.

In-vitro cholesterol trafficking assay
KCs was first treated with NBD-cholesterol for 16Hrs, allowing the fluorescent cholesterol analog
to integrate into their membranes. Next, KCs were washed with RPMI 1640 and cultured for again
2Hrs in the same medium to eliminate any residual fluorescence from the background. These
labelled KCs were infected with LD for 4Hrs and washed to remove loosely attached MΦs. The
KCs were then either immediately fixed with 2% paraformaldehyde (PFA) or left untreated to serve
as a control group. Finally, the fixed MΦs were examined under a confocal microscope.

Flow cytometry to determine NBD-cholesterol
783 KCs were first treated with NBD-cholesterol as previously described. LD parasites were isolated
784 after 2Hrs of attachment to the KCs. The quenching of NBD-cholesterol was measured by flow
785 cytometric analysis, using an unstained control.

Total internal reflection fluorescence (TIRF) microscopy
788 To visualize membrane fluidity, TIRF microscopy was performed [79]. Briefly, KCs were plated on
789 coverslips, and after 24Hrs of LD-infection, KCs were fixed with 2% PFA. KCs were stained with
790 Laurdan dye and visualized with Olympus cell TIRF microscopic system. Representative images
791 of three independent biological experiments have been provided.

RNAseq analysis
794 RNA-Sequencing was performed utilizing external service by Bionivid Project for LD-S and LD-R-
795 infected-PECs, following 24Hrs of infection keeping uninfected MΦs as control (Bioproject. Average
796 expression of differentially expressed genes related to lipid metabolism between LD-S and LD-R-
797 infected-KC is represented as heat map. RAW sequencing reads have been deposited to Indian
798 Nucleotide Data Archive (https://inda.rcb.ac.in/home) with accession number INRP000146.

Lipid profiling from serum
801 Lipid profile analysis of patient serum was conducted using lipid estimation kits employing the
802 enzymatic method for the determination of cholesterol, triglycerides, HDL, and LDL. For BALB/c

803 mice, blood was collected following anesthesia and prior to euthanasia. The blood samples were
804 maintained at 4°C to allow clotting, after which the serum was isolated**.**
805
Evaluation of EC50
KCs infected with LD parasites were treated with SAG or Amp-B 24Hrs post-infection and incubated
for an additional 24Hrs. At the experimental endpoints, coverslips were washed with PBS, air-dried,
and fixed with 100% ice-cold methanol. Subsequently, the coverslips were stained with Giemsa
solution and examined microscopically. To quantify the number of amastigotes, one hundred KCs
per coverslip were counted. [2]. The average of three untreated coverslips was taken as 100%
control, and the percentage inhibition of infected KCs in treated cultures was calculated. EC50
values for each isolate were estimated against each drug [2]. Results were also expressed as
EC50.
815
Aspirin treatment and amastigote load determination
KCs were infected with LD parasites and, after 4Hrs, treated with 5μM Aspirin for a duration of 24
hours. Following this treatment, the KCs were washed and subsequently treated with Amp-B at a
specific dose for an additional 24 hours. Coverslips were then washed with PBS, air-dried, and
fixed with 100% ice-cold methanol. After fixation, the coverslips were stained with Giemsa solution
and examined microscopically. To quantify the number of amastigotes, one hundred KCs per
coverslip were counted. [2]
823
siRNA mediated Knock down (KD)
For all siRNA transfections, Lipofectamine® RNAiMAX Reagent (Life Technologies, 13778100)
specifically designed for knockdown assays in primary cells was used according to the
manufacturer's instructions with slight modifications. PECs were seeded into 24-well plates at a
density of $1x10^5$ per well, and incubated at 37°C with 5% CO2. The transfection complex,
comprising (1μl Lipofectamine® RNAiMAX and 50μl Opti MEM) and (1 μl siRNA and 50μl Opti
MEM) mixed together directly added to the incubated PECs. Gene silencing was checked by IFA
and by Western blot as mentioned previously.
832
Blood samples collection
Blood samples were collected from a total of 22 individuals spanning a diverse age range (8 to 70
years) by RMRI, Bihar, India. Among these, nine samples were obtained from healthy individuals
residing in endemic regions to serve as controls. Serum was isolated from each blood sample
through centrifugation, and the lipid profile was subsequently analysed using a specialized
diagnostic kit (Coral Clinical System) following the manufacturer's protocol.
839
Statistical analysis
All statistical analyses were performed using GraphPad Prism 8 on raw datasets to ensure robust
and reproducible results. For datasets involving comparisons across multiple conditions, one-way
or two-way analysis of variance (ANOVA) was conducted, followed by Tukey's post hoc test to
assess pairwise differences while controlling for multiple comparisons. A 95% confidence interval
(CI) was applied to determine the statistical reliability of the observed differences. For non-
parametric comparisons across multiple groups, Wilcoxon rank-sum tests were employed,
maintaining a 95% confidence interval, which is particularly useful for analysing skewed data
distributions. In cases where only two groups were compared, Student's t-test was used to
determine statistical significance, ensuring an accurate assessment of mean differences. All

quantitative data are represented as mean ± standard error of the mean (SEM) to illustrate variability within experimental replicates. Statistical significance was determined at P ≤ 0.05. Notation for significance levels: *P ≤ 0.05; **P ≤ 0.001; ***P ≤ 0.0001.

Acknowledgments
SP acknowledges CSIR-UGC (2020-2021) and PMRF (2021-2024) fellowship. DD is a recipient of GATE fellowship. DM is a recipient of CSIR fellowship, SC is a recipient of GATE fellowship. Authors would like to acknowledge IIR lab, SMST, IITKGP for their help in kupffer cell isolation and Flow cytometric analysis, Confocal Microscope facility under DST-FIST grant conferred on the School of Bioscience, File no. SR/FST/LS-I/2019/595(C), and Confocal facility of Department of Biotechnology, DST-FIST, Govt. of India. Authors would like to thank Prof. Syamal Roy, IICB, India for providing all the LD strains used in this study. Prof. Yasuyuki Goto, The University of Tokyo, Japan for allowing experiments related to live cell video microscopy in his lab. Authors would like to thank Dr. Praphulla Chandra Shukla for providing Apoe$^{-/-}$ mice. Authors acknowledges CRF SMST, and CRF IITKGP. Authors would like to thanks Dr. Moumita Bhaumik, NICED, India for helping in membrane isolation experiments. Authors would like to thank all the interns of the IDI lab, SMST who help in various experiments related to this project.

Author Contributions: SP conceptualization, experiments, analysis, MS preparation. DD performed Raman spectroscopy experiments. DM generated all the GFP, RFP expressing LD strains, SC perform experiments related to EC50, AB performed Raman spectroscopy experiments and analysis, KC did inhibitor study related to Cytochalasin-D, Latrunculin -A, RM serum collection from VL patients, AS related serum lipid profile, formal analysis of the MS, KP related to serum lipid profile, formal analysis of the MS, SD helped in formal analysis of Raman spectroscopic experiments. B.M conceptualization, analysis, conceived and directed the project and prepared the MS with input from all the authors.

Competing Interest Statement: The authors declare no competing interests.

Data availability: RAW sequencing reads have been deposited to Indian Nucleotide Data Archive (https://inda.rcb.ac.in/home) with accession number INRP000146.

References

1.   Dumetz, F., et al., *Molecular preadaptation to antimony resistance in Leishmania donovani on the Indian subcontinent.* MSphere, 2018. **3**(2): p. 10.1128/msphere. 00548-17.

2.   Mukhopadhyay, R., et al., *Characterisation of antimony-resistant Leishmania donovani isolates: biochemical and biophysical studies and interaction with host cells.* International journal for parasitology, 2011. **41**(13-14): p. 1311-1321.

3.   Mukherjee, B., et al., *Antimony-resistant but not antimony-sensitive Leishmania donovani up-regulates host IL-10 to overexpress multidrug-resistant protein 1.* Proceedings of the National Academy of Sciences, 2013. **110**(7): p. E575-E582.

4.   Verma, S., et al., *Quantification of parasite load in clinical samples of leishmaniasis patients: IL-10 level correlates with parasite load in visceral leishmaniasis.* PloS one, 2010. **5**(4): p. e10107.

5.   Thakur, C., D. Mitra, and S. Narayan, *Skewing of cytokine profiles towards T helper cell type 2 response in visceral leishmaniasis patients unresponsive to sodium antimony gluconate.* Transactions of the Royal Society of Tropical Medicine and Hygiene, 2003. **97**(4): p. 409-412.

6.   Ouakad, M., et al., *Increased metacyclogenesis of antimony-resistant Leishmania donovani clinical lines.* Parasitology, 2011. **138**(11): p. 1392-1399.

7.   Vanaerschot, M., et al., *Antimonial resistance in Leishmania donovani is associated with increased in vivo parasite burden.* PloS one, 2011. **6**(8): p. e23120.

8.   Mukherjee, B., et al., *Probing the molecular mechanism of aggressive infection by antimony resistant Leishmania donovani.* Cytokine, 2021. **145**: p. 155245.

9.   Feingold, K.R. and C. Grunfeld, *Lipids: a key player in the battle between the host and microorganisms1.* Journal of lipid research, 2012. **53**(12): p. 2487-2489.

10.   Sonda, S. and A.B. Hehl, *Lipid biology of Apicomplexa: perspectives for new drug targets, particularly for Toxoplasma gondii.* TRENDS in Parasitology, 2006. **22**(1): p. 41-47.

11.   Goad, L.J., G.G. Holz Jr, and D.H. Beach, *Sterols of Leishmania species, implications for biosynthesis.* Molecular and biochemical parasitology, 1984. **10**(2): p. 161-170.

12.   Pucadyil, T.J., et al., *Cholesterol is required for Leishmania donovani infection: implications in leishmaniasis.* Molecular and biochemical parasitology, 2004. **133**(2): p. 145-152.

13.   Sood, C., et al., *Leishmania highjack host lipid body for its proliferation in macrophages by overexpressing host Rab18 and TRAPPC9 by downregulating miR-1914-3p expression.* Plos Pathogens, 2024. **20**(2): p. e1012024.

14.   Bouazizi-Ben Messaoud, H., et al., *Changes in lipid and fatty acid composition during intramacrophagic transformation of Leishmania donovani complex promastigotes into amastigotes.* Lipids, 2017. **52**: p. 433-441.

15.   Lal, C.S., et al., *Hypocholesterolemia and increased triglyceride in pediatric visceral leishmaniasis.* 2007.

16.   Pradhan, S., et al., *Linking membrane fluidity with defective antigen presentation in leishmaniasis.* Parasite immunology, 2021. **43**(7): p. e12835.

17.   Douanne, N., et al., *Leishmania parasites exchange drug-resistance genes through extracellular vesicles.* Cell reports, 2022. **40**(3).

18. Légaré, D., et al., *ABC proteins of Leishmania.* Journal of bioenergetics and biomembranes,
2001. **33**: p. 469-474.
19. Gourbal, B., et al., *Drug uptake and modulation of drug resistance in Leishmania by an*
aquaglyceroporin. Journal of Biological Chemistry, 2004. **279**(30): p. 31010-31017.
20. Ghosh, J., et al., *Hyperlipidemia offers protection against Leishmania donovani infection:*
role of membrane cholesterol [S]. Journal of lipid research, 2012. **53**(12): p. 2560-2572.
21. Sen, S., et al., *Restoration of IFNγR subunit assembly, IFNγ signaling and parasite*
clearance in Leishmania donovani infected macrophages: role of membrane cholesterol.
PLoS pathogens, 2011. **7**(9): p. e1002229.
22. Toledo, A. and J.L. Benach, *Hijacking and use of host lipids by intracellular pathogens.*
Virulence mechanisms of bacterial pathogens, 2016: p. 635-666.
23. Chang, W., et al., *Trapping toxins within lipid droplets is a resistance mechanism in fungi.*
Scientific reports, 2015. **5**(1): p. 15133.
24. Cao, Y., *Adipocyte and lipid metabolism in cancer drug resistance.* The Journal of clinical
investigation, 2019. **129**(8): p. 3006-3017.
25. Day, N.J., P. Santucci, and M.G. Gutierrez, *Host cell environments and antibiotic efficacy*
in tuberculosis. Trends in Microbiology, 2023.
26. Sheng, W., G. Ji, and L. Zhang, *Role of macrophage scavenger receptor MSR1 in the*
progression of non-alcoholic steatohepatitis. Frontiers in Immunology, 2022. **13**: p.
1050984.
27. Purkait, B., et al., *Mechanism of amphotericin B resistance in clinical isolates of Leishmania*
donovani. Antimicrobial agents and chemotherapy, 2012. **56**(2): p. 1031-1041.
28. Srivastava, P., et al., *An unusual case of Amphotericin B refractoriness in Visceral*
Leishmaniasis from a non-endemic region of India. Journal of clinical microbiology, 2011.
49(8): p. 3088.
29. Ghosh, J., et al., *Human visceral leishmaniasis: decrease in serum cholesterol as a function*
of splenic parasite load. Annals of Tropical Medicine & Parasitology, 2011. **105**(3): p. 267-
271.
30. Haldar, A.K., et al., *Leishmania donovani isolates with antimony-resistant but not-sensitive*
phenotype inhibit sodium antimony gluconate-induced dendritic cell activation. PLoS
Pathogens, 2010. **6**(5): p. e1000907.
31. Beattie, L., et al., *Dynamic imaging of experimental Leishmania donovani-induced hepatic*
granulomas detects Kupffer cell-restricted antigen presentation to antigen-specific CD8+
T cells. PLoS pathogens, 2010. **6**(3): p. e1000805.
32. Hinder, L.M., et al., *Apolipoprotein E knockout as the basis for mouse models of*
dyslipidemia-induced neuropathy. Experimental neurology, 2013. **239**: p. 102-110.
33. Carter, K., et al., *The in vivo susceptibility of Leishmania donovani to sodium*
stibogluconate is drug specific and can be reversed by inhibiting glutathione biosynthesis.
Antimicrobial agents and chemotherapy, 2003. **47**(5): p. 1529-1535.
34. Semini, G., et al., *Changes to cholesterol trafficking in macrophages by Leishmania*
parasites infection. Microbiologyopen, 2017. **6**(4): p. e00469.
35. Vance, J.E., *Cellular itinerary of LDL cholesterol.* Proceedings of the National Academy of
Sciences, 2022. **119**(6): p. e2122584119.
36. Pessoa, C.C., et al., *ATP6V0d2 controls Leishmania parasitophorous vacuole biogenesis via*
cholesterol homeostasis. PLoS pathogens, 2019. **15**(6): p. e1007834.

37.   Feuerer, N., et al., *Lipidome profiling with Raman microspectroscopy identifies*
979   *macrophage response to surface topographies of implant materials.* Proceedings of the
980   National Academy of Sciences, 2021. **118**(52): p. e2113694118.

981   38.   Živanović, V., et al., *Chemical mapping of leishmania infection in live cells by SERS*
982   *microscopy.* Analytical chemistry, 2018. **90**(13): p. 8154-8161.

983   39.   Tiwari, D., et al., *Temporal in vitro Raman spectroscopy for monitoring replication kinetics*
984   *of Epstein–Barr virus infection in glial cells.* ACS omega, 2020. **5**(45): p. 29547-29560.

985   40.   Beattie, J.R., S.E. Bell, and B.W. Moss, *A critical evaluation of Raman spectroscopy for the*
986   *analysis of lipids: Fatty acid methyl esters.* Lipids, 2004. **39**(5): p. 407-419.

987   41.   Goldstein, J.L., et al., *Receptor-mediated endocytosis: concepts emerging from the LDL*
988   *receptor system.* Annu Rev Cell Biol, 1985. **1**(1): p. 1-39.

989   42.   Lucero, D., et al., *Interleukin 10 promotes macrophage uptake of HDL and LDL by*
990   *stimulating fluid-phase endocytosis.* Biochimica et Biophysica Acta (BBA)-Molecular and
991   Cell Biology of Lipids, 2020. **1865**(2): p. 158537.

992   43.   Popow-Woźniak, A., et al., *Cofilin overexpression affects actin cytoskeleton organization*
993   *and migration of human colon adenocarcinoma cells.* Histochemistry and Cell Biology,
994   2012. **138**: p. 725-736.

995   44.   Urano, Y., et al., *Transport of LDL-derived cholesterol from the NPC1 compartment to the*
996   *ER involves the trans-Golgi network and the SNARE protein complex.* Proceedings of the
997   National Academy of Sciences, 2008. **105**(43): p. 16513-16518.

998   45.   Verma, J.K., R. Rastogi, and A. Mukhopadhyay, *Leishmania donovani resides in modified*
999   *early endosomes by upregulating Rab5a expression via the downregulation of miR-494.*
PLoS pathogens, 2017. **13**(6): p. e1006459.

46.   Garver, W.S., et al., *The Niemann-Pick C1 gene is downregulated by feedback inhibition of*
the SREBP pathway in human fibroblasts. Journal of lipid research, 2008. **49**(5): p. 1090-
1102.

47.   Ho, C.-M., et al., *Accumulation of free cholesterol and oxidized low-density lipoprotein is*
associated with portal inflammation and fibrosis in nonalcoholic fatty liver disease.
Journal of Inflammation, 2019. **16**: p. 1-8.

48.   Murray, H., H. Masur, and J. Keithly, *Cell-mediated immune response in experimental*
visceral leishmaniasis. I. Correlation between resistance to Leishmania donovani and
lymphokine-generating capacity. 1982.

49.   Kima, P.E. and L. Soong, *Interferon gamma in leishmaniasis.* Frontiers in immunology,
2013. **4**: p. 156.

50.   Govaere, O., et al., *Macrophage scavenger receptor 1 mediates lipid-induced*
inflammation in non-alcoholic fatty liver disease. Journal of hepatology, 2022. **76**(5): p.
1001-1012.

51.   Lu, M. and O. Gursky, *Aggregation and fusion of low-density lipoproteins in vivo and in*
vitro. Biomolecular concepts, 2013. **4**(5): p. 501-518.

52.   Yano, H., Y. Fujiwara, and Y. Komohara, *Cholesterol metabolism and lipid droplet vacuoles;*
a potential target for the therapy of aggressive lymphoma. Journal of clinical and
experimental hematopathology, 2022. **62**(4): p. 190-194.

53.   Banerjee, S., et al., *Leishmania donovani infection induce Extracellular signal-regulated*
kinase ½ (ERK½) mediated lipid droplet generation in macrophages. Molecular
Immunology, 2022. **141**: p. 328-337.

54. Meester, I., et al., *The roles of lipid droplets in human infectious disease.* Revista
Portuguesa de Cardiologia, 2011. **13**(53): p. 207-216.
55. Bosch, M., et al., *Lipid droplets and the host–pathogen dynamic: FATal attraction?* Journal
of Cell Biology, 2021. **220**(8): p. e202104005.
56. Liffner, B., et al., *Atlas of Plasmodium falciparum intraerythrocytic development using
expansion microscopy.* Elife, 2023. **12**: p. RP88088.
57. Ghosh, S., et al., *A novel bioimpedance-based detection of miltefosine susceptibility
among clinical Leishmania donovani isolates of the Indian subcontinent exhibiting
resistance to multiple drugs.* Frontiers in Cellular and Infection Microbiology, 2021. **11**: p.
768830.
58. D'Avila, H., et al., *Host cell lipid bodies triggered by Trypanosoma cruzi infection and
enhanced by the uptake of apoptotic cells are associated with prostaglandin E2
generation and increased parasite growth.* Journal of Infectious Diseases, 2011. **204**(6): p.
951-961.
59. Thakur, A., H. Mikkelsen, and G. Jungersen, *Intracellular pathogens: host immunity and
microbial persistence strategies.* Journal of immunology research, 2019. **2019**(1): p.
1356540.
60. Kreimendahl, S. and L. Pernas, *Metabolic immunity against microbes.* Trends in Cell
Biology, 2023.
61. Díaz-Varela, M., et al., *The different impact of drug-resistant Leishmania on the
transcription programs activated in neutrophils.* iScience, 2024. **27**(5): p. 109773.
62. Geraci, N.S., J.C. Tan, and M.A. McDowell, *Characterization of micro RNA expression
profiles in L eishmania-infected human phagocytes.* Parasite immunology, 2015. **37**(1): p.
43-51.
63. Tiwari, N., et al., *Identification and characterization of miRNAs in response to Leishmania
donovani infection: delineation of their roles in macrophage dysfunction.* Frontiers in
microbiology, 2017. **8**: p. 314.
64. Moradimotlagh, A., et al., *Leishmania infection upregulates and engages host
macrophage Argonaute 1, and system-wide proteomics reveals Argonaute 1-dependent
host response.* Frontiers in Immunology, 2023. **14**: p. 1287539.
65. Boodhoo, N., N. Kamble, and S. Behboudi, *De novo cholesterol biosynthesis and its
trafficking in LAMP-1-positive vesicles are involved in replication and spread of Marek's
disease virus.* Journal of Virology, 2020. **94**(24): p. 10.1128/jvi. 01001-20.
66. Prakash, S. and A.K. Rai, *Retinoic acid increases cellular cholesterol in Leishmania
donovani-infected macrophages in an mTOR-independent manner.* Microbiology
Spectrum, 2022. **10**(6): p. e02699-22.
67. Kim, H. and S.J. Shin, *Revolutionizing control strategies against Mycobacterium
tuberculosis infection through selected targeting of lipid metabolism.* Cellular and
Molecular Life Sciences, 2023. **80**(10): p. 291.
68. Awadh, A.A., *The Role of Cytosolic Lipid Droplets in Hepatitis C Virus Replication, Assembly,
and Release.* Biomed Res Int, 2023. **2023**: p. 5156601.
69. 2024, L.i.f.b.a.W.T.C.A.Z.Z., *LeishGEM is funded by a Wellcome Trust Collaborative Award*,
in *LeishGEM*. 2024.
70. Downing, T., et al., *Whole genome sequencing of multiple Leishmania donovani clinical
isolates provides insights into population structure and mechanisms of drug resistance.*
Genome Res, 2011. **21**(12): p. 2143-56.

71.　Patole, S., S. Burza, and G.M. Varghese, *Multiple relapses of visceral leishmaniasis in a*
patient with HIV in India: a treatment challenge. International Journal of Infectious
Diseases, 2014. **25**: p. 204-206.
72.　Ray, A. and B.N. Dittel, *Isolation of mouse peritoneal cavity cells.* JoVE (Journal of
Visualized Experiments), 2010(35): p. e1488.
73.　Khanra, S., et al., *In vivo experiments demonstrate the potent antileishmanial efficacy of*
repurposed suramin in visceral leishmaniasis. PLoS Neglected Tropical Diseases, 2020.
14(8): p. e0008575.
74.　Li, P.-z., et al., *An efficient method to isolate and culture mouse Kupffer cells.* Immunology
letters, 2014. **158**(1-2): p. 52-56.
75.　Mukherjee, S., et al., *Short-course treatment with imipramine entrapped in squalene*
liposomes results in sterile cure of experimental visceral leishmaniasis induced by
antimony resistant Leishmania donovani with increased efficacy. Frontiers in Cellular and
Infection Microbiology, 2020. **10**: p. 595415.
76.　Amundson, D.M. and M. Zhou, *Fluorometric method for the enzymatic determination of*
cholesterol. Journal of biochemical and biophysical methods, 1999. **38**(1): p. 43-52.
77.　Real, F., *Isolation of intact Leishmania amazonensis large parasitophorous vacuoles from*
infected macrophages by density gradient fractionation. Experimental Parasitology, 2020.
218: p. 107989.
78.　Alpizar-Sosa, E.A., et al., *Amphotericin B resistance in Leishmania mexicana: Alterations*
to sterol metabolism and oxidative stress response. PLoS Neglected Tropical Diseases,
2022. **16**(9): p. e0010779.
79.　Schirripa Spagnolo, C. and S. Luin, *Setting up multicolour TIRF microscopy down to the*
single molecule level. Biomolecular Concepts, 2023. **14**(1): p. 20220032.

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

Table-1: EC50 against SAG (Antimonial) for the LD strains used in this study

Strain Name	EC50 value for SAG(µg/ml) in PEC	EC50 value for SAG(µg/ml) in KC
MHOM/IN/2009/BHU575/0	22.73±3.7	23.89±2.1
MHOM/IN/10/BHU814/1	30.3±7.2	28.5±5.3
MHOM/IN/83/AG83	1.52±1.2	3.3±2.2
MHOM/IN/80/DD8	2.1±1.2	3.2±1.9
DD8-AmpB	1.8±2.1	2.2±1.3
BHU575-EGFP	21.45±3.2	22.81±3.1
AG83-EGFP	2.1±1.4	2.4±3.1

Table-2: Lipid profile of VL patients

Serum Code	Age	Sex	Time of Blood Collection	Infection status	Treatment received	Patient status	LDL (mg/dl)	HDL (mg/dl)	Cholesterol (mg)	Triglycerides (mg/dl)
RMRI N1	45	F	Before	Control population from endemic	NA	NA	45.3501585	46.6365931	101.6949153	177.258567
RMRI N6	70	F	Before	Control population from endemic	NA	NA	110.855478	50.3762712	111.8644068	206.5420561
RMRI N7	45	F	Before	Control population from endemic	NA	NA	75.5620787	41.8711864	84.40677966	165.4205607
RMRI N8	19	M	Before	Control population from endemic	NA	NA	83.946038	62.9533123	100.3389831	224.9221184
RMRI N11	37	M	Before	Control population from endemic	NA	NA	72.4549921	120.914196	134.2372881	244.5482866
RMRI N12	60	M	Before	Control population from endemic	NA	NA	52.6307448	113.851735	92.54237288	180.6853583
RMRI N13	30	M	Before	Control population from endemic	NA	NA	66.7533281	48.3413249	138.3050847	236.4485981
RMRI N14	22	M	Before	Control population from endemic	NA	NA	47.8939778	49.6807571	100	260.7476636
RMRI N15	37	F	Before	Control population from endemic	NA	NA	46.4904913	45.9059937	108.8135593	219.6261682
RMRI C1	20	F	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	110.389045	44.2700565	112.5423729	200.623053
RMRI C2	62	F	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	121.738904	64.5514124	145.4237288	301.8691589
RMRI C3	15	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	76.1839888	36.7463277	95.59322034	185.6697819
RMRI C4	31	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	72.141573	41.1079096	116.9491525	241.7445483
RMRI C5	52	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	98.1063202	48.1954802	117.2881356	282.2429907
RMRI C6	18	F	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	68.8765449	36.7463277	96.61016949	200.623053
RMRI C7	45	F	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	110.544522	55.6101695	124.4067797	381.9314642
RMRI C8	32	F	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	94.8412921	43.0706215	106.1016949	232.3987539
RMRI C18	66	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	61.4136236	35.9830508	91.86440678	203.4267913
RMRI R14	8	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Relapse	79.2935393	83.6333333	96.94915254	262.9283489
RMRI R16	49	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Relapse	56.9047753	102.279096	63.05084746	209.0342679
RMRI R17	18	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Relapse	59.7033708	101.079661	83.38983051	217.1339564
RMRI R18	59	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Relapse	63.4348315	94.319209	84.40677966	304.6728972

Figure Legends

Figure 1. Severe dyslipidemia is linked with higher intracellular proliferation of LD-R amastigotes. (A) Serum lipid profile of 28 days LD-infected Balb/c mice (N=5). Equal number of sorted metacyclics promastigotes of two representative LD-S (LD-S[1], LD-S[2]) and LD-R (LD-R[1], LD-R[2]) strains were used to perform independent infection. **(B)** Number of attached LDs on KC surface 4Hrs p.i, data represents (N=3) for two LD-S and two LD-R strains. **C (i)** Representative Giemsa stain images of LD-S[1] and LD-R[1]-infected KC at 4Hrs, 24Hrs and 48Hrs p.i Black arrow represents LD amastigotes. N denotes host nucleus. Scale bar 20μM. Giemsa Images are represented in grey scale to clearly represent LD nucleus (black arrow). **C (ii)** Intracellular amastigote count for 4Hrs, 24Hrs and 48Hrs p.i. Each dot represent count from 100 infected-KC (N=6). **D (i)** LD-infected-KC imaged to visualize the PV at 4Hrs and 24Hrs p.i. PV marked by ATP6V0D2 (magenta), KC and LD nucleus (blue), Scale bar 5μm. Yellow arrow represents LD amastigotes. In the merged image, white dotted line marks KC's periphery, and host nucleus is represented with N. **D (ii)** PV counts measured from 30 infected KCs. **D (iii)** Confocal 3D reconstruction illustrating the spatial distribution of parasitophorous vacuoles (PVs) in Kupffer cells (KCs) infected for 24 hours. ATP6V0D2, a lysosomal vacuolar ATPase subunit, is visualized in magenta, while the nucleus is depicted in cyan. The final panel highlights PV structural grooves outlined in red solid lines, with intracellular *Leishmania donovani* (LD) amastigotes indicated by white arrows. Higher magnification of Figure 1D further emphasizes the increased abundance of PVs in LD-R infected cells, suggesting enhanced intracellular replication and adaptation mechanisms of drug-resistant strains. Scale bar 5μM. Both yellow and magenta solid line box represents the same area of the image. *** signifies p value < 0.0001, * signifies p value < 0.05, n.s non-significant.

Figure 2. LD-R-infection results in a higher organ parasite load in hypercholesterolemic *Apoe*[-/-]mice. A (i) Representative Giemsa images of LD-infected-KCs cultured in high and low lipid media. Black arrow represents LD amastigotes; N represents host nucleus. Scale bar 20μM. **A (ii)** Graph representing number of intracellular LD amastigotes in KCs with LD infection performed either in high, low or lipid free media (N=6). **B (i)** Giemsa images of LD-infected-KCs isolated from wild type BL/6 and *Apoe*[-/-] mice. Scale bar 20μm, N representing host nucleus. Giemsa Images are represented in grey scale to clearly represent LD nucleus (black arrow). **B (ii)** Amastigote count from 100 LD-infected-KCs (N=6) as in B(i). Data are presented as mean ± SEM. **C (i)** Schematic representation of *in-vivo* LD-infection in BL/6 and *Apoe*[-/-] mice performed with EGFP-LD-S or EGFP-LD-R **C (ii)** Flow cytometry representing GFP-positive splenocytes isolated from 11 days EGFP-LD infected BL/6 and *Apoe*[-/-] mice. The black box indicates % of GFP-positive splenocytes (LD-infected). **C (iii)** Graph representing data from six infected mice as in C (ii) are presented. *** signifies p value < 0.0001, ** signifies p value <0.001, * signifies p value < 0.05, n.s non-significant.

Figure 3. Endocytosed-LDL is the primary lipid source for intracellular LD-R proliferation. A (i) Live cell images of LDL-endocytosis in 24Hrs post infected KCs with LDL (red), LD (green), and the nucleus (blue). Images were acquired directly from 96 well glass bottom plate and bright field was merged with reduced brightness and increased contrast identically for all experimental conditions. Scale bar 40μm, Yellow arrow represents intracellular LD. **A (ii)** Graph representing Fluorescence plate reader-based quantification of A (i). **B (i)** Co-localization of LDL with the LD-amastigotes in infected KCs 24Hrs p.i. Merged image showing LDL (magenta), LD (green), nucleus (blue), White dotted line in the merged image demarcates infected-KC for co-localization analysis. Scale bar 10μm, yellow arrow represents LD nucleus and N represents host nucleus **B (ii)** Intensity plot representing co-localized signals from B (i), marked with grey transparent box. **C (i)** Confocal image showing the isolated LD-S and LD-R-PV, stained with LAMP1 (magenta) and LD (Green)**.** Scale bar 2μm **C (ii)** Western blot analysis validated the purification of PV fractions, indicated by LAMP-1 positivity, while the presence of LD parasites within the PV was confirmed by KMP-11 detection**. C (iii)** Western blot of cellular fraction without PV showing minimal presence of LAMP1. **D (i)**GC-MS lipid profile of isolated PVs from LD-infected-KCs. Retention time 27.73 corresponding

to the cholesterol and 28.56 corresponding to fatty acids are represented with * for LD-S-PV and
with black arrowheads for LD-R-PV. **D (ii)** Relative abundance of Cholesterol and Fatty acids of D
(i). **E (i)** Representative Confocal Raman spectroscopy image of uninfected or LD-infected-KCs
24Hrs p.i, illustrating distinct lipid-related signal distribution patterns marked with pseudo-colours.
Yellow dotted circle in the left most panel demarcates the host cell nucleus while white arrow
marking intracellular LD nucleus. In merged panel white dotted line demarcates cells periphery.
The right most panel shows dot plot representation of lipid distribution, where each colour
corresponds to different lipid-related peaks with respect to blue (B) indicating the nucleus. **E (ii)**
Comparative Raman spectra from LD-S and LD-R-infected-KCs. Lipid-related peaks are
demarcated with shaded box, representing 1. 540-560 cm-1 (Cholesterol), 2. 1080-1090 cm-1
(phospholipids), 3. 1270-1280 cm-1 (triglycerides), 4. 1300-1340 cm-1 (Amide-III bond), 5. 1440-
1453 cm-1 (Fatty acids and Triglycerides), and 6. 1650-1660 cm-1 (Amide-I bond). *** signifies p
value < 0.0001, ** signifies p value <0.001, * signifies p value < 0.05, n.s non-significant.

Figure 4. Increased LDL uptake in LD-R-Infected-MΦs through receptor independent fluid
phase endocytosis. (A) Immunofluorescence images of LDLr expression in LD-infected-KCs.
Scale bar 40µm. Enlarged view of A showing infected-KC with low LDLr expression (yellow arrow)
compared to neighboring uninfected KC (white arrow) in the merged image. **B (i)** Cofilin and
phosphorylated-Cofilin expression by western blot in LD-infected-KCs. **B (ii)** Graphical
representation of comparative expression of Cofilin and phosphorylated-Cofilin with respect to β-
actin for B (i). **C (i)** SIM images showing the Cofilin distribution in LD-infected-KCs. Yellow arrow
represents intracellular LD, white dotted line in the merged image demarcates cell boundary. Scale
bar 5µm. **C (ii)** Colour spectrum representing Cofilin distribution pattern of C (i) with green
representing maximum and red representing minimum distribution. **D (i)** Confocal images
representing Filamentous actin (F-actin) protrusions in LD-infected-KCs. Scale bar 20µm, with
yellow arrow representing LD amastigotes. Individual experimental condition is represented in
pseudo-colour, with uniform high contrast to clearly represent the F-actin protrusions. **D (ii)** Graph
representing Cofilin and F-actin distribution in KCs infected with LD or left uninfected. **(E)** Still
images from live cell video microscopy representing co-localization of LD-R with LDL in presence
or absence of Latrunculin-A (LAT-A). Scale bar 60µm. Yellow arrow represents LD-infected-KCs
with LDL(red). **(F)** Graphical representation (N=6) of amastigote count in 100 LD-R-infected-KCs
treated with either LAT-A, or Cytochalasin (CYT-D) or left untreated. Data are presented as mean
± SEM. ** signifies p value <0.001, * signifies p value < 0.05, n.s non-significant.

Figure 5. Endocytosed-LDL fuses with LD-R-PV to provide Cholesterol to LD-R amastigotes.
(A) Confocal images of representing LDL-vesicles, with LAMP1 positive lysosomal vesicles in LD-
infected-KCs 4Hrs p.i. Yellow arrow represents LD nucleus, white dotted box marks the region
further cropped and zoomed to show fusion of lysosomal vesicles with LDL and LD-R. Scale bar
10µm. **A (ii)** Schematic representation showing trafficking of LDL-vesicle towards the LD-R-PV. **B**
(i) SIM images representing convergence of LDL-vesicles with LD amastigotes and LAMP1 in
24Hrs with LD-infected-KCs. White dotted box marks the region further cropped and Zoomed to
clearly show the convergence (of LD amastigotes with LDL and LAMP-1 (white vesicles shown by
yellow arrow), Scale bar 5µm. **B (ii)** Intensity plot representing of co-localization signals for B (i). **C**
Western blot analysis showing LAMP1 expression in LD-infected-KCs from early to late hours. **C**
(i), C (ii) representing LAMP1 expression LD-S and LD-R infected-KC **D (i)** Filipin staining of EGFP-
LD-infected-KCs. Filipin (white) and LD (green), yellow arrow represents intracellular LD. Scale bar
10µm. White dotted line marked cells used to generate co-localization plot. **D (ii)** Co-localization
signals of LD (EGFP and Filipin represented by the dot plot. Pearson co-localization coefficient
value (r) represented for LD-S and LD-R-infected-KC. **D (iii)** Graph representing Pearson's
coefficient for EGFP-LD-infected-KCs (N=19). **E** Filipin staining of EGFP-LD-R-infected KCs
incubated with dextran beads (Magenta) revealed minimal colocalization between the beads and
Filipin, while the LD-R amastigotes within same macrophage shows a significant co-localization
with Filipin. Blue arrow demarking beads with limited filipin colocalization, red arrow demarking LD-
R with filipin colocalization. Scale bar 5µM. Statistical significance in the observed differences was

determined through ANOVA utilizing GraphPad Prism software (version 9.0). *** signifies p value
< 0.0001.

Figure 6. LD-R infection suppress NPC-1 to alter Cholesterol mobilization in infected MΦs.
(A) 24Hrs LD-S and LD-R-infected-PECs were subjected to *RNAseq* analysis keeping uninfected-
PECs as control. Heat map representing average expression of differentially expressed genes
related to lipid metabolism between LD-S and LD-R-infected-PEC. Scale represents median-
centered counts (TPM). Row Z score data represented here. Expression of RNAs (right margins)
is presented as centered and ''scaled'' [mean normalized $\log_2(TPM + 1)$]. **B (i)** Expression of
SREBP2 and **B (ii)** NPC-1 in LD-infected and uninfected KCs by Western blot with β-actin as
control. **C** Intracellular localization of SREBP2 in response to LD-infection in KC 24Hrs p.i. White
dotted line represents cell periphery with yellow arrow representing intracellular LD and N
representing host nucleus. 3D image represented below with one quadrant clipped to confirm
nuclear translocation of SREBP2. Scale bar 5µm. **(D)** Expression of HMGCR LD-infected and
uninfected KCs by Western blot with β-actin as control. **E** Immunofluorescence showing NPC-1
expression in LD-infected-KCs. White arrow representing NPC-1 expression in LD-R-infected-KCs
and yellow arrow representing NPC-1 in adjacent uninfected-KCs in the merged panel. White
dotted box marks regions further cropped and shown in the bottom panel with enlarged view
(Zoomed). Scale bar 20µm. **F (i)** TIRF microscopic images of Laurdan stained LD-infected and
uninfected KCs with two different spectra (488nm, 594nm), changes from red to green represents
gel to fluid phase transition of the host plasma membrane. Colour distribution pattern corresponding
to the TIRF images representing each experimental condition is presented in the right most panel.
Scale bar 5µm. **F (ii)** Measurement of membrane cholesterol by Amplex red assay kit. Graphical
representation showing total membrane cholesterol divided by total membrane protein. **G (i)**
Expression of NPC-1 determined by western blot in PECs transfected with NPC-1 siRNA or
scrambled (sc) siRNA. **G (ii)** Expression of NPC-1 determined by immunofluorescence in PECs
transfected with NPC-1 siRNA or scrambled (sc) siRNA. Scale bar 5µm. **G (iii)** Giemsa-stained
images of LD-infected NPC-1 knockdown-PECs with scrambled siRNA control (Sc). Scale bar
20µm. Images are represented in grey scale with increased contrast to represent LD nucleus (black
arrow). **G (iv)** Graphical representation of number of intracellular LD-amastigotes as in F (ii).
Results show the counting of 100 infected-PEC from 6 independent replicate. * signifies p value <
0.05, n.s non-significant.

Figure 7. LD-R-infected-MΦs selectively excludes ox-LDL to suppress inflammatory
response in host. (A) Scheme representing *ex vivo* co-culture experiment performed by isolating
T-cells from LD-infected mice with in vitro LD-infected-KCs **B (i)** IFN-γ level in the supernatant of
LD-infected-KCs co-cultured with the T-cells isolated from spleen of Balb/c mice infected with LD-
S or LD-R 28 days p.i. Data represented as mean ± SD from three independent experiments. **B**
(ii) IFN-γ level in the supernatant of splenocytes cultures from LD-infected-BL/6 and *Apoe*[-/-]-mice
stimulated with SLA (Soluble *Leishmania* antigen) specific for LD-S or LD-R. Data are presented
as mean ± SD from six independent mice. **C (i)** ox-LDL incubated LD-infected-KCs was fixed and
stained with Oil-red-O and Haematoxylin. LD-S[High] represents MOI (1:20) and LD-R[Low] represents
MOI (1:5). Scale bar 10µm. Black arrow represents LD nucleus. **C (ii)** Graph representing % of ox-
LDL positive MΦs under different experimental conditions as represented in C (i). **D (i)** Confocal
image showing MSR-1 in LD-infected-KCs. Yellow arrow represents LD-amastigotes while N
represents host nucleus. Scale bar 20µm **D (ii)** Expression of MSR1 detected by Western blot of
LD-infected-KC with β-actin as loading control. (**E**) Expression of ATP6V0D2 in LD-infected-KCs
by Western blot with β-actin as loading control. **F (i)** Scheme representing the time line of *in vivo*
experiment for IFA. **F (ii)** Expression of MSR1 and neutral lipid droplet accumulation assayed by
Confocal microscopy from the cryo-sectioned liver of 28 days LD-infected-mice. Representative
image showing Lipid droplet marked in red and MSR1 in green, nucleus in blue. Scale bar 40µm.
F (iii) Graphical representation of % positive Lipid droplets accumulation and MSR-1 expression
under different experimental condition as in E (i). Data collected from N=6 mice. *** signifies p value
< 0.0001, ** signifies p value <0.001, * signifies p value < 0.05.

Figure 8. Lipid droplets accumulation around LD amastigotes is inversely related with susceptibility towards Amphotericin-B. (A) Ex-M image showing lipid droplet assimilation in LD-R-amastigotes. White dotted region further cropped and shown in enlarged panel (Zoomed). Nile red (magenta), LD (green), nucleus (cyan). White spots represent Lipid droplets assimilated in LD-R and Host nucleus is represented as N. Scale bar 5μm. **(B)** TEM images showing lipid droplets around the LD-R amastigotes. Black box further magnified and represented in right panel. White * representing LD-R amastigotes, yellow * showing lipid droplets. Scale bar 2μm. (**C**) Schematic representation representing experimental scheme used to determine EC50 against Amphotericin-B under different experimental conditions**. D (i) and (ii)** Determination of EC50 value against Amphotericin-B in low lipid and high lipid media by performing infection with of two independent LD-R strains. **D (iii)** Determination of EC50 value against Amphotericin-B in KC infected with LD-S$^{Amp-B-R}$. Intracellular amastigotes were counted through Giemsa staining. **(E)** Confocal images showing lipid droplets accumulation around intracellular amastigotes with Aspirin (5 μM) or Amp-B (0.36 μM) or with a combination of Aspirin (5μM) and Amp-B (0.11μM)at 48Hrs p.i **(F)** Determination of amastigote load in KCs infected with two independent LD-R isolates in different experimental conditions either untreated or treated with Aspirin (5 μM) or Amp-B (0.36 μM) or with a combination of Aspirin (5μM) and Amp-B (0.11μM) at 48Hrs p.i (**G**) Comparative Lipid profile of VL patients (Cured and Relapsed) in response to Amp-B treatment along with healthy individuals from endemic region of Bihar, India. *** signifies p value < 0.0001, ** signifies p value <0.001, * signifies p value < 0.05.

Figure 9: Scrutinized uptake of host lipids results in breakdown of Amphotericin-B responsiveness in clinical LD isolates with primary resistance to antimony. LD-R reside within membrane-bound parasitophorous vacuoles (PVs) and actively modulate host actin cytoskeleton dynamics through cofilin-mediated depolymerization, inducing high fluid-phase endocytosis of LDL. The internalized LDL-containing vesicles subsequently fuse with LD-R-associated PVs, facilitating increased lipid acquisition essential for their aggressive intracellular proliferation. In these fused membranes, the vesicular cholesterol export protein NPC-1 is downregulated, effectively sequestering LDL-derived cholesterol within the PV and preventing its release into the host cytoplasm. Concurrently, LDL-derived cholesterol esters either directly or getting processed in the endoplasmic reticulum (ER) can lead to the formation of lipid droplets (as presented by **?,** since it needs further investigation. Lipid droplets then accumulate in and around the LD-R-PV. This lipid droplet accumulation is a key factor in rendering LD-R infections unresponsive to Amphotericin-B. Additionally, LD-R regulate inflammatory responses linked to oxidized lipid (Ox-LDL) accumulation by downregulating macrophage scavenger receptor 1 (MSR-1). The elevated LDL influx also leads to a shutdown of *de novo cholesterol* biosynthesis by restricting the nuclear translocation of sterol regulatory element-binding protein 2 (SREBP2).

Supplementary Information

Figure S1. Kupffer cells (KC) as an *in vitro* model for LD-infection. (A) Flow Cytometry representing % Metacyclics LD: **(i)** LD-S (AG83, DD8) and **(ii)** LD-R (BHU814, BHU575) based on FSC and SSC. Left, small box with low FSC and SSC gate for metacyclic parasites and right panel with high FSC and SSC gated procyclic parasites. **(B)** Flow cytometry of F4/80-PE positive KC from the non-parenchymal cell population isolate from murine liver. **C** Representative Giemsa-stained images LD-infected-PEC, 4 and 24Hrs p.i for **(i)** LD-S and **(ii)** LD-R strains. Giemsa Images are represented in grey scale to clearly represent LD nucleus (black arrow). Scale Bar 20µM **C (iii)** Intracellular amastigote count for 4Hrs and 24Hrs p.i. Each dot represent count from 100 infected-KC. N=6. **D** Amastigote count from Giemsa-stained image of KC-infected with LD amastigotes, isolated from the spleen of 28 days LD-infected-mice. Each dot represent mean from 100 infected-KCs. N=6. *** signifies p value < 0.0001, ** signifies p value <0.001, n.s non-significant.

Figure S2. LD-S and LD-R quenches host membrane cholesterol during initial entry. A (i) KCs were pre-incubated with NBD-cholesterol for 16Hrs. Subsequently, KCs were infected with RFP expressing LD-S or LD-R for 4Hrs or left uninfected and imaged. NBD-cholesterol was excited at 488nm, and LDs are presented with magenta pseudo color. White arrow indicates the NBD-cholesterol within LD while yellow arrow represents LD nucleus. Scale bar 5µm. **A (ii)** Co-localization analysis of LD (magenta), and NBD-Cholesterol (green) from A (i), marked in grey transparent box. **B (i)** Representative Flow cytometry-based analysis of NBD in LD-amastigotes isolated from LD-infected-KCs 2Hrs after infection pre-incubated with NBD cholesterol. Black Box representing % NBD positive population. **B (ii)** Graphical representation of MFI for three independent experiments as in B (i). n.s non-significant.

Figure S3. LD-R-PV have higher accumulation of Cholesterol and Fatty acids. (A) Detection of PV Cholesterol, isolated from LD-S or LD-R-infected KC. Data represented as µg of protein/ µg of Cholesterol, (N=6). **(B)** TLC profile of isolated lipids from 24Hrs LD-infected-PVs with Cholesterol standard kept as control. (**C**) Individual Raman spectra from six independent LD-infected-KCs **(i)** LD-S-infected and **(ii)** LD-R-infected. Lipid-related peaks are demarcated within the spectra, including those at 1.540-560 cm-1 (Cholesterol), 2. 1080-1090 cm-1 (phospholipids), 3. 1270-1280 cm-1 (triglycerides), 4. 1300-1340 cm-1 (Amide-(II) I bond), 5. 1440-1453 cm-1 (Fatty acids and Triglycerides), and 6. 1650-1660 cm-1 (Amide-I bond). **(D)** Area under the curve, and shift are calculated and plotted for each experimental condition as represented in C (i) and (i). ** signifies p value <0.001

Figure S4. Latrunculin-A restricts extracellular LDL influx inhibiting LD-R amastigote proliferation. (A) Western blot comparing of LDLr expression between uninfected, LD-S, and LD-R-infected-KC 24Hrs p.i. **B (i)** Expression of LDLr determined by immunofluorescence in PECs treated with LDLr siRNA and scrambled siRNA (Sc). Scale bar 10µm. **B (ii)** Western blot confirming LDLr knockdown upon SiRNA treatment. Scrambled RNA (ScRNA) was used as a negative control, while Small Interfering RNA (SiRNA) specifically targeted LDLr transcripts, TR-1 and TR-2 represent independent experimental trials**.** β-Actin was used as an endogenous loading control for Western blot normalization**. C** Graph representing number of LD-amastigotes in LDLr knockdown per 100 PECs for B (i) (N = 6). Data: mean ± SD. **D (i)** Cofilin and phosphorylated-Cofilin expression by Western blot in LD-infected-PEC. **D (ii)** Graphical representation of comparative expression of abundance of Cofilin and phosphorylated-Cofilin with respect to β-actin for C(i). (**E**) Giemsa-stained images of Latrunculin-A (LAT-A) and Cytochalasin-D (CYT-D) treated LD-infected-KCs in grey scale. LD nucleus represented with Black arrow and host nucleus are represented as N. Scale bar 20µm. n.s non-significant.

Figure S5. Aspirin treatment inhibits Lipid droplets formation and increase responsiveness of LD-R amastigotes towards Amphotericin-B. (A) Determination of lipid droplet accumulation by Nile red staining in LD-S and LD-R-infected-KC or KC left uninfected. Dotted line mark the cell periphery while dotted box cells were used for 3D reconstruction and represent in Movie 4. Scale

Bar 5 µm. (**B**) Graphical representation of Lipid Droplet positive KCs in high lipid and low lipid
media. Data represented as % of lipid droplet positive KCs (N=6) collected from 3 independent
experiments. (**C**) Effect of Aspirin(5µM) on Lipid droplets accumulation in LD-R-infected-KCs.
Independent infection was performed with LD-R[1] and LD-R[2] strains (N=6) and % KCs with high
lipid droplets accumulation are represented. (**D**) Representative Giemsa stained images of LD-R-
infected-KCs in different experimental conditions either untreated or treated with Aspirin (5µM) or
Amp-B (0.36 µM) or with a combination of Aspirin (5µM) and Amp-B (0.11 µM) at 48Hrs p.i. Scale
bar 20µm. *** signifies p value < 0.0001

1381

Movie Legends

Movie 1: Phase contrast video representing attached LD promastigotes to PEC 6Hrs p.i. Black
arrow showing flagella of attached LD parasite. **1A** represent LD-S and **1B** represent LD-R
infection. Scale bar 10µm.

Movie 2: Phase contrast video representing 72Hrs LD infected KCs. scale bar 20µm.

Movie 3: Effect of Latrunculin-A (LAT-A) on LDL uptake in LD-R infected PECs. **3A.** In absence of
LAT-A. **3B.** In presence of LAT-A. LD (green), LDL (red) nucleus (blue). Scale bar 60µm

Movie 4: 3D-representation of lipid droplets around the LD-S amastigotes and LD-R amastigotes
(S5B). Scale bar 5µm

Figure 1

A

Serum lipid profile (mg/dl)

UI
LD-S¹
LD-S²
LD-R¹
LD-R²

LDL   HDL   Cholesterol   Triglycerides

ns
*
*
*

B

% of attached LD/100 KC

ns

LD-S   LD-R

C i

4Hrs   24Hrs   48Hrs

LD-S¹

LD-R¹

C ii

Amastigotes/100 KC

4Hrs   24Hrs   48Hrs

*   *   ***   ***

LD-S 1   LD-S 2   LD-R 1   LD-R 2

D i

DAPI   ATP6V0D2   Merged

UI

LD-S 4Hrs p.i

LD-R 4Hrs p.i

LD-S 24Hrs p.i

LD-R 24Hrs p.i

D ii

Number of PV/KC

ns   ***

LD-S   LD-R   LD-S   LD-R

4Hrs   24Hrs

D iii

Merged   3D PV   Zoomed

ATP6V0D2   DAPI   3D PV

Figure 2

Figure 3

A i

EGFP-LD LDL-red DAPI

UI   LD-S   LD-R

A ii

- UI
- LD-S
- LD-R
- LD-R Killed
- LD-S Killed

Fluorescence Intensity

** ***

4Hrs p.i   24Hrs p.i

B i

DAPI   EGFP-LD   LDL-red   Merged

LD-S

N

LD-R

N

B ii

- LDL-red
- EGFP-LD
- Co-localized Signal

Intensity

Distance

C i

LD-S-PV   LD-R-PV

LAMP-1   LAMP-1
LD   Merge   LD   Merge

C ii

PV Fraction

UI LD-S LD-R

LAMP-1   —130
                                    —100
                                    —75

β-actin   —55
                                    —37

KMP-11   —25
                                    —15
                                    —10

C iii

Cellular Fraction without PV

UI   LD-S   LD-R

LAMP-1   —130
                                    —100

β-actin   —55
                                    —37

D i

LD-S-PV

25.57 25.77 26.48 26.64 * 27.73 * 28.56 29.16 29.89 ** 30.51 30.99

LD-R-PV

25.36 26.07 25.71 26.47 26.59 ◄ 27.74 ◄ 28.56 ◄ 29.53 29.22 29.68 29.78 31.00

Time(min)

Relative abundance

D ii

- LD-S
- LD-R

Relative abundance

*   *

Cholesterol   Fatty acids

E i

Nucleus Cholesterol Lipid Esterified lipid Merged Distribution pattern

UI

LD-S

LD-R

E ii

- LD-R
- LD-S

Intensity

1   2 3   4 5   6

500   1000   1500

Raman Shift (cm⁻¹)

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A **Ex-M**   Zoomed

LD
DAPI
Nile-Red

B

C KC infected with LD-R¹/LD-R²

LD-R¹/LD-R²/LD-S$^{Amp-B-R}$ cultured in M199

Wash after 4Hrs

24Hrs in low lipid medium → Amp-B treatment → After 24Hrs → EC50 value determination

24Hrs in high lipid medium → Amp-B treatment → After 24Hrs → EC50 value determination

D i   LD-R¹   High lipid / Low lipid

D ii   LD-R²   High lipid / Low lipid

D iii   LD-S$^{Amp-B-R}$ High lipid / Low lipid

E   DAPI   Nile Red   Merged

Asp only

Amp-B

Asp +Amp-B

F   LD-R¹   LD-R²   Amastigotes/100 KC   UT, Amp-B, Asp, Asp+Amp-B   ***   **

G   Control   Cured   Relapsed   Lipid profile (mg/dl)   LDL   HDL   Cholesterol   Triglycerides   *   *

Figure 9