Title: Scrutinized lipid utilization disrupts Amphotericin-B responsiveness in clinical
isolates of *Leishmania donovani*

Supratim Pradhan[1], Dhruba Dhar[1], Debolina Manna[1], Shubhangi Chakraborty[1], Arkapriya
Bhattacharyya[1], Khushi Chauhan[1], Rimi Mukherjee[2], Abhik Sen[2], Krishna Pandey[2], Soumen Das[1]
and Budhaditya Mukherjee[1]*.

[1] School of Medical Science and Technology, Indian Institute of Technology Kharagpur, India-
721302

[2] ICMR-Rajendra Memorial Research Institute of Medical Sciences, Patna, India-800007

*Corresponding   author: Budhaditya   Mukherjee,   Email: aditya26884@gmail.com,
bmukherjee@smst.iitkgp.ac.in

Running Title: Neutral Lipid Build-Up Disrupts Amp-B responsiveness

Abstract

The management of *Leishmania donova*ni (LD), responsible for fatal visceral leishmaniasis (VL), faces increasing challenges due to rising drug-unresponsiveness, leading to increasing treatment failures. While hypolipidemia characterizes VL, LD, a cholesterol auxotroph, relies on host lipid scavenging for its intracellular survival. The aggressive pathology, in terms of increased organ parasite load, observed in hosts infected with antimony-unresponsive-LD (LD-R) as compared to their sensitive counterparts (LD-S), highlights LD-R's heightened reliance on host lipids. Here we report that LD-R-infection promotes fluid-phase endocytosis in the host, selectively accumulating neutral lipids while excluding oxidized-LDL. LD-R enhances the fusion of endocytosed LDL-vesicles with its phagolysosomal membrane and inhibits cholesterol mobilization from these vesicles by suppressing NPC-1. This provides LD-R amastigotes with excess lipids, supporting their rapid proliferation and membrane synthesis. This excess LDL-influx leads to an eventual accumulation of neutral lipid droplets around LD-R amastigotes, thereby increasing their unresponsiveness towards Amphotericin-B, a second-line amphiphilic antileishmanial. Notably, VL patients showing relapse with Amphotericin-B treatment exhibited significantly lower serum LDL and cholesterol than cured cases. Treatment with Aspirin, a lipid droplet blocker, reduced lipid droplets around LD-R amastigotes, restoring Amphotericin-B responsiveness.

Keywords: Clinical Leishmania donovani isolates / drug-unresponsiveness / Fluid phase endocytosis /vesicle fusion/LDL-cholesterol / Lipid droplets

Introduction

Antimony, which was mainstay for treating fatal visceral leishmaniasis (VL), has been withdrawn from Indian Subcontinent for more than a decade now due to emergence of resistance. Despite this, recent clinical LD isolates remain unresponsive to antimony (LD-R), indicating persistent phenotype [1]. Notably, LD-R also exhibits more aggressive pathology in clinical and experimental infection featuring increased organ parasite burden as compared to LD-S-infection [2-5]. Increased metacyclogenesis among LD-R was initially thought to be the primary factor contributing to higher infectivity, resulting in organ-parasite overload [6]. However, later studies revealed that LD-R promastigotes have a greater replicative potential than LD-S, possibly offering a selective intracellular survival advantage [7, 8]. This implies LD-R might be more adept at acquiring and metabolizing host derived nutrients within their parasitophorous vacuole (PV), facilitating their heightened intracellular proliferation. Lipids are crucial carbon sources for energy in proliferating intracellular pathogens. Several intracellular bacteria, protozoans and viruses utilizes cellular lipid droplets and plasma lipoproteins for entry, energy acquisition and biomembrane synthesis, demonstrating intricate interactions between intracellular pathogens and host lipid metabolism [9, 10]. *Leishmania* parasites are cholesterol auxotrophs [11], and need (i) host membrane cholesterol for successful attachment and infection [12]. (ii) Intracellular leishmania amastigotes replicating within membrane-bound PVs, accumulates cholesterol and lipid-droplets [13]. (iii) *Leishmania* amastigotes isolated from infected host have been shown to have increased cholesterol content in them [14], all these clearly suggest replicating *Leishmania* amastigotes might co-opt host derived lipids to facilitate their intracellular proliferation and membrane synthesis. In fact, hypocholesteremia is a hallmark of VL, with intracellular LD depleting host cholesterol and lipoproteins [15, 16]. However, the mechanism by which PV-bound LD amastigotes acquire this lipids, and how variations in lipid type, source, and utilization dynamics affect disease pathology and drug responsiveness, remains unclear.

So far, drug responsiveness in leishmaniasis has mostly been studied through in vitro selection of drug-resistant promastigotes, with limited focus on the drug susceptibility profiles of circulating clinical strains of *Leishmania* [17-19]. However, growing evidences indicates that drug responsiveness and treatment failure in VL are multifactorial, influenced by the host's immune-metabolic status [3, 20]. Importantly, recent reports have also linked emergence of drug unresponsiveness among intracellular pathogens with their ability to modulate host lipid metabolism [21, 22]. For instance, increased lipid accumulation can hinder drug penetration by trapping lipophilic drugs, fostering resistance [23-25]. VL treatment in recent years have seen a rise in treatment failure against Amphotericin-B (Amp-B), a second-line lipophilic drug [2], making it critical to investigate these failures in the context of lipid utilization during LD infection.

Using clinical antimony unresponsive LD-strains isolated from patients this study elucidates how LD-R-infection leads to an increased uptake of low-density lipoprotein (LDL) leading to eventual accumulation of neutral lipids droplets within their PV. In vitro infection of liver-resident Kupffer cells (KCs), primary sites for cholesterol synthesis and LD amastigote residence, demonstrated that LD-R infection facilitates fluid-phase endocytosis, increasing LDL-vesicle uptake in infected-KCs cells. LD-R-PV exhibits a higher propensity to fuse with endocytosed LDL-vesicles, coupled with the inhibition of cholesterol recycling from the vesicles back to the plasma membrane of Infected-MΦs. Our observation thus explains the heightened energy source and cholesterol availability for PV and daughter membrane synthesis to accommodate higher number of LD-R amastigotes observed in clinical infection. We also observed a suppressed expression of MSR-1, a scavenger receptor of ox-LDL [26], in LD-R infected murine liver, suggesting LD-R-infection limits the uptake of ox-LDL,

reducing inflammatory responses. Furthermore, our findings indicate that lipid droplet accumulation
around LD-R amastigotes contributes to Amp-B unresponsiveness, explaining recent reports of
Amp-B cross-resistance in clinical LD-R isolates from India [27, 28]. Significant low lipid profiles in
VL patients experiencing treatment failure against liposomal Amp-B, combined with complete
clearance of LD-R amastigotes using a combination treatment of Aspirin (a lipid droplet formation
blocker) and Amp-B, confirmed critical role of assimilating host lipids in developing Amp-B
unresponsiveness among rapidly replicating LD amastigotes exhibiting primary unresponsiveness
towards antimony. This study identifies sequence of events that enables LD-R amastigotes to
redirect host nutrients for their own aggravated sustenance, and how this preferential lipid utilization
might equip them with unresponsiveness against Amp-B contributing to the gradual spread of
cross-resistance among clinical LD-R isolates.
111
112 **Results**
113
1.   LD-R infection caused severe dyslipidaemia as compared to LD-S infection
Intracellular amastigotes of *Leishmania* parasites exhibit lipid auxotrophy [11], with dyslipidaemia
being a hallmark for leishmaniasis [15], including VL patients. Previously, LD amastigote load in
splenic aspirates of VL patients have been shown to exhibit an inverse correlation with serum
cholesterol level [29]. Since infection with antimony-unresponsive-LD (LD-R) has been previously
reported to exhibit higher amastigote load in spleen and liver of infected host, as compared to
infection with their responsive counterparts (LD-S) [7, 30], we wanted to investigate possible extent
of dyslipidaemia in response to LD-R and LD-S-infection and how it might affect disease pathology.
Mice infected with equal number of sorted metacyclics (**Fig S1A**) of natural clinical LD-isolates with
confirmed LD-R phenotype, MHOM/IN/09/BHU575/0 (LD-R[1]) and MHOM/IN/10/BHU814/1 (LD-
R[2]), showed a significantly lower serum levels of LDL, HDL, and cholesterol as compared to
MHOM/IN/83/AG83 (LD-S[1]) and MHOM/IN/80/DD8 (LD-S[2]), infected mice (**Figure 1A).** Liver
serves as a primary site for lipid synthesis and liver-resident MΦs (Kupffer cells, KC) are one of the
primary sites harboring LD amastigotes [31]. Therefore, we decided to test if KCs can be
established as an *in vitro* model for LD-infection to investigate the possible link of host lipid
utilization with respect to aggressive pathology shown by LD-R. Purified KCs (**Figure S1B**) infected
with equal number of sorted metacyclics for LD-S and LD-R-isolates as before displayed no
significant difference in their initial attachment to host membrane (**Figure 1B** and **Movie 1A, 1B).**
However, although initial number of intracellular LD-S and LD-R-amastigotes 4Hrs post infection
(p.i) showed no difference, a significantly higher number of LD-R amastigotes were observed in
24Hrs infected-KC (**Figure 1Ci, ii**). Parallel infection performed in peritoneal exudate macrophages
(PEC), which serves as a long-standing model for *in vitro* LD-infection, under identical experimental
conditions also resulted in similar observation with LD-R strains showing significantly higher
amastigote load at 24Hrs p.i than LD-S strains (**Figure S1Ci, ii, iii**), as previously reported by
multiple groups [1, 4, 6]. Moreover, KCs infected with LD-S or LD-R amastigotes isolated from 28
days infected mouse spleens, also resulted in significantly higher parasite load for LD-R (**Figure
S1D**), clearly indicating that higher parasite load is a conserved LD-R phenotype which does not
depends on initial infectivity or MΦ origin. Notably, infected-KCs had a slightly higher inherent
amastigote load compared to infected-PECs for both LD-S and LD-R-infections, with no significant
difference in parasite load within the two independent LD-S or LD-R strains (**Figure 1C, Figure
S1C, S1D**). Hence, BHU575 was selected as representative LD-R and AG83 as representative LD-
S to conduct all further comparisons with the inclusion of other representative strains as and when
mentioned. Higher intracellular amastigote load in response to LD-R-infection was further

confirmed by imaging LD-S and LD-R-PV at 4Hrs and 24Hrs infected-KCs (**Figure 1D**).
Comparative analysis revealed significantly higher number of LD-R-PVs as compared to LD-S-PVs
at 24Hrs p.i, although this difference was not observed at 4Hrs p.i (**Figure 1Di, Dii**). Interestingly,
3D image reconstruction for LD-R-PV at 24Hrs p.i, presented it as a larger fused structure as
compared to smaller individual segmented appearance of LD-S-PV (**Figure 1Diii**).
152
2.   Lipid deprivation is detrimental for LD-R while inherent hypercholesterolemia
facilitates its proliferation
To determine if extracellular lipids are absolutely essential for proliferation of LD-R amastigotes,
LD-S and LD-R-infections were performed in high, low or lipid-free medium. Reduction in
extracellular lipids severely compromised proliferation of LD-R but not LD-S amastigotes in
infected-KCs, confirming LD-R's dependency on external lipid availability (**Figure 2A i, ii**).
Alternatively, infection performed in KCs isolated from ApoE-knockout (KO)-mice($Apoe^{-/-}$) with
inherent hyperlipidemia [32], resulted in significant increase in amastigote load as compared to
wild type BL/6-infected-KCs, only for LD-R-infection, with LD-S infection resulting in comparable
amastigote load (**Figure 2B i, ii**). Previously, LD-S-infections in $Apoe^{-/-}$ mice have been reported
to typically get cleared by 42 days p.i, due to heightened T-cell activity [20]. Interestingly, however,
we observed $Apoe^{-/-}$ mice infected with GFP expressing LD-S or LD-R resulted in increased splenic
parasite burden as compared to wild type BL/6 mice at an earlier time point,11 days p.i (**Figure 2C**
i, ii), This observation possibly indicates a higher initial infection propensity in $Apoe^{-/-}$ mice, which
might get resolved later on due to increasing T-cell activity. Our observation also showed that
splenic parasite load for LD-R is significantly higher in $Apoe^{-/-}$ than BL/6 (**Figure 2C ii, iii)** again
confirming that higher availability of extracellular lipids might help in increased proliferation of LD-
R amastigotes.
171
3. Quenching of host membrane cholesterol remains consistent during initial infection with
LD-S and LD-R
Host lipids scavenging is essential for any intracellular pathogens exhibiting lipid auxotrophy [22].
Thus, to support their aggressive division, LD-R amastigotes should require significantly higher
amount of host lipids, which in turn will allow excess supply of cholesterol for increased PV and
daughter membrane biogenesis as compared to LD-S amastigotes. *Leishmania* amastigotes are
lipid auxotrophs [11, 14], and it has been previously reported that *L. Mexicana* can acquire host
membrane cholesterol during its initial entry into host MΦs [33]. Thus, host membrane Cholesterol
was investigated as a possible source and potential mechanism by which LD-R acquires host-
derived lipids distinctively from their LD-S counterparts. However, confocal image analysis of NBD-
Cholesterol labeled KCs infected with RFP expressing LD-S or LD-R revealed that both LD-S and
LD-R can extract NBD-Cholesterol from the KC membrane during their initial infection (4Hrs p.i)
with comparable efficiency (**Figure S2A i, ii**). This observation was further confirmed by
quantification of NBD-Cholesterol between freshly invaded LD-S and LD-R parasites isolated from
infected-KC by flow cytometry (**Figure S2B i, ii**).
187
4. LD-R-infection mobilizes extracellular LDL within infected-KCs to support its intracellular
proliferation
Since the quenching of host membrane cholesterol during initial entry of LD promastigotes does
not significantly differ in LD-S and LD-R mediated infections (**Figure S2**), we explored other
potential host lipid sources for supporting LD-R's increased intracellular division and membrane
synthesis. LDL serves as the primary source of cellular lipids which helps to maintain lipid

homeostasis, and gets readily metabolized to produce bioavailable cholesterol [34]. Our previous
observation showing a significant drop in LDL levels between LD-S and LD-R-infected mice (**Figure**
1), led us to investigate LDL as a possible source which can differentially contribute in providing
excess lipids, sterols, and fatty acids to rapidly proliferating intracellular LD-R amastigotes. Live
cell imaging comparing uptake of fluorescent red-LDL in EGFP-LD-S or EGFP-LD-R-infected KCs
at 24Hrs p.i revealed a significantly higher red-LDL accumulation within EGFP-positive LD-R-
infected KCs as compared to LD-S-infection (**Figure 3Ai**). Fluorescence plate reader-based
quantification further supported this observation showing significantly higher LDL accumulation in
LD-R-infected-KCs over time, which was not observed by performing infection with killed parasites
(**Figure 3A ii**), indicating that, LDL-influx is mediated by replicating LD amastigotes. Confocal
microscopic images at 24Hrs p.i. of LD-S and LD-R-infected-KCs confirmed significantly higher
LDL co-localizing with LD-R amastigotes compared to LD-S amastigotes (**Figure 3B i, ii**).
206
5.   LD-R-infection results in a distinct metabolic shift in lipid peaks of infected host
Next, we wanted to confirm if higher LDL-uptake could result in accumulation of excess lipids,
particularly cholesterol, in LD-R-PV that would provide LD-R amastigotes with higher replicative
capabilities and associated enhanced membrane synthesis potential. PV fractions were purified
from LD-S and LD-R-infected-KCs post 24Hrs infection, as reported previously [35]. Quantitative
analysis of total cholesterol using Amplex red kit (**Figure S3A)** and thin layer chromatography (TLC)
(**Figure S3B**) revealed significantly higher cholesterol content in LD-R-PV as compared to LD-S-
PV. Furthermore, GC-MS of LD-S and LD-R-PV under identical experimental condition identified
prominent peaks for cholesterol and fatty acids in LD-R-PV which were missing for LD-S-PV
(**Figure 3C i, ii**). Apart from this biochemical characterization, we also performed biophysical
analysis of lipid related peaks using non-invasive Confocal Raman spectroscopy to capture a
snapshot of lipid distribution in LD-S and LD-R-infected-KCs under native conditions (**Figure 3D i,**
ii). This analysis highlighted a distinct clustering of lipid peaks representing cholesterol,
triglycerides, and fatty acids [36] around LD-R amastigotes, contrasting with their minimal clustering
around LD-S amastigotes which resemble uninfected-KCs more closely (**Figure 3D i**). Raman
Shift analysis within the biologically active region (300cm^{-1}-1800cm^{-1}) [37], further confirmed a
significant difference in relative abundance for lipid related peaks' at wavelengths: 540-560 cm^{-1}
(cholesterol), 1080-1090 cm^{-1} (phospholipids), 1270-1280 cm^{-1} (triglycerides), 1300-1340 cm^{-1}
(amide-III bond), 1440-1453 cm^{-1} (fatty acids and triglycerides) [36, 38], along with olefinic stretch
of C=C double bonds at 1,655 cm^{-1} between LD-S and LD-R-infected-KC, at 24Hrs p.i. **(Figure 3D**
ii and S3C i, ii). We also noticed a distinct shift by plotting area under the curve from (Right to Left)
in response to LD-R-infection (**Figure S3 D**), indicative of distinctly different lipidomes [39].
229
6. LDL-uptake in LD-infected-MΦs occurs through fluid phase endocytosis
As LD-R amastigotes heavy reliance on increased host lipid acquisition for rapid proliferation
became evident, we wanted to investigate molecular basis of this acquisition. Considering that
MΦs primarily uptake LDL via the LDL receptor (LDLr) [40], we checked its status in response to
LD-S and LD-R-infection. However, immunofluorescence analysis showed a decrease in LDLr
expression in infected-KCs harboring LD-amastigotes as compared to adjacent uninfected KCs,
regardless of infection type (**Figure 4A**), along with no significant changes in total LDLr expression
between uninfected, LD-S, and LD-R-infected-KCs, as observed from Western blot (**Figure S4A**).
This indicated that LDL-uptake by LD-infected-KCs might be receptor-independent. Our attempts
to generate siRNA mediated Knock down (KD) in KCs were unsuccessful, hence to convincingly
determine the role of LDLr in LD-infected-MΦs we generated LDLr-KD-PEC (**Figure S4B i**) and

performed infection with LD-S or LD-R. Infecting LDLr-KD-PEC with LD-S or LD-R resulted in no
significant changes in intracellular amastigote numbers for both infection types as compared to
scrambled RNA control (**Figure S4B ii**) confirming that although LD-R relies on increased LDL-
influx for heightened proliferation, this influx is LDLr-independent. Apart from LDLr, MΦs can also
acquire LDL through receptor-independent fluid phase endocytosis [41]. To confirm this possibility,
we checked the status of Cofilin, an actin modulating protein, which serves as a marker for fluid
phase endocytosis. Western blot analysis of infected-KCs revealed an increase in Cofilin levels
with a concurrent decrease in its phosphorylated form as LD-R-infection progressed, indicating
active fluid phase endocytosis (**Figure 4Bi, ii**). In contrast, LD-S-infection did not induce such
changes (**Figure 4B i, ii**). Interestingly, this decrease in Cofilin phosphorylation was also observed
in LD-R-infected-PEC (**Figure S4C i, ii**), signifying that enhanced fluid phase endocytosis may be
a conserved mechanism for facilitating LDL-influx in LD-R-infected-MΦs. This enhanced fluid phase
endocytosis in response to LD-R-infection was further confirmed by structural illumination imaging
(SIM), which showed increased Cofilin turnover with dispersed distribution in LD-R-infected-KCs,
leading to enhanced F-Actin rearrangement and protrusion formation [42], which was absent for
uninfected or LD-S-infected-KCs (**Figure 4Ci, ii and Figure 4Di, ii**). Finally, we infected the KCs
with GFP expressing LD-R for 4Hrs, washed and allowed the infection to proceed in presence of
fluorescent red-LDL and Latrunculin-A, a compound which specifically inhibits fluid phase
endocytosis by inducing actin depolymerization [41]. Real-time fluorescence tracking demonstrated
that Latrunculin-A treatment not only prevented the uptake of fluorescent red-LDL but also severely
impacted intracellular proliferation of LD-R amastigotes (**Movie 2A and 2B and Figure 4E**). In
contrast, treatment with Cytochalasin-D, which alters cellular F-actin organization but does not
affect fluid phase endocytosis [41], had no effect on the intracellular proliferation of LD-R (**Figure
4F and Figure S4D**).
265
7.   Endocytosed-LDL traffics and fuses with and LD-R-PV with increased proficiency
Typically, LDL-vesicles fuses with lysosomes upon entry into MΦs, leading to subsequent transfer
of LDL-derived cholesterol into the endoplasmic reticulum and cytoplasm [43]. As we have already
confirmed an increased influx with higher presence of endocytosed-LDL vesicles around LD-R
amastigotes along with significantly higher accumulation of cholesterol and fatty acids in LD-R-PV
as compared to LD-S-PV (**Figure 2**), we hypothesized that the mode of trafficking of endocytosed-
LDL might differ between LD-S and LD-R-infection. LD-PV are modified phagolysosomes [35, 44],
and we observed an increased convergence of LDL-vesicles with LAMP-1 positive LD-R-PV even
at 4Hrs p.i, while such convergence was not noticed in LD-S-infected-KC (**Figure 5A i, ii**).
Structured Illumination Microscopy confirmed a complete convergence of LDL-vesicles with LD-R-
PV at 24Hrs p.i, while LD-S-PV showed only partial convergence with LDL-vesicles even at this
late infection point (**Figure 5B i, ii).** Interestingly, while, Western blot analysis revealed minimal
expression of LAMP-1 in response to LD-S-infection at 4Hrs p.i, which increased considerably as
the infection progressed (**Figure 5C**), however, a considerable amount of LAMP-1 expression was
observed for LD-R-infected-KCs from 4Hrs p.i, which also increased as infection progresses
(**Figure 5C ii**), indicating that a higher fusion of LDL-vesicles in LD-R-PV may be accompanied by
increased lysosomal biogenesis required for degrading LDL into cholesterol. Importantly, Filipin
staining of EGFP-LD-S and EGFP-LD-R-infected-KCs at 24Hrsp.i confirmed enriched cholesterol
staining around LD-R amastigotes with significantly higher co-localization coefficient for LD-R
amastigotes (**Figure 5D i, ii, iii**).
286
8.   LD-R-infection suppress release of metabolized cholesterol from PV

Once LDL is broken down to cholesterol by lysosomal acid hydrolases, metabolized cholesterol is
released back into cytosol from LDL-vesicles by two transmembrane proteins namely NPC-1 and
NPC-2 [43]. We suspected that LD-R must prevent or delay this process as this will limit cholesterol
accessibility to replicating LD-R amastigotes. Analysis of available *RNAseq* (INRP000146) data for
uninfected, LD-S and LD-R-infected-PEC (**Figure 6A)** revealed a suppression of *npc-1* expression
along with its transcription factor *srebp2* [45], in response to LD-R-infection. Western Blot analysis
of LD-S and LD-R-infected-KCs confirmed this low expression of SREBP2 and NPC-1 in LD-R-
infected-KC 24Hrs p.i (**Figure 6Bi, ii**). More interestingly, Confocal imaging not only confirmed this
low expression of SREBP2, but even in those cases of LD-R-infected-KCs where we did observe
some residual expression of SREBP2, it was entirely restricted to cytoplasm, unlike LD-S or
uninfected-KCs where SREBP2 did get translocated in the nucleus (**Figure 6C**). For NPC-1,
immunofluorescence imaging comparing adjacent uninfected and infected-KCs harboring LD-S or
LD-R, clearly confirmed unlike LD-S, LD-R specifically suppresses NPC-1 to prevent release of
LDL-derived cholesterol from fused vesicles making it accessible to amastigotes (**Figure 6D**). We
confirmed this observation by performing a Total Internal Reflection Microscopy (TIRF) which
clearly showed a low plasma membrane cholesterol for LD-R-infected-KCs as compared to LD-S-
infected-KCs (**Figure 6Ei**) and subsequently revalidated this by quantifying the cholesterol content
in isolated membrane fraction for individual experimental set as mentioned above (**Figure 6E ii**).
Finally, siRNA mediated knock down of NPC-1 in PEC (due to failure of successful KD generation
in KCs), resulted in significant increase in amastigote load in response to LD-R-infection, while no
such change in amastigote load was observed for LD-S-infected NPC-1-KD-PECs (**Figure 6F i, ii,**
iii).
310
9.   LD-R-Infection specifically excludes oxidized-LDL in infected murine liver to limit
inflammation
The increased accumulation of LDL within MΦs typically triggers an inflammatory response [46].
This possessed an inherent conceptual dilemma to our observation as increased inflammatory
response should be detrimental for LD-R amastigotes [47, 48]. To check this, we measured
inflammatory IFN-γ by co-culturing splenic T-cells isolated from LD-S or LD-R-infected mice, with
LD-S and LD-R-infected-KCs, respectively (**Figure 7 upper left panel**). Surprisingly, our ELISA
results showed a low IFN-γ levels in the culture supernatant of LD-R-infected-KCs even after 24Hrs
of co-culture, while for LD-S-infection a significant increase was noted at this point (**Figure 7Ai**).
Similarly, stimulation of LD-infected splenocytes obtained from LD-S and LD-R-infected *Apoe−/−*
mice with specific soluble leishmania antigens (SLA) of LD-S or LD-R also confirmed this restricted
IFN-γ response to LD-R-infection (**Figure 7A ii)**, thus suggesting that although LD-R-infection
results in increased accumulation of lipids in host MΦs, it elicits a suppressed inflammatory
response as compared to LD-S. Notably, accumulation of oxidized-LDL (ox-LDL), and not its
neutral form has been reported to cause inflammatory response [46, 49]. LD-S or LD-R-infected-
KCs incubated with ox-LDL showed a significant purple positive staining of ox-LDL only for LD-S-
infected-KCs (**Figure 7B i**), with LD-R-infected-KCs seemingly inhibiting uptake of ox-LDL.
Infecting KCs with higher number of LD-S or lower number of LD-R did not alter their ox-LDL uptake
capabilities, clearly suggesting LD-R-infection inherently suppress ox-LDL uptake, irrespective of
the number of intracellular amastigotes (**Figure 7B. ii,**). Although, receptor of ox-LDL, MSR-1 [26],
have shown a slightly lower expression in LD-R-infected-PEC in the *RNAseq* analysis (**Figure 6A**),
however owing to reported dominant expression of *msr-1* in the liver [49], we checked expression
of MSR-1 in the LD-S and LD-R-infected-KCs. Confocal images and Western blot analysis
confirmed a significant lower expression of MSR-1 in LD-R-infected-KCs as compared to LD-S-

infected-KCs (**Figure 7C i, ii**), thus explaining reduced staining for ox-LDL in response to LD-R
infection. Furthermore, increased expression of *atp6vod2* transcripts, implicated in LD-PV
acidification and lipid utilization [35], was also noted to be slightly higher for LD-R-infected-PECs
in the *RNAseq* (**Figure 6A**). However, western blot analysis showed a slightly increased expression
of ATP6V0D2 in LD-R-infected-KCs as compared to LD-S-infected-KCs (**Figure 7D**), suggesting
that it might also play a role in maintaining LD-PV acidity, facilitating lipid utilization and limiting ox-
LDL uptake, as previously reported for *L. mexicana* infection [35]. Finally, immunostaining of liver
sections from 28 days LD-S and LD-R-infected-mice confirmed a significant lower expression of
MSR-1 in LD-R-infected liver with higher accumulation of neutral lipid droplets as compared to LD-
S-infected murine liver **(Figure 7E i, ii),** thus confirming LD-R specifically accumulates neutral lipids
by restricting MSR-1 expression.
346
10.   Lipid droplets accumulation decreases Amphotericin-B responsiveness in LD-R
amastigotes
Increased LDL-uptake has been reported to cause high lipid droplets formation [50, 51], and lipid
droplets formation has been reported to play critical role in intracellular proliferation of several
pathogens, including LD-amastigotes [52-54]. As we have already observed significant
accumulation of lipid droplets in LD-R-infected murine liver (**Figure 7**), we further investigated its
role in response to LD-S and LD-R-infection. Lipid droplets staining of LD-S and LD-R-infected-
KCs revealed significantly higher lipid droplets accumulation around LD-R amastigotes (**Figure**
S5A and **Movie 3).** For better visualization**,** we expanded [55] EGFP-LD-R-infected-KCs and again
performed lipid droplets staining on them which confirmed assimilation of neutral lipid into LD-R
amastigotes (**Figure 8A**). This was further confirmed by performing TEM in LD-R-infected-KC
which showed Lipid droplets accumulating around LD-R amastigotes **(Figure 8B)**. Lipid
accumulation has been previously reported to cause Amphotericin-B unresponsiveness among
pathogens by hindering drug penetration [23]. Interestingly, we noted that two LD-R-strains
(BHU575 and BHU814) which we have included in this study also shows reduced responsiveness
against Amp-B, a second-line-antileishmanial [56], and we validate this by determining their EC50
against Amp-B in LD-infected-KCs, which showed an EC50 of 0.39 µM and 0.31 µM for LD-R[1] and
LD-R[2] amastigotes respectively, as compared to EC50 of 0.03 µM for LD-S[1] and 0.032 µM LD-S[2].
We suspected that significant lipid droplets accumulation around LD-R amastigotes might be
responsible for their reduced susceptibility towards Amp-B by limiting drug accessibility. To test this
possibility, we performed independent infection with two LD-R strains in low lipid and high lipid
medium and access their EC50 against Amp-B. Low lipid medium results in significant drop in lipid
droplets accumulation within LD-R-infected-KCs (**Figure S5B**) and also resulted in enhanced
susceptibility for both the LD-R strains against Amp-B with EC50 reduced to 0.39 µM to 0.16 µM
for LD-R[1] and 0.31 µM to 0.06 µM for LD-R[2] as compared to high lipid medium (**Figure 8C i, ii**).
Interestingly, infection with lab-generated Amphotericin-B-unresponsive LD promastigotes (LD-
S[Amp-B-R]) without primary unresponsiveness towards antimony did not result in any significant lipid
droplets accumulation around LD-R amastigotes (**Figure S5A**) and neither showed any significant
alteration in EC50 between low (0.09 µM) and high lipid medium (0.11 µM) (**Figure 8C iii**). This
observation clearly suggests that this high LDL-influx along with higher lipid droplet accumulation
responsible for increased unresponsiveness of LD-R isolates towards Amp-B is a specific attribute
of LD isolates with primary unresponsiveness against antimony and may vary significantly from lab
generated Amp-B unresponsive strains. Previously, Aspirin (Asp), a lipid droplet blocker has been
reported to block lipid droplet formation in *T.cruzi* and LD-infected-MΦs [52, 57]. We also observed
a drastic drop in lipid droplets formation in LD-R-infected-KCs treated with Aspirin as compared

untreated LD-R-infected-KCs (**Figure S5C).** Interestingly, while treatment with Aspirin alone did
not significantly reduce the amastigote load in LD-R-infected-KCs, combining Aspirin with
previously determined low dose of Amp-B (0.1 µM), completely restored Amp-B responsiveness
against intracellular LD-R amastigotes (**Figure 8E, F**). This suggests that the enhanced uptake of
LDL leading to lipid droplet accumulation might be a primarily responsible for increasing Amp-B
unresponsiveness among clinical LD-R isolates. Importantly, this observation is consistent with
findings from human VL patients from the endemic region of Bihar, India, where those who relapsed
after Amp-B treatment had significantly lower serum LDL and cholesterol levels compared to Amp-
B treatment responders (**Figure 8G and Table 2**).
391
Discussion
Our findings revolve around understanding how antimony-resistant *Leishmania donovani* (LD-R)
has evolved as a persistent strain by adopting to a strategy of better utilization of host derived lipids
to sustain their own aggressive intracellular proliferation. Intracellular pathogens often alter the
host's immune-metabolic state for better survival [58, 59]. Our observations pointed out that high
organ parasite load linked with LD-R-infection does not rely only on initial infectivity as commonly
believed to date [7], rather is indicative of a selective higher intracellular proliferation in host. We
observed LD-R amastigotes scrutinize and access specific type of host lipoproteins to support their
aggressive proliferation as seen in clinical infection (**Figure 1, S1**) [15]. Previously increased
amastigote load in VL patients has been inversely linked with their serum cholesterol level [29], and
we also observed that mice infected with LD-R isolates had significantly lower HDL, LDL, and
cholesterol compared to LD-S-infected mice (**Figure 1**).
LD amastigotes are lipid auxtrophs [11, 14], and this severe dyslipidemia in response to LD-R-
infection can be logically interpreted in terms of LD-R's increase need to support their aggravated
intracellular proliferation in the infected-host. In liver-resident macrophages (Kupffer cells), primary
sites for LD infection and host cholesterol synthesis, we observed significantly higher number of
LD-R amastigotes and parasitophorous vacuoles formation (PV), compared to LD-S-infection,
indicating LD-R's better capacity to utilize host cholesterol (**Figure 1, S1**). Also, in hyperlipidemic
Apoe$^{-/-}$ mice, LD-R-infection led to organ parasite overload compared to LD-S-infection, confirming
LD-R's increased dependence on extracellular lipid sources to sustain its aggressive proliferation
(**Figure 2**). Thus, next we focused on investigating LD-R's ability to differentially regulate host's
immune-metabolic status. Previous reports suggested that LD with altered drug-responsiveness
can modulate their host differentially resulting in different disease outcome [3, 60]. Our results
confirmed that while both LD-R and LD-S promastigotes initially quench cholesterol from the host
plasma membrane similarly (**Figure S2**), LD-R amastigotes accumulate significantly more
cholesterol and fatty acids in their PV by specifically promoting increased endocytosis of LDL as
the infection progresses (**Figure 3, S3**). This possibly also explains the similar initial attachment
and infectivity of LD-S and LD-R promastigotes with the host MΦs again indicating towards their
ability to infect their host equally (**Figure 1, S1, Movie 1**).
Interestingly, this LD-R-induced LDL-endocytosis is independent of the host's LDL receptor and
relies on fluid-phase endocytosis, involving significant actin cytoskeleton reorientation in LD-R-
infected-MΦs (**Figure 4, S4**). It is worth mentioning that LD-R-infection has been reported to induce
high levels of IL-10 [3], which is linked with enhanced fluid-phase endocytosis [41].
Importantly, we also noted an increased convergence of endocytosed LDL-vesicles with lysosomal
associated marker protein (LAMP-1), along with higher LAMP-1 expression around LD-R
amastigotes as compared to LD-S ones (**Figure 5**). Lysosomal acid hydrolases metabolize LDL
into bioavailable cholesterol, and higher fusion of lysosomal vesicles with LD-R-PV explains the

significant Filipin-cholesterol staining around LD-R amastigotes (**Figure 5**). These findings not only confirm that endocytosed-LDL can be better utilized by LD-R amastigotes but probably also indicates towards a rapid maturation of LD-R-PV due to increased LDL-vesicle fusion [13, 35].

Although available *RNAseq* analysis **(Figure 6**) did not support this increased expression of *lamp-1* in the transcript level, it did reflect a notable upregulation of vesicular fusion protein (VSP) *vamp8* and *stx1a* in response to LD-R-infection. LD infection can regulate LAMP-1 expression, and the role of VSPs in LDL-vesicle fusion with LD-R-PV is worthy of further investigation. Additionally, RNAseq analysis also showed suppressed expression of *npc-1* in LD-R-infected-MΦs. NPC-1 is crucial for mobilizing vesicular cholesterol to the cytoplasm and plasma membrane [43], and its suppressed expression correlates with higher cholesterol availability for aggressive LD-R proliferation (**Figure 6**).

Besides LDL endocytosis, *de novo* cholesterol synthesis could also serve as a lipid source for intracellular LD amastigotes [61]. Surprisingly, LD-R-infected-MΦs showed suppressed expression of SREBP2, with minimal nuclear translocation compared to LD-S-infection (**Figure 6**), consistent with reports of a shutdown in de novo cholesterol synthesis in MΦs with high endocytosed LDL accumulation [62]. Notably, SREBP2 regulates NPC-1 expression [45] and has been and has been shown to get activated in response to LD-S infection [62].

Surprisingly, LD-R-Infected-MΦs showed minimal inflammatory response despite high lipid accumulation, due to their ability to successfully exclude uptake of oxidized LDL (ox-LDL). Low expression of MSR-1, a scavenger receptor of ox-LDL, in the liver of LD-R-infected mice explained this observation, leading to the accumulation of neutral lipid droplets around LD-R amastigotes (**Figure 7, S5, Movie 3**). Expansion Microscopy (ExM) [55] along with TEM confirmed significant assimilation of these neutral lipids into LD-R amastigotes (**Figure 8**). Several intracellular pathogens like *Mycobacterium*, *Toxoplasma*, HCV have been reported to transfer stored triacylglycerols and sterol esters from lipid droplets to support their proliferation [10, 63, 64]. While the understanding of lipid uptake and metabolizing genes in LD is limited, a recent genome-wide CRISPR screen in *L. Mexicana* [65], have identified such potential candidates "essential" for parasite survival. Close proximity lipid droplets with LD-R amastigotes makes it tempting to speculate existence of lipid droplets contact sites [13], or possible copy number variations in these lipid metabolizing genes [66], providing survival advantages to LD-R. This speculation is further supported by the fact that although clinical LD-R isolates with neutral lipid droplet accumulation in infected MΦs displayed increased unresponsiveness to Amphotericin-B (Amp-B), lab-generated Amp-B unresponsive promastigotes failed to do so in infected MΦs (**Figure 8**). Fact that LD-R amastigotes regained their responsiveness towards Amp-B in low lipid medium or in presence of Aspirin [52], further confirms that eliminating lipid droplets around LD-R amastigotes might be a weapon of choice to combat treatment failure against Amp-B [27, 28, 67]. Our results thus suggest that natural clinical LD-R-isolates have developed an inherent adaptability to gradually develop cross-resistance towards Amp-B, probably as an additional consequence of their ability to accumulate increased host derived lipids as a rapidly proliferating intracellular amastigote (represented graphically in **Figure 9**). Our work explains increasing Amp-B treatment failure among VL patients [27, 28, 67] and correlates with low serum LDL and cholesterol levels in patients relapsing after Amp-B treatment (**Figure 9, Table 2**).

476
Methods
Reagents
Laurdan (Merck), photoRED-LDL (Thermo Fisher Scientific), NBD Cholesterol (Sigma), Nile red
(Sigma-Aldrich), Oil red O (Thermo Fisher Scientific), Filipin (Sigma-Aldrich), Collagenase (Gibco)
were purchased. Tissue culture plasticware were purchased from NUNC (Roskilde, Denmark).
Invitrogen Amplex-Red assay kit was purchased from Thermo Fisher Scientific. Primary antibodies
required for the experiments were anti-ATP6V0D2 (ABclonal), anti-MSR1 (ABclonal), anti-LAMP1
(Thermo Fisher Scientific), anti-SREBP2 (Thermo Fisher Scientific), anti-NPC1 (Santa Cruz
Biotechnology), anti-LDLr (Santa Cruz Biotechnology), goat anti-mouse IgG Alexa Fluor 488
(Thermo Fisher Scientific), goat anti-mouse IgG Alexa Fluor 594 (Thermo Fisher Scientific), goat
anti-Rabbit IgG-HRP (Thermo Fisher Scientific), goat anti-Mouse IgG-HRP (Thermo Fisher
Scientific), β actin (Cell Signalling Technology), CD11b (Cell Signalling Technology), Cofilin (Cell
Signalling Technology), and phospho-Coffilin (Cell Signalling Technology). RhodaminePhalloidin
reagent was purchased from Thermo Fisher Scientific and ELISA Kits for IFN-γ were purchased
from R&D Systems, Aspirin generic composition, SAG (Albert David)
492
Animals
494 BALB/c, C57BL/6 and $Apoe^{-/-}$ mice were maintained and bred under pathogen-free conditions with
495 food supplements and filtered water. All experiments were conducted in accordance to Institutional
496 Animal Ethics Committee (IAEC), IIT Kharagpur guidelines India (IAEC/BI/1 55/2021) and
497 approved by National Regulatory Guidelines issued by Committee for the Purpose of Supervision
498 of Experiments on Animals, Ministry of Environment and Forest, Government of India.
499
Ethical statement
In this study, all procedures involving human participants were conducted in accordance with the
ethical standards of the 1964 Helsinki Declaration and approval of the "Institutional Human Ethical
Committee" of ICMR-Rajendra Memorial Research Institute of Medical Sciences, Patna, India
(Approval No: RMRI/EC/20/2020). Following information about the potential risks, benefits, and the
investigational nature of the study consent from all study participants was obtained in the informed
consent form.
507
Parasite Cultures and Maintenance
Leishmania donovani (LD) isolates used in this study are natural clinical isolates obtained from the
patients MHOM/IN/09/BHU569/0, MHOM/IN/2009/BHU575/0, MHOM/IN/10/BHU814/1,
MHOM/IN/83/AG83, MHOM/IN/09/BHU777/0, MHOM/IN/80/DD8 [2, 3], and were maintained in
BALB/c mice as per institute biosafety guidelines (BT/BS/17/27/97-PID). Susceptibility of each of
these LD isolates against antimony were checked and confirmed against previously reported EC50
value for antimony before they were included in the study (This can be reported in a common table,
Table 1). Lab generated Amp-B resistant strain (DD8-AmpB) strain was kind from Dr. Arun Kumar
Haldar (CSIR-CDRI). Amastigotes were isolated by macerating the spleen of 21 days infected mice
under sterile conditions, and subsequently transformed into promastigotes by maintaining in M199
(Gibco) supplemented with 5% (v/v) heat-inactivated fetal bovine serum (FBS) (Gibco) and 1% (v/v)
Penicillin Streptomycin solution (Gibco) under drug selection.
520
Cell Culture and Infection

Peritoneal exudate cells (PECs) were harvested from BALB/c mice as mentioned in already
published protocol, with slight modifications [68]. After 48Hrs of 4% starch intraperitoneal injection,
PECs were isolated and plated on 35-mm tissue culture petri dishes, sterile 22-mm square
coverslips in 35-mm petri dishes or 24-well plates at a density of 1×10^6, 1.5×10^5 or 1.5×10^5,
respectively, all in RPMI 1640 medium (Gibco) supplemented with 10% (v/v) heat-inactivated FBS,
1% (v/v) Penicillin Streptomycin solution. The cells were allowed to adhere for 48Hrs at 37°C in 5%
CO_2 before the infection process.
529
***In vivo* infection**
For *in vivo* infections, healthy female C57BL/6 mice or *Apoe*-/-mice (4-6 weeks old and 20-25 g
each) were injected via tail vein route with 1×10^7 GFP-expressing *L. donovani* stationary phase
metacyclic promastigotes (suspended in 100 µl 1X PBS) as reported previously [69]. After infection
mice were kept in atherogenic diet till sacrifice. Each group contained five mice that were sacrificed
11 days' post-infection (p.i). Splenocytes were isolated and GFP positive population was measured
through flow cytometry. Each experiment was performed in triplicate.
537
Isolation of Kupffer cells
Kupffer cells (KCs) were harvested from mice livers following a previously published protocol, with
slight modifications [70]. Wild-type BALB/c mice were anesthetized, and warm PBS containing
0.05% collagenase IV was used for liver perfusion. The livers were swiftly excised and minced into
small fragments on an ice bath. Tissue digestion was performed using a 0.1% Type IV Collagenase
solution in RPMI 1640 for 30 minutes at 37°C with intermittent shaking for every 10 minutes.
Following digestion, KCs were separated from hepatocytes and other cellular components through
sequential centrifugation steps and plated on well plates, as required, for 4Hrs in RPMI 1640 media
supplemented with 10% (v/v) Penicillin Streptomycin and 10% (v/v) FBS. After 24Hrs of resting at
37°C in 5% CO_2, the infection studies commenced. To assess the purity of the isolated KCs by flow
cytometry, KCs were plated on a 6 well plate. After 4Hrs, KCs were scraped out, centrifuged at
300g for 5 minutes and stained with monoclonal anti-F4/80 PE conjugated antibody (BD
Biosciences). The KCs were then washed and fixed with 2% paraformaldehyde for 5 minutes,
analyzed using a flow cytometer (ARIA FACS DIVA), and further processed using FLOW JO
software for data analysis. Purified KCs were infected with stationary phase sorted metacyclic LD
promastigotes as mentioned previously. PECs and KCs are collectively designated as MΦs
(macrophages).
555
Sorting of Metacyclic LD and infection in MΦs
Metacyclic parasites were sorted through Beckman Coulter CytoFLEX cell sorter, as per
manufacturer's protocol. The LD promastigotes were washed and resuspended in PBS before
being subjected to light scatter analysis using flow cytometry. Sorted population were checked with
forward scatter (FSC) versus side scatter (SSC) and dot plot was generated to represent the
acquisition of 10,000 events using a FACS Aria II system. Required number of sorted metacyclic
LD were used to perform subsequent MΦ and mice infections. MΦs were infected with stationary-
phase Flow sorted metacyclic LD promastigotes at a MOI 1:10 ratio (in some cases 1:5 and 1:20)
for 4Hrs which was considered as the 0^{th} point of the infection. MΦs was then washed to remove
any extracellular loosely attached parasites and incubated further as per experimental requirement.
566
Intracellular amastigote load determination

To enumerate intracellular amastigote load, resting stage PECs and KCs on glass coverslips were
infected with sorted metacyclic LD promastigotes of various strains at a ratio of 1:10, as described
in previous protocols [3, 71]. Following a 4Hrs incubation, the PECs or KCs were washed and
subsequently cultured for specific durations. Coverslips were retrieved after specific time points,
rinsed with PBS, fixed using methanol, and stained with Giemsa. These prepared coverslips were
then affixed onto glass slides and examined utilizing a light microscope (Olympus). Parasite
quantification was conducted by counting amastigotes per 100 macrophages. Image processing
related to Figure 1A and Figure S1 D was performed using Fiji by separating the spectral overlap
between the light and dark stained region to Giemsa stained MΦs. Next, for better visualization,
pseudo colours were assigned to represent nucleus (white) and cytoplasm (red) with a black
background. Colour separated images were then merged showing LD within PV.
579
Raman Spectroscopy of LD-infected-KCs
KCs were plated on Raman-grade stainless steel plate and kept for 24Hrs in a humidified
environment at 37°C with 5% CO_2, followed by infection with LD as already described. At 24Hrs
p.i., the KCs were washed using PBS, fixed with 2% paraformaldehyde and proceeded to perform
Raman spectroscopy (Witec, alpha300R). The Raman grade stainless steel plates were positioned
on the microscope stage and focused with a 100x air objective and measurements were performed
using a 532 nm laser source producing a laser spot of approximately 0.3 μm with 20mW laser
power. Spectral measurements were obtained through the Project FIVE interface (Witec, Germany)
after setting the grating to 600 gratings/mm. The spectral resolution was set to 2.87 cm^{-1}. The
Raman spectra were taken with integration time set to 5 seconds, and ten accumulations were
selected after confirming its effectiveness on the KCs being studied. Baseline correction and other
spectral denoising procedures were carried out using MATLAB 2017b (MathWorks), and the
processed spectra were analysed as described preciously [36, 38].
593
Quantification of membrane cholesterol with Amplex Red assay kit
The KC membrane preparation was conducted following the method described previously [20],
with slight modifications. KCs were ruptured by repeated freezing and thawing cycles followed by
probe sonicating for 4 cycles with 30secs burst and 30sec gap. The resulting homogenate was then
centrifuged at 900g for 10mins at 4°C. The supernatant was filtered through a nylon mesh (100
μm), while the pellet was discarded. The filtered supernatant was centrifuged at 100,000g for 1hr
at 4°C, and resulting pellet, representing native membrane, was suspended in a buffer containing
50 mM Tris (Ph 7.4), 1 mM EDTA, 0.24 mM PMSF, and 10 mM iodoacetamide. The protein content
of the membrane was quantified using Bradford reagent (Biorad), following manufacturer's protocol.
The total cholesterol content was determined using an Amplex Red reagent kit [72], with the results
expressed in moles of cholesterol per gram of protein.
605
Parasitophorous vacuole (PV) Isolation and cholesterol measurement
24Hrs LD infected KCs were lysed by passing through a 22-gauge syringe needle to release cellular
contents. Parasitophorous vacuoles (PV) were then isolated using a previously outlined protocol
[73]. Cholesterol and protein contents with PV were determined by an Amplex Red assay kit and
Bradford assay, respectively. Resulting data were represented as micrograms of cholesterol per
microgram of protein.
612
GC-MS analysis of LD-S and LD-R-PV

Following a 24Hrs infection period, KCs were harvested, washed with phosphate-buffered saline
(PBS), and pelleted. Subsequent to this, PV isolation was carried out using the previously described
method [74]. The resulting PV pellet was suspended in 20 ml of dichloromethane:methanol (2:1,
vol/vol) and incubated at 4°C for 24hours. After centrifugation (11,000 g, 1 hour, 4°C), the
supernatant was checked through thin layer chromatography (TLC) and subsequently evaporated
under vacuum. The residue and pellet were saponified with 30% potassium hydroxide (KOH) in
methanol at 80°C for 2 hours. Sterols were extracted with n-hexane, evaporated, and dissolved in
dichloromethane. A portion of the clear yellow sterol solution was treated with N, O-
bis(trimethylsilyl)trifluoroacetamide (BSTFA) and heated at 80°C for 1 hour to form trimethylsilyl
(TMS) ethers. Gas chromatography/mass spectrometry (GC/MS) analysis was performed using a
Varian model 3400 chromatograph equipped with DB5 columns (methyl-phenylsiloxane ratio, 95/5;
dimensions, 30 m by 0.25 mm). Helium was used as the gas carrier (1 ml/min). The column
temperature was maintained at 270°C, with the injector and detector set at 300°C. A linear gradient
from 150 to 180°C at 10°C/min was used for methyl esters, with MS conditions set at 280°C, 70
eV, and 2.2 kV.
629
Cytokine measurement
Supernatants from KCs infected with stationary-phase LD co-culture with T cells, isolated from the
mice infected with LD. For detection of IFN-γ from Apoe-/-, splenocytes isolated from LD infected
mice and cultured with SLA. IFN-γ levels in the supernatants were measured using a sandwich
ELISA Kit (R&D Systems) as per manufacturer's protocol. Detection limit for these kits were 8pg/ml
for IFN-γ. Cytokine levels were determined by measuring the OD at 450 nm using a MultiSkan FC
microplate photometer (Thermo Fisher Scientific). Data was demonstrated as mean ± SD of all five
individual experiments.
638
639 **Western Blot Analysis**
640 Lysates of LD-infected PECs and KCs were prepared, and western blotting was performed for
641 different proteins with endogenous control β-actin (Cell Signaling Technology). Blots were probed
642 with specific antibodies. Binding of secondary HRP-labeled goat anti-rabbit or goat anti-mouse Abs
643 was analyzed using SuperSignalR West Pico or West Dura Chemiluminescent substrate (Pierce).
644
Confocal and Structural Illumination Microscopy (SIM)
PECs and KCs were cultured on 13-mm diameter glass coverslips at a density of 2*10^5, and
infected with LD as mentioned previously. After varying time intervals, infected PECs or KCs were
rinsed with PBS, fixed with 2% paraformaldehyde, and permeabilized using 0.2% Triton/PBS for
20mins on an orbital shaker. Non-specific binding was blocked with 2% BSA in 0.2% Triton/PBS
for another 20mins on an orbital shaker. The PECs and KCs were then incubated with primary
antibodies for 1Hr, followed by washing and staining with IgG Alexa Fluor 594 or IgG Alexa Fluor
488, as required. Mounting was performed using Fluromount-G-DAPI (Thermo Scientific). Images
were captured using Olympus confocal microscopy (FV3000) and super-resolution microscopy
ZEISS (SIM) with a 63x objective. 3D image visualization processed through Fiji 3D viewer and 3D
project plugin. Live cell microscopy was performed using ZEISS phase contrast microscope and
ZEISS Apotome. MΦs were plated on confocal dish and incubated with photoRED-LDL for live cell
imaging of LDL uptake (**Movie2 and Movie 3)**.
658
659 **Lipid droplet staining**

LD-infected-KCs were first fixed using 2% paraformaldehyde (PFA) and subsequently permeabilized with 0.25% Triton X-100. Coverslips were then treated with Nile red (Sigma, diluted 1:100) in PBS for at least 1Hr to stain lipids, followed by DNA staining with DAPI. The coverslips were mounted onto glass slides using Fluoromount-G and visualized with confocal microscope (LEICA STELLARIS 5). 3D stacks were acquired to visualize distribution of lipid droplets. Images were processed and analysed, and stacks were merged to reconstruct 3D images using Fiji.

Ultrastructure expansion microscopy

Ultrastructure Expansion Microscopy (U-ExM) was conducted on LD-infected-MΦs following established experimental protocols [55]. Protein crosslinking was initiated by immersing the coverslip in a solution containing 1.4% formaldehyde, 2% acrylamide, and PBS for 5Hrs at 37°C. Monomer solution (19% sodium acrylate, 10% acrylamide, and 0.1% N,N'-methylenbisacrylamide in PBS) was mixed with 10% TEMED and 10% APS solution on ice and added to 6-well pates. Gelation was performed on ice, with cell-plated coverslip placed face down on the gelation solution in a humid chamber for 5 minutes at 37°C for 1 hour. The gel and coverslip were together transferred to a 6-well plate filled with denaturation buffer (200 mM SDS, 200 mM NaCl, and 50 mM Tris pH 9.0) and incubated face up for 15mins at room temperature under agitation, followed by incubation at 95°C for 1.5Hrs. The first expansion phase involved incubating the gel three times in ddH2O for 30mins each. Previously mentioned IFA steps were adopted for antibody staining. Images were captured using an Olympus confocal microscope (FV3000) equipped with a 63X 1.4 NA oil objective and Small Volume Computational Clearing mode was used to obtain deconvolved images. 3D stacks were acquired and analyzed using Fiji software.

Transmission electron microscopy (TEM)

For transmission electron microscopy, PECs infected with the LD-R strain were fixed after 24 hours of infection using 2.5% glutaraldehyde, an electron microscopy (EM) grade fixative. The cells were then processed according to previously established protocols[57]. Following this, the samples were examined using a JEM-F200 transmission electron microscope, allowing for lipid droplet imaging and detailed visualization of the cellular structures.

***In-vitro* cholesterol trafficking assay**

KCs was first treated with NBD-cholesterol for 16Hrs, allowing the fluorescent cholesterol analog to integrate into their membranes. Next, KCs were washed with RPMI 1640 and cultured for again 2Hrs in the same medium to eliminate any residual fluorescence from the background. These labelled KCs were infected with LD for 4Hrs and washed to remove loosely attached MΦs. The KCs were then either immediately fixed with 2% paraformaldehyde (PFA) or left untreated to serve as a control group. Finally, the fixed MΦs were examined under a confocal microscope.

Flow cytometry to determine NBD-cholesterol

KCs were first treated with NBD-cholesterol as previously described. LD parasites were isolated after 2Hrs of attachment to the KCs. The quenching of NBD-cholesterol was measured by flow cytometric analysis, using an unstained control.

Total internal reflection fluorescence (TIRF) microscopy

To visualize membrane fluidity, TIRF microscopy was performed [75]. Briefly, KCs were plated on coverslips, and after 24Hrs of LD-infection, KCs were fixed with 2% PFA. KCs were stained with

706   Laurdan dye and visualized with Olympus cell TIRF microscopic system. Representative images
707   of three independent biological experiments have been provided.
708

RNAseq analysis
RNA-Sequencing was performed utilizing external service by Bionivid Project for LD-S and LD-R-
infected-PECs, following 24Hrs of infection keeping uninfected MΦs as control (Bioproject. Average
expression of differentially expressed genes related to lipid metabolism between LD-S and LD-R-
infected-KC is represented as heat map. RAW sequencing reads have been deposited to Indian
Nucleotide Data Archive (https://inda.rcb.ac.in/home) with accession number INRP000146.
715

716 **Lipid profiling from serum**
717   Lipid profile analysis of patient serum was conducted using lipid estimation kits employing the
718   enzymatic method for the determination of cholesterol, triglycerides, HDL, and LDL. For BALB/c
719   mice, blood was collected following anesthesia and prior to euthanasia. The blood samples were
720   maintained at 4°C to allow clotting, after which the serum was isolated**.**
721

Evaluation of EC50
KCs infected with LD parasites were treated with SAG or Amp-B 24Hrs post-infection and incubated
for an additional 24Hrs. At the experimental endpoints, coverslips were washed with PBS, air-dried,
and fixed with 100% ice-cold methanol. Subsequently, the coverslips were stained with Giemsa
solution and examined microscopically. To quantify the number of amastigotes, one hundred KCs
per coverslip were counted. [2]. The average of three untreated coverslips was taken as 100%
control, and the percentage inhibition of infected KCs in treated cultures was calculated. EC50
values for each isolate were estimated against each drug [2]. Results were also expressed as
EC50.
731

Aspirin treatment and amastigote load determination
KCs were infected with LD parasites and, after 4Hrs, treated with 5µM Aspirin for a duration of 24
hours. Following this treatment, the KCs were washed and subsequently treated with Amp-B at a
specific dose for an additional 24 hours. Coverslips were then washed with PBS, air-dried, and
fixed with 100% ice-cold methanol. After fixation, the coverslips were stained with Giemsa solution
and examined microscopically. To quantify the number of amastigotes, one hundred KCs per
coverslip were counted. [2]
739

siRNA mediated Knock down (KD)
For all siRNA transfections, Lipofectamine 2000 was used according to the manufacturer's
instructions. PECs were seeded into 24-well plates at a density of $1x10^5$ per well, and incubated at
37°C with 5% CO_2. The transfection complex, comprising 1 µl Lipofectamine 2000 (Invitrogen) and
specific siRNAs, was directly added to the incubated PECs. Gene silencing was checked by IFA
as mentioned previously.
746

Statistical analysis
All statistical analyses were performed using GraphPad Prism on raw data. For datasets involving
single parameters or single data groups, one-way or two-way analysis of variance (ANOVA) was
utilized, followed by Tukey's test with 95% confidence interval. Wilcoxon rank sum analyses, with
95% confidence interval, were employed for comparing multiple groups. To check if the difference
between two groups is statistically significant, we used Student t-test. The data are represented as

mean ± SEM. Statistical significance was determined at P ≤ 0.05. Notation for significance levels:
*P ≤ 0.05; **P ≤ 0.001; ***P ≤ 0.0001.
755

Acknowledgments
SP acknowledges CSIR-UGC (2020-2021) and PMRF (2021-2024) fellowship. DD is a recipient of
GATE fellowship. DM is a recipient of CSIR fellowship, SC is a recipient of GATE fellowship.
Authors would like to acknowledge IIR lab, SMST, IITKGP for their help in kupffer cell isolation and
Flow cytometric analysis, Confocal Microscope facility under DST-FIST grant conferred on the
School of Bioscience, File no. SR/FST/LS-I/2019/595(C), and Confocal facility of Department of
Biotechnology, DST-FIST, Govt. of India. Authors would like to thank Prof. Syamal Roy, IICB, India
for providing all the LD strains used in this study and Prof. Yasuyuki Goto, The University of Tokyo,
Japan for allowing experiments related to live cell video microscopy in his lab. Authors
acknowledges CRF SMST, and CRF IITKGP. Authors would like to thanks Dr. Moumita Bhaumik,
NICED, India for helping in membrane isolation experiments. Authors would like to thank all the
interns of the IDI lab, SMST who help in various experiments related to this project.

Author Contributions: SP conceptualization, experiments, analysis, MS preparation. DD
performed Raman spectroscopy experiments. DM generated all the GFP, RFP expressing LD
strains, SC perform experiments related to EC50, AB performed Raman spectroscopy experiments
and analysis, KC did inhibitor study related to Cyt-D, Lat-A, RM serum collection from VL patients,
AS related serum lipid profile, formal analysis of the MS, KP related to serum lipid profile, formal
analysis of the MS, SD helped in formal analysis of Raman spectroscopic experiments. B.M
conceptualization, analysis, conceived and directed the project and prepared the MS with input
from all the authors.
777

778   **Competing Interest Statement:** The authors declare no competing interests.

779   **Data availability:** RAW sequencing reads have been deposited to Indian Nucleotide Data Archive
780   (https://inda.rcb.ac.in/home) with accession number INRP000146.

References

1.   Dumetz, F., et al., *Molecular preadaptation to antimony resistance in Leishmania donovani*
on the Indian subcontinent. MSphere, 2018. **3**(2): p. 10.1128/msphere. 00548-17.
2.   Mukhopadhyay, R., et al., *Characterisation of antimony-resistant Leishmania donovani*
isolates: biochemical and biophysical studies and interaction with host cells. International
journal for parasitology, 2011. **41**(13-14): p. 1311-1321.
3.   Mukherjee, B., et al., *Antimony-resistant but not antimony-sensitive Leishmania donovani*
up-regulates host IL-10 to overexpress multidrug-resistant protein 1. Proceedings of the
National Academy of Sciences, 2013. **110**(7): p. E575-E582.
4.   Verma, S., et al., *Quantification of parasite load in clinical samples of leishmaniasis*
patients: IL-10 level correlates with parasite load in visceral leishmaniasis. PloS one, 2010.
5(4): p. e10107.
5.   Thakur, C., D. Mitra, and S. Narayan, *Skewing of cytokine profiles towards T helper cell*
type 2 response in visceral leishmaniasis patients unresponsive to sodium antimony
gluconate. Transactions of the Royal Society of Tropical Medicine and Hygiene, 2003.
97(4): p. 409-412.
6.   Ouakad, M., et al., *Increased metacyclogenesis of antimony-resistant Leishmania*
donovani clinical lines. Parasitology, 2011. **138**(11): p. 1392-1399.
7.   Vanaerschot, M., et al., *Antimonial resistance in Leishmania donovani is associated with*
increased in vivo parasite burden. PloS one, 2011. **6**(8): p. e23120.
8.   Mukherjee, B., et al., *Probing the molecular mechanism of aggressive infection by*
antimony resistant Leishmania donovani. Cytokine, 2021. **145**: p. 155245.
9.   Feingold, K.R. and C. Grunfeld, *Lipids: a key player in the battle between the host and*
microorganisms1. Journal of lipid research, 2012. **53**(12): p. 2487-2489.
10.   Sonda, S. and A.B. Hehl, *Lipid biology of Apicomplexa: perspectives for new drug targets,*
particularly for Toxoplasma gondii. TRENDS in Parasitology, 2006. **22**(1): p. 41-47.
11.   Goad, L.J., G.G. Holz Jr, and D.H. Beach, *Sterols of Leishmania species, implications for*
biosynthesis. Molecular and biochemical parasitology, 1984. **10**(2): p. 161-170.
12.   Pucadyil, T.J., et al., *Cholesterol is required for Leishmania donovani infection: implications*
in leishmaniasis. Molecular and biochemical parasitology, 2004. **133**(2): p. 145-152.
13.   Sood, C., et al., *Leishmania highjack host lipid body for its proliferation in macrophages by*
overexpressing host Rab18 and TRAPPC9 by downregulating miR-1914-3p expression.
Plos Pathogens, 2024. **20**(2): p. e1012024.
14.   Bouazizi-Ben Messaoud, H., et al., *Changes in lipid and fatty acid composition during*
intramacrophagic transformation of Leishmania donovani complex promastigotes into
amastigotes. Lipids, 2017. **52**: p. 433-441.
15.   Lal, C.S., et al., *Hypocholesterolemia and increased triglyceride in pediatric visceral*
leishmaniasis. 2007.
16.   Pradhan, S., et al., *Linking membrane fluidity with defective antigen presentation in*
leishmaniasis. Parasite immunology, 2021. **43**(7): p. e12835.
17.   Douanne, N., et al., *Leishmania parasites exchange drug-resistance genes through*
extracellular vesicles. Cell reports, 2022. **40**(3).
18.   Légaré, D., et al., *ABC proteins of Leishmania.* Journal of bioenergetics and
biomembranes, 2001. **33**: p. 469-474.
19.   Gourbal, B., et al., *Drug uptake and modulation of drug resistance in Leishmania by an*
aquaglyceroporin. Journal of Biological Chemistry, 2004. **279**(30): p. 31010-31017.
20.   Ghosh, J., et al., *Hyperlipidemia offers protection against Leishmania donovani infection:*
role of membrane cholesterol [S]. Journal of Lipid Research, 2012. **53**(12): p. 2560-2572.
21.   Sen, S., et al., *Restoration of IFNγR subunit assembly, IFNγ signaling and parasite*
clearance in Leishmania donovani infected macrophages: role of membrane cholesterol.
PLoS pathogens, 2011. **7**(9): p. e1002229.
22.   Toledo, A. and J.L. Benach, *Hijacking and use of host lipids by intracellular pathogens.*
Virulence mechanisms of bacterial pathogens, 2016: p. 635-666.

23. Chang, W., et al., *Trapping toxins within lipid droplets is a resistance mechanism in fungi.*
Scientific reports, 2015. **5**(1): p. 15133.

24. Cao, Y., *Adipocyte and lipid metabolism in cancer drug resistance.* The Journal of clinical
investigation, 2019. **129**(8): p. 3006-3017.

25. Day, N.J., P. Santucci, and M.G. Gutierrez, *Host cell environments and antibiotic efficacy*
in tuberculosis. Trends in Microbiology, 2023.

26. Sheng, W., G. Ji, and L. Zhang, *Role of macrophage scavenger receptor MSR1 in the*
progression of non-alcoholic steatohepatitis. Frontiers in Immunology, 2022. **13**: p.
1050984.

27. Purkait, B., et al., *Mechanism of amphotericin B resistance in clinical isolates of Leishmania*
donovani. Antimicrobial agents and chemotherapy, 2012. **56**(2): p. 1031-1041.

28. Srivastava, P., et al., *An unusual case of Amphotericin B refractoriness in Visceral*
Leishmaniasis from a non-endemic region of India. Journal of clinical microbiology, 2011.
49(8): p. 3088.

29. Ghosh, J., et al., *Human visceral leishmaniasis: decrease in serum cholesterol as a*
function of splenic parasite load. Annals of Tropical Medicine & Parasitology, 2011. **105**(3):
p. 267-271.

30. Haldar, A.K., et al., *Leishmania donovani isolates with antimony-resistant but not-sensitive*
phenotype inhibit sodium antimony gluconate-induced dendritic cell activation. PLoS
Pathogens, 2010. **6**(5): p. e1000907.

31. Beattie, L., et al., *Dynamic imaging of experimental Leishmania donovani-induced hepatic*
granulomas detects Kupffer cell-restricted antigen presentation to antigen-specific CD8+ T
cells. PLoS pathogens, 2010. **6**(3): p. e1000805.

32. Hinder, L.M., et al., *Apolipoprotein E knockout as the basis for mouse models of*
dyslipidemia-induced neuropathy. Experimental neurology, 2013. **239**: p. 102-110.

33. Semini, G., et al., *Changes to cholesterol trafficking in macrophages by Leishmania*
parasites infection. Microbiologyopen, 2017. **6**(4): p. e00469.

34. Vance, J.E., *Cellular itinerary of LDL cholesterol.* Proceedings of the National Academy of
Sciences, 2022. **119**(6): p. e2122584119.

35. Pessoa, C.C., et al., *ATP6V0d2 controls Leishmania parasitophorous vacuole biogenesis*
via cholesterol homeostasis. PLoS pathogens, 2019. **15**(6): p. e1007834.

36. Feuerer, N., et al., *Lipidome profiling with Raman microspectroscopy identifies*
macrophage response to surface topographies of implant materials. Proceedings of the
National Academy of Sciences, 2021. **118**(52): p. e2113694118.

37. Živanović, V., et al., *Chemical mapping of leishmania infection in live cells by SERS*
microscopy. Analytical chemistry, 2018. **90**(13): p. 8154-8161.

38. Tiwari, D., et al., *Temporal in vitro Raman spectroscopy for monitoring replication kinetics*
of Epstein–Barr virus infection in glial cells. ACS omega, 2020. **5**(45): p. 29547-29560.

39. Beattie, J.R., S.E. Bell, and B.W. Moss, *A critical evaluation of Raman spectroscopy for*
the analysis of lipids: Fatty acid methyl esters. Lipids, 2004. **39**(5): p. 407-419.

40. Goldstein, J.L., et al., *Receptor-mediated endocytosis: concepts emerging from the LDL*
receptor system. Annu Rev Cell Biol, 1985. **1**(1): p. 1-39.

41. Lucero, D., et al., *Interleukin 10 promotes macrophage uptake of HDL and LDL by*
stimulating fluid-phase endocytosis. Biochimica et Biophysica Acta (BBA)-Molecular and
Cell Biology of Lipids, 2020. **1865**(2): p. 158537.

42. Popow-Woźniak, A., et al., *Cofilin overexpression affects actin cytoskeleton organization*
and migration of human colon adenocarcinoma cells. Histochemistry and Cell Biology,
2012. **138**: p. 725-736.

43. Urano, Y., et al., *Transport of LDL-derived cholesterol from the NPC1 compartment to the*
ER involves the trans-Golgi network and the SNARE protein complex. Proceedings of the
National Academy of Sciences, 2008. **105**(43): p. 16513-16518.

44. Verma, J.K., R. Rastogi, and A. Mukhopadhyay, *Leishmania donovani resides in modified*
early endosomes by upregulating Rab5a expression via the downregulation of miR-494.
PLoS pathogens, 2017. **13**(6): p. e1006459.

45.   Garver, W.S., et al., *The Niemann-Pick C1 gene is downregulated by feedback inhibition*
899   *of the SREBP pathway in human fibroblasts.* Journal of lipid research, 2008. **49**(5): p. 1090-
900   1102.
901   46.   Ho, C.-M., et al., *Accumulation of free cholesterol and oxidized low-density lipoprotein is*
902   *associated with portal inflammation and fibrosis in nonalcoholic fatty liver disease.* Journal
903   of Inflammation, 2019. **16**: p. 1-8.
904   47.   Murray, H., H. Masur, and J. Keithly, *Cell-mediated immune response in experimental*
905   *visceral leishmaniasis. I. Correlation between resistance to Leishmania donovani and*
906   *lymphokine-generating capacity.* 1982.
907   48.   Kima, P.E. and L. Soong, *Interferon gamma in leishmaniasis.* Frontiers in immunology,
908   2013. **4**: p. 156.
909   49.   Govaere, O., et al., *Macrophage scavenger receptor 1 mediates lipid-induced inflammation*
910   *in non-alcoholic fatty liver disease.* Journal of hepatology, 2022. **76**(5): p. 1001-1012.
911   50.   Lu, M. and O. Gursky, *Aggregation and fusion of low-density lipoproteins in vivo and in*
912   *vitro.* Biomolecular concepts, 2013. **4**(5): p. 501-518.
913   51.   Yano, H., Y. Fujiwara, and Y. Komohara, *Cholesterol metabolism and lipid droplet*
914   *vacuoles; a potential target for the therapy of aggressive lymphoma.* Journal of clinical and
915   experimental hematopathology, 2022. **62**(4): p. 190-194.
916   52.   Banerjee, S., et al., *Leishmania donovani infection induce Extracellular signal-regulated*
917   *kinase ½ (ERK½) mediated lipid droplet generation in macrophages.* Molecular
918   Immunology, 2022. **141**: p. 328-337.
919   53.   Meester, I., et al., *The roles of lipid droplets in human infectious disease.* Revista
920   Portuguesa de Cardiologia, 2011. **13**(53): p. 207-216.
921   54.   Bosch, M., et al., *Lipid droplets and the host–pathogen dynamic: FATal attraction?* Journal
922   of Cell Biology, 2021. **220**(8): p. e202104005.
923   55.   Liffner, B., et al., *Atlas of Plasmodium falciparum intraerythrocytic development using*
924   *expansion microscopy.* Elife, 2023. **12**: p. RP88088.
925   56.   Ghosh, S., et al., *A novel bioimpedance-based detection of miltefosine susceptibility*
926   *among clinical Leishmania donovani isolates of the Indian subcontinent exhibiting*
927   *resistance to multiple drugs.* Frontiers in Cellular and Infection Microbiology, 2021. **11**: p.
928   768830.
929   57.   D'Avila, H., et al., *Host cell lipid bodies triggered by Trypanosoma cruzi infection and*
930   *enhanced by the uptake of apoptotic cells are associated with prostaglandin E2 generation*
931   *and increased parasite growth.* Journal of Infectious Diseases, 2011. **204**(6): p. 951-961.
932   58.   Thakur, A., H. Mikkelsen, and G. Jungersen, *Intracellular pathogens: host immunity and*
933   *microbial persistence strategies.* Journal of immunology research, 2019. **2019**(1): p.
934   1356540.
935   59.   Kreimendahl, S. and L. Pernas, *Metabolic immunity against microbes.* Trends in Cell
936   Biology, 2023.
937   60.   Díaz-Varela, M., et al., *The different impact of drug-resistant Leishmania on the*
938   *transcription programs activated in neutrophils.* iScience, 2024. **27**(5): p. 109773.
939   61.   Boodhoo, N., N. Kamble, and S. Behboudi, *De novo cholesterol biosynthesis and its*
940   *trafficking in LAMP-1-positive vesicles are involved in replication and spread of Marek's*
941   *disease virus.* Journal of Virology, 2020. **94**(24): p. 10.1128/jvi. 01001-20.
942   62.   Prakash, S. and A.K. Rai, *Retinoic acid increases cellular cholesterol in Leishmania*
943   *donovani-infected macrophages in an mTOR-independent manner.* Microbiology
944   Spectrum, 2022. **10**(6): p. e02699-22.
945   63.   Kim, H. and S.J. Shin, *Revolutionizing control strategies against Mycobacterium*
946   *tuberculosis infection through selected targeting of lipid metabolism.* Cellular and Molecular
947   Life Sciences, 2023. **80**(10): p. 291.
948   64.   Awadh, A.A., *The Role of Cytosolic Lipid Droplets in Hepatitis C Virus Replication,*
949   *Assembly, and Release.* Biomed Res Int, 2023. **2023**: p. 5156601.
950   65.   2024, L.i.f.b.a.W.T.C.A.Z.Z., *LeishGEM is funded by a Wellcome Trust Collaborative*
951   *Award*, in *LeishGEM*. 2024.

66. Downing, T., et al., *Whole genome sequencing of multiple Leishmania donovani clinical*
isolates provides insights into population structure and mechanisms of drug resistance.
Genome Res, 2011. **21**(12): p. 2143-56.
67. Patole, S., S. Burza, and G.M. Varghese, *Multiple relapses of visceral leishmaniasis in a*
patient with HIV in India: a treatment challenge. International Journal of Infectious
Diseases, 2014. **25**: p. 204-206.
68. Ray, A. and B.N. Dittel, *Isolation of mouse peritoneal cavity cells.* JoVE (Journal of
Visualized Experiments), 2010(35): p. e1488.
69. Khanra, S., et al., *In vivo experiments demonstrate the potent antileishmanial efficacy of*
repurposed suramin in visceral leishmaniasis. PLoS Neglected Tropical Diseases, 2020.
14(8): p. e0008575.
70. Li, P.-z., et al., *An efficient method to isolate and culture mouse Kupffer cells.* Immunology
letters, 2014. **158**(1-2): p. 52-56.
71. Mukherjee, S., et al., *Short-course treatment with imipramine entrapped in squalene*
liposomes results in sterile cure of experimental visceral leishmaniasis induced by
antimony resistant Leishmania donovani with increased efficacy. Frontiers in Cellular and
Infection Microbiology, 2020. **10**: p. 595415.
72. Amundson, D.M. and M. Zhou, *Fluorometric method for the enzymatic determination of*
cholesterol. Journal of biochemical and biophysical methods, 1999. **38**(1): p. 43-52.
73. Real, F., *Isolation of intact Leishmania amazonensis large parasitophorous vacuoles from*
infected macrophages by density gradient fractionation. Experimental Parasitology, 2020.
218: p. 107989.
74. Alpizar-Sosa, E.A., et al., *Amphotericin B resistance in Leishmania mexicana: Alterations*
to sterol metabolism and oxidative stress response. PLoS Neglected Tropical Diseases,
2022. **16**(9): p. e0010779.
75. Schirripa Spagnolo, C. and S. Luin, *Setting up multicolour TIRF microscopy down to the*
single molecule level. Biomolecular Concepts, 2023. **14**(1): p. 20220032.

     Figure and Legends

Figure 1

994
Figure 1. Severe dyslipidemia is linked with higher intracellular proliferation of LD-R
amastigotes. (A) Serum lipid profile of 28 days LD-infected Balb/c mice (N=5). Equal number of

sorted metacyclics promastigotes of two representative LD-S (LD-S[1], LD-S[2]) and LD-R (LD-R[1], LD-
R[2]) strains were used to perform independent infection. **(B)** Number of attached LDs on KC surface
4Hrs p.i, data represents (N=3) for two LD-S and two LD-R strains. **C (i)** Representative Giemsa
stain images of LD-S[1] and LD-R[1]-infected KC at 4Hrs and 24Hrs p.i Black arrow represents LD
amastigotes. N denotes host nucleus. Scale bar 20µM. Giemsa Images are represented in grey
scale to clearly represent LD nucleus (black arrow). **C (ii)** Intracellular amastigote count for 4Hrs
and 24Hrs p.i. Each dot represent count from 100 infected-KC (N=6). **D (i)** LD-infected-KC imaged
to visualize the PV at 4Hrs and 24Hrs p.i. PV marked by ATP6V0D2 (magenta), KC and LD nucleus
(blue), Scale bar 5µm. Yellow arrow represents LD amastigotes. In the merged image, white dotted
line marks KC's periphery, and host nucleus is represented with N. **D (ii)** PV counts measured from
30 infected KCs. **D (iii)** 3D morphometric representation for better visualization of PV in 24Hrs post
infected KC marked with white dotted box as in D (i). ATP6V0D2 (magenta), nucleus (cyan) with
last panel showing PV-grooves marked in red solid line with white arrow representing LD
amastigotes. *** signifies p value < 0.0001, * signifies p value < 0.05, n.s non-significant.

Figure 2

1011

Figure 2. LD-R-infection results in a higher organ parasite load in hypercholesterolemic
***Apoe*[-/-]mice. A (i)** Representative Giemsa images of LD-infected-KCs cultured in high and low
lipid media. Black arrow represents LD amastigotes; N represents host nucleus. Scale bar 20µM.

A (ii) Graph representing number of intracellular LD amastigotes in KCs with LD infection performed either in high, low or lipid free media (N=6). **B (i)** Giemsa images of LD-infected-KCs isolated from wild type BL/6 and *Apoe*$^{-/-}$ mice. Scale bar 20µm, N representing host nucleus. Giemsa Images are represented in grey scale to clearly represent LD nucleus (black arrow). **B (ii)** Amastigote count from 100 LD-infected-KCs (N=6) as in B(i). Data are presented as mean ± SEM. **C (i)** Schematic representation of *in-vivo* LD-infection in BL/6 and *Apoe*$^{-/-}$ mice performed with EGFP-LD-S or EGFP-LD-R **C (ii)** Flow cytometry representing GFP-positive splenocytes isolated from 11 days EGFP-LD infected BL/6 and *Apoe*$^{-/-}$ mice. The black box indicates % of GFP-positive splenocytes (LD-infected). **C** (**iii)** Graph representing data from six infected mice as in C (ii) are presented. *** signifies p value < 0.0001, ** signifies p value <0.001, * signifies p value < 0.05, n.s non-significant.

Figure 3

Figure 3. Endocytosed-LDL is the primary lipid source for intracellular LD-R proliferation. **A (i)** Live cell images of LDL-endocytosis in 24Hrs post infected KCs with LDL (red), LD (green), and the nucleus (blue). Images were acquired directly from 96 well glass bottom plate and bright field was merged with reduced brightness and increased contrast identically for all experimental conditions. Scale bar 40µm, Yellow arrow represents intracellular LD. **A (ii)** Graph representing Fluorescence plate reader-based quantification of A (i). **B (i)** Co-localization of LDL with the LD-amastigotes in infected KCs 24Hrs p.i. Merged image showing LDL (magenta), LD (green), nucleus (blue), White dotted line in the merged image demarcates infected-KC for co-localization analysis.

Scale bar 10µm, yellow arrow represents LD nucleus and N represents host nucleus **B (ii)** Intensity plot representing co-localized signals from B (i), marked with grey transparent box. **C (i)** GC-MS lipid profile of isolated PVs from LD-infected-KCs. Retention time 27.73 corresponding to the cholesterol and 28.56 corresponding to fatty acids are represented with * for LD-S-PV and with black arrowheads for LD-R-PV. **C (ii)** Relative abundance of Cholesterol and Fatty acids of C (i). **D (i)** Representative Confocal Raman spectroscopy image of uninfected or LD-infected-KCs 24Hrs p.i, illustrating distinct lipid-related signal distribution patterns marked with pseudo-colours. Yellow dotted circle in the left most panel demarcates the host cell nucleus while white arrow marking intracellular LD nucleus. In merged panel white dotted line demarcates cells periphery. The right most panel shows dot plot representation of lipid distribution, where each colour corresponds to different lipid-related peaks with respect to blue (B) indicating the nucleus. **D (ii)** Comparative Raman spectra from LD-S and LD-R-infected-KCs. Lipid-related peaks are demarcated with shaded box, representing 1. 540-560 cm-1 (Cholesterol), 2. 1080-1090 cm-1 (phospholipids), 3. 1270-1280 cm-1 (triglycerides), 4. 1300-1340 cm-1 (Amide-III bond), 5. 1440-1453 cm-1 (Fatty acids and Triglycerides), and 6. 1650-1660 cm-1 (Amide-I bond). *** signifies p value < 0.0001, ** signifies p value <0.001, * signifies p value < 0.05, n.s non-significant.

Figure 4

Figure 4. Increased LDL uptake in LD-R-Infected-MΦs through receptor independent fluid phase endocytosis. (A) Immunofluorescence images of LDLr expression in LD-infected-KCs.

Scale bar 40µm. Enlarged view of A showing infected-KC with low LDLr expression (yellow arrow)
compared to neighboring uninfected KC (white arrow) in the merged image. **B (i)** Cofilin and
phosphorylated-Cofilin expression by western blot in LD-infected-KCs. **B (ii)** Graphical
representation of comparative expression of Cofilin and phosphorylated-Cofilin with respect to β-
actin for B (i). **C (i)** SIM images showing the Cofilin distribution in LD-infected-KCs. Yellow arrow
represents intracellular LD, white dotted line in the merged image demarcates cell boundary. Scale
bar 5µm. **C (ii)** Colour spectrum representing Cofilin distribution pattern of C (i) with green
representing maximum and red representing minimum distribution. **D (i)** Confocal images
representing Filamentous actin (F-actin) protrusions in LD-infected-KCs. Scale bar 20µm, with
yellow arrow representing LD amastigotes**.** Individual experimental condition is represented in
pseudo-colour, with uniform high contrast to clearly represent the F-actin protrusions. **D (ii)** Graph
representing Cofilin and F-actin distribution in KCs infected with LD or left uninfected. (**E)** Still
images from live cell video microscopy representing co-localization of LD-R with LDL in presence
or absence of Latrunculin-A (LAT-A). Scale bar 60µm. Yellow arrow represents LD-infected-KCs
with LDL(red). **(F)** Graphical representation (N=6) of amastigote count in 100 LD-R-infected-KCs
treated with either LAT-A, or Cytochalasin (CYT-D) or left untreated. Data are presented as mean
± SEM. ** signifies p value <0.001, * signifies p value < 0.05, n.s non-significant.

1074

Figure 5

1075

Figure 5. Endocytosed-LDL fuses with LD-R-PV to provide Cholesterol to LD-R amastigotes.
(A) Confocal images of representing LDL-vesicles, with LAMP1 positive lysosomal vesicles in LD-infected-KCs 4Hrs p.i. Yellow arrow represents LD nucleus, white dotted box marks the region further cropped and zoomed to show fusion of lysosomal vesicles with LDL and LD-R. Scale bar 10μm. **A (ii)** Schematic representation showing trafficking of LDL-vesicle towards the LD-R-PV. **B (i)** SIM images representing convergence of LDL-vesicles with LD amastigotes and LAMP1 in 24Hrs with LD-infected-KCs. White dotted box marks the region further cropped and Zoomed to clearly show the convergence (of LD amastigotes with LDL and LAMP-1 (white vesicles shown by yellow arrow), Scale bar 5μm. **B (ii)** Intensity plot representing of co-localization signals for B (i). **C** Western blot analysis showing LAMP1 expression in LD-infected-KCs from early to late hours. **D (i)** Filipin staining of EGFP-LD-infected-KCs. Filipin (white) and LD (green), yellow arrow represents intracellular LD. Scale bar 10μm. White dotted line marked cells used to generate co-localization

1088   plot. **D (ii)** Co-localization signals of LD (EGFP and Filipin represented by the dot plot. Pearson co-
1089   localization coefficient value (r) represented for LD-S and LD-R-infected-KC. **D (iii)** Graph
1090   representing Pearson's coefficient for EGFP-LD-infected-KCs (N=19). Statistical significance in the
1091   observed differences was determined through ANOVA utilizing GraphPad Prism software (version
1092   9.0). *** signifies p value < 0.0001.

1093

Figure 6

1094

Figure 6. LD-R infection suppress NPC-1 to alter Cholesterol mobilization in infected MΦs.
(A) 24Hrs LD-S and LD-R-infected-PECs were subjected to *RNAseq* analysis keeping uninfected-PECs as control. Heat map representing average expression of differentially expressed genes

related to lipid metabolism between LD-S and LD-R-infected-PEC. Scale represents median-
centered counts (TPM). Row Z score data represented here. Expression of RNAs (right margins)
is presented as centered and "scaled" [mean normalized $\log_2$(TPM + 1)]. **B (i)** Expression of
SREBP2 and **B (ii)** NPC-1 in LD-infected and uninfected KCs by Western blot with β-actin as
control. (**C**) Intracellular localization of SREBP2 in response to LD-infection in KC 24Hrs p.i. White
dotted line represents cell periphery with yellow arrow representing intracellular LD and N
representing host nucleus. 3D image represented in the right most panel with one quadrant clipped
to confirm nuclear translocation of SREBP2. Scale bar 5µm. (**D**) Immunofluorescence showing
NPC-1 expression in LD-infected-KCs. White arrow representing NPC-1 expression in LD-R-
infected-KCs and yellow arrow representing NPC-1 in adjacent uninfected-KCs in the merged
panel. White dotted box marks regions further cropped and shown in the bottom panel with
enlarged view (Zoomed). Scale bar 20µm. **E (i)** TIRF microscopic images of Laurdan stained LD-
infected and uninfected KCs with two different spectra (488nm, 594nm), changes from red to green
represents gel to fluid phase transition of the host plasma membrane. Colour distribution pattern
corresponding to the TIRF images representing each experimental condition is presented in the
right most panel. Scale bar 5µm. **F (i)** Expression of NPC-1 determined by immunofluorescence in
PECs transfected with NPC-1 siRNA or scrambled (sc) siRNA. Scale bar 5µm. **F (ii)** Giemsa-
stained images of LD-infected NPC-1 knockdown-PECs with scrambled siRNA control (Sc). Scale
bar 20µm. Images are represented in grey scale with increased contrast to represent LD nucleus
(black arrow). **F (iii)** Graphical representation of number of intracellular LD-amastigotes as in F (ii).
Results show the counting of 100 infected-PEC from 6 independent replicate. * signifies p value <
0.05, n.s non-significant.

Figure 7

1120

Figure 7. LD-R-infected-MΦs selectively excludes ox-LDL to suppress inflammatory response in host. (A) Scheme representing ex vivo co-culture experiment performed by isolating T-cells from LD-infected mice with in vitro LD-infected-KCs **B (i)** IFN-γ level in the supernatant of LD-infected-KCs co-cultured with the T-cells isolated from spleen of Balb/c mice infected with LD-S or LD-R 28 days p.i. Data represented as mean ± SD from three independent experiments. **B (ii)** IFN-γ level in the supernatant of splenocytes cultures from LD-infected-BL/6 and *Apoe*[-/-]-mice stimulated with SLA (Soluble *Leishmania* antigen) specific for LD-S or LD-R. Data are presented

as mean ± SD from six independent mice. **C (i)** ox-LDL incubated LD-infected-KCs was fixed and
stained with Oil-red-O and Haematoxylin. LD-S^{High} represents MOI (1:20) and LD-R^{Low} represents
MOI (1:5). Scale bar 10μm. Black arrow represents LD nucleus. **C (ii)** Graph representing % of ox-
LDL positive MΦs under different experimental conditions as represented in C (i). **D (i)** Confocal
image showing MSR-1 in LD-infected-KCs. Yellow arrow represents LD-amastigotes while N
represents host nucleus. Scale bar 20μm **C (ii)** Expression of MSR1 detected by Western blot of
LD-infected-KC with β-actin as loading control. (**D)** Expression of ATP6V0D2 in LD-infected-KCs
by Western blot with β-actin as loading control. **E (i)** Expression of MSR1 and neutral lipid droplet
accumulation assayed by Confocal microscopy from the cryo-sectioned liver of 28 days LD-
infected-mice. Representative image showing Lipid droplet marked in red and MSR1 in green,
nucleus in blue. Scale bar 40μm. **E (ii)** Graphical representation of % positive Lipid droplets
accumulation and MSR-1 expression under different experimental condition as in E (i). Data
collected from N=6 mice. *** signifies p value < 0.0001, ** signifies p value <0.001, * signifies p
value < 0.05.

Figure 8

1142

Figure 8. Lipid droplets accumulation around LD amastigotes is inversely related with susceptibility towards Amphotericin-B. (A) Ex-M image showing lipid droplet assimilation in LD-

R-amastigotes. White dotted region further cropped and shown in enlarged panel (Zoomed). Nile
red (magenta), LD (green), nucleus (cyan). White spots represent Lipid droplets assimilated in LD-
R and Host nucleus is represented as N. Scale bar 5μm. **(B)** TEM images showing lipid droplets
around the LD-R amastigotes. Black box further magnified and represented in right panel. White *
representing LD-R amastigotes, yellow * showing lipid droplets. Scale bar 2μm. (**C**) Schematic
representation representing experimental scheme used to determine EC50 against Amphotericin-
B under different experimental conditions**. D (i) and (ii)** Determination of EC50 value against
Amphotericin-B in low lipid and high lipid media by performing infection with of two independent
LD-R strains. **D (iii)** Determination of EC50 value against Amphotericin-B in KC infected with LD-
$S^{Amp-B-R}$. Intracellular amastigotes were counted through Giemsa staining. **(E)** Confocal images
showing lipid droplets accumulation around intracellular amastigotes with Aspirin (5 μM) or Amp-B
(0.36 μM) or with a combination of Aspirin (5μM) and Amp-B (0.11μM)at 48Hrs p.i   **(F)**
Determination of amastigote load in KCs infected with two independent LD-R isolates in different
experimental conditions either untreated or treated with Aspirin (5 μM) or Amp-B (0.36 μM) or with
a combination of Aspirin (5μM) and Amp-B (0.11μM) at 48Hrs p.i (**G)** Comparative Lipid profile of
VL patients (Cured and Relapsed) in response to Amp-B treatment along with healthy individuals
from endemic region of Bihar, India. *** signifies p value < 0.0001, ** signifies p value <0.001, *
signifies p value < 0.05.

1163

1164 **Graphical Abstract**

1165

Figure 9: Scrutinized uptake of host lipids results in breakdown of Amphotericin-B
responsiveness in clinical LD isolates with primary resistance to antimony. (1-2) LD-R
residing in membrane bound parasitophorous vacuoles (PVs), and (3-5) induce high fluid phase
endocytosis of LDL by modulating actin depolymerization of host membrane through cofilin. (6-7)
LDL containing vesicles fused with LD-R-PV which provides increased lipid access to sustain their
aggressive intracellular proliferation. Vesicular cholesterol export protein, NPC-1 is downregulated
in the fused membrane, preventing release of LDL derived cholesterol back to host cytoplasm. (8-
12) LDL derived cholesterol esters also recycled in the ER and form lipid droplets which ultimately
accumulate in and around the LD-R-PV ultimately leading to the Amphotericin-B
unresponsiveness. (13-15) LD-R control inflammatory response related to accumulation of oxidized
lipids (Ox-LDL) by downregulating MSR-1 (16) High LDL influx also results in shutdown of *de-novo*
cholesterol synthesis in LD-R infected host through restriction of nuclear translocation of SERBP2.

Right panel, representing LD-S infected host with low LDL influx with high NPC-1 expression
leading to compromised lipid droplet accumulation around its PV, thereby exhibiting high
responsiveness against Amphotericin-B.

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

 Supplementary Information

Figure S1

1207

1208 **Figure S1. Kupffer cells (KC) as an *in vitro* model for LD-infection. (A)** Flow Cytometry
1209 representing % Metacyclics LD: **(i)** LD-S (AG83, DD8) and **(ii)** LD-R (BHU814, BHU575) based on
1210 FSC and SSC. Left, small box with low FSC and SSC gate for metacyclic parasites and right panel

 with high FSC and SSC gated procyclic parasites. **(B)** Flow cytometry of F4/80-PE positive KC from
the non-parenchymal cell population isolate from murine liver. **C** Representative Giemsa-stained
images LD-infected-PEC, 4 and 24Hrs p.i for **(i)** LD-S and **(ii)** LD-R strains. Giemsa Images are
represented in grey scale to clearly represent LD nucleus (black arrow). Scale Bar 20µM **C (iii)**
Intracellular amastigote count for 4Hrs and 24Hrs p.i. Each dot represent count from 100 infected-
KC. N=6. **D** Amastigote count from Giemsa-stained image of KC-infected with LD amastigotes,
isolated from the spleen of 28 days LD-infected-mice. Each dot represent mean from 100 infected-
KCs. N=6. *** signifies p value < 0.0001, ** signifies p value <0.001, n.s non-significant.

Figure S2

1219

Figure S2. LD-S and LD-R quenches host membrane cholesterol during initial entry. A (i)
KCs were pre-incubated with NBD-cholesterol for 16Hrs. Subsequently, KCs were infected with
RFP expressing LD-S or LD-R for 4Hrs or left uninfected and imaged. NBD-cholesterol was excited
at 488nm, and LDs are presented with magenta pseudo color. White arrow indicates the NBD-
cholesterol within LD while yellow arrow represents LD nucleus. Scale bar 5µm. **A (ii)** Co-
localization analysis of LD (magenta), and NBD-Cholesterol (green) from A (i), marked in grey
transparent box. **B (i)** Representative Flow cytometry-based analysis of NBD in LD-amastigotes
isolated from LD-infected-KCs 2Hrs after infection pre-incubated with NBD cholesterol. Black Box
representing % NBD positive population. **B (ii)** Graphical representation of MFI for three
independent experiments as in B (i). n.s non-significant.

1230

Figure S3

1231

Figure S3. LD-R-PV have higher accumulation of Cholesterol and Fatty acids. (A) Detection
of PV Cholesterol, isolated from LD-S or LD-R-infected KC. Data represented as μg of protein/ μg
of Cholesterol, (N=6).   **(B)** TLC profile of isolated lipids from 24Hrs LD-infected-PVs with
Cholesterol standard kept as control. (**C**) Individual Raman spectra from six independent LD-
infected-KCs **(i)** LD-S-infected and **(ii)** LD-R-infected. Lipid-related peaks are demarcated within
the spectra, including those at 1.540-560 cm-1 (Cholesterol), 2. 1080-1090 cm-1 (phospholipids),
3. 1270-1280 cm-1 (triglycerides), 4. 1300-1340 cm-1 (Amide-(II) I bond), 5. 1440-1453 cm-1 (Fatty
acids and Triglycerides), and 6. 1650-1660 cm-1 (Amide-I bond). **(D)** Area under the curve, and
shift are calculated and plotted for each experimental condition as represented in C (i) and (i). **
signifies p value <0.001,

Figure S4. Latrunculin-A restricts extracellular LDL influx inhibiting LD-R amastigote proliferation. (A) Western blot comparing of LDLr expression between uninfected, LD-S, and LD-R-infected-KC 24Hrs p.i. **B (i)** Expression of LDLr determined by immunofluorescence in PECs treated with LDLr siRNA and scrambled siRNA (Sc). Scale bar 10µm. **B (ii)** Graph representing number of LD-amastigotes in LDLr knockdown per 100 PECs for B (i) (N = 6). Data: mean ± SD. **C (i)** Cofilin and phosphorylated-Cofilin expression by Western blot in LD-infected-PEC. **C (ii)** Graphical representation of comparative expression of abundance of Cofilin and phosphorylated-Cofilin with respect to β-actin for C(i). **(D)** Giemsa stained images of Latrunculin-A (LAT-A) and Cytochalasin-D (CYT-D) treated LD-infected-KCs in grey scale. LD nucleus are represented with Black arrow and host nucleus are represented as N. Scale bar 20µm. n.s non-significant.

1255

Figure S5. Aspirin treatment inhibits Lipid droplets formation and increase responsiveness of LD-R amastigotes towards Amphotericin-B. (A) Determination of lipid droplet accumulation by Nile red staining in LD-S and LD-R-infected-KC or KC left uninfected. Dotted line mark the cell periphery while dotted box cells were used for 3D reconstruction and represent in Movie 3. Scale Bar 5 μm. (**B)** Graphical representation of Lipid Droplet positive KCs in high lipid and low lipid media. Data represented as % of lipid droplet positive KCs (N=6) collected from 3 independent experiments. (**C)** Effect of Aspirin(5μM) on Lipid droplets accumulation in LD-R-infected-KCs. Independent infection was performed with LD-R[1] and LD-R[2] strains (N=6) and % KCs with high lipid droplets accumulation are represented. (**D)** Representative Giemsa stained images of LD-R-infected-KCs in different experimental conditions either untreated or treated with Aspirin (5μM) or Amp-B (0.36 μM) or with a combination of Aspirin (5μM) and Amp-B (0.11 μM) at 48Hrs p.i. Scale bar 20μm. *** signifies p value < 0.0001

1268

1269

1270

 Table-1: EC50 against SAG (Antimonial) for the LD strains used in this study

Strain Name	EC50 value for SAG(µg/ml) in PEC	EC50 value for SAG(µg/ml) in KC
MHOM/IN/2009/BHU575/0	22.73±3.7	23.89±2.1
MHOM/IN/10/BHU814/1	30.3±7.2	28.5±5.3
MHOM/IN/83/AG83	1.52±1.2	3.3±2.2
MHOM/IN/80/DD8	2.1±1.2	3.2±1.9
DD8-AmpB	1.8±2.1	2.2±1.3
BHU575-EGFP	21.45±3.2	22.81±3.1
AG83-EGFP	2.1±1.4	2.4±3.1

1272

1273   **Table-2: Lipid profile of VL patients**

Serum Code	Age	Sex	Time of Blood Collection	Infection status	Treatment received	Patient status	LDL (mg/dl)	HDL (mg/dl)	Cholesterol (mg)	Triglycerides (mg/dl)
RMRI N1	45	F	Before	Control population from endemi	NA	NA	45.3501585	46.6365931	101.6949153	177.258567
RMRI N6	70	F	Before	Control population from endemi	NA	NA	110.855478	50.3762712	111.8644068	206.5420561
RMRI N7	45	F	Before	Control population from endemi	NA	NA	75.5620787	41.8711864	84.40677966	165.4205607
RMRI N8	19	M	Before	Control population from endemi	NA	NA	83.946038	62.9533123	100.3389831	224.9221184
RMRI N11	37	M	Before	Control population from endemi	NA	NA	72.4549921	120.914196	134.2372881	244.5482866
RMRI N12	60	M	Before	Control population from endemi	NA	NA	52.6307448	113.851735	92.54237288	180.6853583
RMRI N13	30	M	Before	Control population from endemi	NA	NA	66.7533281	48.3413249	138.3050847	236.4485981
RMRI N14	22	M	Before	Control population from endemi	NA	NA	47.8939778	49.6807571	100	260.7476636
RMRI N15	37	F	Before	Control population from endemi	NA	NA	46.4904913	45.9059937	108.8135593	219.6261682
RMRI C1	20	F	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	110.389045	44.2700565	112.5423729	200.623053
RMRI C2	62	F	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	121.738904	64.5514124	145.4237288	301.8691589
RMRI C3	15	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	76.1839888	36.7463277	95.59322034	185.6697819
RMRI C4	31	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	72.141573	41.1079096	116.9491525	241.7445483
RMRI C5	52	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	98.1063202	48.1954802	117.2881356	282.2429907
RMRI C6	18	F	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	68.8765449	36.7463277	96.61016949	200.623053
RMRI C7	45	F	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	110.544522	55.6101695	124.4067797	381.9314642
RMRI C8	32	F	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	94.8412921	43.0706215	106.1016949	232.3987539
RMRI C18	66	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Cured	61.4136236	35.9830508	91.86440678	203.4267913
RMRI R14	8	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Relapse	79.2935393	83.6333333	96.94915254	262.9283489
RMRI R16	49	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Relapse	56.9047753	102.279096	63.05084746	209.0342679
RMRI R17	18	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Relapse	59.7033708	101.079661	83.38983051	217.1339564
RMRI R18	59	M	Before	Confirmed VL, Hospitalization	Ambisome Single dose	Relapse	63.4348315	94.319209	84.40677966	304.6728972