1 **The transcription factor Capicua maintains the oocyte polarity in the panoistic**

2 **ovary of the German cockroach.**

5 Nashwa Elshaer[1][§], Jorge Escudero[1], Maria-Dolors Piulachs[1]*

7 *[1] Institute of Evolutionary Biology (CSIC-University Pompeu Fabra). Passeig Marítim de*
8 *la Barceloneta, 37. 08003 Barcelona, Spain*
9
10 [§] Present address: *Department of Plant Protection, Faculty of Agriculture, Zagazig*
11 *University, 44511 Zagazig, Egypt*

13 * Corresponding author: Maria-Dolors Piulachs (mdolors.piulachs@ibe.upf-csic.es)

Abstract

The establishment of the symmetry axis is crucial for the development of all organisms.

In insects, this process begins early in oogenesis with the correct distribution of the

mRNAs and proteins in the oocyte. One protein that plays a role in organizing this

distribution is the transcription factor Capicua (Cic). Cic has been studied in the context

of oogenesis and embryonic development in *Drosophila melanogaster*. It is maternally

expressed, begins essential for establishing the dorsoventral axis, and functions as a

transcriptional repressor. Although the Cic sequences are conserved across species, their

function in other types of insect ovaries is still little known. We wondered whether the

function of Cic in insects has been maintained through evolution despite the ovary type or

if it has been modified in parallel to the ovary evolution. To address this, we studied the

Cic function in a phylogenetically basal insect, the cockroach *Blattella germanica*, a

species with panoistic ovaries. Our findings show that *B. germanica* Cic is essential for

oocyte development and the maturation of ovarian follicles. A loss of Cic function leads

to disrupted cytoskeletal organization, defects in anterior-posterior polarity, and

compromised follicle integrity. The conservation and functional divergence of Cic across

different species suggest evolutionary adaptations in the mechanisms of insect oogenesis.

Keywords

Oocyte polarity, *Drosophila*, oogenesis, panoistic ovary, embryo development

1. Introduction

Oogenesis is a crucial process in the animal kingdom, ensuring species continuity. Therefore, it must be regulated with great precision. A good model for studying oogenesis is the insect ovary, especially, the panoistic ovary, characterized by the absence of nurse cells escorting the germinal cell or oocyte. Instead of this organization, each ovarian follicle only consists of an oocyte surrounded by a monolayered follicular epithelium (Büning, 1994). The panoistic ovary is characteristic of phylogenetically basal insects like the cockroach *Blattella germanica*, our insect model (Belles et al., 2024), and it is the most common ovarian type among invertebrates and vertebrates (Telfer, 1975). A distinguishing feature of *B. germanica* oogenesis is that the basal ovarian follicle (BOF), from each ovariole, is the only one that matures in each reproductive cycle, while the other ovarian follicles remain in the vitellarium, awaiting their chance to occupy the basal position in subsequent cycles (Belles et al., 2024; Irles & Piulachs, 2014; Tanaka & Piulachs, 2012) (Fig. 1). The BOFs begin to grow during the last nymphal instar. When vitellogenesis starts in 3-day-old adults, the BOF grows exponentially (Fig. 1), increasing its size tenfold. During this process, the follicular cells in the BOF proliferate, encompassing the oocyte growth. Coinciding with vitellogenesis, the follicular cells arrest cytokinesis, becoming binucleate and entering into endoreplication. By the end of the cycle, there is an amplification of the chorion genes, which are synthesized in a few hours to complete the egg (Fig. 1).

During oocyte growth, the organization of the future embryo is determined. The positioning of the nucleus and cell organelles, as well as the localization of specific proteins and RNAs within the oocyte, determine the establishment of oocyte polarity. This polarity will be essential later, during the embryogenesis, to reach a correct body plan development (Bashirullah et al., 1998; Bradley et al., 2000; Wilson et al., 2011). *Drosophila melanogaster*, with meroistic polytrophic ovaries, is the species in which oocyte polarity has been most thoroughly studied. The ovarian follicles in a polytrophic ovary are asymmetric, as they are formed by a cluster of nurse cells and the oocyte, all of them, surrounded by a monolayer of follicular cells (Büning, 1994). In this species, oocyte asymmetry is established in mid-oogenesis by the cell-cell interactions between the oocyte and follicular cells, which ultimately determine the final position of the oocyte

nucleus. The displacement of the oocyte nucleus to a dorsal-anterior position is facilitated

by the distribution of microtubules in the ooplasm, ensuring the proper localization of

mRNAs, like *Oskar* or *bicoid,* in the ooplasm (Roth & Lynch, 2012; Wilson et al., 2011).

Conversely, in panoistic ovaries, like those in *B. germanica*, the ovarian follicles exhibit

a bilateral symmetry. The changes in symmetry that occur during oocyte development are

subtle and occurs in adults, and information about this process remains limited. In *Acheta*

domesticus (Netzel 1968; Bradley et al., 2000) and *B. germanica* (Tanaka 1973; Zhang &

Kunkel, 1992), changes in the position of the oocyte nucleus, facilitate distinguishing the

dorsoventral axis. In both species, the nucleus moves from a central to the ventral

position. In *B. germanica*, the oocyte nucleus is maintained in a mid-ventral position

(Tanaka, 1973). In addition, the asymmetric distribution of F-actins in the oocyte

membrane, together with the maintenance of patency in the ventral follicular cells,

facilitate the identification of the ventral side of the BOF, when the oocyte reaches its

final size (Zhang & Kunkel, 1992).

The formation of the anteroposterior axis in oocytes of panoistic ovaries is even less well

known. One of the first indications of the presence of an anteroposterior axis was given

by Neztle (1968) in *A. domesticus* when he described the displacement of the nucleus

towards a ventral, slightly posterior position. Later, Bradlez and collaborators (Bradlez et

al., 2000), also in *A. domesticus*, described the formation of Balbani bodies in both poles,

anterior and posterior. Initially in the anterior pole and later in the posterior, this

determines the special localization of the RNAs and proteins. In the case of *B. germanica*,

we described a population of follicular cells in the apical pole of the BOF, which

differentiate at the end of the cycle to synthesize the egg structures (Pascual et al., 1992;

Irles et al., 2009). This limited available information requires further investigation about

the organization of oocytes and eggs in insects with panoistic ovaries.

We have adopted the panoistic ovary of *B. germanica* as a model, to study the function of

the transcription factor Capicua (Cic), which acts as a repressor and is structurally

conserved from cnidarians to vertebrates (Jiménez et al., 2000; Lam et al., 2006;

Rodrıguez-Muñoz et al., 2022). In *D. melanogaster*, two Cic isoforms have been

described. One short, Ci*c*-S, considered the canonical isoform (Rodrıguez-Muñoz et al.,

2022) reported as a maternally expressed gene that is required for the establishment of

dorsoventral patterning of the eggshell and the embryonic termini during embryogenesis
(Goff et al., 2001; Jiménez et al., 2000), the other longer isoform Cic-L, the most
common between metazoan, is expressed in all tissues (Rodrıguez-Muñoz et al., 2022). In
Tribolium castaneum, with meroistic telotrophic ovaries, the function of Cic in the
terminal patterning is conserved, although its depletion does not affect the dorsoventral
axis (Prido□lh. et al., 2017)
Our findings indicate that Cic is necessary for maintaining polarity in the ovarian follicles
of *B. germanica* and for stabilizing the distribution of F-actin in oocytes, a function that
seems to be performed early during oocyte development.
134
135

2. Materials and Methods

2.1. Insects
Blattella germanica (L.) specimens were obtained from a colony reared in the dark at 29
$\pm 1°C$ and 60–70% relative humidity (Bellés et al., 1987). Adult females were maintained
with males during all the first gonadotrophic cycle, and mating was confirmed at the end
of the experiments by assessing the presence of spermatozoa in the spermathecae. All
dissections and tissue sampling were carried out on carbon dioxide-anesthetized
specimens. Tissues were frozen on liquid nitrogen and stored at -80ºC until use.

2.2. Cloning and Sequencing
A fragment of *B. germanica capicua* (*cic*) sequence (493 bp) was obtained from an
mRNA library available in the laboratory. The sequence of *cic* was completed by 3´- and
5´- rapid amplification of cDNA ends (RACE), according to the kit protocol (Ambion,
Huntingdon, Cambridgeshire, UK). The amplified fragments were cloned into the
pSTBlue-1 vector (Novagen, Madison, WI, USA) and sequenced by Sanger (accession
number in NCBI: LN623700).

2.3. Sequence comparisons and phylogenetic analysis

Sequences of insect Cic protein, used in the phylogenetic analysis, were obtained from
GenBank. The search was enlarged by Blast to NCBI, using the *B. germanica* and *D.*
melanogaster Cic as queries. Cic sequences were selected taking representatives of
diverse insect taxa. Priority was given to species with well-annotated genomes to ensure
high-quality data. The accession numbers of the sequences used are listed in
Supplementary Table 1. The amino acid sequences were aligned using MAFFT (Katoh et
al., 2019). The resulting alignment was analyzed by the PHYML 3.0 program (Guindon
et al., 2010) based on the maximum-likelihood principle, using a JTT matrix, four
substitution rate categories with a gamma shape parameter of 1.444, and using empirical
base frequencies and estimating proportions. The data was bootstrapped for 100
replicates.
The domains NLS, TUDOR, N1, HMG, C2, and C1 in the Cic sequences of the studied
species were localized based on the description done in *H. sapiens* and *D. melanogaster*
in Webb et al, 2022, and Rodriguez-Muñoz et al., 2022 (See Fig. S1).
169

2.4. RNA extraction and expression studies

Total RNA was isolated using the GenElute Mammalian Total RNA kit (Sigma, Madrid,
Spain). An amount of 400 ng from each RNA extraction was DNAse treated (Promega,
Madison, WI, USA) and reverse transcribed with Superscript II reverse transcriptase
(Invitrogen, Carlsbad, CA, USA) and random hexamers (Promega). RNA quantity and
quality were estimated by spectrophotometric absorption at 260/280 nm in a Nanodrop
Spectrophotometer ND-1000® (NanoDrop Technologies, Wilmington, DE, USA).
Expression of *cic* was determined by quantitative real-time PCR (qRT-PCR) in the last
nymphal instar and the adult stage during the first gonadotrophic cycle. PCR primers
used in qRT-PCR expression studies were designed using the Primer Express 2.0
software (Applied Biosystems, Foster City, CA, USA), and are detailed in Supplementary
Table 2. The *B. germanica actin-5c* mRNA (Accession number AJ862721), was used as a
reference. qRT-PCR reactions were performed using the SYBR Green Supermix (BioRad)
containing 200 nM of each specific primer. Amplification reactions were carried out at
95ºC for 2 min, and 40 cycles of 95ºC for 15 s and 60ºC for 30s, using MyIQ Single
Color RTPCR Detection System (BioRad). After the amplification phase, a dissociation

curve was carried out to ensure that there was only one product. Levels of mRNA were

calculated relative to the respective reference gene, using the $2^{-\Delta\Delta Ct}$ method and $2^{-\Delta Ct}$

method (Irles et al., 2009; Livak & Schmittgen, 2001; Maestro et al., 2005). Results are

given as mRNA copies per 1000 copies of *actin-5c* mRNA and correspond to three

biological replicates.

2.5. RNA interference

To deplete *cic* mRNA, and asses the specificity of the phenotype, two different dsRNAs

(ds*cic*-1 and ds*cic*-2) were designed and synthesized as previously described (Ciudad et

al., 2006). The ds*cic*-1 was 324 bp in length (from nucleotides 5,874 to 6,197) and the

second one, ds*cic*-2, 392 bp (from nucleotides 4,366 to 4,757). A dsRNA, corresponding

to 307 bp of the *Autographa californica* nucleopolyhedrovirus sequence (GenBank:

K01149), was used as a control (dsMock). The corresponding cDNAs were amplified by

PCR and cloned into the pSTBlue-1 vector. Newly-emerged sixth instar nymph females,

or newly-emerged adult females, were independently treated with 1 µg of ds*cic*-1 or 1 µg

ds*cic*-2. Since the same ovary phenotype was observed regardless of which dsRNA was

used, we will refer to RNAi treatments collectively as ds*cic*. The same dose and

conditions applied in ds*cic* treatments were applied to dsMock-treated females.

2.6. Immunohistochemistry

After dissection, ovaries (n = 4, for each experiment and treatment) were fixed

immediately for 2 h with 4% paraformaldehyde in PBS, washed in PBT (PBS; 0.3%

Triton-X100), treated with 50 µg/ml proteinase K for 2 min, washed for 2 min in 2

mg/mL glycine in PBT, washed for 10 min in PBT and fixed again for 20 min in the

same solution. Then the ovaries were washed three times for 10 min with PBT (PBS;

0.1% Triton-X100, 0.1% BSA). After three washes with PBT the ovaries were saturated

for 1 h at room temperature in PBTBN (PBS; 0.1% Triton-X100, 0.5% BSA and 5%

normal goat serum), and incubated overnight at 4ºC with the primary antibody. The

primary antibodies used were rat anti-Cic (1:200, generated against the C-terminal half of

D. melanogaster Cic protein, kindly provided by Jordi Casanova; Molecular Biology

Institute of Barcelona, CSIC), mouse anti-EGFR (15µg/mL) and mouse anti-Eya (1:50).

Tissues were washed with PBTBN three times and incubated for 2 h with Alexa-Fluor

647 goat anti- mouse or anti-rat IgG (Molecular Probes, Carlsbad, CA, USA) diluted

1:400 in PBTBN. For F-actin visualization, ovaries were incubated for 20 min in 1µg/mL

phalloidin-TRITC (Sigma), and after washing they were mounted in UltraCruz™

Mounting Medium (Santa Cruz Biotechnology®, inc., Delaware CA, USA), which

contains DAPI for DNA staining. Samples were observed by epifluorescence microscopy

using a Zeiss AxioImager.Z1 microscope (ApoTome), using the software Zen 2012, blue

edition (Carl Zeiss MicroImaging). The relative fluorescence intensities of the Cic

labeling, DAPI, and phalloidin-TRITC, were graphically quantified using the *Zen 2.5D*

analysis tool (Carl Zeiss Microscopy).

2.7. Statistics

Data are expressed as mean ± standard error of the mean (S.E.M.). Statistical analysis of

gene expression values was carried out using the REST 2008 program (Relative

Expression Software Tool V 2.0.7; Corbett Research) (Pfaffl et al., 2002). This program

makes no assumptions about the distributions, evaluating the significance of the derived

results by Pair-Wise Fixed Reallocation Randomization Test tool in REST (Pfaffl et al.,

2002). The sample group was considered significantly different from the control group

when $P(H1) = 0.000$.

Results

3.1. *Blattella germanica* has a structurally conserved Capicua ortholog

capicua (*cic*) cDNA from *B. germanica*, was amplified, cloned, and sequenced from

ovarian tissues. The complete sequence of *B. germanica cic* has 7,631 bp with an ORF

encoding for a protein of 2,279 amino acids (nucleotide positions 553-7,392). It is similar

to the Cic-L isoform from *D. melanogaster*, and similar to the sequences of other insect

species available in databanks (Fig. S1). From the sequence comparisons, the Dipteran

Cic included in the analysis, could be an exception since they have a sequence

corresponding to the short isoform. *B. germanica* Cic possesses the HMG-box domain

(from amino acid 1,206 to 1,274) characteristic of Cic proteins highly conserved between

insect species (Fig. S1). In addition, the NLS, TUDOR, N1 and N2 domains, the helical

C1 module, and the C2 MAPK docking site, which helps DNA binding (Webb et al.,

2022), are also present in the insect species analyzed, showing also a high degree of

conservation (Fig. S1).

Maximum-likelihood analysis of Cic sequences from insects available in the literature

and databases gave a phylogenetic tree (Fig. 2) with two clades, clearly separated

corresponding to hemimetabolous (purple) and holometabolous (blue) insect species. The

topology of the phylogenetic tree of Cic sequences overlaps with the currently accepted

phylogeny of the included species (Misof et al., 2014). All insect orders appeared

grouped with high consistency in the nodes. Interestingly, the Cic sequences from

Lepidoptera and Diptera form two clusters with longer branches, indicating more

extensive changes and, consequently, a faster divergence rate compared to the Cic

proteins from other insect orders. The Cic-S and Cic-L from *D. melanogaster* appear to

be grouped in the same node (Fig. 2), suggesting they share the same origin (Fig. 2).

3.2. *cic* is highly expressed in the ovary during the last nymphal instar

The expression of *cic* mRNA was studied in different tissues from 3-day-old adult

females, namely the ovary, thoracic ganglia, colleterial glands, fat body, brain, and

midgut (Fig. 3A). *cic* is expressed in all tissues analyzed, with the highest levels in the

brain and ovary. In ovaries, the expression pattern of *cic* does not show significant

changes (Fig. 3B). In general, in ovaries from sixth instar nymphs, *cic* mRNA levels are

higher compared to the levels of the last day of the fifth nymphal instar. After the

imaginal molt, *cic* expression tends to decrease, reaching the lowest levels just before

oviposition (Fig. 3B). These expression patterns suggest that the main function of Cic in

the ovary occurs during the last nymphal instar.

3.3. Cic protein localizes in somatic and germ cells

A heterologous antibody against the *D. melanogaster* Cic protein was utilized to localize

Cic in *B. germanica* ovaries. Cic labeling was detected in ovaries of sixth instar nymphs,

in all ovarian follicles throughout the ovariole, encompassing germinal and somatic cells

(Fig. 3C). However, there are differences in the distribution of Cic in the growing oocyte.

In the basal follicle of 0-day-old sixth instar nymphs, Cic labeling appeared as a thin

layer adjacent to the oocyte membrane (Fig. 3D), with lower intensity at both the anterior

and posterior poles of the oocyte. Later, in 6-day and early 8-day-old sixth instar nymphs

(Fig. 3E and F, respectively), Cic labeling became more widespread through the ooplasm

and accumulated as a thick layer at the oocyte boundary. This labeling is displayed in all

ovarian follicles, from the basal to the most distal, as well as in the germarium (Fig. 3C).

In 8-day-old sixth instar nymphs, as the molt to adult approaches, Cic labeling appears in

the oocyte nucleus, spreading gradually with a faint signal throughout the cytoplasm,

becoming more prominent at the oocyte membrane (Fig. 3F'). After the imaginal molt,

Cic labeling in the oocyte becomes very weak (Fig. 3G). Concerning follicular cells, in 8-

day-old sixth instar nymphs, Cic appears abundant in the cytoplasm (Fig. 3H). In contrast,

in 0-day-old adults, the labeling in the follicular cells became fainter and is concentrated

at the basal pole (Fig. 3I). Overall, these findings indicate that Cic localization in the

ovary undergoes sequential and significant changes, suggesting distinct functional roles

of this transcription factor during oocyte maturation.

3.4. Cic is involved in the oocyte development of *B. germanica*.

To unveil the functions of Cic in the panoistic ovaries of *B. germanica*, 1µg of ds*cic* was

injected into two different experimental models: 0-day-old adult (ds*cic*-treated n = 45;

dsMock n = 20) and 0-day-old sixth instar nymphs (ds*cic*-treated n = 65; dsMock n = 20).

In ds*cic*-treated adult females just after the emergence, the ovarian mRNA levels for *cic*

were measured 48h after the treatment, exhibiting 27% reduction. This depletion was

sufficient to perturb oviposition and embryogenesis in the early stages. From the 45

treated females, 18% did not oviposit and 4% died before oviposition. Those treated

females that oviposited (78%, n = 35), 60% dropped the ootheca 48 – 72 hours after its

formation. The remaining females maintained the ootheca attached to the genital pouch,

which gave rise to nymphs that emerged normally without delays.

The females treated as 0-day-old sixth instar nymphs (n = 65), molted normally to the

adult 8 days later, similar to the dsMock-treated nymphs. At the end of the gonadotrophic

cycle, all dsMock-treated females oviposited and had a normal ootheca, whereas none of

the ds*cic*-treated females did. Ovarian *cic* mRNA levels were measured every 48 h during

the nymphal instar and just after the adult emergence (Fig. 4A). The results indicated that

in 8-day-old sixth instar nymphs treated with ds*cic*, the expression of *cic* mRNA tended

to decrease compared to those treated with dsMock (Fig. 4A). In addition, in these 8-day-

old ds*cic*-treated nymphs (Fig. 4B), no Cic protein labeling was detected in the BOF.

Only a faint signal was observed at the basal pole of the follicular cells (Fig. 4B and Fig.

S2). Conversely, in 8-day-old dsMock-treated nymphs, Cic labeling appeared in the BOF

ooplasm and the basal pole of the follicular cells (Fig. 4C, and Fig. S2).

The most evident effect observed in these ds*cic*-treated females was the change of the

BOF shape, which appeared elongated with defects affecting mainly the anterior and

posterior poles (Fig. 4B) a morphology that is very different from that of the dsMock-

treated females, which have the typical elliptical shape (Fig. 4C). A new batch of 0-day-

old ds*cic*-treated (n = 20) and dsMock-treated (n = 20), sixth instar nymphs, was prepared

to observe the ovarian follicles at different ages, in the last instar nymphs and adults.

In 8-day-old ds*cic*-treated sixth instar nymphs, the BOF appeared elongated (Fig. 4B)

compared to dsMock-treated nymphs (Fig. 4C). The anterior pole of the oocyte became

narrow extending into the lumen of the stalk (Fig. 5A and B, arrowheads) whereas, the

posterior pole showed a "fishtail-like" morphology (Fig. 5C and D, arrowheads). After

the molt, in 3-day-old adults, the BOF of ds*cic*-treated females appeared even more

elongated (Fig. 5E) and the apical pole of the oocyte that was entering the lumen of the

stalk, appeared engulfed by the follicular cells located in the anterior pole of the ovarian

follicle (Fig. 5F and G, arrowheads). Later, in 7-day-old adults, it is observed that the

ovarioles of females treated with ds*cic* have lost their basal and subbasal oocytes, leaving

only the sheath of the follicular epithelium that surrounded them (Fig. 5H, arrowheads).

Additionally, the remaining distal oocytes in the vitellarium, display an unusual hourglass

shape, probably due to the generalized enrichment of F-actin in the oocyte membrane and

ooplasm (Fig. 5H arrow, and I).

Usually, in last instar nymphs, at both poles of the younger oocytes in the vitellarium, is

observed an accumulation of F-actin, constituting the microtubule organization center

(MTOC) (Fig. 6A, arrowheads), with a network of F-actins connecting both MTOCs (Fig.

6A), which usually disappears in the basal oocytes. However, in the basal oocyte of 8-

day-old ds*cic*-treated nymphs, the MTOC remains present and strongly labeled (Fig. 6B

and C). It is also preserved and significantly larger in the BOF of 3-day-old adults from

the ds*cic*-treated group (Fig. 6D). The presence of the MTOCs at non-corresponding ages

may be the cause of changes in the shape of the ovarian follicles.

3.5. Depletion of *cic* mRNA affects the immature ovarian follicles

In the young ovarian follicles from 8-day-old ds*cic*-treated sixth instar nymphs (Fig. 7),

the nucleus in these oocytes is not in a central position, as occurs in controls (Fig. 7A).

Rather, it takes a posterior position, in the oocytes within the ovariole (Fig. 7B-D,

arrowheads). It seems that the nucleus was pushing the oocyte towards the posterior pole

of the ovarian follicle, and as a consequence, many oocytes can be encapsulated in one

ovarian follicle (Fig. 7D). This is facilitated by the absence of follicular cells in the poles

and the lack of stalk cells separating young ovarian follicles, which determined the fusion

of the oocyte membranes, thus giving the aspect of a single oocyte with more than one

nucleus (Fig. 7B-D, arrowheads). Moreover, the absence of the stalk connecting

successive ovarian follicles leads to a general disorganization of the ovariole, especially

affecting recently formed ovarian follicles, that are ready to leave the germarium. As a

consequence of these dysfunctions, in older adult females (7-day-old), the malformed

BOF has fallen, and is lost into the oviduct, before completing maturation (Fig. 5H).

3.6. Cic interacts with other signaling pathways in the ovary

The complex phenotypes produced by the depletion of *cic* mRNA suggests that Cic

protein could interact with other pathways related to cell proliferation and stalk formation.

As Cic is a downstream effector of EGFR signaling, we analyzed the EGFR in 8-day-old

sixth instar ds*cic*-treated nymphs. The expression of *EGFR* in ovaries tended to decrease,

although the differences compared with the control group were not statistically

significant (Fig. 8A). However, the localization of EGFR protein in ovaries of ds*cic*-

treated nymphs, appeared modified compared to dsMock-treated nymphs (Fig. 8B-D). In

ds*cic*-treated females, EGFR labeling was more restricted to the mid-ventral region of the

basal oocyte and around the nucleus (Fig. 8C-D). In contrast, in dsMock-treated nymphs,

EGFR localization is less restricted and spreads over the cytoplasm and within the
nucleus, with a higher concentration close to the nucleus membrane (Fig. 8B).
The absence of stalk between ovarian follicles in ds*cic*-treated specimens, and the
reduced number of follicular cells in some areas of the ovarian follicles, suggest us to
measure the expression of some components of the Hippo and Notch pathways. In *cic*
knockdown nymphs, *hippo* (*hpo*) was significantly down-regulated (0.351-fold change)
(Fig. 8A). We also measured the expression of *Notch* (*N*) and *Delta* (*Dl*), from the Notch
pathway (Irles et al., 2016; Irles & Piulachs, 2014), and as expected, the expression of *N*
tended to decrease, while its ligand *Dl* was significantly up-regulated (1.161-fold change;
Fig. 8A), suggesting that the Hippo and Notch pathways respond to Cic signaling.
Finally, the modifications observed in the anterior-posterior poles in ovarian follicles
from ds*cic*-treated sixth instar nymphs suggested looking for genes that drive the terminal
position in *D. melanogaster* egg chambers and embryos. Therefore, we examined the
response of *tailless* (*tll*) and *Dorsal* (DL) (Fig. 8A) in ds*cic*-treated nymphs. The
expression levels of *tll* were significantly down-regulated (0.355-fold change). While the
expression of the transcription factor DL, which is required for the determination of
dorsal-ventral polarity, decreased significantly (0.408-fold change) (Fig. 8A), reinforcing
the role of *cic* in the panoistic ovary, maintaining the polarity of the ovarian follicles.
390

4. Discussion
Cic is a transcription factor that is highly conserved among animals, from cnidarians to
vertebrates (Lam et al., 2006; C. J. Lee et al., 2002). The function of Cic has been
extensively studied in *D. melanogaster*, where two isoforms have been identified
(Jiménez et al., 2012; Rodrıguez-Muñoz et al., 2022). One is the Cic-S, considered the
canonical version, which appears mainly in dipterans (Fig. S1). The longer isoform,
found in most insect species, including *B. germanica*, was described in *D. melanogaster*
as responsible for maintaining the correct somatic and germinal cell cycle in the ovary
(Rodrıguez-Muñoz et al., 2022). These insect Cic sequences contain the typical domains
for the long Cic isoform (Forés et al., 2015), including the N2 motif, described previously
as Diptera exclusive (Forés et al., 2015). The phylogenetic analysis of the insect Cic

sequences suggests that the long isoforms could be ancestral, and the Cic-S was an
innovation in some insect groups.

The *B. germanica cic* is expressed in the brain and ovaries, similar to the *D.*
melanogaster Cic-L isoform, affecting brain size and development (Yang et al., 2016).
The high levels of *cic* expression in the *B. germanica* brain suggest that Cic may
participate in developing the central nervous system. However, we focused our research
on the expression and function of Cic in the panoistic ovary of *B. germanica*.

The role of Cic in insect oogenesis and embryo development has been thoroughly
documented in *D. melanogaster*. In this species, Cic participates in the specification of
terminal regions within the embryo; the loss of Cic function results in dorsalization of
both the eggshell and the embryo (Atkey et al., 2006; Goff et al., 2001; Jiménez et al.,
2000; Rodríguez-Muñoz et al., 2022), affecting the anterior-posterior embryonic axis
(Andreu et al., 2012). In *T. castaneum*, a species with meroistic telotrophic ovaries and a
short germband embryo, Cic is essential for anterior-posterior patterning without any
significant impact on dorsoventral pattern formation, and its depletion causes the terminal
anlagen to enlarge (Pridöhl et al., 2017). In the panoistic ovary of *B. germanica*, Cic
protein accumulates in the basal oocytes of sixth-instar nymphs as they progress in the
instar. However, after the imaginal molt, Cic labeling in the basal oocytes becomes weak,
suggesting that, in the panoistic ovary, the most important steps for oogenesis and further
oocyte maturation occur during the last nymphal instar. Thus, when the female reaches
the adult stage, the basal follicles are ready to incorporate storage proteins into the oocyte
and to complete growth (Belles et al., 2024; Ciudad et al., 2006; Herraiz et al., 2014; Irles
and Piulachs, 2014; Tanaka and Piulachs, 2012). According to our findings, Cic plays a
crucial role in oogenesis, which occurs before the ovarian follicle reaches maturity, and
its depletion causes female sterility. Equivalent observations have been reported for *D.*
melanogaster (Jiménez et al., 2000), where Cic has a fundamental function in oocyte
development, determining female sterility when depleted. However, the mechanisms
underlying this macroscopic phenotype differ between the two species with two ovary
types.

D. melanogaster Cic mutants can oviposit, although eggs have defects in the eggshell,
and the embryos begin to develop. However, they do not complete the embryogenesis

and cannot hatch (Goff et al., 2001; Rodríguez-Muñoz et al., 2022). Conversely, in *B.*
germanica, *cic*-depleted females do not lay eggs correctly because the oocyte cannot
complete its development. In *B. germanica*, Cic is required to maintain the anterior-
posterior axis since depletion of *cic* mRNA triggers morphological modifications in the
anterior-posterior poles of the BOFs, resulting in significant changes in the distribution of
F-actin microfilaments, in both the oocytes and the follicular cells.

Actin fibers are essential components of the cytoskeleton and play a crucial role in
various cellular processes, such as cell division, cell shape maintenance, oocyte growth,
and establishing oocyte polarity (Riparbelli & Callaini, 1995; Sun & Schatten, 2006). In
last instar nymphs, the MTOCs in the growing oocytes are located at the anterior and
posterior poles. These MTOCs disappear from the basal oocytes in the days leading up to
the molt into adulthood and are never detected in adult oocytes. However, in females
treated with ds*cic*, the MTOCs remain at the poles of the basal oocyte during the
transition from nymph to adult. The role of the MTOC in organizing the microtubule
network is crucial for the distribution of mRNAs and proteins within the oocyte, as
shown in studies involving *D. melanogaster* (Cooley & Theurkauf, 1994; Roth & Lynch,
2012). Although a network of F-actin can be observed connecting the two MTOCs at the
poles in BOFs from adult ds-*cic*-treated, we cannot provide information regarding the
function of MTOCs in the distribution of proteins or mRNAs in panoistic ovaries. The
varying prevalence of MTOCs at different stages suggests that this fact may influence the
shape changes observed in ovarian follicles.

Cic is a downstream effector of EGFR signaling that controls cell growth, differentiation,
and survival in different contexts (Goff et al., 2001; Elshaer and Piulachs, 2015). In *B.*
germanica EGFR-depleted nymphs, the anterior-posterior polarity is disrupted, leading to
ovarian follicles, and ovariole disorganization (Elshaer & Piulachs, 2015). The
restriction of EGFR protein to the mid-ventral region might explain the fusion of
immature oocyte membranes in ds*cic* specimens of *B. germanica*, determining the fusion
of some oocytes and suggesting that Cic in the panoistic ovary of *B. germanica* is
necessary to transduce the EGFR signaling, as happens in *D. melanogaster*, where Cic is
required to translate the EGFR signaling to pipe and establishing the nuclear dorsal
gradient (Moussian & Roth, 2005). In addition, the relationship between EGFR signaling

and cell proliferation has been established in different species. In *D. melanogaster*, the inhibition of Cic by EGFR leads to the de-repression of hpo and subsequently influences cell proliferation (Herranz et al., 2012). Conversely, our findings in *B. germanica* indicate that the depletion of *cic* results in the down-regulation of *hpo*. However, the follicular epithelium continues to proliferate, likely because Notch signaling within the ovarian follicle sustains an adequate level of proliferation, as previously reported (Irles & Piulachs, 2014). This indicates that the follicular epithelium can continue to proliferate even with low Hpo levels, due to the ongoing activity of N in the follicular cells. Previously, we demonstrated that Dl in young females is accumulated at the anterior and posterior poles of basal ovarian oocytes (Irles et al., 2016), interacting with N to determine specific cell populations in de BOF. The depletion of Dl in ds*cic*-treated females could also be responsible for the phenotypes observed in the BOFs.

Depletion of Cic in *B. germanica* seems to modify the expression of those genes that in *D. melanogaster* are required to establish the anterior-posterior polarity or the DV axis (like tailless and dorsal) (Lynch & Roth, 2011; Roth & Lynch, 2009), as they were down-regulated in the cockroach. It will be necessary to investigate the function of all these genes in a panoistic ovary, where the polarity of the oocyte differs greatly from that of a meroistic oocyte, and unveil how its function has been modified through evolution.

Acknowledgments

The work was funded with the grant PID2021-122316OB-I00 from the MICIU/AEI /10.13039/501100011033 and ERDF, a way of making Europe. We are grateful to Xavier Belles for the critical reading of the manuscript.

References

Andreu, M. J., González-Pérez, E., Ajuria, L., Samper, N., González-Crespo, S., Campuzano, S., & Jiménez, G. (2012). Mirror represses pipe expression in follicle cells to initiate dorsoventral axis formation in *Drosophila*. *Development (Cambridge, England)*, *139*(6), 1110–1114. https://doi.org/10.1242/DEV.076562

Atkey, M. R., Lachance, J. F. B., Walczak, M., Rebello, T., & Nilson, L. A. (2006). Capicua regulates follicle cell fate in the *Drosophila* ovary through repression of

mirror. *Development (Cambridge, England)*, *133*(11), 2115–2123.
https://doi.org/10.1242/DEV.02369

Bashirullah, A., Cooperstock, R.L., and Lipshitz, H.D. (1998). RNA localization in
development. *Annual Review of Biochemistry, 67*: 335-394.
https://doi.org/10.1146/annurev.biochem.67.1.335

Belles, X., Casas, J., Messeguer, A., & Piulachs, M. D. (1987). In vitro biosynthesis of
JH III by the corpora allata of adult females of *Blattella germanica* (*L*). *Insect*
Biochemistry, *17*(7), 1007–1010. https://doi.org/10.1016/0020-1790(87)90111-9

Belles, X., Maestro, J. L., & Piulachs, M. D. (2024). The German cockroach as a model
in insect development and reproduction in an endocrine context. *Advances in Insect*
Physiology, *66*(1), 1–47. https://doi.org/https://doi.org/10.1016/bs.aiip.2024.03.001

Bradley, J.T., Kloc, M., Wolfe, K.G., Estridge, B.H., Bilinski, S.M. (2000). Balbiani
bodies in cricket oocytes: Development, ultrastructure, and presence of localized
RNAs. *Differentiation, 67*: 117-127

Büning, J. (1994). The Insect Ovary. *The Insect Ovary*. https://doi.org/10.1007/978-94-
011-0741-9

Ciudad, L., Piulachs, M. D., & Bellés, X. (2006). Systemic RNAi of the cockroach
vitellogenin receptor results in a phenotype similar to that of the *Drosophila* yolkless
mutant. *The FEBS Journal*, *273*(2), 325–335. https://doi.org/10.1111/J.1742-
4658.2005.05066.X

Cooley, L., & Theurkauf, W. E. (1994). Cytoskeletal Functions During *Drosophila*
Oogenesis. *Science*, *266*(5185), 590–596.
https://doi.org/10.1126/SCIENCE.7939713

Elshaer, N., & Piulachs, M. D. (2015). Crosstalk of EGFR signalling with Notch and
Hippo pathways to regulate cell specification, migration and proliferation in
cockroach panoistic ovaries. *Biology of the Cell*, *107*(8), 273–285.
https://doi.org/10.1111/BOC.201500003

Forés, M., Ajuria, L., Samper, N., Astigarraga, S., Nieva, C., Grossman, R., González-
Crespo, S., Paroush,Z., Jiménez, G. (2015). Origins of context-dependent gene
repression by capicua. PLoS Genet 11(1): e1004902.
doi:10.1371/journal.pgen.1004902

Goff, D. J., Nilson, L. A., & Morisato, D. (2001). Establishment of dorsal-ventral polarity
of the Drosophila egg requires capicua action in ovarian follicle cells. *Development*,
128(22), 4553–4562. https://doi.org/10.1242/DEV.128.22.4553

Guindon, S., Dufayard J.F., Lefort V., Anisimova M., Hordijk W., Gascuel, O. (2010).
New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies:
Assessing the Performance of PhyML 3.0. *Systematic Biology*, *59*(3):307-21.
https://doi.org/10.1093/sysbio/syq010

Herraiz, A., Belles, X., & Piulachs, M. D. (2014). Chorion formation in panoistic ovaries
requires windei and trimethylation of histone 3 lysine 9. *Experimental Cell Research*,
320(1), 46–53. https://doi.org/10.1016/J.YEXCR.2013.07.006

Herranz, H., Hong, X., & Cohen, S. M. (2012). Mutual repression by bantam miRNA and
Capicua links the EGFR/MAPK and Hippo pathways in growth control. *Current
Biology: CB*, *22*(8), 651–657. https://doi.org/10.1016/J.CUB.2012.02.050

Irles, P., Bellés, X., & Piulachs, M. D. (2009). Brownie, a Gene Involved in Building
Complex Respiratory Devices in Insect Eggshells. *PLOS ONE*, *4*(12), e8353.
https://doi.org/10.1371/JOURNAL.PONE.0008353

Irles, P., Elshaer, N., & Piulachs, M. D. (2016). The Notch pathway regulates both the
proliferation and differentiation of follicular cells in the panoistic ovary of *Blattella
germanica*. *Open Biology*, *6*(1). https://doi.org/10.1098/rsob.150197

Irles, P., & Piulachs, M. D. (2014). Unlike in Drosophila Meroistic Ovaries, Hippo
Represses Notch in *Blattella germanica* Panoistic Ovaries, Triggering the Mitosis-
Endocycle Switch in the Follicular Cells. *PLOS ONE*, *9*(11), e113850.
https://doi.org/10.1371/JOURNAL.PONE.0113850

Jiménez, G., Guichet, A., Ephrussi, A., & Casanova, J. (2000). Relief of gene repression
by Torso RTK signaling: role of capicua in *Drosophila* terminal and dorsoventral
patterning. *Genes & Development*, *14*(2), 224. https://doi.org/10.1101/gad.14.2.224

Jiménez, G., Shvartsman, S. Y., & Paroush, Z. (2012). The Capicua repressor – a general
sensor of RTK signaling in development and disease. *Journal of Cell Science*,
125(6), 1383. https://doi.org/10.1242/JCS.092965

Katoh, K., Rozewicki, J., Yamada, K.D. (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization, *Briefings in Bioinformatics, 20* (4): 1160–1166, https://doi.org/10.1093/bib/bbx108

Lam, Y. C., Bowman, A. B., Jafar-Nejad, P., Lim, J., Richman, R., Fryer, J. D., Hyun, E. D., Duvick, L. A., Orr, H. T., Botas, J., & Zoghbi, H. Y. (2006). ATAXIN-1 interacts with the repressor Capicua in its native complex to cause SCA1 neuropathology. *Cell*, *127*(7), 1335–1347. https://doi.org/10.1016/J.CELL.2006.11.038

Lee, C. J., Chan, W. I., Cheung, M., Cheng, Y. C., Appleby, V. J., Orme, A. T., & Scotting, P. J. (2002). CIC, a member of a novel subfamily of the HMG-box superfamily, is transiently expressed in developing granule neurons. *Molecular Brain Research*, *106*(1–2), 151–156. https://doi.org/10.1016/S0169-328X(02)00439-4

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2-ΔΔCT method. *Methods*, *25*(4), 402–408. https://doi.org/10.1006/meth.2001.1262

Lynch, J. A., & Roth, S. (2011). The evolution of dorsal–ventral patterning mechanisms in insects. *Genes & Development*, *25*(2), 107. https://doi.org/10.1101/GAD.2010711

Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R, Butcher S. (2024). The EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024, *Nucleic Acids Research, 52*(W1): W521-W525. doi: 10.1093/nar/gkae241

Maestro, O., Cruz, J., Pascual, N., Martín, D., & Bellés, X. (2005). Differential expression of two RXR/ultraspiracle isoforms during the life cycle of the hemimetabolous insect *Blattella germanica* (*Dictyoptera, Blattellidae*). *Molecular and Cellular Endocrinology*, *238*, 27–37. https://doi.org/10.1016/j.mce.2005.04.004

Misof, B., Liu, S., Meusemann, K., Peters, R. S., Donath, A., Mayer, C., Frandsen, P. B., Ware, J., Flouri, T., Beutel, R. G., Niehuis, O., Petersen, M., Izquierdo-Carrasco, F., Wappler, T., Rust, J., Aberer, A. J., Aspöck, U., Aspöck, H., Bartel, D., … Zhou, X. (2014). Phylogenomics resolves the timing and pattern of insect evolution. *Science*, *346*(6210), 763–767. https://doi.org/10.1126/SCIENCE.1257570

586   Moussian, B., Roth, S. (2005). Dorsoventral Axis Formation in the Drosophila Embryo—
587   Shaping and Transducing a Morphogen Gradient, *Current Biology*, *15*(21), R887-
588   R899. https://doi.org/10.1016/j.cub.2005.10.026.

589   Netzel, H., (1968), The establishment of polarity and bilateral symmetry in the oocytes of
590   Gryllus domesticus L. *Wilhelm Roux' Archiv für Entwicklungsmechanik der*
591   *Organismen. 160* (1): 119-166. DOI: 10.1007/BF00573649

592   Pfaffl, M. W., Horgan, G. W., & Dempfle, L. (2002). Relative expression software tool
593   (REST) for group-wise comparison and statistical analysis of relative expression
594   results in real-time PCR. *Nucleic Acids Research*, *30*(9).
595   https://doi.org/10.1093/NAR/30.9.E36

596   Pridöhl, F., Weißkopf, M., Koniszewski, N., Sulzmaier, A., Uebe, S., Ekici, A. B., &
597   Schoppmeier, M. (2017). Transcriptome sequencing reveals maelstrom as a novel
598   target gene of the terminal system in the red flour beetle *Tribolium castaneum*.
599   Development 144 (7): 1339–1349. https://doi.org/10.1242/dev.136853

600   Riparbelli, M. G., & Callaini, G. (1995). Cytoskeleton of the *Drosophila* egg chamber:
601   new observations on microfilament distribution during oocyte growth. *Cell Motility*
602   *and the Cytoskeleton*, *31*(4), 298–306. https://doi.org/10.1002/CM.970310406

603   Rodrıguez-Muñoz, L., Lagares, C., Gonzalez-Crespo, S., Castel, P., Veraksa, A., &
604   Jimenez, G. (2022a). Noncanonical function of Capicua as a growth termination
605   signal in *Drosophila* oogenesis. Proceedings of the National Academy of Sciences
606   of the United States of America, 119 (31): 1-10.
607   https://doi.org/10.1073/PNAS.2123467119

608   Roth, S., & Lynch, J. (2012). Axis Formation: Microtubules Push in the Right Direction.
609   *Current Biology*, *22*(13), R537–R539. https://doi.org/10.1016/j.cub.2012.05.038

610   Roth, S., & Lynch, J. A. (2009). Symmetry breaking during *Drosophila* oogenesis. *Cold*
611   *Spring Harbor Perspectives in Biology*, *1*(2).
612   https://doi.org/10.1101/CSHPERSPECT.A001891

613   Sun, Q.-Y., & Schatten, H. (2006). Regulation of dynamic events by microfilaments
614   during oocyte maturation and fertilization. *Reproduction (Cambridge, England)*,
615   *131*(2), 193–205. https://doi.org/10.1530/REP.1.00847

Tanaka, A. (1973). General accounts on the oocyte growth and the identification of
vitellogenin by means of immunospecificity in the cockroach, *Blattella germanica*
(l.). *Development, Growth & Differentiation*, *15*(3), 153–168.
https://doi.org/10.1111/J.1440-169X.1973.00153.X

Tanaka, E.D., & Piulachs, M. D. (2012). Dicer-1 is a key enzyme in the regulation of
oogenesis in panoistic ovaries. *Biology of the Cell*, *104*(8), 452–461.
https://doi.org/10.1111/BOC.201100044

Telfer, W. H. (1975). Development and Physiology of the Oöcyte-Nurse Cell Syncytium.
Advances in Insect Physiology, *11*(C), 223–319. https://doi.org/10.1016/S0065-
2806(08)60164-2

Webb, J., Liew, J.J.M., Gnann, A.D., Patterson, M., Paul, S., Forés, M., Jiménez, G.,
Veraksa, A., Dowling, D.P. (2022). Molecular basis of DNA recognition by the
HMG-box-C1 module of Capicua. *bioRxiv* 2022.03.28.485992; doi:
https://doi.org/10.1101/2022.03.28.485992

Wilson, M. J., Abbott, H., & Dearden, P. K. (2011). The evolution of oocyte patterning in
insects: multiple cell-signaling pathways are active during honeybee oogenesis and
are likely to play a role in axis patterning. *Evolution & Development*, *13*(2), 127–
137. https://doi.org/10.1111/J.1525-142X.2011.00463.X

Yang, L., Paul, S., Trieu, K.G., Dent, L.G., Froldi, F., Forés, M., Webster, K., Siegfried,
K.R., Kondo, S., Harvey, K., Cheng, L., Jiménez, G., Shvartsman, S.Y., Veraksa, Y.
(2016). Minibrain and Wings apart control organ growth and tissue patterning
through down-regulation of Capicua. *PNAS 113* (38) 10583-10588.
https://doi.org/10.1073/pnas.1609417113

Zhang, Y., & Kunkel, Joseph. G. (1992). Program of F-actin in the follicular epithelium
during oogenesis of the german cockroach, *Blattella germanica*. *Tissue and Cell*,
24(6), 905–917. https://doi.org/10.1016/0040-8166(92)90025-3

642
643
644
645 **Figure Legends**
646

Figure 1. Changes in the *Blattella germanica* basal ovarian follicle, during the
gonadotropic cycle. Geometric growth of the basal ovarian follicle (BOF), in key
moments of the gonadotropic cycle. The BOFs grow linearly during the last instar
nymphs, until 3-day-old adults. During this period, the follicular cells (FCs) surrounding
the basal oocyte proliferate. In 3-day-old females, the vitellogenesis begins, and the BOF
grows exponentially, increasing ten times its size. The oocyte nucleus migrates to the
ventral pole, where the germinal band will be formed. The FCs arrest cytokinesis
becoming binucleated, and enter endoreplication. In the last days of the cycle, these cells
amplify some genome regions to improve the synthesis of the chorion proteins.

Figure 2. Phylogenetic analysis of the insect transcription factor Capicua, The
analysis is based on the maximum-likelihood principle, with the amino acid substitution
model, and was carried out using insect Cic sequences available in public databases. The
Cic sequence of *B. germanica* is highlighted in bold grey. Blue branches indicate species
belonging to holometabolous insects, while purple branches represent species belonging
to hemimetabolous insects. Only bootstrap with values greater than 60 are shown. The
complete species names and accession numbers of the sequences are listed in Table S1.
The scale bar indicates the number of substitutions per site.

Figure 3. ***B. germanica capicua* (*cic*). Expression and localization**. (A) Relative
expression of *cic* mRNA in ovary (Ov), thoracic ganglia (TG), colleterial glands (CG), fat
body (FB), brain (Br), and midgut (MG) from adult females. (B) Relative expression of
cic in the ovary during the sixth (last) instar nymph and in the adult. The expression on
the last day of the fifth nymphal instar ovaries was also measured. Data in (A) and (B) are
expressed as copies per 1000 copies of *actin-5c* (n = 3-4). (C) Immunolocalization of Cic
in an ovariole from a 6-day-old sixth instar nymph showing the asymmetric distribution
of the actin in the basal ovarian follicle (BOF) and sub-basal ovarian follicle (sBOF), TF:
terminal filament. (D-F) Immunolocalization of Cic in basal follicles from 0-day-old (D),
6-day-old (E), and 8-day-old (F and F') sixth instar nymphs, showing the changes of Cic
distribution related to the age of the female, from an accumulation in the oocyte
membrane (D, E) to be extended by the cytoplasm and nucleus (F'). In these images, the

exposure time is the same, to allow comparisons. (G) Weakly Cic labeling in the
follicular cells of the anterior and posterior poles in the BOF from 6-day-old sixth instar
nymphs. (H-I) Optical section of BOFs from 8-day-old sixth instar nymph (H) and 0-day-
old adult (I), showing differences in Cic distribution and labeling intensity within the
ooplasm and the follicular cells. The arrows, in both images, indicate the layer of
bacteriocyte endosymbionts. Oo: ooplasm. The anterior pole in C and G is towards the
right, and in D-F it is to the top of the image. Cic: green, F-actin: red, DAPI: blue.

Figure 4. **Capicua (Cic) in ovaries of ds*cic*-treated females**. (A) Relative expression of
cic mRNA in ovaries of sixth instar nymphs and newly emerged adult ds*cic*-treated. ds*cic*
was applied to nymphs just after the molt to last instar and the RNA levels were
measured at different developmental stages (N6D2, N6D4, N6D6, N6D8 and AdD0, n =
3 for each age and treatment). Data are expressed as copies of *cic* per 1000 copies of
actin-5c. (B) Immunolocalization of Cic protein in basal follicles of 8-day-old ds*cic*-
treated (n = 3) and (C) in dsMock-treated sixth instar nymph (n = 3). The anterior pole of
the basal follicle in A and B is towards the right. Cic: green, F-actin: red, DAPI: blue. In
Figure S2 the labeling intensity for each image is show.

Figure 5. Depletion of *cic* in basal ovarian follicles of different ages. Last instar
nymphs were treated the day of emergence, and the phenotypes analyzed at different ages.
(A, B) Basal ovarian follicles (BOF) from 8-day-old ds*cic*-treated sixth instar nymphs,
showing different degrees of narrowing in the anterior pole (arrowheads). (C, D) BOFs
exhibiting the "fishtail-like" phenotype, with a concentration of F-actin in the posterior
pole (arrowheads). (E-G) Ovarioles from 3-day-old ds*cic*-treated adults, showing the
elongated shape of the BOF (E), and the engulfment of the anterior pole of the basal
oocyte by the follicular epithelia (F, G; arrowheads). (H) Ovarioles from a 7-day-old
ds*cic*-treated adult. The basal and sub-basal oocytes were prematurely oviposited
(arrowheads) before completing their development, leaving the follicular epithelia that
surround those oocytes. In these ovarioles, the new first oocyte shows an hourglass shape
(arrows). (I) detail of the new first oocyte showing an hourglass shape. In addition, it was

not possible to detect the stalk separating the ovarian follicles. The anterior pole is

towards the left. F-actin: red (except in I, which is white), DAPI: blue.

Figure 6. MTOC in ovarian follicles from N6D8 and AdD3 treated with ds*cic* newly-

emerged sixth instar nymph. (A) Sub-basal ovarian follicle from an 8-day-old-dsMock

treated sixth instar nymphs, showing the MTOCs placed in the anterior and posterior

oocyte poles (arrowheads), interconnected by and F-actin network. (B, C) BOF from 8-

day-old ds*cic*-treated sixth instar nymph, showing an abnormally enlarged MTOC in the

anterior pole of the oocytes. (D) BOF from a 3-day-old ds*cic*-treated adult showing the

network of F-actin, with a big MTOC in the anterior pole of the oocyte, when, usually at

this age, the MTOC and the F-actins network, are not detected. The anterior pole is

towards the left. F-actin: red (except in D that is white), DAPI: blue.

Figure 7. Localization of the oocyte nucleus in ds*cic*-treated sixth instar nymphs. (A)

Ovariole from an 8-day-old dsMock-treated nymph, showing the normal localization of

oocyte nucleus in the different ovarian follicles along the ovarioles (BOF: basal ovarian

follicle; sBOF: sub-basal ovarian follicle). (B, C) Disorganization of the ovarian follicles

in the vitellarium due to the stalk loss showing the mislocalization of the oocyte nucleus

(The arrowhead indicates the expected position of the stalk, which is absent). (D) Fusion

of two oocytes resulting in a single ovarian follicle containing two nuclei. In B the

anterior pole is up towards the left. The oocyte nucleus is labeled by anti-Eya antibody

(green); F-actin: red; DAPI: Blue.

Figure 8. Genes involved in oocyte growth and development. (A) Relative expression

of *EGFR, notch* (*N*)*, delta* (*Dl*)*, hippo* (*hpo*)*, tailless* (*tll*)*, and *dorsal* (*DL*) in ovaries from

8-day-old ds*cic*-treated sixth instar nymphs. Data represent normalized values against

dsMock controls (reference value = 1) (n = 3) (*) indicates statistically significant change

P(H1) = 0.0001. (B) Basal ovarian follicle from 8-day-old dsMock-treated sixth instar

nymph, showing the EGFR protein with high labeling on nuclear membrane and inside

the oocyte nucleus. (C-D) Basal ovarian follicles of a ds*cic*-treated nymph showing

changes in EGFR localization. In these follicles EGFR labeling appeared restricted to the

central region of the oocyte around the nucleus and towards the oocyte ventral side (ar-

rowhead). F-actin: red; DAPI: Blue, EGFR protein: Green.

DNA
Actin
Cic

8-day-old sixt instar nymphs | 3-day-old adult

sBOF dsMock | ds*cic*-treated | ds*cic*-treated

A. sBOF, BOF. 50 μm

B. BOF. 50 μm

C. BOF. 50 μm

D. BOF. 50 μm

A

Relative expression

EGFR   N   Dl   Hpo   tll   DL

B

DNA
Actin
EGFR

BOF

50 μm   dsMock

C

DNA
Actin
EGFR

BOF

50 μm

ds*cic*

D

ds*cic*

DNA
Actin
EGFR

BOF

50 μm

Figure S1. Multiple sequence alignment of the Capicua protein across different insect species, using MAFFT (Katoh et al., 2013). Key functional domains—NLS, Tudor, N1, N2, HMG, C1, and C2—are highlighted in distinct colors (red, blue, green, violet, orange, purple, and fluorescent green, respectively) to indicate their locations and relative conservation. Highly conserved regions are marked, emphasizing the evolutionary stability of these functional motifs. The characteristic phenylalanine residues within the C1 domain are specifically indicated (red arrows). Accession numbers from different species are provided in Supplementary Table 1

Figure S2. **Cic protein in basal ovarian follicles of ds*cic* treated females**. (A) Cic protein in basal follicles of 8-day-old ds*cic*-treated and (B) in dsMock-treated sixth instar nymph (Images from Figure 4). The relative fluorescence intensities of the different fluorophores in the Figure 4B and C,are shown in B' and C'. It was graphically quantified using the Zen 2.5D analysis tool (Carl Zeiss Microscopy).The anterior pole of the basal follicle in A and B is towards the right. Cic: green, F-actin: red, DAPI: blue.