1 **Loss of O-specific antigen shapes *Pseudomonas aeruginosa* population microbiogeography in**
2 **murine preclinical pulmonary infection model**

3 Fraser H.[1], Moustafa D. A.[2,3], Goldberg J. B.[2,3], Azimi S.[1]

4
5
6 1. Biology Department, College of Arts and Sciences, Georgia State University, Atlanta, GA, 30303.
7 2. Division of Pulmonary, Asthma, Cystic Fibrosis, and Sleep, Department of Pediatrics, Emory University
8 School of Medicine, Atlanta, GA 30322.
9 3. Emory-Children's Cystic Fibrosis Center, Atlanta, GA 30322.
10
11 To whom correspondence may be addressed. Email: sazimi@gsu.edu or dina.moustafa@emory.edu

12
13
Abstract

Chronic *Pseudomonas aeruginosa* infections are the hallmark of late-stage lung disease in individuals
with cystic fibrosis. During chronic infection *P. aeruginosa* becomes the dominant bacteria in the
airway. Within-host adaptation of *P. aeruginosa* leads to vast phenotypic and genetic population
heterogeneity. *In vitro* studies show mutations in lipopolysaccharide (LPS) O-specific antigen
changes the aggregate formation in *P. aeruginosa*, however role of these changes in aggregate
assembly *in vivo* is not understood. Using a synthetic CF sputum media and a preclinical murine
infection model we assessed how the PAO1 wildtype and O-specific antigen mutants interact with
each other, and if *P. aeruginosa* population heterogeneity affects the colonization of the murine lungs.
Our findings suggest that the presence of variants lacking O-specific antigen does not impact the
population fitness and size in both *in vitro* and *in vivo,* however it can influence the aggregate volume
in vivo.

Chronic infections are usually polymicrobial, where bacteria live in biofilms or as suspended
aggregates of $10–10^4$ cells with defined spatial patterning and distances relative to host cells and
other microorganism[1-3]. In chronic infections, within-host adaptation, genetic drift, and antimicrobial
selection are key drivers of intrastrain genetic heterogeneity of bacterial populations. Although the
intrastrain genetic heterogeneity is known, its impact on pathogenesis, and microbiogeography of
infections remains unclear.
In chronic pulmonary infection with *Pseudomonas aeruginosa* in people with cystic fibrosis (pwCF)
is the hallmark of loss of lung function. Divergent evolution of *P. aeruginosa,* and accumulation of
different genetic variants in CF airways and its impact on population functional phenotypes is well
documented[3-6], but how this intrastrain heterogeneity influences polymicrobial community and
airway focal injury remains unknown. We recently showed that mutations in genes encoding
Lipopolysaccharide (LPS) O-specific antigen (OSA) biosynthesis and assembly such as *ssg* and *wbpL*
increase *P. aeruginosa* cell surface hydrophobicity, and alter *P. aeruginosa* aggregate assembly in *in*
vitro preclinical CF sputum model (SCFM2)[7-9]. Considering *P. aeruginosa* variants with mutations in
OSA commonly isolated from chronic CF airway infections[10-13], along with LPS smooth phenotypes, it
is important to determine the role of OSA genetic and phenotypic heterogeneity on spatial
organization, and *P. aeruginosa* population fitness.
43
44   To address the influence of genetic heterogeneity on spatial organization of *P. aeruginosa* population
45   in the airways, we examined whether presence of variants with mutations in OSA biosynthesis and
46   assembly influence the aggregate assembly of *P. aeruginosa* populations *in vitro*. We constructed
47   mixed populations of PAO1WT with either PAO1Δ*ssg* or, PAO1Δ*wbpL* at 1:1 initial ratio (5×10^6: $5 \times$

Fig. 1. Loss of OSA due to mutations in *ssg* and *wbpL* genes, alters the biogeography and aggregate assembly of *P. aeruginosa* populations. (a) Presence of both PAO1Δ*ssg* and PAO1Δ*wbpL* lead to increase of total population size, but (b) not the population structure, as the relative abundance of PAO1, PAO1Δ*ssg* and, PAO1Δ*wbpL* in mixed populations remains the same. (c) Changes of aggregate assembly from Stacked to Clumped due to loss of OSA and increases cell surface hydrophobicity, and presence of variants without OSA alters the biogeography of mixed populations (lower panel), (data presented from three independent experiments).

10⁶ CFU) in SCFM2. We grew each variant and the parental PAO1 strain in isolation and mixed
populations for 24 hours and evaluated the changes in total population size, and changes in the ratio
of PAO1Δ*ssg* and PAO1Δ*wbpL* compared to the wild type PAO1 strain. We assessed the abundance of
each variant by qPCR and by enumerating colony forming unites (CFUs). Although we detected a
significant change in aggregate assembly of WT cells in presence of ΔOSA variants (PAO1Δ*ssg* and
PAO1Δ*wbpL*), we did not detect any significant changes in proportion of each variant in mixed
populations *in vitro* (Fig. 1 and b). Interestingly, we observed that presence of ΔOSA variants
significantly disrupts the stacked aggregate assembly of parental strain (Fig. 1c).

Although presence of ΔOSA variants alters the population biogeography *in vitro,* it is not clear how
the changes in population structure affect the spatial organization in airways. To examine how
presence of ΔOSA variants influence airway colonization, we modified an established preclinical
acute airway infection model[14, 15] of six-week-old female BALB/c mice. We prepared cultures of PAO1,
PAO1Δ*ssg,* PAO1Δ*wbpL*, and synthetic mixed populations of PAO1: PAO1Δ*ssg* and PAO1: PAO1Δ*wbpL*
in SCFM2 and incubated the cultures 4-6 hours at 37 °C. We then infected the mice with 25 μL of the
standard inoculum 1x 10⁷ CFU of each bacterial population. We then sacrificed the mice 24 hours
post infection for further analysis of colonization rate and bacterial spatial organization in the
airways (All animal procedures were conducted according to the guidelines of the Emory University
Institutional Animal Care and Use Committee, under approved protocol number PROTO201700441).
Briefly, we determined the abundance of each *P. aeruginosa* variants by homogenizing the lungs, and
enumerating CFUs[15], and qPCR (Fig. 2 a and b). In brief, we perfused the lungs with 15 ml of 20Uml
of heparin in PBS. We then fixed the lungs in 500 ml of 4% PFA/PBS (v/v) for future

Fig. 2. Presence of PAO1Δ*ssg* increase the PAO1 aggregate size in murine airway. (a) Presence of ΔOSA variants in a population does not significantly impact colonization and population size in murine airways. (b) Loss of OSA does not influence *P. aeruginosa* fitness in murine airway. (c) Use of iDISCO clearing method allows for whole lung imaging and localization of bacteria in the airways. (d) Presence of PAO1Δ*ssg* does not affect the abundance of WT cells but (e) leads to formation of larger aggregates (df= 1011.2, *p*< 0.001).

immunohistochemistry, hybridization chain reaction (HCR)-probing[16, 17], and tissue-clearing
following iDISCO[18](Fig. 2c).
We measured the colonization level of each variant and mixed populations in the airways by
homogenizing the lungs in 1ml of PBS, followed by enumerating the CFUs. We further extracted the
genomic DNA from lung homogenates and measured the abundance of each variant in mixed
populations using qPCR. We found that similar to what we observed in *in vitro,* loss of OSA due to
deletion of *ssg* and *wbpL* does not influence *P. aeruginosa* fitness in murine airways either in single
or in mixed populations (Fig. 2 b and c). To determine whether presence of ΔOSA variants in *P.*
aeruginosa populations alter the spatial organization and microbiogeography of the airways, we used
confocal and light sheet microscopy. To visualize the bacteria in the lungs, we infected the mice with
PAO1Δ*ssg*: *gfp* and used HCR-probes targeting *ssg* mRNA to detect and visualized PAO1 WT cells. We
used Imaris.10 software to measure the aggregate volumes and observed a significant change in
PAO1WT aggregate volumes in presence of PAO1Δ*ssg* cells from average 178 μm^3 to 898 μm^3 (Fig. 2
d and e).
In summary, our study highlights the significant role that genetic heterogeneity within *P. aeruginosa*
populations plays in shaping bacterial spatial organization, particularly in chronic infections such as
those found in CF. Despite no measurable impact on overall population fitness, the presence of
variants with OSA mutations notably alters aggregate assembly both *in vitro* and *in vivo*. These
findings suggest that population genetic heterogeneity have the potential to influence the structural
organization of bacterial communities, which can subsequently impact the infection progression,
host immune responses at micron scale, and treatment outcomes. Future studies assessing the
impacts of genetic heterogeneity in clinically sourced populations, on host pathogen interactions and
pathogenesis, will be crucial for understanding the broader implications of *P. aeruginosa* genetic
diversity in chronic infections and focal airway injury.
94
95

Acknowledgement: We would like to acknowledge members of the Azimi, and Goldberg labs for
their assistance in setting up the animal experiments. This study was supported by Georgia State
University startup to SA, and CFF-WHITELE20A0 to JBG.

f. Literature Cited

1.   Mark Welch JL, Dewhirst FE, Borisy GG. Biogeography of the Oral Microbiome: The Site-
Specialist Hypothesis. Annu Rev Microbiol. 2019;73:335-58. Epub 2019/06/11. doi:
10.1146/annurev-micro-090817-062503. PubMed PMID: 31180804; PMCID: PMC7153577.
2.   Kolpen M, Kragh KN, Enciso JB, Faurholt-Jepsen D, Lindegaard B, Egelund GB, Jensen AV,
Ravn P, Mathiesen IHM, Gheorge AG, Hertz FB, Qvist T, Whiteley M, Jensen PO, Bjarnsholt T. Bacterial
biofilms predominate in both acute and chronic human lung infections. Thorax. 2022. doi:
10.1136/thoraxjnl-2021-217576. PubMed PMID: WOS:000741509200001.
3.   Azimi S, Lewin GR, Whiteley M. The biogeography of infection revisited. Nat Rev Microbiol.
2022. Epub 2022/02/10. doi: 10.1038/s41579-022-00683-3. PubMed PMID: 35136217.
4.   Azimi S, Roberts AEL, Peng S, Weitz JS, McNally A, Brown SP, Diggle SP. Allelic polymorphism
shapes community function in evolving *Pseudomonas aeruginosa* populations. ISME J. 2020. Epub
2020/04/29. doi: 10.1038/s41396-020-0652-0. PubMed PMID: 32341475.
5.   Vanderwoude J, Fleming D, Azimi S, Trivedi U, Rumbaugh KP, Diggle SP. The evolution of
virulence in *Pseudomonas aeruginosa* during chronic wound infection. Proc Biol Sci.
2020;287(1937):20202272. Epub 20201021. doi: 10.1098/rspb.2020.2272. PubMed PMID:
33081616; PMCID: PMC7661302.
6.   Rossi E, La Rosa R, Bartell JA, Marvig RL, Haagensen JAJ, Sommer LM, Molin S, Johansen HK.
Pseudomonas aeruginosa adaptation and evolution in patients with cystic fibrosis. Nat Rev
Microbiol. 2021;19(5):331-42. Epub 20201119. doi: 10.1038/s41579-020-00477-5. PubMed PMID:
33214718.
7.   Azimi S, Thomas J, Cleland SE, Curtis JE, Goldberg JB, Diggle SP. O-Specific Antigen-
Dependent Surface Hydrophobicity Mediates Aggregate Assembly Type in *Pseudomonas aeruginosa*.
mBio. 2021:e0086021. Epub 2021/08/11. doi: 10.1128/mBio.00860-21. PubMed PMID: 34372703.
8.   Cornforth DM, Diggle FL, Melvin JA, Bomberger JM, Whiteley M. Quantitative Framework for
Model Evaluation in Microbiology Research Using *Pseudomonas aeruginosa* and Cystic Fibrosis
Infection as a Test Case. mBio. 2020;11(1). Epub 2020/01/16. doi: 10.1128/mBio.03042-19.
PubMed PMID: 31937646; PMCID: PMC6960289.
9.   Lewin GR, Kapur A, Cornforth DM, Duncan RP, Diggle FL, Moustafa DA, Harrison SA, Skaar
EP, Chazin WJ, Goldberg JB, Bomberger JM, Whiteley M. Application of a quantitative framework to
improve the accuracy of a bacterial infection model. Proc Natl Acad Sci U S A.
2023;120(19):e2221542120. Epub 20230501. doi: 10.1073/pnas.2221542120. PubMed PMID:
37126703; PMCID: PMC10175807.
10.   Cigana C, Curcuru L, Leone MR, Ierano T, Lore NI, Bianconi I, Silipo A, Cozzolino F, Lanzetta
R, Molinaro A, Bernardini ML, Bragonzi A. *Pseudomonas aeruginosa* exploits lipid A and
muropeptides modification as a strategy to lower innate immunity during cystic fibrosis lung
infection. PLoS One. 2009;4(12):e8439. Epub 2009/12/29. doi: 10.1371/journal.pone.0008439.
PubMed PMID: 20037649; PMCID: PMC2793027.
11.   Malhotra S, Hayes D, Jr., Wozniak DJ. Mucoid *Pseudomonas aeruginosa* and regional
inflammation in the cystic fibrosis lung. J Cyst Fibros. 2019;18(6):796-803. Epub 20190426. doi:
10.1016/j.jcf.2019.04.009. PubMed PMID: 31036488; PMCID: PMC6815243.
12.   Mathee K, Ciofu O, Sternberg C, Lindum PW, Campbell JI, Jensen P, Johnsen AH, Givskov M,
Ohman DE, Molin S, Hoiby N, Kharazmi A. Mucoid conversion of *Pseudomonas aeruginosa* by
hydrogen peroxide: a mechanism for virulence activation in the cystic fibrosis lung. Microbiology.
1999;145 (Pt 6):1349-57. Epub 1999/07/20. doi: 10.1099/13500872-145-6-1349. PubMed PMID:
10411261.
13.   Tognon M, Kohler T, Gdaniec BG, Hao Y, Lam JS, Beaume M, Luscher A, Buckling A, van
Delden C. Co-evolution with *Staphylococcus aureus* leads to lipopolysaccharide alterations in
Pseudomonas aeruginosa. ISME J. 2017. Epub 2017/05/27. doi: 10.1038/ismej.2017.83. PubMed
PMID: 28548661.

14.   Mukherjee S, Moustafa DA, Stergioula V, Smith CD, Goldberg JB, Bassler BL. The PqsE and
RhlR proteins are an autoinducer synthase-receptor pair that control virulence and biofilm
development in *Pseudomonas aeruginosa*. Proc Natl Acad Sci U S A. 2018;115(40):E9411-E8. Epub
20180917. doi: 10.1073/pnas.1814023115. PubMed PMID: 30224496; PMCID: PMC6176596.
15.   Duncan RP, Moustafa DA, Lewin GR, Diggle FL, Bomberger JM, Whiteley M, Goldberg JB.
Improvement of a mouse infection model to capture *Pseudomonas aeruginosa* chronic physiology in
cystic fibrosis. Proc Natl Acad Sci U S A. 2024;121(33):e2406234121. Epub 20240805. doi:
10.1073/pnas.2406234121. PubMed PMID: 39102545; PMCID: PMC11331117.
16.   Choi HM, Calvert CR, Husain N, Huss D, Barsi JC, Deverman BE, Hunter RC, Kato M, Lee SM,
Abelin AC, Rosenthal AZ, Akbari OS, Li Y, Hay BA, Sternberg PW, Patterson PH, Davidson EH,
Mazmanian SK, Prober DA, van de Rijn M, Leadbetter JR, Newman DK, Readhead C, Bronner ME,
Wold B, Lansford R, Sauka-Spengler T, Fraser SE, Pierce NA. Mapping a multiplexed zoo of mRNA
expression. Development. 2016;143(19):3632-7. Epub 2016/10/06. doi: 10.1242/dev.140137.
PubMed PMID: 27702788; PMCID: PMC5087610.
17.   DePas WH, Starwalt-Lee R, Van Sambeek L, Ravindra Kumar S, Gradinaru V, Newman DK.
Exposing the Three-Dimensional Biogeography and Metabolic States of Pathogens in Cystic Fibrosis
Sputum via Hydrogel Embedding, Clearing, and rRNA Labeling. mBio. 2016;7(5). Epub 2016/09/30.
doi: 10.1128/mBio.00796-16. PubMed PMID: 27677788; PMCID: PMC5040109.
18.   Habart M, Lio G, Soumier A, Demily C, Sirigu A. An optimized iDISCO+ protocol for tissue
clearing and 3D analysis of oxytocin and vasopressin cell network in the developing mouse brain.
STAR Protoc. 2023;4(1):101968. Epub 20230103. doi: 10.1016/j.xpro.2022.101968. PubMed PMID:
36598854; PMCID: PMC9826885.

172