Apical clathrin-coated endocytic pits control the growth and size of epithelial microvilli

Olivia L. Perkins[1], Alexandra G. Mulligan[1], Evan S. Krystofiak[1], K. Elkie Peebles[1], Leslie M. Meenderink[1,3], Bryan A. Millis[1,2], Matthew J. Tyska[1*]

[1]Department of Cell and Developmental Biology, Vanderbilt University School of Medicine, Nashville, TN 37232
[2]Department Biomedical Engineering, Vanderbilt University School of Engineering, Nashville, TN 37232
[3]United States Department of Veterans Affairs Tennessee Valley Health Care System, Nashville, TN, 37232

*To whom correspondence should be addressed: matthew.tyska@vanderbilt.edu

____

ABSTRACT **Epithelial cells from diverse contexts assemble apical specializations to serve tissue-specific functions. In virtually all cases, these features consist of arrays of microvilli: micron-scale, actin bundle-supported protrusions that mediate biochemical and physical interactions with the external environment. Despite their importance for epithelial physiology, how microvilli grow during cellular differentiation remains poorly understood. Using genetic and small molecule perturbations, we found that an epithelial cell's potential for growing microvilli of normal size is limited by an adjacent actin-dependent process: apical clathrin-mediated endocytosis. Timelapse imaging of individual microvillar growth events revealed tight spatial and temporal coupling to sites of clathrin-mediated endocytosis. Finally, ultrastructural analysis of undifferentiated epithelial monolayers showed that most nascent microvilli are immediately adjacent to or in contact with an apical endocytic pit. Based on these discoveries, we conclude that clathrin-mediated endocytosis controls the timing and location of microvillar growth, as well as the dimensions of the resulting protrusions.**

____

INTRODUCTION

Actin-based membrane protrusions extend from the surface of virtually all animal cells, where they mediate biochemical and physical interactions with the extracellular environment. Microvilli are canonical examples; as finger-like protrusions that are found in large arrays on the apical/luminal surfaces of solute transporting epithelial tissues, they significantly increase the plasma membrane surface area available for channels and transporters that are central to physiological function[1-4]. Variations on this morphological theme are found throughout the animal kingdom, on organisms ranging from ancient single-celled protists to mammals[5-7]. Interestingly, previous evolutionary studies suggested that as mammalian cell functions became more specialized, building arrays of microvilli enabled epithelial cells to increase surface area for maximizing solute uptake potential[4,6]. In specific tissues, such as the intestinal tract and kidney tubules, epithelial cells build highly ordered arrays of microvilli known as brush borders, where the apical surface is nearly saturated with protrusions, characterized by hexagonal spacing when viewed _en face_[8].

The ultrastructure and proteome of intestinal microvilli have been well characterized in studies spanning decades[9]. Driven by an initiation complex containing epidermal growth factor pathway substrate 8 (EPS8) and insulin-receptor tyrosine kinase substrate (IRTKS), microvilli first appear stochastically on the apical surface of epithelial cells during the early stages of differentiation, subsequently undergoing rapid population expansion and packing to form full brush borders in mature cells[10-14]. Interestingly, microvillus structure is highly stereotyped: each protrusion is supported by 20-30 actin filaments bundled in parallel such that their fast-growing barbed ends are oriented towards the distal tips[8,15,16]. Parallel filament bundling is supported by fimbrin (also known as plastin-1), villin, espin, and mitotic spindle positioning protein (MISP)[17-20]. Recent studies revealed that these factors occupy distinct regions along the length of the core bundle, suggesting a potential mechanism for regional tuning of physical properties[20].The core actin bundle is connected to the encapsulating membrane through membrane-cytoskeleton linkers, including ezrin and myosin-1a, which maintain membrane wrapping and increase long-term protrusion stability[21,22]. Despite

significant progress toward defining the structure and composition of microvilli, mechanisms that control the
growth of these protrusions remain poorly understood.
Microvillus morphology is tightly regulated by a balance of force-generating cytoskeletal elements
and the force-resisting physical properties of the plasma membrane[2, 11, 23, 24]. Endocytosis, the process by
which cells internalize membrane and extracellular materials, is well positioned to impact this force balance.
Indeed, the actin cytoskeleton plays a crucial role in apical clathrin-mediated endocytosis (CME) in epithelial
cells, as actin filament assembly at endocytic patches generates the necessary inward force to deform the
membrane[25-27]. The series of molecular events that drive CME is well-studied; local synthesis of
phosphatidylinositol 4,5-bisphosphate ($PI_{4,5}P_2$) at clathrin coated pits (CCPs) facilitates recruitment of actin-
binding proteins including neuronal-Wiscott-Aldrich syndrome protein (N-WASp) and actin related protein
2/3 complex (Arp2/3), capping protein, and actin-bundling and severing proteins such as cofilin. These
proteins work together to promote the polymerization and turnover of actin networks to drive the formation
and scission of CCPs[26, 28-35]. Intriguingly, CCPs and microvilli share several molecular components such as
actin-binding and adaptor molecules, suggesting a potential mechanistic link between these functionally
and topologically distinct apical processes. For example, in solute transporting epithelial cells, $PI_{4,5}P_2$ is
enriched in the apical membrane and has been shown to recruit and activate ezrin, a key membrane-
cytoskeletal linker in the brush border[36]. Actin-bundling proteins like fimbrin, necessary for microvillar
formation and stabilization of the microvillar core actin bundle, are also recruited to endocytic patches[18, 37-
39]. Mammalian actin binding protein 1 (mAbp1), with binding domains for both actin and clathrin, acts as a
molecular bridge by physically linking the actin cytoskeleton to endocytic machinery[40]. Notably, mAbp1 has
also been shown to bind two molecules implicated in building microvilli pacsin2 (also known as syndapin2)
and cordon bleu (COBL)[41]. Previous studies also established that knockout of pacsin2 in mice resulted in
significantly shorter and fewer microvilli with reduced membrane wrapping on the surface of mature villus
enterocytes[42]. Notably, endocytic intermediates were found to be stalled between microvilli on the apical
surface, suggesting a defect in endocytic activity. Although the mechanisms underlying these phenotypes
remain unclear, when combined with the molecular links alluded to above, these studies collectively suggest
that endocytic activity at the apical surface may be needed for normal microvillar growth.
Does clathrin-mediated endocytic activity control the growth of microvilli? To address this question,
we used a combination of genetic and small molecule perturbations coupled with live confocal and
ultrastructural imaging to examine microvillar dynamics and structure on the apical surface of brush border-
building epithelial cell lines. Perturbation of CME in actively differentiating cells led to dramatic remodeling
of microvilli; activation of CME depleted microvilli, whereas inhibition resulted in exaggerated protrusions
containing more actin filaments and actin-binding proteins. Interestingly, volumetric live-cell imaging
revealed that microvilli grow directly from CCPs, suggesting that endocytic events mark future growth sites.
Scanning electron microscopy (SEM) also revealed a tight spatial association between apical surface pits
and microvilli in both kidney- and intestine-derived transporting epithelial cells. Together, these data
suggest that, across multiple epithelial model systems, CME plays a direct role controlling the timing and
location of microvillar growth as well as the dimensions of resulting protrusions. These findings redefine our
understanding of microvillar biogenesis and hold broad implications for deciphering the impact of
endocytosis on cellular morphology across diverse epithelial and non-epithelial contexts.
92
RESULTS
Endocytic markers localize to the apical surface of actively differentiating epithelial
cells
While previous studies established that the endocytic machinery localizes to the apical surface of
functionally mature transporting epithelial cells, whether endocytosis plays a role in microvillar growth during
epithelial differentiation remains unexplored [42-45]. We first turned to SEM to examine the surface of cells
lining intestinal crypts, the site of microvillar growth in intestinal tissues, where nutrient transporting
enterocytes comprise ~95% of the epithelium. The intestinal crypt-villus axis offers a unique opportunity for
imaging epithelial cells across all stages of differentiation; crypts house the transit amplifying cells (actively
differentiating) while the villus is populated by mature epithelial cells (fully differentiated) [1, 2, 46]. SEM of
crypts in fractured samples of mouse small intestine revealed a gradual increase in microvillar density on
the apical surface, suggesting that the brush border continuously packs with protrusions as cells migrate
from the base of the crypt to the crypt-villus transition (Fig. 1a,b). We confirmed this point using
immunofluorescent staining to probe for the microvillar-specific actin-bundler, villin, as a proxy for microvillar

density. Consistent with the SEM data, confocal images revealed the apical villin signal increased nearly 2-
fold from crypt base (magenta) to crypt-villus transition (green) (Fig. 1c,d), indicative of microvillar
accumulation on the surface of these cells. We next stained for dynamin2 and pacsin2 to assess the
abundance of these canonical endocytic markers in the crypts. The resulting confocal images revealed that,
similar to villin levels, both markers demonstrated a gradual increase in signal, from the crypt base
(magenta) to crypt-villus transition (Fig. 1c,d). Collectively, these data suggest that the accumulation of
microvilli at the apex of actively differentiating cells spatially and temporally coincides with an increase in
endocytic marker enrichment at the apical surface, before cells are exposed to the contents of the lumen.
115

Clathrin-mediated endocytic activity is higher early in epithelial differentiation

While the localization of endocytic markers at the apical surface suggests increased endocytic activity, we
sought to confirm this using a functional assay for endocytic uptake in a cell culture model that would be
more amenable to perturbation studies than native intestinal epithelia[47]. The solute transporting LLC-PK1-
CL4 (CL4) cell line provides a powerful model for studying apical morphogenesis due to their quick
differentiation time course (3-4 days) in culture[48-50]. We used fluorescent transferrin to measure clathrin-
mediated endocytic activity in differentiating CL4 cells, which exhibit robust microvillar growth at 0 days
post confluence (0-DPC), and fully differentiated CL4 cells with established brush borders at 3 days post
confluence (3-DPC). After incubating CL4 monolayers with transferrin for 15 minutes, confocal imaging
revealed robust transferrin uptake by 0-DPC CL4 cells, although this activity was significantly reduced by
3-DPC (Fig. 2a,b). The reduction in CME activity at later time points was also observed using the human
intestinal epithelial cell line, CACO-2$_{BBE}$ (Extended Data Fig. 1). These findings suggest that CME activity
decreases coincident with the reduction of microvillar growth that takes place later in differentiation.
129

Induction of clathrin-mediated endocytosis depletes microvilli

After confirming robust clathrin-mediated endocytic activity at the apical surface early in differentiation, we
sought to determine if perturbing CME impacted microvillar growth. Here we employed the previously
established "hook and anchor" tool to induce clathrin-mediated endocytosis[51]. In this system, FKBP-tagged
β2 subunit of AP-2 (β2) is recruited to an FRB-tagged CD8 membrane anchor upon addition of rapalog,
resulting in robust formation and internalization of CCPs in a temporally controlled manner (Fig. 2c,d). We
expressed this system in undifferentiated CL4 cells along with Halo-tagged espin, a microvillar actin
bundling protein that specifically labels microvilli[19]. Following endocytic induction, we performed live
spinning disk confocal (SDC) imaging to monitor the impact on microvillar growth and structure (Fig. 2e).
To measure the impact on core actin bundle structure, we generated linescans orthogonal to individual
protrusions visualized in the Halo-espin channel, and calculated areas under the resulting intensity curves
(AUC); the resulting metric is expected to reflect the diameter of the core bundle and the number of
constituent actin filaments (Fig. 2f). Performing this analysis on cells before and 45 min after rapalog
treatment revealed that inducing CME resulted in a ~50% reduction in Halo-espin signal (Fig. 2g),
suggesting that induction of CME negatively impacts microvillar growth and stability.
145

Inhibition of clathrin-meditated endocytosis leads to the formation of exaggerated microvilli

We next sought to determine the impact of endocytic inhibition on microvillar assembly. For these studies,
we used well established inhibitors that disrupt events within the CME pathway and employed SDC
microscopy to visualize protrusion growth and dynamics on the surface of CL4 cells expressing mCherry-
espin[19, 52]. We inhibited CME by targeting both the initiation and scission steps of the pathway through
chlorpromazine (CPZ) and dyngo, respectively. CPZ disrupts clathrin lattices on the plasma membrane and
targets AP-2 to endosomes, while dyngo inhibits dynamin2, a GTPase critical for vesical scission during
the final step of CCP invagination[53-55] (Fig. 3a). CPZ treatment of subconfluent CL4 cells resulted in
dramatic exaggeration of microvillar structure (Supplementary Movie 1); following 45 minutes of CPZ
treatment, protrusions demonstrated a ~1.5-fold increase in AUC measured from the espin signal and ~2-
fold increase in length (Fig. 3b-d and Supplementary Movie 1). This effect was titrated across a range of
CPZ concentrations in CL4 cells and resulted in a similar phenotype in CACO-2$_{BBE}$ cells treated with CPZ
(Extended Data Fig. 2). In contrast, 45 minutes of dyngo treatment did not significantly impact microvillar
morphology (Fig. 3b-d). To ensure that the observed phenotypic differences were not due to disparities in

drug efficacy, we performed a transferrin uptake assay to assess CME activity following 45 minutes of
treatment with either CPZ or dyngo. As expected, both endocytosis inhibitors significantly reduced
transferrin uptake (Extended Data Fig. 3). Thus, blocking endocytosis early in the pathway (Fig. 3a)
promotes the growth of larger microvilli.
The increase in mCherry-espin signal in microvilli following CPZ treatment (Fig. 3b-d) suggests that
core bundles incorporate more actin filaments during growth in the absence of normal endocytic activity.
To determine if this is the case, CL4 cells expressing mNeon-Green-ß-actin were treated with CPZ as
described above (Fig. 3e). Consistent with the mCherry-espin readout, we observed a ~2-fold increase in
AUC, indicating that CPZ treatment indeed leads to higher levels of actin in exaggerated microvilli post-
CPZ treatment (Fig. 3f). Labeling CPZ-treated cells with CellMask-DeepRed (CellMask), a plasma
membrane dye, further revealed that exaggerated microvilli on these cells retained membrane wrapping[56,]
[57], with the length of the wrapped region increasing with the longer core bundles formed under these
conditions (Fig. 4a,b). We also examined the localization of ezrin one of the major membrane-actin linkers
in these protrusions[58]. AUC analysis of EGFP-ezrin signal in CPZ-treated cells further confirmed the
increased diameter of exaggerated microvilli (Fig. 4c,d). Finally, we examined the impact of endocytic
inhibition on factors that reside at the distal tips of microvilli, such as EPS8 and IRTKS[13]. Distal tip
fluorescence intensity of EGFP-EPS8 or EGFP-IRTKS, quantified using Trackmate, revealed an
enrichment of both EPS8 and IRTKS after CPZ treatment (Fig. 4e-h). Collectively, these data indicate that,
under normal conditions, CME plays a critical role in limiting microvillar dimensions and actin content.
180
Exaggerated microvilli that form following CPZ treatment exhibit perturbed motility
Microvilli extending from differentiating epithelial cells exhibit active motility across the apical surface, which
is driven by the treadmilling of the supporting core actin bundles[48]. Given their exaggerated dimensions and
increased filament content following CPZ treatment, we sought to determine if the exaggerated microvilli
exhibit abnormal motility. To address this question, we employed live SDC imaging to track the motility of
individual microvilli on CL4 cells co-expressing mCherry-espin and EGFP-EPS8 (not shown) (Fig. 5a).
Mean square displacement analysis of 50 randomly selected microvillar trajectories per condition revealed
parabolic curves consistent with a mixture of robust active motion and limited diffusion (Fig. 5d). Further,
while we observed no significant changes in velocity nor track displacement after endocytosis inhibition,
exaggerated microvilli did exhibit higher persistence of motion, which was reflected in the extended reach
of individual trajectories when viewed in rose plots (Fig. 5b,c,e-g). Collectively, these data suggest that
while the increased actin density in microvilli does not alter overall speed or displacement, it promotes more
persistent active motility. Based on these findings, we conclude that endocytic activity controls not only
microvillar architecture, but also the dynamics of these structures.
195
Global reduction of plasma membrane tension does not lead to the formation of
exaggerated microvilli
Endocytic activity plays a global role in decreasing plasma membrane surface area and thus, increasing
tension in this bilayer[59, 60]. Moreover, previous studies on CPZ suggest this compound intercalates into the
plasma membrane, which is expected to reduce membrane tension[61, 62]. To determine if global
perturbations in membrane tension are responsible for the growth of exaggerated microvilli post-CPZ
treatment, we treated CL4 cells with other compounds that are known to reduce membrane tension
including 400 µM deoxycholate, 1% ethanol, and addition of 125 mM NaCl; all of these treatments decrease
membrane tension as measured in tether force assays[63-65]. However, none of these global perturbations
induced the formation of microvilli with exaggerated dimensions (Extended Data Fig. 4). These results
suggest that endocytic events might instead locally regulate the growth and dimensions of microvilli.
207
Microvilli grow from apical clathrin-coated pits
We next sought to test the hypothesis that endocytic sites are spatially and temporally linked to microvillar
growth events on the apical surface. To this end, we leveraged a recently developed SDC-based live
imaging approach which allows for the visualization of individual microvillar growth events on the surface
of sub-confluent CL4 cells[12]. To screen for potential spatiotemporal links between microvillar growth and
CME, we imaged mCherry-espin expressing CL4s co-expressing a range of EGFP-tagged endocytic
components including pacsin2, clathrin light chain A (LCA), dynamin2, mAbp1, and PI$_{4,5}$P$_2$ (Fig. 6a,c,e and

Extended Data Fig. 5)[28, 40, 66, 67]. To quantify the localization of endocytic markers at sites of microvillar
growth, we analyzed the maximum intensity projected fluorescence signals of CME marker puncta that
colocalized with elongating espin signals (representing the growth of new microvillar core bundles) over
time. Strikingly, most endocytic markers we examined localized to the apical surface at future sites of
microvillar growth ~2-3 minutes prior to espin accumulation and elongation (Fig. 6b,d,f). One exception to
this pattern was dynamin2, which briefly localized ~1 minute after espin accumulation began, presumably
marking the disassembly of the CCP (Extended Data Fig. 5c). Following their localization to sites of
microvillar growth, most endocytic components disassembled within a few minutes, a timeframe consistent
with previously reported dynamics of CME[68, 69]. Together, these timelapse recordings suggest that the early
steps of CCP formation create a specialized domain at the apical plasma membrane that is permissive for
microvillar growth.
226
Ultrastructural imaging reveals juxtaposition of nascent microvilli and endocytic
pits
To further examine the relationship between CCP formation and the growth of new microvilli, we employed
SEM to visualize the fine structural details of the apical surface of undifferentiated CL4 cells. Strikingly,
these data showed that 51.1% of the pits that we could visualize were in close proximity to microvilli (<250
nm) (Fig. 7a,c)). To determine if the observed spacing was closer than expected based on randomly
distributed pits and microvilli, we performed a nearest neighbor analysis, measuring from the center of every
pit to the center of the base of the nearest microvillus. We also simulated cell surfaces and nearest neighbor
measurements with the same number of pits and microvilli randomly distributed over the same area
(Extended Data Fig. 6). Remarkably, experimental nearest neighbor measurements exhibited a skewed
distribution with a peak at ~200 nm; in contrast, simulated random nearest neighbor measurements
demonstrated a Gaussian distribution with a wider peak at ~450 nm (Fig. 7c). Thus, the observed spacing
between endocytic pits and microvilli cannot be explained by random co-occurrence and is instead
consistent with two processes that are linked in space and time. We also noted that microvilli associated
with endocytic pits were significantly shorter than microvilli further away, suggesting that core bundles
mature as they as they disassociate from the CCPs (Fig. 7d). Expanding the significance of this finding,
we also observed the same close juxtaposition of endocytic pits and microvilli on the surface of CACO2$_{BBE}$
cells (67.1% of pits associated with microvillus) (Fig. 7b,e). These ultrastructural findings provide a
compelling complement to our timelapse data, and strongly indicate that CME and microvillar biogenesis
are spatially and temporally coordinated processes at the apical cell surface.
247
DISCUSSION
Despite the extensive characterization of microvillus architecture and composition in studies spanning more
than 75 years, our understanding of how cells control the growth and resulting dimensions of these
protrusions remains limited. Here, we establish apical endocytic activity as a critical regulator of the location,
timing, and dimensions of nascent microvilli. Given the overlapping molecular machinery and the role of the
actin cytoskeleton and membrane bending in both apical CME and microvilli, we sought to test the
hypothesis that microvillar growth is linked to or, in some way, informed by apical endocytic activity. Indeed,
our analysis of endocytic markers (pacsin2 and dynamin2) in native intestinal crypts revealed that
differentiating cells experiencing rapid microvillar growth also exhibit robust apical localization of endocytic
machinery. Consistent with this, CL4 and CACO-2$_{BBE}$ cells, two well-established cell culture models for
solute transporting epithelia, revealed elevated transferrin uptake in undifferentiated cells relative to their
differentiated counterparts. Together, these findings revealed that apical endocytic activity is well positioned
to influence microvillar assembly early in the time course of epithelial differentiation.
261
How might CME impact microvillar growth on the apical surface of differentiating epithelial cells? Previous
studies using genetic manipulation of microvillar components or pharmacological inhibition of actin
polymerization collectively indicated that experimental depletion of microvilli is difficult, underscoring the
robustness of this evolutionarily ancient process[48, 70-73]. However, we found that inducing high levels of CME
through the recruitment of the b2 subunit of AP2 to the plasma membrane led to a striking depletion of
microvillar actin content across the cell surface. Our results echo those from previous studies which showed
that overexpression of clathrin heavy chain perturbed the morphology of microvilli on the surface of Madin-
Darby Canine Kidney (MDCK) cells, as assessed using transmission electron microscopy[25]. A superficial

interpretation of these findings is that apical CME influences microvillar growth by impacting apical
membrane tension. Yet, in the current study, global perturbations of membrane tension using orthogonal
methods failed to disrupt microvillar architecture. An appealing alternative model is that apical CCPs and
microvillar assembly mechanisms share resources, such as specific protein machinery, membrane lipids,
or actin filaments, and over- or underutilization of these resources by CME compromises the growth of
protrusions from the apical surface. We elaborate more on the experimental support for this idea below.
276
If CCPs and microvilli share resources during their assembly, then inhibition of CME would be expected to
divert those components into the microvillar growth pathway, and potentially result in more protrusions
and/or larger structures. Indeed, when we inhibited CME using CPZ, a compound that inhibits clathrin
recycling to the plasma membrane, new microvilli grew with exaggerated dimensions, appearing longer and
with increased diameter, suggesting they contained more actin filaments. Protrusions assembled following
CPZ treatment retained key structural features of canonical microvilli, including ezrin-mediated membrane
wrapping, and distal tip localization of EPS8 and IRTKS. Additionally, these factors increased in a manner
proportional to the size of the microvillus after endocytosis inhibition, suggesting that stoichiometric binding
was retained in the exaggerated structures. Interestingly, another endocytic inhibitor, dyngo, did not have
the same effect, although this may be explained by the distinct mechanisms by which CPZ and dyngo block
CME (Fig 3a)[53-55]. CPZ, prevents the formation of coated pits by disrupting AP-2 recruitment to the plasma
membrane. In contrast, dyngo treatment allows for pit formation and leads to stabilization of pits on the
plasma membrane by inhibiting vesicle scission. The stalled endocytic intermediates created by dyngo
might still sequester resources such as actin filaments, as suggested in previous studies[74, 75], potentially
explaining why dyngo treatment does not lead to microvillar overgrowth.
292
The apical surface phenotypes associated with gain and loss of CME function – depletion and exaggeration
of microvilli, respectively - indicate that CCP formation and microvillar growth are mechanistically linked,
potentially through a shared requirement for a molecular resource such as actin. This link could operate
globally with these pathways engaging in long-range competition for the same cytoplasmic actin resources,
as suggested in previous studies on actin allocation[76]. Alternatively, this link could operate locally, with both
pathways sharing resources at the same time *and* in the same location at the apical membrane. This latter
possibility is supported by our live imaging studies of epithelial cells early in differentiation. Using SDC
microscopy, we defined a time course for the recruitment of endocytic machinery, which appeared in
diffraction limited puncta on the apical surface in events that lasted several minutes as expected[69, 77].
Remarkably, these puncta also marked future sites of microvillar growth, with some CME markers arriving
2-3 minutes in advance of microvillar growth initiation. All CME markers examined in this assay
demonstrated significant temporal and spatial overlap with the microvillar core bundle marker, espin,
suggesting that resource sharing could occur at discrete sites that are scaled at or below the diffraction limit
(< 200 nm). Interestingly, we noted that not all CCPs progressed to form microvilli. Given the variation in
composition of CCPs, it stands to reason that only a specific subset of pits might be equipped to support
microvillar growth[78-80]. What distinguishes these pits from their counterparts? One intriguing possibility lies
in the mechanism of pit closure. Studies have identified three distinct closure types for CME: symmetric,
asymmetric, and undefined, with asymmetric being the most prevalent (accounting for roughly 70% of pits)
[75, 81]. During asymmetric closure, membrane bulges outward on one side of the CCP propelled by actin
polymerization, progressively covering and closing the pit. This contrasts with symmetric closure (~20% of
events), where the pit slowly closes in an isotropic manner[75, 81]. The specific mode of pit closure and/or the
presence of asymmetric F-actin accumulation, might dictate whether that pit ultimately leads to microvillus
assembly[75, 82].
316
We also surveyed the apical surfaces of fixed epithelial cell monolayers using SEM, with the goal of
examining large numbers of endocytic pits and characterizing their spatial relationship with microvilli.
Strikingly, analysis of undifferentiated CL4 and CACO-2$_{BBE}$ monolayers revealed that >50% of pits were
within 250 nm of a microvillus (CL4: 51.1%, CACO-2 $_{BBE}$: 67.1%), a distance comparable to the diameter of
the diffraction limited puncta that we observed in our live confocal imaging studies. Mathematical
simulations revealed that such close juxtaposition of endocytic pits and microvilli could not be explained by
the random positioning of these objects across the apical surface. We also found that pit-associated
microvilli were significantly shorter relative to more distal protrusions. This observation could indicate that
pit-associated microvilli represent younger growth intermediates, whereas longer and more distal microvilli

could represent fully formed protrusions at steady-state length. In combination with the CME gain and loss
of function studies and the timelapse observations described above, these data lead us to propose a model
whereby microvilli form at sites that are immediately adjacent to (within 250 nm) pre-existing CCPs (Fig.
6f).
330
Why do CCPs serve as preferential sites for microvillar assembly? From several perspectives, the local
biochemical and physical landscape generated during endocytic pit formation appears well-suited for
supporting protrusion growth. For example, CCPs contain a significant F-actin coat generated by the Arp2/3
branched nucleation complex, and the protein machinery that drives microvillar growth could harness those
polymers, elongating and bundling them to form core actin bundles. Interestingly, molecular counting
studies revealed that ~200 Arp2/3 complexes localize to CCPs, and simulations suggest that this population
would be sufficient to generate ~150 filaments for driving pit internalization[83]. Based on ultrastructural
studies of the enterocyte brush border, we know that microvillar core actin bundles contain only ~25
filaments, so the scale of the CCP-associated actin filament network is large enough to support the growth
of potentially a few microvilli[8, 24]. The general concept of co-opting filaments from an existing network to
form a new structure with distinct architecture is similar to the 'convergent elongation' model that was
originally proposed by Svitkina and colleagues to explain the formation of filopodia on the surface of
crawling cells[84]. In that scenario, the actin filaments that support filopodia originate deep in the lamellipodial
meshwork, which is generated by Arp2/3 activity. During filopodial growth, a small number of lamellipodial
filaments are protected from capping by barbed end elongation factors (Ena/VASP, mDia) and therefore
allowed to elongate and become crosslinked by fascin to create a mechanically robust core bundle[84]. In the
context of microvillar growth, EPS8 and/or IRTKS might function as elongation factors[12, 13, 85], whereas
espin and other factors (MISP, fimbrin, and villin) could participate in filament bundling[17-20]. Whether a
similar mechanism is operational at the smaller scale of CCPs and leads to formation of microvilli will require
deep ultrastructural investigation using sophisticated cryo preservation and imaging approaches.
351
Interestingly, our SEM images revealed that when an endocytic pit is adjacent to a microvillus, the
stoichiometry is almost always 1:1, i.e. one microvillus per pit. Thus, in addition to the availability of actin
filaments, other constraints probably serve to limit the number of new microvilli that form during these
events. In our studies, we also observed enrichment of $PI_{4,5}P_2$ before microvilli grow, suggesting a potential
role for specific lipids in microvillar biogenesis. Additionally, the local accumulation of specific lipids or
membrane-associated proteins is known to impact membrane curvature and tension[86], and these physical
factors are likely to impact microvillar growth. Future investigations taking advantage of next generation
membrane tension probes might explore this possibility.
360
The link between CME and microvillar growth revealed in this work points to a fascinating evolutionary
adaptation in epithelial biology. While CME is a ubiquitous cellular process shared by all eukaryotes,
microvilli are specialized structures that emerged later in evolution, with their earliest known appearance in
choanoflagellates[6, 87]. The evolution of endocytosis provided a crucial foundation for membrane
maintenance, cell signaling, and nutrient uptake[88, 89]. We propose that, over time, this actin-dependent
process was co-opted to support the formation of actin-based protrusions such as microvilli. If true, this
would represent an elegant example of evolutionary efficiency, where existing processes are repurposed
to meet new functional demands. The model we propose here might also offer an explanation for growth of
more elaborate microvilli-like structures, including the stereocilia that extend from the apical surface of
mechanosensory hair cells found in the cochlear and vestibular systems[90].

 FIGURES

Figure 1. Endocytic proteins are apically localized in the crypts. (A) Scanning electron microscopy (SEM) of native mouse small intestine along full crypt-villus axis. Scale bar = 50 μm. (B) High magnification of the small intestinal crypt (scale bar = 10 μm) with zooms (base, middle, and transition (scale bars = 1 um (solid box zooms) and 500 nm (dashed box zooms)) showing the accumulation of microvilli along the transit amplifying zone. (C) Native mouse small intestine stained for villin, dynamin2, and pacsin2. Panels show single z-plane confocal image of intestinal crypts. Scale bar = 10 μm. (D) Mean brush border intensity measurements for each marker at the base of the crypt (magenta) and at the crypt-villus transition zone (green). n = 20-25 crypts per marker, 3 WT mice. Multiple Mann-Whitney tests; ****p = <0.0001, **p = 0.002.

Figure 2. Induction of clathrin-mediated endocytosis depletes microvilli. (A) Maximum intensity projection (MIP) of F-actin (Alexa Fluor 647 Phalloidin) and transferrin (Alexa Fluor 488 Conjugate) after 15-minute incubation with human purified transferrin in 0 days post confluency (DPC) and 3-DPC LLC-PK1/CL4 (CL4) cells. Scale bar = 10 μm. (B) Superplot quantification of whole cell mean transferrin intensity measurements. Small transparent points represent individual cells, larger opaque points show experimental replicates. n = 3 replicates, 50 cells per replicate. Paired t-test; **p = 0.0036 (C) Cartoons depicting constructs used to induce clathrin-mediated endocytosis. Numbers represent amino acids. (D) Schematic depicting the rapalog-inducible system to dock the β2 subunit of AP-2 on the plasma membrane to induce endocytosis. (E) Confocal MIP of CL4 cell expressing Halo-espin (shown), CD8-FRB (not shown), and GFP-FKBP (not shown) or FKBP-β2-GFP (not shown) before and after 45 min rapalog incubation. Scale bar = 5 μm. (F) Schematic showing area under the curve (AUC) quantification. (G) Fold change in AUC of Halo-espin signal of microvilli post- vs. pre- rapalog induction. Area under the curve (AUC) was calculated as shown in F. 11-12 cells per condition taken from at least three experimental replicates, 120-150 microvilli per condition. Mann-Whitney test; ****p =<0.0001.

A

1. Initiation   2. Invagination   3. Constriction   4. Fission

AP-2
CPZ
Actin   Clathrin
Dynamin2
Dyngo

B

	DMSO	30 µM CPZ	30 µM Dyngo
0 min	Espin	Espin	Espin
45 min + Drug	Espin	Espin	Espin

103 — 700   103 — 700   103 — 450

C

ns
****

Fold Change in Length

DMSO   CPZ   Dyngo

D

ns
****

Fold Change in AUC

DMSO   CPZ   Dyngo

E

	DMSO	30 µM CPZ
0 min	β-Actin	β-Actin
45 min + Drug	β-Actin	β-Actin

130 — 580   105 — 160

F

****

Fold Change in AUC

DMSO   CPZ

Figure 3. Inhibition of endocytosis results in exaggerated microvilli. (A) Schematic depicting chlorpromazine and dyngo mechanisms of action. (B) Confocal maximum intensity projection (MIP) of representative CL4 cells at 0 min and 45 min post DMSO (left), chlorpromazine (CPZ) (middle), and dyngo (right) treatment. Drug was added immediately after 0 min image was captured. Scale bar = 5 μm. (C) Fold change in length of microvilli post- vs. pre- drug treatments. Microvillar length was measured using mCherry-espin signal. n = 13-16 cells from at least three experimental replicates, 150-200 microvilli per condition. Kruskal-Wallis test with multiple comparisons; ****p = <0.0001. (D) Fold change in area under the curve of mCherry-espin signal of microvilli post- vs. pre- drug treatments. 14-16 cells from at least three experimental replicates, 150-200 microvilli per condition. Kruskal-Wallis test with multiple comparisons; ****p = <0.0001. (E) Confocal MIP of representative CL4 cells expressing mCherry-espin (not shown) and mNeonGreen-βActin at 0 min and 45 min post-DMSO or CPZ treatment. Scale bar = 5 μm. (F) Fold change in area under the curve of mNeonGreen-βActin signal of microvilli post- vs. pre- drug treatments. n = 15 cells from at least three experimental replicates, 150-200 microvilli per condition. Mann-Whitney test; ****p = <0.0001.

Figure 4. Microvilli on CPZ-treated cells retain microvillar characteristics. (A) Confocal maximum
intensity projection (MIP) of CL4 cells expressing mCherry-espin stained with CellMask-DeepRed. Cells
were first treated with either DMSO or CPZ, then incubated with CellMask for 15 minutes. Scale bar = 5
µm. (B) Superplot quantification of the length of membrane coverage along the microvillus after DMSO or
CPZ treatment. Small, transparent points represent individual microvilli, larger opaque points show whole
cell averages. (C, E, G) Confocal MIP of CL4 cells expressing mCherry-espin and (C) EGFP-ezrin, (E)
EGFP-EPS8, and (E) EGFP-IRTKS at 0 min and 45 min. Single channel zooms show intensity-coded color
profile. Intensity scales from low (dark blue) to high (yellow). Scale bar = 3 µm. (D) Quantification of ezrin
AUC before and after CPZ treatment. N = 11 cells, 130-160 microvilli per condition. Mann Whitney U test;
***p =0.0002. (F,H) Superplot quantification of (F) Eps8 or (H) IRTKS intensity pre/post CPZ treatment.
Small, transparent data points are individual puncta and larger, opaque data points are individual cells. N
= 13-15 cells, 4000-6000 microvilli per condition. Mann Whitney U test; ***p = 0.0005.

Figure 5. Exaggerated microvilli exhibit active, persistent motility. (A) Representative montage of microvillar motility and elongation before (left) and after (right) CPZ treatment. Single channels show intensity-coded color profile. Intensity scales from low (white) to high (yellow). Column width = 2 μm. (B) (left) Rose plot of 50 microvillar tracks over 10 min (μm units) in cells pre CPZ treatment. B. (right) 16 representative tracks isolated from B. Scale bar = 1 μm. (C) (left) Rose plot of 50 microvillar tracks over 10 min (μm units) in cells post CPZ treatment. C. (right) 16 representative tracks isolated from C. Scale bar = 1 μm. (D) Mean squared displacement analysis of 100 microvillar tracks imaged for 20 minutes every 1 minute. (E-G). Superplot quantification of persistence, displacement, and velocity, respectively in untreated and CPZ treated cell populations. Only tracks lasting exactly 10 minutes were analyzed. Small transparent points show individual microvilli, larger opaque points show per cell averages. n = 10-12 cells sampled from at least three independent experiments, 60-100 microvilli per cell. Mann Whitney test; P* = 0.0439.

480
Figure 6. Microvilli grow from apical clathrin coated pits. (A) Montage of a single microvillus growth
event in CL4 cells expressing EGFP-pPLC-ΔPH-GFP (PIP$_{4,5}$P$_2$ marker) and mCherry-espin. Every column
represents 30 seconds. Arrows mark the analyzed core bundle. Column width = 2.09 μm. (B) Average
fluorescence intensity measurements over time. n = 20 growth events from 10 cells across 3 independent
experiments. (C) Montage of a single microvillus growth event in CL4 cells expressing EGFP-pacsin2 and
mCherry-espin. Every column represents one minute. Arrows mark the analyzed core bundle. Column width
= 2.09 μm. (D) Average fluorescence intensity measurements over time. n = 20 growth events from 10
cells across 3 independent experiments. (C) Montage of a single microvillus growth event in CL4 cells
expressing EGFP-LCA and mCherry-espin. Every column represents 30 seconds. Arrows mark the
analyzed core bundle. Column width = 2.09 μm. (D) Average fluorescence intensity measurements over
time. n = 21 growth events from 8 cells across 3 independent experiments.

Figure 7. Nascent microvilli are closely associated with endocytic pits across the cell surface. (A) Scanning electron microscopy image of 0-DPC CL4 cells. Scale bar = 10 µm. Zooms 1 and 2 noted by bounding boxes. Zoom scale bar = 2 µm. (B) Scanning electron microscopy image of 0-DPC CACO-2$_{BBE}$ cells. Scale bar = 2 µm. Zooms 1 and 2 noted by bounding box. Zoom scale bar = 1 µm. (C) Nearest neighbor distance measurements of from microvilli to closest endocytic pits on the surface of CL4 cells. Blue line denotes simulated nearest neighbor distances assuming random distribution of the same number of microvilli and endocytic pits across the same cell surface area. (D). Length of microvilli associated (<250 nm) vs not associated (>300 nm) with endocytic pits on the surface of CL4 cells. N = 175 microvilli associated with pits and 95 microvilli not associated with pits. Mann Whitney test P**** = <0.0001. (E) Nearest neighbor distance measurements of from microvilli to closest endocytic pits on the surface of CACO-2$_{BBE}$ cells. Blue line denotes simulated nearest neighbor distances assuming random distribution of same number of microvilli and endocytic pits across the same cell surface area. (F) Schematic depicting proposed model of clathrin mediated endocytosis and microvillar biogenesis sharing materials during epithelial cell differentiation.

EXTENDED DATA

Supplemental Movie 1, related to Figure 3. Inhibition of endocytosis results in exaggerated microvilli. Live cell movie of maximum intensity projection (MIP) confocal time series of 3 CL4 cells expressing mCherry-espin. CPZ was added at T = 30 min. 6.8 μm MIPs are composed of 17 x 0.4 μm confocal slices. Images were captured at 1 frame per 1 minute. Time hour: min: sec. Single channels show intensity-coded color profile. Intensity scales from low (white) to high (yellow). Scale bar = 5 μm.

Extended Data Figure 1. Clathrin-mediated endocytic activity decreases as epithelial cells differentiate. (A) Maximum intensity projection (MIP) of F-actin (Alexa Fluor 647 Phalloidin) and transferrin (Alexa Fluor 488 Conjugate) after 15-minute incubation with human purified transferrin in 0 days post confluency (DPC) and 14-DPC CACO$_2$-BBE cells. Scale bar = 10 μm. (B) Superplot quantification of whole cell mean transferrin intensity measurements. Small transparent points represent individual cells, larger opaque points show experimental replicates. n = 3 replicates, 50 cells per replicate. Paired t-test; p= 0.0886.

Extended Data Figure 2. **CPZ treatment phenotype is titratable and robust across model systems.**
(A) Confocal maximum intensity projection (MIP) of representative CL4 cells at 0 min and 45 min after
varying concentrations of CPZ treatment. Drug was added immediately after 0 min image was captured.
Single channels show intensity-coded color profile. Intensity scales from low (white) to high (yellow). Scale
bar = 5 µm. (B) Fold change in area under the curve of mCherry-espin signal of microvilli post- vs. pre- CPZ
treatment. 8-10 cells from at least three experimental replicates, 100-150 microvilli per condition. Images
and data for the 30 µM CPZ treatment were taken from figure 3. (C) MIPs of 3 representative CACO$_2$-BBE
cells at 0 min and 45 min after treatment with 30 µM CPZ. Drug was added immediately after 0 min image
was captured. Single channels show intensity-coded color profile. Intensity scales from low (white) to high
(yellow). Scale bar = 5 µm.

559 **Extended Data Figure 3**. **CPZ and dyngo equally inhibit transferrin uptake.** (A) Maximum intensity
560 projection (MIP) of CL4 cells transfected with mCherry-espin incubated with human purified transferrin after
561 45-minute treatment of DMSO, 30 µM CPZ, or 30 µM Dyngo. Scale bar = 10 µM (B) Superplot quantification
562 of whole cell mean transferrin intensity measurements. Small transparent points represent individual cells,
563 larger opaque points show experimental replicates. n = 100 cells sampled from two experimental replicates.

582
583
Extended Data Figure 4. Global reduction in plasma membrane tension does not lead to the
formation of exaggerated microvilli. (A) Confocal maximum intensity projection (MIP) of representative
CL4 cells before and after treatment with CPZ, 1% EtOH, 400 μM Deoxycholate, and 125 mM NaCl. Cells
were treated with CPZ and Deoxycholate for 45 minutes, and with 1% EtOH and 125 mM NaCl for 5-10
minutes. Drug was added immediately after 0 min image was captured. Single channels show intensity-
coded color profile. Intensity scales from low (white) to high (yellow). Scale bar = 5 μm. (B) Fold change in
area under the curve of mCherry-espin signal of microvilli post- vs. pre- drug treatments. 8-10 cells from 2
experimental replicates, 100-150 microvilli per condition. Images and data for the 30 μM CPZ treatment
were taken from figure 3.

610
Extended Data Figure 5. Microvilli grow from apical clathrin coated pits. (A) Montage of a single
microvillus growth event in CL4 cells expressing EGFP-mAbp1 and mCherry-espin. Every column
represents 30 seconds. Arrows mark the analyzed core bundle. Column width = 2.09 μm. (B) Average
fluorescence intensity measurements over time. n = 9 growth events from 4 cells across 2 independent
experiments. (C) Montage of a single microvillus growth event in CL4 cells expressing mCherry-dynamin2
and GFP-espin. Every column represents one minute. Arrows mark the analyzed core bundle. Column
width = 2.09 μm. (D) Average fluorescence intensity measurements over time. n = 10 growth events from
4 cells across 3 independent experiments.

Extended Data Figure 6. Simulation of randomly distributed endocytic pits and microvilli leads to nearest neighbor distances with gaussian distributions. (A) 10 runs of simulated nearest neighbor distance measurements from endocytic pits to nearest microvillus on the surface of CL4 cells. Simulated densities were matched with experimental measurements of 516 pits, 2570 microvilli over 1601.25 μm^2. (B) 10 runs of simulated nearest neighbor distance measurements from endocytic pits to nearest microvillus on the surface of CACO$_2$-BBE cells. Simulated densities were matched with experimental measurements of 415 pits, 2039 microvilli over 1828.7 μm^2.

METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to the lead contact, Matthew J. Tyska (matthew.tyska@vanderbilt.edu).

Materials availability

Plasmids generated in this study will be made available from the lead contact on request.

Data code availability

No large-scale datasets or new code were generated in this study.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cell culture models

LLC-PK1-CL4 (porcine kidney proximal tubule) cells were grown in 1X high glucose DMEM containing 2mM L-glutamine (Corning #10-0130CV) supplemented with 10% fetal bovine serum (FBS) (R&D Systems) and 1% L-Glutamine (Corning # 25-005-CI). CACO-2$_{BBE}$ cells were grown in 1x high glucose DMEM containing 2mm L-glutamine supplemented with 20% FBS. Cells were maintained in treated plastic culture flasks incubated at 37 °C and 5% CO_2. Cells were tested for mycoplasma monthly using the MycoStrip Mycoplasma Detection Kit (Invivogen #rep-mys-50).

METHOD DETAILS

Cloning and Constructs

All overexpression constructs listed in this paper were previously generated and/or obtained as noted in the key resources table.

Cell line generation

LLC-PK1-CL4 cells stably expressing pmCherry-espin were previously generated using the stable cell line generation protocol as described in Gaeta et al., 2022. Briefly, cells were transfected with Lipofectamine 2000 following manufacturer protocol in a T25 cell culture flask. The next day, cells were split up to a T75 flask and treated with 1 mg/mL G418 sulfate for antibiotic selection. Cells were maintained in culture under constant G418 selection to create the stably expressing cell line.

Transient transfection

LLC-PK1-CL4 cells were seeded in 6 well plates and transfected at ~90% confluency using Lipofectamine2000 with 3 µg DNA. 4 hrs after incubation with the lipid complex, cells were replated on CellVis 35mm 1.5# glass bottom dishes for imaging the following day.

Immunofluorescence staining

Tissue: Tissue sections were deparaffinized and rehydrated through a descending ethanol series and subjected to antigen retrieval in a heated buffer at pH 9. Following cooling, sections were washed and blocked with 10% normal goat serum (NGS) and probed for villin, pacsin2, or dynamin2 in 1% NGS overnight at 4 °C. The following day, sections were washed, incubated at room temperature with secondary antibodies, dehydrated through an ascending ethanol series, and mounted in ProLong Gold mounting medium.

Live imaging microscopy

Nikon Ti2 inverted light microscope equipped with a Yokogawa CSU-X1 spinning disk head, a Photometrics Prime 95B sCMOS camera, and 488 nm, 561 nm, and 647 nm excitation lasers. A 60x/1.49 NA or 100x/1.49 NA TIRF oil immersion objective was used for all experiments. Cells were maintained in a stage top incubator at 37 °C with 5% CO2 (Tokai Hit). Z stacks of 5-8 µm using a 0.2-0.4 µm step size were collected using a triggered NIDAQ piezo Z stage.

Scanning electron microscopy (SEM)

Tissue: 5 mm murine duodenal sections were fixed in 2.5% glutaraldehyde and 4% paraformaldehyde in
SEM, washed, and then embedded in OCT. To ensure stability of the explant lumens, samples were gently
passed through 3 rounds of fresh OCT. Samples were then frozen in cryomolds and stored at -80 °C .
Frozen explants were sectioned, melted onto stainless steel AFM discs, and processed through OsO_4,
ddH_2O, and subsequently dehydrated through graded ethanol series. Detached sections were recovered
and adhered to aluminum SEM specimen stubs. SEM imaging was performed on a Quanta 250
environmental SEM or Zeiss Crossbeam 550. For more detailed methods, see Meenderink et al[48].

Cells: Caco-2$_{BBE}$ cells were plated at 70% confluence on Thermanox coverslips in a 6-well dish. 24 hrs
post-plating, cells were washed 3X with PBS to remove all media and were fixed in a buffer of 2.5%
glutaraldehyde and 0.1 M $CaCl_2$ free cacodylate overnight at room temperature. Coverslips were washed
3X in 0.1 M $CaCl_2$ free cacodylate buffer and incubated in pre-warmed 1% tannic acid in cacodylate.
Following washes with 0.1 M cacodylate, coverslips were incubated for 1 hour in 1% OsO_4. After washes
with ddH_2O, they were incubated for 30 min with 1% uranyl acetate in ddH_2O. Finally, coverslips were
washed with ddH_2O 3X for 5 min to remove uranyl acetate before drying with a series of graded EtOH
washes. After samples were dried using critical point drying and vented for 20 min, coverslips were mounted
on SEM specimen stubs via a conductive adhesive tab and sputter coated prior to imaging. SEM imaging
was performed on a Zeiss Crossbeam 550 with a voltage of 2 keV and an SE2 detector. All reagents were
purchased from Electron Microscopy Sciences.

Transferrin uptake assay
LLC-PK1-CL4 cells expressing mCherry-espin were seeded on acid-washed #1.5 coverslips at ~60%
confluence in a cell culture 6-well treated plate on day 0. The next day, cells were treated with 30 μM CPZ,
30 μM Dyngo, or DMSO and incubated for 45 min. At the 45-min mark, transferrin conjugate was added to
the cells at 25 μg/mL and incubated for 20 min. After 20 min, the cells were fixed in 4% paraformaldehyde
diluted in PBS and incubated at 37 °C for 15 min. Coverslips were subsequently mounted on slides in
Prolong Gold antifade reagent and imaged using spinning disk microscopy.

QUANTIFICATION AND STATISTICAL ANALYSIS:
Statistical testing
722 All statistical testing was performed in GraphPad Prism 9 by running a Mann Whitney test or a paired t-test
723 for pairwise comparisons or Kruskal-Wallis to compare more than two conditions. For all figures, exact n
724 values and their definitions are reported in the figure legends.

Time series analysis of microvillus growth
727 Microvillar growth was analyzed as previously described[12]. Briefly, a bounding ROI was manually drawn
728 around the signal of interest and mean intensity was measured for each timeframe. Mean intensity values
729 were normalized, with the smallest value equal to 0 and the largest equal to one. Normalized values were
730 then averaged and plotted with SEM.

Area under the curve analysis
733 To measure the impact on core actin bundle structure, we generated linescans (5 pixels in width, 1.54 μm
734 in length) orthogonal to individual protrusions visualized in the Halo-EPSN channel. Linescans were drawn
735 to ensure lines only crossed through one protrusion per line. Areas under the resulting intensity curves
736 (AUC) were then calculated in prism, averaged and plotted; the resulting metric is expected to reflect the
737 diameter of the core bundle.

EGFP-EPS8 and EGFP-IRTKS puncta intensities
740 Mean puncta intensities were measured using Nikon Elements' "Spot Detection" function.

Microvilli tracking using EGFP-EPS8 puncta
Raw 3D movies were converted into maximum intensity projections, and the EPS8 channel (488) was
isolated. The isolated channel movie was loaded into Trackmate and puncta were identified using a DoG
detector with an estimated object diameter of 500 nm. Identified spots were thresholded based on
background intensity and track length. Tracks beginning on the first frame or ending on the last frame were
excluded. Track and spot data were then exported to Excel and analyzed and plotted in Prism as radial XY

positions over time by subtracting each position in X or Y from their respective point position at time 0,
making the first position (0,0). Persistence, track displacement, and linear velocity were taken from tracks
lasting exactly 10 min.
751
752 **Mean squared displacement analysis**
753 EPS8 puncta trajectories as a proxy for microvillar motion were analyzed using mean squared displacement
754 (MSD) analysis as previously described[48]. MSD curves were subsequently fit with an active motion model
755 in the form $MSD(nDt) = 4Dt + V^2 * nDt^2$, where D is the diffusion coefficient and V is the velocity of active
756 motion[91, 92]. Curve fitting and sum-of-squares F-tests were performed using PRISM v.9.
757
Nearest neighbor expected distribution simulation
To determine the expected nearest neighbor distance between pits and microvilli, a random population of
microvilli and endocytic pits were simulated within a rectangular area using a custom python script. The
number of objects in each population was set to match the observed experimental densities. Euclidean
distances between each endocytic pit and every microvillus were measured and the subsequent minimum
distance (nearest neighbor) for each endocytic pit was identified. Each simulation (CACO-2$_{BBE}$ and CL4)
was run 10 times and then averaged (Extended Data Fig. 6).

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and virus strains		
E. coli DH5-Alpha competent cells	Molecular Cell Biology Resource Core, Vanderbilt Medical Center	Item# DH5 Alpha
Chemicals, peptides, and recombinant proteins		
G418 Sulfate	Gold Biotechology	Cat# G-418-1
Janelia Fluor635 (JF635)	Janelia	N/A
Lipofectamine 2000	Thermo Fischer Scientific	Cat# 11668019
A/C Heterodimerizer	Takara	Cat# 635057
488 Transferrin	Thermo Fischer Scientific	T13342
CellMask-Deep Red	Thermo Fischer Scientific	C10046
Alexa Fluor 647 Phalloidin	Thermo Fischer Scientific	A22287
16% Paraformaldehyde	Electron Microscopy Sciences	Cat# 15710
ProLong Gold Antifade Reagent	Thermo Scientific	P36930
Triton X-100	Sigma	Cat# T8787
DMSO	Sigma	Cat# D8418
Chlorpromazine	Selleck Chemicals	S5749
Dyngo-4a	Selleck Chemicals	S7163
Deoxycholate		
Glutaraldehyde, 25%	Electron Microscopy Sciences	Cat# 16220
Tannic Acid	Electron Microscopy Sciences	Cat# 21700
Osmium Tetroxide	Electron Microscopy Sciences	Cat# 19112
Tissue-Tek OCT	Sakura Finetek	#4583
Cell Lines		
LLC-PK1-CL4	Gift from Dr. Carolyn Slayman (Yale University)	N/A
Antibodies		
anti-villin (mouse) [1:100]	Santa Cruz Biotechnology	sc-66022
dynamin2 (rabbit) [1:200]	Sigma-Aldrich	HPA054246
pacsin2 (rabbit) [1:200]	Sigma-Aldrich	HPA049854
Recombinant DNA		
mCherry-C1-syndapin2 (mouse)	Addgene	#27681
GFP-ß2 (616-951)-FKBP (human)	Adgene	#59353100729
DNMII-pmCherry-N1 (dynamin2) (human)	Addgene	#27689
pmCherry-espin (rat)	Gift from Dr. James Bartles, NWU	N/A
mNeonGreen-βActin (human)	Tyska Lab	N/A
pEGFP-C1-EPS8 (human)	Tyska Lab	N/A
pEGFP-C1-IRTKS (human)	Tyska Lab	N/A
pEGFP-N1-ezrin (human)	Addgene	#202684
pGFP-C1-espin (rat)	Tyska Lab	N/A
Halo-C1-espin (rat)	Tyska Lab	N/A
pGFP-FKBP	Addgene	#67529
pCD8-mCherry-FRB (human)	Addgene	#100738
pEGFP-C2-mAbp1 (human)	Addgene	#26719
pPLC-ΔPH-GFP	Addgene	#21179
Software and Algorithms		
FIJI		
NIS AR Elements Analysis	Nikon	N/A
Prism 9	Graphpad	N/A

BioRender	BioRender	N/A
Procreate	Savage Interactive	N/A
Other		
Mycostrip™ – Myocplasma Detection Kit	InvivoGen	Cat# rep-mys50
35mm #1.5 glass bottom dishes	CellVis	Cat# D35-20-1.5N

804
805

REFERENCES

1.   Crawley, S.W., Mooseker, M.S. & Tyska, M.J. Shaping the intestinal brush border. *The Journal of cell biology* **207**, 441-451 (2014).
2.   Creamer, B., Shorter, R.G. & Bamforth, J. The turnover and shedding of epithelial cells. I. The turnover in the gastro-intestinal tract. *Gut* **2**, 110-118 (1961).
3.   Helander, H.F. & Fandriks, L. Surface area of the digestive tract - revisited. *Scand J Gastroenterol* **49**, 681-689 (2014).
4.   Sebe-Pedros, A. *et al.* Insights into the origin of metazoan filopodia and microvilli. *Mol Biol Evol* **30**, 2013-2023 (2013).
5.   BSC, K.S.a.L. Cytoskeleton Structure and Composition in Choanoflagellates. *Journal of Eukaryotic Microbiology* **45**, 361-367 (1998).
6.   Pena, J.F. *et al.* Conserved expression of vertebrate microvillar gene homologs in choanocytes of freshwater sponges. *Evodevo* **7**, 13 (2016).
7.   Adl, S.M. *et al.* The revised classification of eukaryotes. *J Eukaryot Microbiol* **59**, 429-493 (2012).
8.   Ohta, K., Higashi, R., Sawaguchi, A. & Nakamura, K. Helical arrangement of filaments in microvillar actin bundles. *J Struct Biol* **177**, 513-519 (2012).
9.   McConnell, R.E., Benesh, A.E., Mao, S., Tabb, D.L. & Tyska, M.J. Proteomic analysis of the enterocyte brush border. *Am J Physiol Gastrointest Liver Physiol* **300**, G914-926 (2011).
10.   Cencer, C.S. *et al.* Adhesion-based capture stabilizes nascent microvilli at epithelial cell junctions. *Dev Cell* **58**, 2048-2062 e2047 (2023).
11.   Crawley, S.W. *et al.* Intestinal brush border assembly driven by protocadherin-based intermicrovillar adhesion. *Cell* **157**, 433-446 (2014).
12.   Gaeta, I.M., Meenderink, L.M., Postema, M.M., Cencer, C.S. & Tyska, M.J. Direct visualization of epithelial microvilli biogenesis. *Curr Biol* **31**, 2561-2575 e2566 (2021).
13.   Postema, M.M., Grega-Larson, N.E., Neininger, A.C. & Tyska, M.J. IRTKS (BAIAP2L1) Elongates Epithelial Microvilli Using EPS8-Dependent and Independent Mechanisms. *Curr Biol* **28**, 2876-2888 e2874 (2018).
14.   Specian, R.D. & Neutra, M.R. The surface topography of the colonic crypt in rabbit and monkey. *Am J Anat* **160**, 461-472 (1981).
15.   Mooseker, M.S. & Tilney, L.G. Organization of an actin filament-membrane complex. Filament polarity and membrane attachment in the microvilli of intestinal epithelial cells. *J Cell Biol* **67**, 725-743 (1975).
16.   Pollard, T.D. & Mooseker, M.S. Direct measurement of actin polymerization rate constants by electron microscopy of actin filaments nucleated by isolated microvillus cores. *J Cell Biol* **88**, 654-659 (1981).
17.   Bretscher, A. & Weber, K. Villin: the major microfilament-associated protein of the intestinal microvillus. *Proceedings of the National Academy of Sciences of the United States of America* **76**, 2321-2325 (1979).

18. Bretscher, A. & Weber, K. Fimbrin, a new microfilament-associated protein present in
microvilli and other cell surface structures. *The Journal of cell biology* **86**, 335-340
(1980).
19. Bartles, J.R., Zheng, L., Li, A., Wierda, A. & Chen, B. Small espin: a third actin-bundling
protein and potential forked protein ortholog in brush border microvilli. *The Journal of cell*
biology **143**, 107-119 (1998).
20. Morales, E.A., Arnaiz, C., Krystofiak, E.S., Zanic, M. & Tyska, M.J. Mitotic Spindle
Positioning (MISP) is an actin bundler that selectively stabilizes the rootlets of epithelial
microvilli. *Cell Rep* **39**, 110692 (2022).
21. Howe, C.L. & Mooseker, M.S. Characterization of the 110-kdalton actin-calmodulin-, and
membrane-binding protein from microvilli of intestinal epithelial cells. *J Cell Biol* **97**, 974-
985 (1983).
22. Berryman, M., Franck, Z. & Bretscher, A. Ezrin is concentrated in the apical microvilli of
a wide variety of epithelial cells whereas moesin is found primarily in endothelial cells. *J*
Cell Sci **105 (Pt 4)**, 1025-1043 (1993).
23. Delacour, D., Salomon, J., Robine, S. & Louvard, D. Plasticity of the brush border - the
yin and yang of intestinal homeostasis. *Nat Rev Gastroenterol Hepatol* **13**, 161-174
(2016).
24. Tilney, L.G. & Cardell, R.R. Factors controlling the reassembly of the microvillous border
of the small intestine of the salamander. *J Cell Biol* **47**, 408-422 (1970).
25. Hyman, T., Shmuel, M. & Altschuler, Y. Actin is required for endocytosis at the apical
surface of Madin-Darby canine kidney cells where ARF6 and clathrin regulate the actin
cytoskeleton. *Mol Biol Cell* **17**, 427-437 (2006).
26. Svitkina, T. The Actin Cytoskeleton and Actin-Based Motility. *Cold Spring Harb Perspect*
Biol **10** (2018).
27. Armstrong, G. & Olson, M.F. Bending over backwards: BAR proteins and the actin
cytoskeleton in mammalian receptor-mediated endocytosis. *Eur J Cell Biol* **101**, 151257
(2022).
28. Antonescu, C.N., Aguet, F., Danuser, G. & Schmid, S.L. Phosphatidylinositol-(4,5)-
bisphosphate regulates clathrin-coated pit initiation, stabilization, and size. *Mol Biol Cell*
22, 2588-2600 (2011).
29. Sept, D. & McCammon, J.A. Thermodynamics and kinetics of actin filament nucleation.
Biophys J **81**, 667-674 (2001).
30. Takenawa, T. Phosphoinositide-binding interface proteins involved in shaping cell
membranes. *Proc Jpn Acad Ser B Phys Biol Sci* **86**, 509-523 (2010).
31. Cullen, P.J., Cozier, G.E., Banting, G. & Mellor, H. Modular phosphoinositide-binding
domains--their role in signalling and membrane trafficking. *Curr Biol* **11**, R882-893
(2001).
32. Jost, M., Simpson, F., Kavran, J.M., Lemmon, M.A. & Schmid, S.L. Phosphatidylinositol-
4,5-bisphosphate is required for endocytic coated vesicle formation. *Curr Biol* **8**, 1399-
1402 (1998).
33. Svitkina, T.M. & Borisy, G.G. Arp2/3 complex and actin depolymerizing factor/cofilin in
dendritic organization and treadmilling of actin filament array in lamellipodia. *J Cell Biol*
145, 1009-1026 (1999).
34. Pavlov, D., Muhlrad, A., Cooper, J., Wear, M. & Reisler, E. Actin filament severing by
cofilin. *J Mol Biol* **365**, 1350-1358 (2007).
35. Chen, Q. & Pollard, T.D. Actin filament severing by cofilin dismantles actin patches and
produces mother filaments for new patches. *Curr Biol* **23**, 1154-1162 (2013).
36. Al-Momany, A., Li, L., Alexander, R.T. & Ballermann, B.J. Clustered PI(4,5)P(2)
accumulation and ezrin phosphorylation in response to CLIC5A. *J Cell Sci* **127**, 5164-
5178 (2014).

37.   Gerke, V. & Weber, K. Isolation and characterization of mammalian villin and fimbrin, the
two bundling proteins of the intestinal microvilli. *Eur J Cell Biol* **31**, 249-255 (1983).
38.   Kubler, E. & Riezman, H. Actin and fimbrin are required for the internalization step of
endocytosis in yeast. *EMBO J* **12**, 2855-2862 (1993).
39.   Skau, C.T. *et al.* Actin filament bundling by fimbrin is important for endocytosis,
cytokinesis, and polarization in fission yeast. *J Biol Chem* **286**, 26964-26977 (2011).
40.   Kessels, M.M., Engqvist-Goldstein, A.E., Drubin, D.G. & Qualmann, B. Mammalian
Abp1, a signal-responsive F-actin-binding protein, links the actin cytoskeleton to
endocytosis via the GTPase dynamin. *J Cell Biol* **153**, 351-366 (2001).
41.   Haag, N. *et al.* The actin nucleator Cobl is crucial for Purkinje cell development and
works in close conjunction with the F-actin binding protein Abp1. *J Neurosci* **32**, 17842-
17856 (2012).
42.   Postema, M.M., Grega-Larson, N.E., Meenderink, L.M. & Tyska, M.J. PACSIN2-
dependent apical endocytosis regulates the morphology of epithelial microvilli. *Mol Biol*
Cell **30**, 2515-2526 (2019).
43.   Danielsen, E.M. Probing endocytosis from the enterocyte brush border using fluorescent
lipophilic dyes: lipid sorting at the apical cell surface. *Histochem Cell Biol* **143**, 545-556
(2015).
44.   Matter, K., Stieger, B., Klumperman, J., Ginsel, L. & Hauri, H.P. Endocytosis, recycling,
and lysosomal delivery of brush border hydrolases in cultured human intestinal epithelial
cells (Caco-2). *J Biol Chem* **265**, 3503-3512 (1990).
45.   Michael Danielsen, E. & Hansen, G.H. Small molecule pinocytosis and clathrin-
dependent endocytosis at the intestinal brush border: Two separate pathways into the
enterocyte. *Biochim Biophys Acta* **1858**, 233-243 (2016).
46.   Krndija, D. *et al.* Active cell migration is critical for steady-state epithelial turnover in the
gut. *Science* **365**, 705-710 (2019).
47.   Gammella, E., Buratti, P., Cairo, G. & Recalcati, S. The transferrin receptor: the cellular
iron gate. *Metallomics* **9**, 1367-1375 (2017).
48.   Meenderink, L.M. *et al.* Actin Dynamics Drive Microvillar Motility and Clustering during
Brush Border Assembly. *Dev Cell* **50**, 545-556 e544 (2019).
49.   Hull, R.N., Cherry, W.R. & Weaver, G.W. The origin and characteristics of a pig kidney
cell strain, LLC-PK. *In Vitro* **12**, 670-677 (1976).
50.   Tyska, M.J. & Mooseker, M.S. MYO1A (brush border myosin I) dynamics in the brush
border of LLC-PK1-CL4 cells. *Biophys J* **82**, 1869-1883 (2002).
51.   Wood, L.A., Larocque, G., Clarke, N.I., Sarkar, S. & Royle, S.J. New tools for "hot-
wiring" clathrin-mediated endocytosis with temporal and spatial precision. *J Cell Biol*
216, 4351-4365 (2017).
52.   Loomis, P.A. *et al.* Espin cross-links cause the elongation of microvillus-type parallel
actin bundles in vivo. *J Cell Biol* **163**, 1045-1055 (2003).
53.   Onaya, T., Solomon, D.H. & Davidson, W.D. Effect of chlorpromazine on thyrotropin-
stimulated endocytosis and glucose oxidation in canine thyroid slices. *Endocrinology* **85**,
150-154 (1969).
54.   Bouley, R., Yui, N., Terlouw, A., Cheung, P.W. & Brown, D. Chlorpromazine Induces
Basolateral Aquaporin-2 Accumulation via F-Actin Depolymerization and Blockade of
Endocytosis in Renal Epithelial Cells. *Cells* **9** (2020).
55.   McCluskey, A. *et al.* Building a better dynasore: the dyngo compounds potently inhibit
dynamin and endocytosis. *Traffic* **14**, 1272-1289 (2013).
56.   Bretscher, A. & Weber, K. Localization of actin and microfilament-associated proteins in
the microvilli and terminal web of the intestinal brush border by immunofluorescence
microscopy. *The Journal of cell biology* **79**, 839-845 (1978).

57.   Cramer, L.P. Organization and polarity of actin filament networks in cells: implications for
948   the mechanism of myosin-based cell motility. *Biochem Soc Symp* **65**, 173-205 (1999).
949   58.   Pelaseyed, T. & Bretscher, A. Regulation of actin-based apical structures on epithelial
950   cells. *J Cell Sci* **131** (2018).
951   59.   Willy, N.M. *et al.* Membrane mechanics govern spatiotemporal heterogeneity of
952   endocytic clathrin coat dynamics. *Mol Biol Cell* **28**, 3480-3488 (2017).
953   60.   Djakbarova, U., Madraki, Y., Chan, E.T. & Kural, C. Dynamic interplay between cell
954   membrane tension and clathrin-mediated endocytosis. *Biol Cell* **113**, 344-373 (2021).
955   61.   Romer, J. & Bickel, M.H. Interactions of chlorpromazine and imipramine with artificial
956   membranes investigated by equilibrium dialysis, dual-wavelength photometry, and
957   fluorimetry. *Biochem Pharmacol* **28**, 799-805 (1979).
958   62.   Jutila, A., Soderlund, T., Pakkanen, A.L., Huttunen, M. & Kinnunen, P.K. Comparison of
959   the effects of clozapine, chlorpromazine, and haloperidol on membrane lateral
960   heterogeneity. *Chem Phys Lipids* **112**, 151-163 (2001).
961   63.   Raucher, D. & Sheetz, M.P. Cell spreading and lamellipodial extension rate is regulated
962   by membrane tension. *J Cell Biol* **148**, 127-136 (2000).
963   64.   Raucher, D. & Sheetz, M.P. Characteristics of a membrane reservoir buffering
964   membrane tension. *Biophys J* **77**, 1992-2002 (1999).
965   65.   Raucher, D. & Sheetz, M.P. Membrane expansion increases endocytosis rate during
966   mitosis. *J Cell Biol* **144**, 497-506 (1999).
967   66.   Qualmann, B. & Kelly, R.B. Syndapin isoforms participate in receptor-mediated
968   endocytosis and actin organization. *J Cell Biol* **148**, 1047-1062 (2000).
969   67.   Itoh, T. *et al.* Dynamin and the actin cytoskeleton cooperatively regulate plasma
970   membrane invagination by BAR and F-BAR proteins. *Dev Cell* **9**, 791-804 (2005).
971   68.   Mettlen, M., Chen, P.H., Srinivasan, S., Danuser, G. & Schmid, S.L. Regulation of
972   Clathrin-Mediated Endocytosis. *Annu Rev Biochem* **87**, 871-896 (2018).
973   69.   Ehrlich, M. *et al.* Endocytosis by random initiation and stabilization of clathrin-coated
974   pits. *Cell* **118**, 591-605 (2004).
975   70.   Ferrary, E. *et al.* In vivo, villin is required for Ca(2+)-dependent F-actin disruption in
976   intestinal brush borders. *J Cell Biol* **146**, 819-830 (1999).
977   71.   Grimm-Gunter, E.M. *et al.* Plastin 1 binds to keratin and is required for terminal web
978   assembly in the intestinal epithelium. *Mol Biol Cell* **20**, 2549-2562 (2009).
979   72.   Pinson, K.I., Dunbar, L., Samuelson, L. & Gumucio, D.L. Targeted disruption of the
980   mouse villin gene does not impair the morphogenesis of microvilli. *Dev Dyn* **211**, 109-
981   121 (1998).
982   73.   Revenu, C. *et al.* A new role for the architecture of microvillar actin bundles in apical
983   retention of membrane proteins. *Mol Biol Cell* **23**, 324-336 (2012).
984   74.   Larsson, E., Moren, B., McMahon, K.A., Parton, R.G. & Lundmark, R. Dynamin2
985   functions as an accessory protein to reduce the rate of caveola internalization. *J Cell Biol*
986   **222** (2023).
987   75.   Jin, M. *et al.* Branched actin networks are organized for asymmetric force production
988   during clathrin-mediated endocytosis in mammalian cells. *Nat Commun* **13**, 3578 (2022).
989   76.   Burke, T.A. *et al.* Homeostatic actin cytoskeleton networks are regulated by assembly
990   factor competition for monomers. *Curr Biol* **24**, 579-585 (2014).
991   77.   Loerke, D. *et al.* Cargo and dynamin regulate clathrin-coated pit maturation. *PLoS Biol* **7**,
992   e57 (2009).
993   78.   Mettlen, M., Loerke, D., Yarar, D., Danuser, G. & Schmid, S.L. Cargo- and adaptor-
994   specific mechanisms regulate clathrin-mediated endocytosis. *J Cell Biol* **188**, 919-933
995   (2010).
996   79.   Mundell, S.J., Luo, J., Benovic, J.L., Conley, P.B. & Poole, A.W. Distinct clathrin-coated
997   pits sort different G protein-coupled receptor cargo. *Traffic* **7**, 1420-1431 (2006).

80. Keyel, P.A. *et al.* A single common portal for clathrin-mediated endocytosis of distinct
cargo governed by cargo-selective adaptors. *Mol Biol Cell* **17**, 4300-4317 (2006).
81. Yu, Y. & Yoshimura, S.H. Self-assembly of CIP4 drives actin-mediated asymmetric pit-
closing in clathrin-mediated endocytosis. *Nat Commun* **14**, 4602 (2023).
82. Boulant, S., Kural, C., Zeeh, J.C., Ubelmann, F. & Kirchhausen, T. Actin dynamics
counteract membrane tension during clathrin-mediated endocytosis. *Nat Cell Biol* **13**,
1124-1131 (2011).
83. Akamatsu, M. *et al.* Principles of self-organization and load adaptation by the actin
cytoskeleton during clathrin-mediated endocytosis. *Elife* **9** (2020).
84. Svitkina, T.M. *et al.* Mechanism of filopodia initiation by reorganization of a dendritic
network. *J Cell Biol* **160**, 409-421 (2003).
85. Croce, A. *et al.* A novel actin barbed-end-capping activity in EPS-8 regulates apical
morphogenesis in intestinal cells of Caenorhabditis elegans. *Nat Cell Biol* **6**, 1173-1179
(2004).
86. McMahon, H.T. & Boucrot, E. Membrane curvature at a glance. *J Cell Sci* **128**, 1065-
1070 (2015).
87. Dergai, M., Iershov, A., Novokhatska, O., Pankivskyi, S. & Rynditch, A. Evolutionary
Changes on the Way to Clathrin-Mediated Endocytosis in Animals. *Genome Biol Evol* **8**,
588-606 (2016).
88. Schmid, S.L., Sorkin, A. & Zerial, M. Endocytosis: Past, present, and future. *Cold Spring
Harb Perspect Biol* **6**, a022509 (2014).
89. de Duve, C. The origin of eukaryotes: a reappraisal. *Nat Rev Genet* **8**, 395-403 (2007).
90. Tilney, L.G., Cotanche, D.A. & Tilney, M.S. Actin filaments, stereocilia and hair cells of
the bird cochlea. VI. How the number and arrangement of stereocilia are determined.
Development **116**, 213-226 (1992).
91. Sheetz, M.P., Turney, S., Qian, H. & Elson, E.L. Nanometre-level analysis demonstrates
that lipid flow does not drive membrane glycoprotein movements. *Nature* **340**, 284-288
(1989).
92. Qian, H., Sheetz, M.P. & Elson, E.L. Single particle tracking. Analysis of diffusion and
flow in two-dimensional systems. *Biophys J* **60**, 910-921 (1991).
1028