Apical clathrin-coated pits control the location, timing, and scale of microvillar growth

Olivia L. Perkins[1], Alexandra G. Mulligan[1], Evan S. Krystofiak[1], K. Elkie Peebles[1], Rekha N. Nagarajan, Leslie M. Meenderink[1,3], Bryan A. Millis[1, 2], Matthew J. Tyska[1*]

[1]Department of Cell and Developmental Biology, Vanderbilt University School of Medicine, Nashville, TN 37232
[2]Department Biomedical Engineering, Vanderbilt University School of Engineering, Nashville, TN 37232
[3]United States Department of Veterans Affairs Tennessee Valley Health Care System, Nashville, TN, 37232

*To whom correspondence should be addressed: matthew.tyska@vanderbilt.edu

____

SUMMARY

Epithelial cells from diverse contexts assemble apical specializations to serve tissue-specific functions. In virtually all cases, these features consist of arrays of microvilli: micron-scale, actin bundle-supported protrusions that mediate biochemical and physical interactions with the external environment. Despite their importance for epithelial physiology, how microvilli grow during cellular differentiation remains poorly understood. Using genetic and small molecule perturbations, we found that an epithelial cell's potential for growing microvilli of normal size is limited by an adjacent actin-dependent process: apical clathrin-mediated endocytosis. Unexpectedly, timelapse imaging of individual microvillar growth events revealed tight spatial and temporal coupling to sites of clathrin-mediated endocytosis. Ultrastructural characterization of undifferentiated epithelial monolayers also showed that most nascent microvilli are in contact with an apical endocytic pit. Finally, inhibition of the Arp2/3 branched nucleation complex, which drives actin polymerization on coated pits, significantly reduced the accumulation of new microvilli on the surface of differentiating epithelial cells. Based on these discoveries, we conclude that clathrin-mediated endocytosis and its associated Arp2/3-based actin nucleation activity control the timing and location of microvillar growth, as well as the dimensions of the resulting protrusions.

____

INTRODUCTION

Actin-based membrane protrusions extend from the surface of virtually all animal cells, where they mediate biochemical and physical interactions with the extracellular environment. Microvilli are canonical examples; as finger-like protrusions that are found in large arrays on the apical surfaces of solute transporting epithelial tissues, they significantly increase the plasma membrane surface area available for holding the channels and transporters that are central to physiological function [1, 2]. Variations on this morphological theme are found throughout the animal kingdom, on organisms ranging from ancient single-celled protists to mammals [3, 4]. Interestingly, previous evolutionary studies suggested that as mammalian cell functions became more specialized, building arrays of microvilli enabled epithelial cells to increase surface area for maximizing solute uptake potential [3, 5]. In specific tissues, such as the intestinal tract and

kidney tubules, epithelial cells build highly ordered arrays of microvilli known as brush borders,
where the apical surface is nearly saturated with protrusions, characterized by hexagonal
spacing when viewed *en face* [6].

The ultrastructure and proteome of intestinal microvilli have been well characterized in
studies spanning decades [7, 8]. Microvilli first appear on the apical surface of epithelial cells
during the early stages of differentiation, and subsequently undergo rapid population expansion
and packing to form mature brush borders [9, 10]. Microvillus structure is highly stereotyped: each
protrusion is supported by 30-40 actin filaments bundled in parallel such that their fast-growing
barbed ends are oriented towards the distal tips [6, 11], which are marked by a complex containing
epidermal growth factor pathway substrate 8 (EPS8) and insulin-receptor tyrosine kinase
substrate (IRTKS) [12, 13]. Parallel filament bundling is supported by fimbrin (also known as
plastin-1), villin, espin, and mitotic spindle positioning protein (MISP) [14-17]. Recent studies
revealed that these factors occupy distinct regions along the length of the core bundle,
suggesting a potential mechanism for regional tuning of physical properties [17]. The core actin
bundle is connected to the encapsulating membrane through membrane-cytoskeleton linkers,
including ezrin and myosin-1a, which maintain membrane wrapping and increase long-term
protrusion stability [18, 19]. Despite significant progress toward defining the structure and
composition of microvilli, mechanisms that control the growth of these protrusions remain poorly
understood.

The morphologies of surface features such as microvilli are regulated by a balance of
force-generating cytoskeletal elements and the force-resisting physical properties of the plasma
membrane [20, 21]. Endocytosis, the process by which cells internalize membrane and extracellular
materials, is well positioned to impact this force balance as it involves localized membrane
deformation that is potentiated by actin polymerization. The series of molecular events that drive
actin assembly during clathrin-mediated endocytosis (CME) is well-studied in both yeast and
mammalian cells [22]. Synthesis of phosphatidylinositol 4,5-bisphosphate (PI(4,5)P$_2$) at clathrin
coated pits (CCPs) [23] triggers the recruitment of actin-binding proteins including neuronal-
Wiscott-Aldrich syndrome protein (N-WASP)[24] and actin related protein 2/3 complex (Arp2/3)[25],
capping protein [26], and actin-bundling [27] and severing proteins such as cofilin [28]; these proteins
work together to promote the polymerization and turnover of actin networks to drive the
formation and scission of CCPs [22]. Although CCPs can internalize without actin assembly, in
high membrane tension scenarios such as the epithelial apical surface, actin polymerization
counteracts the additional resistance to membrane bending [29, 30]. Intriguingly, CCPs and
microvilli share several molecular components such as actin-binding and adaptor molecules,
suggesting a potential mechanistic link between these functionally and topologically distinct
apical processes. For example, in solute transporting epithelial cells, PI(4,5)P$_2$ is enriched in the
apical membrane and has been shown to recruit and activate ezrin, a key membrane-
cytoskeletal linker in the brush border [31]. Actin-bundling proteins like fimbrin, which crosslink
filaments in the microvillar core actin bundle, are also recruited to endocytic patches [15, 27].
Mammalian actin binding protein 1 (mAbp1), with binding domains for both actin and clathrin,
acts as a molecular bridge by physically linking the actin cytoskeleton to endocytic machinery [32].
Notably, mAbp1 has also been shown to bind two molecules implicated in building microvilli:
pacsin2 (also known as syndapin2) [33] and cordon bleu (COBL) [34]. Previous studies established
that knockout of pacsin2 in mice resulted in significantly shorter and fewer microvilli with

reduced membrane wrapping on the surface of mature villus enterocytes [35]. That work also
revealed stalled endocytic intermediates, which accumulated between microvilli, suggesting a
defect in endocytic activity. Although the mechanisms underlying these phenotypes remain
unclear, when combined with the molecular links alluded to above, these studies collectively
suggest that endocytic activity at the apical surface may be needed for normal microvillar
growth.
Does CME control the growth of microvilli? To address this question, we used a
combination of genetic and small molecule perturbations coupled with live confocal and
ultrastructural imaging to examine microvillar dynamics and structure on the apical surface of
brush border-building epithelial cell lines. Perturbation of CME in actively differentiating cells led
to dramatic remodeling of microvilli; activation of CME depleted microvilli, whereas inhibition
resulted in exaggerated protrusions containing more actin filaments and actin-binding proteins.
Interestingly, volumetric live-cell imaging revealed that microvilli grow directly from CCPs, and
scanning electron microscopy (SEM) also revealed a tight spatial association between apical
surface pits and microvilli in both kidney- and intestine-derived transporting epithelial cells.
Finally, inhibition of the Arp2/3 branched nucleation complex, which drives actin polymerization
on coated pits, significantly reduced the accumulation of new microvilli on the surface of
differentiating epithelial cells. Together, these data suggest that, across multiple epithelial
models, CME, and its associated Arp2/3-based actin nucleation activity, controls the timing and
location of microvillar growth as well as the dimensions of resulting protrusions. These findings
redefine our understanding of microvillar biogenesis and, given the widespread appearance of
CME in diverse biological systems, hold broad implications for understanding how cell surface
morphologies are controlled.
111

RESULTS

Endocytic markers localize to the apical surface of actively differentiating epithelial cells
113
While previous studies established that the endocytic machinery localizes to the apical surface
of functionally mature transporting epithelial cells, whether endocytosis plays a role in microvillar
growth during epithelial differentiation remains unexplored [36, 37]. We first turned to SEM to
examine the surface of cells lining intestinal crypts, the site of microvillar growth in intestinal
tissues, where nutrient transporting enterocytes comprise ~95% of the epithelium. The intestinal
crypt-villus axis offers a unique opportunity for imaging epithelial cells across all stages of
differentiation; crypts house the transit amplifying cells (actively differentiating) while the villus is
populated by mature epithelial cells (fully differentiated) [38]. SEM of crypts in fractured samples
of mouse small intestine revealed a gradual increase in microvillar density on the apical surface,
suggesting that the brush border continuously packs with protrusions as cells migrate from the
base of the crypt to the crypt-villus transition (Fig. 1A,B). We confirmed this point using
immunofluorescent staining to probe for the microvillar-specific actin-bundler, villin, as a proxy
for microvillar density. Consistent with the SEM data, confocal images revealed the apical villin
signal increased nearly 2-fold from crypt base (magenta) to crypt-villus transition (green) (Fig.
1C,D), indicative of microvillar accumulation on the surface of these cells. We next stained for
dynamin2 and pacsin2 to assess the abundance of these canonical endocytic markers in the
crypts. The resulting confocal images revealed that, similar to villin, both markers demonstrated
a gradual increase in signal, from the crypt base (magenta) to crypt-villus transition (Fig. 1C,D).

Collectively, these data suggest that the accumulation of microvilli at the apex of actively
differentiating cells spatially and temporally coincides with an increase in endocytic marker
enrichment at the apical surface, well before cells are exposed to the contents of the lumen.
135

Induction of clathrin-mediated endocytosis depletes microvilli
Given the localization of clathrin-mediated endocytic components to the apical surface of
differentiating epithelial cells in native intestinal crypts, we sought to determine if perturbing
CME impacted microvillar growth. Here we employed the previously established "hook and
anchor" tool to induce clathrin-mediated endocytosis [39]. In this system, FKBP-tagged β2 subunit
of AP-2 (β2) is recruited to an FRB-tagged CD8 membrane anchor upon addition of rapalog,
resulting in robust formation and internalization of CCPs in a temporally controlled manner (Fig.
2A,B). We expressed this system in undifferentiated CL4 cells along with Halo-tagged espin, a
microvillar actin bundling protein that specifically labels microvilli [16]. Following endocytic
induction, we performed live spinning disk confocal (SDC) imaging to monitor any effects on
microvillar growth and structure (Fig. 2C). To measure the impact on core actin bundle
structure, we generated linescans orthogonal to individual protrusions visualized in the Halo-
espin channel, and calculated areas under the resulting intensity curves (AUC); the resulting
metric is expected to reflect the diameter of the core bundle and the number of constituent actin
filaments (Fig. 2D). Performing this analysis on cells before and 45 min after rapalog treatment
revealed that inducing CME resulted in a ~50% reduction in Halo-espin signal (Fig. 2E).
Additionally, total microvillar surface coverage after endocytosis induction was also reduced
relative to the GFP control (Fig. 2F). Together these data suggest that induction of CME
negatively impacts microvillar growth and stability.
155

Inhibition of clathrin-meditated endocytosis leads to the formation of exaggerated
microvilli
We next sought to determine the impact of endocytic inhibition on microvillar assembly. For
these studies, we used well established inhibitors to disrupt specific steps in the CME pathway
and employed SDC microscopy to visualize the impact on protrusion growth and dynamics on
the surface of CL4 cells expressing mCherry-espin[16, 40]. We inhibited CME by targeting both the
initiation and scission steps of the pathway through chlorpromazine (CPZ) and dyngo,
respectively. CPZ disrupts clathrin lattices on the plasma membrane and targets AP-2 to
endosomes, while dyngo inhibits dynamin2, a GTPase critical for vesical scission during the
final step of CCP invagination [41, 42] (Fig. 3A). CPZ treatment of undifferentiated CL4 cells
resulted in dramatic exaggeration of microvillar structure (Movie S1); following 45 minutes of
CPZ treatment, protrusions demonstrated a ~1.5-fold increase in AUC measured from the espin
signal and ~2-fold increase in length (Fig. 3B-D, Movie S1). This effect was titrated across a
range of CPZ concentrations in CL4 cells and resulted in a similar phenotype in CACO-2$_{BBE}$ cells
treated with CPZ (Fig. S1). In contrast, 45 minutes of dyngo treatment did not significantly
impact microvillar morphology (Fig. 3B-D). To ensure that the observed phenotypic differences
were not due to disparities in drug efficacy, we performed a transferrin uptake assay to assess
CME activity following 45 minutes of treatment with either CPZ or dyngo. As expected, both
endocytosis inhibitors significantly reduced transferrin uptake (Fig. S2). Thus, blocking
endocytosis early in the pathway (Fig. 3A) promotes the growth of larger microvilli.

The increase in mCherry-espin signal in microvilli following CPZ treatment (Fig. 3B-D)
suggests that growing core bundles incorporate more actin filaments in the absence of normal
endocytic activity. To determine if this is the case, CL4 cells expressing mNeon-Green-ß-actin
were treated with CPZ as described above (Fig. 3E). Consistent with the mCherry-espin
readout, we observed a ~2-fold increase in AUC, indicating that CPZ treatment indeed leads to
higher levels of actin in exaggerated microvilli post-CPZ treatment (Fig. 3F). Labeling CPZ-
treated cells with CellMask-DeepRed (CellMask), a plasma membrane dye, further revealed that
exaggerated microvilli on these cells retained membrane wrapping, with the length of the
wrapped region increasing with the longer core bundles formed under these conditions (Fig.
4A,B). We also examined the localization of ezrin, a major membrane-actin linker in these
protrusions [43]. AUC analysis of EGFP-ezrin signal in CPZ-treated cells further confirmed the
increased diameter of exaggerated microvilli (Fig. 4C,D). Finally, we examined the impact of
endocytic inhibition on two factors that reside at the distal tips of microvilli: EPS8 and IRTKS [12].
Distal tip fluorescence intensity of EGFP-EPS8 or EGFP-IRTKS, quantified using Trackmate,
revealed an enrichment of both EPS8 and IRTKS after CPZ treatment (Fig. 4E-H). Collectively,
these data indicate that, under normal conditions, CME plays a role in limiting microvillar
dimensions and actin content.
193
Exaggerated microvilli that form following CPZ treatment exhibit abnormal motility
Microvilli that extend from differentiating epithelial cells exhibit active motility across the apical
surface, which is driven by the treadmilling of the supporting core actin bundles; this activity
drives collisions between nascent protrusions, which in turn promote the formation of
intermicrovillar adhesion complexes and microvilli packing [44]. Given the increased length and
filament content of microvilli following CPZ treatment, we sought to determine if these
exaggerated protrusions exhibit aberrant motility. To address this question, we employed live
SDC imaging to track the motility of individual microvilli on CL4 cells co-expressing mCherry-
espin and EGFP-EPS8 (Fig. 5A). Mean square displacement analysis of 50 randomly selected
microvillar trajectories per condition revealed parabolic curves consistent with a mixture of
robust active motion and limited diffusion (Fig. 5D). Further, while we observed no significant
changes in velocity nor track displacement after endocytosis inhibition, exaggerated microvilli
did exhibit higher persistence of motion, which was reflected in the extended reach of individual
trajectories when viewed in rose plots (Fig. 5B,C,E-G). Thus, inhibition of CME with CPZ leads
to the growth of exaggerated microvilli that exhibit abnormal motility.
209
Global reduction of plasma membrane tension does not lead to the formation of
exaggerated microvilli
Endocytic activity plays a global role in decreasing plasma membrane surface area and thus,
increasing tension in this bilayer [45]. Moreover, previous studies on CPZ suggest this compound
intercalates into the plasma membrane, which is expected to reduce membrane tension [46, 47]. To
determine if global perturbations in membrane tension are responsible for the growth of
exaggerated microvilli post-CPZ treatment, we treated CL4 cells with other compounds that are
known to reduce membrane tension including 400 μM deoxycholate, 1% ethanol, and addition
of 125 mM NaCl; all of these treatments decrease membrane tension as measured in tether
force assays [48-50]. However, none of these global perturbations induced the formation of

microvilli with exaggerated dimensions (Fig. S3). These results suggest that endocytic events
might instead locally regulate the growth and dimensions of microvilli.
222
Microvilli grow from apical clathrin-coated pits
We next sought to determine if endocytic sites are spatially and temporally linked to microvillar
growth events on the apical surface. To this end, we leveraged a recently developed SDC-
based live imaging approach which allows for the visualization of individual microvillar growth
events on the surface of sub-confluent CL4 cells [9]. To screen for potential spatiotemporal links
between microvillar growth and CME, we imaged mCherry-espin expressing CL4s co-
expressing a range of EGFP-tagged endocytic components including pacsin2, clathrin light
chain A (LCA), dynamin2, mAbp1, and $PI(4,5)P_2$ (Fig. 6A,C,E and Fig. S4) [32, 51-53]. To quantify
the localization of endocytic markers at sites of microvillar growth, we analyzed the maximum
intensity projected fluorescence signals of CME marker puncta that colocalized with elongating
espin signals (representing the growth of new microvillar core bundles) over time. Strikingly,
most endocytic markers we examined localized to the apical surface at future sites of microvillar
growth ~2-3 minutes prior to espin accumulation and elongation (Fig. 6B,D,F). One exception to
this pattern was dynamin2, which briefly localized ~1 minute after espin accumulation began,
presumably marking the disassembly of the CCP (Fig. S4C). Following their localization to sites
of microvillar growth, most endocytic components disassembled within a few minutes, a
timeframe consistent with previously reported dynamics of CME [54]. Together, these timelapse
recordings suggest that the early steps of CCP formation create a specialized apical membrane
domain that is permissive for microvillar growth.
242
Ultrastructural imaging reveals juxtaposition of nascent microvilli and endocytic pits
To further examine the relationship between CCP formation and the growth of new microvilli, we
employed SEM to visualize fine structural details of the apical surface of undifferentiated CL4
cells. Strikingly, these data showed that 51.1% of the pits that we could visualize were in
proximity to microvilli (<250 nm) (Fig. 7A,C). To determine if the observed spacing was closer
than expected based on randomly distributed pits and microvilli, we performed a nearest
neighbor analysis, measuring from the center of every pit to the center of the base of the
nearest microvillus. We also simulated cell surfaces and nearest neighbor measurements with
the same number of pits and microvilli randomly distributed over the same area (Fig. S5).
Remarkably, experimental nearest neighbor measurements exhibited a skewed distribution with
a peak at ~200 nm; in contrast, simulated random nearest neighbor measurements
demonstrated a Gaussian distribution with a wider peak at ~450 nm (Fig. 7C). Thus, the
observed spacing between endocytic pits and microvilli cannot be explained by random co-
occurrence and is instead consistent with two processes that are linked in space and time. We
also noted that microvilli associated with endocytic pits were significantly shorter than microvilli
further away, suggesting that core bundles mature as they as they disassociate from CCPs (Fig.
7D). Expanding the significance of this finding, we observed the same close juxtaposition of
endocytic pits and microvilli on the surface of $CACO2_{BBE}$ cells (67.1% of pits associated with
microvillus) (Fig. 7B,E). These ultrastructural findings provide a compelling complement to our
timelapse data, and strongly indicate that CME and microvillar biogenesis are spatially and
temporally coordinated processes at the apical cell surface.

Arp2/3 complex inhibition decreases the growth of new microvilli

Actin filament assembly on the surface of CCPs, driven by the Arp2/3 branched nucleation
complex, is critical for promoting pit closure and internalization under conditions of high
membrane tension, like that experienced at the apical surface [29, 30]. Given the close association
of nascent microvilli and endocytic pits revealed in our confocal and SEM imaging studies, we
wondered if Arp2/3-generated actin filaments on the surface of CCPs directly support the
formation of microvillar core actin bundles. Because the actin nucleators that support microvillar
growth have yet to be identified, we were eager to test this idea. Toward this end, we used
SDC-based live imaging to visualize individual microvillar growth events on the surface of sub-
confluent CL4 cells [9] in the presence of CK-666, a potent inhibitor of Arp2/3 nucleation activity
[55]. Time-lapse acquisitions over the course of 30 min revealed a 10-fold reduction in the rate of
microvillar growth initiation following CK-666 treatment (Fig. 7A,B). When acquisitions were run
for longer periods of time (90 min), analysis of total microvillar content on the apical surface also
revealed a striking decrease in protrusions, suggesting that the reduced growth rates following
CK-666 treatment also led to reduced accumulation of protrusions long-term (Fig. 7C,D).
Together, these results suggest that actin filaments generated by CCP-associated Arp2/3 are
remodeled to form the parallel, polarized core bundles that support microvilli.

DISCUSSION

Despite the extensive characterization of microvillus architecture and composition, our
understanding of how cells control the growth and resulting dimensions of these protrusions
remains limited. Here, we establish apical endocytic activity as a critical regulator of the location,
timing, and dimensions of nascent microvilli. Given the overlapping molecular machinery and
the role of the actin cytoskeleton and membrane bending in both apical CME and microvilli, we
sought to test the hypothesis that microvillar growth is linked to apical endocytic activity. Indeed,
our analysis of endocytic markers (pacsin2 and dynamin2) in native intestinal crypts revealed
that differentiating cells experiencing rapid microvillar growth also exhibit robust apical
localization of endocytic machinery. This finding revealed that apical endocytic activity is well
positioned to influence microvillar assembly early in the time course of epithelial differentiation.

How might CME impact microvillar growth on the apical surface of differentiating epithelial cells?
Previous studies collectively indicated that experimental depletion of microvilli is difficult,
underscoring the robustness of this evolutionarily ancient process [56-59]. However, we found that
inducing high levels of CME through the recruitment of the β2 subunit of AP2 to the plasma
membrane led to a striking depletion of microvillar actin content and coverage across the cell
surface. Our results echo those from previous studies which showed that overexpression of
clathrin heavy chain perturbed the morphology of microvilli on the surface of Madin-Darby
Canine Kidney (MDCK) cells, as assessed using transmission electron microscopy [30]. A
superficial interpretation of these findings is that apical CME influences microvillar growth by
impacting apical membrane tension. Yet, in the current study, global perturbations of membrane
tension using orthogonal methods failed to disrupt microvillar architecture. An appealing
alternative model is that apical CCPs and microvillar assembly mechanisms share resources,
such as specific protein machinery, membrane lipids, or actin filaments, and over- or

underutilization of these resources by CME compromises the growth of protrusions from the
apical surface. We elaborate more on the experimental support for this idea below.
310
If CCPs and microvilli share resources during their assembly, then inhibition of CME would be
expected to divert those components into the microvillar growth pathway, and potentially result
in more protrusions and/or larger structures. Indeed, when we inhibited CME using CPZ, a
compound that inhibits clathrin recycling to the plasma membrane, new microvilli grew with
exaggerated dimensions, appearing longer and with increased diameter, suggesting they
contained more actin filaments. Protrusions assembled following CPZ treatment retained key
structural features of canonical microvilli, including ezrin-mediated membrane wrapping, and
distal tip localization of EPS8 and IRTKS. Interestingly, another endocytic inhibitor, dyngo, did
not have the same effect, although this may be explained by the distinct mechanisms by which
CPZ and dyngo block CME (Fig. 3A)[41, 42, 60]. CPZ prevents the formation of coated pits by
disrupting AP-2 recruitment to the plasma membrane. In contrast, dyngo treatment allows for pit
formation and leads to stabilization of pits on the plasma membrane by inhibiting vesicle
scission. The stalled endocytic intermediates created by dyngo still sequester resources such as
actin filaments, as suggested in previous studies [61, 62], potentially explaining why this treatment
does not lead to microvillar overgrowth.
326
The apical surface phenotypes associated with gain and loss of CME function – depletion and
exaggeration of microvilli, respectively - indicate that CCP formation and microvillar growth are
mechanistically linked, potentially through a shared requirement for a molecular resource such
as actin. This link could operate globally with these pathways engaging in long-range
competition for the same pool of actin monomers, as suggested in previous studies on actin
allocation [63]. Alternatively, this link could operate locally, with both pathways sharing resources
at the same time *and* in the same location at the apical membrane. This latter possibility is
supported by our live imaging studies of epithelial cells early in differentiation. Using SDC
microscopy, we defined a time course for the recruitment of endocytic machinery, which
appeared in diffraction limited puncta on the apical surface in events that lasted several minutes
as expected[54, 64]. Remarkably, these puncta also marked future sites of microvillar growth, with
some CME markers arriving 2-3 minutes in advance of growth initiation. All CME markers
examined in this assay demonstrated significant temporal and spatial overlap with the
microvillar core bundle marker, espin, suggesting that resource sharing could occur at discrete
sites that are scaled at or below the diffraction limit (< 200 nm). Interestingly, we noted that not
all CCPs progressed to form microvilli. Given the variation in composition of CCPs, it may be
that only a subset of pits are equipped to support microvillar growth[65-67]. What distinguishes
these pits from their counterparts? One intriguing possibility lies in the mechanism of pit closure.
Studies have identified three distinct closure types for CME: symmetric, asymmetric, and
undefined, with asymmetric being the most prevalent (accounting for roughly 70% of pits) [62, 68].
During asymmetric closure, membrane bulges outward on one side of the CCP propelled by
actin polymerization, progressively covering and closing the pit. This contrasts with symmetric
closure (~20% of events), where the pit slowly closes in an isotropic manner[62, 68]. The specific
mode of pit closure and/or the presence of asymmetric F-actin accumulation might dictate
whether that pit ultimately leads to microvillus assembly[29, 62].

We also surveyed the apical surfaces of fixed epithelial cell monolayers using SEM, with the
goal of examining large numbers of endocytic pits and characterizing their spatial relationship
with microvilli. Analysis of undifferentiated CL4 and CACO-2$_{BBE}$ monolayers revealed that >50%
of pits were within 250 nm of a microvillus (CL4: 51.1%, CACO-2 $_{BBE}$: 67.1%), a distance
comparable to the diameter of the diffraction limited puncta that we observed in our live confocal
imaging studies. Mathematical simulations revealed that such close juxtaposition of endocytic
pits and microvilli could not be explained by the random positioning of these objects across the
apical surface. We also found that pit-associated microvilli were significantly shorter relative to
more distal protrusions. This observation could indicate that pit-associated microvilli represent
younger intermediates, whereas longer and more distal microvilli represent fully formed
protrusions at steady-state length. In combination with the CME gain and loss of function
studies and the timelapse observations described above, these data lead us to propose a model
whereby microvilli form at sites that are immediately adjacent (within 250 nm) to pre-existing
CCPs (Fig. 6F).

Why do CCPs serve as preferential sites for microvillar assembly? From several perspectives,
the local biochemical and physical landscape generated during endocytic pit formation appears
well-suited for supporting protrusion growth. For example, CCPs contain a significant F-actin
coat generated by the Arp2/3 branched nucleation complex, and the protein machinery that
drives microvillar growth could harness those polymers, elongating and bundling them to form
core actin bundles. Interestingly, molecular counting studies revealed that ~200 Arp2/3
complexes localize to CCPs, and simulations suggest that this population would be sufficient to
generate ~150 filaments for driving pit internalization[69]. Based on ultrastructural studies of the
enterocyte brush border, we know that microvillar core actin bundles contain only ~30-40
filaments [11], so the scale of the CCP-associated actin filament network is large enough to
support the growth of a few microvilli. Inhibition of the Arp2/3 branched nucleation complex in
CL4 cells significantly reduced the growth and accumulation of new microvilli on the surface of
differentiating epithelial cells, suggesting that the actin filaments in the microvillus core bundle
are derived from Arp2/3 generated networks (Fig. 7). The general concept of co-opting filaments
from an existing network to form a new structure with distinct architecture is similar to the
'convergent elongation' model that was originally proposed by Svitkina and colleagues to
explain the formation of filopodia on the surface of crawling cells[70]. In that scenario, the actin
filaments that support filopodia originate deep in the lamellipodial meshwork, which is generated
by Arp2/3 activity. During filopodial growth, a small number of lamellipodial filaments are
protected from capping by barbed end elongation factors (Ena/VASP, mDia) and therefore
allowed to elongate and become crosslinked by fascin to create a mechanically robust core
bundle[70]. In the context of microvillar growth, EPS8 and/or IRTKS might function as elongation
factors[12], whereas espin and other factors (MISP, fimbrin, and villin) could participate in filament
bundling[14-17]. Whether a similar mechanism is operational at the smaller scale of CCPs and
leads to formation of microvilli will require deeper ultrastructural investigation.

SEM images revealed that when an endocytic pit is adjacent to a microvillus, the stoichiometry
is almost always 1:1, i.e. one microvillus per pit. Thus, in addition to the availability of actin

filaments, other constraints probably limit the number of new microvilli that form during these
events. In our studies, we also observed enrichment of PI(4,5)P$_2$ before microvilli begin to grow,
suggesting a potential role for specific lipids in microvillar biogenesis. The local accumulation of
specific lipids or membrane-associated proteins is known to impact membrane curvature and
tension[71], and these physical factors likely impact microvillar growth. Future investigations
taking advantage of next generation membrane tension probes might explore this possibility.
402
The link between CME and microvillar growth revealed in this work points to a fascinating
evolutionary adaptation in epithelial biology. While CME is a ubiquitous cellular process shared
by all eukaryotes, microvilli are specialized structures that emerged later in evolution, with their
earliest known appearance in choanoflagellates[3, 72]. The evolution of endocytosis provided a
crucial foundation for membrane maintenance, cell signaling, and nutrient uptake[73, 74]. We
propose that, over time, this actin-dependent process was co-opted to support the formation of
actin-based protrusions such as microvilli. If true, this would represent an elegant example of
evolutionary exaptation, where existing processes are repurposed to meet new functional
demands. The model we propose here might also offer an explanation for growth of more
elaborate microvilli-like structures, including the stereocilia that extend from the apical surface of
mechanosensory hair cells found in the cochlear and vestibular systems.
414
415

416 ## ACKNOWLEDGEMENTS
417 The authors thank all members of the Tyska laboratory for their constructive feedback.
418 Microscopy was performed in part by the Vanderbilt Cell Imaging Shared Resource. This study
419 was supported by the NIH grants R01 DK095811 (M.J.T.), R01 DK125546 (M.J.T.), and R01
420 DK111949 (M.J.T.).

METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to the lead contact, Matthew J. Tyska (matthew.tyska@vanderbilt.edu).

Materials availability

Plasmids generated in this study will be made available from the lead contact on request.

Data code availability

No large-scale datasets or new code were generated in this study.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cell culture models

LLC-PK1-CL4 (porcine kidney proximal tubule) cells were grown in 1X high glucose DMEM containing 2mM L-glutamine (Corning #10-0130CV) supplemented with 10% fetal bovine serum (FBS) (R&D Systems) and 1% L-Glutamine (Corning # 25-005-Cl). CACO-2$_{BBE}$ cells were grown in 1x high glucose DMEM containing 2mm L-glutamine supplemented with 20% FBS. Cells were maintained in treated plastic culture flasks incubated at 37 °C and 5% CO_2. Cells were tested for mycoplasma monthly using the MycoStrip Mycoplasma Detection Kit (Invivogen #rep-mys-50).

METHOD DETAILS

Cloning and Constructs

All overexpression constructs listed in this paper were previously generated and/or obtained as noted in the key resources table.

Cell line generation

LLC-PK1-CL4 cells stably expressing pmCherry-espin were previously generated using the stable cell line generation protocol as described in Gaeta et al., 2022. Briefly, cells were transfected with Lipofectamine 2000 following manufacturer protocol in a T25 cell culture flask. The next day, cells were split up to a T75 flask and treated with 1 mg/mL G418 sulfate for antibiotic selection. Cells were maintained in culture under constant G418 selection to create the stably expressing cell line.

Transient transfection

478 LLC-PK1-CL4 cells were seeded in 6 well plates and transfected at ~90% confluency using
479 Lipofectamine2000 with 3 μg DNA. 4 hrs after incubation with the lipid complex, cells were
480 replated on CellVis 35mm 1.5# glass bottom dishes for imaging the following day.
481
Immunofluorescence staining
Tissue: Tissue sections were deparaffinized and rehydrated through a descending ethanol
series and subjected to antigen retrieval in a heated buffer at pH 9. Following cooling, sections
were washed and blocked with 10% normal goat serum (NGS) and probed for villin, pacsin2, or
dynamin2 in 1% NGS overnight at 4 °C. The following day, sections were washed, incubated at
room temperature with secondary antibodies, dehydrated through an ascending ethanol series,
and mounted in ProLong Gold mounting medium.
489
Live imaging microscopy
Nikon Ti2 inverted light microscope equipped with a Yokogawa CSU-X1 spinning disk head, a
Photometrics Prime 95B sCMOS camera, and 488 nm, 561 nm, and 647 nm excitation lasers. A
60x/1.49 NA or 100x/1.49 NA TIRF oil immersion objective was used for all experiments. Cells
were maintained in a stage top incubator at 37 °C with 5% CO2 (Tokai Hit). Z stacks of 5-8 μm
using a 0.2-0.4 μm step size were collected using a triggered NIDAQ piezo Z stage.
496
Scanning electron microscopy (SEM)
Tissue: 5 mm murine duodenal sections were fixed in 2.5% glutaraldehyde and 4%
paraformaldehyde in SEM, washed, and then embedded in OCT. To ensure stability of the
explant lumens, samples were gently passed through 3 rounds of fresh OCT. Samples were
then frozen in cryomolds and stored at -80 °C . Frozen explants were sectioned, melted onto
stainless steel AFM discs, and processed through OsO4, ddH2O, and subsequently dehydrated
through graded ethanol series. Detached sections were recovered and adhered to aluminum
SEM specimen stubs. SEM imaging was performed on a Quanta 250 environmental SEM or
Zeiss Crossbeam 550. For more detailed methods, see Meenderink et al[44].
506
Cells: Caco-2$_{BBE}$ cells were plated at 70% confluence on Thermanox coverslips in a 6-well dish.
24 hrs post-plating, cells were washed 3X with PBS to remove all media and were fixed in a
buffer of 2.5% glutaraldehyde and 0.1 M CaCl$_2$ free cacodylate overnight at room temperature.
Coverslips were washed 3X in 0.1 M CaCl$_2$ free cacodylate buffer and incubated in pre-warmed
1% tannic acid in cacodylate. Following washes with 0.1 M cacodylate, coverslips were
incubated for 1 hour in 1% OsO$_4$. After washes with ddH$_2$O, they were incubated for 30 min with
1% uranyl acetate in ddH$_2$O. Finally, coverslips were washed with ddH$_2$O 3X for 5 min to
remove uranyl acetate before drying with a series of graded EtOH washes. After samples were
dried using critical point drying and vented for 20 min, coverslips were mounted on SEM
specimen stubs via a conductive adhesive tab and sputter coated prior to imaging. SEM
imaging was performed on a Zeiss Crossbeam 550 with a voltage of 2 keV and an SE2
detector. All reagents were purchased from Electron Microscopy Sciences.
519
Transferrin uptake assay

LLC-PK1-CL4 cells expressing mCherry-espin were seeded on acid-washed #1.5 coverslips at
~60% confluence in a cell culture 6-well treated plate on day 0. The next day, cells were treated
with 30 µM CPZ, 30 µM Dyngo, or DMSO and incubated for 45 min. At the 45-min mark,
transferrin conjugate was added to the cells at 25 µg/mL and incubated for 20 min. After 20 min,
the cells were fixed in 4% paraformaldehyde diluted in PBS and incubated at 37 °C for 15 min.
Coverslips were subsequently mounted on slides in Prolong Gold antifade reagent and imaged
using spinning disk microscopy.
528
529 *QUANTIFICATION AND STATISTICAL ANALYSIS*
530 **Statistical testing**
531 All statistical testing was performed in GraphPad Prism 9 by running a Mann Whitney test or a
532 paired t-test for pairwise comparisons or Kruskal-Wallis to compare more than two conditions.
533 For all figures, exact n values and their definitions are reported in the figure legends.
534
535 **Time series analysis of microvillus growth**
536 Microvillar growth was analyzed as previously described[9]. Briefly, a bounding ROI was manually
537 drawn around the signal of interest and mean intensity was measured for each timeframe. Mean
538 intensity values were normalized, with the smallest value equal to 0 and the largest equal to
539 one. Normalized values were then averaged and plotted with SEM.
540
Area under the curve analysis
To measure the impact on core actin bundle structure, we generated linescans (5 pixels in
width, 1.54 µm in length) orthogonal to individual protrusions visualized in the Halo-EPSN
channel. Linescans were drawn to ensure lines only crossed through one protrusion per line.
Areas under the resulting intensity curves (AUC) were then calculated in prism, averaged and
plotted; the resulting metric is expected to reflect the diameter of the core bundle.
547
548 **EGFP-EPS8 and EGFP-IRTKS puncta intensities**
549 Mean puncta intensities were measured using Nikon Elements' "Spot Detection" function.
550
Microvilli tracking using EGFP-EPS8 puncta
Raw 3D movies were converted into maximum intensity projections, and the EPS8 channel
(488) was isolated. The isolated channel movie was loaded into Trackmate and puncta were
identified using a DoG detector with an estimated object diameter of 500 nm. Identified spots
were thresholded based on background intensity and track length. Tracks beginning on the first
frame or ending on the last frame were excluded. Track and spot data were then exported to
Excel and analyzed and plotted in Prism as radial XY positions over time by subtracting each
position in X or Y from their respective point position at time 0, making the first position (0,0).
Persistence, track displacement, and linear velocity were taken from tracks lasting exactly 10
min.
561
Mean squared displacement analysis
EPS8 puncta trajectories as a proxy for microvillar motion were analyzed using mean squared
displacement (MSD) analysis as previously described[44]. MSD curves were subsequently fit with

an active motion model in the form MSD(nDt) = 4Dt + V^2* nDt^2, where D is the diffusion
coefficient and V is the velocity of active motion [75]. Curve fitting and sum-of-squares F-tests
were performed using PRISM v.9.
568
Nearest neighbor expected distribution simulation
To determine the expected nearest neighbor distance between pits and microvilli, a random
population of microvilli and endocytic pits were simulated within a rectangular area using a
custom python script. The number of objects in each population was set to match the observed
experimental densities. Euclidean distances between each endocytic pit and every microvillus
were measured and the subsequent minimum distance (nearest neighbor) for each endocytic pit
was identified. Each simulation (CACO-2$_{BBE}$ and CL4) was run 10 times and then averaged
(Extended Data Fig. 5).

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and virus strains		
E. coli DH5-Alpha competent cells	Molecular Cell Biology Resource Core, Vanderbilt Medical Center	Item# DH5 Alpha
Chemicals, peptides, and recombinant proteins		
G418 Sulfate	Gold Biotechology	Cat# G-418-1
Janelia Fluor635 (JF635)	Janelia	N/A
Lipofectamine 2000	Thermo Fischer Scientific	Cat# 11668019
A/C Heterodimerizer	Takara	Cat# 635057
488 Transferrin	Thermo Fischer Scientific	T13342
CellMask-Deep Red	Thermo Fischer Scientific	C10046
Alexa Fluor 647 Phalloidin	Thermo Fischer Scientific	A22287
16% Paraformaldehyde	Electron Microscopy Sciences	Cat# 15710
ProLong Gold Antifade Reagent	Thermo Scientific	P36930
Triton X-100	Sigma	Cat# T8787
DMSO	Sigma	Cat# D8418
Chlorpromazine	Selleck Chemicals	S5749
Dyngo-4a	Selleck Chemicals	S7163
Deoxycholate		
Glutaraldehyde, 25%	Electron Microscopy Sciences	Cat# 16220
Tannic Acid	Electron Microscopy Sciences	Cat# 21700
Osmium Tetroxide	Electron Microscopy Sciences	Cat# 19112
Tissue-Tek OCT	Sakura Finetek	#4583
Cell Lines		
LLC-PK1-CL4	Gift from Dr. Carolyn Slayman (Yale University)	N/A
Antibodies		
anti-villin (mouse) [1:100]	Santa Cruz Biotechnology	sc-66022
dynamin2 (rabbit) [1:200]	Sigma-Aldrich	HPA054246
pacsin2 (rabbit) [1:200]	Sigma-Aldrich	HPA049854
Recombinant DNA		
mCherry-C1-syndapin2 (mouse)	Addgene	#27681
GFP-ß2 (616-951)-FKBP (human)	Adgene	#59353100729
DNMII-pmCherry-N1 (dynamin2) (human)	Addgene	#27689
pmCherry-espin (rat)	Gift from Dr. James Bartles, NWU	N/A
mNeonGreen-βActin (human)	Tyska Lab	N/A
pEGFP-C1-EPS8 (human)	Tyska Lab	N/A
pEGFP-C1-IRTKS (human)	Tyska Lab	N/A
pEGFP-N1-ezrin (human)	Addgene	#202684
pGFP-C1-espin (rat)	Tyska Lab	N/A
Halo-C1-espin (rat)	Tyska Lab	N/A

pGFP-FKBP	Addgene	#67529
pCD8-mCherry-FRB (human)	Addgene	#100738
pEGFP-C2-mAbp1 (human)	Addgene	#26719
pPLC-∆PH-GFP	Addgene	#21179
Software and Algorithms		
FIJI		
NIS AR Elements Analysis	Nikon	N/A
Prism 9	Graphpad	N/A
BioRender	BioRender	N/A
Procreate	Savage Interactive	N/A
Other		
Mycostrip^{TM} – Myocplasma Detection Kit	InvivoGen	Cat# rep-mys50
35mm #1.5 glass bottom dishes	CellVis	Cat# D35-20-1.5N

616
617
618

REFERENCES

1. Helander, H.F. & Fandriks, L. Surface area of the digestive tract - revisited. *Scand J Gastroenterol* **49**, 681-689 (2014).
2. Delacour, D., Salomon, J., Robine, S. & Louvard, D. Plasticity of the brush border - the yin and yang of intestinal homeostasis. *Nat Rev Gastroenterol Hepatol* **13**, 161-174 (2016).
3. Pena, J.F. *et al.* Conserved expression of vertebrate microvillar gene homologs in choanocytes of freshwater sponges. *Evodevo* **7**, 13 (2016).
4. Adl, S.M. *et al.* The revised classification of eukaryotes. *J Eukaryot Microbiol* **59**, 429-493 (2012).
5. Sebe-Pedros, A. *et al.* Insights into the origin of metazoan filopodia and microvilli. *Mol Biol Evol* **30**, 2013-2023 (2013).
6. Mooseker, M.S. & Tilney, L.G. Organization of an actin filament-membrane complex. Filament polarity and membrane attachment in the microvilli of intestinal epithelial cells. *J Cell Biol* **67**, 725-743 (1975).
7. Morales, E.A., Gaeta, I. & Tyska, M.J. Building the brush border, one microvillus at a time. *Curr Opin Cell Biol* **80**, 102153 (2023).
8. Crawley, S.W., Mooseker, M.S. & Tyska, M.J. Shaping the intestinal brush border. *J Cell Biol* **207**, 441-451 (2014).
9. Gaeta, I.M., Meenderink, L.M., Postema, M.M., Cencer, C.S. & Tyska, M.J. Direct visualization of epithelial microvilli biogenesis. *Curr Biol* **31**, 2561-2575 e2566 (2021).
10. Specian, R.D. & Neutra, M.R. The surface topography of the colonic crypt in rabbit and monkey. *Am J Anat* **160**, 461-472 (1981).
11. Ohta, K., Higashi, R., Sawaguchi, A. & Nakamura, K. Helical arrangement of filaments in microvillar actin bundles. *J Struct Biol* **177**, 513-519 (2012).
12. Postema, M.M., Grega-Larson, N.E., Neininger, A.C. & Tyska, M.J. IRTKS (BAIAP2L1) Elongates Epithelial Microvilli Using EPS8-Dependent and Independent Mechanisms. *Curr Biol* **28**, 2876-2888 e2874 (2018).
13. Croce, A. *et al.* A novel actin barbed-end-capping activity in EPS-8 regulates apical morphogenesis in intestinal cells of Caenorhabditis elegans. *Nat Cell Biol* **6**, 1173-1179 (2004).
14. Bretscher, A. & Weber, K. Villin: the major microfilament-associated protein of the intestinal microvillus. *Proceedings of the National Academy of Sciences of the United States of America* **76**, 2321-2325 (1979).
15. Bretscher, A. & Weber, K. Fimbrin, a new microfilament-associated protein present in microvilli and other cell surface structures. *J Cell Biol* **86**, 335-340 (1980).
16. Bartles, J.R., Zheng, L., Li, A., Wierda, A. & Chen, B. Small espin: a third actin-bundling protein and potential forked protein ortholog in brush border microvilli. *J Cell Biol* **143**, 107-119 (1998).
17. Morales, E.A., Arnaiz, C., Krystofiak, E.S., Zanic, M. & Tyska, M.J. Mitotic Spindle Positioning (MISP) is an actin bundler that selectively stabilizes the rootlets of epithelial microvilli. *Cell Rep* **39**, 110692 (2022).
18. Tyska, M.J. *et al.* Myosin-1a is critical for normal brush border structure and composition. *Mol Biol Cell* **16**, 2443-2457 (2005).
19. Liu, Y. *et al.* Constitutively active ezrin increases membrane tension, slows migration, and impedes endothelial transmigration of lymphocytes in vivo in mice. *Blood* **119**, 445-453 (2012).
20. Mogilner, A. & Rubinstein, B. The physics of filopodial protrusion. *Biophysical Journal* **89**, 782-795 (2005).
21. Theriot, J.A. The polymerization motor. *Traffic* **1**, 19-28 (2000).

669   22.   Kaksonen, M. & Roux, A. Mechanisms of clathrin-mediated endocytosis. *Nat Rev Mol*
670   *Cell Biol* **19**, 313-326 (2018).
671   23.   Jost, M., Simpson, F., Kavran, J.M., Lemmon, M.A. & Schmid, S.L. Phosphatidylinositol-
672   4,5-bisphosphate is required for endocytic coated vesicle formation. *Curr Biol* **8**, 1399-
673   1402 (1998).
674   24.   Merrifield, C.J., Qualmann, B., Kessels, M.M. & Almers, W. Neural Wiskott Aldrich
675   Syndrome Protein (N-WASP) and the Arp2/3 complex are recruited to sites of clathrin-
676   mediated endocytosis in cultured fibroblasts. *Eur J Cell Biol* **83**, 13-18 (2004).
677   25.   Winter, D., Podtelejnikov, A.V., Mann, M. & Li, R. The complex containing actin-related
678   proteins Arp2 and Arp3 is required for the motility and integrity of yeast actin patches.
679   *Curr Biol* **7**, 519-529 (1997).
680   26.   Kim, K. *et al.* Actin-based motility during endocytosis in budding yeast. *Mol Biol Cell* **17**,
681   1354-1363 (2006).
682   27.   Kubler, E. & Riezman, H. Actin and fimbrin are required for the internalization step of
683   endocytosis in yeast. *EMBO J* **12**, 2855-2862 (1993).
684   28.   Idrissi, F.Z., Wolf, B.L. & Geli, M.I. Cofilin, but not profilin, is required for myosin-I-
685   induced actin polymerization and the endocytic uptake in yeast. *Mol Biol Cell* **13**, 4074-
686   4087 (2002).
687   29.   Boulant, S., Kural, C., Zeeh, J.C., Ubelmann, F. & Kirchhausen, T. Actin dynamics
688   counteract membrane tension during clathrin-mediated endocytosis. *Nat Cell Biol* **13**,
689   1124-1131 (2011).
690   30.   Hyman, T., Shmuel, M. & Altschuler, Y. Actin is required for endocytosis at the apical
691   surface of Madin-Darby canine kidney cells where ARF6 and clathrin regulate the actin
692   cytoskeleton. *Mol Biol Cell* **17**, 427-437 (2006).
693   31.   Niggli, V., Andreoli, C., Roy, C. & Mangeat, P. Identification of a phosphatidylinositol-4,5-
694   bisphosphate-binding domain in the N-terminal region of ezrin. *FEBS Lett* **376**, 172-176
695   (1995).
696   32.   Kessels, M.M., Engqvist-Goldstein, A.E., Drubin, D.G. & Qualmann, B. Mammalian
697   Abp1, a signal-responsive F-actin-binding protein, links the actin cytoskeleton to
698   endocytosis via the GTPase dynamin. *J Cell Biol* **153**, 351-366 (2001).
699   33.   Da Costa, S.R. *et al.* Impairing actin filament or syndapin functions promotes
700   accumulation of clathrin-coated vesicles at the apical plasma membrane of acinar
701   epithelial cells. *Mol Biol Cell* **14**, 4397-4413 (2003).
702   34.   Haag, N. *et al.* The actin nucleator Cobl is crucial for Purkinje cell development and
703   works in close conjunction with the F-actin binding protein Abp1. *J Neurosci* **32**, 17842-
704   17856 (2012).
705   35.   Postema, M.M., Grega-Larson, N.E., Meenderink, L.M. & Tyska, M.J. PACSIN2-
706   dependent apical endocytosis regulates the morphology of epithelial microvilli. *Mol Biol*
707   *Cell* **30**, 2515-2526 (2019).
708   36.   Danielsen, E.M. Probing endocytosis from the enterocyte brush border using fluorescent
709   lipophilic dyes: lipid sorting at the apical cell surface. *Histochem Cell Biol* **143**, 545-556
710   (2015).
711   37.   Matter, K., Stieger, B., Klumperman, J., Ginsel, L. & Hauri, H.P. Endocytosis, recycling,
712   and lysosomal delivery of brush border hydrolases in cultured human intestinal epithelial
713   cells (Caco-2). *J Biol Chem* **265**, 3503-3512 (1990).
714   38.   Kolev, H.M. & Kaestner, K.H. Mammalian Intestinal Development and Differentiation-The
715   State of the Art. *Cell Mol Gastroenterol Hepatol* **16**, 809-821 (2023).
716   39.   Wood, L.A., Larocque, G., Clarke, N.I., Sarkar, S. & Royle, S.J. New tools for "hot-
717   wiring" clathrin-mediated endocytosis with temporal and spatial precision. *J Cell Biol*
718   **216**, 4351-4365 (2017).

719   40.   Loomis, P.A. *et al.* Espin cross-links cause the elongation of microvillus-type parallel
720   actin bundles in vivo. *J Cell Biol* **163**, 1045-1055 (2003).
721   41.   Onaya, T., Solomon, D.H. & Davidson, W.D. Effect of chlorpromazine on thyrotropin-
722   stimulated endocytosis and glucose oxidation in canine thyroid slices. *Endocrinology* **85**,
723   150-154 (1969).
724   42.   McCluskey, A. *et al.* Building a better dynasore: the dyngo compounds potently inhibit
725   dynamin and endocytosis. *Traffic* **14**, 1272-1289 (2013).
726   43.   Pelaseyed, T. & Bretscher, A. Regulation of actin-based apical structures on epithelial
727   cells. *J Cell Sci* **131** (2018).
728   44.   Meenderink, L.M. *et al.* Actin Dynamics Drive Microvillar Motility and Clustering during
729   Brush Border Assembly. *Dev Cell* **50**, 545-556 e544 (2019).
730   45.   Willy, N.M. *et al.* Membrane mechanics govern spatiotemporal heterogeneity of
731   endocytic clathrin coat dynamics. *Mol Biol Cell* **28**, 3480-3488 (2017).
732   46.   Romer, J. & Bickel, M.H. Interactions of chlorpromazine and imipramine with artificial
733   membranes investigated by equilibrium dialysis, dual-wavelength photometry, and
734   fluorimetry. *Biochem Pharmacol* **28**, 799-805 (1979).
735   47.   Jutila, A., Soderlund, T., Pakkanen, A.L., Huttunen, M. & Kinnunen, P.K. Comparison of
736   the effects of clozapine, chlorpromazine, and haloperidol on membrane lateral
737   heterogeneity. *Chem Phys Lipids* **112**, 151-163 (2001).
738   48.   Raucher, D. & Sheetz, M.P. Cell spreading and lamellipodial extension rate is regulated
739   by membrane tension. *J Cell Biol* **148**, 127-136 (2000).
740   49.   Raucher, D. & Sheetz, M.P. Characteristics of a membrane reservoir buffering
741   membrane tension. *Biophys J* **77**, 1992-2002 (1999).
742   50.   Raucher, D. & Sheetz, M.P. Membrane expansion increases endocytosis rate during
743   mitosis. *J Cell Biol* **144**, 497-506 (1999).
744   51.   Qualmann, B. & Kelly, R.B. Syndapin isoforms participate in receptor-mediated
745   endocytosis and actin organization. *J Cell Biol* **148**, 1047-1062 (2000).
746   52.   Antonescu, C.N., Aguet, F., Danuser, G. & Schmid, S.L. Phosphatidylinositol-(4,5)-
747   bisphosphate regulates clathrin-coated pit initiation, stabilization, and size. *Mol Biol Cell*
748   **22**, 2588-2600 (2011).
749   53.   Itoh, T. *et al.* Dynamin and the actin cytoskeleton cooperatively regulate plasma
750   membrane invagination by BAR and F-BAR proteins. *Dev Cell* **9**, 791-804 (2005).
751   54.   Ehrlich, M. *et al.* Endocytosis by random initiation and stabilization of clathrin-coated
752   pits. *Cell* **118**, 591-605 (2004).
753   55.   Nolen, B.J. *et al.* Characterization of two classes of small molecule inhibitors of Arp2/3
754   complex. *Nature* **460**, 1031-1034 (2009).
755   56.   Ferrary, E. *et al.* In vivo, villin is required for Ca(2+)-dependent F-actin disruption in
756   intestinal brush borders. *J Cell Biol* **146**, 819-830 (1999).
757   57.   Grimm-Gunter, E.M. *et al.* Plastin 1 binds to keratin and is required for terminal web
758   assembly in the intestinal epithelium. *Mol Biol Cell* **20**, 2549-2562 (2009).
759   58.   Pinson, K.I., Dunbar, L., Samuelson, L. & Gumucio, D.L. Targeted disruption of the
760   mouse villin gene does not impair the morphogenesis of microvilli. *Dev Dyn* **211**, 109-
761   121 (1998).
762   59.   Revenu, C. *et al.* A new role for the architecture of microvillar actin bundles in apical
763   retention of membrane proteins. *Mol Biol Cell* **23**, 324-336 (2012).
764   60.   Bouley, R., Yui, N., Terlouw, A., Cheung, P.W. & Brown, D. Chlorpromazine Induces
765   Basolateral Aquaporin-2 Accumulation via F-Actin Depolymerization and Blockade of
766   Endocytosis in Renal Epithelial Cells. *Cells* **9** (2020).
767   61.   Larsson, E., Moren, B., McMahon, K.A., Parton, R.G. & Lundmark, R. Dynamin2
768   functions as an accessory protein to reduce the rate of caveola internalization. *J Cell Biol*
769   **222** (2023).

62. Jin, M. *et al.* Branched actin networks are organized for asymmetric force production
during clathrin-mediated endocytosis in mammalian cells. *Nat Commun* **13**, 3578 (2022).
63. Burke, T.A. *et al.* Homeostatic actin cytoskeleton networks are regulated by assembly
factor competition for monomers. *Curr Biol* **24**, 579-585 (2014).
64. Loerke, D. *et al.* Cargo and dynamin regulate clathrin-coated pit maturation. *PLoS Biol* **7**,
e57 (2009).
65. Mettlen, M., Loerke, D., Yarar, D., Danuser, G. & Schmid, S.L. Cargo- and adaptor-
specific mechanisms regulate clathrin-mediated endocytosis. *J Cell Biol* **188**, 919-933
(2010).
66. Mundell, S.J., Luo, J., Benovic, J.L., Conley, P.B. & Poole, A.W. Distinct clathrin-coated
pits sort different G protein-coupled receptor cargo. *Traffic* **7**, 1420-1431 (2006).
67. Keyel, P.A. *et al.* A single common portal for clathrin-mediated endocytosis of distinct
cargo governed by cargo-selective adaptors. *Mol Biol Cell* **17**, 4300-4317 (2006).
68. Yu, Y. & Yoshimura, S.H. Self-assembly of CIP4 drives actin-mediated asymmetric pit-
closing in clathrin-mediated endocytosis. *Nat Commun* **14**, 4602 (2023).
69. Akamatsu, M. *et al.* Principles of self-organization and load adaptation by the actin
cytoskeleton during clathrin-mediated endocytosis. *Elife* **9** (2020).
70. Svitkina, T.M. *et al.* Mechanism of filopodia initiation by reorganization of a dendritic
network. *J Cell Biol* **160**, 409-421 (2003).
71. McMahon, H.T. & Boucrot, E. Membrane curvature at a glance. *J Cell Sci* **128**, 1065-
1070 (2015).
72. Dergai, M., Iershov, A., Novokhatska, O., Pankivskyi, S. & Rynditch, A. Evolutionary
Changes on the Way to Clathrin-Mediated Endocytosis in Animals. *Genome Biol Evol* **8**,
588-606 (2016).
73. Schmid, S.L., Sorkin, A. & Zerial, M. Endocytosis: Past, present, and future. *Cold Spring
Harb Perspect Biol* **6**, a022509 (2014).
74. de Duve, C. The origin of eukaryotes: a reappraisal. *Nat Rev Genet* **8**, 395-403 (2007).
75. Qian, H., Sheetz, M.P. & Elson, E.L. Single particle tracking. Analysis of diffusion and
flow in two-dimensional systems. *Biophys J* **60**, 910-921 (1991).
799

FIGURES

Figure 1. Endocytic proteins are apically localized in the crypts. (A) Scanning electron microscopy (SEM) of native mouse small intestine along full crypt-villus axis. Scale bar = 50 μm. (B) High magnification of the small intestinal crypt (scale bar = 10 μm) with zooms (base, middle, and transition (scale bars = 1 um (solid box zooms) and 500 nm (dashed box zooms)) showing the accumulation of microvilli along the transit amplifying zone. (C) Native mouse small intestine stained for villin, dynamin2, and pacsin2. Panels show single z-plane confocal image of intestinal crypts. Scale bar = 10 μm. (D) Mean brush border intensity measurements for each marker at the base of the crypt (magenta) and at the crypt-villus transition zone (green). n = 20-25 crypts per marker, 3 WT mice. Multiple Mann-Whitney tests; ****p = <0.0001, **p = 0.002.

Figure 2. Induction of clathrin-mediated endocytosis depletes microvilli. (A) Cartoons depicting constructs used to induce clathrin-mediated endocytosis. Numbers represent amino acids. (B) Schematic depicting the rapalog-inducible system to dock the β2 subunit of AP-2 on the plasma membrane to induce endocytosis. (C) Confocal MIP of CL4 cell expressing Halo-espin (shown), CD8-FRB (not shown), and GFP-FKBP (not shown) or FKBP-β2-GFP (not shown) before and after 45 min rapalog incubation. Scale bar = 5 μm. (D) Schematic showing area under the curve (AUC) quantification. (E) Fold change in AUC of Halo-espin signal of microvilli post- vs. pre- rapalog induction. Area under the curve (AUC) was calculated as shown in D. 11-12 cells per condition taken from at least three experimental replicates, 120-150 microvilli per condition. Mann-Whitney test; ****p =<0.0001. (F) Fold change in area of microvillar coverage before and after 45 min rapalog incubation. N = 11-13 cells per condition, taken from at least three experimental replicates. Mann-Whitney test; *p=0.0218.

Figure 3. Inhibition of endocytosis results in exaggerated microvilli. (A) Schematic depicting chlorpromazine and dyngo mechanisms of action. (B) Confocal maximum intensity projection (MIP) of representative CL4 cells at 0 min and 45 min post DMSO (left), chlorpromazine (CPZ) (middle), and dyngo (right) treatment. Drug was added immediately after 0 min image was captured. Scale bar = 5 µm. (C) Fold change in length of microvilli post- vs. pre- drug treatments. Microvillar length was measured using mCherry-espin signal. n = 13-16 cells from at least three experimental replicates, 150-200 microvilli per condition. Kruskal-Wallis test with multiple comparisons; ****p = <0.0001. (D) Fold change in area under the curve of mCherry-espin signal of microvilli post- vs. pre- drug treatments. 14-16 cells from at least three experimental replicates, 150-200 microvilli per condition. Kruskal-Wallis test with multiple comparisons; ****p = <0.0001. (E) Confocal MIP of representative CL4 cells expressing mCherry-espin (not shown) and mNeonGreen-βActin at 0 min and 45 min post-DMSO or CPZ treatment. Scale bar = 5 µm. (F) Fold change in area under the curve of mNeonGreen-βActin signal of microvilli post- vs. pre- drug treatments. n = 15 cells from at least three experimental replicates, 150-200 microvilli per condition. Mann-Whitney test; ****p = <0.0001.

Figure 4. Exaggerated microvilli that assemble post-CPZ treatment retain canonical features. (A) Confocal maximum intensity projection (MIP) of CL4 cells expressing mCherry-espin stained with CellMask-DeepRed. Cells were first treated with either DMSO or CPZ, then incubated with CellMask for 15 minutes. Scale bar = 5 μm. (B) Superplot quantification of the length of membrane coverage along the microvillus after DMSO or CPZ treatment. Small, transparent points represent individual microvilli, larger opaque points show whole cell averages. (C, E, G) Confocal MIP of CL4 cells expressing mCherry-espin and (C) EGFP-ezrin, (E) EGFP-EPS8, and (E) EGFP-IRTKS at 0 min and 45 min. Single channel zooms show intensity-coded color profile. Intensity scales from low (dark blue) to high (yellow). Scale bar = 3 μm. (D) Quantification of ezrin AUC before and after CPZ treatment. N = 11 cells, 130-160 microvilli per condition. Mann Whitney U test; ***p =0.0002. (F,H) Superplot quantification of (F) Eps8 or (H) IRTKS intensity pre/post CPZ treatment. Small, transparent data points are individual puncta and larger, opaque data points are individual cells. N = 13-15 cells, 4000-6000 microvilli per condition. Mann Whitney U test; ***p = 0.0005.

Figure 5. Exaggerated microvilli exhibit highly persistent motility. (A) Representative montage of microvillar motility and elongation before (left) and after (right) CPZ treatment. Single channels show intensity-coded color profile. Intensity scales from low (white) to high (yellow). Column width = 2 μm. (B) (left) Rose plot of 50 microvillar tracks over 10 min (μm units) in cells pre CPZ treatment. B. (right) 16 representative tracks isolated from B. Scale bar = 1 μm. (C) (left) Rose plot of 50 microvillar tracks over 10 min (μm units) in cells post CPZ treatment. C. (right) 16 representative tracks isolated from C. Scale bar = 1 μm. (D) Mean squared displacement analysis of 100 microvillar tracks imaged for 20 minutes every 1 minute. (E-G). Superplot quantification of persistence, displacement, and velocity, respectively in untreated and CPZ treated cell populations. Only tracks lasting exactly 10 minutes were analyzed. Small transparent points show individual microvilli, larger opaque points show per cell averages. n = 10-12 cells sampled from at least three independent experiments, 60-100 microvilli per cell. Mann Whitney test; P* = 0.0439.

Figure 6. Microvilli grow from apical clathrin coated pits. (A) Montage of a single microvillus growth event in CL4 cells expressing EGFP-LCA and mCherry-Espin. Every column represents 30 seconds. Arrows mark the analyzed core bundle. Column width = 2.09 µm. (B) Average fluorescence intensity measurements over time. n = 21 growth events from 8 cells across 3 independent experiments. (C) Montage of a single microvillus growth event in CL4 cells expressing mCherry-Dynamin2 and GFP-Espin. Every column represents one minute. Arrows mark the analyzed core bundle. Column width = 2.09 µm. (D) Average fluorescence intensity measurements over time. n = 10 growth events from 4 cells across 3 independent experiments. (E) Scanning electron microscopy image of 0-DPC CL4 cells. Scale bar = 10 µm. Zooms 1 and 2 noted by bounding boxes. Zoom scale bar = 2 µm. (F) Scanning electron microscopy image of 0-DPC CACO-2$_{BBE}$ cells. Scale bar = 2 µm. Zooms 1 and 2 noted by bounding box. Zoom scale bar = 1 µm. (G) Nearest neighbor distance measurements of from microvilli to closest endocytic pits on the surface of CL4 cells. Blue line denotes simulated nearest neighbor distances assuming random distribution of the same number of microvilli and endocytic pits across the same cell surface area. (H). Length of microvilli associated (<250 nm) vs not associated (>300 nm) with endocytic pits on the surface of CL4 cells. N = 175 microvilli in >300 category and 95 microvilli in <250 nm category. Mann Whitney test ****p = <0.0001. (I) Nearest neighbor distance measurements of from microvilli to closest endocytic pits on the surface of CACO-2$_{BBE}$ cells. Blue line denotes simulated nearest neighbor distances assuming random distribution of same number of microvilli and endocytic pits across the same cell surface area.

Figure 7. Arp2/3 complex inhibition decreases microvillar growth. (A) Confocal maximum intensity projection (MIP) of representative CL4 cells at 0 min and 45 min post DMSO (left) or 100 µM CK666. Cartoons beneath MIPs highlight growth events seen in Supplemental movie 2, color coded according to when growth event occurred during 30 minute imaging window. (B) Fold change in the rate of microvillar growth initiation. To define a fold change value for cells that had zero growth events following CK666 treatment, a value of 0.001 was added to all measurements. N= 5-6 cells per condition. Mann Whitney test **p = 0.0043. (C) Confocal MIPs of representative CL4 cells at 0 min and 90 min post DMSO (left) or 60 µM CK666 (right). Bounding boxes denote zoom insets. (D) Fold change in mean apical espin intensity of medial (middle) and marginal (junctional) microvilli. N = 23 cells in DMSO condition and 35 cells in CK666 condition. Ordinary one-way ANOVA with Šídák's multiple comparisons test; ****p = <0.0001

Figure 8. Schematic depicting proposed model of clathrin mediated endocytosis and microvillar biogenesis sharing materials during epithelial cell differentiation.

SUPPLEMENTAL DATA

Movie S1, related to Figure 3. Inhibition of endocytosis results in exaggerated microvilli. Live cell movie of maximum intensity projection (MIP) confocal time series of 3 CL4 cells expressing mCherry-espin. CPZ was added at T = 30 min. Time hour: min: sec. Single channels show intensity-coded color profile. Intensity scales from low (white) to high (yellow). Scale bar = 5 µm.

Movie S2, related to Figure 7. Arp2/3 complex is required for microvillar growth. Live cell movie of maximum intensity projection (MIP) of confocal time series of 2 CL4 cells expressing mCherry-Espin. DMSO (left) or 100 µM CK666 (right) was added at T = 30 min. Cyan spots which accumulate highlight microvillar growth events. Images were captured at 1 frame per 1 minute. Time min: sec. Single channels show intensity-coded color profile. Intensity scales from low (white) to high (yellow).

Figure S1. **CPZ treatment phenotype is titratable and robust across model systems.** (A) Confocal maximum intensity projection (MIP) of representative CL4 cells at 0 min and 45 min after varying concentrations of CPZ treatment. Drug was added immediately after 0 min image was captured. Single channels show intensity-coded color profile. Intensity scales from low (white) to high (yellow). Scale bar = 5 µm. (B) Fold change in area under the curve of mCherry-espin signal of microvilli post- vs. pre- CPZ treatment. 8-10 cells from at least three experimental replicates, 100-150 microvilli per condition. Images and data for the 30 µM CPZ treatment were taken from figure 3. (C) MIPs of 3 representative $CACO_2$-BBE cells at 0 min and 45 min after treatment with 30 µM CPZ. Drug was added immediately after 0 min image was captured. Single channels show intensity-coded color profile. Intensity scales from low (white) to high (yellow). Scale bar = 5 µm.

Figure S2. **CPZ and dyngo equally inhibit transferrin uptake.** (A) Maximum intensity projection (MIP) of CL4 cells transfected with mCherry-espin incubated with human purified transferrin after 45-minute treatment of DMSO, 30 μM CPZ, or 30 μM Dyngo. Scale bar = 10 μM (B) Superplot quantification of whole cell mean transferrin intensity measurements. Small transparent points represent individual cells, larger opaque points show experimental replicates. n = 100 cells sampled from two experimental replicates.

Figure S3. Global reduction in plasma membrane tension does not lead to the formation of exaggerated microvilli. (A) Confocal maximum intensity projection (MIP) of representative CL4 cells before and after treatment with CPZ, 1% EtOH, 400 µM Deoxycholate, and 125 mM NaCl. Cells were treated with CPZ and Deoxycholate for 45 minutes, and with 1% EtOH and 125 mM NaCl for 5-10 minutes. Drug was added immediately after 0 min image was captured. Single channels show intensity-coded color profile. Intensity scales from low (white) to high (yellow). Scale bar = 5 µm. (B) Fold change in area under the curve of mCherry-espin signal of microvilli post- vs. pre- drug treatments. 8-10 cells from 2 experimental replicates, 100-150 microvilli per condition. Images and data for the 30 µM CPZ treatment were taken from figure 3.

Supplemental Figure 4. Microvilli grow from apical clathrin coated pits. (A) Montage of a single microvillus growth event in CL4 cells expressing EGFP-mAbp1 and mCherry-Espin. Every column represents 30 seconds. Arrows mark the analyzed core bundle. Column width = 2.09 µm. (B) Average fluorescence intensity measurements over time. n = 9 growth events from 4 cells across 2 independent experiments. (C) Montage of a single microvillus growth event in CL4 cells expressing EGFP-pPLC-ΔPH-GFP (PIP(4,5)P₂ marker) and mCherry-Espin. Every column represents 30 seconds. Arrows mark the analyzed core bundle. Column width = 2.09 µm. (D) Average fluorescence intensity measurements over time. n = 20 growth events from 10 cells across 3 independent experiments. (E) Montage of a single microvillus growth event in CL4 cells expressing EGFP-pacsin2 and mCherry-espin. Every column represents one minute. Arrows mark the analyzed core bundle. Column width = 2.09 µm. (F) Average fluorescence intensity measurements over time. n = 20 growth events from 10 cells across 3 independent experiments.

Figure S5. Simulation of randomly distributed endocytic pits and microvilli leads to nearest neighbor distances with gaussian distributions. (A) 10 runs of simulated nearest neighbor distance measurements from endocytic pits to nearest microvillus on the surface of CL4 cells. Simulated densities were matched with experimental measurements of 516 pits, 2570 microvilli over 1601.25 μm^2. (B) 10 runs of simulated nearest neighbor distance measurements from endocytic pits to nearest microvillus on the surface of CACO$_2$-BBE cells. Simulated densities were matched with experimental measurements of 415 pits, 2039 microvilli over 1828.7 μm^2.