A circuit involving the lncRNA MB3 and the driver genes MYC and OTX2 inhibits apoptosis in Group 3 Medulloblastoma by regulating the TGF-β pathway via HMGN5

Grandioso Alessia[1], Tollis Paolo[2], Pellegrini Francesca Romana[3], Falvo Elisabetta[3], Palma Alessandro[1], Migliaccio Francesco[4], Belvedere Alessandro[1], Rea Jessica[2], Tisci Giada[1,5], Carissimo Annamaria[6], Bozzoni Irene[1,2], Trisciuoglio Daniela[3], Ballarino Monica[1], Ceci Pierpaolo[3], Laneve Pietro[1,3,#]

1. Department of Biology and Biotechnologies Charles Darwin, Sapienza University of Rome, Piazzale Aldo Moro 5, 00185, Rome, Italy.
2. Center for Life Nano-& Neuro-Science, Fondazione Istituto Italiano di Tecnologia, Viale Regina Elena 291, 00161, Rome, Italy.
3. Institute of Molecular Biology and Pathologies, National Research Council of Italy, Piazzale Aldo Moro 7, 00185, Rome, Italy.
4. Department of Electrical Engineering and Information Technology, University Federico II, Via Claudio 21, 80125, Naples, Italy.
5. Department of Biochemical Sciences, Sapienza University of Rome, Piazzale Aldo Moro 5, 00185, Rome, Italy.
6. Institute for Applied Mathematics "Mauro Picone", National Research Council of Italy, Via Pietro Castellino 111, 80131 Naples, Italy.

correspondence to: Pietro Laneve, IBPM-CNR, c/o Sapienza University of Rome, Piazzale Aldo Moro 5, 00185, Rome, Italy. E-mail: pietro.laneve@cnr.it; Phone: +390649912205

Running title: *LncMB3* controls TGF-β pathway in Medulloblastoma

ABSTRACT

Background. Group 3 (G3) is one of the most common, aggressive and fatal subtypes of the paediatric cerebellar tumour Medulloblastoma (MB), primarily driven by the MYC oncogene. Targeting MYC has long been challenging and this, combined with our incomplete understanding of G3 MB molecular bases, has hindered the development of effective targeted therapies. Long noncoding RNAs (lncRNAs), with their extensive oncogenic roles, cancer-specific expression, and connection to MYC biology, offer opportunities for unravelling this complexity and providing new insights and therapeutic targets.

Methodology. Using genome-wide, molecular and cellular assays, we characterised the activity of the MYC-dependent, anti-apoptotic lncRNA *lncMB3* in G3 MB cells.

Results. Through transcriptomic and interactomic analyses, we clarified *lncMB3* function and mode-of-action. *LncMB3* controls the TGF-β pathway, critically altered in G3 medulloblastomagenesis. This regulation occurs via the direct coding-noncoding RNA interaction between *lncMB3* and the mRNA for the epigenetic factor HMGN5, with both sharing targets in the TGF- β cascade. This axis converges on apoptosis through OTX2, another G3 MB driver gene, and photoreceptor lineage genes. Synergistic effects between *lncMB3* targeting and cisplatin treatment underscores the relevance of this regulatory network *in vitro*. Finally, we propose novel ferritin-based nanocarriers as efficient delivery tools for antisense oligonucleotides targeting *lncMB3*.

Conclusions. *LncMB3* emerges as a central node linking MYC amplification to apoptosis inhibition through a circuit involving RNA-based mechanisms, G3 MB key drivers and underexplored factors.

47 This integrated framework deepens our understanding of G3 MB molecular underpinnings and lay
48 the foundation for translating lncRNA research into potential applications.
49

INTRODUCTION

Medulloblastoma (MB) is the most prevalent malignant paediatric tumour in the central nervous

system, accounting for ~20% of all brain cancers in children [1], [2]. Although all MBs originate in the

cerebellum, share histomorphological features, and possess signatures of embryonic cerebellar

lineages [3], they exhibit high heterogeneity. MB is classified into 4 subtypes with distinct biological,

clinical and therapeutical implications [4]: Wingless (WNT), Sonic Hedgehog (SHH), Group 3 (G3) and

Group 4 (G4). G3 and G4, collectively, account for 60% of MB cases and represent the most

aggressive subtypes, posing significant treatment challenges due to their poor biochemical

characterisation. Understanding the molecular mechanisms behind these subgroups is critical for

advancing precision medicine strategies. Specifically, G3 MB is associated with a dismal prognosis,

often presenting with metastases at diagnosis and exhibiting high recurrence rates [2]. It is driven

primarily by the oncogenic transcription factor (TF) MYC, whose gene copy number amplification or

elevated expression are distinctive of this subtype [5]. Targeting MYC and its related

pathways/complexes is essential for managing G3 MB, despite the challenges associated with

addressing canonical MYC circuits [6].

The discovery of novel classes of noncoding RNAs (ncRNAs) is expanding our understanding of

genome output and function [7], offering new avenues for cancer therapeutics. Among ncRNAs, the

highly versatile long noncoding RNAs (lncRNAs) have emerged as key regulators of cancer hallmarks

[8], showing potential as biomarkers and targets due to their cell-specific expression and regulatory

activities. Over the years, we have unravelled the roles of ncRNAs in various nervous system cancer

models, including MB [9]. Recently, we identified in G3 MB a class of MYC-dependent lncRNAs, with

three of them showing a potential function as oncogenes [10], including *lncMB3*.

In this study, we present a detailed functional analysis of *lncMB3*, which is overexpressed in the tumour condition and acts as anti-apoptotic factor. We show that *lncMB3* regulates the TGF-β pathway, a key player known to be altered in G3 MB [11]. This is achieved by a functional interaction with the mRNA of the epigenetic factor HMGN5, whose protein level is upregulated and impacts on a subset of TGF-β pathway genes also affected by *lncMB3* targeting. Notably, coding-noncoding RNA interplays are increasingly recognised as important mechanisms in cancer biology. This regulatory axis influences apoptosis through OTX2, an additional crucial G3 MB driver gene [12], and its downstream cascade. The relevance of this circuit is underscored by the synergistic effects observed when *lncMB3* targeting is combined with cisplatin administration in G3 MB cells. Lastly, we demonstrate that recombinant human ferritin-based nanovectors, a promising platform for drug administration in disease conditions [13], serve as effective carriers for delivering antisense oligonucleotides (ASOs) targeting *lncMB3* in G3 MB cells.

RESULTS

Identification of *lncMB3*-dependent transcriptome in G3 MB cells

To identify the molecular network downstream of *lncMB3* in MB, we performed a transcriptome analysis following its knockdown (KD) in the MYC-amplified D283 Med cell line where the lncRNA is highly expressed (Fig. S1A) and initially identified [10]. According to GTEx (https://gtexportal.org/home/), *lncMB3* is minimally expressed in healthy tissues (Fig. S1B). The KD of *lncMB3* reduced its expression by ~70% (Fig. S2A) 72 hours after multi-pulse transfections of a Locked Nucleic Acid (LNA)-based ASO [14], named GapmeR #1, which targets the lncRNA at its 5' region. Initial transcriptional data analysis yielded positive results regarding the number of detected genes (Fig. S2B, left panel), clustering of transcriptomes (Fig. S2B, right panels) and consistency between *lncMB3* KD normalised read counts and previous qRT-PCR assessments (Dataset 1).

RNA-Seq analysis (Dataset 1) identified 14.655 transcripts expressed in at least one sample, and

2.995 differentially expressed genes (DEGs) between *lncMB3* KD and control (SCR), with an FDR <

0,05 (Fig. 1A, left panel). Among these, 1.395 were downregulated and 1.600 were upregulated

following *lncMB3* silencing (Fig. 1A, right panel, blue and red dots). A dozen of these upregulated or

downregulated genes were randomly selected, with padj ranging from 0,001 to 0,05, for qRT-PCR

validation, which confirmed RNA-Seq expression data (Fig. S2C and Dataset 1).

To functionally explore the transcriptomic landscape, DEGs were filtered using quantitative criteria,

such as fold change (FC) robustness, statistical significance, and expression levels (Fig. S2D). We

identified 13 RNAs regulated by *lncMB3* (12 protein-coding transcripts and 1 lncRNA, being 7

significantly downregulated and 6 upregulated RNAs, Dataset 2). qRT-PCR validated RNA-Seq data

trends for the selected DEGs, demonstrating statistical significance (for 10 out of 13 genes) or, at

least, discernible variance trends (Fig. S3A). Notably, the top gene identified was *INHBE*, a

component of the TGF-β pathway [15], associated with G3 MB pathogenesis [11].

Gene Ontology (GO) analysis was performed on DEGs to identify the full set of gene functions

affected by *lncMB3* depletion (Fig. 1B, left panel). Concerning the ontological annotations, among

the top categories, GO revealed pathways such as "PIK3-Akt signalling" known to affect

chemosensitivity in various cancers, including MB [16], and "MAPK signalling" possibly reflecting

DNA damage response to apoptosis induced by *lncMB3* KD. Our focus, however, was again on the

TGF-β pathway, which emerged as the second top gene category and is deregulated in G3 MB

compared to other subgroups. Moreover, the TGF-β pathway was a category present in both the up

and downregulated gene sets (Fig. S3B), suggesting deregulation of activatory and inhibitory

effectors in the same pathway. Finally, Gene Set Enrichment Analysis (GSEA) further confirmed the

TGF-β pathway as significantly affected by *lncMB3* KD in G3 MB cells (Fig. S3C).

LncMB3 regulates the TGF-β pathway in G3 MB

Aberrant TGF-β signalling plays a key role in the development of various diseases, including cancer

[17]. In G3 MB, its deregulation is believed to drive processes such as cell proliferation,

differentiation, and survival [11].

The importance of this pathway in our system was further highlighted by filling the category with

additional DEGs involved in the pathway. We finally identified a set of 9 genes, including TGF-

β/Activin signalling ligands (*INHBE*, [15]; *THBS1*, *THBS3*, [18]), receptors (*ACVR1B*,[19]; *BAMBI*, [20]),

effectors (*SMAD3*, *SMAD6*, *SMAD7*, [19]), and downstream regulators (*TFDP1*, [21]) (Fig. S3D). qRT-

PCR upon *lncMB3* KD revealed the repression of genes with activatory roles in the pathway (*THBS1*,

THBS3, *INHBE*, *ACVR1B*, *SMAD3*, *SMAD6*, *SMAD7*), with downregulations ranging from 20% to 80%

(Fig. 1B, right panel). Conversely, inhibitory components (*BAMBI* and *TFDP1*) were upregulated

under the same conditions (1.5- to 3-fold, Fig. 1B, right panel), suggesting a coherent contribution

of both positive and negative regulators that leads to an overall downregulation of the TGF-β/Activin

branch. To ensure specificity in targeting, we employed an additional GapmeR (GapmeR #2), which

recognises the 3' end of *lncMB3* (Fig. S4A). This achieved a 70% depletion of the transcript in D283

Med cells (Fig. S4B). To further corroborate these findings, we also used GapmeR #2 to target *lncMB3*

in HD-MB03, a G3 MB cell line where MYC and *lncMB3* are upregulated (Fig. S1A). This led to a 50%

reduction of the lncRNA levels (Fig. S4B). Validation data showed similar expression alterations in all

the condition tested for 5 TGF-β network genes: *BAMBI*, *ACVR1B*, *INHBE*, *SMAD3*, and *SMAD6* (Fig.

1C). Since GapmeR #1 targets the 5' end of *lncMB3* and results in a more efficient downregulation

(Fig. S4B), it was preferentially used in subsequent experiments. Overall, the RNA-Seq analysis clearly

points to a regulatory role of *lncMB3* in the TGF-β pathway in G3 MB cells.

Insights into apoptosis regulation by *lncMB3*

In the developing nervous system, a relevant target of the TGF-β cascade is the brain-specific TF

OTX2, recognised as a G3 MB-enriched oncogene [12] and upregulated in both D283 Med and HD-

MB03 cells (Fig. S1A). During retinal development, OTX2 occupies a pivotal position in a hierarchy of

photoreceptor lineage genes, including the basic leucine zipper TFs *NRL* and *CRX* [22]. These factors

are overexpressed in G3 MB, where they mediate cancer survival by inducing the expression of the

anti-apoptotic factor *BCL2L1* ([23] and Fig. S3D).

We queried an available transcriptomic dataset (GSE164677) from 59 patient-derived samples

representing all MB subgroups and 4 control samples to evaluate the expression of these genes in

primary MB tumours. The expression of *NRL*, *CRX*, and *BCL2L1* was significantly upregulated in G3

compared to either healthy tissues or other MB subgroups. Although the upregulation of *OTX2* in

G3 MB compared to cerebellum samples did not reach statistical significance, it exhibited an upward

trend (Fig. 2A). Figure 2B (left panel) outlines the connections among these factors. Consistent with

this model, photoreceptor gene program was categorised in several downregulated annotations in

lncMB3 KD-dependent GSEA (Fig. 2B, middle panel).

Given this context, we observed the downregulation of *OTX2*, *NRL*, *CRX*, and *BCL2L1* RNAs in D283

Med cells following *lncMB3* depletion (Fig. 2B, right panel). Protein downregulation was also

assessed for OTX2 and BCL2L1 (Fig. 2C), as well as for PARP-1 (Fig. 2D) [24], whose cleavage (C-PARP)

is a hallmark of apoptosis. The involvement of the lncRNA in this axis was further confirmed using

GapmeR #1 in HD-MB03 cells, where *lncMB3* depletion led to i) decreased RNA and protein levels

for OTX2, NRL, CRX, and BCL2L1 (Fig. 3A and B); ii) reduced cell viability (Fig. 3C); and iii) coherent

modulation of C-PARP and PARP-1 levels (Fig. 3D).

Consistent with *lncMB3* being regulated by MYC [10], the *lncMB3*-dependent RNA alterations

highlighted here were also observed in the transcriptome of MYC-inhibited D283 Med cells *vs*

untreated cells (Dataset 3). To further demonstrate that *lncMB3* and the TGF-β pathway are part of

the same regulatory network, we utilised a TGF-β Receptor 1 Kinase inhibitor, which prevents

Smad2/3 phosphorylation. After 48 hours of treatment, we observed a dose-dependent reduction

in the number of viable cells (Fig. 3E). This cell phenotype was correlated with increased apoptosis,

as confirmed through the Annexin V binding assay (Fig. 3F). This shows that direct inhibition of the

TGF-β cascade leads to effects converging with those observed upon *lncMB3* silencing.

To definitively establish the function of *lncMB3* in counteracting apoptosis, we performed a rescue

assay in G3 MB. We generated a plasmid carrying the lncRNA sequence lacking the first 16

nucleotides (Δ16), which are necessary for recognition by GapmeR #1 (Fig. 3G, left panel). When

overexpressed in D283 Med cells along with the ASO, this construct successfully expressed a mutant

lncRNA derivative (Fig. 3G, middle panel), and restored cell viability compared to cells transfected

with GapmeR #1 alone (Fig. 3G, right panel).

These findings allow us to reconstruct a molecular network initiated by MYC amplification and

involving a non-coding RNA, which leads to the suppression of apoptosis in G3 MB.

Insights into the RNA interactome of *lncMB3*

RNA-RNA pairing is a crucial mechanism for lncRNA functionality. Given this and *lncMB3*

nucleocytoplasmic localisation in MB cells, we investigated whether the lncRNA could modulate the

expression of TGF-β and apoptotic pathway genes through such interactions.

Initially, we hypothesized a straightforward interaction between *lncMB3* and candidate mRNAs

downregulated upon its KD. However, preliminary analyses using the IntaRNA algorithm

(http://rna.informatik.uni-freiburg.de/), which maps transcript binding regions, yielded low

interaction propensity (Table 1). To experimentally explore possible indirect associations, we

performed RNA pull-down assays as in [25] (Fig. 4A). qRT-PCR analysis comparing *lncMB3*-enriched

pull-down fractions with EVEN and ODD probe sets with mock-precipitated LacZ fractions (Table 2)

showed no significant enrichment for any tested candidate (Table 3), excluding molecular

interactions between *lncMB3* and transcripts of the TGF-β and downstream pathways.

We then focused on the competing endogenous RNA network, where long RNAs modulate

microRNA activity by sponging them [26]. Preliminary interactions between *lncMB3* and the core

RNA-induced silencing complex component Argonaute 2 (AGO2) were predicted via the *cat*RAPID

algorithm (http://s.tartaglialab.com/page/catrapid_group) (Dataset 4) and experimentally validated

through a CLIP assay (Fig. S5A). Based on these results, we employed Targetscan

(https://www.targetscan.org/vert_80/) to identify microRNAs predicted to bind both *lncMB3* and

candidate mRNAs (Table 4). Among the six candidates (*hsa-miR-135b-5p*, *hsa-miR-216a-3p*, *hsa-miR-*

3681-3p, *hsa-miR-613*, *hsa-miR-1197*, *hsa-miR-196b-5p*), the top three were tested by qRT-PCR on

lncMB3 pull-down fractions, but none of them showed significant enrichment, ruling out

competition for microRNA association in the context investigated (Table 5).

An alternative scenario suggests that *lncMB3* could regulate its target genes via interactions with

other mRNAs [27]. To explore this, we identified *lncMB3* interactome crossing RNA-Seq data from

replicate independent *lncMB3* pull-down experiments (Fig. 4B and Dataset 5) and performing a

stringent analysis to highlight the predominant binding partners. We found two mRNAs (*HMGN5*,

EIF5B) and one lncRNA (*ANKDDA1*) associated to *lncMB3* (Fig. 4B and Fig. S6A), with the mRNA for

the nuclear factor HMGN5 (High Mobility Group Nucleosome Binding Domain 5) showing the highest

FC and the most significant FDR. Given the broad epigenetic role of HMGN5 in transcriptional

regulation [28], its interplay with *lncMB3* underlined this factor as a candidate for further analysis.

LncMB3 interacts with HMGN5 mRNA

The chromatin factor HMGN5 regulates gene expression by modulating interactions between

nucleosomes and DNA [28], [29]. *HMGN5* is expressed ubiquitously (Fig. S6B), and, among brain

tissues, its expression is low in cerebellum. Interestingly, RNA-Seq data from the primary MB dataset GSE164677 revealed *HMGN5* upregulation, with a more pronounced tendency in G3 (Fig. 4C).

Co-enrichment of *HMGN5* mRNA with *lncMB3* was validated in an additional pull-down assay (Fig. 4D). IntaRNA predicted an interaction site (energy score: -30,1 kcal/mol) spanning ~120 nucleotides over the 5' region of *lncMB3* (nt 189-307) and the coding sequence of *HMGN5* mRNA (nt 933-1059) (Fig. 4E). This prediction was experimentally verified using a 4'-aminomethyl-4,5',8-trimethylpsoralen (AMT)-crosslinked RNA pull-down assay [25], which highlights direct RNA-RNA pairings *in vivo* (Fig. 4F). Interestingly, digital PCR-based absolute quantifications of *lncMB3* and *HMGN5* abundance showed that their expression levels were stoichiometrically equivalent (Fig. S6C).

To assess the functional consequences of this interaction, we investigated whether *lncMB3* affects *HMGN5* expression. *HMGN5* mRNA levels were unchanged by *lncMB3* KD, as indicated by RNA-Seq data (Dataset 1) and confirmed by qRT-PCR (Fig. S7A). However, HMGN5 protein levels decreased by ~80% upon *lncMB3* KD (Fig. 4G), consistently with the oncogenic activity attributed to HMGN5 in a spectrum of cancers [30], [31]. Rescue of *lncMB3* levels through transient exogenous expression of Δ16 construct (Fig. 3G, left panel) showed a recovery trend for HMGN5 levels (Fig. 4G), indicating that HMGN5 protein levels are regulated by *lncMB3* in a specific, quantitative manner.

The impact of *lncMB3* on HMGN5 protein levels, along with their RNA interaction, indicates a role for the lncRNA as a translational activator. This was confirmed using a Ribo-Tag strategy [32] with transient expression of a FLAG-tagged version of the 60S ribosomal protein L22 (RPL22) in D283 Med cells (Fig. S7B). Western blot analysis of FLAG-RPL22-precipitated fractions showed comparable levels of FLAG-RPL22 in gap_SCR *vs* gap_*lncMB3* protein extracts (Fig. S7C). At variance, qRT-PCR analysis showed a consistent decrease of *HMGN5* mRNA association to ribosomes in the absence of

lncMB3, compared to *GAPDH* as a control (Fig. 4H). These results indicate that *lncMB3* is required

for *HMGN5* mRNA association with ribosomes.

HMGN5 regulates the TGF-β pathway and apoptosis in G3 MB cells

Given HMGN5 documented activity in the nucleus, we re-examined the expression of genes altered

by *lncMB3* KD to assess whether their up or downregulation was primarily due to transcriptional

regulation. qRT-PCR analysis was performed using exonic primers *vs* primers spanning exon-intron

junctions, to distinguish mature mRNAs from primary transcripts. As shown in Fig. S7D, *lncMB3*

silencing led to the downregulation of *ACVR1B*, *INHBE*, *SMAD3*, *OTX2*, *NRL*, and *CRX* pre-mRNAs,

highlighting a nuclear control over several genes regulated by *lncMB3* (compare Figs 1B, 1C, 2B, 3A).

A role for HMGN5 in suppressing apoptosis had not previously been assessed in MB. To establish a

functional connection between *lncMB3* and HMGN5 activities, we sought to mimic the molecular

effects of *lncMB3* KD by silencing *HMGN5*. We designed a short interfering RNA targeting *HMGN5*

with VectorBuilder (https://en.vectorbuilder.com/). Transient transfection of this construct in D283

Med cells resulted in a ~30% downregulation of *HMGN5* mRNA (Fig. 4I) and a ~60% reduction in

protein levels (Fig. S7E). This led to a 20-50% RNA decrease for *INHBE*, *SMAD3*, *OTX2*, and *BCL2L1*

(Fig. 4I), which are among the targets identified upon *lncMB3* KD (see Figs 1B, 1C, 2B, 3A). This

indicates that the achieved levels of *HMGN5* KD appreciably phenocopy the molecular effects of

lncMB3 downregulation (Fig. S7F). Regarding cell death, after *HMGN5* KD we also observed a

threefold increase in c-PARP1 protein (Fig. 4J, left panel) and a 30% decrease in full-length PARP1

(Fig. 4J, right panel). Reduction in HMGN5 levels also phenocopied the decrease in the number of

viable D283 Med cells, with a drop of ~30% (Fig. 4K), similarly with the effects of *lncMB3* targeting.

Overall, these findings indicate that *lncMB3* and HMGN5 functionally participate into the same

molecular and cellular phenotypes. Interacting with *lncMB3*, *HMGN5* mRNA decreased translation

contributes to the nuclear regulation of the TGF-β pathway and of the downstream gene module in

G3 MB cells.

***LncMB3* inhibition synergizes with DDP treatment in G3 MB cells**

Cisplatin (DDP) is a first-line anti-cancer agent and a cornerstone of MB treatment [33]. Combining

therapies has been shown to reduce DDP side effects and overcome both intrinsic and acquired

cancer resistance. Based on the shared influence of *lncMB3* and DDP on apoptosis-related signalling

pathways [34], we explored whether *lncMB3* KD could enhance DDP cytotoxicity in G3 MB cells.

Initial tests of DDP administration to D283 Med cells at concentrations of 1-10μM reduced the

number of viable cells by 10%-20% after 48 hours (Fig. 5A, left panel). We then combined 10μM DDP

with transfection of 100nM GapmeR #1 or GapmeR #2 and counted the surviving cells.

Normalisation of cell counts from double-treated *vs* drug-untreated samples showed a reduced

relative cell survival rate in gap_*lncMB3*-treated cells compared to scramble-transfected controls,

indicating a synergistic effect between GapmeR and DDP treatments (Fig. 5A, right panel). This was

further confirmed through an apoptotic assay based on Annexin V staining (Fig. 5B, left panel).

Quantification of non-viable *vs* viable cells showed that combined treatments caused an increase of

late apoptotic cell number compared to other cell populations (Fig. 5B, middle panel), confirming a

synergistic effect (Fig. 5B, right panel).

We also examined whether this synergistic activity could be explained by crosstalk between the two

treatments. Testing the effect of DDP on the genes of interest revealed that cisplatin partially

deregulates both the TGF-β (Fig. 5C, left panel) and apoptotic pathway (Fig. 5C, right panel), similarly

to *lncMB3* targeting. Notably, *lncMB3* expression was not affected by chemotherapy (Fig. S8A),

suggesting that the two treatments represent independent triggers with overlapping outcomes.

It is worth noting that the synergistic enhancement of anticancer effects achieved through *lncMB3* targeting was not indiscriminate. When we tested the concurrent use of GapmeR #1 and vincristine, another drug used in MB therapy [35], we observed only additive, not synergistic effects on viable cell counts (Fig. S8B) and apoptosis (Fig. S8C). These data suggest that *in vitro*, *lncMB3* targeting synergistically enhances the efficacy of DDP treatment in reducing G3 MB cell proliferation and inducing apoptosis.

Production of the ferritin variant HFt-HIS-PASE

Targeting *lncMB3* impacts the TGF-β pathway, enhances apoptosis, and improves chemotherapeutic efficiency in cancer cell lines. These findings encouraged us to explore the use of recombinant human ferritin nanoparticles (NPs) to formulate innovative RNA/protein complexes for *lncMB3*-directed GapmeR delivery. Human ferritin H-type (HFt) is a promising NP for drug delivery, especially chemotherapeutics, due to its ability to be internalised by the transferrin receptor 1 (TfR1 or CD71), often overexpressed in tumour cells [13].

We designed and synthesized HFt-HIS-PASE, a recombinant ferritin variant derived from a previously described version of HFt, i.e. HFt-MP-PASE [36]. HFt-HIS-PASE contains a tumour-selective sequence (MP) responsive to proteolytic cleavage by specific tumour proteases (MMP 2 and 9), and a shielding polypeptide (PASE) that increases stability, masks the surface, and improves target specificity. Additionally, HFt-HIS-PASE includes a five-histidine motif reported to enhance endosomal escape of nucleic acids [37] (Fig. S9A).

Preliminary assessments of *MMP 2*, *MMP 9*, and *CD71* mRNA abundance in D283 Med and HD-MB03 cells via qRT-PCR (Fig. S9B) led to the selection of D283 Med cells for further experiments. We then evaluated the targeting capability of a Fluorescein-5-Maleimide-labeled HFt-HIS-PASE nanovector (HFt-HIS-PASE-fluo). D283 Med cells were cultured with HFt-HIS-PASE-fluo (1mg/mL) and 12 hours

post-administration ~85% of cells exhibited green fluorescence, confirming their affinity for HFt-HIS-PASE (Fig. 6A). Immunofluorescence performed on these samples 24 hours later, with nuclear (DAPI) and cytoplasmic (Phalloidin) staining, and confocal laser microscopy analysis, revealed intracellular uptake of HFt, with a nuclear-cytoplasmic distribution (Fig. 6B).

Characterisation of HFt-HIS-PASE-GapmeR complexes and *lncMB3* targeting

To investigate the potential of targeting *lncMB3* using HFt-ASO complexes, we generated and characterised HFt-HIS-PASE-GapmeR nucleoprotein particles. The *lncMB3* GapmeR #1 was encapsulated within the internal cavity of HFt-HIS-PASE using a pH-dependent dissociation/reassociation strategy. Successful loading of the GapmeR payload was verified via electrophoresis of the purified HFt-HIS-PASE-GapmeR complexes. Distinct migration patterns were observed for the HFt-HIS-PASE-GapmeR complex compared to the free GapmeR. When visualised by DNA-specific staining (Fig. 6C, left panel), the nucleic acid displayed a significant delay when associated with HFt-HIS-PASE (lane HFt + gap), compared to the free GapmeR. Protein-specific staining showed that the HFt-HIS-PASE-GapmeR complex co-migrated with ferritin (Fig. 6C, right panel).

Concerning complex stability, no evidence of nucleic acid degradation was observed after 1 hour and overnight incubations of HFt-HIS-PASE-GapmeR with DNA/RNA nuclease (Denarase) at 37°C (Fig. S9C). Additionally, no degradation was observed after 3 months of storage at 4°C (data not shown), confirming that GapmeR was encapsulated and protected within the protein cavity and not exposed to external surface.

To evaluate the efficiency of GapmeR binding to HFt NPs, the relative intensities of agarose bands corresponding to DNA or protein molecules were quantified. Comparisons were made to free GapmeR and HFt-HIS-PASE molecules in standard concentration as references. The final GapmeR-

HFt-HIS-PASE molecular ratio was determined to be 0,5:1. Purity and hydrodynamic volume of the

HFt-HIS-PASE-GapmeR complexes were assessed via size-exclusion chromatography. The HFt-HIS-

PASE-GapmeR sample displayed an elution volume and size similar to non-encapsulated HFt-HIS-

PASE protein, with no free GapmeR observed, indicating that GapmeR molecules in solution co-

eluted with ferritin (Fig. 6D).

The biological activity of the complex was tested in D283 Med cells treated with a single dose of HFt-

HIS-PASE alone or HFt-HIS-PASE-GapmeR complex (200nM). Steady-state *lncMB3* expression levels,

assessed 48 hours after administration, showed a 30% reduction following encapsulated GapmeR #1

delivery compared to mock-treated cells (Fig. 6E). Similar results were obtained after re-testing HFt-

HIS-PASE-GapmeR samples stored at 4°C for 3 months. This setup simulated the effects of a standard

single-pulse transfection of GapmeR #1 at 100nM for 48 hours (Fig. S9D). Likely due to insufficient

lncMB3 downregulation, we observed no modulation of TGF-β pathway genes, no reduction in cell

proliferation, nor any increase in apoptosis under these conditions.

However, to further explore the cellular effects of HFt-HIS-PASE-mediated *lncMB3* targeting, we

leveraged our earlier findings on the synergistic interplay between *lncMB3* KD and DDP treatment.

D283 Med cells were treated with 5μM DDP alone or in combination with HFt-HIS-PASE-GapmeR. In

the latter condition, we observed by qRT-PCR that combined DDP and GapmeR #1 administration

led to reduction in TGF-β pathway and downstream genes, ranging from 20% to 50% (Fig. 6F).

Consistent with previous observations (Fig. 5A), there was also a decrease in cancer cell survival

following the combined treatments, compared to DDP treatment alone (Fig. 6G). These results

confirm the efficacy of *lncMB3* targeting, facilitated by an HFt-based delivery system, in sensitising

G3 MB cells to anticancer drugs.

DISCUSSION

Integrated multi-omics has enabled the combination of genetic and histological analyses within a framework for understanding the inherent heterogeneity and complexity of MB [1]. Despite unique genomic, biochemical and clinical traits of each MB subgroup [38], current therapeutic strategies still rely on general interventions, including surgical resection, chemotherapy, and radiotherapy, with 5-year survival rates ranging from 60% to 80% [39]. This underscores the need for targeted therapies and deep explorations of dysfunctional pathways, particularly in high-risk and poorly characterised subgroups, such as G3 and G4. G3 MB, which accounts for 25% of cases, exhibits the highest recurrence rate and a strong propensity for metastasis, correlating with significantly worse outcomes [40]. The discovery of MYC amplification/overexpression as a major pathogenic factor [41], [42] is consistent with estimates of aberrant MYC activity in nearly 70% of human cancers [42]. However, targeting MYC has long been challenging due to its disordered structure, which has led to candidate therapeutic molecules with low potency, suboptimal pharmacological properties, and considerable off-target effects [43]. Furthermore, MYC plays a critical role in physiological processes, including cell differentiation, proliferation, and cell cycle regulation, which are central to development and post-natal life. This makes it essential to develop strategies for MYC inhibition particularly in paediatric cancers. Several pre-clinical studies and clinical trials are currently exploring direct MYC inhibitors [43], [44], such as the dominant-negative OMOMYC, that showed promising results in solid cancers [10], [44] and that we used to map the MYC transcriptome in G3 MB cells [10]. Indirect strategies that target MYC cofactors or regulators represent a viable alternative for functional MYC inhibition. Along this direction, a comprehensive understanding of MYC-dysregulated gene pathways, including non-coding RNA networks, may diversify and enhance opportunities for MYC-targeted therapies [45].

Over the past two decades, the rise of non-coding RNAs has revolutionised cancer biology [46]. LncRNAs have gained attention as modulators of cancer hallmarks due to their tissue- and cell-type-

specific expression and structural flexibility, making them attractive biomarkers and targets for molecular medicine. However, their potential as master regulators of biological processes require comprehensive functional characterisation.

To date, only 3 lncRNAs have been mechanistically described in G3 MB. This study elucidates the role of the MYC-regulated *lncMB3* in G3 medulloblastomagenesis and lays the foundation for its potential applicative use. *LncMB3* is upregulated in G3 MB cell lines and primary tumours and exhibits a strong activity in evading cell death *in vitro*. Through transcriptomic, molecular, and cellular assays, we demonstrate that *lncMB3* plays a crucial role in controlling the TGF-β cascade, a key player in G3 MB pathology, influencing cancer cell proliferation and survival [11]. Notably, we identified a cluster of *lncMB3* target genes in this pathway, whose aberrant expression leads to overexpression of the retinal TF OTX2, an oncogenic event in G3 MB that promotes cell proliferation, inhibits apoptosis, and contributes to tumour aggressiveness [47]. OTX2 and other photoreceptor TFs, including NRL and CRX, drive the expression of anti-apoptotic factors like BCL2L1 [23], linking this gene program to the sustained viability of cancer cells. *LncMB3* regulatory activity orchestrates critical pathways, genes, and processes in G3 MB, forming a nexus among the *MYC* driver gene, the TGF-β signalling, photoreceptor gene network, and apoptosis regulators. This interconnection offers an integrated framework for understanding the molecular underpinnings of this tumour subgroup.

While our coding transcriptome analysis provided insights into *lncMB3* function, the exploration of its interactome revealed its mode of action. *LncMB3* directly interacts with the mRNA encoding the epigenetic regulator HMGN5, a member of the HMGN nucleosome-binding and nuclear architecture family that extensively regulates chromatin decondensation and transcription [30], [31]. *HMGN5*, which plays a role in embryonic gene regulation [48] and displays oncogenic properties in several cancers [30], [31], is upregulated in G3 MB. Our findings suggest that RNA-RNA interactions between coding and non-coding molecules, an emerging theme in cancer regulation, result in the functional

upregulation of HMGN5 protein levels, explaining why *HMGN5* RNA interference phenocopies the effects of *lncMB3* targeting. Since several *lncMB3* targets are dysregulated in the nucleus, where HMGN5 operates, the mechanism we propose adequately accounts for the observed molecular and cellular phenotypes. This does not exclude the possibility of additional regulative cascades or secondary effects, which could align with the recently suggested moonlighting role for HMGN5 in RNA binding [48]. Finally, *lncMB3* and *HMGN5* co-expression, confined to pathological cerebellum, proposes their interaction as a platform for developing innovative inhibitors to compete with disease-specific complexes.

Due to the aggressive nature of G3 MB and its resistance to treatments, integrated therapies offer promising solutions. Meanwhile, ASOs have gained prominence in RNA-based treatments due to their flexible design and favourable pharmacological properties [46]. We investigated the interplay between *lncMB3*-targeting LNA GapmeRs and standard MB chemotherapeutic approaches, revealing synergistic effects that enhance drug cytotoxicity. This combinatorial approach likely stems from both treatments acting on the intrinsic apoptotic pathway [34] and the TGF-β cascade. These findings suggest that modulating the *lncMB3* molecular network could amplify the efficacy of anticancer agents, paving the way for RNA-based combinatorial strategies to reduce tumour resistance and improve treatment outcomes. However, one of the major challenges in RNA-based therapies is ensuring efficient delivery of effectors to the intended sites. We explored the application of HFt-based nanocarriers for deploying LNA GapmeRs to target *lncMB3*. HFt unique properties, from high biocompatibility and stability to low toxicity and production costs, make it a powerful tool for drug delivery [13]. Moreover, HFt ability to cross the blood-brain barrier [13] positions it as a promising tool for brain tumour-targeting therapies. Finally, ferritin symmetrical self-assembly, small uniform size, and versatile surface functionalisation make it ideal for bioactive compound delivery, including small nucleic acids. Though HFt has been employed to administer siRNAs [49] and

microRNAs [50], this work represents, to our knowledge, the first example of gene silencing via HFt-encapsulated ASOs. The combined administration of *lncMB3*-targeting NPs and cisplatin reproduced the synergistic effects observed with ASO transfection, impacting on the TGF-β pathway and reducing cell survival. These results indicate that HFt-based strategies hold promise for targeting oncogenic pathways at the RNA level through biocompatible delivery of tumour-specific antisense oligomers.

In conclusion, this study highlights the critical role of *lncMB3* in regulating apoptosis in G3 MB cells through modulation of the TGF-β pathway and interaction with underexplored molecular players such as HMGN5. *LncMB3* stands out as a regulatory node in the G3 MB pathway, linking MYC amplification to increased tumour cell survival by cell death inhibition (Fig. 7). Future efforts will focus on further investigating *lncMB3* interactions and regulatory mechanisms, as well as validating the therapeutic potential of targeting the lncRNA network in preclinical G3 MB models using the HFt nanocarriers.

MATERIALS AND METHODS

Cell Cultures

All cell lines were obtained from ATCC. Semi-adherent D283 Med cells were cultured in MEM medium (M2279, Sigma-Aldrich) supplemented with 20% of heat-inactivated foetal bovine serum (USA origin), 1% Sodium Pyruvate, 1% MEM Non-Essential Amino Acids Solution, 1% Glutamax, and 1x Penicillin/Streptomycin. Adherent HD-MB03 cells were cultured in RPMI medium (R0883, Sigma-Aldrich) supplemented with 10% of heat-inactivated foetal bovine serum (USA origin), 1% Penicillin/Streptomycin.

Cell Propagation

D283 Med floating cell subpopulation was collected, whereas adherent cell fraction was washed

with 1x PBS w/o Ca^{++}/Mg^{++} and detached by incubation in 1x Trypsin/EDTA (T4299, Sigma-Aldrich)

for 5 min. Cell fractions were centrifuged at 800 RPM for 5 min and resuspended in fresh culture

medium. HD-MB03 cells were washed, detached, centrifuged and resuspended as described above.

Cells were counted on a CytoSMART Cell Counter (CLS6749, Corning, Sigma-Aldrich) and replated at

the desired concentration.

Evaluation of Viable Cell Number

D283 Med cells and HD-MB03 cells were seeded in plates, treated according to experimental plans

(GapmeR/plasmid transfection and/or chemotherapeutic treatment) and counted at different time

points. Specifically, 10µL cell aliquots were thoroughly suspended, mixed to 10µL of trypan blue dye

and counted by CytoSMART Cell Counter for cell population quantifications.

Plasmids construction

Δ16: D283 Med cell cDNA was used as template to amplify with DNA oligos MB3 FW and MB3 REV

(Table 2) a mutant version of *lncMB3* sequence lacking GapmeR #1 terminal binding site. EPB-PURO-

TT backbone was linearised with DNA oligos Delta16 FW - T7 REV (Table 2). Construct was generated

using In-Fusion Snap Assembly Cloning Kit (Takara Bio).

sh_HMGN5: was obtained by linearising the EPB-BSD-PGK backbone with DNA oligos

shRNA_TOOL_fw and shRNA_TOOL_rev (Table 2). Construct was generated using In-Fusion Snap

Assembly Cloning Kit (Takara Bio). Control plasmid was obtained by [25].

sh_HMGN5 sequence: GTTGTTGAAGAAGACTACAAT

Transfections

LNA GapmeRs: negative control (LG00000002, Qiagen) or specific LNA GapmeRs (Qiagen) were

transfected 1-3 times (every 24 hours) in D283 Med or HD-MB03 cells at a concentration of 100 nM

with Lipofectamine 2000 (11668-019, Invitrogen, ThermoFisher Scientific), in opti-MEM medium

(31985070, Gibco, ThermoFisher Scientific), according to manufacturer's instructions.

Control (empty EPB-PURO-TT) and Δ16 plasmids were transfected in D283 Med cells at 100 ng with

Lipofectamine 2000 (11668-019, Invitrogen, ThermoFisher Scientific), in opti-MEM medium

(31985070, Gibco, ThermoFisher Scientific), according to manufacturer's instructions. Doxycycline

100 ng/mL (D9891, Sigma Aldrich, Merck KGaA) was added in the complete growing medium.

Five µg of sh_SCR (negative control) or sh_HMGN5 plasmids were transfected twice (24 and 48 hours

after cell seeding) in D283 Med cells with Lipofectamine 2000 (11668-019, Invitrogen), in opti-MEM

medium (31985070, Gibco, ThermoFisher Scientific), according to manufacturer's instructions.

RNA Extraction and Analysis by qRT-PCR

Total RNA was extracted by Direct-zol RNA MiniPrep (R2052, Zymo Research). For quantitative real-

time PCR (qRT-PCR) assay, cDNA was synthesised by Takara PrimeScript RT Reagent Kit (RR037A,

Takara-bio). qRT-PCR detection was performed using SensiFAST SYBR Lo-ROX Kit (BIO-94020, Bioline)

on a 7500 Fast Real-Time PCR (Applied Biosystem). *GAPDH* or *ATP5O* were used as a reference target.

Digital PCR analysis

Digital PCR analysis was performed on the QIAcuity® nanoplate digital PCR 8.5K 24-well (Qiagen).

The PCR reaction mix was prepared following the manufacturer's instructions. Twelve µl of the final

PCR mix were loaded on each nanoplate well, and then automatically sub-aliquoted in 8500

partitions. The EVA green signal acquisition was performed with an exposure duration of 500 ms and

a gain of 6.

Immunoblotting

Protein samples for immunoblotting were extracted from MB cells in RIPA Buffer (50 mM Tris–HCl (pH 8), 150 mM EGTA, 150 mM NaCl, 50 mM NaF, 10% glycerol, 1.5 mM $MgCl_2$, 1% Triton). Lysates were separated on gradient poly-acrylamide gels and transferred to Amersham Protran 0.45 µm nitrocellulose membrane (10600002, GE Healthcare Life Sciences), through the NuPAGE System (EI0002, Invitrogen). Immunoblots were incubated with the following antibodies: anti-GAPDH (sc-32233, Santa Cruz Biotechnology); anti-FLAG (F1804, Sigma-Aldrich); anti-cleaved PARP-1 (bsm-52408R, Bioss); anti-PARP-1 (PA5-34803); anti-HMGN5 (720391, Invitrogen); anti-OTX2 (ab183951, Abcam); anti-BCL2L1 (2764S, Cell Signaling); anti-ATP5O (A305-419A, Bethyl Laboratories). Protein staining was performed by WesternBright ECL (K-12045-D50, Advansta), detected by ChemiDoc XRS+ Molecular Imager (Bio-Rad) and quantified through the Image Lab Software (release 3.0.1).

RNA-Seq and differential expression analysis

TruSeq Stranded mRNA Library Prep Kit (Illumina, San Diego, CA, USA) was used to obtain sequencing libraries from polyA+ RNA extracted from D283 Med cells (3 independent biological replicates). The sequencing reaction produced 100 nucleotide-long paired-end reads and was performed on a Novaseq 6000 sequencing system (Illumina) with a depth of more than 20M. To remove adapter sequences and low-quality end bases, Trim Galore188 (version 0.6.4_dev) software was used; the minimum read length after trimming was set to 20. Alignment to human GRCh38 genome primary assembly was performed using STAR (version 2.7.9a189). Quality of the sequencing data was assessed with FastQC. More than 85% of the reads were uniquely mapped to the human genome. The "--quantMode TranscriptomeSAM" option was used to generate alignments translated into transcript coordinates. The expression levels of transcripts and genes were quantified by using

RSEM. Differential expression analysis (DEA) was performed using the Bioconductor package "DESeq2". Only those genes that showed an absolute expression level of at least 10 in minimum three of the six samples were retained for the DEA. Shrinked logFC values were obtained by the function "lfcShrink". FDR < 0,05 and |logFC| > 0,6 were considered differentially expressed and used for further analysis.

Gene Ontology

Gene Ontology (GO) overrepresentation analysis was performed using the DAVID tool (https://david.ncifcrf.gov/tools.jsp) on Wikipathways databases. GO analysis was conducted on both significantly upregulated and downregulated genes.

Gene Set Enrichment Analysis

Gene Set Enrichment Analysis (GSEA) was performed by using the Bioconductor package "fgsea". The whole gene set of the experiment was employed and, specifically, logFC and padj values. Using this information, genes were ranked in a sample dataset by multiplying -$\log_{10}$(FDR) * logFC according to their differential expression between two experimental groups. Ensemble Gene IDs were mapped to Entrez Gene IDs using BioMart, and the "fgsea" function was run with minSize = 15, maxSize = 500 and nPermSimple = 10000. The analysis was performed on the gene sets belonging to the Molecular Signatures Database (MsigDb v7.4). For each gene set, an enrichment score was computed and assigned to the gene set of the ranked gene lists. Additionally, GSEA calculated a normalised enrichment score to account for variations in pathway size, ensuring robustness across different gene sets and datasets. An FDR of 25% was used for this analysis.

Treatments with chemotherapeutics

Chemotherapeutics (cisplatin, HY-17394, MedChem; vincristine HY-N0488, MedChem) were

resuspended in DMSO and administered at different concentrations according to the experimental

plan. For combined treatments, the selected amount of drug was administered 5 hours after

transfections. Cells were counted/analysed 24/48 hours after treatment.

Flow Cytometry Analyses

Apoptosis induction was analysed by flow cytometry determination of Annexin V-FITC staining (Enzo

Life Sciences, ALX-850-020), to discriminate viable (FITC-/PI-), early apoptotic (FITC+/PI-) and late or

necrotic cells (FITC+/PI+). Briefly, cells were collected and resuspended in binding buffer containing

Annexin V-FITC and incubated at RT for 15 min. PI was added before reading the sample to flow

cytometer (BD FACSCalibur; Beckman Coulter - CytoFLEX SRT). Data were analysed using the Flowing

software. Ten thousand events were collected from each sample.

Native RNA pull-down assay

Native RNA pulldown experiments were performed as described in [25]. Briefly, 1×10^8 proliferating

D283 Med cells were centrifuged at 400 g for 5 min and pellets were lysed for 30 min on rotating

wheel at 4°C in Cell Lysis Buffer (Tris-HCl pH 7,5 50 mM, NaCl 150 mM, MgCl2 3 mM, NP40 0,5%,

EDTA 2 mM, DTT 1 mM; 1× PIC and RNase inhibitors). The lysate was cleared by centrifugation at

15000 g for 15 min at 4°C. One mg of supernatant (whole cell extract) was diluted in a 1:2 ratio with

Hybridization Buffer (Tris-HCl pH 7,5 100 mM, NaCl 300 mM, MgCl2 1 mM, SDS 0,2%, Formamide

15%, NP40 0,5%, EDTA 10 mM, DTT 1 mM, 1× PIC and RNase inhibitor).

10% of the total extract was collected for the Input sample. 100 pmol of biotinylated probes (EVEN,

ODD and LacZ sets, see Table probes) were heated at 80°C for 3 minutes and added to the mix. After

4-hour incubation on rotating wheel at 4°C, 0,1 ml of washed streptavidin Magnasphere

paramagnetic beads (Promega) were added, and the mixture was incubated for as before. After pull-

down, beads were collected using a magnetic rack and washed 4 times with Hybridization Buffer.

Then 500 µL triZOL (ThermoFisher Scientific) were added to each sample and RNA was extracted

(RNeasy Mini Kit Qiagen), DNase-treated and retro transcribed using SuperScript Vilo cDNA

Synthesis Kit (ThermoFisher Scientific). Pull-down qRT-PCR results were represented as a percentage

of enrichment related to INP.

AMT-crosslinked RNA pull-down assay

AMT-crosslinked RNA pull-down experiments were performed as described in [25]. Briefly, 1×10^8

D283 Med cells were centrifuged at 400 g for 5 min, resuspended in 10 mL of ice-cold complete PBS

supplemented with 0,5 mg/ml 4'-aminomethyl-4,5',8-trimethylpsoralen (AMT, Sigma-Aldrich), and

crosslinked at 365 nm five times for 2-minutes cycles. Guanidine Hydrochloride 6M was added 1:1

to AMT volume. The lysate was split into 250µL aliquots and 25µL of a 20mg/mL solution of

Proteinase K (Ambion) and 6,5µL of 20% SDS were added to each aliquot and incubated at 65°C for

1 h. Then 750µL triZOL (ThermoFisher Scientific) were added to each sample and RNA was extracted

(RNeasy Mini Kit Qiagen). 50µg of purified AMT-crosslinked RNA were used for each RNA pull-down

sample. 500µL of RNAse-free water were added to each sample and heated at 95°C for 2 minutes

and then put on ice. 1 volume of 2x Binding Buffer (containing Tris-HCl pH 7,5 20mM, NaCl 1M, EDTA

2mM, SDS 0,1%) was added to 1 volume of sample with RNAse inhibitors. 10% of the total extract

was collected for the Input sample. 500 pmol of biotinylated probes (EVEN, ODD, LacZ sets, see Table

2) were added to the mix and samples were heated at 65°C for 3 minutes. After 4-hour incubation

on rotating wheel at 4°C, 0,5ml of washed streptavidin Magnasphere paramagnetic beads (Promega)

were added, and the mixture was incubated for 2h at 4°C on rotating wheel. After pull-down, beads

were collected using a magnetic rack and washed 4 times with hybridization buffer. Then 500µL

triZOL (ThermoFisher) were added to each sample and RNA was extracted (RNeasy Mini Kit Qiagen),

UV-decrosslinked at 254nm for 10 minutes, DNase-treated and retro transcribed using SuperScript

Vilo cDNA Synthesis Kit (ThermoFisher Scientific). Pull-down (PD) qRT-PCR results were represented

as a percentage of enrichment related to INP.

Pull down-RNAseq and pre-processing

Raw reads in FASTQ format were retrieved from Illumina BaseSppace and adaptors removed with

fastp tool. Clean reads were aligned to a GRCh38 assembly using STAR aligner software. Gene *loci*

fragment quantification was performed on a customised gene annotation file (GTF file) based on

Ensemble (release 87), using the *–quantMode GeneCounts* parameter. The GTF file was previously

edited adding the *lncMB3* gene genomic coordinates and annotations. Read counts were combined

into a count matrix and processed with DESeq2 R package for normalisation and differential

expression analysis. The padj threshold was set to 0,05 to consider DEGs as statistically significant.

Pull down-RNAseq analysis

edgeR was used to quantify significant genes as putative interactors on *lncMB3*, using a generalised

linear model (GLM) to test for each gene. Statistically significant genes were obtained by separately

comparing and intersecting EVEN and ODD samples *vs* Input samples, and then subtracting genes

obtained in the comparison between LacZ *vs* Input samples. Only genes with positive FC were used

and retained as putative interactors.

RNA–RNA interaction prediction

RNA–RNA interaction prediction was computed using IntaRNA 3.3.2 giving as input the full FASTA

sequences of transcripts of interest, using the default parameters.

microRNA *in silico* predictions

TargetScan Human database, release 8, was exploited to obtain candidates microRNA families targeting genes of interest. MicroRNA families were filtered by choosing those conserved among vertebrates and mammals, then we selected those with conserved sites on the target transcripts, and finally we filtered out the families having more than -0,20 cumulative weighted context++ score, considered as the beginning of moderate confidence level of interaction. Interaction predictions were run by using a locally downloaded version of the interaction prediction tool RNA22.

Crosslinking Immunoprecipitation assay

For Crosslinking Immunoprecipitation (CLIP) assay $3,5\times10^7$ D283 Med cells were grown, UV-crosslinked at 4000×100 μj/cm2 (energy) and collected in NP-40 lysis buffer (50 mM HEPES-KOH, 150mM KCl, 2mM EDTA, 1mM NaF, 0,5% (v/v) NP40 pH 7,4, 0,5mM DTT) by low sonication and centrifugation at 20.000 rcf. For immunoprecipitation, 2mg of cytoplasmic extracts were incubated with 10μg of AGO2 antibody or IgG as a control, and coupled to ProteinG Dynabeads resin (10004D, Invitrogen) and 10% was collected as Input. After washing with high-salt buffer (50mM HEPES-KOH, pH7,5, 500mM KCl, 0,05%(v/v) NP40, 0,5mM DTT), immunoprecipitated proteins (20% of the sample) were collected in RIPA buffer and analysed by immunoblotting. Immunoprecipitated RNA (80% of the sample) was treated by Proteinase K, extracted by Direct-zol RNA MiniPrep (Zymo Research), retro-transcribed by Superscript Vilo cDNA synthesis Kit (11754050, Invitrogen) and analysed by qRT-PCR using specific primers. RNA enrichments were reported as percentage of the input.

Ribotagging and RNA Immunoprecipitation

1,5x10^7 D283 Med cells were transfected with 15µg of FLAG-RPL22 expressing plasmid. On the day

after, cell populations were splitted in two fractions and each sample was treated with 100nM

gap_SCR or gap_*lncMB3*, respectively. 24 hours after GapmeRs administration, lysates were

prepared in 350µL of POLYSOME LYSIS BUFFER (HEPES 10mM pH 7, KCl 100mM, MgCl2 5mM, NP-40

0,5%, dH20) and NT2 BUFFER, Tris 50mM pH 7,5, NaCl 150mM, MgCl2 1mM, NP-40 0,05%, dH2O).

150µL of extracts were incubated with 10µg of FLAG antibody (SLBD6976, Sigma Aldrich) or IgG

(ultrapure mouse IgG, MU-003-N, Immunoreagents) as control and precipitated through ProteinG-

Dynabeads resin (10004D, Invitrogen). 30µL were collected as Input. Immunoprecipitated proteins

(20% of the sample) were prepared in loading dye for immunoblot. Immunoprecipitated RNA (80%

of the sample) was extracted by Direct-zol RNA MiniPrep (Zymo Research), retro-transcribed by

Superscript Vilo cDNA synthesis Kit (11754050, Invitrogen) and analysed by qRT-PCR using specific

primers. RNA enrichments were reported as percentage of the input.

HFt-HIS-PASE cloning, expression and purification

The design of HFt-HIS-PASE protein, a variant of the HFt-MP-PASE gene previously reported [36],

includes, C- to N-terminus: (a) a first domain, comprising the amino acid sequence of the native

human ferritin HFt modified through the inclusion of a 5-histidine peptide sequence, inserted in the

external loop (referred as CD loop) of the ferritin surface; (b) a second N-terminal domain comprising

the amino acid sequence of the matrix metalloproteinase (MMP) cleavage site (PLGLAG) followed

by a Proline, Serine, Alanine, and Glutamate (PASE) sequence (ASPAAPAPASPAEPAPSAPA). The

expression vector pET-27b containing the HFt-HIS-PASE gene was assembled by GENEART AG

(Thermo Fisher Scientific), considering codon optimisation for high expression levels in *E. coli*. BL21

(DE3) strain (New England BioLabs).

HFt-HIS-PASE was produced with a typical yield of 50mg of pure protein per litre of culture. Cells

harbouring the recombinant gene were grown o.n. at 23°C and then at 37°C up to OD600: 1,0, in 1

L of terrific broth medium (Grisp), containing 0,03mg/mL kanamycin (Serva). After induction with

1mM Isopropil-β-D-tiogalattopiranoside (PanReac) bacteria were incubated at 30°C for 2 hours,

harvested by centrifugation at 5000 rpm for 20 minutes at 4°C, resuspended in phosphate-buffered

saline (PBS) complemented with 150mM NaCl and 20mM imidazole, pH 7,4, protease inhibitors

(ThermoFisher Scientific) and 1mM Phenylmethylsulfonyl fluoride (ThermoFisher Scientific). DNAse

(0,1 mg/mL final concentration, Sigma-Aldrich) was added before cell disruption by sonication. The

lysate was incubated at 37°C for 40 minutes and centrifuged at 14000 rpm for 50 minutes at 4°C.

The supernatant was recovered and loaded in a buffer-equilibrated His-Trap column (Cytiva) for

metal affinity chromatography. HFt-HIS-PASE was purified by 100mM imidazole-containing elution

buffer, dialysed o.n. against PBS, pH 7,5, and concentrated by 100kDa Amicon Ultra-15 centrifugal

filter devices (Millipore). Finally, HFt-HIS-PASE was sterile filtered and stored at 4 °C (short-term) or

-20°C (long-term). Preparation purity was assessed by protein staining with Coomassie brilliant blue

after a run on 15% SDS-PAGE. Protein concentrations were spectrophotometrically determined at

280 nm, using a molar extinction coefficient (on a 24-mer basis) of $4,56 \times 10^5$ M-1cm-1 (ProtParam

sofware, http://www.expasy.org).

HFt-HIS-PASE characterisation

Size-Exclusion Chromatography experiments were performed using a Superose 6 gel-filtration

column equilibrated with PBS at pH 7,4. All samples were prepared at 1mg/mL in filtered ddH2O. All

the traces for SEC experiments were analysed with QtiPlot (IONDEV SRL, Bucuresti, Romania).

HFt-HIS-PASE fluorescent labelling

The HFt-HIS-PASE protein (1mg/mL) was incubated with 1mM Fluorescein-5-Maleimide (λex 491 nm,

λem 518 nm; ThermoFisher Scientific) in PBS, pH 7 for 2 hours at R.T., under stirring in the dark. The

sample was filtered, dialysed, and exchanged with PBS as described above. The sample was sterile

filtered and stored at 4 °C in the dark. The number of dyes linked per protein was determined by

absorbance spectroscopy, in accordance with the manufacturer's instructions, applying the

Lambert–Beer law.

Synthesis of HFt-HIS-PASE-GapmeR complex

GapmeR #1 was encapsulated within the internal cavity of the HFt-HIS-PASE using reversible, pH-

dependent cage dissociation (acid pH) and re-association (neutral pH), in the presence of bioactive

payload. HFt-HIS-PASE dissociation into subunits was achieved by lowering pH to 2,8 by 0,1M HCl.

GapmeR #1 was added to the disassembled protein solution at a molar ratio of 1:1. The pH was then

adjusted to 7,4 by 0,1M NaOH, the solution was ultra-filtered at pH 7,4 and exchanged with PBS by

100kDa Amicon Ultra-15 centrifugal filter devices to remove unbound molecules.

Characterisation of HFt-HIS-PASE-GapmeR complex

Electrophoresis on 1,8% agarose gel (AGE) of the purified ferritin-GapmeR complexes was used to

demonstrate the loading of GapmeRs in the protein cavity. The gel was run at 90V for 60 min stained

with Nucleic Acid SYBR Gold for DNA/RNA visualisation and imaged by ChemiDoc™ (Bio-Rad,

America). The amount of GapmeR bound to the ferritin NPs was evaluated by the Image J software

(https://imagej.net/ij/), by calculating the area and statistics value of pixels in user-defined selection-

objects based on intensity. The same analysis was used to evaluate the concentration of the protein

using standards of HFt-HIS-PASE as reference after gel staining with Coomassie dye. Purity and

hydrodynamic volume of the HFt-HIS-PASE-GapmeR complexes were determined by SEC

experiments. To verify GapmeR encapsulation and protection, HFt-HIS-PASE-GapmeR sample was

incubated at 37°C for 1 hour and overnight with 12,5x103 U/mL Denarase (c-LEcta GmbH) to digest

DNA and RNA impurities. AGE was used as previously described to determine the GapmeR intensity

of each sample. AGE was also employed to check the stability of HFt-HIS-PASE-GapmeR NPs stored

at 4 °C for 3 months.

HFt-HIS-PASE-GapmeR delivery

HFt-HIS-PASE-empty and HFt-HIS-PASE-GapmeR complexes were added to D283 Med cell complete

growing medium at a concentration of 200nM and maintained for 48 hours. In HFt/DDP double

treatment experiments 5µM DDP was added to cells 2 hours after HFt-HIS-PASE administration.

Cytoplasm and nucleus staining

D283 Med cells were cultured as previously described on 0,01% poly-L-ornithine/Murine Laminin

20µg/ml (Sigma Aldrich)-coated coverslips, fixed in 4% paraformaldehyde (Electron Microscopy

Sciences) for 20 min at 4 °C and washed with PBS. Fixed cells were then permeabilised and blocked

with 0,2% Triton X-100/3% BSA/PBS for 15 min at RT. Subsequently, cells were incubated for 45 min

at RT in a solution containing 1% donkey and 1% goat serum in PBS, Alexa Fluor™ 555 Phalloidin

1:100 (A34055, Invitrogen) to mark actin filaments. After washing with PBS, nuclei were

counterstained with DAPI solution (1 µg/ml/PBS, D9542, Sigma Aldrich) and the coverslips were

mounted using ProLong Diamond Antifade Mountant (Thermo Fisher Scientific, P-36961).

Confocal Microscopy

Samples were imaged using an Olympus iX83 FluoView1200 laser scanning confocal microscope.

Acquisitions were performed with a 60X oil objective (NA 1,35) and a 20x objective (NA 0,75) and

images were acquired in 16-bit with a resolution of 800 x 800 pixels. For each acquisition with the

60X objective, at least 30 focal planes were imaged with a step size of 0,3μm. For each acquisition

with the 20X objective, at least 50 focal planes were imaged with a step size of 0,5μm.

Cell Count Analysis

Cell counting was performed using an Olympus iX83 FluoView1200 laser scanning confocal

microscope with a 20X objective (NA 0,75). Images were acquired in 16-bit with a step size of 0,5μm

over at least 50 focal planes. A threshold was set for DAPI (blue, nuclear staining) and for FTH (green,

wavelength 488) signals. A maximum intensity projection was created from the Z-stack using Fiji,

resulting in a binarised image.

The Fiji "Analyse Particles" tool was employed with a minimum size of 25^2 pixels per field. This

approach allowed for the counting of particles in each field for both the green and DAPI signals. The

number of green cells was calculated relative to the total number of DAPI-marked cells, providing a

percentage of green signal delivery.

Colocalisation Analysis

Colocalisation between signals was analysed using the "JACoP" Fiji plugin. The green signal

(wavelength 488 nm), representing FTH, and the red signal (wavelength 555 nm), provided by

Phalloidin for cytoplasmic staining, were assessed for colocalisation. Cell nuclei were identified using

DAPI (blue). JACoP thresholds of 290 for the green signal and 181 for the blue (DAPI) or red

(Phalloidin) signals were applied to all the acquisitions to evaluate the percentage of colocalisation.

Manders' coefficients were used to determine the overlap between the signals, with the

colocalisation percentages indicating the extent of the signal overlap.

 Statistical Analyses

Results are expressed as mean ± SEM from at least 3 biological replicates. In specific cases single

experiments were analysed (biological replicates are indicated in figure legends). Statistical

differences between groups were analysed by two-tailed Student's *t*-test (performed with GraphPad

QuickCalcs). A *p*-value < 0,05 was considered statistically significant. * *p* < 0,05, ** *p* < 0,01, *** *p* <

0,001, **** p < 0,0001. One-way analysis of variance (ANOVA) was applied to multi-group data

comparisons, followed by Fisher's LSD test.

Oligonucleotides and probes

Oligonucleotide, probe and GapmeR sequences are listed in Table 2.

MAIN FIGURE LEGENDS

Figure 1. Differential gene expression analysis upon *lncMB3* KD in G3 MB cells

A. Left Panel: heatmap showing the relative levels of DEGs according to RNA-Seq analyses, along

with genes hierarchical clustering (3 biological replicates for each condition). The heatmap

represents only the DEGs. Right Panel: volcano plot showing DEG distribution. Genes were plotted

based on statistical significance $-\log_{10}$(FDR) and differential expression $\log_2$(FC). Downregulated

genes (logFC < -0,6) are indicated by the blue dots; upregulated genes (logFC > 0,6) by red dots.

Invariant genes are indicated in black.

B. GO (wikipathways) showing the distribution in clusters of the DEGs in gap_lncMB3 *vs* gap_SCR-

treated D283 Med cells, according to RNA-seq data. Categories are listed considering $-\log_{10}$(pvalue)

(left panel). qRT-PCR validation analysis of TGF-β pathway genes in D283 Med cells treated for 72

hours with gap_SCR or gap_*lncMB3* (GapmeR #1, right panel). Expression levels were compared to

gap_SCR sample as control, set as 1. Data (means ± SEM) are expressed in arbitrary units and are

relative to *GAPDH* mRNA levels. *N* = 3, * *p* ≤ 0,05, ** *p* ≤ 0,01, *** *p* ≤ 0,001 (two- tailed Student's *t*-

test).

C. qRT-PCR analysis of TGF-β pathway genes in D283 Med cells treated for 72 hours with gap_SCR or

gap2_*lncMB3* (GapmeR #2, left panel), or in HD-MB03 cells treated as above (GapmeR #2, right

panel). Expression levels were compared to gap_SCR sample as control, set as 1. Data (means ± SEM)

are expressed in arbitrary units and are relative to *GAPDH* mRNA levels. *N* = 3 to 5, * *p* ≤ 0,05, ** *p*

≤ 0,01, *** *p* ≤ 0,001 (two- tailed Student's *t*-test).

Figure 2. Expression of *OTX2*, *NRL*, *CRX* and *BCL2L1* in MB

A. *OTX2*, *NRL*, *CRX* and *BCL2L1* mRNA expression in 4 healthy cerebella (normal samples) and 59

primary MB samples according to the transcriptomic dataset GSE164677. Results are expressed in

log_2(FPKM). * p ≤ 0,05, ** p ≤ 0,01, *** p ≤ 0,001 **** p ≤ 0,0001 (two-tailed Student's *t*-test).

B. Schematic representation of *OTX2* network on apoptosis (left panel). GSEA biological process (BP)

domains performed on deregulated genes. Categories are listed considering -log_{10}(pvalue) (middle

panel). qRT-PCR analysis of *OTX2*, *NRL*, *CRX* and *BCL2L1* in D283 Med cells treated for 72 hours with

gap_SCR or gap_*lncMB3* (GapmeR #1, right panel). Expression levels were compared to gap_SCR

sample as control, set as 1. Data (means ± SEM) are expressed in arbitrary units and are relative to

GAPDH mRNA levels. *N* = 3, ** p ≤ 0,01, *** p ≤ 0,001 (two- tailed Student's *t*-test).

C. Western blot analysis of OTX2 and BCL2L1 protein levels in D283 Med cells treated for 72 hours

with gap_SCR or gap_*lncMB3* (GapmeR #1). Normalisations were performed relative to GAPDH

protein levels. N = 3, * *p* ≤ 0,05, *** *p* ≤ 0,001, (two-tailed Student's t-test).

D. Western blot analysis of PARP1 protein levels in D283 Med cells treated for 72 hours with gap_SCR

or gap_*lncMB3* (GapmeR #1). Details as in panel C.

Figure 3. Analysis of TGF-β pathway and apoptosis in G3 MB cells

A. qRT-PCR analysis of *OTX2*, *NRL*, *CRX* and *BCL2L1* in HD-MB03 cells treated for 72 hours with gap_SCR or gap_*lncMB3* (GapmeR #1). Expression levels were compared to gap_SCR sample as control, set as 1. Data (means ± SEM) are expressed in arbitrary units and are relative to *GAPDH* mRNA levels. N = 3, * p ≤ 0,01 (two- tailed Student's *t*-test).

B. Western blot analysis of OTX2 (left panel) and BCL2L1 (right panel) protein levels in HD-MB03 cells treated for 72 hours with gap_SCR or gap_*lncMB3* (GapmeR #1). Normalisations were performed relative to GAPDH protein levels. N = 3, * p ≤ 0,05, *** p ≤ 0,001, (two-tailed Student's t-test).

C. Time course analysis of the number of viable HD-MB03 cells depleted for *lncMB3* (gap_*lncMB3*). Scramble-transfected cells (gap_SCR) were used as control. Cells were transfected at day 0 (D0) and counted at each timepoint. Data (means ± SEM) are expressed as the number of viable cells, counted by an automated cell counter. N = 4, ** p ≤ 0,01, (two-tailed Student's *t*-test).

D. Western blot analysis of c-PARP1 (left panel) and PARP1 (right panel) protein levels in HD-MB03 cells treated for 72 hours with gap_SCR or gap_*lncMB3* (GapmeR #1). N = 3, ** p ≤ 0,01, *** p ≤ 0,001. Details as in panel B.

E. Dose-response analysis of the number of viable D283 Med cells after TGF-β inhibition. Control cells were treated with vehicle (DMSO). Cell counts started 24 hours after treatment. Data (means ± SEM) are expressed as the number of viable cells, counted by an automated cell counter. N = 4, * p ≤ 0,05, ** p ≤ 0,01 (two-tailed Student's *t*-test).

F. Annexin V staining of D283 Med cells upon TGF-β inhibition. Left panel: representative flow cytometry analysis of propidium iodide- and Annexin V-stained cells. Right panel: quantification of the fractions (expressed as percentage of total cell number) of viable (Annexin V-/PI-), early apoptotic (Annexin V+/PI-), late apoptotic (Annexin V+/PI+) and necrotic (Annexin V-/PI+) cells. For early+late apoptotic cells, N = 3, * p ≤ 0,05 (two-way ANOVA). **G.** Left panel: schematic

representation of the construct overexpressing *lncMB3* Δ16. Middle panel: qRT-PCR analysis of *lncMB3* expression in D283 Med cells transfected with gap_SCR, gap_*lncMB3* or gap_*lncMB3* + Δ16 construct. *N* = 5. Details as in panel A. Right panel: number of viable D283 Med cells upon gap_*lncMB3* transfection (*lncMB3* KD) or Δ16 co-expression, compared to gap_SCR condition. Normalisation was performed on *lncMB3* KD condition. *N* = 5, ** *p* ≤ 0,01 (two- tailed Student's *t*-test).

Figure 4. Analysis of *lncMB3*-*HMGN5* mRNA interaction

A. qRT-PCR analysis of RNA enrichment in EVEN, ODD and LacZ fractions over Input, from native *lncMB3* RNA pull-down experiments of D283 Med cell extracts. Data expressed as percentage of Input, *N* = 2.

B. Venn diagram showing intersections of mRNAs retrieved and sequenced from two independent native *lncMB3* pull-down fractions (EVEN, ODD, LacZ).

C. *HMGN5* mRNA expression in 4 healthy cerebella (normal samples) and 59 primary MB samples according to the transcriptomic dataset GSE164677. Results are expressed in $\log_2$(FPKM).

* *p* ≤ 0,05, ** *p* ≤ 0,01, *** *p* ≤ 0,001, **** p ≤ 0,0001 (two-tailed Student's *t*-test).

D. qRT-PCR analysis of *HMGN5* mRNA enrichment in native *lncMB3* RNA pull-down fractions. Details as in panel A.

E. IntaRNA energy map representing the predicted stability of RNA-RNA interaction between *lncMB3* and *HMGN5* mRNA. mRNA and lncRNA sequences are positioned along the x- and y-axes, respectively. The free energy of predicted intramolecular pairs ranges from red (higher energy, unstable pairing) to blue (minimal energy, stable pairing).

F. qRT-PCR analysis of *lncMB3* (left panel) and *HMGN5* mRNA (right panel) enrichment in EVEN, ODD and LacZ fractions over Input, from AMT-crosslinked *lncMB3* RNA pull-down experiment in D283 Med cells. Data expressed as percentage of Input, *N* = 1.

G. Western blot analysis of HMGN5 protein levels in D283 Med cells treated for 72 hours with gap_SCR or gap_*lncMB3* (GapmeR #1) or gap_*lncMB3* + Δ16 construct for *lncMB3* recovery. Normalisations were performed relative to GAPDH protein levels. N = 3, * $p \leq 0,05$, (two-tailed Student's t-test).

H. qRT-PCR analysis of *HMGN5* and *GAPDH* upon RPL22-FLAG RIP assay in gap_SCR and gap_*lncMB3* (GapmeR #1) conditions. Analysis of RNA enrichment in IP and IgG fractions over Input. *N* = 1. Data expressed as percentage of Input.

I. qRT-PCR analysis of *HMGN5*, *INHBE*, *SMAD3*, *OTX2*, *BCL2L1* mRNAs in D283 Med cells treated for 72 hours with sh_SCR or sh_HMGN5. Expression levels were compared to sh_SCR sample as control, set as 1. Data (means ± SEM) are expressed in arbitrary units and are relative to *GAPDH* mRNA levels. *N* = 3, * $p \leq 0,05$, ** $p \leq 0,01$ (two- tailed Student's *t*-test).

J. Western blot analysis of c-PARP1 (left panel) and PARP1 (right panel) protein levels in D283 Med cells treated for 72 hours with sh_SCR or sh_HMGN5. N = 3, * $p \leq 0,05$ (two-tailed Student's t-test).

K. Analysis of number of viable D283 Med cells treated for 72 hours with sh_SCR or sh_HMGN5. Data (means ± SEM) are expressed as the number of viable cells, counted by an automated cell counter. *N* = 3, * $p \leq 0,05$ (two-tailed Student's *t*-test).

Figure 5. Analysis of *lncMB3* targeting and DDP treatment on D283 Med cell viability and apoptosis

A. Left Panel: Dose-response analysis of the number of viable D283 Med cells upon DDP treatment. Control cells (NT) were treated with the DDP vehicle (DMSO). Cell counts were conducted 24 hours

after DDP administration. *N* = 1. Middle Panel: percentage of viable D283 Med cells following DDP

administration and *lncMB3* GapmeR transfection (GapmeR #1, middle panel; GapmeR #2, right

panel). Cell counts were conducted 48 hours after DDP/GapmeR treatments.

Double-treated samples (gap_SCR+DDP and gap_*lncMB3*+DDP) were normalised on the viable cell

number of the corresponding transfected-only sample (gap_scr and gap_*lncMB3*, respectively). DDP

concentration used was 10μM. *N* = 4, * p ≤ 0,05 (two- tailed Student's t-test).

B. Apoptosis analysis of D283 Med cells following DDP administration [1μM] and GapmeR #1

transfection. Left panel: representative flow cytometry analysis of PI- and Annexin V-stained D283

Med cells 24 hours after *lncMB3* KD, with or without DPP treatment. Middle panel: quantification of

viable (Annexin V-/PI-), early apoptotic (Annexin V+/PI-), late apoptotic (Annexin V+/PI+) and

necrotic (Annexin V-/PI+) fractions. *N* = 3, * p ≤ 0,05 (comparison between early + late apoptotic

cells). Right panel: percentage of viable D283 Med cells. Data normalisations as in (A). *N* = 3, * p ≤

0,05 (two-way ANOVA).

C. qRT-PCR analysis of TGF-β pathway genes (left panel) and downstream pathway genes (right

panel) in D283 Med cells treated for 24 hours with DDP at different doses (reported on the x-axis).

Control cells were treated with vehicle (DMSO) and set as 1. Data (means ± SEM) are expressed in

arbitrary units and are relative to *GAPDH* mRNA levels. *N* = 3, * *p* ≤ 0,05, ** *p* ≤ 0,01, *** *p* ≤ 0,001

(two- tailed Student's *t*-test).

Figure 6. Synthesis, analysis and application of HFt-HIS-PASE/GapmeR complexes

A. Left panel: representative image of colocalisation analysis between fluorescein-HFt-HIS-PASE and

DAPI in D283 Med cells, 12 hours after administration (magnification 20x). Fluorescein-HFt-HIS PASE

(green) and DAPI (blue, nuclear staining) are on the left, object counts after binarisation are in the

middle and the merge of the two channels is on the right. Right panel: percentage of colocalisation

between fluorescein-HFt-HIS-PASE and DAPI in D283 Med cells. Data (mean ± SEM) were obtained from 8 different fields.

B. Left panel: representative immunofluorescence of fluorescein-HFt-HIS PASE delivered for 24 hours (green), DAPI (blue, nuclear staining) and phalloidin (red, cytoplasmic staining) (magnification 60x). Right panel: percentage of nucleus/cytoplasm subcellular localisation of HFt-HIS PASE in D283 Med cells. Data (mean + SEM) were obtained from 5 different fields.

C. Band migration profiles on agarose gel electrophoresis. Gel was double stained with SYBR Gold for GapmeR visualisation (left panel) and Coomassie Blue for ferritin visualisation (right panel). *Lane 1*: GapmeR standard 2µM; *Lane 3*: GapmeR standard 1µM; *Lane 5*: GapmeR standard 0,5µM; *Lane 6*: HFt-HIS-PASE-GapmeR complex.

D. Size Exclusion Chromatography profile analysis of HFt-HIS-PASE (in black), HFt-HIS-PASE-GapmeR #1 complexes (in red) and GapmeR #1 alone (in blue).

E. qRT-PCR analysis of *lncMB3* levels in D283 Med cells upon treatment with HFt-HIS-PASE-GapmeR #1 complexes, 48 hours after delivery, compared with empty HFt-HIS-PASE, set as 1. Data (means ± SEM) are expressed in arbitrary units and are relative to *GAPDH* mRNA levels. $N = 3$, *** $p ≤ 0,001$ (two- tailed Student's *t*-test).

F. qRT-PCR analysis of TGF-β pathway genes in D283 Med cells treated for 48 hours with HFt-HIS-PASE-GapmeR #1 + 5µM DDP. Expression levels were compared to HFt-HIS-PASE-empty + 5µM DDP, set as 1. Data (means ± SEM) are expressed in arbitrary units and are relative to *GAPDH* mRNA levels. $N = 3$, * $p ≤ 0,05$, ** $p ≤ 0,01$ (two- tailed Student's *t*-test).

G. Analysis of number of viable D283 Med cells treated for 48 hours with HFt-HIS-PASE-GapmeR #1 + 5µM DDP. HFt-HIS-PASE empty + 5µM DDP was used as control condition. Data (means ± SEM) are expressed as the percentage of viable cells, counted by an automated cell counter. $N = 4$, ** $p ≤ 0,01$ (two-tailed Student's *t*-test).

Figure 7. Schematic representation of the MYC-dependent *lncMB3* mechanism of action. Image

made with Biorender (https://biorender.com).

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. *LncMB3* expression – Referred to Figure 1 and 2

A. qRT-PCR analysis of *MYC, lncMB3 and OTX2* in a pool of 10 healthy cerebella (cereb) set as 1, in

D283 Med cells or in HD-MB03 cells. Data (means ± SEM) are expressed in arbitrary units and are

relative to *GAPDH* mRNA levels. N = 3, * p ≤ 0,05, ** p ≤ 0,01 (two- tailed Student's *t*-test).

B. Expression of *lncMB3* from 54 non-diseased tissues, according to Genotype-Tissue Expression

(GTEx) Analysis, Release V8. Levels are shown in Transcripts per Million (TPM). Circles represent

outliers. Details at https://gtexportal.org/home/gene/RP11-95I16.6.

Figure S2. *LncMB3*-dependent transcriptomic analysis in G3 MB cells – Referred to Figure 1

A. qRT-PCR analysis of *lncMB3* in gap_SCR samples *vs* gap_*lncMB3* (GapmeR #1) samples. *LncMB3*

expression levels upon KD was compared to gap_SCR condition, set as 1. Data (means ± SEM) are

expressed in arbitrary units and are relative to *GAPDH* mRNA levels. N = 3, *** p ≤ 0,001 (two- tailed

Student's *t*-test).

B. Left panel: the plot reports, for each sequenced sample, the number of detected genes obtained

from RNA-Seq analysis. Right Panels: Euclidean distance heatmap showing the similarity in gene

expression within the SCR and KD (GapmeR #1) samples. Pearson correlation heatmap showing the

similarity in gene expression within the SCR and KD (GapmeR #1) samples.

C. qRT-PCR analysis of RNA-Seq deregulated transcripts with padj spanning from 0,001 to 0,05 in

gap_SCR *vs* gap_*lncMB3* (GapmeR #1) samples. Expression levels upon KD were compared to

gap_SCR condition, set as 1. Data (means ± SEM) are expressed in arbitrary units and are relative to

GAPDH mRNA levels. N = 3, * p ≤ 0,05, ** p ≤ 0,01, *** p ≤ 0,001 (two- tailed Student's *t*-test).

D. Schematic representation of DEG quantitative selection.

Figure S3. RNA-Seq data mining – Referred to Figure 1

A. qRT-PCR validation of RNA-Seq data in *lncMB3* KD compared to gap_SCR condition, set as 1.

Validated candidates (10 RNAs) are indicated in histograms. Data (means ± SEM) are expressed in

arbitrary units and are relative to *GAPDH* mRNA levels. N = 3, * p ≤ 0,05, ** p ≤ 0,01, *** p ≤ 0,001

(two- tailed Student's *t*-test).

B. GO showing the distribution in clusters of the DEGs in gap_lncMB3 *vs* gap_SCR-treated D283 Med

cells, according to RNA-Seq data: Downregulated (BLUE) and Upregulated (RED) sets referring to

WikiPathways Database. Categories are listed considering -$\log_{10}$(pvalue).

C. GSEA showing the differentially expressed TGF-β pathway components in gap_lncMB3 *vs*

gap_SCR-treated D283 Med cells, according to RNA-seq data.

D. Schematic representation of the relevant validated genes in the tumour (G3 MB, left panel) and

in the lncMB3 KD condition (right panel). Orange squares highlight the TGF-β pathway components.

Figure S4. Validation of *lncMB3* KD specificity – Referred to Figure 1, 2 and 3

A. Schematic representation of GapmeR #1 and GapmeR #2 (blue lines) binding to *lncMB3* (black

line).

B. Left histograms: qRT-PCR analysis of *lncMB3* in D283 Med and HD-MB03 cells upon *lncMB3* KD

(GapmeR #1). Right histograms: qRT-PCR analysis of *lncMB3* D283 Med and HD-MB03 cells upon

lncMB3 KD (GapmeR #2). N = 3, * p ≤ 0,05, ** p ≤ 0,01, *** p ≤ 0,001 (two- tailed Student's).

Figure S5. AGO2 CLIP assay in D283 Med cells – Referred to Figure 4

A. Left Panel: Western blot analysis of AGO2 levels in AGO2- (IP), IgG-immunoprecipitated cell

extracts, and in Input sample, as control. Right panel: qRT-PCR analysis of RNA enrichment over Input

in IP and IgG fractions. *N* = 1. Data expressed as percentage of Input. *GAPDH* and *PTBP1* mRNAs were

used as negative and positive control, respectively.

Figure S6. Analysis of *lncMB3* mechanism of action – Referred to Figure 4

A. Table showing the genes enriched according to the RNA-Seq analysis of two independent native

lncMB3 pull-down assays performed on total extracts of D283 Med cells.

B. Expression of *HMGN5* from 54 healthy tissues, according to GTEx Analysis, Release V8. Levels are

shown in RPKM. Circles represent outliers. Details at https://gtexportal.org/home/gene/HMGN5

C. Table showing the absolute number of molecules per µL for *lncMB3*, *HMGN5* and *GAPDH*,

according to the evaluation performed through digital PCR analysis.

Figure S7. Analysis of *lncMB3* mechanism of action – Referred to Figure 4

A. qRT-PCR analysis of *HMGN5* mRNA in D283 Med cells upon *lncMB3* KD (GapmeR #1). Data (means

± SEM) are expressed in arbitrary units and are relative to *GAPDH* mRNA levels. *N* = 3, (two- tailed

Student's *t*-test).

B. Western blot analysis of RPL22 protein levels in mock-treated or overexpressing FLAG-RPL22

construct D283 Med cells. Normalisations were performed relative to GAPDH protein levels. *N* = 1.

C. Western blot analysis of RPL22 protein levels in fractions (Inp, IP, IgG) from RIP assay performed

in gap_SCR, or gap_*lncMB3* conditions. Normalisations were performed relative to GAPDH protein

levels. *N* = 1.

D. qRT-PCR analysis of pre-mRNAs of the TGF-β and following pathways upon *lncMB3* KD (GapmeR #1) relative to gap_SCR, set as 1. Data (means ± SEM) are expressed in arbitrary units and are relative to *GAPDH* mRNA levels. *N* = 3, * *p* ≤ 0,05, ** *p* ≤ 0,01, *** *p* ≤ 0,001 (two- tailed Student's *t*-test).

E. Western blot analysis of HMGN5 protein levels in D283 Med cells treated with sh_SCR or sh_HMGN5 after 72 hours. Normalisations were performed relative to GAPDH protein levels. *N* = 3, * *p* ≤ 0,05 (two- tailed Student's *t*-test).

F. Table showing the % of downregulation of *INHBE, SMAD3, OTX2 and BCL2L1* upon gap_lncMB3 or sh_HMGN5 in D283 Med cells.

Figure S8. Analysis of *lncMB3* targeting and chemotherapeutic treatments on D283 Med cells – Referred to Figure 5

A. qRT-PCR analysis of *lncMB3* in D283 Med cells treated for 24 hours with DDP at different doses (reported on the x-axis). Control cells were treated with vehicle (DMSO) and set as 1. Data (means ± SEM) are expressed in arbitrary units and are relative to *GAPDH* mRNA levels. *N* = 3, (two- tailed Student's *t*-test).

B. Right panel: dose-response analysis of the number of viable D283 Med cells upon vincristine treatment. Cells were treated with the vincristine vehicle (DMSO), 5, 20 or 50μg/mL vincristine, without GapmeRs or with gap_SCR or gap_*lncMB3* (GapmeR #1). Cell counts were conducted 48 hours after vincristine/GapmeR treatments. *N* = 3, * *p* ≤ 0,05, ** *p* ≤ 0,01 (two- tailed Student's). Left panel: double-treated samples (gap_SCR+vincristine and gap_*lncMB3*+vincristine) were normalised on the viable cell number of the corresponding transfected-only sample (gap_scr and gap_*lncMB3*, respectively). Vincristine concentration was 20μg/mL. *N* = 3, (two- tailed Student's t-test).

1025   **C.** Left panel: quantification of c-PARP protein levels in D283 Med cells upon vincristine treatment in

1026   untreated (NT), gap_SCR or gap_*lncMB3* samples, in the same conditions as in panel B. GAPDH was

1027   used as control. Right panel: western blot analysis of c-PARP protein levels from the experiment

1028   described in left panel. Normalisation was performed on GAPDH protein levels. $N = 1$ for each

1029   condition (drug concentration).

1030

Figure S9 – Molecular analysis related to Hft-HIS-PASE-GapmeR complex delivery – Referred to

Figure 6

A. Schematic representation of Hft-HIS-PASE and Hft-HIS-PASE-GapmeR encapsulation in the protein

cavity. Single domains and molecules are described in the legend. To allow the internal surface of

the protein to be visualised (lighter colours), only 18 monomers out of the 24 are shown. In addition,

for clarity purposes, only 4 out of the 24 histidine tags are shown. The picture has been generated

with Pymol and GNU Image Manipulation Programs.

B. qRT-PCR analysis of *CD71*, *MMP 2* and *MMP 9* mRNA levels in D283 Med (left panel) or HD-MB03

(right panel) cells. Data (means ± SEM) are expressed in arbitrary units and are relative to *GAPDH*

mRNA levels.

C. HFt-HIS-PASE-GapmeR complexes before and after Denarase treatment at 37°C visualised as band

migration profiles on agarose gel electrophoresis. *Lane 1:* HFt-HIS-PASE-GapmeR complex; *Lane 2:*

HFt-HIS-PASE-GapmeR complexes after overnight incubation with Denarase; *Lane 3*: HFt-HIS-PASE-

GapmeR complexes after 1-hour incubation with Denarase. Gel was stained with SYBR Gold.

D. qRT-PCR analysis of *lncMB3* levels upon KD with GapmeR #1 (gap_*lncMB3*) or Hft-HIS-PASE-

GapmeR complexes (HFt-gap_*lncMB3*), relative to gap_SCR, set as 1. Data are expressed in arbitrary

units and are relative to *GAPDH* mRNA levels. $N = 1$.

1048

ACKNOWLEDGEMENTS

We acknowledge M. Marchioni and J. Genovese for their experimental assistance, C. Grelloni and A. Giuliani for their technical and analytical suggestions and E. Caffarelli for valuable discussions and laboratory support. FRP was supported by an AIRC fellowship. Confocal microscopy acquisitions were conducted at the Imaging Facility of CLN²S@Sapienza, Istituto Italiano di Tecnologia, Rome. Cytofluorimeters were available at IBPM-CNR. RNA-Seq were performed by DNA Link (Seoul, KR) and by Eclipse Bioinnovation (San Diego, CA).

In memory of M. Arceci, beloved friend and colleague.

FUNDING

This work was supported by grants: from MUR (PRIN 2017, id. 2017P352Z4) to PL; from the European Union - Next Generation EU, within the MUR National Recovery and Resilience Plan (NRRP) - M4C2 - Action 1.4 - Call "Potenziamento strutture di ricerca e di campioni nazionali di R&S", Project CN3 "National Center for Gene Therapy and Drugs based on RNA Technology", no. CN00000041 (Spoke #6 "RNA Drug Development", CUP B83C22002860006 to PL, and Spoke 3 "Neurodegeneration", CUP B83C22002870006 to IB and MB); from the European Union - Next-Generation EU within the MUR National Recovery and Resilience Plan [NRRP] - M4C2 - Action 1.1, Call "PRIN 2022" (id. 2022BYB33L, CUP: B53D23016090006 to MB and PL, and id. 2022WC7BL2, CUP B53D23016520006 to EF); from the European Union - Next GenerationEU within the MUR National Recovery and Resilience Plan [NRRP] - M4C2, Action. 1.1, Call "PRIN 2022 PNRR" – id. P2022FFEWN RNA2FUN, CUP B53D23026140001 to MB; from the National Research Council of Italy (projects DBA.AD005.225-NUTRAGE-FOE2021 and DSB.AD006.371-InvAt-FOE2022) to PL; from Sapienza University (RM12117A5DE7A45B and RM123188F6B80CE4) to MB; from AIRC (IG 2020 N° 24942) to DT and AIRC (IG 2019 Id.23053) to IB and ERC-2019-SyG (855923-ASTRA) to IB.

AUTHOR CONTRIBUTIONS

CRedIT author statement. Conceptualisation: AG, PL; Investigation: AG (gene expression analysis, cloning, molecular and cellular assays), PT (interaction assays, imaging and microscopy), FRP (flow cytometry), EF and GT (nanovector synthesis and characterisation), AB (gene expression analysis), AP, FM and AC (bioinformatic analysis); Formal Analysis: AG, PT, FRP, AP, FM, AB and AC; Validation: AG, PT, FRP, EF, AB and GT; Methodology: AG, PT, FRP, EF, JR and GT; Data Curation: AG, PT, AP, FM and AC; Software; AP, FM and AC; Visualisation: AG and JR; Resources: AC, IB, DT, MB and EF; Supervision: AC, DT, MB, PC and PL; Writing – original draft (main text): AG, MB and PL; Funding Acquisition and Project administration: PL.

DATA AVAILABILITY

The RNA-Seq data presented in this study are available in GEO, accession numbers GSE277976 and GSE278666.

COMPETING INTERESTS

The authors declare no competing interests.

INFORMED CONSENT

Manuscript is approved by all authors for publication.

REFERENCES

[1]   P. A. Northcott *et al.*, "Medulloblastoma," *Nat Rev Dis Primers*, vol. 5, no. 1, p. 11, Feb. 2019, doi: 10.1038/s41572-019-0063-6.

1097   [2]   Q. T. Ostrom *et al.*, "CBTRUS Statistical Report: Primary Brain and Other Central Nervous
1098   System Tumors Diagnosed in the United States in 2012–2016," *Neuro Oncol*, vol. 21, no.
1099   Supplement_5, pp. v1–v100, Nov. 2019, doi: 10.1093/neuonc/noz150.

1100   [3]   M. C. Vladoiu *et al.*, "Childhood cerebellar tumours mirror conserved fetal transcriptional
1101   programs," *Nature*, vol. 572, no. 7767, pp. 67–73, Aug. 2019, doi: 10.1038/s41586-019-
1102   1158-7.

1103   [4]   F. M. G. Cavalli *et al.*, "Intertumoral Heterogeneity within Medulloblastoma Subgroups,"
1104   *Cancer Cell*, vol. 31, no. 6, pp. 737-754.e6, Jun. 2017, doi: 10.1016/j.ccell.2017.05.005.

1105   [5]   M. F. Roussel and J. L. Stripay, "Epigenetic Drivers in Pediatric Medulloblastoma," *The
1106   Cerebellum*, vol. 17, no. 1, pp. 28–36, Feb. 2018, doi: 10.1007/s12311-017-0899-9.

1107   [6]   M. F. Roussel and G. W. Robinson, "Role of MYC in Medulloblastoma," *Cold Spring Harb
1108   Perspect Med*, vol. 3, no. 11, pp. a014308–a014308, Nov. 2013, doi:
1109   10.1101/cshperspect.a014308.

1110   [7]   J. S. Mattick *et al.*, "Long non-coding RNAs: definitions, functions, challenges and
1111   recommendations," *Nat Rev Mol Cell Biol*, vol. 24, no. 6, pp. 430–447, Jun. 2023, doi:
1112   10.1038/s41580-022-00566-8.

1113   [8]   O. Beylerli, I. Gareev, A. Sufianov, T. Ilyasova, and Y. Guang, "Long noncoding RNAs as
1114   promising biomarkers in cancer," *Noncoding RNA Res*, vol. 7, no. 2, pp. 66–70, Jun. 2022,
1115   doi: 10.1016/j.ncrna.2022.02.004.

1116   [9]   P. Laneve and E. Caffarelli, "The Non-coding Side of Medulloblastoma," *Front Cell Dev Biol*,
1117   vol. 8, May 2020, doi: 10.3389/fcell.2020.00275.

1118   [10]   J. Rea *et al.*, "Identification and Functional Characterization of Novel MYC-Regulated Long
1119   Noncoding RNAs in Group 3 Medulloblastoma," *Cancers (Basel)*, vol. 13, no. 15, p. 3853, Jul.
1120   2021, doi: 10.3390/cancers13153853.

1121   [11]   M. Morabito *et al.*, "An autocrine ActivinB mechanism drives <scp>TGF</scp> β/Activin
1122   signaling in Group 3 medulloblastoma," *EMBO Mol Med*, vol. 11, no. 8, Aug. 2019, doi:
1123   10.15252/emmm.201809830.

1124   [12]   D. C. Adamson *et al.*, "OTX2 Is Critical for the Maintenance and Progression of Shh-
1125   Independent Medulloblastomas," *Cancer Res*, vol. 70, no. 1, pp. 181–191, Jan. 2010, doi:
1126   10.1158/0008-5472.CAN-09-2331.

1127   [13]   F. Marrocco *et al.*, "Nose-to-brain selective drug delivery to glioma via ferritin-based
1128   nanovectors reduces tumor growth and improves survival rate," *Cell Death Dis*, vol. 15, no.
1129   4, p. 262, Apr. 2024, doi: 10.1038/s41419-024-06653-2.

1130   [14]   J. Kurreck, "Design of antisense oligonucleotides stabilized by locked nucleic acids," *Nucleic
1131   Acids Res*, vol. 30, no. 9, pp. 1911–1918, May 2002, doi: 10.1093/nar/30.9.1911.

[15] A. Weiss and L. Attisano, "The TGFbeta Superfamily Signaling Pathway," *WIREs*
Developmental Biology, vol. 2, no. 1, pp. 47–63, Jan. 2013, doi: 10.1002/wdev.86.

[16] Y. He *et al.*, "Targeting PI3K/Akt signal transduction for cancer therapy," *Signal Transduct*
Target Ther, vol. 6, no. 1, p. 425, Dec. 2021, doi: 10.1038/s41392-021-00828-5.

[17] A. B. Baba *et al.*, "Transforming Growth Factor-Beta (TGF-β) Signaling in Cancer-A Betrayal
Within," *Front Pharmacol*, vol. 13, Feb. 2022, doi: 10.3389/fphar.2022.791272.

[18] D. Vanhoutte *et al.*, "Thbs1 regulates skeletal muscle mass in a TGFβ-Smad2/3-ATF4-
dependent manner," *Cell Rep*, vol. 43, no. 5, p. 114149, May 2024, doi:
10.1016/j.celrep.2024.114149.

[19] B. Schmierer and C. S. Hill, "TGFβ–SMAD signal transduction: molecular specificity and
functional flexibility," *Nat Rev Mol Cell Biol*, vol. 8, no. 12, pp. 970–982, Dec. 2007, doi:
10.1038/nrm2297.

[20] D. Onichtchouk *et al.*, "Silencing of TGF-β signalling by the pseudoreceptor BAMBI," *Nature*,
vol. 401, no. 6752, pp. 480–485, Sep. 1999, doi: 10.1038/46794.

[21] C.-R. Chen, Y. Kang, P. M. Siegel, and J. Massagué, "E2F4/5 and p107 as Smad Cofactors
Linking the TGFβ Receptor to c-myc Repression," *Cell*, vol. 110, no. 1, pp. 19–32, Jul. 2002,
doi: 10.1016/S0092-8674(02)00801-2.

[22] A. Nishida *et al.*, "Otx2 homeobox gene controls retinal photoreceptor cell fate and pineal
gland development," *Nat Neurosci*, vol. 6, no. 12, pp. 1255–1263, Dec. 2003, doi:
10.1038/nn1155.

[23] A. Garancher *et al.*, "NRL and CRX Define Photoreceptor Identity and Reveal Subgroup-
Specific Dependencies in Medulloblastoma," *Cancer Cell*, vol. 33, no. 3, pp. 435-449.e6, Mar.
2018, doi: 10.1016/j.ccell.2018.02.006.

[24] C. Soldani and A. I. Scovassi, "Poly(ADP-ribose) polymerase-1 cleavage during apoptosis: An
update," *APOPTOSIS*, vol. 7, no. 4, pp. 321–328, 2002, doi: 10.1023/A:1016119328968.

[25] P. Tollis *et al.*, "The long noncoding RNA nHOTAIRM1 is necessary for differentiation and
activity of iPSC-derived spinal motor neurons," *Cell Death & Disease 2023 14:11*, vol. 14, no.
11, pp. 1–14, Nov. 2023, doi: 10.1038/s41419-023-06196-y.

[26] P. Laneve *et al.*, "The long noncoding RNA linc-NeD125 controls the expression of
medulloblastoma driver genes by microRNA sponge activity," *Oncotarget*, vol. 8, no. 19, pp.
31003–31015, May 2017, doi: 10.18632/oncotarget.16049.

[27] C. Grelloni *et al.*, "BRCA1 levels and DNA-damage response are controlled by the
competitive binding of circHIPK3 or FMRP to the BRCA1 mRNA," *Mol Cell*, Oct. 2024, doi:
10.1016/j.molcel.2024.09.016.

[28] M. Rochman, C. Malicet, and M. Bustin, "HMGN5/NSBP1: A new member of the HMGN
protein family that affects chromatin structure and function," *Biochimica et Biophysica Acta*
(BBA) - Gene Regulatory Mechanisms, vol. 1799, no. 1–2, pp. 86–92, Jan. 2010, doi:
10.1016/j.bbagrm.2009.09.012.

[29] C. Malicet, M. Rochman, Y. Postnikov, and M. Bustin, "Distinct Properties of Human HMGN5
Reveal a Rapidly Evolving but Functionally Conserved Nucleosome Binding Protein," *Mol Cell*
Biol, vol. 31, no. 13, pp. 2742–2755, Jul. 2011, doi: 10.1128/MCB.05216-11.

[30] Y. Gan *et al.*, "Knockdown of HMGN5 increases the chemosensitivity of human urothelial
bladder cancer cells to cisplatin by targeting PI3K/Akt signaling," *Oncol Lett*, Sep. 2017, doi:
10.3892/ol.2017.7045.

[31] J. Mou *et al.*, "HMGN5 Escorts Oncogenic STAT3 Signaling by Regulating the Chromatin
Landscape in Breast Cancer Tumorigenesis," *Molecular Cancer Research*, vol. 20, no. 12, pp.
1724–1738, Dec. 2022, doi: 10.1158/1541-7786.MCR-22-0241.

[32] E. Sanz, J. C. Bean, D. P. Carey, A. Quintana, and G. S. McKnight, "RiboTag: Ribosomal Tagging
Strategy to Analyze Cell-Type-Specific mRNA Expression In Vivo," *Curr Protoc Neurosci*, vol.
88, no. 1, Jun. 2019, doi: 10.1002/cpns.77.

[33] J. Enríquez Pérez, S. Fritzell, J. Kopecky, E. Visse, A. Darabi, and P. Siesjö, "The effect of locally
delivered cisplatin is dependent on an intact immune function in an experimental glioma
model," *Sci Rep*, vol. 9, no. 1, p. 5632, Apr. 2019, doi: 10.1038/s41598-019-42001-7.

[34] S. Ghosh, "Cisplatin: The first metal based anticancer drug," *Bioorg Chem*, vol. 88, p. 102925,
Jul. 2019, doi: 10.1016/j.bioorg.2019.102925.

[35] C. Seidel, S. Heider, P. Hau, A. Glasow, S. Dietzsch, and R.-D. Kortmann, "Radiotherapy in
Medulloblastoma—Evolution of Treatment, Current Concepts and Future Perspectives,"
Cancers (Basel), vol. 13, no. 23, p. 5945, Nov. 2021, doi: 10.3390/cancers13235945.

[36] E. Falvo *et al.*, "The presence of glutamate residues on the PAS sequence of the stimuli-
sensitive nano-ferritin improves in vivo biodistribution and mitoxantrone encapsulation
homogeneity," *Journal of Controlled Release*, vol. 275, pp. 177–185, Apr. 2018, doi:
10.1016/j.jconrel.2018.02.025.

[37] J. He, S. Xu, and A. J. Mixson, "The Multifaceted Histidine-Based Carriers for Nucleic Acid
Delivery: Advances and Challenges," *Pharmaceutics*, vol. 12, no. 8, p. 774, Aug. 2020, doi:
10.3390/pharmaceutics12080774.

[38] D. N. Louis *et al.*, "The 2007 WHO Classification of Tumours of the Central Nervous System,"
Acta Neuropathol, vol. 114, no. 2, pp. 97–109, Aug. 2007, doi: 10.1007/s00401-007-0243-4.

[39] E. M. Thompson *et al.*, "Prognostic value of medulloblastoma extent of resection after
accounting for molecular subgroup: a retrospective integrated clinical and molecular

analysis," *Lancet Oncol*, vol. 17, no. 4, pp. 484–495, Apr. 2016, doi: 10.1016/S1470-
2045(15)00581-1.

[40] K. Juraschka and M. D. Taylor, "Medulloblastoma in the age of molecular subgroups: a
review," *J Neurosurg Pediatr*, vol. 24, no. 4, pp. 353–363, Oct. 2019, doi:
10.3171/2019.5.PEDS18381.

[41] O. Menyhárt, F. Giangaspero, and B. Győrffy, "Molecular markers and potential therapeutic
targets in non-WNT/non-SHH (group 3 and group 4) medulloblastomas," *J Hematol Oncol*,
vol. 12, no. 1, p. 29, Dec. 2019, doi: 10.1186/s13045-019-0712-y.

[42] Y. Dong, R. Tu, H. Liu, and G. Qing, "Regulation of cancer cell metabolism: oncogenic MYC in
the driver's seat," *Signal Transduct Target Ther*, vol. 5, no. 1, p. 124, Jul. 2020, doi:
10.1038/s41392-020-00235-2.

[43] V. Llombart and M. R. Mansour, "Therapeutic targeting of 'undruggable' MYC,"
EBioMedicine, vol. 75, p. 103756, Jan. 2022, doi: 10.1016/j.ebiom.2021.103756.

[44] E. Garralda *et al.*, "MYC targeting by OMO-103 in solid tumors: a phase 1 trial," *Nat Med*,
vol. 30, no. 3, pp. 762–771, Mar. 2024, doi: 10.1038/s41591-024-02805-1.

[45] A. Dent *et al.*, "HAVOC: Small-scale histomic mapping of cancer biodiversity across large
tissue distances using deep neural networks," *Sci Adv*, vol. 9, no. 39, Sep. 2023, doi:
10.1126/sciadv.adg1894.

[46] M. Coan, S. Haefliger, S. Ounzain, and R. Johnson, "Targeting and engineering long non-
coding RNAs for cancer therapy," *Nat Rev Genet*, Feb. 2024, doi: 10.1038/s41576-024-
00693-2.

[47] V. Ferrucci *et al.*, "Metastatic group 3 medulloblastoma is driven by PRUNE1 targeting
NME1–TGF-β–OTX2–SNAIL via PTEN inhibition," *Brain*, vol. 141, no. 5, pp. 1300–1319, May
2018, doi: 10.1093/brain/awy039.

[48] "HMGN5, an RNA or Nucleosome binding protein - potentially switching between the
substrates to regulate gene expression", doi: 10.1101/2022.07.15.500188.

[49] L. Li *et al.*, "Ferritin-mediated siRNA delivery and gene silencing in human tumor and
primary cells," *Biomaterials*, vol. 98, pp. 143–151, Aug. 2016, doi:
10.1016/j.biomaterials.2016.05.006.

[50] F. R. Liberati *et al.*, "Combined Delivery of miR-15/16 through Humanized Ferritin Nanocages
for the Treatment of Chronic Lymphocytic Leukemia," *Pharmaceutics*, vol. 16, no. 3, p. 402,
Mar. 2024, doi: 10.3390/pharmaceutics16030402.

1233

FIGURE 1

A

B

C

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

FIGURE 7