DprA recruits ComM to facilitate recombination during natural transformation in Gram-negative bacteria

Triana N. Dalia[1], Mérick Machouri[2,3], Céline Lacrouts[2,3], Yoann Fauconnet[4], Jessica Andreani[4,*], Raphaël Guerois[4,*], J. Pablo Radicella[2,3,*], and Ankur B. Dalia[1,*]

[1]Department of Biology, Indiana University, Bloomington, Indiana, United States of America
[2]Université Paris-Saclay, iRCM/IBFJ, CEA, Genetic Stability, Stem Cells and Radiation, F-92260 Fontenay-aux-Roses, France
[3]Université de Paris Cité, iRCM/IBFJ, CEA, Genetic Stability, Stem Cells and Radiation, F-92260 Fontenay-aux-Roses, France
[4]Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198, Gif-sur-Yvette, France

*Authors for correspondence: jessica.andreani@cea.fr, raphael.guerois@cea.fr, pablo.radicella@cea.fr, and ankdalia@iu.edu

ABTRACT

Natural transformation (NT) represents one of the major modes of horizontal gene transfer in bacterial species. During NT, cells can take up free DNA from the environment and integrate it into their genome by homologous recombination. While NT has been studied for >90 years, the molecular details underlying this recombination remain poorly understood. Recent work has demonstrated that ComM is an NT-specific hexameric helicase that promotes recombinational branch migration in Gram-negative bacteria. How ComM is loaded onto the post-synaptic recombination intermediate during NT, however, remains unclear. Another NT-specific recombination mediator protein that is ubiquitously conserved in both Gram-positive and Gram-negative bacteria is DprA. Here, we uncover that DprA homologs in Gram-negative species contain a C-terminal winged helix domain that is predicted to interact with ComM by AlphaFold. Using *Helicobacter pylori* and *Vibrio cholerae* as model systems, we demonstrate that ComM directly interacts with the DprA winged-helix domain, and that this interaction is critical for DprA to recruit ComM to the recombination site to promote branch migration during NT. These results advance our molecular understanding of recombination during this conserved mode of horizontal gene transfer. Furthermore, they demonstrate how structural modeling can help uncover unexpected interactions between well-studied proteins to provide deep mechanistic insight into the molecular coordination required for their activity.

SIGNIFICANCE STATEMENT

Bacteria can acquire novel traits like antibiotic resistance and virulence through horizontal gene transfer by natural transformation. During this process, cells take up free DNA from the environment and integrate it into their genome by homologous recombination. Many of the molecular details underlying this process, however, remain incompletely understood. In this study, we identify a new protein-protein interaction between ComM and DprA, two factors that promote homologous recombination during natural transformation in Gram-negative species. Through a combination of bioinformatics, structural modeling, cell biological assays, and complementary genetic approaches, we demonstrate that this interaction is required for DprA to recruit ComM to the site of homologous recombination.

INTRODUCTION

52
53
Natural transformation (NT) is a broadly conserved mechanism of horizontal gene transfer
present in diverse Gram-positive and Gram-negative eubacteria, as well as some archaeal
species[1,2]. During this process, cells translocate single-stranded transforming DNA (tDNA)
across the cytoplasmic membrane[3-5], which can then be integrated into the genome by
homologous recombination. This can promote the acquisition of novel traits including antibiotic
resistance and virulence factors in bacterial pathogens.
60
During NT, the primary bacterial recombinase, RecA, binds to tDNA and both (1) facilitates a
homology search and (2) initiates the recombination reaction by promoting strand invasion of
the bacterial chromosome to generate a 3-stranded D-loop structure. This 3-stranded
intermediate is distinct from the more canonical recombination intermediates that form between
two dsDNA substrates during DNA repair. For example, the 3-stranded intermediate during NT
can never form a classical Holliday junction. As a result, the recombination reaction that occurs
during NT must be distinct from more canonical forms of dsDNA recombination. Consistent with
this, there are a number of proteins that have been characterized that are uniquely required for
recombination during NT.
70
71 One of those proteins is DprA, which is present in all naturally transformable species[2]. DprA is a
72 ssDNA-binding recombination mediator protein. During NT, DprA has been implicated in both (1)
73 protecting ssDNA that has been internalized into the cytoplasm and (2) facilitating recombination
74 by aiding in the recruitment of RecA[6-10]. Thus, to date, DprA has primarily been associated with
75 the early pre-synaptic steps of recombination that lead to formation of the D-loop.
76
77 Another NT-specific recombination protein is ComM, which is only conserved in Gram-negative
78 naturally transformable species[11]. ComM is a NT-specific helicase that binds to the crossover
79 junctions (*i.e.*, branches) of the D-loop to expand them via recombinational branch migration[11,12].
80 This activity facilitates the integration of heterologous sequences during NT.
81
82 While most of the proteins involved in recombination during NT have been identified and
83 characterized to some degree, the molecular coordination between these factors remains poorly
84 understood. Here, we uncover that an extended C-terminal winged-helix domain of Gram-
85 negative DprA homologs helps to recruit ComM to the site of tDNA recombination.
86
87

RESULTS

88
89
ComM-DprA interactions promote NT in Helicobacter pylori
90
DprA is broadly conserved in bacterial species and is always found in species capable of NT[2].
Prior work characterized the structure of the C-terminal winged helix domain of *Helicobacter*
pylori DprA, which is present in a subset of DprA proteins[13]. The DprA$_{Hp}$ C-terminal domain is
not involved in DNA-binding despite displaying a winged helix fold[13,14]. Because this domain is
required for efficient NT, it was proposed that its function could be to establish interactions with
other NT-specific proteins[13]. We therefore analyzed DprA in a diverse set of bacterial species
and uncovered that the DprA homologs among Gram-negative species commonly encode this
extended C-terminal winged helix domain, but it is absent in the DprA homologs of Gram-
positive species (**Fig. S1, Dataset S1**). This distribution largely mirrors the distribution of ComM,
where species that contain a ComM homolog also encode a DprA homolog with a C-terminal
winged-helix domain (**Fig. S1, Dataset S1**).

102
Thus, we hypothesized that the DprA winged-helix domain may directly interact with ComM to
facilitate NT. To test this, we first used *H. pylori* as a model system. Consistent with our
hypothesis, AlphaFold 3 (AF3)[15] predicts a high-confidence interaction between the DprA$_{Hp}$
winged-helix domain and ComM$_{Hp}$ (ipTM = 0.76; pTM = 0.86)(**Fig. 1A**). To test whether these
proteins interact empirically, we performed bacterial adenylate cyclase two-hybrid (BACTH)
assays between the DprA$_{Hp}$$^{C\text{-}ter}$ (winged-helix domain) and full length ComM. In these assays,
LacZ activity is directly linked to the interaction of these proteins[16]. This BACTH analysis
revealed that DprA$_{Hp}$ and ComM$_{Hp}$ interact with one another (**Fig. 1B**, **Fig. S2**).
111
To test whether the interaction between ComM$_{Hp}$ and DprA$_{Hp}$ is functionally relevant, we sought
to assess the impact of disrupting these interactions on NT frequency in *H. pylori*. For these
assays, *dprA$_{Hp}$* and *comM$_{Hp}$* alleles were expressed ectopically at the *ureA* locus. Importantly,
ectopic expression of the wildtype alleles (*comM$_{Hp}$* and *dprA$_{Hp}$*) restored NT in the strains where
the native gene was deleted (**Fig. 1C**). When cells were complemented with an allele of *dprA*
lacking the C-terminal winged-helix domain (*dprA$_{Hp}$$^{\Delta C\text{-}ter}$*) the NT frequency was indistinguishable
from Δ*comM$_{Hp}$* (**Fig. 1C**). Importantly, *dprA$_{Hp}$$^{\Delta C\text{-}ter}$* transformed at rates significantly higher than
Δ*dprA*, suggesting that deleting the DprA$_{Hp}$ winged-helix did not abolish all DprA activity. Thus,
we hypothesized that the reduced NT frequency observed in *dprA$_{Hp}$$^{\Delta C\text{-}ter}$* was due to ablation of
DprA$_{Hp}$-ComM$_{Hp}$ interactions. To test this more rigorously, we assessed the phenotype of a
dprA$_{Hp}$$^{\Delta C\text{-}ter}$ Δ*comM$_{Hp}$* double mutant. If the reduced NT observed in *dprA$_{Hp}$$^{\Delta C\text{-}ter}$* is due to ablation
of DprA$_{Hp}$-ComM$_{Hp}$ interactions, we hypothesized that these mutations should be epistatic.
Indeed, we found that the phenotype of *dprA$_{Hp}$$^{\Delta C\text{-}ter}$* Δ*comM$_{Hp}$* double mutant was not
significantly different from either the *dprA$_{Hp}$$^{\Delta C\text{-}ter}$* or Δ*comM$_{Hp}$* single mutants (**Fig. 1C**). This result
is consistent with DprA$_{Hp}$-ComM$_{Hp}$ interactions facilitating NT.
127
To test this further, we analyzed the AF3 model of DprA$_{Hp}$-ComM$_{Hp}$ interactions and found that
DprA$_{Hp}$^{L252} is buried within the interface of these two proteins (**Fig. 1A**). Thus, we hypothesized
that introducing a charged amino acid at this site should disrupt DprA$_{Hp}$-ComM$_{Hp}$ interactions.
Consistent with this, we found that DprA$_{Hp}$^{L252E} no longer interacts with ComM$_{Hp}$ in BACTH
assays (**Fig. 1B**). Moreover, *dprA$_{Hp}$^{L252E}* phenocopies *dprA$_{Hp}$$^{\Delta C\text{-}ter}$* in NT frequency assays (**Fig.**
1C). All together, these data strongly suggest that DprA$_{Hp}$-ComM$_{Hp}$ interactions are critical for
ComM$_{Hp}$ activity in *H. pylori*.
135
ComM-DprA interactions promote recombination of tDNA during NT in V. cholerae
The DprA C-terminal winged-helix domain is widely distributed among Gram-negative bacteria
(**Fig. S1**). This includes *Vibrio cholerae*, the organism in which ComM was first characterized as
a branch migration factor, and a species in which many tools have already been established for
studying ComM activity[11,17]. So next, we sought to use *V. cholerae* as a model system to
elucidate (1) whether ComM-DprA interactions are broadly conserved and if so, (2) how DprA-
ComM interactions mechanistically promote NT.
143
144 First, we found that AF3 predicted a high confidence interaction between the DprA$_{Vc}$ winged-
145 helix domain and ComM$_{Vc}$ (ipTM = 0.92; pTM = 0.85) (**Fig. 2A**). Consistent with this, we found
146 that the DprA$_{Vc}$ winged-helix domain and ComM$_{Vc}$ interact with one another in BACTH assays
147 (**Fig. 2B**).
148
To test whether the interaction between these proteins is functionally relevant, we mutated 4
residues that lie at the predicted DprA$_{Vc}$-ComM$_{Vc}$ interface (**Fig. 2A**). Two of these residues
represented a potential electrostatic interaction (*comM$_{Vc}$^{R497}*, *dprA$_{Vc}$^{E353}*), so we mutated these
residues by "flipping" their charge: *comM$_{Vc}$^{R497E}* and *dprA$_{Vc}$^{E353R}*. We also mutated two leucine

residues buried within the interface ($comM_{Vc}^{L455}$, $dprA_{Vc}^{L350}$) to a charged amino acid:
$comM_{Vc}^{L455E}$, $dprA_{Vc}^{L350E}$. Three of these alleles – $DprA_{Vc}^{E353R}$, $ComM_{Vc}^{L455E}$, $DprA_{Vc}^{L350E}$ –
disrupted $DprA_{Vc}$-$ComM_{Vc}$ interactions, while $ComM_{Vc}^{R497E}$ did not (**Fig. 2B**). Consistent with
$DprA_{Vc}$-$ComM_{Vc}$ interactions playing a functional role in *V. cholerae*, the 3 alleles that disrupted
interactions ($dprA_{Vc}^{E353R}$, $comM_{Vc}^{L455E}$, $dprA_{Vc}^{L350E}$) all phenocopied $\Delta comM$ in NT frequency
assays, while the allele that did not disrupt interactions ($comM_{Vc}^{R497E}$) had no effect on NT (**Fig.**
2C). Importantly, the point mutations in *dprA* ($dprA_{Vc}^{E353R}$ and $dprA_{Vc}^{L350E}$), did not completely
inhibit NT (as in $\Delta dprA$), indicating that some functions of DprA remain intact (**Fig. 2C**).
Furthermore, $dprA_{Vc}^{E353R}$ and $dprA_{Vc}^{L350E}$ were both epistatic with $\Delta comM$ (**Fig. 2C**). Together,
this suggests that disrupting $DprA_{Vc}$-$ComM_{Vc}$ interactions disrupts $ComM_{Vc}$ activity during NT,
which is consistent with what was observed above in *H. pylori* (**Fig. 1**).
164
To more rigorously test this model, we took advantage of the potential electrostatic interaction
between $ComM_{Vc}^{R497}$-$DprA_{Vc}^{E353}$. Our results demonstrate that flipping the charge on DprA
($dprA_{Vc}^{E353R}$) disrupts $DprA_{Vc}$-$ComM_{Vc}$ interactions and ComM activity, which may be due to
electrostatic repulsion between $DprA_{Vc}^{E353R}$ and $ComM_{Vc}^{R497}$ based on the AF3 model. If so, we
hypothesized that flipping the charge of the interacting residue on ComM ($ComM_{Vc}^{R497E}$) should
restore ComM activity by restoring $DprA_{Vc}$-$ComM_{Vc}$ interactions. Indeed, we found that NT in
$dprA_{Vc}^{E353R}$ $comM_{Vc}^{R497E}$ was partially restored when compared to the $dprA_{Vc}^{E353R}$ single mutant
(**Fig. 2C**). Importantly, NT of $dprA_{Vc}^{L350E}$ $comM_{Vc}^{R497E}$ was not restored when compared to
$dprA_{Vc}^{L350E}$ (**Fig. S3A**). This indicates that there is specificity to the recovery observed in
$dprA_{Vc}^{E353R}$ $comM_{Vc}^{R497E}$, and demonstrates that $comM_{Vc}^{R497E}$ is not simply a gain-of-function
allele of ComM. This data strongly suggests that $DprA_{Vc}^{E353}$ and $ComM_{Vc}^{R497}$ interact, which
further validates the AF3 model.
177
178 While we see partial recovery for NT in $dprA_{Vc}^{L350E}$ $comM_{Vc}^{R497E}$, we did not see recovery of
179 interaction with these proteins in BACTH assays (**Fig. 2B**). One possibility is that the recovered
180 protein-protein interaction is below the detection limit for this assay. This is supported, in part, by
181 the fact that NT is only partially restored in $dprA_{Vc}^{E353R}$ $comM_{Vc}^{R497E}$.
182
Prior work shows that ComM is not required for the uptake of tDNA into the cytoplasm, instead,
it is only required for the recombination of tDNA into the genome[11,18]. Consistent with this, NT of
$\Delta comM$ is indistinguishable from the parent when a replicating plasmid is used as the tDNA[11],
which does not require recombination (**Fig. 2D**). DprA, however, has a presynaptic role in
protecting tDNA from being degraded in the cytoplasm prior to tDNA integration[6]. Consistent
with that, NT of $\Delta dprA$ is significantly reduced when a replicating plasmid is used as tDNA (**Fig.**
2D). The point mutations in $DprA_{Vc}$ that disrupt $DprA_{Vc}$-$ComM_{Vc}$ interactions ($dprA_{Vc}^{L350E}$ and
$dprA_{Vc}^{E353R}$), however, do not affect NT with a replicating plasmid. This suggests that these DprA
alleles retain their presynaptic function; namely, the ability to protect tDNA in the cytoplasm.
192
$DprA_{Vc}$-$ComM_{Vc}$ interactions are required for branch migration during NT
Our data thus far suggest that DprA-ComM interactions are critical for ComM activity because
mutations that disrupt this interaction phenocopy a $\Delta comM$ mutant. Also, these mutations are
epistatic with a $\Delta comM$ mutation. ComM is required for recombinational branch migration during
NT. Thus, we next sought to test whether $DprA_{Vc}$-$ComM_{Vc}$ interactions are critical for ComM-
dependent branch migration. To do this, we used a previously established assay to evaluate
branch migration activity *in vivo*[11]. This assay indirectly measures branch migration by
assessing the comigration of two linked genetic markers: a Trimethoprim antibiotic resistance
marker (Tm[R]) and a premature stop codon in *lacZ* (*lacZ**) (**Fig. 3A**). We employed two different
tDNA products, one where Tm[R] and *lacZ** are separated by 243 bp and another where they are
separated by 820 bp. Consistent with ComM playing an important role in branch migration

during NT, $\Delta comM$ exhibits significantly reduced comigration for these markers with both tDNA
products (**Fig. 3B**). In this assay, we found that the mutations that disrupt DprA$_{Vc}$-ComM$_{Vc}$
interactions ($dprA_{Vc}^{L350E}$, $dprA_{Vc}^{E353R}$, and $comM_{Vc}^{L455E}$) had a significant reduction in comigration
that was similar to $\Delta comM$ (**Fig. 3B**).
208
Furthermore, we found that comigration in $dprA_{Vc}^{E353R}$ $comM_{Vc}^{R497E}$ was restored when
compared to the $dprA_{Vc}^{E353R}$ single mutant (**Fig. 3B**). In fact, comigration of $dprA_{Vc}^{E353R}$
$comM_{Vc}^{R497E}$ was restored to parental levels, despite the fact that NT is only partially restored in
this background (**Fig. 3B**, **Fig. 2C**). This is consistent with DprA$_{Vc}$-ComM$_{Vc}$ interactions being
restored when the charge on both sides of this electrostatic interaction are flipped, which further
supports the validity of the AF3 model. Importantly, comigration was not restored in $dprA_{Vc}^{L350E}$
$comM_{Vc}^{R497E}$ when compared to $dprA_{Vc}^{L350E}$ indicating that the recovery observed was specific
and not due to $comM_{Vc}^{R497E}$ being a gain-of-function ComM allele (**Fig. S3B**).
217
218
DprA$_{Vc}$ recruits ComM$_{Vc}$ to the recombination site during NT
DprA has predominantly been characterized for the presynaptic role it plays during NT. In
particular, it plays an important role in recruiting RecA to the tDNA via a direct protein-protein
interaction[6-10]. Our data suggest that DprA-ComM interactions are critical for ComM-dependent
branch migration. And while it is clear that ComM activity is required for NT and not for other
forms of recombination, the mechanism by which this specificity is determined is unclear. Thus,
we hypothesized that DprA recruits ComM to the recombination site during NT.
226
We have previously shown that in single cells undergoing NT, a fluorescent fusion of ComM
(GFP-ComM$_{Vc}$) forms foci that are (1) dependent on RecA, (2) only formed in the presence of
exogenous tDNA, (3) specifically formed at the chromosomal locus homologous to the tDNA,
and (4) transient and highly dynamic, which all together indicates that these foci represent
active ComM complexes acting at the recombination site during NT[17]. So, to test whether
DprA$_{Vc}$-ComM$_{Vc}$ interactions help recruit ComM$_{Vc}$ to the recombination site, we assessed GFP-
ComM$_{Vc}$ focus formation in cells incubated with tDNA (**Fig. 4A**). Consistent with prior work, we
found that the parent strain forms many GFP-ComM$_{Vc}$ foci (**Fig. 4B-C**), with ~60% of cells
exhibiting at least 1 GFP-ComM$_{Vc}$ focus within the timelapse imaging window. By contrast, point
mutants that disrupt DprA$_{Vc}$-ComM$_{Vc}$ interactions ($dprA_{Vc}^{L350E}$ and $comM_{Vc}^{L455E}$) reduce GFP-
ComM$_{Vc}$ focus formation to the limit of detection (**Fig. 4B-C**). These data are consistent with a
model in which DprA helps recruit ComM to the recombination site during NT.
239
240
DprA promotes ComM-dependent branch migration for NT in species-specific manner
Despite DprA-ComM interactions being preserved in both *H. pylori* and *V. cholerae*, the
interaction interfaces between these proteins are distinct. For example, the electrostatic
interaction that we demonstrate is critical for DprA$_{Vc}$-ComM$_{Vc}$ interactions, is absent in the
DprA$_{Hp}$-ComM$_{Hp}$ model (**Fig. 1A, 2A**). Consistently, ectopically expressed ComM$_{Hp}$ in *V.*
cholerae could not complement the defect in NT and branch migration of a *V. cholerae* $\Delta comM$
mutant (**Fig. 5A-B**). Thus, we hypothesized that DprA-ComM interactions may be species
specific (*i.e.*, DprA$_{Vc}$ cannot recruit ComM$_{Hp}$ and *vice versa*). If this is the case, ectopic
expression of both DprA$_{Hp}$ and ComM$_{Hp}$ in *V. cholerae* should be sufficient to rescue ComM
activity. First, we find that ectopic expression of DprA$_{Hp}$ in *V. cholerae* results in a slight
decrease in NT frequency (**Fig. S4**), but deletion of native *comM* in this background still
diminishes NT ~100-fold as expected (**Fig. 5**). When DprA$_{Hp}$ and ComM$_{Hp}$ are both ectopically
expressed in *V. cholerae*, we found that NT was unchanged when native *comM* is deleted in this
background (**Fig. 5A**) and branch migration activity was restored (**Fig. 5B**). This result is

consistent with DprA$_{Hp}$ recruiting ComM$_{Hp}$ to facilitate ComM-dependent branch migration
activity in *V. cholerae*. These results also suggest that DprA and ComM represent the "functional
unit" that facilitates branch migration during NT in Gram-negative species.
258
259
260 **DISCUSSION**
261
262 The results presented above advance our understanding of the molecular coordination
263 underlying tDNA integration during horizontal gene transfer by NT. Specifically, they
264 demonstrate that in Gram-negative species, the winged-helix domain of DprA directly binds to
265 ComM to recruit this branch migration factor to the recombination site during NT.
266
Work from our group and others has characterized the molecular and structural basis of ComM-
dependent branch migration of D-loops during NT in Gram-negative species[11,12,19]. D-loop
structures are also generated as intermediates during DNA repair, however, ComM does not
facilitate this process and is instead uniquely required for tDNA integration during NT[11]. This
suggests that ComM may be specifically loaded onto the D-loops generated during NT.
However, the basis for ComM loading has remained elusive. Our results demonstrate DprA is
likely required to load ComM onto D-loops during NT. Thus, DprA-dependent loading of ComM
may explain why this branch migration factor is exclusively associated with recombination
during NT.
276
Our prior work has demonstrated that ComM is a bidirectional helicase that can translocate both
5' → 3' and 3' → 5' on model branched DNA substrates *in vitro*[11]. If this directionality is
unregulated, then ComM would facilitate (*i.e.*, expand the D-loop) and inhibit (*i.e.*, collapse the
D-loop) tDNA integration with equal probability. Indeed, this is exactly what is observed *in vitro*
with purified ComM[11]. Single cell tracking of the spatiotemporal dynamics of tDNA integration,
however, suggest that the recombination reaction is highly efficient[17], which suggests that there
may be additional factors that regulate ComM directionality *in vivo*. Here, we show that DprA-
ComM interactions are required for branch migration activity *in vivo*. Thus, it is tempting to
speculate that DprA-dependent loading of ComM onto the D-loop regulates the directionality of
branch migration, which is something we will test moving forward.
287
This discovery of DprA-dependent recruitment of ComM is notable for a number of reasons.
DprA has primarily been characterized for its presynaptic roles during NT. This includes its role
in protecting single-stranded tDNA in the cytoplasm, eviction of SSB, and recruitment of RecA[6-
10]. Together, these activities help promote RecA-mediated strand invasion to initiate the
recombination process. Our results demonstrate that DprA is also required to initiate branch
migration, which indicates that it also plays a postsynaptic role during NT by recruiting ComM.
DprA has also been implicated in playing a regulatory role in some systems like *Streptococcus*
pneumoniae, where DprA directly interacts with the master regulator of competence, ComE, to
shut-off competence[20]. Due to its varied roles during NT, most of which are facilitated by direct
protein-protein interactions, it is tempting to speculate that DprA serves as the master "hub" that
coordinates this complex process. On a related note, ComM has previously only been
implicated in the postsynaptic step of branch migration. The recruitment of ComM by DprA may
also suggest that ComM aids in a presynaptic step of recombination (*e.g.*, the homology search
or initiation of strand invasion).
302
The highly varied roles for DprA during NT also highlight the value of the approach used in this
study. Because of the critical presynaptic roles that DprA performs during NT, deletion of DprA
completely inhibits NT. Thus, uncovering its postsynaptic role would not have been possible

using classical genetic approaches like transposon screens. While genetic screens can help
identify the genes and proteins necessary for a particular phenotype, they may fail to reveal the
underlying molecular coordination necessary for their activity – especially in cases where factors
have more than one function. The structural modeling approach used here, however, helped
reveal unexpected interactions that led to testable hypotheses, which ultimately uncovered how
these factors coordinate to carry out their activity. Advances in structural modeling are
developing at breakneck speed, and recent work demonstrates that genome-wide unbiased
searches for novel protein-protein interactions are now highly feasible[21,22]. These unbiased
genome-wide *in silico* screens hold great promise in accelerating the discovery of these types of
unexpected interactions moving forward.
316
317
318   **METHODS**
319
Bacterial strains and culture conditions
All *V. cholerae* strains used throughout this study are derivatives of strain E7946[23]. Cells were
routinely grown in LB Miller broth and agar supplemented with spectinomycin (200 µg/mL),
kanamycin (50 µg/mL), trimethoprim (10 µg/mL), tetracycline (0.5 µg/mL), sulfamethoxazole
(100 µg/mL), chloramphenicol (1 µg/mL), erythromycin (10 µg/mL), carbenicillin (20 µg/mL),
and/or zeocin (100 µg/mL) as appropriate.
326
All *H. pylori* strains used in this study are derivatives of strain 26695[24]. Cultures were grown
under microaerophilic conditions (5% O_2, 10% CO_2, using the Don Whitley system) at 37°C.
Blood agar base medium (BAB) supplemented with 10% defibrillated horse blood
(ThermoFisher Scientific) was used for plate cultures. An antibiotic mix containing polymyxin B
(0.155 mg/ml), vancomycin (6.25 mg/ml), trimethoprim (3.125 mg/ml), and amphotericin B
(1.25 mg/ml) was added to both plate and liquid cultures. Additional antibiotics were added as
required: kanamycin (20 µg/ml), apramycin (12.5 µg/ml), and chloramphenicol (8 µg/ml)
streptomycin (10 µg/ml).
335
336
337   ***Construction of mutant strains***
338   For *V. cholerae*, all strains were generated by natural transformation using splicing-by-overlap-
339   extension (SOE) PCR mutant constructs exactly as previously described[25,26]. Ectopic
340   expression constructs were generated by SOE PCR and integrated onto the *V. cholerae*
341   genome exactly as previously described[27].
342
For generating mutant variants in *H. pylori* the specific gene regions (with 200bp of flanking
sequences) were amplified from genomic DNA (gDNA) of *H. pylori* 26695 using sequence
specific primers and cloned into pjET1.2 plasmid using sequence- and ligation-independent
cloning (SLIC)[28]. Replacement of the ORF by antibiotic resistant cassettes (either
chloramphenicol, kanamycin or apramycin) was carried out by SLIC. The constructs were
introduced in *H. pylori* by natural transformation and at least two independent clones were
selected for each construct. For complementation by expression from the *ureA* locus the
corresponding gene ORFs were amplified by PCR and cloned by SLIC into a pjET2.1 plasmid
carrying an antibiotic resistance cassette flanked by 200 bp upstream and downstream of the
ureA gene. The resulting construct was introduced in *H. pylori* by natural transformation.
353
354   All strains were confirmed by PCR and/or sequencing. For a detailed list of all strains, see **Table
355   S1.** For a complete list of oligos used to generate mutant constructs, see **Table S2**.
356

357
Bioinformatic analysis of DprA homologs
Using BLAST[29], we searched for DprA homologs in a set of bacterial genomes where ComM homologs had previously been searched[11]. For all identified DprA homologs, we performed protein structure prediction using AlphaFold v2.3[30] as implemented in the ColabFold v1.5.2 pipeline[31]. We ran MMseqs2[32] against the UniRef30_2202 database to generate multiple sequence alignments used as input to the AlphaFold2 structure prediction module. We then generated 5 structural models for each DprA homolog using 3 recycles. We structurally superimposed the best models (highest pTM score for each species) with PyMOL (v3.0) and aligned them to the experimental structure of the DprA$_{Hp}$ C-terminal domain[13] (PDB identifier: 6GW7). We visually identified the presence of a C-terminal winged helix domain in all species with an identified ComM homolog, apart from *B. burgdoferi* (Fig S1 and Dataset S2). To validate the presence of a C-terminal winged helix domain, we annotated Pfam[33] domains in all DprA homolog sequences by HH-suite homology search[34]: we ran 5 iterations of HHblits against the UniRef30_2202 database to generate multiple sequence alignments, used as input for HHsearch against Pfam v35.0 (with 80% maximum pairwise sequence identity). We identified a winged helix Pfam domain (PF17782, PF10771.12 or PF15977.8) with HHsearch probability above 90% in all species with ComM homologs, except for *E. coli* (probability 88.2%), *F. nucleatum* (probability 89.7%)*, H. pylori* (probability 23%)*, C. jejuni* (probability 31.7%), and *L. interrogans* (probability 59%). Note that the DprA$_{Hp}$ C-terminal domain is not annotated in Pfam despite being structurally characterized.

Natural transformation assays
For *V. cholerae*, strains were engineered to bypass the native signals required for NT. Specifically, strains had P_{tac}-*tfoX* and Δ*luxO* mutations, which ectopically express the master competence regulator (TfoX) and genetically lock cells in a high cell density state as previously described[35]. Chitin-independent transformation assays to determine NT frequency were performed exactly as previously described[35]. Briefly, cells were grown at 30ºC rolling in LB supplemented with 100 µM IPTG, 20 mM MgCl$_2$, 10 mM CaCl$_2$ to late-log phase. Then, ~10^8 CFUs were diluted into instant ocean medium (7g/L; Aquarium Systems). Then, 100 ng of transforming DNA (tDNA) was added to each reaction. The tDNA used to assess transformation frequency was a linear ΔVC1807::Erm^R PCR product. VC1807 is a frame-shifted transposase gene. Control reactions were also performed for each strain where no tDNA was added. Cells were incubated with tDNA overnight at 30ºC static. Then, reactions were outgrown by adding 1mL of LB and shaking at 37ºC for 3 hours. Reactions were then plated for quantitative culture by dribble plating on LB agar supplemented with erythromycin to quantify transformants, and plain LB to quantify total viable counts. Transformation frequency is defined as the CFU/mL of transformants divided by the CFU/mL of total viable counts.

For assays where branch migration was indirectly assessed via linkage analysis in *V. cholerae*, transformation reactions were performed exactly as described above except that they were plated onto LB agar supplemented with trimethoprim and X-gal (40 µg/mL). On these plates, all cells integrated the Tm^R marker, and if cells integrated the genetically linked *lacZ* nonsense mutation they would make white colonies. If they did not integrate the linked lacZ nonsense mutation, they would make blue colonies on these plates. Thus, the comigration frequency is defined as the percent of Tm^R colonies that were white on these plates.

Natural transformation frequencies for *H. pylori* were determined as follows. Total chromosomal DNA (600 ng) from a streptomycin resistant 26695 strain was mixed with 45 µL of a culture at OD$_{600}$ = 4.0. Then, three 15µL drops of the mix were spotted on solid medium and incubated

408 overnight at 37°C. Next day, cells were resuspended in 300 μL of peptone water and 10 μl drops
409 of serial dilutions were deposited on BAB plates with and without streptomycin (10 μg/mL).
410 Transformation frequency is defined as the CFU/mL of streptomycin resistant transformants
411 divided by the CFU/mL of total viable counts.
412
Microscopy for GFP-ComM focus formation
Assays to assess GFP-ComM focus formation in the presence of tDNA were performed exactly
as previously described[17]. Briefly, cells were grown as described above for natural
transformation assays. Then, ~10^7 cells were mixed with ~650 ng of a linear ~6 kb tDNA product
that is homologous to the host chromosome, and placed under an 0.2% gelzan pad. Samples
were then subjected to timelapse imaging. Phase contrast and fluorescence images were
collected every 3 min for a total of 90 mins on a Nikon Ti-2 microscope using a Plan Apo 60X
objective, a GFP filter cube, a Hamamatsu ORCA Flash 4.0 camera and Nikon NIS Elements
imaging software. The number of GFP-ComM foci was determined by analyzing 60 cells per
replicate.
423
Bacterial Adenylate Cyclase Two-Hybrid (BACTH) Assays
Gene inserts were amplified by PCR and cloned into the pUT18C, pUT18, pKT25, or pKNT25
vectors to generate N- and C- terminal fusions to the T18 and T25 fragments of adenylate
cyclase. Miniprepped vectors were then cotransformed into *E. coli* BTH101. Then, single
colonies were picked and grown in LB supplemented with kanamycin 50 μg/mL and carbenicillin
100 μg/mL. Next, 3 μL of each strain was spotted onto LB agar plates supplemented with
kanamycin 50 μg/mL, carbenicillin 100μg/mL, 0.5 mM IPTG and 40 μg/mL X-gal. Plates were
incubated at 30°C for 24 hours prior to imaging.
432
To quantify the strength of BACTH interactions, five colonies were collected and used to
inoculate 3 mL of LB containing both antibiotics and allowed to grow overnight at 30°C. The
OD_{600} of the culture was determined and cells from 1mL were pelleted and resuspended in 1mL
of buffer Z (60 mM Na_2HPO_4, 40 mM NaH_2PO_4, 10 mM KCl, 1 mM $MgSO_4$, 50 mM β-
mercaptoethanol, pH 7). Next, 100 μL of chloroform was added and reactions were vortexed for
20 s. Then, 20 μL of the upper phase was mixed in 96-well microplates with 180 μL of buffer Z
with 0.7 mg/mL of ortho-nitrophenyl-β-galactoside (ONPG) but without β-mercaptoethanol.
Absorbance measures at 540 and 414 nm were carried out every 5 minutes for 2 h. β-
galactosidase activity (A) was defined as: $A = 1000 * (OD_{414} (t) − (1.75 * OD_{540} (t)) / (t * V_{sample} *$
initial OD_{600}) and expressed as Miller units ($A414$/min mL OD_{600})
443
444
445 ***Statistics***
446 Statistical differences were assessed using GraphPad Prism software. The statical tests
447 employed are indicated in figure legends. Descriptive statistics for all samples, and a
448 comprehensive list of statistical comparisons can be found in **Dataset S2**.
449
450
ACKNOWLEDGEMENTS

We thank Carrie Shaffer and Prashant Damke for providing *H. pylori* genomic DNA to ABD. We
also thank Anne Marie Di Guilmi for setting up the initial BACTH experiments and Juliette
Bernardin for technical assistance in JPR's laboratory. This work was supported by grant
R35GM128674 from the National Institutes of Health to ABD and grant ANR-22-CE44-0044
from Agence Nationale de la Recherche to JPR, JA and RG. We thank the BIOI2 platform for
making the ColabFold pipeline easily accessible at the I2BC for YF, JA and RG.

REFERENCES

1   Dubnau, D. & Blokesch, M. Mechanisms of DNA Uptake by Naturally Competent
Bacteria. *Annu Rev Genet* **53**, 217-237 (2019).
2   Johnston, C., Martin, B., Fichant, G., Polard, P. & Claverys, J. P. Bacterial
transformation: distribution, shared mechanisms and divergent control. *Nat Rev*
Microbiol **12**, 181-196 (2014).
3   Fox, M. S. & Allen, M. K. On the Mechanism of Deoxyribonucleate Integration in
Pneumococcal Transformation. *Proc Natl Acad Sci U S A* **52**, 412-419 (1964).
4   Dubnau, D. & Davidoff-Abelson, R. Fate of transforming DNA following uptake by
competent Bacillus subtilis. I. Formation and properties of the donor-recipient complex. *J*
Mol Biol **56**, 209-221 (1971).
5   Mejean, V. & Claverys, J. P. Use of a cloned DNA fragment to analyze the fate of donor
DNA in transformation of Streptococcus pneumoniae. *J Bacteriol* **158**, 1175-1178 (1984).
6   Mortier-Barriere, I. *et al.* A key presynaptic role in transformation for a widespread
bacterial protein: DprA conveys incoming ssDNA to RecA. *Cell* **130**, 824-836 (2007).
7   Lisboa, J. *et al.* Molecular determinants of the DprA-RecA interaction for nucleation on
ssDNA. *Nucleic Acids Res* **42**, 7395-7408 (2014).
8   Quevillon-Cheruel, S. *et al.* Structure-function analysis of pneumococcal DprA protein
reveals that dimerization is crucial for loading RecA recombinase onto DNA during
transformation. *Proc Natl Acad Sci U S A* **109**, E2466-2475 (2012).
9   Yadav, T. *et al.* Bacillus subtilis DprA recruits RecA onto single-stranded DNA and
mediates annealing of complementary strands coated by SsbB and SsbA. *J Biol Chem*
288, 22437-22450 (2013).
10   Bakhlanova, I. *et al.* Mechanisms of RecA filament nucleation on ssDNA by the DprA
protein. *bioRxiv*, 2024.2005.2007.592916 (2024).
11   Nero, T. M. *et al.* ComM is a hexameric helicase that promotes branch migration during
natural transformation in diverse Gram-negative species. *Nucleic Acids Res* **46**, 6099-
6111 (2018).
12   Hardy, L. *et al.* YraN is a helicase-associated nuclease fostering extended recombination
events by natural transformation. *bioRxiv*, 2024.2002.2006.579203 (2024).
13   Lisboa, J. *et al.* The C-terminal domain of HpDprA is a DNA-binding winged helix domain
that does not bind double-stranded DNA. *FEBS J* **286**, 1941-1958 (2019).
14   Dwivedi, G. R. *et al.* Insights into the Functional Roles of N-Terminal and C-Terminal
Domains of Helicobacter pylori DprA. *PLoS One* **10**, e0131116 (2015).
15   Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493-500 (2024).
16   Karimova, G., Pidoux, J., Ullmann, A. & Ladant, D. A bacterial two-hybrid system based
on a reconstituted signal transduction pathway. *Proc Natl Acad Sci U S A* **95**, 5752-5756
(1998).
17   Dalia, A. B. & Dalia, T. N. Spatiotemporal Analysis of DNA Integration during Natural
Transformation Reveals a Mode of Nongenetic Inheritance in Bacteria. *Cell* **179**, 1499-
1511 e1410 (2019).
18   Gwinn, M. L., Ramanathan, R., Smith, H. O. & Tomb, J. F. A new transformation-deficient
mutant of Haemophilus influenzae Rd with normal DNA uptake. *J Bacteriol* **180**, 746-748
(1998).
19   Rosa, L. T., Vernhes, E., Soulet, A. L., Polard, P. & Fronzes, R. Structural insights into
the mechanism of DNA branch migration during homologous recombination in bacteria.
EMBO J (2024).

20   Mirouze, N. *et al.* Direct involvement of DprA, the transformation-dedicated RecA loader,
in the shut-off of pneumococcal competence. *Proc Natl Acad Sci U S A* **110**, E1035-1044
(2013).
21   Humphreys, I. R. *et al.* Protein interactions in human pathogens revealed through deep
learning. *Nat Microbiol* **9**, 2642-2652 (2024).
22   Gao, M., Nakajima An, D., Parks, J. M. & Skolnick, J. AF2Complex predicts direct
physical interactions in multimeric proteins with deep learning. *Nat Commun* **13**, 1744
(2022).
23   Miller, V. L., DiRita, V. J. & Mekalanos, J. J. Identification of toxS, a regulatory gene
whose product enhances toxR-mediated activation of the cholera toxin promoter. *J
Bacteriol* **171**, 1288-1293 (1989).
24   Tomb, J. F. *et al.* The complete genome sequence of the gastric pathogen Helicobacter
pylori. *Nature* **388**, 539-547 (1997).
25   Dalia, A. B. Natural Cotransformation and Multiplex Genome Editing by Natural
Transformation (MuGENT) of Vibrio cholerae. *Methods Mol Biol* **1839**, 53-64 (2018).
26   Dalia, A. B., McDonough, E. & Camilli, A. Multiplex genome editing by natural
transformation. *Proc Natl Acad Sci U S A* **111**, 8937-8942 (2014).
27   Dalia, T. N., Chlebek, J. L. & Dalia, A. B. A modular chromosomally integrated toolkit for
ectopic gene expression in Vibrio cholerae. *Sci Rep* **10**, 15398 (2020).
28   Li, M. Z. & Elledge, S. J. SLIC: a method for sequence- and ligation-independent
cloning. *Methods Mol Biol* **852**, 51-59 (2012).
29   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment
search tool. *J Mol Biol* **215**, 403-410 (1990).
30   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021).
31   Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**,
679-682 (2022).
32   Steinegger, M. & Soding, J. MMseqs2 enables sensitive protein sequence searching for
the analysis of massive data sets. *Nat Biotechnol* **35**, 1026-1028 (2017).
33   Paysan-Lafosse, T. *et al.* InterPro in 2022. *Nucleic Acids Res* **51**, D418-D427 (2023).
34   Steinegger, M. *et al.* HH-suite3 for fast remote homology detection and deep protein
annotation. *BMC Bioinformatics* **20**, 473 (2019).
35   Ellison, C. K. *et al.* Retraction of DNA-bound type IV competence pili initiates DNA
uptake during natural transformation in Vibrio cholerae. *Nat Microbiol* **3**, 773-780 (2018).
543

544
545
Fig. 1. **ComM-DprA interactions promote NT in *H. pylori*.** (**A**) AlphaFold 3 model of ComM$_{Hp}$-
DprA$_{Hp}$$^{C-term}$ interactions. The inset shows the DprA$_{Hp}$^{L252} interaction interface residue targeted
for mutagenesis in yellow. (**B**) BACTH analysis to assess the interaction between the indicated
alleles of DprA and ComM. All DprA constructs for BACTH consisted of C-terminal fusions of
T25 to the terminal 53 amino acids of the indicated DprA$_{Hp}$ allele. The ComM construct for
BACTH consisted of C-terminal fusions of T18 to full length ComM$_{Hp}$. The T25-zip and T18-zip
constructs represent a known interacting pair and are a positive control in this assay. Data are
representative of two independent experiments. (**C**) Natural transformation assays of the
indicated *H. pylori* strains using linear tDNA that integrates onto the genome. Data are from at
least 5 independent biological replicates and shown as the mean ± SD. Statistical comparisons
were made by one-way ANOVA with Tukey's multiple comparison test of the log-transformed
data. NS, not significant. *** = $p < 0.001$. Statistical identifiers directly above bars represent
comparisons to the *wt*.

559
560
Fig. 2. **ComM-DprA interactions promote recombination of tDNA during NT in *V. cholerae*.**
(**A**) AlphaFold 3 model of $ComM_{Vc}$-$DprA_{Vc}^{C-term}$ interactions. The inset shows the interaction
interface residues targeted for mutagenesis. The residues that form a putative electrostatic
interaction ($ComM_{Vc}^{R497}$ and $DprA_{Vc}^{E353}$) are in gray with charged atoms in blue (positive) or red
(negative), while the two leucine residues within the interface ($DprA_{Vc}^{L350}$ and $ComM_{Vc}^{L455}$) are
colored yellow. (**B**) BACTH analysis to assess the interaction between the indicated alleles of
DprA and ComM. All DprA constructs for BACTH consisted of N-terminal fusions of T18 to the
terminal 71 amino acids of the indicated DprA allele. All ComM constructs for BACTH consisted
of C-terminal fusions of T25 to full length ComM allele indicated. Data in **B** are representative of
two independent experiments. (**C**) Natural transformation assays of the indicated strains using
linear tDNA that integrates onto the genome. (**D**) Natural transformation assays of the indicated
strains using a replicating plasmid as the tDNA. Data in **C** and **D** are from at least 3 independent
biological replicates and shown as the mean ± SD. Statistical comparisons were made by one-
way ANOVA with Tukey's multiple comparison test of the log-transformed data. NS, not
significant. *** = $p < 0.001$. LOD, limit of detection. Statistical identifiers directly above bars
represent comparisons to the parent.

577
578
Fig. 3. **ComM$_{Vc}$-DprA$_{Vc}$ interactions are required for branch migration during NT.** (**A**)
Schematic of the experimental setup to assess branch migration via linkage analysis between a
trimethoprim resistance marker (Tm^R) and a nonsense mutation in *lacZ*. Two different tDNAs
were used: one where the *lacZ* nonsense mutation was 243 bp away from the Tm^R cassette,
and another where the spacing was 820 bp. (**B**) The two tDNA products described in **A** were
transformed into the indicated strains and the comigration frequency (i.e., the linkage frequency
between the Tm^R and *lacZ* nonsense mutation) is reported. All strains in these assays contained
a Δ*mutS* mutation to prevent mismatch repair from confounding the results. Data are from at
least 3 independent biological replicates and shown as the mean ± SD. Statistical comparisons
were made by one-way ANOVA with Tukey's multiple comparison test of the log-transformed
data for each tDNA product. NS, not significant. *** = $p < 0.001$, ** = $p < 0.01$. Statistical
identifiers directly above bars represent comparisons to the parent.

591
592
Fig. 4. **DprA$_{Vc}$ recruits ComM$_{Vc}$ to the recombination site during NT.** (**A**) Montage of
timelapse imaging showing that the recruitment of ComM$_{Vc}$ to tDNA can be easily observed as
fluorescent foci (white arrows) in a parent strain harboring a functional GFP-ComM$_{Vc}$ fusion at
the native locus. Scale bar, 4µm. (**B**) Quantification of timelapse images for (**B**) the percent of
cells that exhibited at least one GFP-ComM$_{Vc}$ focus and (**C**) the number of GFP-ComM$_{Vc}$ foci
observed per cell. The indicated strains were imaged every 3 min for 1.5 hours. Data are from 3
independent replicates and shown as the mean ± SD. n = 60 cells analyzed per replicate.

600
601
Fig. 5. **DprA-dependent recruitment of ComM is species-specific**. (**A**) Relative
transformation frequency of *V. cholerae* strains where native *comM* is intact (parent; gray bars)
or deleted (Δ*comM*; black bars) using linear tDNA that integrates onto the genome. Strains
express *H. pylori* ComM (ComM$_{Hp}$) and/or DprA (DprA$_{Hp}$) from an ectopic locus as indicated (+
denotes that the strain expresses the indicated protein). (**B**) Comigration frequency in the
indicated strain backgrounds to assess branch migration via linkage analysis between a
trimethoprim resistance marker (Tm^R) and a nonsense mutation in *lacZ* as depicted in **Fig. 2A**.
For the tDNA used is this experiment the *lacZ* nonsense mutation was 820 bp away from the
Tm^R cassette. All data are from at least 3 independent biological replicates and shown as the
mean ± SD. Statistical comparisons were made by one-way ANOVA with Tukey's multiple
comparison test of the log-transformed data. NS, not significant. *** = $p < 0.001$, ** = $p < 0.01$.

613
614
Fig. S1. **Most species with ComM encode a DprA homolog with a C-terminal winged-helix**
domain. Estimated maximum likelihood phylogeny of the indicated species based on a
concatenated alignment of 36 conserved proteins identified from whole genome sequences
(Genome Accession numbers are available in **Dataset S1**). Species that encode a ComM
homolog are highlighted in gray. The architecture of the DprA homolog (if present) is indicated to
the right of each species name. The core DprA smf domain is indicated by a black box, while the
presence of a C-terminal winged helix domain is demarcated by a green box (details of C-
terminal domain identification are presented in Methods and **Dataset S1**). Species that have
been shown to have the capacity for NT are designated by a black star. Major taxa are labeled
along their nodes. Pro: Proteobacteria (Greek letters denote subdivisions); Bac: Bacilli; Mol:
Mollicutes; Cya: Cyanobacteria; Arc: Archaea.

626
627
628 **Fig. S2. Quantification of BACTH for ComM$_{Hp}$-DprA$_{Hp}$ interactions.** The indicated
629 interactions were quantitatively assessed for interactions by performing Miller Assays. Statistical
630 identifiers directly above bars represent comparisons to the parent containing the empty vectors
631 (T18 / T25). NS, not significant. *** = $p < 0.001$.

632
633
Fig. S3. **ComM[R497E] is not a gain-of-function allele.** Branch migration was assessed via
linkage analysis between a trimethoprim resistance marker (Tm[R]) and a nonsense mutation in
lacZ as depicted in **Fig. 2A**. For the tDNA used is this experiment the *lacZ* nonsense mutation
was 820 bp away from the Tm[R] cassette. (**A**) Transformation frequency and (**B**) comigration
frequency are reported for strains with the indicated genotypes. Data are from 3 independent
biological replicates. Statistical comparisons were made by one-way ANOVA with Tukey's
multiple comparison test of the log-transformed data. NS, not significant. *** = $p < 0.001$, * = $p <$
0.05. Statistical identifiers directly above bars represent comparisons to the parent.

642
643
Fig. S4. DprA-dependent recruitment of ComM is species-specific. Transformation
frequency assays of the indicated *V. cholerae* strains where native *comM* is intact (parent; gray
bars) or deleted ($\Delta comM$; black bars). Strains also express $ComM_{Hp}$ and/or $DprA_{Hp}$ from an
ectopic locus as indicated (+ denotes that the strain expresses the indicated allele). These are
the raw transformation frequencies for the data presented in **Fig. 5A**.

Table S1 - *Strains used in this study*

Strain #*	Genotype	Figures	Use in manuscript
***Helicobacter pylori* strains**			
LR1	*wild-type*	Fig. 1C	wt for NT frequency assays
LR1187	*ΔcomM::Kan^R*	Fig. 1C	*ΔcomM* for NT frequency assays
LR1241	*ΔcomM::Kan^R, P$_{ureA}$::comM::Cm^R*	Fig. 1C	*ΔcomM; ureA::comM* for NT frequency assays
LR827	*ΔdprA::Cm^R*	Fig. 1C	*ΔdprA* for NT frequency assays
LR853	*ΔdprA::Kan^R, P$_{ureA}$::dprA::Apra^R*	Fig. 1C	*ΔdprA; ureA::dprA* for NT frequency assays
LR918	*ΔdprA::Kan^R, P$_{ureA}$::dprA^{C-ter}::Apra^R*	Fig. 1C	*ΔdprA; ureA::dprA^{C-ter}* for NT frequency assays
LR1373	*ΔdprA::kan^R, ΔcomM::Cm^R, P$_{ureA}$::dprA^{C-ter}::Apra^R*	Fig. 1C	*ΔcomM; ΔdprA; ureA::dprA^{C-ter}* for NT frequency assays
LR1255	*ΔdprA::Kan^R, P$_{ureA}$::dprA^{L252E}::Cm^R*	Fig. 1C	*ΔdprA; ureA::dprA^{L252E}* for NT frequency assays
***Vibrio cholerae* strains**			
TND0905 / SAD2468	*P$_{tac}$-tfoX, ΔluxO, lacZ::lacI^q, pilA^{S67C}, ΔVC1807::Zeo^R*	Fig. 2C, 2D	parent for NT frequency assays
TND2245 / SAD3567	*ΔcomM::Carb^R, P$_{tac}$-tfoX, ΔluxO, lacZ::lacI^q, pilA^{S67C}, ΔVC1807::Zeo^R*	Fig. 2C, 2D	*ΔcomM* for NT frequency assays
TND1072 / SAD3569	*ΔdprA::Cm^R, P$_{tac}$-tfoX, ΔluxO, lacZ::lacI^q, pilA^{S67C}, ΔVC1807::Zeo^R*	Fig. 2C, 2D	*ΔdprA* for NT frequency assays
TND4143 / SAD3589	*dprA^{L350E}, P$_{tac}$-tfoX, ΔluxO, lacZ::lacI^q, pilA^{S67C}, ΔVC1807::Cm^R*	Fig. 2C, 2D	*dprA^{L350E}* for NT frequency assays
TND4148 / SAD3590	*ΔcomM::Carb^R, dprA^{L350E}, P$_{tac}$-tfoX, ΔluxO, lacZ::lacI^q, pilA^{S67C}, ΔVC1807::Cm^R*	Fig. 2C	*dprA^{L350E} ΔcomM* for NT frequency assays
TND4145 / SAD3591	*dprA^{E353R}, P$_{tac}$-tfoX, ΔluxO, lacZ::lacI^q, pilA^{S67C}, ΔVC1807::Cm^R*	Fig. 2C, 2D	*dprA^{E353R}* for NT frequency assays
TND4150 / SAD3592	*ΔcomM::Carb^R, dprA^{E353R}, P$_{tac}$-tfoX, ΔluxO, lacZ::lacI^q, pilA^{S67C}, ΔVC1807::Cm^R*	Fig. 2C	*dprA^{E353R} ΔcomM* for NT frequency assays
TND4146 / SAD3593	*comM^{L455E}, P$_{tac}$-tfoX, ΔluxO, lacZ::lacI^q, pilA^{S67C}, ΔVC1807::Cm^R*	Fig. 2C, 2D	*comM^{L455E}* for NT frequency assays
TND4147 / SAD3594	*comM^{R497E}, P$_{tac}$-tfoX, ΔluxO, lacZ::lacI^q, pilA^{S67C}, ΔVC1807::Cm^R*	Fig. 2C, 2D	*comM^{R497E}* for NT frequency assays
TND4152 / SAD3595	*dprA^{E353R}, comM^{R497E}, P$_{tac}$-tfoX, ΔluxO, lacZ::lacI^q, pilA^{S67C}, ΔVC1807::Kan^R*	Fig. 2C	*dprA^{E353R}, comM^{R497E}* for NT frequency assays
SAD1063	*ΔmutS, Kan^R-P$_{tac}$-tfoX*	Fig. 3B, 5B, S3	parent for branch migration assays
SAD1071	*ΔcomM::Spec^R, ΔmutS, Kan^R-P$_{tac}$-tfoX*	Fig. 3B, 5B, S3	*ΔcomM* for branch migration assays
TND4153 / SAD3596	*dprA^{L350E}, ΔVC1807::Cm^R, ΔmutS, Kan^R-P$_{tac}$-tfoX*	Fig. 3B, S3	*dprA^{L350E}* for branch migration assays
TND4151 / SAD3597	*dprA^{E353R}, ΔVC1807::Cm^R, ΔmutS, Kan^R-P$_{tac}$-tfoX*	Fig. 3B	*dprA^{E353R}* for branch migration assays
TND4155 / SAD3598	*comM^{L455E}, ΔVC1807::Cm^R, ΔmutS, Kan^R-P$_{tac}$-tfoX*	Fig. 3B	*comM^{L455E}* for branch migration assays
TND4156 / SAD3599	*comM^{R497E}, ΔVC1807::Cm^R, ΔmutS, Kan^R-P$_{tac}$-tfoX*	Fig. 3B, S3	*comM^{R497E}* for branch migration assays
TND4169 / SAD3600	*dprA^{E353R}, comM^{R497E}, ΔVC1807::Cm^R, ΔmutS, Kan^R-P$_{tac}$-tfoX*	Fig. 3B	*dprA^{E353R} comM^{R497E}* for branch migration assays
SAD1924	P$_{tac}$-tfoX, ΔluxO::Spec^R, comEA-mCherry, ΔVC1807::Cm^R, pilA^{S67C}, 0.11 Mbp::parS$_{MT1}$, gfp-comM, ΔlacZ::P$_{lac}$-cyPet-parB$_{MT1}$ Zeo^R	Fig. 4A-C	parent for GFP-ComM focus formation assay
TND4178 / SAD3601	*dprA^{L350E}, P$_{tac}$-tfoX, ΔluxO::Spec^R, comEA-mCherry, ΔVC1807::Kan^R,*	Fig. 4A-C	*dprA^{L350E}* for GFP-ComM focus formation assay

	$pilA^{S67C}$, 0.11 Mbp::$parS_{MT1}$, gfp-$comM$, $\Delta lacZ$::P_{lac}-$cyPet$-$parB_{MT1}$ Zeo^R		
TND4177 / SAD3602	$comM^{L455E}$, P_{tac}-$tfoX$, $\Delta luxO$::$Spec^R$, $comEA$-$mCherry$, $\Delta VC1807$::Kan^R, $pilA^{S67C}$, 0.11 Mbp::$parS_{MT1}$, gfp-$comM$, $\Delta lacZ$::P_{lac}-$cyPet$-$parB_{MT1}$ Zeo^R	Fig. 4A-C	$comM^{L455E}$ for GFP-ComM focus formation assay
TND4441 / SAD3603	$igVCA0265$-$VCA0266$::Cm^R-P_{tac}-gfp, $\Delta VCA0692$-$Carb^R$-P_{tac}-gfp, P_{tac}-$tfoX$, $\Delta luxO$, $lacZ$::$lacI^q$, $pilA^{S67C}$, $\Delta VC1807$::Zeo^R	Fig. 5A, S4	parent for NT frequency assays
TND4450 / SAD3604	$\Delta comM$::$Spec^R$, $igVCA0265$-$VCA0266$::Cm^R-P_{tac}-gfp, $\Delta VCA0692$-$Carb^R$-P_{tac}-gfp, P_{tac}-$tfoX$, $\Delta luxO$, $lacZ$::$lacI^q$, $pilA^{S67C}$, $\Delta VC1807$::Zeo^R	Fig. 5A, S4	$\Delta comM$ for NT frequency assays
TND4406 / SAD3605	$igVCA0265$-$VCA0266$::Cm^R-P_{tac}-gfp, $\Delta VCA0692$-$Carb^R$-P_{tac}-$comM_{Hp}$, P_{tac}-$tfoX$, $\Delta luxO$, $lacZ$::$lacI^q$, $pilA^{S67C}$, $\Delta VC1807$::Zeo^R	Fig. 5A, S4	Ectopic $comM_{Hp}$ for NT frequency assays
TND4426 / SAD3606	$\Delta comM$::$Spec^R$, $igVCA0265$-$VCA0266$::Cm^R-P_{tac}-gfp, $\Delta VCA0692$-$Carb^R$-P_{tac}-$comM_{Hp}$, P_{tac}-$tfoX$, $\Delta luxO$, $lacZ$::$lacI^q$, $pilA^{S67C}$, $\Delta VC1807$::Zeo^R	Fig. 5A, S4	$\Delta comM$ native + ectopic $comM_{Hp}$ for NT frequency assays
TND4410 / SAD3607	$igVCA0265$-$VCA0266$::Cm^R-P_{tac}-$dprA_{Hp}$, $\Delta VCA0692$-$Carb^R$-P_{tac}-gfp, P_{tac}-$tfoX$, $\Delta luxO$, $lacZ$::$lacI^q$, $pilA^{S67C}$, $\Delta VC1807$::Zeo^R	Fig. 5A, S4	Ectopic $dprA_{Hp}$ for NT frequency assays
TND4427 / SAD3608	$\Delta comM$::$Spec^R$, $igVCA0265$-$VCA0266$::Cm^R-P_{tac}-$dprA_{Hp}$, $\Delta VCA0692$-$Carb^R$-P_{tac}-gfp, P_{tac}-$tfoX$, $\Delta luxO$, $lacZ$::$lacI^q$, $pilA^{S67C}$, $\Delta VC1807$::Zeo^R	Fig. 5A, S4	$\Delta comM$ native + ectopic $dprA_{Hp}$ for NT frequency assays
TND4353 / SAD3609	$igVCA0265$-$VCA0266$::Cm^R-P_{tac}-$dprA_{Hp}$, $\Delta VCA0692$-$Carb^R$-P_{tac}-$comM_{Hp}$, P_{tac}-$tfoX$, $\Delta luxO$, $lacZ$::$lacI^q$, $pilA^{S67C}$, $\Delta VC1807$::Zeo^R	Fig. 5A, S4	Ectopic $comM_{Hp}$ and $dprA_{Hp}$ for NT frequency assays
TND4357 / SAD3610	$\Delta comM$::$Spec^R$, $igVCA0265$-$VCA0266$::Cm^R-P_{tac}-$dprA_{Hp}$, $\Delta VCA0692$-$Carb^R$-P_{tac}- $comM_{Hp}$, P_{tac}-$tfoX$, $\Delta luxO$, $lacZ$::$lacI^q$, $pilA^{S67C}$, $\Delta VC1807$::Zeo^R	Fig. 5A, S4	$\Delta comM$ native + ectopic $comM_{Hp}$ and $dprA_{Hp}$ for NT frequency assays
TND4370 / SAD3611	$\Delta VCA0692$-$Carb^R$-P_{tac}-$comM_{Hp}$, $\Delta mutS$, Kan^R-P_{tac}-$tfoX$	Fig. 5B	Ectopic $comM_{Hp}$ for branch migration assays
TND4385 / SAD3612	$\Delta comM$::$Spec^R$, $\Delta VCA0692$-$Carb^R$-P_{tac}-$comM_{Hp}$, $\Delta mutS$, Kan^R-P_{tac}-$tfoX$	Fig. 5B	$\Delta comM$ native + ectopic $comM_{Hp}$ for branch migration assays
TND4388 / SAD3613	$igVCA0265$-$VCA0266$::Cm^R-P_{tac}-$dprA_{Hp}$, $\Delta VCA0692$-$Carb^R$-P_{tac}- $comM_{Hp}$, $\Delta mutS$, Kan^R-P_{tac}-$tfoX$	Fig. 5B	Ectopic $comM_{Hp}$ and $dprA_{Hp}$ for branch migration assays
TND4396 / SAD3614	$\Delta comM$::$Spec^R$, $igVCA0265$-$VCA0266$::Cm^R-P_{tac}-$dprA_{Hp}$, $\Delta VCA0692$-$Carb^R$-P_{tac}- $comM_{Hp}$, $\Delta mutS$, Kan^R-P_{tac}-$tfoX$	Fig. 5B	$\Delta comM$ native + ectopic $comM_{Hp}$ and $dprA_{Hp}$ for branch migration assays
TND4318 / SAD3615	$dprA^{L350E}$, $comM^{R497E}$, $\Delta VC1807$::Cm^R, $\Delta mutS$, Kan^R-P_{tac}-$tfoX$	Fig. S3	$dprA^{L350E}$ $comM^{R497E}$ for branch migration assays
***Escherichia coli* strains**			
PR1577	pKNT25-$dprA_{Hp}$	Fig. 1B	Harbors the DprA$_{Hp}$-T25 (only the C-terminal 53 residues) for BACTH analysis
PR1650	pKNT25-$dprA_{Hp}^{L350E}$	Fig. 1B	Harbors the DprA$_{Hp}^{L350E}$-T25 (only the C-terminal 53 residues) for BACTH analysis
PR1550	pUTN18-$comM_{Hp}$	Fig. 1B	Harbors the ComM$_{Hp}$-T18 vector for BACTH analysis

TND4189 / SAD3616	pUT18C-*dprA$_{Vc}$*	Fig. 2B	Harbors the T18-DprA$_{Vc}$ (only the C-terminal 71 residues) for BACTH analysis
TND4230 / SAD3617	pUT18C-*dprA$_{Vc}$^{L350E}*	Fig. 2B	Harbors the T18-DprA$_{Vc}$^{L350E} (only the C-terminal 71 residues) for BACTH analysis
TND4235 / SAD3618	pUT18C-*dprA$_{Vc}$^{E353R}*	Fig. 2B	Harbors the T18-DprA$_{Vc}$^{E353R} (only the C-terminal 71 residues) for BACTH analysis
TMN0427 / SAD3619	pKTN25-*comM$_{Vc}$*	Fig. 2B	Harbors the ComM$_{Vc}$-T25 vector for BACTH analysis
TND4231 / SAD3620	pKTN25-*comM$_{Vc}$^{L455E}*	Fig. 2B	Harbors the ComM$_{Vc}$^{L455E}-T25 vector for BACTH analysis
TND4232 / SAD3621	pKTN25-*comM$_{Vc}$^{R497E}*	Fig. 2B	Harbors the ComM$_{Vc}$^{R497E}-T25 vector for BACTH analysis
SAD2236	pKT25	Fig. 2B	Harbors the T25 empty vector for BACTH analysis
SAD2240	pUT18C	Fig. 2B	Harbors the T18 empty vector for BACTH analysis
SAD2241	pUT18C-zip	Fig. 2B	Harbors the T18-zip vector for BACTH analysis
SAD2238	pKT25-zip	Fig. 2B	Harbors the T25-zip vector for BACTH analysis

651
652 *Multiple Strain# designations indicate that the strain was stocked in two independent strain
653 collections.

Table S2 – *Primers used in this study*

Primer	Sequence (5' → 3')	Description
Helicobacter pylori **mutant constructs**		
Op329	gagaatattgtaggagatcttctagaaagataaggttaagatagagtatcaaaaccg	Amplify c*omM* locus F
Op330	ccggatggctcgagtttttcagcaagatggttcattctgcttgatgaaagc	Amplify c*omM* locus R
Op345	gtaacaaccatgattaacacgatgtttt<u>gatggctaaaatgagaatatcaccg</u>	Insertion *Kan^R* in *comM* locus F
Op346	cttataaaaatccttttaagaaatctttctaaaac<u>ctaaaacaattcatccagtaaaat</u>	Insertion *Kan^R* in *comM* locus R
Op478	gaaaaacactttaagaataggagaataaggaattcatgattaacacgatgttttgcg	Insertion *ComM* in *ureA* locus for ectopic expression F
Op479	gccatttattcctcctagttagtcacccg<u>ggtacc</u>ttaagaaatctttctaaaac	Insertion *ComM* in *ureA* locus for ectopic expression R
Op839	ggggataagcttttagaatacgaggagttgggtaagatcaagcgcatcaatc	*DprA^{L252E}* site-directed mutagenesis F
Op840	gattgatgcgcttgatcttacccaactcctcgtattctaaaagcttatcccc	*DprA^{L252E}* site-directed mutagenesis R
Op607	ggataacaatttcacacaggaaacagctatgtaccatttaaaagaaatgcctgaaatgg	Amplify C-terminal 53 residues of *dprA$_{Hp}$* to clone into BACTH vector F
Op608	ccatttcaggcatttcttttaaatggtacatagctgtttcctgtgtgaaattgttatcc	Amplify C-terminal 53 residues of *dprA$_{Hp}$* to clone into BACTH vector R
Op493	ggataacaatttcacacaggaaacagct<u>at</u>gatgattaacacgatgttttgcgcg	Amplify *comM$_{Hp}$* to clone into BACTH vector F
Op495	cgcgcaaaacatcgtgttaatcatcatagctgtttcctgtgtgaaattgttatcc	Amplify *comM$_{Hp}$* to clone into BACTH vector R
Vibrio cholerae **mutant constructs***		
ABD820	CGCTCTTATCTGCTTGGATAATGG	Δ*dprA$_{Vc}$* (*i.e.*, ΔVC0048) or *dprA* point mutants F1
ABD998	gtcgacggatccccggaatCATTAACTGGCATCATCAACC	Δ*dprA$_{Vc}$* (*i.e.*, ΔVC0048) R1
ABD999	gaagcagctccagcctacaTAGCTATGATGATGGATATTTTGATG	Δ*dprA$_{Vc}$* (*i.e.*, ΔVC0048) F2
ABD996	GGTTTGGTTATCCGGATCACAC	Δ*dprA$_{Vc}$* (*i.e.*, ΔVC0048) or *dprA$_{Vc}$* point mutants R2
ABD855	CATGAATCACTTTGGCATGAGG	Δ*comM$_{Vc}$* (*i.e.*, ΔVC0032) or *comM$_{Vc}$* point mutants F1
ABD856	gtcgacggatccccggaatCATTGCTTCCCTTAGTATTTGATC	Δ*comM$_{Vc}$* (*i.e.*, ΔVC0032) R2
ABD857	gaagcagctccagcctacaTAGTACTCTGACCTGCAGAGTTC	Δ*comM$_{Vc}$* (*i.e.*, ΔVC0032) F2
ABD858	AAATTCCAGAAAAACCACGTC	Δ*comM$_{Vc}$* (*i.e.*, ΔVC0032) or *comM$_{Vc}$* point mutants R2
DOG0886	GCTCAAGCTCttcGAGCTGCATCATGATATCTTGCACCGGTATCTG	*dprA$_{Vc}^{L350E}$* R1
DOG0887	GATGCAGCTCgaaGAGCTTGAGCTCCTTGGGCATGTGGTTGCAGTA	*dprA$_{Vc}^{L350E}$* F2
DOG0888	GCCCAAGGAGacgAAGCTCCAAGAGCTGCATCATGATATCTTGCAC	*dprA$_{Vc}^{E353R}$* R1
DOG0889	CTTGGAGCTTcgtCTCCTTGGGCATGTGGTTGCAGTACCTGGTGGC	*dprA$_{Vc}^{E353R}$* F2

DOG0891	TTGATAAACCttcGCGATGCAGCGCGCTTTCTAGAAATTCCGCATC	$comM_{Vc}^{L455E}$ R1
DOG0892	GCTGCATCGCgaaGGTTTATCAATTCGGGCTTACCATCGCATCATT	$comM_{Vc}^{L455E}$ F2
DOG0893	GCTGTTTGAGTAAttcATCCATGGCTCGATAGCCCAACGCCTCGGCCAA	$comM_{Vc}^{R497E}$ R1
DOG0894	TCGAGCCATGGATgaaTTACTCAAACAGCTCAGTGCGCAGAACGTCTAG	$comM_{Vc}^{R497E}$ F2
BBC4911	gcaggtggagcaggtggaTATCAAGCAACACTTTTTTCTGCTG	Amplify C-terminal 71 residues of $dprA_{Vc}$ to clone into BACTH vector F
TMN0211	tgtaggctggagctgcttcCTAGCCTCTCCCCTTTCTAATATAGC	Amplify C-terminal 71 residues of $dprA_{Vc}$ to clone into BACTH vector R
TMN0202	gcaggtggagcaggtggaGGACTTGCGATCATTCATAGC	Amplify $comM_{Vc}$ to clone into BACTH vector F
TMN0204	tccaccacttccacctgcGACGTTCTGCGCACTGAG	Amplify $comM_{Vc}$ to clone into BACTH vector R
BBC5044	caatttcacacaggatcccgggAGGAGGTCAACGAATGAAAAGCCACTTCC	Amplify $dprA_{Hp}$ to clone into ectopic expression construct F
BBC5045	tgtaggctggagctgcttcTCACGCTAACACCACAATGTG	Amplify $dprA_{Hp}$ to clone into ectopic expression construct R
BBC5047	caatttcacacaggatcccgggAGGAGGTACAACCATGATTAACACGATGTT	Amplify $comM_{Hp}$ to clone into ectopic expression construct F
BBC5048	tgtaggctggagctgcttcTTAAGAAATCTTTCTAAAACTCAGCGC	Amplify $comM_{Hp}$ to clone into ectopic expression construct R

657 *Lower case nucleotides on the 5' end demarcate overlap sequences for SOE PCR and/or
658 isothermal assembly with BACTH vectors. While internal lower case nucleotides demarcate
659 point mutations.

660