1 # RDAforest: identifying environmental drivers of polygenic adaptation.

2 Mikhail V .Matz[1*] and Kristina L. Black[1,2]
3
[1] Department of Integrative Biology, University of Texas at Austin
[2] Current affiliation: the National Center for Atmospheric Research, Boulder, CO, USA
*Corresponding author, matz@utexas.edu

Abstract

Identifying environmental gradients driving genetic adaptation is one of the major goals of ecological
genomics. We present RDAforest, a methodology that leverages the predominantly polygenic nature of
adaptation and harnesses the versatility of random forest regression to solve this problem. Instead of
computing individual SNP-environment associations, RDAforest seeks to explain the overall genetic
covariance structure based on multiple environmental predictors. By relying on random forest instead of
linear regression, this method can detect non-linear and non-monotonous dependencies as well as all
possible interactions between predictors. It also incorporates a novel procedure to select the best
predictor out of several correlated ones, and uses jackknifing to model uncertainty of genetic structure
determination. Lastly, our methodology incorporates delineation and plotting of "adaptive
neighborhoods" - areas on the landscape that are predicted to harbor differentially adapted individuals.
Such maps can be used as a guide for planning conservation and ecological restoration efforts. We
demonstrate the use of RDAforest in two simulated scenarios and one real dataset (North American gray
wolves).

Introduction

Landscape genomics seeks to plot genetic adaptation on a spatial map, by associating environmental
parameters with allele frequencies across the genome (Balkenhol et al. 2017). Landscape genomics
methodologies are still under active development, but thus far most of them relied upon linear
associations between allele frequencies at individual SNPs and environmental predictors (Forester et al.
2018; Caye et al. 2019; Duruz et al. 2019; Privé et al. 2020). The major problem with such an approach is
that most adaptive responses in nature are polygenic, based on small allele frequency changes across a
great number of loci (Barghi, Hermisson, and Schlötterer 2020), and may be undetectable at single-SNP
level. SNP-by-SNP analysis is unavoidable when one wants to identify SNPs that are putatively involved in
adaptation, but that is but one of the three major goals of landscape genomics. The other two goals are
to identify likely environmental drivers of adaptation, and to predict where differentially adapted
individuals are located on the landscape. Here we show that these two goals can be achieved without
computing individual SNP-by-environment correlations.

Our first idea is that polygenic responses to natural selection are reflected in the overall genetic covariance
structure (Berg and Coop 2014). In simple terms, this means that multiple SNPs driving polygenic

adaptation match in their allelic state more often among individuals adapted to the same conditions,
which makes such individuals more genetically similar to each other than to the rest of the population.
Imagining all individuals as a multivariate cloud of points defined by genetic distances between them, such
co-adaptation would look like a bulge or extension of the cloud, likely aligned with one of its principal
components. So, to find such signals, it is possible to analyze the principal components of the genetic
distance matrix - genetic principal components (gPCs) - which would simultaneously leverage the
polygenic nature of adaptation to boost sensitivity and dodge the need to rely upon individual SNP-
environment associations. In support of this idea, a study in maize recently demonstrated that individual
SNP-environment association information did not improve prediction of yield under different
environmental conditions, compared to just the population structure captured by the first five gPCs (Li et
al. 2024). Likewise, in poplar, the prediction of genetic mis-adaptation calculated from randomly chosen
SNPs correlated with common garden observations just as well or better than the mis-adaptation
predicted from putative SNPs under environmental selection (Fitzpatrick et al. 2021). (We must
acknowledge that by historical convention, principal components of a distance matrix are called "principal
coordinates", but since their mathematical meaning is exactly the same (Legendre and Legendre 2012)
we will call them genetic principal components, gPCs, throughout this paper).
57
In asking how the overall covariance structure of a multivariate dataset aligns with external variables, our
method is conceptually similar to the redundancy analysis (RDA, (Borcard, Gillet, and Legendre 2011)),
which has the same goal. However, the second idea of our method makes it different from the classical
RDA: associations with environmental predictors are detected by random forest regressions instead of
linear regression. The advantage is twofold. First, random forest can detect non-linear and non-
monotonous associations, for example, peak-like dependencies that may be entirely flat from the point
of view of linear regression. Second, random forest seamlessly accounts for all possible interactions
between predictors without the need to specify them and without loss of power. Of particular use to us
is the random forest implementation called gradient forest (Ellis, Smith, and Pitcher 2012), which extends
random forest regression to multiple response variables (in our case, multiple gPCs). It also introduces
two innovations that greatly help visualization and interpretation of the results. The first one is the
concept of turnover curves, which are graphs to visualize the magnitude of change in the response matrix
as one moves across the scale of a given predictor. The other innovation is scaling of predictor
importances, typically measured as increase in mean square error when the predictor is permuted, in
terms of proportion of variation explained by the predictor. For technical details, we refer the reader to
the excellent summaries of how RDA (Borcard, Gillet, and Legendre 2011), regression trees (De'Ath 2002),
and random forest (Ellis, Smith, and Pitcher 2012) algorithms work.
75
Existing methodologies have explored either the former or the latter idea, but did not put the two
together. There is a landscape genomics method that uses the overall genetic covariance structure to find
associations with the environment, the RDA approach by Forester et al (Capblancq and Forester 2021;
Forester et al. 2018). However, it still relies on linear regressions to detect associations. On the other
hand, gradient forest has been previously used to analyze landscape genomics data (Fitzpatrick and Keller
2015; Bay et al. 2018), but it was applied to individual SNP genotypes instead of gPCs, thus missing the

opportunity to capitalize on the overall covariance structure. Also, such an implementation is limited to
just a few hundred SNPs, because of limitations of the gradient forest algorithm.
84	
There are two more innovations in our methodology. The first one is the predictor selection procedure,
to identify the true driver of genetic structure out of many cross-correlated predictors. It is a common
problem with environmental predictors: for example, many standard bioclimatic variables that are
routinely used in landscape genomics (e.g., (Bay et al. 2018)) are correlated with each other. Our solution
is based on the observation that predictors that don't actually drive the response but are simply correlated
with the true driver diminish in importance at a higher *mtry* setting (Strobl et al. 2008). *mtry* is the number
of predictors randomly chosen for tree construction in a specific random forest replicate. Correlated
predictors appear important at lower *mtry* because they "stand in" for the true driver that was not chosen
in that replicate. At higher *mtry*, the true driver is more often present among the chosen predictors and
is then preferred for tree construction, which leads to diminishing importance of "stand-in" predictors.
Our procedure reruns the analysis at two *mtry* settings and highlights predictors that do not decrease in
importance at higher *mtry*.
97	
98	The second innovation is the jackknifing procedure when computing gPCs, which models uncertainty of
99	their determination due to finite sample size. In each jackknifing replicate we withhold some proportion
100	(20% by default) of all samples from the analysis, compute gPCs without them, and then project the
101	withheld samples onto the ordination to obtain their scores. This models how different the gPCs would
102	be if we had 20% fewer samples.
103	
104	**Methods**
105	
106	Inputs
107	
108	*Genetic distances*
109	
RDAforest operates on a square matrix of genetic distances between individuals. This can be computed
from a matrix of genotypes (*G*) in [0,1,2] format (i.e. listing the number of derived or non-reference alleles
at the given SNP in a given individual) by computing genotypic correlations using R function *cor()* and
converting them to distances by subtracting them from 1. For non-model species with Restriction site
Associated DNA (RAD) sequencing or shallow whole-genome sequencing (WGS) data, a robust way to
compute genetic distances is provided by ANGSD software (Korneliussen, Albrechtsen, and Nielsen 2014):
it is identity-by-state (IBS) calculation based single-read resampling. For each base in the genome that is
covered by at least one read in each of the two compared individuals, the procedure randomly samples
one read from each individual and records either match or mismatch. The proportion of observed matches
is the IBS, and the distance returned by ANGSD is 1-IBS (note that the IBS-distance of an individual against
itself or its clone is not 0 but half the individual's heterozygosity). The method does not require genotype
calling and is robust to variation in coverage depth between samples. Another promising method for
computing genetic distances from raw low-coverage read mapping data is PCAngsd (Meisner and

Albrechtsen 2018), which takes genotype likelihoods from ANGSD output and computes the genetic
covariance matrix while adjusting genotype likelihoods based on the emerging population structure. The
covariance matrix can then be then converted to distances based on genotypic correlations using
cov2cor() function in R.
127
128 *Environmental variables*
129
130 The core *RDAforest* analysis uses a table of environmental variables (*E*) with rows corresponding to each
131 individual in *G* (in the same order) and columns corresponding to environmental data. *E* could contain
132 hundreds of variables, some correlated with each other. While *RDAforest* is designed to deal with
133 correlated predictors to a degree, we recommend pruning the most strongly correlated ones, to leave
134 only one representative variable for a group that shows cross-correlations of $r = 0.9$ or higher.
135
136 *Spatial coordinates of genotyped individuals*
137
This is a table (*S*) with two columns giving spatial locations of sampled individuals (with rows in the same
order as *G*). It is needed to check whether the genetic structure follows the isolation-by-distance (IBD)
pattern and to remove that trend, if present. Typically the spatial coordinates are longitude and latitude.
To avoid distortions across the latitudinal range, we convert them to "great circle coordinates", which are
the first two principal components of the great circle distance matrix that takes the curvature of the Earth
into account. Great circle distances are used to investigate the presence of IBD, and great circle
coordinates are used to correct for IBD (if it is detected) during ordination construction.
145
146 *Prediction space*
147
148 This is a large table (*P*) containing latitude and longitude coordinates of points where adaptation is to be
149 predicted and their associated environmental values. To keep things simple, we recommend specifying a
150 grid of points on the map at a desired final resolution of the predictions, which would merge into a
151 continuous layer when plotted with a resolution-size symbol.
152
Workflow

153
154
155 *Exploring and countering isolation-by-distance trend*
156
Genetic structure across distances exceeding per-generation dispersal range tends to follow the isolation-
by-distance (IBD) pattern, with closely located individuals being more similar genetically (Wright 1943).
The easiest way to visually detect this trend is to plot genetic distances against geographic distances,
looking for a sloping cline, such as the one on Fig. 3B (or multiple clines, if there are additional drivers of
genetic structure besides IBD). To formally test the significance of association between genetic and
geographical distances one can use a Procrustes test (function *vegan::protest(), (Oksanen et al. 2007)*) or
the classical Mantel test. In a theoretically extreme case of IBD, when genetic distances perfectly

correspond with geographic distances, the genetic PCA would look like a geographical map of sampled
individuals (assuming per-generation dispersal does not vary across the study area, i.e. there are no
physical barriers to migration - please see Discussion for other situations). Interestingly, humans provide
an example approaching such a case: the European genetic PCA looks very much like the map of Europe
(Novembre et al. 2008). IBD is a nuisance trend for us because it is not driven by adaptation to specific
environmental gradients. Since strong IBD causes the genetic PCA to approach a two-dimensional map,
to collapse the IBD trend it is sufficient to linearly regress the two spatial coordinates ("great circle
coordinates") out of genetic data, i.e. construct a partial genetic ordination conditional on the spatial
coordinates. This approach is quite conservative since some meaningful environmental gradients may be
associated with longitude and especially latitude; regressing spatial coordinates out of the data would de-
power our ability to detect the influence of these gradients. Yet, we feel that this should be a default way
of dealing with IBD - but only if it is actually evident in the plot of genetic vs geographic distances and in
the Procrustes or Mantel test.
177
178   *Constructing genetic ordination*
179
We use the function *vegan::capscale()* (Oksanen et al. 2007) to build the ordination, which accommodates
the "Condition" argument to specify nuisance variables to be regressed out of the data (by linear
regression). These can be spatial "great circle coordinates" in case we need to account for IBD,
designations of physically isolated regions on the landscape (see Discussion), or technical covariates that
might affect the ordination, such as sequencing batch or read coverage. It is highly recommended to
explore the shape of the ordination for possible artifacts prior to running the RDAforest analysis, and
including all the nuisance variables that appear to have an effect into the "Condition" argument.
187
188   *Exploring the full* RDAforest *model*
189
The goal of this step is to evaluate how good a model we can build with all our predictors in *E*, which gPCs
are predictable, which variables in *E* are likely to be the most influential, and which gPCs they are most
associated with. At this stage we will deliberately use more leading gPCs that are likely informative (up to
all gPCs in the ordination). We do not yet perform any resampling to evaluate uncertainty, this will come
later. The function *makeGF()* takes an ordination (partial ordination if IBD was regressed out), table of
predictors, and runs *gradientForest()* analysis on it (Ellis, Smith, and Pitcher 2012). This function returns a
gradientForest model object that contains "importance" (predictive power) of each predictor scaled as
proportion of variation explained (R-squared) for each gPC. To compute proportions of total variation in
the whole dataset explained by each predictor, per-gPC R-squared values must be summed up while
weighing them by the proportion of total variation attributable to each gPC, which is what the function
importance_RDAforest() does. Summing these values over all predictors gives the cumulative proportion
of variation explained by the model. The *$result* slot of the *gradientForest* model contains the list of gPCs
that are predicted with a non-zero R-squared; only these gPCs will be used in subsequent stages. The
gradientForest model can also be used to plot turnover curves for each predictor and for each predictable
gPC, which is a good way to visualize which gPCs are responsive to which predictor(s) and where in the
predictor's range major gPC transitions happen.

Note that here as well as everywhere in the *RDAforest* pipeline the R-squared values are computed based
on cross-validation, using "out-of-bag" samples that were withheld from model building. This is different
from the classical linear regression, where the custom is to report goodness-of-fit measures based on the
same data that were used for model building. Such measures are always inflated compared to the true
predictive power of the model if it was applied to new data. Cross-validation R-squared values may seem
low to new users, but they are a more honest reflection of the model's predictive power.

Predictor selection based on mtry

gradientForest models are constructed using package *extendedForest*, which already handles correlated
predictors to a degree: it employs conditional permutation (Strobl et al. 2008) to avoid over-inflating the
importance of correlated predictors. Still, the predictive power remains somewhat spread over several
correlated predictors, even if only one of them is truly influential. To identify which one of the group of
correlated predictors is the most likely true driver (or most closely correlated with the unidentified true
driver), RDAforest uses the *mtry* criterion: at higher *mtry*, true predictors rise in importance, while non-
influential correlated predictors diminish in importance, compared to the lower *mtry* setting. This was
first seen in simulations that led to the development of the conditional permutation algorithm (Strobl et
al. 2008) and confirmed in simulations described below. Additional selection criterion is the relative
predictor's importance: we discard predictors whose importance is less than specified fraction (default
0.1) of the importance of the best predictor. Here, we use *gradientForest*-style importance (R-squared for
the predictor averaged over all predictable gPCs) rather than RDAforest-style importance introduced in
the previous paragraph (proportion of total variation predicted), because the latter would create bias
towards predictors associated with lower-number gPCs.

Ordination jackknifing

Predictor selection is where we use ordination jackknifing for the first time. The motivation for this
procedure is that with finite sample size the gPC axes are identified with some error, which might affect
the apparent importance of predictors. To model the uncertainty of gPCs determination, we run multiple
jackknifing replicates where we withhold a certain fraction of the data (default 20%) when constructing
gPC axes, thus modeling their uncertainty if we had 20% less samples. The hold-out samples are projected
back to the ordination before running the *gradientForest* analysis so we don't lose power due to smaller
sample size. The *mtry*-based selection procedure builds two *gradientForest* models on each replicate, with
higher and lower *mtry* setting, and records the importance of each predictor. These results are then
summarized as the proportion of replicates in which the importance or a given predictor increases at
higher *mtry*.

Measuring amount of variation attributable to most important predictors

After the most important predictors are identified, we use them in another series of ordination jackknifing
replicates with the function *ordinationJackknife()*, this time recording the importance (proportion of total

variation explained) for each passing predictor in each replicate. Boxplot of these values is the end-result
of the analysis of the influence of the environment on genetic structure.
Predicting adaptation at unsampled locations
251
252 We form two types of predictions for the new environmental dataset (*P*): gPC scores predicted by
253 *randomForest()*, and turnover curve values predicted by *gradientForest()*. These predictions are generated
254 by *ordinationJackknife()*, averaging across jackknifing replicates. We recommend running a relatively high
255 number (50 or more) replicates to ensure that predictions are robust across the landscape.
256
257 *Plotting adaptive neighborhoods*
258
Earlier papers (Fitzpatrick and Keller 2015; Fitzpatrick et al. 2021; Bay et al. 2018) followed the vignette
for the *gradientForest* package (Ellis, Smith, and Pitcher 2012) and plotted turnover curve values as
indicators of differential adaptation (or different community composition) on the landscape. However, as
we show in simulations below, using turnover curves for this purpose would imply that every transition
across a given predictor's range leads to a new (progressively diverging) state and happens regardless of
values of other predictors. Thus, it is appropriate only when changes across the range of each predictor
are monotonous, and when there is no interaction between predictors. In contrast, gPC scores predicted
by *randomForest()* are much more appropriate to plot since they retain both non-monotony and
interactions; however, they may be noisy across space since every point on the landscape is predicted on
its own. One way to improve this is by clustering landscape points based on turnover curves, which do
integrate over multiple data points to infer boundaries where major transitions happen. Such clusters
would robustly group landscape points containing similar adaptation, with a caveat that adaptation within
a given cluster can be the same as in some other clusters. To solve this problem, we generate clusters
based on turnover curves and then merge clusters that contain similar *randomForest()* predictions.
273

274 Results

275
276 *"Thresholds" simulation*
277
In this "toy" simulation (Fig. 1) the distribution of entities (species or alleles) on the landscape follows
simple thresholding rules. This is deliberately unrealistic; the primary goal of this simulation is to illustrate
how core algorithms of the *RDAforest* work. The environment is a 50 x 50 square grid with two important
predictors, *a* and *b*, forming orthogonal gradients. There are additional four predictors correlated with *a*
and two predictors correlated with *b*, plus three random predictors (Fig. 1 A, B). The landscape is
populated by five "species" that occupy certain regions defined by *a*, *b*, or their combinations (Fig. 1 C).
Note that two of these "species" ("a:b" and "b") show non-monotonous (peak-like) dependence on the
predictors and would not show any linear association with them. The distribution of the "species" on the
landscape can be summarized as the PCA map (Fig. 1 D), which shows scores of the first three PCs of the
"species" community on the landscape as color mixtures. The higher the color contrast, the more distinct

the communities are. To the best of our knowledge, this visualization was introduced in the vignette to
the *gradientForest* package, (Ellis, Smith, and Pitcher 2012).
290
The input for *RDAforest* analysis is a random selection of 10% of the landscape points (Fig. 1 E). The
gradientForest model indicated that predictors *a* and *b* are the most important (in terms of proportion of
total variance explained), with *b* less so than *a* (Fig. 1 F). This difference in importance makes sense since
there are four transitions along the range of *a* versus three transitions along the range of *b*; moreover, all
four regions along *a* are highly distinct while the first and third regions along *b* are similar (compare colors
on Fig. 1 D). The *mtry*-based selection procedure robustly identified *a* and *b* as the only viable predictors
(Fig. 1 G). Turnover curves for them (Fig. 1 H) reflect the number of transitions ("steps") along each
predictor's range, which values of the predictor correspond to transition boundaries (*x*-axis coordinates
of "steps"), amount of variation attributable to each transition (height of each "step"), and the proportion
of total variance explained (the maximum *y*-axis value reached).
301
The next stage is to use the model based on *a* and *b* to predict (in this case, recreate) the distribution of
distinct communities across the remaining 90% of the landscape. Two types of predictions are formed.
randomForest (RF) predictions (Fig. 1 I) are a direct attempt to predict gPC scores based on the model, for
each point individually. Note that RF does a decent job (compare RF predictions on panel I to the true
community picture on panel D) but is somewhat uncertain about the exact gPC values, especially at the
boundaries of "species" ranges. Clustering these predictions (Fig. 1 K) improves the situation but some
mis-predictions along the boundaries remain. In contrast, predictions based on turnover curves made by
gradientForest (Fig. 1 L) reconstruct the boundaries perfectly, but in many cases the distinctions between
bounded regions are incorrect (compare color contrasts on panel L to panel D). There are two problems
here. The first one is that turnover-based predictions make it look like the difference always increases
with each new transition, which is not necessarily the case: for example, the second transition along the
range of *b* leads to almost the same community as the initial state (Fig. 1 D), but in turnover-based
predictions it looks like the difference keeps accumulating (Fig. 1 L). The second problem is that turnover
predictions assume that each transition for a given predictor happens across the whole landscape,
irrespective of other predictors' values (i.e., predictors don't interact). For example, in reality the fourth
transition along predictor *a* does not happen unless *b* is higher than 15 (Fig. 1 D), but turnover-based
predictions miss that. Still, the fact that boundaries between *potentially distinct* communities are
reconstructed quite precisely (unlike RF predictions) makes turnover-based predictions useful: we first
cluster the turnover-based predictions to identify all the bounded regions, and then merge them based
on RF-based predictions contained within the regions. This two-step procedure leads to the perfect
recreation of the true community picture (Fig. 1 M).
323
324

Figure 1. "Thresholds" simulation: a toy example to illustrate how algorithms included in the RDAforest work. A: Environment is a 50 x 50 square with truly influential predictors *a* and *b*. There are four additional predictors (*c*, *e*, *f*, *g*) correlated with *a* and two more (*d*, *h*) correlated with *b*, plus three random predictors (*i-k*). Spatial coordinates are *x* and *y*. B: Hierarchical clustering tree of predictors based on their correlation across the landscape. C: Occurrences of five "species" across the landscape (black - occupied, gray - unoccupied), which in this simulation follows simple threshold rules. Each species occupies regions defined by predictors *a* and *b*. D: PCA map of "communities". Colors reflect the scores along the first three PCs of the community composition, based on species presences-absences. E: Samples (10% of total) for model building. F: Importances of predictors inferred by a simple *gradientForest* model (no jackknifing). G: Result of *mtry*-based selection procedure, based on 50 jackknifing replicates. Asterisks mark the passing predictors. H: Turnover curves for the passing predictors; steps are the boundaries where the community transitions to a different state. I: Predictions of gPCs of the community based on randomForest (RF), for each point separately. K: Clustering of the RF predictions. L: Clustered predictions of turnover curve values across the landscape. Note that while the boundaries between *potentially distinct* communities are reconstructed quite precisely, the values within each such partition do not reflect their actual differences (compare the color distribution to panel D). M: Clustering of turnover curve predictions followed by merging of the clusters based on RF predictions that they contain. N: *randomForest* predictions of the community based exclusively on spatial coordinates (*x* and *y*, panel A), using twice denser sampling than shown on panel E.

Last but not least, this simulation is useful to illustrate that one should not attempt to remove variation
that can be explained by spatial coordinates alone (*x* and *y*, Fig. 1 A). The random forest algorithm is
versatile enough to reconstruct *any* configuration of patches based on just the two linear coordinates (Fig.
1 N), provided dense enough sampling (we have used 20% of all points to generate Fig. 1 N). Clearly,
removing this signal would strongly (and perhaps even completely) de-power the main analysis since it
also aims to detect *the same patches* on the landscape.
350
351   *"Frequencies" simulation*
352
This simulation is more realistic in that the probability of finding certain species (or allele) depending on
the predictor changes smoothly across the predictor's range, instead of simply flipping between 0 and 1
like in the "thresholds" simulation. The environment (Fig. 2 A, B) remains the same as in the "thresholds"
simulation, but the "species" (Fig. 2 C) communities now show smooth transitions instead of sharp
boundaries. There is also one "species" that is found everywhere, to avoid having points with no species
present. Note that the spatial distribution of true PCA values of the community (Fig. 2 D) is very noisy,
reflecting the graininess of the simulation and the stochasticity of species occurrences. Our goal will be to
draw the most reasonable community boundaries despite this noise.
361
Using 10% of all points (Fig. 2 E) to build an *RDAforest* model, we first see that, once again, *a* and *b* are
identified as the most important predictors (Fig. 2 F). However, the proportion of total variance explained
by them is notably lower than in the "thresholds" simulation, because of the stochasticity in species
occurrences (Fig. 2 C). The *mtry*-based predictor selection procedure identifies *a* and *b* as the only two
viable predictors (Fig. 2 G); note that several additional predictors (marked by "x" symbols on Fig. 2 G)
pass the *mtry* criterion but are discarded due to low importance relative to the best predictor. Turnover
curves (Fig. 2 H) have smoothened steps reflecting the fact that transitions between communities are
gradual, not abrupt like in "thresholds" simulation. The RF predictions (Fig. 2 I, K) recapitulate the
noisiness of the original community (Fig. 2 D). With clustering the communities are sometimes reasonably
well separated (such as blue and purple in the lower right side of the landscape), but more typically the
boundaries between them remain undefined (Fig. 2 K). Turnover-based predictions show better promise
in terms of drawing community boundaries (Fig. 2 L) although it is often incorrect in predicting differences
between communities, just like in the "thresholds" simulation. Clustering based on turnover predictions
followed by merging the clusters based on RF predictions generates a map that demarcates regions
containing different communities reasonably well (Fig. 2 M). And once again, *randomForest* based on just
the spatial coordinates *x* and *y* (Fig. 2 A) is quite good at predicting the whole *a* and *b*-driven community
patchiness (Fig. 2 N), so removing all detectable spatial signal from the data prior to the analysis would
not be a good idea.
380
381

Simulation 2: frequencies

Figure 2. "Frequencies" simulation: a more realistic example. A and B: Environment is the same as in "thresholds" simulation. C: Occurrences of "species" across the landscape (black - occupied, gray - unoccupied). The probability of finding a species changes smoothly across the predictor range. D: PCA map of "communities". Colors reflect the scores along the first three PCs of the community composition. The noisiness reflects the stochasticity of species' occurrence. E: Samples (10% of total) for model building. F: Importances of predictors inferred by a simple *gradientForest* model (no jackknifing). G: Result of *mtry*-based selection procedure, based on 50 jackknifing replicates. Asterisks mark the passing predictors, "x" marks predictors that pass the *mtry* criterion but fail the importance criterion (they explain less than 0.1 of the variation explained by the best predictor). H: Turnover curves for passing predictors. Note that steps here are smoother than in "thresholds" simulation, because transitions between communities are gradual. I: Predictions of gPCs of the community based on randomForest (RF), for each point separately. K: Clustering of RF predictions. L: Clustered predictions of turnover curve values across the landscape. Note that the boundaries between communities are inferred quite reasonably, despite the noise in actual data (panel D). M: Clustering of turnover curve predictions followed by merging of the clusters based on RF predictions that they contain. N: *randomForest* predictions of the community based exclusively on spatial coordinates (*x* and *y*, panel A), using twice denser sampling than shown on panel E.

Real data analysis (North American wolves)

This dataset comes from (Schweizer et al. 2016) and has been used multiple times to illustrate the power
of landscape genomics methods (Capblancq and Forester 2021; Smith et al. 2024). For *G*, there are 94
wolves sampled across Canada and Alaska genotyped for 29,437 SNPs. For environmental predictors for
each wolf (*E*), we have a standard set of bioclimatic variables from
https://www.worldclim.org/data/bioclim.html, plus tree cover data for the approximate year or sampling
(2010) from   http://www.cec.org/north-american-environmental-atlas/land-cover-2010-modis-250m/.
Each wolf has been assigned an "ecotype" in the original paper (Schweizer et al. 2016); we expect that
our analysis will recover those ecotypes as differentially adapted groups. For *P* (prediction space) we have
a raster of points covering the whole Canada and USA at one-third of a latitudinal degree resolution
(approximately 37 km) and associated with the same variables as in *E*. These data and the Rmarkdown
script for the analysis described below are available at https://github.com/z0on/RDA-forest, the
Rmarkdown output can be viewed at https://rpubs.com/cmonstr/1224249.

415
<u>Figure 3</u>. Identifying factors driving genetic adaptation in North American wolves. A: raw PCA of genetic structure,
based on genotype correlations. B: Visual examination of the isolation-by-distance (IBD) trend: genetic distances
clearly increase with geographical (great circle) distances. C: Partial genetic PCA conditional on the "great circle
coordinates", to correct for IBD. Scores from this PCA become the input for further analysis. D: Correlations between
bioclimatic variables and tree cover used as potential predictors. E: Predictor selection based on *mtry*, with
somewhat relaxed passing criteria - requiring the predictor to show increased importance at higher *mtry* in only 0.25
of all replicates; red line. F: Boxplots of proportion of variation explained by the passing predictors, based on 50
ordinationJaccknife replicates.

The first step is exploration of genetic structure and isolation-by-distance (IBD). Wolves appear to be
highly genetically distinct across the landscape, apparently with respect to the assigned ecotype (Fig. 3 A).
Plotting genetic distances against geographic (great circle) distances revealed a clear sloping cline (Fig. 3

B), signifying prominent IBD; which was also supported by the Procrustes test of association between
scores of genetic ordination (Fig. 3 A) and geographic coordinates (function *vegan::protest*, p < 0.001). To
remove the IBD trend from the data, we formed a partial ordination conditional on the great circle
coordinates, which are the first two principal coordinates of the great circle distance matrix (Fig. 3 C). The
ecotype distinction largely remained, although Atlantic forest wolves now appear quite similar to West
forest and Boreal forest wolves. The gPCs from this partial ordination are used in the subsequent steps.
434

435   The next preparatory stage involves exploration and pruning (if necessary) of predictor variables. We
436   plotted a hierarchical clustering tree of our predictors, based on their correlation, and observed that many
437   bioclimatic variables were strongly correlated with each other. For this analysis, we have removed all but
438   one representative variable out of groups forming clades below 0.1 in the hierarchical clustering tree (i.e.,
439   groups of variables correlated with each other with *r* > 0.9, Fig. 3 D).
440

441
<u>Figure 4.</u> Plotting predicted patterns of differential adaptation for North American wolves. A: PCA of *randomForest*
predictions across the landscape, each point being a location within a 37-km-resolution raster covering the region
of interest. The four most-important predictors are projected onto the PCA using the function *vegan::envfit*. B:
Legend to ecotypes on subsequent panels. C: Unclustered randomForest (RF) predictions, overlaid with the 94 points
corresponding to the wolves used for model building (colors identify ecotypes according to panel B). Similarity of
colors on the underlying map implies similarity of inferred adaptation. D: Inferring boundaries between adaptive
neighborhoods by clustering RF predictions. E: Inferring boundaries between adaptive neighborhoods by clustering
turnover curve predictions followed by merging the resulting clusters based on RF predictions.
450

Next we ran the *mtry*-based variable selection procedure, relaxing the default passing criteria a bit: we
required that a predictor should increase in importance in more that a quarter of all replicates (the default
is more than one-third). We relaxed the criteria because our top priority in the end is to obtain the most

accurate predictions of adaptation across the landscape. If our main goal was to identify the most
important predictors and measure their importance, we would have used more stringent passing criteria
(see Discussion). Only three variables did not pass our lenient criteria (Fig. 3 E).

457
458   Next we ran 50 *ordinationJackknife* replicates to measure the importance of passing predictors (Fig. 3 F)
459   and to form robust adaptation predictions (averaged across jackknifing replicates) across the landscape.
460   Just like in the simulations above, we formed two types of predictions: direct gPC scores predicted by
461   *randomForest* ("RF predictions") and turnover-based predictions based on *gradientForest*.

462
The next step is plotting adaptive neighborhoods and exploring which environmental variables they
correspond to (Fig. 4). We begin with exploring unclustered gPC scores predicted by *randomForest* ("RF
predictions"). First we plot the PCA of these scores, with points colored according to the values of three
PCs (Ellis, Smith, and Pitcher 2012), and project the four most-important environmental predictors onto
it (Fig. 3 A). We see that blue and green points (landscape locations) are predicted to host wolves adapted
to colder and drier conditions compared to warm-colored points, and that wolves at blue points are likely
adapted to the high seasonality. The reddest points are associated with the wettest conditions during the
dry quarter. We then plotted these predicted points on the landscape (they merge into a continuous layer)
and overlayed the points showing the original 94 wolves, colored according to their assigned ecotype (Fig.
3 B, C). We see that distinct colors of the RF predictions tend to align with different ecotypes, with the
exception of West forest and Atlantic forest ecotypes that (as our model infers) appear to be adapted to
the same conditions.

475
We next explore the two different options for delineating the boundaries between adaptive
neighborhoods. The first one is to cluster the RF predictions directly, which corresponds to clustering of
simulation results shown on panels I and K of Fig. 1 and 2. We form 12 clusters and then merge them to
get seven or eight. The results look reasonable, cleanly delimiting all the ecotypes with the exception of
the High Arctic one, which remains spread across two closely related clusters (Fig. 4 D). Lastly, we do
clustering based on turnover curves followed by merging based on RF predictions, like on panels I, L, and
M of Fig 1 and 2. Visually, this clustering solution is almost indistinguishable from direct clustering based
on RF predictions (Fig. 4 E). Notably, the High Arctic ecotype remains problematic, only this time we see
that half of its representatives are attributed to the Arctic cluster.

485
486   **Discussion**

487
488   *Space: the perpetual annoyance*

489
The question how to deal with signals of genetic structure that is exclusively due to spatial separation
continues to be a hotly debated issue in the field of genotype-environment association analysis. One such
signal is isolation-by-distance (IBD), which is easy to detect by plotting genetic distances against
geographic distances (Fig. 3 B). It is also easy to deal with by linear regression since IBD drives one-to-one
correspondence between the spatial map and genetic structure as seen in PCA (Novembre et al. 2008).

This is a valid solution when per-generation dispersal distance does not vary across the land/seascape.
But what if it does? The problem is, there is no general way to tell whether such variation is due to physical
barriers to migration (which are of no interest to us) or selection against maladapted immigrants (which
is what we are looking for): both would generate similar distortions to the genetic structure in PCA space.
This means that one should not seek to remove just any signal of spatial patchiness from genetic data.
Although it is a relatively common practice to regress any signal of spatial structure that can be captured
by Moran eigenvector maps (MEMs) out, we believe that this amounts to splashing the baby out with the
bathwater. As we have shown in our simulations (Fig. 1 N and Fig. 2 N), with random forest regression and
enough sampling any spatial genetic structure (including instances of true local adaptation) can be
captured based on just spatial coordinates. So, if we remove all the detectable spatial signals from the
genetic data, we will lose the true adaptation signal as well. The only "space-correction" beyond
correction for IBD that we believe is prudent would be for cases when the dataset involves clearly defined
physically separated regions, such as different aquifers, mountain tops, or islands, between which
migration is likely restricted. These regions should be coded in metadata as dummy quantitative variables
(using a column of ones and zeros for each region) and regressed out of the data at the stage of ordination
construction (i.e. become additional constraints for the ordination).
511
512 *Dealing with collinear predictors*
513
Collinear (correlated) environmental variables are another perpetual problem of ecology. Some prior
landscape genomics papers took an ordination-based approach to recode the matrix of correlated
environmental variables into uncorrelated environmental PC scores (Bay et al. 2018). This is definitely a
valid approach with just one downside: one loses track of the actual variables and so the results become
harder to interpret. But should we really strive to minimize correlations among predictors? In classical
linear regression the answer is definitely yes, because the regression coefficients for correlated predictors
can swing wildly depending on the data and on which predictors are included in the model. In our method
this problem is greatly mitigated via extensive resampling of both data and predictors (the staple of the
random forest regression) and correlations-aware conditional permutation (Strobl et al. 2008; Ellis, Smith,
and Pitcher 2012). Thanks to these features, in our method the inferred predictor importances are stable
and the most important predictors are identified correctly, but still their importance remains partially
"shared" with correlated predictors. For example, consider the situation when there are two correlated
predictors and only one of them is truly influential. This predictor would explain 10% of variation on its
own, but including both predictors into the model would spread these 10% among the two depending on
the strength of their correlation (maybe 7% for the true predictor and 3% for the correlated one). Whether
or not this is a problem depends on the goal of the analysis. If the goal is to identify the drivers of genetic
variation across the landscape and measure their importance, then yes, one should try to remove all the
correlated but likely non-influential predictors. Conversely, if the goal is to build the most accurate model
to predict adaptation across the landscape, retaining correlated predictors will not hurt and may actually
help improve the model's accuracy. The default thresholds in the *mtrySelJack()* function for keeping a
predictor are 0.333 for the fraction of replicates in which its importance must increase at higher *mtry*, and
0.1 for the minimum importance relative to the best predictor. More conservative *mtry* selection
thresholds (higher importance in 50% or even 75% of high-*mtry* replicates and 0.2 for the minimum

allowed importance compared to the best predictor) would be preferred for identifying the most
important predictors and measuring the amount of variation they explain. The alternative "lenient mode"
for the most accurate model building would be to discard only the most hopeless predictors, those that
never increase in importance at higher *mtry* and explain less than 0.01 fraction of variation compared to
the best predictor. The possible downside is that the "guided clustering" of neighborhoods based on
turnover curves will likely stop working efficiently with too many predictors: a high number of intersecting
turnover curves will be splitting the landscape into too many small areas of potential differential
adaptation, countering the benefit of integrative nature of each turnover curve. Yet, direct gPC predictions
based on *randomForest* are expected to improve, so the lenient mode is definitely worth considering
when delineating adaptive neighborhoods. It is advisable to try several levels of predictor selection
stringency to see whether relaxing the selection criteria (and retaining more predictors) actually helps
improve the overall model accuracy.
549
550 *When the true predictor is missing*
551
It is quite possible that the actual driver of adaptation is missing from the variables in *E*, although there
might be one or more predictors correlated with it. Since correlated predictors could "stand-in" for the
true one, the RDAforest procedure will work, although its predictive power will be limited by the tightness
of correlation between true driver and existing predictors. Importantly, there is no formal way to tell
whether the most important predictor is indeed the true driver of adaptation or is only correlated with
the unknown true driver. Because of this, one should never claim that the true driver has been identified,
unless it is validated experimentally in a common garden experiment.
559
560 It is also possible to be missing the true driver and not having any predictors correlated with it. In that
561 case, some clearly visible separations of individuals in PCA space would remain unexplained. We seem to
562 have such a situation in the case of Arctic and High Arctic wolves: although they are clearly distinct in the
563 PCA space (Fig. 3 C), our model struggles to predict this (Fig. 4 C-E).
564
565 *Clustering and neighborhood boundaries*
566
Clustering works great to draw neighborhood boundaries in simulations (Fig. 1 and Fig 2), but there is a
downside: clustering involves three arbitrary decisions: which clustering method to use (RF-only or
turnover+RF), how many clusters to make initially, and at which similarity level to merge them. The
function *plot_adaptation()* that does the clustering/merging returns a hierarchical clustering tree with the
merging threshold indicated and plots a simple map with resulting clusters numbered on the landscape,
both of which can be used as a guide while exploring alternative clustering settings (Fig. 5). However, with
no prior data such as ecotype designations that we have for wolves, there is hardly any reliable guidance
for choosing clustering settings. We therefore recommend that the practitioners use unclustered maps
(such as on Fig. 4 C) as much as possible and avoid making decisions based on the boundaries produced
by clustering, unless there is a way to cross-validate the clusters.
577
578

579
580 Figure 5. Example clustering-guiding output of the function *plot_adaptation()*. A: Hierarchical clustering tree of RF
581 predictions for cluster medoids. The y-axis indicates how disparate the clusters are; it is scaled in terms of the fraction
582 of maximum observed difference between clusters. The red line is the merging threshold used. Numbers correspond
583 to clusters; merged clusters are assigned the same number. B: Merged clusters on the map. Numbers correspond to
584 clusters on panel A.

585
586 *Practical relevance*

587
The most important practical application of our methodology that we see is in conservation and ecological
restoration. Identifying the most important environmental drivers of adaptation essentially answers the
question "Which environmental changes pose the greatest challenge to our focal species", which is
directly useful for modeling species distributions under climate change. One way to approach this is to
calculate the "genetic offset" between current genetic makeup of populations and future conditions,
which can serve as a predictor of their future maladaptation and decline (Fitzpatrick and Keller 2015). The
genetic offset is simply the euclidean distance between corresponding points in the present-day
adaptation map (such as on Fig. 4 C) and the same kind of map generated for future values of the same
predictors. We must emphasize that one must use direct *randomForest* predictions rather than turnover-
based predictions for this calculation, because in a general case involving non-monotony and/or
interactions between predictors, turnover-based predictions do not accurately reconstruct the
distribution of differentially-adapted genotypes on the landscape (Fig. 1 D, L). It is also possible to plan
assisted migration interventions based on maps of adaptive neighborhoods (Fig. 4), such as importing
individuals from a like-adapted population for a genetic rescue of a locally-extirpated one. For example,
according to our model, if Atlantic Forest wolves were in decline, importing West Forest wolves could be
a solution: even though these populations are highly genetically distinct (Fig. 3 A), they appear to adapt
to the same environmental conditions (Fig. 4 D, E). It is also possible to identify areas on landscape that
currently don't host the focal species but are environmentally suitable for it (or becoming suitable due to
climate change), as potential sites for assisted colonization. For example, our model suggests that the
British Columbia ecotype might do well at the easternmost locations on our map (southern shore of
Newfoundland, Nova Scotia, and Prince Edward Island, Fig. 4 D).

609
610
611

612   *Need for validation*
613
Unfortunately, the results of ecological genomic analyses are rarely validated experimentally, leaving
open the question of how reliable are the rankings of putative adaptation drivers and how much can we
trust the maps of differential adaptation (such as the one on Fig. 4). A common garden experiment would
test the former: representatives of allegedly differentially adapted populations can be placed under the
same condition, where their fitness ranking is expected to reflect the difference between the common-
garden environment and the optimal environment predicted for them. Common garden experiments
should also be used to validate candidate drivers of adaptation, by measuring fitness of supposedly
differentially adapted individuals in two common gardens that are identical except for the focal driver.
Maps of adaptive neighborhoods can be validated in a series of reciprocal transplantations between
locations that are predicted to require increasingly divergent adaptation. Such experiments are very
challenging in animals, but there are few examples of such studies in plants. Two studies, in sorghum
(Lasky et al. 2015) and maize (Gates et al. 2019) have successfully validated that environment-associated
SNPs do indeed predict how genotypes will fare under specific environmental stressors in a common
garden experiment. In poplar, the genetic offset that accounts for unequal importance of environmental
variables for adaptation predicted the performance of individuals in a common garden better than "naive"
climate difference, which weighs all environmental parameters equally (Fitzpatrick et al. 2021). The most
interesting validation thus far has been offered not in plants but in a bird (yellow warbler), where
population declines predicted by the future genetic offset corresponded with population declines
observed in the field (Bay et al. 2018). This claim was rather extraordinary so it is hardly surprising that it
drew criticism (Fitzpatrick, Keller, and Lotterhos 2018). Indeed, genotype-environment association
inferences are not always experimentally supported: in European beech, the genomic region supposedly
associated with winter temperature adaptation did not show association with fitness in a common garden
(Lazic et al. 2024). Overall, the ecological genomics field is still under active development both in terms of
experimental design and analytical methods, and experimental validation could greatly help solidify the
methodological base, by showing what works and what doesn't.
639

640   ## Acknowledgements
641

642   This work was supported by the National Science Foundation grant OCE- 2318775 to M. V. M. and the
643   NatureNet Science Fellowship by The Nature Conservancy to K. L. B.

645   ## References
646

Balkenhol, Niko, Rachael Y. Dudaniec, Konstantin V. Krutovsky, Jeremy S. Johnson, David M. Cairns,
Gernot Segelbacher, Kimberly A. Selkoe, et al. 2017. "Landscape Genomics: Understanding
Relationships between Environmental Heterogeneity and Genomic Characteristics of Populations."
In *Population Genomics*, 261–322. Cham: Springer International Publishing.

Barghi, Neda, Joachim Hermisson, and Christian Schlötterer. 2020. "Polygenic Adaptation: A Unifying
Framework to Understand Positive Selection." *Nature Reviews. Genetics* 21 (12): 769–81.

Bay, Rachael A., Ryan J. Harrigan, Vinh Le Underwood, H. Lisle Gibbs, Thomas B. Smith, and Kristen
Ruegg. 2018. "Genomic Signals of Selection Predict Climate-Driven Population Declines in a
Migratory Bird." *Science* 359 (6371): 83–86.

Berg, Jeremy J., and Graham Coop. 2014. "A Population Genetic Signal of Polygenic Adaptation." *PLoS*
Genetics 10 (8): e1004412.

Borcard, Daniel, Francois Gillet, and Pierre Legendre. 2011. *Numerical Ecology with R*. Springer New
York.

Capblancq, Thibaut, and Brenna R. Forester. 2021. "Redundancy Analysis: A Swiss Army Knife for
Landscape Genomics." *Methods in Ecology and Evolution / British Ecological Society* 12 (12): 2298–
2309.

Caye, Kevin, Basile Jumentier, Johanna Lepeule, and Olivier François. 2019. "LFMM 2: Fast and Accurate
Inference of Gene-Environment Associations in Genome-Wide Studies." *Molecular Biology and*
Evolution 36 (4): 852–60.

De'Ath, Glenn. 2002. "Multivariate Regression Trees: A New Technique for Modeling Species–
environment Relationships." *Ecology* 83 (4): 1105–17.

Duruz, Solange, Natalia Sevane, Oliver Selmoni, Elia Vajana, Kevin Leempoel, Sylvie Stucki, Pablo Orozco-
terWengel, et al. 2019. "Rapid Identification and Interpretation of Gene-Environment Associations
Using the New R.SamBada Landscape Genomics Pipeline." *Molecular Ecology Resources* 19 (5):
1355–65.

Ellis, Nick, Stephen J. Smith, and C. Roland Pitcher. 2012. "Gradient Forests: Calculating Importance
Gradients on Physical Predictors." *Ecology* 93 (1): 156–68.

Fitzpatrick, Matthew C., Vikram E. Chhatre, Raju Y. Soolanayakanahally, and Stephen R. Keller. 2021.
"Experimental Support for Genomic Prediction of Climate Maladaptation Using the Machine
Learning Approach Gradient Forests." *Molecular Ecology Resources* 21 (8): 2749–65.

Fitzpatrick, Matthew C., and Stephen R. Keller. 2015. "Ecological Genomics Meets Community-Level
Modelling of Biodiversity: Mapping the Genomic Landscape of Current and Future Environmental
Adaptation." *Ecology Letters* 18 (1): 1–16.

Fitzpatrick, Matthew C., Stephen R. Keller, and Katie E. Lotterhos. 2018. "Comment on 'Genomic Signals
of Selection Predict Climate-Driven Population Declines in a Migratory Bird.'" *Science (New York,*
N.Y.) 361 (6401). https://doi.org/10.1126/science.aat7279.

Forester, Brenna R., Jesse R. Lasky, Helene H. Wagner, and Dean L. Urban. 2018. "Comparing Methods
for Detecting Multilocus Adaptation with Multivariate Genotype--Environment Associations."
Molecular Ecology 27 (9): 2215–33.

Gates, Daniel J., Dan Runcie, Garrett M. Janzen, Alberto Romero Navarro, Martha Willcox, Kai Sonder,
Samantha J. Snodgrass, et al. 2019. "Single-Gene Resolution of Locally Adaptive Genetic Variation
in Mexican Maize." *bioRxiv*. bioRxiv. https://doi.org/10.1101/706739.

Korneliussen, Thorfinn Sand, Anders Albrechtsen, and Rasmus Nielsen. 2014. "ANGSD: Analysis of Next
Generation Sequencing Data." *BMC Bioinformatics* 15 (November):356.

Lasky, Jesse R., Hari D. Upadhyaya, Punna Ramu, Santosh Deshpande, C. Tom Hash, Jason Bonnette,
Thomas E. Juenger, et al. 2015. "Genome-Environment Associations in Sorghum Landraces Predict
Adaptive Traits." *Science Advances* 1 (6): e1400218.

Lazic, Desanka, Cornelia Geßner, Katharina J. Liepe, Isabelle Lesur-Kupin, Malte Mader, Céline Blanc-
Jolivet, Dušan Gömöry, et al. 2024. "Genomic Variation of European Beech Reveals Signals of Local
Adaptation despite High Levels of Phenotypic Plasticity." *Nature Communications* 15 (1): 8553.

Legendre, P., and L. Legendre. 2012. *Numerical Ecology*. Elsevier.

Li, Forrest, Daniel J. Gates, Edward S. Buckler, Matthew B. Hufford, Garrett M. Janzen, Rubén Rellán-
Álvarez, Fausto Rodríguez-Zapata, et al. 2024. "The Utility of Environmental Data from Traditional
Varieties for Climate-Adaptive Maize Breeding." *bioRxiv*.
https://doi.org/10.1101/2024.09.19.613351.

Meisner, Jonas, and Anders Albrechtsen. 2018. "Inferring Population Structure and Admixture
Proportions in Low-Depth NGS Data." *Genetics* 210 (2): 719–31.

Novembre, John, Toby Johnson, Katarzyna Bryc, Zoltán Kutalik, Adam R. Boyko, Adam Auton, Amit
Indap, et al. 2008. "Genes Mirror Geography within Europe." *Nature* 456 (7218): 98–101.

Oksanen, Jari, Roeland Kindt, Pierre Legendre, Bob O'Hara, M. Henry H. Stevens, Maintainer Jari
Oksanen, and Mass Suggests. 2007. "The Vegan Package." *Community Ecology Package* 10 (631-
637): 719.

Privé, Florian, Keurcien Luu, Bjarni J. Vilhjálmsson, and Michael G. B. Blum. 2020. "Performing Highly
Efficient Genome Scans for Local Adaptation with R Package Pcadapt Version 4." *Molecular Biology*
and Evolution 37 (7): 2153–54.

Schweizer, Rena M., Bridgett M. vonHoldt, Ryan Harrigan, James C. Knowles, Marco Musiani, David
Coltman, John Novembre, and Robert K. Wayne. 2016. "Genetic Subdivision and Candidate Genes
under Selection in North American Grey Wolves." *Molecular Ecology* 25 (1): 380–402.

Smith, Chris C. R., Gilia Patterson, Peter L. Ralph, and Andrew D. Kern. 2024. "Estimation of Spatial
Demographic Maps from Polymorphism Data Using a Neural Network." *Molecular Ecology*
Resources 24 (7): e14005.

Strobl, Carolin, Anne-Laure Boulesteix, Thomas Kneib, Thomas Augustin, and Achim Zeileis. 2008.
"Conditional Variable Importance for Random Forests." *BMC Bioinformatics* 9 (July):307.

Wright, S. 1943. "Isolation by Distance." *Genetics* 28 (2): 114–38.

720 **Data and code availability**

722 *RDAforest* method is implemented as a fully documented R package and is available from gitHub at
723 https://github.com/z0on/RDA-forest. The repository also includes installation instructions, summary of
724 main functions, and script and data for the wolf analysis. The output of the wolf Rmarkdown script can be
725 viewed at https://rpubs.com/cmonstr/1224249.

Author contributions

Both authors contributed ideas that constitute the method. *RDAforest* package was developed by M.V.M, who also wrote the first version of the manuscript. Both authors contributed to finalizing the manuscript.