WRAD core perturbation impairs DNA replication fidelity promoting immunoediting in pancreatic cancer

Francesca Citron[1,20], I-Lin Ho[1], Chiara Balestrieri[2,3], Zhaoliang Liu[1], Er-Yen Yen[1,4], Luca Cecchetto[1], Luigi Perelli[1,5], Li Zhang[1,5], Luis Castillo Montanez[6], Nicholas Blazanin[1,7], Charles A. Dyke[1], Rutvi Shah[1,4], Sergio Attanasio[1], Sanjana Srinivasan[8], Ko-Chien Chen[9], Ziheng Chen[1], Iolanda Scognamiglio[1], Nhung Pham[1], Hania Khan[1,5], Shan Jiang[1,8], Jing Pan[6], Ben Vanderkruk[10,11,12], Cecilia S. Leung[1], Mahinur Mattohti[1,13], Kunal Rai[1], Yanshuo Chu[1], Linghua Wang[1], Sisi Gao[1,8], Angela K. Deem[1,8], Alessandro Carugo[8,14], Huamin Wang[15], Wantong Yao[16], Giovanni Tonon[2,17], Yun Xiong[18], Philip L. Lorenzi[18], Chiara Bonini[3,17], M. Anna Zal[19], Brad G. Hoffman[10,11], Tim Heffernan[8], Virginia Giuliani[8], Collene R. Jeter[19], Yonathan Lissanu[1,7], Giannicola Genovese[1,2], Mauro Di Pilato[6], Andrea Viale[1,20], Giulio F. Draetta[1,20].

Affiliations:
[1] Department of Genomic Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.
[2] Center for Omics Sciences, IRCCS San Raffaele Scientific Institute, Milan 20132, Italy.
[3] Experimental Hematology Unit, IRCCS San Raffaele Scientific Institute, Milan 20132, Italy.
[4] MD Anderson UTHealth Graduate School of Biomedical Sciences, Houston, TX 77030, USA.
[5] Department of Genitourinary Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.
[6] Department of Immunology, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.
[7] Department of Thoracic and Cardiovascular Surgery, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.
[8] TRACTION, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.
[9] Department of Cancer Biology, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA
[10] Department of Surgery, University of British Columbia, Vancouver, BC, V5Z 4E3, Canada
[11] Diabetes Research Group, British Columbia Children's Hospital Research Institute, 950 West 28th Avenue, Vancouver, BC, V5Z 4H4, Canada
[12] Department of Cell and Developmental Biology, University of Pennsylvania, Perelman School of Medicine, Philadelphia, PA 19194, USA

[13] Immunobiology and Transplant Science Center and Department of Surgery, Houston Methodist Hospital, Houston, TX 77030, USA

Department of Epigenetics and Molecular Carcinogenesis, Division of VP, Research

[14] Department of Oncology, IRBM Spa, Via Pontina Km 30 600, Pomezia, Rome 00071, Italy.

[15] Department of Pathology, Division of Pathology/Lab Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX

[16] Department of Translational Molecular Pathology, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.

[17] Vita-Salute San Raffaele University, Milan 20132, Italy

[18] Proteomics Core Facility, Department of Bioinformatics and Computational Biology, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.

[19] Department of Epigenetics and Molecular Carcinogenesis, Division of VP, Research, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.

[20] Corresponding authors

Correspondence:

Francesca Citron, Pharm.D., Ph.D.
Department of Genomic Medicine
The University of Texas MD Anderson Cancer Center
1901 East Road,
Houston, Texas 77054
713-745-9338
FCitron@mdanderson.org

Andrea Viale, M.D.
Department of Genomic Medicine
The University of Texas MD Anderson Cancer Center
1901 East Road,
Houston, Texas 77054
713-792-8494
AViale@mdanderson.org

Giulio F. Draetta, M.D., Ph.D.
Department of Genomic Medicine
The University of Texas MD Anderson Cancer Center
1901 East Road
Houston, Texas 77054
713-792-6390
GDraetta@mdanderson.org

Abstract

It is unclear how cells counteract the potentially harmful effects of uncoordinated DNA replication in the context of oncogenic stress. Here, we identify the WRAD (WDR5/RBBP5/ASH2L/DPY30) core as a modulator of DNA replication in pancreatic ductal adenocarcinoma (PDAC) models. Molecular analyses demonstrated that the WRAD core interacts with the replisome complex, with disruption of DPY30 resulting in DNA re-replication, DNA damage, and chromosomal instability (CIN) without affecting cancer cell proliferation. Consequently, in immunocompetent models, DPY30 loss induced T cell infiltration and immune-mediated clearance of highly proliferating cancer cells with complex karyotypes, thus improving anti-tumor efficacy upon anti-PD-1 treatment. In PDAC patients, DPY30 expression was associated with high tumor grade, worse prognosis, and limited response to immune checkpoint blockade. Together, our findings indicate that the WRAD core sustains genome stability and suggest that low intratumor DPY30 levels may identify PDAC patients who will benefit from immune checkpoint inhibitors.

Keywords: Pancreatic cancer, WRAD core, DPY30, DNA replication, chromosomal instability, immunoediting, immunotherapy

Introduction

The occurrence of aneuploidy, a condition characterized by a deviation from the normal chromosome number, is differentially sensed in normal and tumor cells (1). In normal cells, aneuploidy triggers molecular checkpoint activation and immune cells recruitment for their own destruction (2, 3). In contrast, in transformed cells, aneuploidy modulates the expression of cancer driver genes (4), favors tumor-intrinsic heterogeneity, phenotypic plasticity (1), and induces genomic instability (5), which can fuel tumor evolution (6, 7). While cell-intrinsic mechanisms have been implicated in contributing to genomic instability in tumors (8, 9), the molecular mechanisms determining the evolutionary trajectory of genomic instability in cancer cells are poorly understood.

To ensure faithful genome maintenance, precise replication of the entire genome occurs only once per cell cycle through a highly conserved and tightly regulated process that begins with the licensing of DNA replication origins (10). Replication licensing is restricted to the G1 phase of the cell cycle and is defined by the loading of the pre-replicative complex (pre-RC), which comprises the origin of replication complex (ORC1-6 complex), Cdt1 (chromatin licensing and DNA replication factor 1), Cdc6 (cell division cycle 6), and DNA helicase (minichromosome maintenance complex; MCM), onto DNA (11-13). As cells progress through the division cycle, DNA replication origin firing is regulated by the phosphorylation of the pre-RC by CDKs and DKK (Cdc7/Dbf4 kinase), which in turn recruit Cdc45 and GINS as accessory factors to fully initiate DNA replication (10, 13-16).

Inaccurate genome duplication, including under- or over-replication of DNA in late S or G2/M phase, can trigger deleterious consequences, such as DNA damage, chromosomal instability (CIN), and aneuploidy. This ultimately leads to aberrant genome architecture (17-19), dysplasia and malignant transformation when surveillance mechanisms fail to clear the damaged cells (12, 20-22). Although the mechanisms that prevent uncoordinated DNA replication in normal cells are well characterized (10, 13, 23-25), it remains unclear how cancer cells limit unscheduled DNA replication, especially given its increased frequency in the context of aberrant oncogenic signaling (e.g. *KRAS* mutation), loss of cell cycle checkpoints (e.g. *TP53*, *CDKN2A*), and cumulative alterations in epigenetic regulators (e.g. *KMT2B*, *KMT2C*, *KMT2D*, *SETD1A*, and *SETD1B*) – all of

which are commonly observed in tumors such as pancreatic ductal adenocarcinoma
(PDAC) (26-30).

Previously, we discovered that multiple members of the COMPASS complex
(WDR5, ASH2L, Menin) are critical for PDAC tumorigenesis and cancer progression in
KRAS/TP53-mutated backgrounds (31, 32). In mammals, COMPASS complexes are
composed of one of six methyltransferase catalytic subunits of the SET1/MLL family
(KMT2A, KMT2B, KMT2C, KMT2D, SET1A, SET1B) and the WRAD core (WDR5,
RbBP5, ASH2L, DPY30). The full complement of these proteins is required for efficient
histone-3-lysine-4 (H3K4) methylation as well as the regulation of transcriptional initiation
and elongation of target genes (33-36).

While WDR5 has been extensively characterized as a core scaffolding component
of SET1/MLL methyltransferases and demonstrated to be essential for gene transcription,
DPY30 is relatively understudied compared to other WRAD components. It is known that
DPY30 must bind ASH2L to enable histone H3K4 trimethylation (37), with loss of DPY30
expression leading to global reductions in H3K4 trimethylation levels in multiple model
systems (38-45). Studies in yeast suggest that, within the WRAD core, DPY30 may
stabilize the COMPASS complex to ensure the correct positioning of H3K4 methylation
at promoters, especially for low-transcribed genes. Indeed, in human and mouse models,
DPY30 expression appears to be essential for the differentiation of embryonic stem cells
as well as for hematopoietic, pancreatic, and neural progenitor/stem cells (38-44).

Here, in cancer cells, we identify a role for the WRAD core in maintaining genomic
stability that is independent of its role in SET/MLL activity. We demonstrate that the
WRAD core interacts with members of the DNA helicase complex, and, upon loss of
DPY30, tumor cells undergo DNA re-replication, resulting in genomic instability. *In vivo*,
loss of DPY30 increases the tumor mutational burden (TMB) and CIN, triggering a robust
CD8-dependent anti-tumor immune response. Our findings implicate the WRAD core in
preserving genome stability in cancer cells and suggest that DPY30 inhibition merits
further exploration as a therapeutic concept to kill tumor cells by genome destabilization
and sensitization to immune-checkpoint inhibitor therapy.

Results

Loss of DPY30 is dispensable for PDAC cell proliferation and H3K4 methylation.

We first assessed the impact of silencing the expression of proteins comprising the WRAD core on PDAC cell proliferation. Consistent with our previous observations (31, 32), doxycycline-induced silencing of *WDR5* (*shWDR5*), *ASH2L* (*shASH2L*), or *RBBP5* (*shRBBP5*) strongly restrained cell proliferation *in vitro* (**Fig. 1a, Extended Data Fig. 1a-e**). Unexpectedly, shRNA-mediated silencing of *DPY30* (*shDPY30*) in human PDAC cells, as well as sgRNA-mediated *Dpy30* knockout (*Dpy30*-KO) in mouse PDAC cells from KPC (*Ptf1a^{Cre};KRas$^{lsl-G12D/wt}$;Trp53$^{R172H/wt}$*) (46) and KP$^{-/-}$C (*Ptf1a^{Cre};KRas^{lsl-fl};Trp53$^{fl/fl}$*) (47)mouse models did not affect cell proliferation (**Fig. 1b-c, Extended Data Fig. 1f-n**). We confirmed the lack of any effect of DPY30 loss on cell proliferation *in vivo*, showing that human *DPY30*-KO PDAC cells injected into immunodeficient NSG mice did not grow differently compared to controls (**Fig. 1d-g**, **Extended Data Fig. 1o-p**). Hence, our data show that PDAC cells are vulnerable to the loss of the WRA (WDR5, RBBP5, ASH2L) core, but not to DPY30 depletion.

Because the three-component WRA core has been shown to be the minimal required cofactor for supporting SET/MLL methyltransferase activity on histone H3K4, while the presence of DPY30 is only needed to enhance the specificity of the enzymatic activity (33), we explored whether inhibiting *DPY30* or *WDR5* expression in PDAC cells could differentially affect H3K4 methylation. To this end, we compared the acute effects of *DPY30* or *WDR5* loss on chromatin profiles in patient-derived PDAC cell lines transduced with doxycycline-inducible *shDPY30* or *shWDR5* by performing transposase-accessible chromatin using sequencing (ATAC-seq) and H3K4 tri-methylation (H3K4me3) and H3K27 acetylation (H3K27ac) chromatin immunoprecipitation sequencing (ChIP-seq), the latter of which provides readouts of COMPASS activity at gene regulatory regions (31, 48). Because WDR5 loss has a detrimental impact on the cell cycle (31, 39) (**Fig. 1a**), we excluded any confounding effects of WDR5 loss on proliferation by collecting cells at 48 hours after shRNA induction *via* doxycycline treatment, a timepoint sufficient for protein downregulation (**Extended Data Fig. 2a-b**). Cluster analysis of open genomic regions, as well as genomic regions marked by H3K4me3 and H3K27ac, revealed that *WDR5*-silenced cells exhibited chromatin states

that were distinct from *DPY30*-silenced cells, which clustered with control cells. This
suggests that, unlike the effects observed upon WDR5 loss, the acute and short depletion
of *DPY30* is not sufficient to alter chromatin accessibility and states (**Fig. 1h-j**).
Consistently, WDR5, but not DPY30, downregulation affected the total levels of H3K4me3
and H3K4me1 histone marks (**Extended Data Fig. 2b, f**).

Although genome-wide analysis of chromatin at regulatory regions proximal to
gene transcriptional start sites (TSS) did not reveal significant differences between
DPY30 and *WDR5* silenced cells (**Extended Data Fig. 2c-e**), the H3K4me3 profiles of
these two groups appear markedly different. Specifically, *DPY30* silencing did not affect
the pattern of H3K4me3, indicating that DPY30 loss likely does not broadly alter the
transcriptome of these cancer cells (**Fig. 1k**). In contrast, *WDR5* silencing significantly
altered H3K4me3 patterns, resulting in the repression of thousands of genomic regions
(**Fig. 1k, Extended Data Table S1**). To determine the functional annotations for the
regions affected by WDR5 depletion, we conducted pathway analysis of genes located in
these genomic regions and identified pathways involved in interferon alpha and gamma
response, inflammatory response, and DNA damage (**Extended Data Fig. 2g, Extended
Data Table S1**).

To dissect the biological role of the WRAD core in PDAC cells, we next performed
bulk RNA-sequencing (RNAseq) analysis of patient-derived PDAC cells stably expressing
shDPY30, *shWDR5*, or shCTR and treated with doxycycline for 48 hrs as described
above (**Extended Data Fig. 2a**). *WDR5*, but not *DPY30*, silencing profoundly modulated
the transcriptome of PDAC cells (**Extended Data Fig. 3a**), which is consistent with the
essential role of WDR5 in regulating gene transcription (31, 36, 39, 49). Specifically, and
in line with our epigenomics analysis (**Fig. 1k, Extended Data Table 1**), *WDR5* silencing
induced a remarkable downmodulation of genes involved in the interferon alpha and
gamma response as well as a dysregulation of pathways involved in mRNA translation
and RNA processing (**Extended Data Fig. 3b, Extended Data Table S2**). Although
DPY30 silencing did not deregulate the transcriptome of PDAC cells as strongly as *WDR5*
silencing, it induced a positive regulation of pathways involved in interferon signaling,
inflammatory response, and DNA damage (**Extended Data Fig. 3e-g, Extended Data
Table 2**). Overall, our data indicate that *DPY30* or *WDR5* silencing leads to distinct

epigenomic and transcriptomic changes in PDAC, thus demonstrating that the WRAD

core has non-redundant roles in maintaining cell proliferation and transcriptional activity

in PDAC.

WRAD core directly interacts with the MCM complex.

To further investigate the discrepancies in the proliferative, epigenetic, and

transcriptional phenotypes that emerged upon depletion of different WRAD core members

in PDAC cells, we expanded our investigation into the biological function of DPY30 in

PDAC. To this end, we immunoprecipitated endogenous DPY30 protein in two human

(PATC53 and PATC66) and one mouse (KPC) PDAC-derived cell models, all of which

carry *TP53* and *KRAS* mutations (31, 32). Unbiased proteomic analysis showed that

multiple top-scoring interactors of DPY30 are involved in chromatin remodeling, RNA

metabolism, splicing, and gene transcription (**Extended Data Fig. 4a-d, Extended Data**

Table S3), consistent with the role of DPY30 as a cofactor supporting histone methylation

and gene transcription regulation by COMPASS complexes (38-44). Proteins involved in

DNA replication, DNA damage, and cell cycle regulation were also identified as DPY30

interactors (**Extended Data Fig. 4a-d, Extended Data Table S3**). These included

proteins previously unknown to interact with DPY30, such as multiple members of the

DNA replisome, including the MCM complex and WDHD1 (**Fig. 2a, Extended Data Fig.**

4d). We validated these findings by expressing either mouse or human HA-tagged DPY30

with mouse or human MCM subunits and found that DPY30 bound most strongly to MCM6

compared to others (**Fig. 2b, Extended Data Fig. 4e**). Notably, proximity ligation analysis

(PLA) revealed that endogenous DPY30 lies in proximity (<40nm) of both MCM6 and

WDHD1, strongly suggesting that DPY30 may be a core component of the DNA replisome

(**Fig. 2c**). To explore whether the interaction of DPY30 with the replisome was cell cycle-

dependent, we expressed the fluorescent ubiquitination-based cell cycle indicator

(FUCCI) system in human PDAC cells (50). By enabling the real-time tracking of cell cycle

progression in live cells (54), this approach allowed us to sort PDAC cells by cell cycle

phase and then perform immunoprecipitation (IP) assays for endogenous DPY30

(**Extended Data Fig. 4f-g)**. Unbiased proteomic analysis of affinity-purified samples

showed that members of the COMPASS complex were, as expected, enriched in DPY30

pull-downs and that these interactions were independent of cell cycle phase (**Extended**

Data Fig. 4h). Furthermore, this analysis confirmed that WDHD1 and all MCM complex

members, including MCM10, MCM3AP, and MCMBP, interacted with DPY30, with all

replisome components except for MCM2, MCM4, and MCM10 interacting with DPY30 in

a cell cycle-independent fashion **(Extended Data Fig. 4i).** Additionally, we found that

WDR5 also interacted with MCM complex members (**Fig. 2d)**. In cell nuclei, WDR5 was

shown to reside topographically close to MCM6 and WDHD1 (**Fig. 2e-f, Extended Data**

Fig. 4j), similar to what was seen for DPY30 (**Fig 2c**). Importantly, endogenous WDR5,

ASH2L, and RBBP5 could be precipitated with MCM6, suggesting that the whole WRAD

participates in this interaction (**Extended Data Fig. 4k-m)**. Interestingly, destabilization

of the WRAD core through silencing of either *WDR5* or *DPY30* impeded the endogenous

binding of DPY30 to MCM6 (**Fig. 2g-h).**

To explore whether WRAD core integrity is necessary for MCM6 binding, we

overexpressed a mutant isoform of DPY30 (L69D) known to hinder the DPY30:ASH2L

interaction and the proper assembly of the WRAD core (51). Remarkably, DPY30^{L69D}

expression prevented the DPY30:MCM6 interaction as well as the assembly of the WRAD

core **(Fig. 2i, Extended Data Fig. 4n)**. We employed a similar approach by

overexpressing two WDR5 mutant isoforms (L240K and V268E) that have altered WDR5-

binding motif (WBM) domain and are known to disrupt the interaction of WDR5 with

RBBP5 and cMyc (31). We consistently found that disrupting the WRAD core assembly

strongly reduced the interaction between WDR5 and MCM6 **(Fig. 2j, Extended Data Fig.**

4o). Altogether these observations suggest that the stability of the WRAD core is

necessary for the interaction between the WRAD core and the MCM6 subunit of the MCM

complex.

The WRAD core participates in replication fork dynamics

Because the MCM complex represents the DNA helicase activated at the start of

DNA replication (10, 12-15, 23), we tested whether WRAD contributes to replication fork

progression. Using an iPOND assay to recover proteins bound to newly synthesized DNA,

we found that WRAD core members localized at DNA replication fork initiation sites (2

min of EdU pulse), persisted during DNA elongation (5-10 min of EdU pulse), but

disappeared at later time points (chase) (**Fig. 3a, Extended Data Fig. 5a**). Notably,
histone H3, a marker for nucleosome incorporation and an indicator of SET/MLL
canonical activity, was barely detectable at 2 min of EdU pulse, gradually incorporated at
5 and 10 minutes, and strongly detected during the chase (**Fig. 3a, Extended Data Fig.**
5a). This contrasting behavior between WRAD and H3 serves as an internal control,
confirming that WRAD members detected at DNA replication forks are not associated
with chromatin-bound proteins (**Fig. 3a, Extended Data Fig. 5a**). These data strongly
indicate that, in the context under study, the WRAD core directly interacts with the DNA
replication fork in a nucleosome-independent manner. Importantly, analysis of the
interactions between proteins and the nascent DNA using a single cell-based method
(SIRF assay) (52, 53) showed that WDR5, but not DPY30, directly interacts with nascent
DNA, further supporting a pivotal role for the WRAD core in DNA replication dynamics
(**Fig. 3b**).

Because WDR5 deregulation affects distinct biochemical pathways and results in
rapid induction of cell death (**Fig. 1**) (31), we sought to avoid any confounding effects by
focusing further mechanistic studies on DPY30. First, iPOND analysis found that *Dpy30*-
KO did not destabilize MCM6 nor WDR5 binding at replication forks, suggesting that
DPY30 is not directly involved in regulating fork assembly (**Fig. 3c, Extended Data Fig.**
5b). Second, to examine the effect of DPY30 on replication fork efficiency, we performed
a DNA fiber assay in *Dpy30*-KO and control cells. *Dpy30*-KO resulted in slower fork
progression rates and reduced symmetry between incorporation of the two different
nucleotide analogues, suggesting that *Dpy30* loss may promote replication stress (**Fig.**
3d-e). Additionally, we observed increased frequency of overlap between the two
incorporated nucleotide analogues in *Dpy30*-KO cells compared to controls, indicating
DNA re-replication in these cells, and further supporting the hypothesis that *Dpy30* loss
may contribute to replication stress (**Fig. 3f-g**).

To corroborate our findings, we next developed an unbiased DNA re-replication
sequencing assay ("Re-repli-seq", see methods section) (**Extended Data Fig. 5c**).
Specifically, S phase synchronized KPC controls or Dpy30-KO cells were released in
presence of CldU (45 min) followed by a second pulse of IdU (45 min) (**Extended Data**
Fig. 5c). The cells were then split, and a ChIP assay was performed using antibodies

against the incorporated CldU and IdU separately, followed by next generation sequencing. After, sequence alignment to reference genome, we represented the data as a ratio of incorporated CldU:IdU (38). In case of DNA re-replication, sequencing of double-labelled immunoprecipitated genomic regions should result in an increased overlap between CldU and IdU sequencing reads. Data analysis showed that, compared to controls, the Dpy30-depleted KPC cells showed broader genomic domains with CldU:IdU ratio close to zero, which means the nucleotide analogues are incorporated equally and overlap in the same genomic regions (**Figure 3h-i**). As an example, regions of different chromosomes (Chr1, Chr2, Chr4, Chr6, Chr10 and Chr16) displayed domains of replicated DNA with an overlapped pattern of CldU:IdU ratio more prominently distributed in Dpy30-depleted cells compared to control cells, that instead, showed a more coordinated replication profile (**Extended Data Fig 5d-e**). Our genome-wide analysis of CldU:IdU incorporation strongly supports that Dpy30 and the WRAD core are required to sustain a coordinated DNA replication during the S phase in tumor cells.

Because excessive DNA replication can trigger DNA damage, we found that during unperturbed growth condition, *Dpy30*-KO KPC cells showed stronger γH2AX (pS139) and pRPA (pS4-8) staining compared to control cells (**Fig. 3j, Extended Data Fig. 5f-h**). Consistently, γH2AX levels *in vivo* were dramatically increased in both *Dpy30*-KO KPC and human-derived *DPY30*-KO or knockdown (KD) tumors compared to controls (**Extended Data Fig. 6a-b**). Further, bulk RNAseq analyses performed in mouse-derived Dpy30-KO and control PDAC cells revealed a deregulation of pathways involved in DNA damage and G2/M checkpoint (**Extended Data Fig. 6c-d, Extended Data Table S4**), Interestingly, Dpy30-depleted cells showed an extensive upregulation of inflammatory pathways (**Extended Data Fig. 6c-d, Extended Data Table S4)**, which might have been activated as a consequence of persistent DNA damage (2, 55, 56). These data align with our previous work (31) showing that the WDR5 silencing strongly increased DNA damage and induced a sustained G2/M accumulation in PDAC cells. Further, these observations suggest that PDAC cells can tolerate the incremental DNA damage caused by *DPY30* loss, but cannot survive the extensive DNA injuries and transcriptomic rewiring upon WDR5 depletion. This suggests that the intact WRAD core sustains genomic stability by counteracting excessive DNA replication.

DPY30 maintains genome integrity by limiting excessive DNA replication and stabilizing the S-to-G2/M transition

The risk of DNA re-replication increases at the S-to-G2/M boundary, during which Cdc6 and Cdt1 are in their active forms and the expression of geminin, an inhibitor of DNA replication, has not yet peaked (10, 13, 23). To investigate whether *DPY30* loss disrupts cell cycle dynamics in these phases specifically, we synchronized PDAC cells in S phase using a double-thymidine block and collected cells at different time points after release (**Extended Data Fig. 6e-f**). In line with our observations in proliferation assays (**Fig. 1a-d, Extended Data Fig. 1a-n**), we observed that *Dpy30* KO did not grossly affect cell cycle distribution in non-synchronized cells (**Extended Data Fig. 6e-f**). However, in synchronized cells, a larger proportion of *Dpy30*-KO cells compared to control cells advanced towards the G2/M boundary by 2 hours after release from S phase (**Extended Data Fig. 6e-f**). Additionally, at this time point, we detected an increased percentage of aneuploid cells (DNA content >4N), higher DNA damage, increased BrdU incorporation, as well as higher phosphorylation of pS139 MCM2, pS54 Cdc6, and pS10 histone H3 in Dpy30-KO cells compared to controls, all of which support an uncoupling of DNA replication and cell cycle progression upon *Dpy30* loss (**Extended Data Fig. 6e-h**). We further observed that, while there were no significant differences in the expression of geminin in the arrested cells (**Extended Data Fig. 6h-i**), geminin expression was strongly downmodulated upon release in *Dpy30*-KO cells. We next synchronized *Dpy30*-KO or control PDAC cells in S phase using a double thymidine block, and in G2/M phase using a thymidine-nocodazole block (**Extended Data Fig. 6j)**. Strikingly, Western blot analysis showed that the dysregulation of Cdc6 and geminin expression was restricted to the G2/M phase rather than the early S phase (**Extended Data Fig. 6j**). Overall, our data indicate that, upon DPY30 depletion, the risk of DNA re-replication increases at the S-to-G2/M boundary due to the uncoupled regulation of the Cdc6/Geminin axis (20, 24, 57-59).

During the G2/M and G1 phases, Geminin stability is tightly regulated by the anaphase promoting complex/cyclosome (APC/C), leading to its proteasome-dependent degradation (60). As a member of the E3 ubiquitin ligase family, APC/C requires the modification of the cullin ring with NEDD8, an-ubiquitin-like protein, for its activity (61).

Chemical inhibition of NEDD8 promotes DNA re-replication through the stabilization of

Cdt1 and Cdc6 protein expression (24, 61-63). Hence, we hypothesized that NEDD8

inhibitors, such as MLN4924, may exacerbate the uncoupling between DNA replication

and G2/M transition in Dpy30-depleted cells by. As expected, human nearly diploid RPE-

1 cells treated with MLN4924 showed increased expression of Cdc6 in both control and

DPY30-KO cells compared to untreated ones (**Extended Data Fig. 7a**). Further, when

compared to control RPE-1 cells, geminin expression was remarkably lower upon DPY30

depletion and MLN4924 treatment, and the spontaneous activation of γH2AX was

observed upon DPY30 depletion and then boosted by MLN4924 treatment (**Extended**

Data Fig. 7a). Finally, we observed increased BrdU incorporation in cells treated with

MLN4924, with the highest incorporation in *DPY30*-KO cells (**Extended Data Fig. 7b**).

These data were confirmed in KPC cells, showing dysregulated geminin and γH2AX

expression, as well as increased percentage of aneuploid cells, in *Dpy30*-KO compared

to control cells (**Extended Data Fig. 7c-d**). Overall, our data suggested that *Dpy30* loss

leads to DNA re-replication through a potent perturbation of the Cdc6/geminin balance

during the G2/M phase.

Finally, because uncoordinated replication can contribute to chromosomal

instability (CIN), we investigated whether DPY30 may induce this phenomenon (16, 22,

25, 64). Indeed, *DPY30*-KO in both human and mouse PDAC-derived models increased

the percentage of cells with chromosome mis-segregation and micronuclei *in vitro* and *in*

*vivo (***Fig. 3k, Extended Data Fig. 7e-g***)*. Consistently, metaphase spreads showed an

increased in the number of chromosomal breaks and fusions *in vitro* (**Fig. 3l, Extended**

Data Fig. 7h-i). Collectively, these data support a critical role for DPY30 in maintaining

genome stability and suppressing DNA re-replication at the G2/M transition.

***DPY30*-KO cells display complex karyotypes and elicit a T cell-mediated**

immune response.

To assess whether the genome instability induced by DPY30 loss may impact

tumor immunogenicity *in vivo*, we injected control or *Dpy30*-KO KPC and KP$^{-/-}$C cells into

cohorts of NSG, nude, and C57BL/6 mice. Strikingly, *Dpy30*-KO strongly impaired tumor

growth compared to controls in immune-competent C57BL/6 mice but had no effect on tumor growth in immunodeficient mice (**Fig. 4a-d, Extended Data Fig. 8a-d**). Moreover, histological analysis revealed that *Dpy30*-KO tumors grown in C57BL/6 mice appeared more differentiated and characterized by an increased number of mucin-positive ductal-like structures compared to their respective controls (**Fig. 4e-f**). Of note, in models of orthotopic transplantation, we observed that Dpy30 depletion significantly prolonged animal lifespan (**Fig. 4g, Extended Data Fig. 8e**).

To eliminate potential confounding effects due to CRISPR/Cas9-related immunogenicity (65), we injected mouse cells derived from $KP^{-/-}C$ tumors stably expressing doxycycline-inducible shDpy30 or shCTR into NSG or C57BL/6 mouse backgrounds. Also in this setting, we confirmed that Dpy30-depleted tumors grown in immunocompetent mice were smaller and more differentiated compared to controls, while in NSG mice no major differences were detected (**Extended Data Fig. 8f-i**). Further, we confirmed that *Dpy30* silencing strongly prolonged animal lifespan (**Extended Data Fig. 8j**).

Intriguingly, the cell proliferative index, evaluated as Ki67- and pS10-H3-positive staining, was remarkably reduced in *Dpy30*-KO cells compared to controls in immune-competent, but not NSG, backgrounds (**Figure 4h, Extended Data Fig. 5g and 8k-l**). This suggests that, in the absence of DPY30, the immune system may inhibit growth of or preferentially clear proliferating tumor cells. Interestingly, gene set enrichment analysis (GSEA) from bulk RNAseq data of *Dpy30*-KO and control $KP^{-/-}C$ tumors showed a positive enrichment score for pathways involved in DNA damage and repair process in immunodeficient mice, while the same cancer cells showed upregulation of gene pathways involved in inflammation, including interferon signaling, the complement cascade, and PD-1 signaling, in an immunocompetent background (**Extended Data Fig. 8m-n, Extended Data Table S5**). Because CIN can enhance the activity of the cGAS/STING pathway, which in turn regulates the expression of the type I interferons and related gene products (56), we sought to evaluate the expression of STING upon Dpy30 depletion. Immunostaining analysis of KPC showed a potent activation of STING in *Dpy30*-KO cells compared to controls both *in vitro* and *in vivo* (**Fig. 4i, Extended Data Fig. 8o).** Altogether, these data suggest that *Dpy30* loss can trigger a T cell-mediated

immune response *in vivo* through DNA damage and CIN-dependent activation of the type

I interferons, the latter of which occurs through the cGAS/STING pathway.

Next, we performed single-cell RNAseq (scRNAseq) on *Dpy30*-KO and control

KPC tumors (n=3 per condition) grown in immunocompetent mice. Dimensionality

reduction analysis of sub-clusters of tumor cells revealed only a partial overlap between

the transcriptomes of *Dpy30*-KO and control tumor cells (**Fig. 4j**). Specifically, we

identified a population of highly proliferative tumor cells that was expanded in control

tumors but minimally represented in *Dpy30*-KO tumors (**Fig. 4j-k**). Expression analysis of

cell cycle regulators corroborated our observation, showing a reduced expression of

proliferative genes (*Ccnb2, Cdc20, Top2a, Aurka, Aurkb, Wee1*) (**Fig. 4k**) and,

consequently, supporting the hypothesis that highly proliferative tumor cells are

selectively cleared by the immune response in the absence of Dpy30. Further, GSEA

confirmed an upregulation of genes involved in interferon signaling and inflammatory

response pathways (Il6/Stat3, allograft rejection, and complement) upon Dpy30 depletion

(**Fig. 4l**). Remarkably, *B2m*, *Cd74*, *H2-K1*, and *H2-D1*, genes involved in antigen

presentation, were among the top-scoring genes in *Dpy30*-KO cells (**Fig. 4m**). Indeed,

when comparing *Dpy30*-KO tumors to controls, immunostaining analysis confirmed an

increased infiltration of CD3+ tumor-infiltrating lymphocytes (TILs) as well as an increase

of cancer associated-fibroblasts (CAFs) (αSMA+), the latter likely the result of the

depletion of the epithelial compartment (**Fig. 4n, Extended Data Fig. 8p**). Accordingly,

examination of the tumor microenvironment (TME) at the single cell level revealed an

increased number of CAFs, B cells, NK cells, and T cells in *Dpy30*-KO tumors grown in

an immunocompetent background when compared to controls (**Fig. 5a-c, Extended Data

Fig. 9a-b**). These findings were corroborated by flow cytometry-based

immunophenotyping, which demonstrated a significant increase in the number of all

infiltrating T cell subsets and NK cells in *Dpy30*-KO tumors. Among the immune cells, the

CD8+ T cells were the predominant population, suggesting that they may be the primary

contributor to the anti-tumor response (**Fig. 5d, Extended Data Fig. 9c-e**). Importantly,

compared to control tumors, CD8+ T cells in *Dpy30*-KO tumors displayed lower

expression of exhaustion marker *Tox* as well as higher expression of cytotoxic markers

such as interferon gamma (*Ifng*), granzyme B (*Gzmb*), and natural killer cell granule

protein 7 (*Nkg7*) (**Fig. 5e, Extended Data Fig. 9f-h**). Altogether, these findings suggest that CD8+ T cells are activated and may selectively clear *Dpy30*-KO tumor cells with increased immunogenicity due to genomic instability.

To test this hypothesis, whole-exome sequencing (WES) was performed on *Dpy30*-KO or control tumors grown in NSG or C57BL/6 mice. We found that, in a permissive immune-deficient environment, *Dpy30*-KO tumors grew without restraint and tolerated a higher mutational burden (TMB) compared to *Dpy30*-proficient controls (**Extended Data Fig. 9i**). However, the same cells transplanted in a selective immune-competent background displayed lower TMB (**Extended Data Fig. 9i**), thus suggesting that cancer cells undergoing uncoordinated DNA replication and CIN were being cleared from tumors. To corroborate our findings and quantify the extent of this depletion, we generated barcoded KPC cells using our established platform (66, 67). We then transiently infected barcoded KPC cells with sgRNAs targeting *Dpy30* or a scramble control and injected them into NSG or C57BL/6 mice. Explanted tumors were subjected to next-generation sequencing (NGS) for barcode detection. Pairwise comparison of subclonal composition of *Dpy30*-depleted versus control KPC cells grown in immunodeficient or immunocompetent mice showed a dramatic depletion of barcodes in the *Dpy30*-KO cells transplanted in C57BL/6 mice compared to an immunodeficient counterpart (**Fig. 5f**). Further, we injected *Dpy30*-KO or control KPC cells into immunocompetent C57BL/6 mice and collected tumors at different time points (day 10, day 15 and day 25). Multiplexed immunostaining analysis showed that, over time, the number of TILs strongly increased exclusively in *Dpy30*-KO tumors compared to controls (**Fig. 5g-h**). This immune infiltration, in the *Dpy30*-KO tumors, was paralleled by a net increase in the cleaved caspase 3 (CC3) staining, a *bone fide* marker of apoptotic cell death, as well as a pronounced reduction in the number of Ki67+ tumor cells (**Fig. 5i-j**). Altogether these data indicate that, in a mutated *TP53/KRAS* genetic background, increases in CIN and TMB due to DPY30 loss enhance the immunogenicity of PDAC cells and a consequent T cell-mediated response.

DPY30 expression correlates with poor response to immune checkpoint blockade and tumor staging.

To determine whether CD8+ T cells are responsible for clearing cells harboring a

high TMB (high-TMB) due to DPY30 depletion, we injected *Dpy30*-KO or control KPC

cells into C57BL/6 mice and depleted the CD8+ T cell population using an anti-CD8

monoclonal antibody (**Fig. 6a**). We found that CD8+ T cell depletion was sufficient to

rescue the phenotype induced by *Dpy30* loss, considerably accelerating tumor growth,

and reducing lifespan when compared to that of control mice (**Fig. 6b-c**). To explore

whether the activation of cytotoxic T cells upon *Dpy30* loss may synergize with

immunotherapy, we randomized transplanted mice to receive either 8 mg/kg of anti-PD-

1 monoclonal antibody or isotype control intraperitoneally three times a week for two

consecutive weeks. Anti-PD-1 treatment dramatically reduced tumor growth and greatly

extended the median survival of mice bearing *Dpy30*-KO tumors, with 60% of these

animals showing complete tumor regression (**Fig. 6a-c**).

To explore the translational relevance of our findings, we next evaluated *DPY30*

expression in PDACs by performing immunohistochemical (IHC) analysis on a tissue

microarray (TMA) comprising of human PDAC (*n* = 128) and matched tumor-adjacent

normal pancreas tissue samples (*n* = 128) collected at the University of Texas MD

Anderson Cancer Center. We found that *DPY30* expression was low in normal pancreatic

tissues but high in PDAC, particularly in poorly differentiated tumors (**Fig. 6d**). Moreover,

we showed that low *DPY30* expression was associated with better survival in the cohort

of patients included in the TMA (**Fig. 6e**). To further investigate this association, we

interrogated The Cancer Genome Atlas (TCGA) PDAC database and found that,

compared to other WRAD core members, high *DPY30* expression was significantly

associated with shorter overall patient survival (**Fig. 6f, Extended Data Fig. 10a-c**).

Based on these clinical findings, we hypothesized that DPY30 expression in PDAC may

be modulated over tumor progression. To test this, we collected pancreas tissues from

the KPC mouse model at different time points (46, 68, 69). In support of our hypothesis,

IHC analysis showed low DPY30 expression in normal pancreas and metaplastic lesions

but a striking increase in expression in transformed cells that became increasingly higher

as the lesions progressed from pancreatic intraepithelial neoplasia (PanIN) to high-grade

PDAC (**Extended Data Fig. 10d**). We also noticed that, in normal pancreas, DPY30 was

mostly expressed in the endocrine compartment, in line with its role in endocrine cell

functional maturation (38, 39), while being weakly detectable in pancreatic ductal and
acinar cells (**Extended Data Fig. 10d**). These data provide evidence that DPY30
expression is associated with PDAC progression in both human and mouse.

To examine whether DPY30 may similarly affect genome integrity and the immune
response in human tumors, we queried the transcriptomic data of the TCGA PDAC patient
cohort. GSEA analysis of 70 samples with at least 50% purity distributed according to
high or low *DPY30* expression showed that low *DPY30* expression was associated with
increased expression of genes involved in the complement response as well as in
interferon and IL6/STAT3 signaling pathways (**Fig. 6g**). We corroborated our data by
querying the transcriptomic data of 47 PDAC patient-derived xenografts (PDXs)
established at UT MD Anderson Cancer Center PDACs. GSEA analysis showed that
PDAC PDXs expressing low levels of *DPY30* are enriched in genes belonging to the
interferon alpha and gamma response pathways (**Extended Data Fig. 10e**). Finally, we
interrogated the dataset of the phase 2 PRINCE trial, in which the efficacy of the PD-1
inhibition (nivolumab) was assessed in patients with metastatic PDAC (70). Strikingly,
PDAC patients expressing low DPY30 levels showed a prolonged progression-free and
overall survival compared to PDAC patients with high DPY30 expression (**Fig. 6h,**
Extended Data Fig. 10f-g). Taken together, our data strongly suggest that DPY30 has a
role in regulating DNA replication in PDAC cells, and that low levels of DPY30 favor
aberrant DNA replication events that activate an interferon-dependent response that
recruits effectors of the adaptive immune response.
486

Discussion
While many studies have focused on the genetics and transcriptional fingerprint of
PDAC to designate subtype classification schemes for this highly heterogeneous disease,
these molecular subtypes have not yet consistently translated to clinical implications (28,
71-73). Cumulative evidence sheds light on the intricate mechanisms underpinning PDAC
tumorigenesis and disease progression, which involve context- and time-dependent
interactions between genomic and epigenomic scars (74, 75). Recently, a study of the
epigenome of PDAC PDXs demonstrated that epigenetic fingerprints are associated with

discrete transcriptional signatures and clinical features, suggesting a convergence
between genetics, epigenetics, tumor-intrinsic, and tumor-extrinsic features (76).

The disruption of evolutionarily conserved epigenetic complexes, including
COMPASS, has been implicated in developmental disorders and cancer, as strict
regulation of the activity of these complexes are critical for developmental and
physiological processes (37, 48). For instance, in PDAC, the cumulative incidence of
mutations in epigenetic regulators is approximately 35%, including recurrent mutations in
KMT2B, *KMT2C*, *KMT2D*, *SETD1A*, and *SETD1B* (26-30). Interestingly, *DPY30*
mutations have not been observed in any cancer.

While the WRAD core of COMPASS complexes is known to regulate the
maintenance of epigenetic information in normal and cancer cells, numerous studies have
demonstrated non-canonical functions of the WRAD subunits that include transcriptional
regulation-independent functions that regulate cell cycle progression (33-35, 49, 77-81).
Here, we demonstrate a novel function for the WRAD core as an effector in DNA
replication and fork stability, and we show that this function is independent of the core's
function as an MLL/Set methyltransferase (38-40, 42-44). Although loss of DPY30 did not
affect the proliferative ability of PDAC cells compared to the inactivation of other WRAD
core members, we show that DPY30 integrity is required to prevent excessive DNA
replication and maintain genome integrity. Specifically, molecular analyses revealed that
DPY30 loss leads to uncoordinated DNA replication, resulting in increased replication
stress and, ultimately, promoting genome instability. Moreover, *DPY30*-depleted cells
displayed a rapid and unchecked transition from S phase to G2/M and were observed to
progress through the cell cycle with mis-segregated chromosomes and persistent DNA
damage.

The S-to-G2 cell cycle checkpoint is enforced by the ATR kinase, which governs
the coupling of DNA replication with mitotic progression by repressing the activity of CDK1
and ensuring proper timing of G2-specific events (82-84). DNA re-replication can itself
activate ATR, which in turn arrests DNA replication and recruits DNA repair effectors (13,
15, 25). However, the biological mechanisms by which this ATR-mediated checkpoint can
sense the completion of DNA replication and distinguish this event from DNA over-
replication remain unknown. Our findings shed light on how tumor cells with oncogenic

signaling and deficient cell cycle checkpoints employ independent, novel pathways to restrain DNA replication to evade cell death mechanisms. Clarifying the interplay between DNA re-replication control, checkpoint activation, and repair of re-replication-associated DNA lesions is fundamental for expanding our knowledge of the biology driving cancer cells.

Whole-genome duplication and CIN frequently occur during the progression of PDAC, and especially at late and metastatic stages. In particular, chromothripsis, a catastrophic mitotic event driving multiple concurrent chromosome structural alterations, is frequently observed in advanced PDAC (30). Interestingly, *DPY30* expression is upregulated during tumorigenesis, and poorly differentiated PDACs displayed the highest *DPY30* expression. This suggests that DPY30 may somehow buffer PDAC cells from oncogene-driven uncontrolled replication and the detrimental effects of CIN. All models evaluated in our study were *P53* deficient, thus warranting additional investigations to determine whether *P53* inactivation represents a specific genetic context in which DPY30 function is essential for regulating DNA homeostasis. Further, because aneuploid cells are characterized by deregulated metabolic programs and protein networks, future studies should assess whether DPY30 loss, through induction of aneuploidy, can evoke cellular adaptive responses, which may ultimately be exploited as co-lethality and druggable targets (1, 6, 85).

Overall, our study demonstrates that DPY30 loss leads to the accumulation of PDAC cells with complex karyotypes and increased TMB, which, in turn, results in the selective clearance of damaged proliferating cells by CD8+ T cells. Considering the paucity of treatment options for patients with PDAC, our work provides evidence that PDAC with low *DPY30* expression may respond to combined therapy of immune-checkpoint inhibitors and conventional chemotherapy.

Methods

Patient-derived samples. Patient-derived samples were obtained from consented patients under an Institutional Review Board (IRB)-approved protocol LAB00-0396 chaired by Dr. Michael Kim. (UTMDACC).

Cell Biology Experiments

Cell culture. Human 293T and RPE-1 (ATCC), PDX-derived PATC53 (PDX1),

PATC66 (PDX2), and mouse-derived PDAC cells KP$^{-/-}$C (internally registered as WY5549

and derived from *Ptf1aCre;LSL-Kras$^{G12D/+}$;LSL-Trp53$^{L/L}$* mouse model) and KPC

(internally registered as N150F and derived from *Ptf1aCre;LSL-Kras$^{G12D/+}$;LSL-*

Trp53$^{R172H/R172H}$ mouse model, and named as N150F, see below "mouse strains were

cultured in Dulbecco modified Eagle medium (DMEM, Sigma) supplemented with 10%

fetal bovine serum (FBS, Gibco), 100IU/mL Penicillin (Gibco) and 100µg/mL

Streptomycin (Gibco).

Lentiviral production. Viral particles were produced using packaging plasmids

psPAX2 and pMD2.G. Briefly, 293T cells (ATCC) were seeded in 15cm diameter plate in

DMEM supplemented with 10% FBS (Gibco), 100IU/mL Penicillin (Gibco), 100µg/mL

Streptomycin (Gibco), 2 mM Glutamine (Gibco) and allowed to adhere. Transfection mix

was prepared by adding the vectors of interest in Opti-MEM (Gibco) in presence of

Polyethyleneimine (PEI). Supernatant was collected 72 hours after transfection, filtered

through 0.45µm low-protein binding filters (Corning) and concentrated in sterile PBS

(Sigma Aldrich) after ultracentrifugation at 25,000 rpm for 2 hours. Concentrated virus

was freshly used to transduce target cells.

Cell transduction with lentiviral particles. Human and mouse cells were transduced

overnight with concentrated lentiviral particles (see Virus preparation) in the presence of

8 µg/mL polybrene. After 72 hrs from infection, the culture medium was replaced with a

fresh medium supplemented with antibiotic selection (G418 50 µg/mL or puromycin 1-5

µg/mL). Overexpression, down-regulation, or knockout of the targets were evaluated by

western blot after one week of antibiotic selection. For doxycycline-inducible shRNAs,

after one week of antibiotic selection, cells were pulsed with doxycycline hyclate

(Millipore-Sigma, D5207) 1 µg/mL at different time points, as indicated in figures and gene

expression was evaluated by western blot.

Cell transfection. Cells were seeded at 50% confluency in 10 cm diameter plate in

complete medium and allow to adhere overnight. Transfection mix was prepared by

adding the 10 µg of the indicated plasmids in 600 µL Opti-MEM (Gibco) in presence of 40

µL polyethyleneimine (PEI). Cells were then transfected for 8-12 hrs and protein

expression was evaluated using Western Blot.

Generation of barcoded KPC (N150F) cell line. KPC cells were infected

overnight with CloneTracker™ XP Lentiviral Expressed Barcode Library (Cellecta)

in presence of polybrene (8μg/ml) (MOI < 0.1). After infection and puromycin

selection, cells were passaged for 10 generations before injection into NSG or

C57BL/6 mice.

Cell proliferation and clonogenic assay. Human and mouse cells were

seeded in a 96-well plate (500-1000 cells per well) in medium supplemented with

doxycycline 1 μg/mL or vehicle. After cell adhesion, cells were incubated at at 37°C

5%CO_2 in the IncuCyte (Essenbioscience) incubator. Cell confluency was

measured and analyzed over a period of 5-8 days using and medium was changed

every 48 hrs.

For clonogenic assay, 100-500 cells were seeded in a 6-well plate in

medium supplemented with doxycycline 1 μg/mL or vehicle and maintained at

37°C 5%CO_2 (medium was changed every 48 hrs). After 8-15 days, clones were

fixed and stained with crystal violet (0.25% crystal violet in methanol 20%).

Colonies with more than approximately 50 cells were counted manually and

clonogenic survival fraction was expressed as the relative plating efficiencies of

the irradiated cells to the control cells.

Molecular Biology Experiments:

Short-hairpin RNAs (shRNAs) cloning. For gene silencing, shRNAs from

Millipore-Sigma were cloned into the constitutive pLKO.1-TRC cloning vector

according to the manufacturer's protocol. These same shRNA sequences were

cloned into pENTR-THT and integrated into the inducible Tet-Puro backbones.

Single-guide RNAs (sgRNAs) design and cloning. sgRNAs were designed

with "GenScript CRISPR sgRNA Design Tool" (https://www.genscript.com/gRNA-

design-tool.html?a=post). First, 5`-phosphorilated oligos were annealed and

diluted 1:20. Then 1 μL of each annealed and diluted sgRNA was cloned in

digested lentiCRISPR-V2 according to Dr. Feng Zhang's protocol

(https://media.addgene.org/cms/files/Zhang_lab_LentiCRISPR_library_protocol.p

df). NEB® Stable Competent E. coli (C3040I) colonies resistant to ampicillin

antibiotic selection were amplified, and presence of sgRNA was confirmed by

Sanger sequencing. Positive clones were transfected individually in 293T cells along with vectors for lentiviral packaging production. Human or mouse cells were infected by lentiviral particles carrying a specific sgRNA and selected for puromycin resistance. Cut efficiency of sgRNA was tested by T7 Endonuclease I (NEB #M0302L) assay on the DNA of infected cells, according to the manufacturer's protocol (https://www.neb.com/protocols/2014/08/11/determining-genome-targeting-efficiency-using-t7-endonuclease-i). The sgRNA sequences for knocking out DPY30 are the following: GAGCCAGAGCAGATGCTGGA (sgDPY30#1), GGGACTTGCTGTGCTTGCAA (sgDPY30#2) for human and GAGTCGGAGCAGATGCTGGA (sgDpy30) for mouse.

Plasmids. The following vectors were purchased from Addgene: psPAX2 (#12260), pMG2.D (#12259), lentiCRISPR-V2 (#52961), pLKO.1-TRC (#8453), pENTR-THT (#55790), Tet-Puro (#21915), pLenti-PGK-Neo-PIP-FUCCI (#118616), emEGFP-Mcm2 (mouse, #54163), emEGFP-Mcm3 (mouse, #54165), emEGFP-Mcm4 (mouse, #54167), emEGFP-Mcm5 (mouse, #54169), emEGFP-Mcm6 (mouse, #54171), emEGFP-Mcm7 (mouse, #54173), pcDNA3 Flag-hDPY30 (#15554), pcDNA3 Flag WDR5 (#15552), pCMV-8xHis-Ub (#107392). The following vectors were purchased from Genscript: pcDNA3.1 WDR5_L240K(+)-N-HA (L240K WDR5_OHu17370C), pcDNA3.1 WDR5_V268E(+)-N-HA (V268E WDR5_OHu17370C), pcDNA3.1 WDR5-(+)-N-HA (WDR5_OHu17370C), pcDNA3.1(+)-C-HA-L69D-hDPY30 (L69D DPY30_OHu17211C), pcDNA3.1(+)-C-HA-hDPY30 (DPY30_OHu17211C), pcDNA3.1(+)-C-HA-mdpy30 (Dpy30_OMu07274C), pcDNA3.1(+)-N-6His-hMCM2 (MCM2_OHu09867C), pcDNA3.1(+)-N-6His-hMCM3 (MCM3_OHu107378C), pcDNA3.1(+)-N-6His-hMCM4 (MCM4_OHu11591C), pcDNA3.1(+)-N-6His-hMCM5 (MCM5_OHu02170C), pcDNA3.1(+)-N-6His-hMCM6 (MCM6_OHu20108C), pcDNA3.1(+)-N-6His-hMCM7 (MCM7_OHu28012C). The following vectors were purchased from Sigma-Millipore: shCTR (SHC002), shDPY30#1 (TRCN0000297810), shDPY30#2 (TRCN0000280267), shWDR5#1 (TRCN0000118047), shWDR5#2 (TRCN0000118049), shRBBP5#1 (TRCN0000165777), shRBBP5#2 (TRCN0000165100), shASH2L#1 (TRCN0000019274), shASH2L#2 (TRCN0000019276), shDpy30#1 (TRCN0000034420), shDpy30#2 (TRCN0000034419).

650   *Protein extraction, Immunoprecipitation and Western Blot Analyses.* For

651 cellular protein lysates, cells were scraped on ice using cold Ripa lysis buffer (150

652 nM NaCl, 50 mM Tris HCl pH 8, 1% Igepal, 0.5% sodium deoxycholate, 0.1% SDS

653 - Boston BioProducts, BP-115) supplemented with a HALT protease and

654 phosphatase inhibitor cocktail (ThermoFisher). Cell lysates centrifuged at 13,200

655 rpm for 20 min at 4°C and supernatants were collected.

656   For immunoprecipitates (IPs), cells were scraped using NP40 buffer (10 mM

657 NaCl, 10 mM Tris HCl pH 7.5, 0.5% Nonidet P-40 - Boston BioProducts, BP-119)

658 supplemented with a HALT protease and phosphatase inhibitor cocktail (#78440,

659 ThermoFisher Scientific). Cell lysates were sonicated twice at 30% amplitude for

660 15 sec on ice and centrifuged at 13,200 rpm for 20 min at 4°C and supernatants

661 were collected. Immunoprecipitation was performed by binding the primary

662 antibody to protein G magnetic beads (Dynabeads, ThermoFisher Scientific,

663 10004D) for 30-60 min at RT. Bs3 (A39266, ThermoFisher Scientific) was used to

664 crosslink primary antibodies to protein G. Cell lysates were diluted with PBS-T

665 buffer (PBS-0.1% Tween20) supplemented with protease/phosphatase inhibitor,

666 added to antibody-bead complexes and incubated overnight at 4°C. IPs were then

667 washed in PBS-T buffer, resuspended in 4X NuPage Sample Buffer

668 (Invitrogen) supplemented with 1% β-mercaptoethanol.

669   IPs and proteins were separated in 4-20% SDS-PAGE (Criterion Precast

670 Midi Gel, Bio-Rad) and transferred to nitrocellulose membranes (Trans-Blot Turbo

671 Midi 0.2 μm nitrocellulose transfer pack, Bio-Rad). Membranes were blocked with

672 5% non-fat dried milk in PBS-T and incubated at 4°C overnight with primary

673 antibodies.

674   Membranes were washed in PBS-T and incubated 1 hour at RT with the

675 appropriate horseradish peroxidase-conjugated secondary antibodies (Cell

676 Signaling Technology) for ECL detection (SuperSignal™ WEST Pico PLUS

677 Chemiluminescent Substrate, ThermoFisher) or with the appropriate IRDye

678 800CW or IRDye 680RD Dye-Labeled secondary antibodies (LICORbio). Band

679 quantification was performed using the QuantiONE software (Bio-Rad

Laboratories). The Re-Blot Plus Strong Solution (#2404, EMD Millipore) was used to strip

the membranes, when reblotting was needed.

Purification of proteins on newly synthesized DNA (iPOND). iPOND was performed

essentially as described previously (86). Briefly, 293T or KPC cells were seeded in a

square at 70% confluency (one square per each iPOND sample). Cells were pulsed with

10 μM EdU (5-Ethynyl-2'-deoxyuridine, Millipore-Sigma, #900584), for 5 and 10 min to

enable the detection of protein bound to active DNA replication forks. For each

experiment, two negative controls were included: one sample was not pulsed with EdU

and one sample was pulsed with EdU followed by 60 min thymidine chase. All samples

were fixed with 1% formaldehyde (ThermoFisher Scientific) for 20 min at RT, and

quenched using 125 mM Glycine (Millipore-Sigma) solution for 5 min at RT. After fixation,

samples were collected in 50-ml tube and washed (once with PBS+1% BSA and once

with PBS). Samples were permeabilized using PBS+0.1% Tryton-X100 for 10 min in

gentle shaking at RT and transferred in a 15-ml tube. Copper(ii) Sulfate (Millipore, Sigma

#451657), (+)-sodium L-ascorbate (Millipore, Sigma #A7631) and Biotin Azide (Click

chemistry tools, Fisher) were used to perform the Click-reaction for biotin-conjugate the

incorporated EdU. After washing steps, cell pellets were resuspended in SDS 1% for

protein extraction and sonicated 3 times at 30% amplitude for 30 sec on ice. After

centrifugation, cleared lysates were diluted 1:2 with PBS and % 1% of protein lysate was

kept as input control. Cleared lysates were then incubated with Pierce™ Streptavidin

Magnetic Beads (ThermoFisher Scientific, 88817) for 2 hr at RT in gentle shaking. Protein

complexes were washed three time with PBS-T and purified proteins were resuspended

in 4X NuPage Sample Buffer (Invitrogen) supplemented with 1% β-mercaptoethanol.

In vitro immunofluorescent staining. Cells were plated at 50% of confluency over

a round glass coverslip (Chemglass Life Sciences, cat. n. CLS1763012), allowed to

adhere overnight, treated with doxycycline as appropriate to induce gene silencing and

fixed using 4% paraformaldehyde for 15 min at RT. Cells were then washed twice in PBS

and permeabilized (0.1% triton X100 in PBS) for 15 min at RT. After a second round of

washes in PBS, cells were blocked using blocking buffer (3% BSA, 3% goat serum, 0.01%

triton X100 in PBS) 1h at RT. Primary antibodies were diluted in blocking buffer and

incubated overnight at 4°C. After washes, secondary antibodies (conjugated with Alexa-

488, Alexa-594, Alexa-647 or Alexa-750) were diluted in blocking buffer and incubated 1h at RT. DAPI was used to counterstain nuclei. Images were captured with a Hamamatsu C11440 digital camera, using a wide-field Nikon Ni-E fluorescent microscope.

Proximity Ligation Assay (PLA) and SIRF assay. For PLA, Duolink® In Situ Detection Reagents Red (Millipore-Sigma) were used and cells were stained according to the manufacturer protocol. Briefly, cells were plated, fixed and blocked as described above. After primary antibody incubation, cells were washed with Buffer A and then the Duolink® In Situ PLA® probe anti-rabbit or anti-mouse were added as appropriate. After ligation and amplification reactions, cells were imaged using Zeiss LSM880 Airyscan confocal microscope (63X objective) at the Advanced Microscopy Core (MD Anderson Cancer Center). SIRF assay was performed essentially as described in (52, 53) with minor modifications. Cells were plated on coverslips as described above and were pulsed with 20 µM BrdU for 5 min. The primary antibodies used were anti-WDR5 (rabbit), anti-DPY30 (rabbit, Bethyl), anti-PCNA (rabbit) and anti-BrdU (mouse). Click-IT reaction was not performed.

DNA Fiber Assay. Fiber stretching was performed essentially as described in (87). Briefly, cells were double-pulsed precisely for 20 min with 25 µM CldU (Chloro-2'-deoxyuridine (Millipore-Sigma, C6891) followed by 250 µM IdU (5-Iodio-2'-deoxyuridine, Millipore-Sigma, I7125). Cells were harvested and diluted at $2x10^5$cells/ml. A 3-µL drop of cell suspension was allowed to air-dry in a glass slide and fiber spreading was carried out by adding 10 µL spreading buffer (0.5% SDS in Tris-HCl EDTA). Fibers were fixed in Methanol/Acetic Acid 3:1 10 min, allowed to completely dried and rinsed with deionized water. DNA was denatured with HCl 2N for 1.5 hr ar RT. After washing steps, slides were blocked in blocking buffer (PBS 2% BSA, 2% goat serum, 0.1% Tween20) 1 hr at RT. CldU and IdU were recognized by using the following primary antibodies: rat monoclonal anti-BrdU clone BU1/75 and mouse monoclonal anti-BrdU clone B44. The antibodies were diluted in blocking buffer and incubated 4 hr at RT, followed by secondary antibody (goat anti-rat IgG (H+L) Alexa Fluor® 488, goat anti-mouse IgG (H+L) Alexa

Fluor® 594, ThermoFisher Scientific). Imaging was performed on the Leica SP8 laser scanning confocal microscope at the Advanced Microscopy Core (MD Anderson Cancer Center).

Chromosome mis-segregation. For the *in vitro* analysis, cells were plated at 50% of confluency over a round glass coverslip (Chemglass Life Sciences, cat. n. CLS1763012), allowed to adhere overnight and fixed using 4% paraformaldehyde for 10 min at RT. DNA was stained using DAPI 10 min at RT. Chromosome mis-segregation were evaluated using a wide-field Nikon Ni-E fluorescent microscope (40X objective). For *in vivo* analysis, H&E samples from human or mouse-derived tumors grown in recipient mice were evaluated for anaphases with chromosome mis-segregation events using a wide-field Nikon Ci-L plus light microscope (100X objective). Analyses was performed by evaluating chromosome mis-segregation in H&E samples (5 mice per group, one section per mouse), and for each mouse at least 20 anaphases were counted.

Preparation of Metaphase Spreads. Analysis of chromosomal aberrations was performed at the Molecular Cytogenetics Facility (MD Anderson Cancer Center) as previously described (32). Cells were incubated with Colcemid (KaryoMAX™, ThermoFisher Scientific, #15212012) at 0.04 µg/mL for 3 hours, trypsinized, washed, and spun down. Cells were incubated at RT for 15 minutes in hypotonic solution (0.075M KCl). Cells were fixed by adding three times 2 mL of fixative solution (glacial MeOH + acetic acid 3:1), followed by centrifugation. Cell suspension was dropped on glass slide. Slides were allowed to air-dry, stained with 4% Giemsa staining solution (KaryoMAX™, ThermoFisher Scientific, #10092013) for 5 min and mounted. At least 35 metaphases were analyzed from each sample for chromosomal aberrations. Images were captured using a Nikon 80i microscope equipped with karyotyping software from Applied Spectral Imaging, Inc.

Cell cycle synchronization and analysis by flowcytometry. Human and mouse cells were seeded in a 6-cm diameter plate (1×10^5 cells per plate) in complete medium and allowed to adhere to plate for 16 hr. To synchronize cells in early S phase, a double thymidine (Millipore-Sigma, T9250) pulse was used. Briefly, cells were cultured in presence of 2 mM thymidine for 18 hrs, released in complete medium for 8 hrs and arrested in S phase with a second thymidine-pulse for 18 hrs. Cells were then released

at different time point as indicated in figure using complete medium. To
synchronize cells in G2/M phase, cells were cultured in presence of 2 mM
thymidine for 18 hrs, released in complete medium for 8 hrs and arrested in G2/M
phase with 100 ng/ml Nocodazole (Millipore-Sigma, M1404) for 16 hrs. For DNA
content analyses cells were harvested at specific time point, resuspended in 250
μL of ice-cold PBS and fixed by adding 750 μL of absolute ethanol dropwise while
vortexing. Cells were kept an overnight in fixative solution, washed twice with PBS
and DNA stained using DAPI. Data were acquired using BD Celesta analyzer and
analyzed by FlowJo.

BrdU incorporation assay and Re-replication analyses using flow cytometry.
Human and mouse cells were seeded in a 6-cm diameter plate (1×10^5 cells per
plate) in complete medium and allowed to adhere to the plate for 16 hr. Cells were
pulsed with 15 μM BrdU for 2 hrs, harvested and fixed using APC or FITC BrdU kit
(#552598 or #559619 BD bioscienses) following manufacturer protocol.
Alternatively, human PDAC cells were fixed with ethanol as described above and
DNA was denatured using 2N HCl solution for 20 min. Cells were then washed
three time with PBS, blocked in PBS 5% BSA and PE-conjugated anti-BrdU
antibody was incubated 1 hr at RT. After washing with PBS, nuclei were
counterstained using DAPI. Data were acquired using BD Celesta analyzer and
analyzed by FlowJo.

Cell sorting. Human cells were seeded in a 15-cm diameter plate at 70%
confluency in complete medium and allowed to adhere to the plate for 24 hr. Cells
were harvested and resuspended in FACS buffer (3% FBS, 1mM EDTA) and DAPI
was used as live/dead reagent. Cells were sorted using Influx Silver sorter and
data were analyzed by FlowJo.

Immunophenotyping using flow cytometry. After tumor dissociation and
single cells extraction (see Cell isolation from tumors), staining was performed
essentially as described in (88). Briefly, cell surface antigens were stained with the
appropriate primary antibodies for 20 minutes at 4°. For the staining of intracellular
proteins, samples were washed, fixed, permeabilized for 30 min at RT and
subsequently incubated with the appropriate primary antibodies for 1 hr at RT.

Dead cells were stained using the fixable viability violet dye Zombie Red (Invitrogen) for

10 min at RT followed by the blocking of Fc receptors with TruStain fcX (Biolegend) for

15 min at 4°C. Cells were analyzed on LSRFortessa X-20 flow cytometers (BD

Biosciences) and data were analyzed with FlowJo software version 10.8.2.

Ex vivo stimulated cytokine secretion. Single-cell suspensions from tumors were

resuspended in RPMI 1640 with 10% FCS, were added to anti-CD3 (clone 145-

2C11)/anti-CD28 (clone 37.51) antibody-coated (overnight at 10 µg/ml antibody) culture

plates for 4hr at 37°C in the presence of 1 µg/ml Golgiplug (BD Bioscience) and Monensin

(Biolegend) and processed for TNF and IFNγ intracellular cytokine staining.

Histopathology, immunohistochemistry, and immunofluorescence. Tumor

specimens were fixed overnight in 4% buffered paraformaldehyde, transferred to 70%

ethanol, and then embedded in paraffin using Leica ASP300S processor. For

histopathological analysis, tumors were sectioned (Leica RM2235, 5µm thickness) and

serial slides were collected. For every series one section was stained with hematoxylin

and eosin and remaining sections were kept for either immunofluorescence or

immunohistochemical analysis. For Alcian Blue (Vector Laboratories, H3501) and

Masson's Trichrome staining (VitroView™ Masson's Trichrome Stain Kit, VB-3016),

samples were processed according to the manufacturers' protocols.

Histological samples were processed as previously described (74). Briefly, after

cutting, baking, and deparaffinization, sections underwent antigen retrieval using Citra-

Plus Solution (BioGenex) according to specifications.

For immunohistochemistry (IHC) staining, endogenous peroxidases were

inactivated by 3% hydrogen peroxide and non-specific signals were blocked using 5%

BSA, 5% goat serum, and 0.1% Triton X-100. Primary antibodies were applied and

incubated overnight at 4°C. ImmPRESS HRP IgGs were used as secondary antibodies

and ImmPACT NovaRED® (Vector Laboratories, # SK-4805) was used for detection.

Images were captured with a Nikon DS-Fi1 digital camera using a wide-field Nikon Ci-L

plus light microscope.

Tissue Microarray slides of PDAC samples with available survival data (n=128)

collected at MD Anderson Cancer Center were stained for DPY30 as described above.

DPY30 nuclear expression was evaluate and categorized in blind as follow: Low (score

0=negative or score 1=weak), medium (score 2=moderate) and high (score

3=strong) expression. Kaplan-Meier survival curve was generated by clustering

PDAC patients according to DPY30 expression and log-rank test was used to

compare the survival difference among different groups.

For immunofluorescence staining, secondary antibodies conjugated with

Alexa-488, Alexa-594, Alexa-647 or Alexa-750 were used. DAPI nuclear staining

(#40043, Biotium) was used to counterstain nuclei. Images were captured with a

Hamamatsu C11440 digital camera, using a wide-field Nikon Ni-E fluorescent

microscope or Vectra Polaris (Akoya Biosciences).

Cyclic immunofluorescence. After the first staining and image acquisition

using PE Vectra polaris, coverslips were removed by incubating slides with 37°C

PBS for 20 min. Slides were then incubated with stripping buffer (62.5 mM Tris

Base / Tris-HCl, 2% Sodium Dodecyl Sulfate, 0.8% Beta ME, 55°C, 30 min), and

then stained with another round of antibodies. Slides were washed with PBS 0.1%

Triton-X100 between each step. DAPI nuclear staining (#40043, Biotium) was

used every cycle.

Antibodies. Primary antibodies purchased from Cell Signaling Technology:

anti-WDR5 (clone D9E1I, #13105), anti-Cleaved Caspase-3 (Asp175, #9661),

anti-STING (clone D2P2F, #13647), anti-STING (S365) (clone D1C4T, #62912),

anti-ASH2L (clone D93F6, #5019), anti-RbBP5 (#8037), anti-MCM2 (S139) (clone

D1Z8X, #12958), anti-MCM2 (clone D7G11, #3619), anti-MCM3 (#4012), anti-

MCM4 (clone D3H6N, #12973), Rabbit monoclonal anti-MCM7 (clone D10A11,

#3735), anti-Histone H3 (pS10) (#9701), anti-Cdc6 (clone C42F7, #3387), anti-

H3K4Me3 (clone C42D8, #9751), anti-His Tag (clone 27E8, #2366). Primary

antibodies purchased from Millipore-Sigma: anti-DPY30 (#HPA043761), anti-

γH2AX (pS139) (clone JBW301, #05-636), anti-beta-Actin (clone AC-74, #A5316).

Primary antibodies purchased from Bethyl Laboratories: anti-DPY30 (#A304-

296A), anti-MCM5 (#A300-195A), anti-MCM6 (#A300-194A), anti-WDHD1

(#A301-140A), anti-RPA2 (pS4-8) (#A300-245A). Primary antibodies purchased

from Santa Cruz Biotechnology: anti-WDR5 (clone G-9, #sc-393080), anti-Cyclin

E (clone HE12, #sc-247), anti-Histone H3 (clone 1G1, #sc-517576), anti-Geminin

(clone FL-209, #sc-13015), anti-Luciferase (clone C-12, #sc-74548). Primary antibodies

purchased from Abcam: anti-BrdU (clone BU1/75 – ICR1, #ab6326), anti-H3K4Me1

(#ab8895), anti-H3K4Me3 (#ab8580), anti-H3K4Ac (#ab4729), Alexa Fluor® 594 anti-

αSMA (clone 1A4, #ab202368), Alexa Fluor® 488 anti-αSMA (clone 1A4, #ab184675).

Primary antibodies purchased from ThermoFisher Scientific: anti-PCNA (#PA5-16797),

anti-Cyclin A2 (clone E23.1, #MA1-154), anti-GFP (clone GF28R, #MA5-15256), anti-

Ki67 (#MA5-14520), anti-HA tag (clone 2-2.2.14, #26183), anti-CD8a (clone 4SM15, #14-

0808-82), PE-conjugated anti-BrdU (clone BU20A, #12-5071-42). Other primary

antibodies: anti-BrdU (clone B44, #347580, BD Laboratories), anti-Cdc6 (pS54) (#102-

15286, RayBiotech), anti-Ki67 (#14965-1-AP, Proteintech), anti-Flag tag (clone OTI4C5,

#TA50011-100, OriGene Technologies), anti-CD3 (#A045201-2, Agilent Technology),

Rabbit gamma globulin (#31887, Invitrogen). Antibodies purchased from Biolegend: anti-

mouse Ki67 (clone 16A8, #652402), BV 510 anti-mouse CD8a (clone 53-6.7, #100751),

PE anti-CD279 (PD-1) (clone RMP1-30, #109104), PB anti-Granzyme B (clone GB11,

#515408), PE-Cy5 anti-CD3 (clone 17A2, #100274), Pe-Cy-7 anti-Foxp3 (clone FJK-16S,

#25-5773-82), APC-Cy7 anti-CD4 (clone GK1.5, #100414), Alexa Fluor® 700 anti-CD45

(clone 30-F11, #103128), PerCP-Cy5.5 anti-Thy1.2 30-H12 (clone 30.H12, #105338),

BV785 anti-NK1.1 (clone PK136, #108749), Alexa Fluor® 647 anti-Tnf (clone MP6-XT22,

#506314), Alexa Fluor® 700 anti-Infγ (clone XMGG1.2, #505824). Secondary antibodies

purchased from Vector Laboratories: ImmPRESS® HRP goat anti-rabbit IgG (#MP-

7451), ImmPRESS® HRP goat anti-mouse IgG (#MP-7451). Secondary antibodies

purchased from ThermoFisher Scientific: Goat anti-rabbit IgG (H+L) Alexa Fluor® 488

(#A-11008), Goat anti-rabbit IgG (H+L) Alexa Fluor® 594 (#A-11012), Goat anti-rabbit

IgG (H+L) Alexa Fluor® 647 (#A-21244), Goat anti-rabbit IgG (H+L) Alexa Fluor® 750

(#A-21039), Goat anti-mouse IgG (H+L) Alexa Fluor® 488 (#A-11001), Goat anti-mouse

IgG (H+L) Alexa Fluor® 594 (#A-11005), Goat anti-mouse IgG (H+L) Alexa Fluor®

647(#A-21235), Goat anti-rat IgG (H+L) Alexa Fluor® 488 (#A-11006), Goat anti-rat IgG

(H+L) Alexa Fluor® 647 (#-21247).

RNA extraction. RNA extraction was achieved using Qiagen RNeasy Micro kit

following manufacturer's instruction. Briefly, RNA extraction was performed from KPC and

KP$^{-/-}$C cells or tumors stably knocked out for Dpy30 or control and from PATC53 stably

expressing doxycycline-inducible scramble control (shCTR) or shRNAs targeting
WDR5 or DPY30 pulsed for 48 hrs with doxycycline. Cells grown *in vitro* were
scraped using 350 µL of RLT buffer supplemented with β-mercaptoethanol. For
RNA extraction from tumors, tissue specimens (approximately 2x2mm) were snap-
frozen and kept at -80°C until further processing. Tumors were then weighted and
dissociated in RLT buffer supplemented with β-mercaptoethanol using
gentleMAX™ M tubes (Miltenyi Biotec) and RNA_02 program. DNA digestion was
performed by adding recombinant DNAse I (Millipore-Sigma, #4536282001). to
purification column. Total RNA was eluted in RNAse-free water, quantified, and
submitted for sequencing.

Genomic DNA extraction. DNA extraction was achieved using Qiagen
DNaeasy Blood and Tissue kit following manufacturer's instruction. Briefly, cells
grown *in vitro* were harvested and cell pellets were snap-frozen and kept at -80°C.
For DNA extraction from tumors, tissue specimens (approximately 2x2mm) were
weighted and dissociated by mechanical digestion in 180 µL Buffer ATL
supplemented with 20 µL proteinase K stock solution. Total DNA was eluted in
DNAse-free water, quantified, and submitted for sequencing.

Chromatin immunoprecipitation (ChIP). ChIP was performed essentially as
described in (89) with optimized DNA shearing condition and few modifications.
Briefly, $9x10^6$ PATC53 cells stably expressing doxycycline-inducible shRNAs
against DPY30, WDR5 or scramble control were seeded in a 15-cm diameter plate
in complete medium in the presence or not of 1 µg/ml doxycycline for 48 hrs. Cells
were then harvested, resuspended in 3 mL of complete medium, cross-linked with
200µL of 16% formaldehyde for 10 min at 37°C in gentle stirring and quenched
with 200 µL Glycine 2M for 5 min at 37°C in gentle stirring. Tissue lysis, sonication
and chromatin immunoprecipitation were performed as described in (31, 74).

ATAC-seq. Cells were processed as described in (74). Briefly, 100,000
human PATC53 cells were lysed in lysis buffer for 5 minutes on ice. Tn5
tagmentation was carried out for 30 minutes at 37 °C. After indexing and PCR
amplification, DNA libraries were multiplexed and sequenced on HiSeq X with
150nt PE configuration.

Immuno-precipitate mass spectrometry. KPC, PATC53 and PATC66 cells were
seeded in 15cm diameter plate at 50% confluency in complete medium. Cells were
scraped in RIPA buffer as described above after 24 hr from cell seeding. For each sample,
2 mg of lysate were immunoprecipitated for DPY30 or control IgG and immunocomplexes
were resuspend in 100μL of NuPage sample buffer, and 20μL were resolved using SDS-
PAGE for quality control. For proteomics, 40μL of each sample were processed by SDS-
PAGE using 10% Bis-Tris NuPage Mini-gel (Invitrogen) with the MES buffer system. The
gel was run 1cm and the mobility region excised into 10 equally sized bands. Each band
was processed by in-gel digestion with trypsin using a robot (ProGest, DigiLab) with the
following protocol: i) washed with 25mM ammonium bicarbonate followed by acetonitrile,
ii) reduced with 10mM dithiothreitol at 60°C followed by alkylation with 50mM
iodoacetamide at RT, iii) digested with sequencing grade trypsin (Promega) at 37°C for
4h, iv) quenched with formic acid and the supernatant was analyzed directly without
further processing. For mass spectrometry, half of each digested sample was analyzed
by nano LC-MS/MS with a Waters M-Class HPLC system interfaced to a ThermoFisher
Fusion Lumos mass spectrometer. Peptides were loaded on a trapping column and eluted
over a 75μm analytical column at 350nL/min; both columns were packed with Luna C18
resin (Phenomenex). The mass spectrometer was operated in data-dependent mode,
with the Orbitrap operating at 60,000 FWHM and 15,000 FWHM for MS and MS/MS
respectively. The instrument was run with a 3s cycle for MS and MS/MS. 5hrs of
instrument time was used for the analysis of each sample.

Quantitative real time PCR. RNA from cells was extracted using RNeasy Mini kit
(Qiagen, #74104). Total RNA was quantified using the Nanodrop Spectrophotometer
(ThermoFisher). The synthesis of cDNA was performed using the SuperScript VILO
Master Mix (ThermoFisher) in an PCR Thermocycler (Bio-Rad). cDNAs were amplified
using PowerUp™ SYBR™ Green Master Mix (Invitrogen, A25742) and Real-time PCR
Detection System (Applied Biosystems). Normalization of the data was performed using
housekeeping genes (*ACTB, GAPDH, 18S*). The following primers (Sigma) were used (5'
– 3'): RFC3 Forward TTCGTGGAAGGCTGTATGAGC, RFC3 Reverse
CCTCCCCTTTCAGTTGTCCAT; *GINS2* Forward CAATGCCCAGCCCTTACTACA,
GINS2 Reverse GCCTTCGGGATGTTGTCTGA; *FEN1* Forward

GTTCCTGATTGCTGTTCGCC, *FEN1* Reverse ATGCGAATGGTGCGGTAGAA;

IFNL1   Forward   GGAGGCATCTGTCACCTTCAA,   *IFNL1*   Reverse

GGACTCAGGGTGGGTTGAC; *IRF-AS1* Forward CACCGCGTCTGTGTCTGTA,

IRF-AS1   Reverse   GCCACCTACGATATACCACCC;   *TRIM10*   Forward

TGGCTAACGTGGTGGAGAAC,   *TRIM10*   *Reverse*

TGCTCTTGGCAGACATCCTC; ACTB Forward CCAGAGGCGTACAGGGATAG,

ACTB   Reverse   CCAACCGCGAGAAGATGA;   *18S*   Forward

CTCAACACGGGAAACCTCAC, *18S* Reverse CGCTCCACCAACTAAGAACG;

GAPDH   Forward   GAAGGTGAAGGTCGGAGTC,   *GAPDH*   Reverse

GAAGATGGTGATGGGATTTC.

NGS library production for barcode detection in bulk tumors. Frozen tumors

were minced and suspended in Buffer P1 (Qiagen, 1 mL buffer per 100 mg tumor)

and a scale for cell number quantification was added. This mixture was

homogenized in a gentleMACS homogenzier (Miltenyi Biotec). After

homogenization, the samples were transferred into 15 mL polypropylene tubes

(Falcon) and RNase A (100 µg/mL) was added (10 µL per 100 mg tumor). After 10

minutes, 10% SDS (Promega) and Proteinase K (Qiagen) were added (both 50 µL

per 100 mg tumor) and incubated at 56°C for 20 minutes. The DNA was then

sheared by passing the lysate through a 23G syringe needle 10 times. The lysates

were transferred into 1 mL polypropylene tubes and DNA was purified using

Phenol:Chloroform:Isoamyl alcohol (25:14:1 pH 8.0, Sigma Aldrich) and

Chloroform:Isoamyl alcohol (24:1, Sigma Aldrich). The aqueous DNA was

precipitated by adding 3M sodium acetate (Sigma Aldrich, 90uL per 1 mL sample)

and isopropanol (Fisher Scientific, 720 µL per 1 mL sample). DNA was centrifuged

at 14,000 rpm for 20 minutes at 4°C and washed with 70% ethanol (Fisher

Scientific). Once dried, the DNA pellets were dissolved in Ultrapure distilled water

(Thermo Fisher Scientific) and DNA concentration was quantified using NanoDrop

2000 (Thermo Fisher Scientific).

Barcodes were amplified through 2 rounds of PCR on the DNA samples

using Titanium Taq DNA polymerase (Clontech, Takara). The first PCR was

performed using the following protocol: 3 minutes at 94°C followed by 16 cycles of

30 seconds 94°C, 10 seconds 60°C, and 20 seconds 68°C, followed by final extension at

72°C for 2 minutes then hold at 4°C. The sequences of the primers used are:

XPg_1stF (5'-ACCGAACGCAACGCACGCA-3')

XPg_1stR (5'-ACGACCACGACCGACCCGAACCACGA-3').

A second PCR was performed on the PCR product using the following protocol: 3

minutes at 94°C followed by 12 cycles of 30 seconds 94°C, 10 seconds 66°C, and 10

seconds 72°C, followed by final extension at 72°C for 2 minutes then hold at 4°C. The

sequences of the primers used are:

P7_XPg_2ndF   (5'-

AGCAGAAGACGGCATACGAGATAGCACCGAACGCAACGCACGCA -3')

unique   index   reverse   primer   P5_XPg_2ndR   (5'-

AGATACGGCGACCACCGAGATCTACACGCACGACGAGACGCAGACGAAXXXXXXA

CGACGACCGACCCGAACCACGA-3', X bases refers to index barcode sequence).

Second PCR products were then identified through agarose gel electrophoresis

(2.5%, Lonza) at expected size of 227 bp and extracted using PureLink Quick Gel

Extraction Kit (Thermo Fisher Scientific). The purified PCR product was quantified using

High Sensitivity D1000 ScreenTape and Agilent 4200 TapeStation System (Agilent

Technologies). Barcode representation was measured by NGS using Illumina HiSeq2000

with Seq_XPg_BC30 (5'- AGACGACCTGCTCCAGCTGCACCA-3') as read 1

sequencing primer and RSeq-IND-XP NGS (5'- ACACGCACGACGAGACGCAGACGAA

-3') as i5 index primer.

Quantitative Scale. A known quantity of barcoded cells was used as normalization

scale as previously described (67). In brief, the barcoded cells are infected with the same

lentivirus that carries a unique barcode that are not shared in the lineage tracing library

with a low Multiplicity of Infection (MOI < 0.1). Resulting infected cells are then selected

by puromycin and expanded for preparing the spike-in mixture with following cells:

50   cells   carrying   the   bc14-001.bc30-100002   barcode

(CCAAAGATAGATCATGGTACACCATGACACGTCACATGTGTGCAGTCA);

500   cells   carrying   the   bc14-001.bc30-100003   barcode

(CCAAAGATAGATCATGGTACTGACACTGCAGTACCACATGCACACAAC);

5,000   cells   carrying   the   bc14-001.bc30-100001   barcode
(CCAAAGATAGATCATGGTCATGGTACACACGTGTCAACTGACCACAAC);

50,000   cells   carrying   the   bc14-001.bc30-100004   barcode
(CCAAAGATAGATCATGGTTGACACACACTGACTGACTGACTGACACCA);

500,000   cells   carrying   the   bc14-001.bc30-100005   barcode
(CCAAAGATAGATCATGGTGTACACGTACGTACTGACCAACTGACGTCA).

During the tissue and cell dissociation step of barcode library preparation,

the cell mixtures were added and served as a reference for spike-in normalization

of the sequencing reads. After normalizing read counts of each library by the size

factor (DNA content) and library size, a high correlation coefficient (r=0.96) was

observed between normalized read counts and cell number in log scale. This

calibration curve was then used to predict cell count of each barcode in tumors.

***In vivo* Experiments**

Mouse strains. Conditional *Ptf1a*Cre;*LSL-Trp53$^{fl/+}$* and *LSL-Kras$^{G12D/+}$*

strains were interbred to obtain *Ptf1a*Cre;*LSL-Kras$^{G12D/+}$;LSL-Trp53$^{fl/fl}$;*double

mutant animals (hereafter KP$^{-/-}$C mouse model) and kept in a C57BL/6 background

(47). Conditional *Ptf1a*Cre; *LSL-Kras$^{G12D/+}$* (obtained from Dr. Haoquang Ying,

UTMDACC) and (46) *LSL-Trp53$^{R172H/+}$* (obtained by Dr. Guillermina Lozano,

UTMDACC) strains were interbred to obtain *Ptf1a*Cre;*LSL-Kras$^{G12D/+}$;LSL-*

*Trp53$^{R172H/+}$;*double mutant animals (hereafter KPC mouse model) and kept in pure

C57BL/6 background. Nude, NSG and C57BL/6 female mice (6 weeks old) were

purchased from Experimental Radiation Oncology at MD Anderson Cancer

Center. For treatment with anti-CD8 and anti-PD1 monoclonal antibodies,

C57BL/6J female mice (6 weeks old) were purchased from Jackson Laboratory.

All animal studies and procedures were approved by the UTMDACC Institutional

Animal Care and Use Committee. All experiments conformed to the relevant

regulatory standards and were overseen by the institutional review board. No sex

bias was introduced during the generation of experimental cohorts. Littermates of

the same sex were assigned randomly to experimental arms.

Pancreas orthotopic injections and surgical procedures. C57BL/6 female

mice (8 weeks old) were shaved and anesthetized using isoflurane (Henry Schein

Animal Health). Analgesia was achieved with buprenorphine SR (0.1 mg/Kg BID) (Par

Parmaceutical) *via* subcutaneous injection, and shaved skin was disinfected with

Chlorhexidine and 70% ethanol. A 0.5-cm incision was performed on the left flank through

the skin/subcutaneous and muscular/peritoneal layers. Pancreas was exposed, and 20

µL of cell resuspension (total 1×10^5 KPC or KP$^{-/-}$C cells per mouse) were injected using a

27 Gauge Hamilton syringe. Hemostasis was controlled with a bipolar cautery (Bioseb) if

needed. The pancreas was carefully repositioned into the abdominal cavity, and

muscular/peritoneal planes were sutured individually by absorbable sutures. The

skin/subcutaneous planes were closed using metal clips. Mice were monitored daily for

the first three days, and twice/week thereafter for signs of tumor growth by manual

palpation or MRI when appropriate.

Subcutaneous transplantation, tumor volume measurement and survival study.

Tumor single cell suspension was resuspended in DMEM without phenol Red (Sigma)

and Matrigel (BD Biosciences, 356231) (1:1 dilution) at a density of 1×10^5 cells/100µL for

KPC and KP-/-C cells and 1×10^6 cells/100µL for PATC53 and PATC66 cells. Cell

suspensions were injected subcutaneously into the flank of 8-week-old NSG, nude or

C57BL/6 female mice as indicated in figures. Mice were monitored every 5 days, and

tumor measurements was calculated according to the formula: Tumor volume (mm^3) =

(W^{2}xL)/2, where W is Width and L is length.

Mice that underwent orthotopic cell transplantation were monitored over time for

tumor development by magnetic resonance imaging and survival annotated.

Treatments. The following antibodies were purchased from Bio X cell: InVivoMab

anti-mouse PD-1 (CD279) (clone 29F.1A12, # BE0273), InVivoMab anti-mouse CD8

(clone YTS 169.2, # BP0117) and InVivoMab rat IgG2a isotype control (clone 2A3, #

BE0089). Treatments with monoclonal antibody against PD-1 or CD8 or control were

started 7 days after KPC cell subcutaneous injection in C57BL/6 mice. Monoclonal

antibodies were diluted in PBS to a final concentration of 200µg/100µL and 50µg/100µL

respectively for anti-PD-1 and anti-CD8. The treatments were performed by

intraperitoneal injection of 100µL of antibody per 25g mouse, three times a week for two

consecutive weeks.

Euthanasia, necropsy, and tissues collection. Mice were euthanized by
exposure to CO_2 followed by cervical dislocation. A necropsy form was filled in with
mouse information, tumor size and weight, infiltrated organ annotations, and
metastasis number and location.

Non-invasive MRI imaging. A 7T Bruker Biospec (BrukerBioSpin), equipped
with 35mm inner diameter volume coil and 12cm inner-diameter gradients, was
used for MRI imaging. A fast acquisition with relaxation enhancement sequence
with 2,000/39 ms TR/TE, 256x192 matrix size, r156uM resolution, 0.75mm slice
thickness, 0.25mm slice gap, 40x30cm FOV, 101kHz bandwidth, and 4 NEX was
used for acquired in coronal and axial geometries a multi-slice T2-weighted
images. To reduce respiratory motion, the axial scan sequences were respiratory
gated. All animal imaging, preparation, and maintenance was carried out in
accordance with MD Anderson's Institutional Animal Care and Use Committee
policies and procedures.

Cell Isolation from tumors. Tumors from mouse xenografts were harvested
in HBSS (Gibco) and kept on ice. Isolation of pancreatic tumor cells was performed
by mechanical dissociation (mincing the tumor tissue with sterile scissors) and
enzymatic digestion 1 hr at 37°C in in gentle rotation (Dispase II 1 mg/ml,
Collagenase IV 2 mg/ml in DMEM high glucose supplemented with 1% FBS). The
single-cell populations were seeded at in DMEM high glucose (Gibco)
supplemented with 10% FBS (Gibco), 100 IU/ml penicillin (Gibco), and 100 μg/ml
streptomycin (Gibco). In order to get rid of the murine fibroblasts in the culture, we
performed brief trypsinization cycle (0.25% Trypsin-EDTA, Gibco). For single-cell
RNA and flow-cytometry analyses, tumors from KPC cells were minced into small
fragments and digested with 1.5 mg/mL collagenase IV and 50 U/mL DNase I for
30min at 37°C under agitation. For flow cytometry analysis, the lymph nodes were
collected and passed through 40 μm cell strainers and used for compensation.

***In Silico* Analyses**

Analysis of the TCGA dataset. For TCGA GSEA analysis, the TCGA
pancreatic adenocarcinoma mRNA dataset is downloaded from Broad GDAC
website (http://firebrowse.org/?cohort=PAAD&download_dialog=true). The mRNA

data from the Illumina Genome Analyzer platform in the website was used here

(PAAD.rnaseqv2__illuminahiseq_rnaseqv2__unc_edu__Level_3__RSEM_genes

_normalized__data.data.txt). There was a total of 179 tumor samples and 4 normal

adjacent tissues with their mRNA available. The clinical survival profile of 179 samples

was downloaded from cbio-portal website (http://www.cbioportal.org). The tumor purity

of 179 TCGA RNA samples was computed by the ABSOLUTE developed by Broad

Institute. Only those sample with purity above 50 percent were selected. These 70 high-

purity tumor samples were classified into high and low abundant groups for survival

analysis through applying median value as the cut off. Survival was analyzed according

to the Kaplan-Meier method, and differences in their distribution were evaluated by

means of the log-rank test.

Analysis of the PRINCE dataset (70). Median DPY30 mRNA expression from

metastatic sites was used to segregate patient into high- or low-DPY30 expression and

generate progression-free and overall survival Kaplan-Meier plots.

Automatic image segmentation and quantification. Fluorescence quantification

was done using MATLAB (The MathWorks, Inc.). Specifically, Otsu's thresholding method

and marker-controlled watershed algorithm was used to segment and specify nuclear

regions of individual cells. Expression of each marker in the individual segmented area

was then quantified by pixel intensity.

ChIP-Seq data analysis. Samples were sequenced using NextSeq500 at

UTMDACC Advanced Technology Genomics Core. Raw reads for each ChIP-seq were

processed using Trimmomatic (v.0.39) (90) to trim off adaptor sequences incorporated in

the read and to remove low quality bases, using the following parameters 'TRAILING:3

LEADING:3 ILLUMINACLIP:adapters.fasta:2:30:10; SLIDINGWINDOW:4:15

MINLEN:50'. Then, the resulting reads were mapped to the to human genome build

GRCh38/hg38 using Bowtie2 v.2 (91) with the '–very-sensitive' preset of parameters.

Reads that did not align to the nuclear genome or aligned to the mitochondrial genome

were removed and duplicate reads were marked and removed using SAMtools (v.1.12)

(92). Peak calling versus the corresponding input genomic DNA was performed using

MACS2 (v.2.2.7.1) (93), for each sample separatly, with the '–nomodel', '–extsize 200', '–

keep-dup all' and '–qvalue 0.05' flags and arguments. Peaks blacklisted by the ENCODE

consortium analysis of artifactual signals in human cells (https://sites.
google.com/site/anshulkundaje/projects/blacklists) were removed using bedtools (v.2.30)
(94). To obtain a confidence peak set, only intersections between biological
replicates were retained and shown as a heat map (generated using deepTools
v.2.5.3) (95). Finally, differential binding analysis was performed using DiffBind
(v.3.0) (96) with settings to use DESeq2 method, full library size and with
subtraction of control input read counts. Binding events with an FDR cutoff of 0.05
were defined as significantly different among experimental conditions. Mean
normalized reads across the three replicates obtained from DiffBind analysis were
used for scatter plot representation using the ggplot2 R package (97).

Proteomic data analysis. Mass spectrometry data were searched using a
local copy of Mascot (Matrix Science) with the following parameters: *enzyme:*
Trypsin/P, *database:* SwissProt Human or Mouse (concatenated forward and
reverse plus common contaminants), *fixed modifications:* Carbamidomethyl (C),
variable modifications: Oxidation (M), Acetyl (N-term), Pyro-Glu (N-term Q),
Deamidation (N,Q), *mass values:* Monoisotopic, *peptide Mass Tolerance:* 10 ppm,
fragment mass tolerance: 0.02 Da, *Max Missed Cleavages:* 2. Mascot DAT files
were parsed into Scaffold (Proteome Software) for validation, filtering and to create
a non-redundant list per sample. Data were filtered using 1% protein and peptide
FDR and requiring at least two unique peptides per protein.

ATAC-seq data analysis. Paired-end ATAC-seq fragments of human cells
were aligned to mouse reference build GRCh38/hg38 using Bowtie2 v.2. (91).
Reads that did not align to the nuclear genome or aligned to the mitochondrial
genome were removed and duplicate reads were marked and removed using
SAMtools (92). Read positions were corrected for transposon insertion offset as
described (98), then only uniquely mapped paired-end reads were kept for further
analysis. All ATAC-seq peaks that were no more than 3 kb away from annotated
gene TSS (GENCODE v19) were selected as promoter (proximal) regions, using
annotatePeaks script from HOMER package (99).

Whole exome sequencing data analysis. DNA sequencing of samples was
performed on the Illumina HiSeq2000 sequencing platform at the UTMDACC

Advanced Technology and Genomics Core. Sequencing fasta files were applied to

FastQC (100) for quality control, adapters were trimmed by trimmomatic (90), and the

genomic fragments were aligned to mm10 using bwa mem (101), then sorted and indexed

by samtools (92). Following Base Quality Score Recalibration (BQSR) by GATK(102).

Somatic mutations were called by Mutect2 and copy number segments were called by

CNVkit (103).

RNA-Seq data analysis. RNA sequencing of samples was performed on the

Illumina NovaSeq6000 sequencing platform at the UTMDACC Advanced Technology and

Genomics Core. FastQC v0.11.9 (https://github.com/s-andrews/FastQC) was used to

perform quality control. Raw reads were first aligned to the human genome GRCh38.p14

or mouse genome GRCm38.p6 using STAR2.7.10b v3.0.0 (104) and sorted using

samtools v1.16 (92). Gene expression was quantified using Rsubread v2.14.2 (105).

Differential expression analysis was performed using edgeR v3.42.4 (106) and limma

v3.56.2 (107). Pathway enrichment analysis was performed using R package fgsea

v1.26.0, clusterProfiler v4.8.2, msigdbr v7.5.1, ReactomePA v1.44.0 (108).

Single-cell RNA-Seq analysis. Tumors deriving from KPC cells were digested and

single cell suspension was diluted in 2% FBS in PBS for further processing. Chromium

single-cell sequencing technology from 10x Genomics was used to perform single-cell

separation, complementary DNA amplification and library construction following the

manufacturer's guidelines. Briefly, the cellular suspensions were loaded on a 10x

Chromium Single Cell Controller to generate single-cell gel bead-in-emulsions. The

scRNA-seq libraries were constructed using the Chromium Single Cell 3' Library & Gel

Bead Kit v.2 (PN-120237, 10x Genomics). The HS dsDNA Qubit Kit was used to

determine the concentrations of both the cDNA and the libraries. The HS DNA

Bioanalyzer was used for quality-tracking purposes and size determination for cDNA and

lower-concentrated libraries. Sample libraries were normalized to 7.5 nM and equal

volumes were added of each library for pooling. The concentration of the library pool was

determined using the Library Quantification qPCR Kit (KAPA Biosystems) before

sequencing. The barcoded library at the concentration of 275 pM was sequenced on the

NovaSeq6000 (Illumina) S2 flow cell (100 cycle kit) using a 26 × 91 run format with 8 bp

index (read 1). To minimize batch effects, the libraries were constructed using the same

versions of reagent kits and following the same protocols, and the libraries were
sequenced on the same NovaSeq6000 flow cell and analyzed together.

The raw scRNA-seq data were pre-processed (demultiplex cellular
barcodes, read alignment, and generation of gene count matrix) using Cell Ranger
Single Cell Software Suite (v7.1.0,10x Genomics). Scrublet v0.2.3 (109), an
algorithm to predict doublets in scRNA-seq data, was used to further clean
doublets. Seurat (version 5.0.2) (110) was applied to the normalized gene-cell
matrix to identify highly variable genes (HVGs) for unsupervised cell clustering.
Principal component analysis (PCA) was performed on the top 2000 HVGs. The
FindNeighbors function of Seurat was used to construct the Shared Nearest
Neighbor (SNN) Graph, based on unsupervised clustering performed with Seurat
function FindClusters and top 60 PCs. Different resolution parameters for
unsupervised clustering were then examined, and cluster marker genes were
checked to determine the optimal number of clusters with distinct transcriptional
profiles. For visualization, the dimensionality was further reduced using Uniform
Manifold Approximation and Projection (UMAP) method (111) with Seurat function
RunUMAP. The PCs used to calculate the embedding were the same as those
used for clustering. Differentially expressed genes (DEGs) were identified for each
cluster using the FindMarkers function in Seurat (110). The top-ranked DEGs and
the enrichment of canonical marker genes were integrated, together with global
cluster distribution and prior knowledge to infer major cell types and cellular states.

Re-replication sequencing. Briefly, 30×10^6 control or Dpy30-KO KPC cells
were seeded in a square bioassay dish (Corning®) and were synchronized in S
phase using a double thymidine block ad described above. After synchronization,
cells were released in complete medium supplemented with 25 μM CldU for 45
minutes. Cells were then washed twice with PBS and a second pulse in complete
medium supplemented with 250 μM IdU was added and incubated for 45 minutes.
Cells were then collected, and cell pellets were lysed in 350 μL of lyses buffer
(12mM Trish-HCl, 0.1x PBS, 6 mM EDTA, 0.5% SDS) supplemented with 40 μL
Proteinase k (20mg/ml) and 10 μL RNAse A (10mg/ml) for 10 minutes at RT and
then 15 minutes at 56°C. Samples were sonicated to obtain chromatin fragments

of ~200 - 600 base pairs (bp). After collecting the 10% of the sample as input control,
samples were diluted using 1.6 mL of ChIP dilution buffer (10 mM Tris-HCl, 140 mM NaCl,
0.1% DOC, 1% Triton-X, 1 mM EDTA). Diluted DNA samples were then denatured for 10
min at 95°C and cool down at 4°C to preserve ssDNA. Next, ssDNA samples were
immunoprecipitated as described above using Dynabeads Protein G (Invitrogen) pre-
conjugated with antibodies against CldU (rat monoclonal anti-BrdU clone BU1/75) and
IdU (mouse monoclonal anti-BrdU clone B44). Eluted ssDNA samples and inputs were
purified with PCR Clean UP kit and eluted in water. ChIP DNA was prepared with xGen
ssDNA & low input DNA library preparation kit and sequenced using NovaSeq X Plus 10
Billion 2x150 PE (142M PE reads per sample).

The raw fastq files were processed using a pipeline of Nextflow. In short, the raw
reads were trimmed to remove adapter sequences, aligned to mouse mm10 genome,
and filtered as described (54). The filtered alignments were then merged, and the
CldU/IdU ratios were calculated with deepTools and normalized with the default window
size of 5000 and loess span of 300000. The coverage tracts of individual and merged
bam files were generated with deepTools (v3.5.4).

Statistical Analysis. *In vitro* and *in vivo* data are presented as the median values
or mean ± S.D. (standard deviation). Statistical analyses were performed using Student's
t-test, one-way or two-way ANOVA test as specified in figure legends. Results from
survival experiments were analyzed with a Log-rank test and expressed as Kaplan–Meier
survival curves.

1257

References

1.   Beach RR, Ricci-Tam C, Brennan CM, Moomau CA, Hsu P-h, Hua B, et al.
Aneuploidy causes non-genetic individuality. Cell. 2017;169(2):229-42. e21.

2.   Bakhoum SF, Ngo B, Laughney AM, Cavallo J-A, Murphy CJ, Ly P, et al.
Chromosomal instability drives metastasis through a cytosolic DNA response. Nature.
2018;553(7689):467-72.

3.   Santaguida S, Richardson A, Iyer DR, M'Saad O, Zasadil L, Knouse KA, et al.
Chromosome Mis-segregation Generates Cell-Cycle-Arrested Cells with Complex Karyotypes
that Are Eliminated by the Immune System. Developmental Cell. 2017;41(6):638-51.e5.

4.   Davoli T, Xu AW, Mengwasser KE, Sack LM, Yoon JC, Park PJ, et al. Cumulative
haploinsufficiency and triplosensitivity drive aneuploidy patterns and shape the cancer genome.
Cell. 2013;155(4):948-62.

5.   Passerini V, Ozeri-Galai E, De Pagter MS, Donnelly N, Schmalbrock S,
Kloosterman WP, et al. The presence of extra chromosomes leads to genomic instability. Nature
communications. 2016;7(1):1-12.

6.   Gordon DJ, Resio B, Pellman D. Causes and consequences of aneuploidy in cancer.
Nature Reviews Genetics. 2012;13(3):189-203.

7.   Holland AJ, Cleveland DW. Boveri revisited: chromosomal instability, aneuploidy
and tumorigenesis. Nature reviews Molecular cell biology. 2009;10(7):478-87.

8.   Sack LM, Davoli T, Li MZ, Li Y, Xu Q, Naxerova K, et al. Profound Tissue
Specificity in Proliferation Control Underlies Cancer Drivers and Aneuploidy Patterns. Cell.
2018;173(2):499-514.e23.

9.   Beroukhim R, Mermel CH, Porter D, Wei G, Raychaudhuri S, Donovan J, et al.
The landscape of somatic copy-number alteration across human cancers. Nature.
2010;463(7283):899-905.

10.   Fragkos M, Ganier O, Coulombe P, Méchali M. DNA replication origin activation
in space and time. Nature Reviews Molecular Cell Biology. 2015;16(6):360-74.

11.   Arias EE, Walter JC. Strength in numbers: preventing rereplication via multiple
mechanisms in eukaryotic cells. Genes & development. 2007;21(5):497-518.

12.   Petropoulos M, Tsaniras SC, Taraviras S, Lygerou Z. Replication licensing
aberrations, replication stress, and genomic instability. Trends in biochemical sciences.
2019;44(9):752-64.

13.   Zhou Y, Pozo PN, Oh S, Stone HM, Cook JG. Distinct and sequential re-replication
barriers ensure precise genome duplication. PLOS Genetics. 2020;16(8):e1008988.

14.   Berti M, Cortez D, Lopes M. The plasticity of DNA replication forks in response
to clinically relevant genotoxic stress. Nature Reviews Molecular Cell Biology. 2020;21(10):633-
51.

15.   Frouin I, Montecucco A, Spadari S, Maga G. DNA replication: a complex matter.
EMBO reports. 2003;4(7):666-70.

16.   Iyer DR, Rhind N. Replication fork slowing and stalling are distinct, checkpoint-
independent consequences of replicating damaged DNA. PLOS Genetics. 2017;13(8):e1006958.

17.   Mulla WA, Seidel CW, Zhu J, Tsai HJ, Smith SE, Singh P, et al. Aneuploidy as a
cause of impaired chromatin silencing and mating-type specification in budding yeast. Elife.
2017;6.

18.   Giunta S, Hervé S, White RR, Wilhelm T, Dumont M, Scelfo A, et al. CENP-A
chromatin prevents replication stress at centromeres to avoid structural aneuploidy. Proceedings
of the National Academy of Sciences. 2021;118(10):e2015634118.

19.   Lambuta RA, Nanni L, Liu Y, Diaz-Miyar J, Iyer A, Tavernari D, et al. Whole-
genome doubling drives oncogenic loss of chromatin segregation. Nature. 2023.

20.   Muñoz S, Búa S, Rodríguez-Acebes S, Megías D, Ortega S, de Martino A, et al.
In Vivo DNA Re-replication Elicits Lethal Tissue Dysplasias. Cell Reports. 2017;19(5):928-38.

21.   Técher H, Koundrioukoff S, Nicolas A, Debatisse M. The impact of replication
stress on replication dynamics and DNA damage in vertebrate cells. Nat Rev Genet.
2017;18(9):535-50.

22.   Flach J, Bakker ST, Mohrin M, Conroy PC, Pietras EM, Reynaud D, et al.
Replication stress is a potent driver of functional decline in ageing haematopoietic stem cells.
Nature. 2014;512(7513):198-202.

23.   Tsunematsu T, Takihara Y, Ishimaru N, Pagano M, Takata T, Kudo Y. Aurora-A
controls pre-replicative complex assembly and DNA replication by stabilizing geminin in mitosis.
Nature Communications. 2013;4(1):1885.

24.   Liontos M, Koutsami M, Sideridou M, Evangelou K, Kletsas D, Levy B, et al.
Deregulated overexpression of hCdt1 and hCdc6 promotes malignant behavior. Cancer research.
2007;67(22):10899-909.

25.   Fu H, Redon CE, Thakur BL, Utani K, Sebastian R, Jang S-M, et al. Dynamics of
replication origin over-activation. Nature Communications. 2021;12(1):3448.

26.   Connor AA, Gallinger S. Pancreatic cancer evolution and heterogeneity:
integrating omics and clinical data. Nature Reviews Cancer. 2022;22(3):131-42.

27.   Ryan DP, Hong TS, Bardeesy N. Pancreatic Adenocarcinoma. New England
Journal of Medicine. 2014;371(11):1039-49.

28.   Bailey P, Chang DK, Nones K, Johns AL, Patch A-M, Gingras M-C, et al. Genomic
analyses identify molecular subtypes of pancreatic cancer. Nature. 2016;531(7592):47-52.

29.   Cao L, Huang C, Cui Zhou D, Hu Y, Lih TM, Savage SR, et al. Proteogenomic
characterization of pancreatic ductal adenocarcinoma. Cell. 2021;184(19):5031-52.e26.

30.   Hayashi A, Hong J, Iacobuzio-Donahue CA. The pancreatic cancer genome
revisited. Nature Reviews Gastroenterology & Hepatology. 2021;18(7):469-81.

31.   Carugo A, Genovese G, Seth S, Nezi L, Rose JL, Bossi D, et al. In Vivo
Functional Platform Targeting Patient-Derived Xenografts Identifies WDR5-Myc Association as
a Critical Determinant of Pancreatic Cancer. Cell Reports. 2016;16(1):133-47.

32.   Giuliani V, Miller MA, Liu C-Y, Hartono SR, Class CA, Bristow CA, et al.
PRMT1-dependent regulation of RNA metabolism and DNA damage response sustains pancreatic
ductal adenocarcinoma. Nature Communications. 2021;12(1):4626.

33.   Ali A, Tyagi S. Diverse roles of WDR5-RbBP5-ASH2L-DPY30 (WRAD) complex
in the functions of the SET1 histone methyltransferase family. Journal of biosciences.
2017;42(1):155-9.

34.   Ali A, Veeranki SN, Tyagi S. A SET-domain-independent role of WRAD complex
in cell-cycle regulatory function of mixed lineage leukemia. Nucleic acids research.
2014;42(12):7611-24.

35.   Ernst P, Vakoc CR. WRAD: enabler of the SET1-family of H3K4
methyltransferases. Briefings in functional genomics. 2012;11(3):217-26.

36.   Wang H, Fan Z, Shliaha PV, Miele M, Hendrickson RC, Jiang X, et al. H3K4me3
regulates RNA polymerase II promoter-proximal pause-release. Nature. 2023;615(7951):339-48.

37.   Cenik BK, Shilatifard A. COMPASS and SWI/SNF complexes in development and
disease. Nature Reviews Genetics. 2021;22(1):38-58.

38.   Campbell SA, Bégin J, McDonald CL, Vanderkruk B, Stephan TL, Hoffman BG.
H3K4 trimethylation is required for postnatal pancreatic endocrine cell functional maturation.
bioRxiv. 2020:2020.11.29.402990.

39.   Campbell SA, McDonald CL, Krentz NAJ, Lynn FC, Hoffman BG. TrxG Complex
Catalytic and Non-catalytic Activity Play Distinct Roles in Pancreas Progenitor Specification and
Differentiation. Cell Reports. 2019;28(7):1830-44.e6.

40.   Dixit D, Prager BC, Gimple RC, Miller TE, Wu Q, Yomtoubian S, et al.
Glioblastoma stem cells reprogram chromatin in vivo to generate selective therapeutic
dependencies on DPY30 and phosphodiesterases. Science Translational Medicine.
2022;14(626):eabf3917.

41.   Simboeck E, Gutierrez A, Cozzuto L, Beringer M, Caizzi L, M Keyes W, et al.
DPY30 regulates pathways in cellular senescence through ID protein expression. The EMBO
journal. 2013;32(16):2217-30.

42.   Yang Z, Shah K, Khodadadi-Jamayran A, Jiang H. Control of Hematopoietic Stem
and Progenitor Cell Function through Epigenetic Regulation of Energy Metabolism and Genome
Integrity. Stem Cell Reports. 2019;13(1):61-75.

43.   Zhao T, Hong Y, Ming G-l, Song H. Loss of chromatin modulator Dpy30
compromises proliferation and differentiation of postnatal neural stem cells. Journal of Molecular
Cell Biology. 2019;12(1):2-3.

44.   Zhao Y, Hao X, Li Z, Feng X, Katz J, Michalek SM, et al. Role of chromatin
modulator Dpy30 in osteoclast differentiation and function. Bone. 2022;159:116379.

45.   Pferdehirt RR, Kruesi WS, Meyer BJ. An MLL/COMPASS subunit functions in
the C. elegans dosage compensation complex to target X chromosomes for transcriptional
regulation of gene expression. Genes & development. 2011;25(5):499-515.

46.   Hingorani SR, Wang L, Multani AS, Combs C, Deramaudt TB, Hruban RH, et al.
Trp53R172H and KrasG12D cooperate to promote chromosomal instability and widely metastatic
pancreatic ductal adenocarcinoma in mice. Cancer Cell. 2005;7(5):469-83.

47.   Yan L, Tu B, Yao J, Gong J, Carugo A, Bristow CA, et al. Targeting Glucose
Metabolism Sensitizes Pancreatic Cancer to MEK Inhibition. Cancer Research. 2021;81(15):4054-
65.

48.   Dou Y, Milne TA, Ruthenburg AJ, Lee S, Lee JW, Verdine GL, et al. Regulation
of MLL1 H3K4 methyltransferase activity by its core components. Nature Structural & Molecular
Biology. 2006;13(8):713-9.

49.   Bryan AF, Wang J, Howard GC, Guarnaccia AD, Woodley CM, Aho ER, et al.
WDR5 is a conserved regulator of protein synthesis gene expression. Nucleic Acids Research.
2020;48(6):2924-41.

50.   Grant GD, Kedziora KM, Limas JC, Cook JG, Purvis JE. Accurate delineation of
cell cycle phase transitions in living cells with PIP-FUCCI. Cell Cycle. 2018;17(21-22):2496-516.

51.   Tremblay V, Zhang P, Chaturvedi C-P, Thornton J, Brunzelle Joseph S, Skiniotis
G, et al. Molecular Basis for DPY-30 Association to COMPASS-like and NURF Complexes.
Structure. 2014;22(12):1821-30.

52.   Roy S, Luzwick JW, Schlacher K. SIRF: Quantitative in situ analysis of protein
interactions at DNA replication forks. J Cell Biol. 2018;217(4):1521-36.

53.   Roy S, Schlacher K. SIRF: A Single-cell Assay for in situ Protein Interaction with
Nascent DNA Replication Forks. Bio Protoc. 2019;9(18):e3377.

54.   Marchal C, Sasaki T, Vera D, Wilson K, Sima J, Rivera-Mulia JC, et al. Genome-
wide analysis of replication timing by next-generation sequencing with E/L Repli-seq. Nature
Protocols. 2018;13(5):819-39.

55.   Bakhoum SF, Cantley LC. The Multifaceted Role of Chromosomal Instability in
Cancer and Its Microenvironment. Cell. 2018;174(6):1347-60.

56.   Decout A, Katz JD, Venkatraman S, Ablasser A. The cGAS–STING pathway as a
therapeutic target in inflammatory diseases. Nature Reviews Immunology. 2021;21(9):548-69.

57.   Chuang L-C, Teixeira LK, Wohlschlegel JA, Henze M, Yates JR, Méndez J, et al.
Phosphorylation of Mcm2 by Cdc7 Promotes Pre-replication Complex Assembly during Cell-
Cycle Re-entry. Molecular Cell. 2009;35(2):206-16.

58.   Vaziri C, Saxena S, Jeon Y, Lee C, Murata K, Machida Y, et al. A p53-Dependent
Checkpoint Pathway Prevents Rereplication. Molecular Cell. 2003;11(4):997-1008.

59.   Kalfalah FM, Berg E, Christensen MO, Linka RM, Dirks WG, Boege F, et al.
Spatio-temporal regulation of the human licensing factor Cdc6 in replication and mitosis. Cell
Cycle. 2015;14(11):1704-15.

60.   McGarry TJ, Kirschner MW. Geminin, an Inhibitor of DNA Replication, Is
Degraded during Mitosis. Cell. 1998;93(6):1043-53.

61.   Blank JL, Liu XJ, Cosmopoulos K, Bouck DC, Garcia K, Bernard H, et al. Novel
DNA Damage Checkpoints Mediating Cell Death Induced by the NEDD8-Activating Enzyme
Inhibitor MLN4924. Cancer Research. 2013;73(1):225-34.

62.   Li A, Blow JJ. Cdt1 downregulation by proteolysis and geminin inhibition prevents
DNA re-replication in Xenopus. Embo j. 2005;24(2):395-404.

63. Walter D, Hoffmann S, Komseli E-S, Rappsilber J, Gorgoulis V, Sørensen CS.
SCFCyclin F-dependent degradation of CDC6 suppresses DNA re-replication. Nature
Communications. 2016;7(1):10530.

64. Neelsen KJ, Zanini IM, Mijic S, Herrador R, Zellweger R, Chaudhuri AR, et al.
Deregulated origin licensing leads to chromosomal breaks by rereplication of a gapped DNA
template. Genes & development. 2013;27(23):2537-42.

65. Mehta A, Merkel OM. Immunogenicity of Cas9 Protein. Journal of Pharmaceutical
Sciences. 2020;109(1):62-7.

66. Seth S, Li CY, Ho IL, Corti D, Loponte S, Sapio L, et al. Pre-existing Functional
Heterogeneity of Tumorigenic Compartment as the Origin of Chemoresistance in Pancreatic
Tumors. Cell Rep. 2019;26(6):1518-32.e9.

67. Ho I-L, Li C-Y, Wang F, Zhao L, Liu J, Yen E-Y, et al. Clonal dominance defines
metastatic dissemination in pancreatic cancer. Science Advances. 2024;10(11):eadd9342.

68. Ying H, Kimmelman Alec C, Lyssiotis Costas A, Hua S, Chu Gerald C, Fletcher-
Sananikone E, et al. Oncogenic Kras Maintains Pancreatic Tumors through Regulation of Anabolic
Glucose Metabolism. Cell. 2012;149(3):656-70.

69. Yao W, Rose JL, Wang W, Seth S, Jiang H, Taguchi A, et al. Syndecan 1 is a
critical mediator of macropinocytosis in pancreatic cancer. Nature. 2019;568(7752):410-4.

70. Padrón LJ, Maurer DM, O'Hara MH, O'Reilly EM, Wolff RA, Wainberg ZA, et
al. Sotigalimab and/or nivolumab with chemotherapy in first-line metastatic pancreatic cancer:
clinical and immunologic analyses from the randomized phase 2 PRINCE trial. Nature Medicine.
2022;28(6):1167-77.

71. Moffitt RA, Marayati R, Flate EL, Volmar KE, Loeza SGH, Hoadley KA, et al.
Virtual microdissection identifies distinct tumor- and stroma-specific subtypes of pancreatic ductal
adenocarcinoma. Nature Genetics. 2015;47(10):1168-78.

72. Puleo F, Nicolle R, Blum Y, Cros J, Marisa L, Demetter P, et al. Stratification of
Pancreatic Ductal Adenocarcinomas Based on Tumor and Microenvironment Features.
Gastroenterology. 2018;155(6):1999-2013.e3.

73. Collisson EA, Sadanandam A, Olson P, Gibb WJ, Truitt M, Gu S, et al. Subtypes
of pancreatic ductal adenocarcinoma and their differing responses to therapy. Nature Medicine.
2011;17(4):500-3.

74.   Del Poggetto E, Ho I-L, Balestrieri C, Yen E-Y, Zhang S, Citron F, et al. Epithelial
memory of inflammation limits tissue damage while promoting pancreatic tumorigenesis. Science.
2021;373(6561):eabj0486.

75.   Alonso-Curbelo D, Ho Y-J, Burdziak C, Maag JLV, Morris JP, Chandwani R, et
al. A gene–environment-induced epigenetic program initiates tumorigenesis. Nature.
2021;590(7847):642-8.

76.   Lomberk G, Blum Y, Nicolle R, Nair A, Gaonkar KS, Marisa L, et al. Distinct
epigenetic landscapes underlie the pathobiology of pancreatic cancer subtypes. Nature
Communications. 2018;9(1):1978.

77.   Dias J, Van Nguyen N, Georgiev P, Gaub A, Brettschneider J, Cusack S, et al.
Structural analysis of the KANSL1/WDR5/KANSL2 complex reveals that WDR5 is required for
efficient assembly and chromatin targeting of the NSL complex. Genes & development.
2014;28(9):929-42.

78.   Li Q, Mao F, Zhou B, Huang Y, Zou Z, denDekker AD, et al. p53 Integrates
Temporal WDR5 Inputs during Neuroectoderm and Mesoderm Differentiation of Mouse
Embryonic Stem Cells. Cell Rep. 2020;30(2):465-80.e6.

79.   Siladi AJ, Wang J, Florian AC, Thomas LR, Creighton JH, Matlock BK, et al. WIN
site inhibition disrupts a subset of WDR5 function. Scientific Reports. 2022;12(1):1848.

80.   Zhao J, Chen W, Pan Y, Zhang Y, Sun H, Wang H, et al. Structural insights into
the recognition of histone H3Q5 serotonylation by WDR5. Science Advances.
2021;7(25):eabf4291.

81.   Morgan MAJ, Shilatifard A. Epigenetic moonlighting: Catalytic-independent
functions of histone modifiers in regulating transcription. Science Advances.
2023;9(16):eadg6593.

82.   Saldivar JC, Hamperl S, Bocek MJ, Chung M, Bass TE, Cisneros-Soberanis F, et
al. An intrinsic S/G$_2$ checkpoint enforced by ATR. Science. 2018;361(6404):806-10.

83.   Saldivar JC, Cortez D, Cimprich KA. The essential kinase ATR: ensuring faithful
duplication of a challenging genome. Nature reviews Molecular cell biology. 2017;18(10):622-36.

84.   Zeng J, Hills SA, Ozono E, Diffley JFX. Cyclin E-induced replicative stress drives
p53-dependent whole-genome duplication. Cell. 2023;186(3):528-42.e14.

85.   Donnelly N, Storchová Z. Dynamic karyotype, dynamic proteome: buffering the
effects of aneuploidy. Biochimica et Biophysica Acta (BBA) - Molecular Cell Research.
2014;1843(2):473-81.

86.   Dungrawala H, Cortez D. Purification of proteins on newly synthesized DNA using
iPOND. Methods Mol Biol. 2015;1228:123-31.

87.   Halliwell JA, Gravells P, Bryant HE. DNA Fiber Assay for the Analysis of DNA
Replication Progression in Human Pluripotent Stem Cells. Current Protocols in Stem Cell Biology.
2020;54(1):e115.

88.   Di Pilato M, Kfuri-Rubens R, Pruessmann JN, Ozga AJ, Messemaker M, Cadilha
BL, et al. CXCR6 positions cytotoxic T cells to receive critical survival signals in the tumor
microenvironment. Cell. 2021;184(17):4512-30.e22.

89.   Terranova C, Tang M, Orouji E, Maitituoheti M, Raman A, Amin S, et al. An
Integrated Platform for Genome-wide Mapping of Chromatin States Using High-throughput ChIP-
sequencing in Tumor Tissues. J Vis Exp. 2018(134).

90.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014;30(15):2114-20.

91.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature
Methods. 2012;9(4):357-9.

92.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.

93.   Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-
based Analysis of ChIP-Seq (MACS). Genome Biology. 2008;9(9):R137.

94.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 2010;26(6):841-2.

95.   Ramírez F, Dündar F, Diehl S, Grüning BA, Manke T. deepTools: a flexible
platform for exploring deep-sequencing data. Nucleic Acids Res. 2014;42(Web Server
issue):W187-91.

96.   Ross-Innes CS, Stark R, Teschendorff AE, Holmes KA, Ali HR, Dunning MJ, et
al. Differential oestrogen receptor binding is associated with clinical outcome in breast cancer.
Nature. 2012;481(7381):389-93.

97.   Wickham H. ggplot2: elegant graphics for data analysis. New York: Springer New York, NY; 2016. Available from: https://link.springer.com/book/10.1007/978-0-387-98141-3.

98.   Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. Nature Methods. 2013;10(12):1213-8.

99.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. Molecular Cell. 2010;38(4):576-89.

100.   Andrews S. FastQC. A quality control tool for high throughput sequence data 2010. Available from: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

101.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 2009;25(14):1754-60.

102.   Geraldine A. Van der Auwera BDOC. Genomics in the Cloud: O'Reilly Media, Inc.; 2020.

103.   Talevich E, Shain AH, Botton T, Bastian BC. CNVkit: Genome-Wide Copy Number Detection and Visualization from Targeted DNA Sequencing. PLOS Computational Biology. 2016;12(4):e1004873.

104.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.

105.   Liao Y, Smyth GK, Shi W. The R package Rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. Nucleic Acids Research. 2019;47(8):e47-e.

106.   McCarthy DJ, Chen Y, Smyth GK. Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. Nucleic Acids Research. 2012;40(10):4288-97.

107.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research. 2015;43(7):e47-e.

108.   Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. Fast gene set enrichment analysis. bioRxiv. 2021:060012.

109.   Wolock SL, Lopez R, Klein AM. Scrublet: Computational Identification of Cell
Doublets in Single-Cell Transcriptomic Data. Cell Syst. 2019;8(4):281-91.e9.

110.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, et al.
Integrated analysis of multimodal single-cell data. Cell. 2021;184(13):3573-87.e29.

111.   Leland McInnes JHaJM. UMAP: Uniform Manifold Approximation and Projection
for Dimension Reduction: arXiv; 2018. Available from: https://arxiv.org/abs/1802.03426.

1545

Acknowledgments: We thank Michael Kim for PDX-derived PDAC cell lines; all
Draetta lab members and Guoxin Feng for scientific discussion; R. Nguyen and J.
Martinez in the Department of Genomic Medicine at MD Anderson for lab management;
Michael Peoples and Vandhana Ramamoorthy with the MD Anderson TRACTION
platform for valuable support with cloning; Alejandro Hernandez Martinez, Edward Q.
Chang and Ningping Feng with the MD Anderson TRACTION platform for support with
tissue imaging acquisition and drug formulation; and Vivien Van, Charles Kingsley and
Jorge Delacerda at the MD Anderson Small Animals Imaging Facility; The Animal Support
Facility (RASF), which was partially funded by the MD Anderson Support Grant
CA016672; the MD Anderson Department of Veterinary Medicine; the MD Anderson
Advanced Microscopy Core, which was funded by NIH S10 RR029552; Asha Multani and
the MD Anderson Cytogenetics and Cell Authentication Core; Erika Thompson, Hongli
Tang, David Pollock and the MD Anderson Advanced Technology Genomics Core
(ATGC), which was funded by the CA016672 Core Grant; Karen Ramirez, David Dwyer,
Veena Papanna, and the MD Anderson Advanced Cytometry and Sorting Facility, which
was funded by the NCI Cancer Center Support Grant P30CA16672; and Genscript for
support and service. The mEmerald-MCM vectors were a gift from Michael Davidson, the
lentiCRISPR v2 vector was a gift from Feng Zhang. The pcDNA3.1 Flag-hDPY30 vector
was a gift from Kai Ge. The psPAX2 and pMD2G vectors were a gift from Didier Trono.
Schematics were created using BioRender.com. This research was supported by the
2022 AACR–AstraZeneca START Grant, Grant Number 22-40-12-CITR to F.C.; AIRC
and the European Union's Horizon 2020 research and innovation program under the
Marie Skłodowska-Curie grant agreement no. 800924 to F.C.; CPRIT Training Award
(RP210028) to E.Y.; N.P. is a TRIUMPH Fellow in the CPRIT Training Program

(RP210028); Cancer Research and Prevention Institute of Texas (CPRIT) grant

RP190599, NIH/NCI R01 CA258917, the Andrew Sabin Family Foundation grant, and

the V Foundation grant (V2020-018) to A.V.; NIH/NCI SPORE in gastrointestinal cancer

grant P50CA221707 to G.F.D and A.V.; UT MD Anderson Cancer Center Start-Up Funds

to A.V.; UT MD Anderson Cancer Center Pancreatic Cancer Moon Shot and Pancreatic

Cancer Action Network (PanCAN) to G.F.D.; NIH/NCI P01 CA117969-12 to A.V. and

G.F.D. G.F.D. was also supported by the Sewell Family Distinguished University Chair in

Genomic Medicine.

1578

Author contributions: Conceptualization: F.C., A.V., G.F.D.; Methodology: F.C.,

L.P., I.H., C.B., E.Y., Z.L., L.Z., L.C.M, P.J., S.A., M.D.P., W.Y., A.Z., C.J.; Software: Y.C.,

C.B., L.Z., Z.L., E.Y., S.S., L.W., G.T., C.Bo.; Validation: F.C., L.P, I.H, E.Y., L.C, H.K.,

R.S., Z.C., I.S., N.P., S.A., K.C.C, C.A.D., C.L., M.M., K.R., C.J.; Formal Analysis: F.C.,

L.P., Z.L., C.B., L.Z., E.Y., S.S., M.D.P.; Investigation: F.C., L.P, I.H, E.Y., K.C.C., L.C.M.,

M.D.P.; Resources: F.C., L.P., I.H., C.B., Z.L., L.Z., E.Y., R.S., L.C.M., S.A., S.J., B.G.H.,

B.V., C.L., M.M., K.R., H.W., W.Y., V.G., T.H., G.T., L.W., M.D.P.; Data curation: F.C.,

L.P., I.H., Z.L., C.B., L.Z., E.Y.; Writing – Original draft: F.C., A.V., G.F.D.; Writing –

Review and Editing: F.C., A.V., G.F.D., S.G.; Visualization: F.C., C.B., Z.L., E.Y.;

Supervision: F.C., A.V., G.F.D; Project Administration: F.C., A.V., G.F.D; Funding

Acquisition: F.C., A.V., G.F.D;

1590

1591 **Competing interest declaration**

1592 The authors declare that they have no conflict of interest.

1593

1594 **Figure legends**

Figure 1

Figure 1. *DPY30* is dispensable for cancer cell proliferation and H3K4 trimethylation.

    a. Growth curve analysis of PATC53 cells carrying shRNAs targeting WDR5 (shWDR5), DPY30 (shDPY30) or control (shCTR) and treated with vehicle (off) or

doxycycline (dox) to induce the gene silencing. **b.** Growth curve (left) and western blot

(right) of control (sgCTR) or *Dpy30*-KO (sgDPY30) KPC cells. **c.** Clonogenic assay of

KPC cells, as described in b, is shown (left) and quantified (right). **d-e.** Volume (d) and

weight (e) of tumors from control (sgCTR) or DPY30-KO (sgDPY30) PATC53 cells grown

in NSG mice. **f-g**. Micrographs showing the histology and Ki67 labeling in tumors

described in (d). Hematoxylin and eosin (H&E) (f) staining, Ki67 IHC staining (g, left) and

quantification (g, right) are shown. **h-j.** Heatmap showing Spearman correlations between

samples and biological replicates of ATACseq (h), and H3K27ac (i) or H3K4me3 (j) Chip-

seq in shCTR, shDPY30 or shWDR5 PATC53 cells. Samples are clustered by Euclidean

distance and single-linkage clustering. **k.** Scatter plot showing mean normalized signals

of H3K4me3 from samples described in j. Regions that significantly (FDR < 0.05) gained

or lost H3K4me3 peaks in shDPY30 or shWDR5 versus shCTR are depicted in pink and

blue, respectively. Not significant peaks are indicated as gray dots. In (a-e, g), data are

expressed as mean $\pm$ SD, and dots represent biological replicates. Statistical significance

was calculated using one-way ANOVA and Tukey's multiple comparison tests (a, d, e),

or Student t test (b, c, g). In panels f-g, scale bar = 200 μm.

Figure 2

1617 **Figure 2. The WRAD core interacts with WDHD1 and the MCM complex.**

1618   **a.** Schematic representation of DPY30 interactors involved in DNA repair,

1619 replisome, and proteosome signaling pathways in PATC53, PATC66, and KPC cells. **b.**

Western blot analysis of the indicated proteins in 293T cells co-overexpressing HA-tagged human *DPY30* and His-tagged human MCM complex subunits. Control IgG and input are shown. **c.** Proximity Ligation Assay (PLA) staining detecting DPY30 interaction with WDHD1, MCM6, Ash2L (positive control) or Luciferase (negative control). **d.** Western blot analysis of the indicated proteins in 293T cells co-overexpressing Flag-tagged human *WDR5* and His-tagged human MCM complex subunits. Control IgG and input are shown. **e.** PLA staining detecting WDR5 interaction with WDHD1, MCM6, RbBP5 (positive control) or Luciferase (negative control). **f.** PLA staining detecting WDR5 interaction with WDHD1 and MCM6 in PATC53 control (shCTR) or silenced for WDR5 (shWDR5). **g-h.** Western blot analysis of the indicated proteins in PATC53 cells control (shCTR) or silenced for WDR5 (shWDR5, g) or DPY30 (shDPy30, h). Control IgG and input are shown, and H3 was used as a loading control. **i-j.** Western blot analyses of the indicated proteins in KPC cells control or overexpressing DPY30 wild type (WT) or mutant (L69D) isoforms (i), and in KPC cells control or overexpressing WDR5 wild type (WT) or mutant (L240K, V268E) isoforms (j). Control IgG and input are shown, and H3 was used as a loading control. In figure, scale bar = 5 μm.

Figure 3

Figure 3. DPY30 loss leads to DNA re-replication and CIN.

a. Western blot analysis of proteins bound to nascent DNA retrieved using iPOND or input control in KPC cells. Cells were treated as follows: control (negative control, no biotin in the click reaction), EdU pulse (EdU treatment for 2, 5 or 10 min), chase (EdU treatment for 10 min, followed by a 60 min thymidine chase). Histone H3 was used as a loading control. **b.** Modified PLA (SIRF) assay detecting the interaction between the incorporated BrdU (5 min pulse) and WDR5 (left) or DPY30 (right). BrdU:PCNA and BrdU:luciferase were used as positive and negative control, respectively. Scale bar = 5 μm. **c.** Western blot analysis of iPOND or input control in control (sgCTR) or *Dpy30*-KO (sgDpy30) KPC cells. Cells were treated as described in a. Histone H3 was used as a loading control. **d.** Schematic illustration and confocal images of DNA fiber assay in sgCTR and sgDpy30 KPC cells. **e.** Violin plots reporting mean fork progression rate (left) and fork symmetry (right) in cells as in d. **f.** Confocal images of DNA fiber assay in sgCTR or sgDpy30 KPC cells. White arrows indicate a fiber with non-aberrant DNA replication, yellow arrows indicate overlapped CldU and IdU incorporation (DNA re-replication). **g.** Violin plots reporting the frequency of DNA re-replication (left) and the fiber re-replication ratio (right) in cells as in f. **h.** Trend curves showing the dual CldU:IdU incorporation (score) calculated from the density of ChIPseq reads of IdU and CldU IP in sgCTR and sgDpy30 KPC cells. **h.** Bar plot showing genome length of single (IdU or CldU) and double nucleotide (CldU+IdU) incorporation in sgCTR and sgDpy30. **j.** immunofluorescence images of sgCTR or sgDpy30 KPC cells stained for pS4-8 RPA (red), pS139 H2AX (γH2AX, green), and nuclei (blue). Right, graphs reporting the percentage of γH2AX-positive or pS4-8 RPA-positive sgCTR or sgDpy30 KPC cells. Each dot represents a different analyzed field, scale bar = 100 μm. **k.** Left, immunofluorescence images of mitotic KPC cells stained with DAPI (DNA, white). Yellow arrows indicate lagging chromosomes and chromosome bridges. Right, violin plot reporting the mean percentage of sgCTR or sgDpy30 KPC cells with aberrant mitotic outcomes *in vitro*. **k.** Left panel, images of metaphase spreads in sgCTR or sgDpy30 KPC cells stained with Giemsa (chromosome). Red arrow indicates chromosome fusion and fragmentation. Right panel, percentage of KPC cells with normal (gray) or aberrant (yellow) mitotic outcomes. In the figure, n indicates the number of analyzed events. Data are expressed as mean values ±

 SD. Statistical significance was calculated using the Mann-Whitney test (e, g), Student's

 t test (j, k) or Fisher's exact Chi square test (i, l).

Figure 4

1670

Figure 4. DPY30 loss results in a distinct immune context-dependent phenotype

a–**c.** Volume of control (sgCTR) and *Dpy30*-KO (sgDpy30) KPC tumors grown in the flank of immune-deficient NSG (a), nude (b), or in immune-competent C57BL/6 mice (c). **d.** Weight (mg) of tumors described in (a–c). **e.** Hematoxylin and eosin (H&E) staining of tumors from experiments described in (a–d). **f.** Alcian blue staining of tumors from sgCTR and sgDpy30 KPC cells grown in C57BL/6 mice. **g.** Left, T2 Magnetic Resonance Imaging (MRI) scans of tumors from sgCTR and sgDpy30 KPC cells grown orthotopically in the pancreas of C57BL/6 mice. Right, Kaplan-Meier survival curve of tumor-bearing mice. Statistical significance and Hazard Ratio (HR) were calculated using the Log-rank test. **h.** Immunofluorescence staining of Ki67 and DPY30 in sgCTR and sgDpy30 KPC cells grown in C57BL/6 mice. **i.** Left, STING immunohistochemical (IHC) staining of tumors from sgCTR and sgDpy30 KPC cells grown in C57BL/6 mice Right, violin plot showing the percentage of STING positive cells per field. **j.** UMAP plots showing bi-dimensional cluster distribution of sgCTR (n=3) or sgDpy30 (n=3) KPC tumor epithelial cells after filtering and quality control. **k.** Bubble plot showing expression levels of the genes involved in cell cycle analyzed from the single-cell RNA sequencing data as described in j. Size of dots represents the percentage of cells expressing the gene; color scale shows the average expression level **l.** Bubble plot showing the over-representation analysis of genes enriched in sgDpy30 KPC cells. Gene expression was calculated using Seurat. **m.** Violin plot showing the expression of *B2m, H2-K1, H2-D1 and Cd74* per cell calculated using Seurat in sgCTR and sgDpy30 KPC cells. **n.** Immunofluorescence staining of CD3 and αSMA in sgCTR or sgDpy30 KPC cells grown in C57BL/6 mice. In the figure, n indicates the number of mice analyzed. In (a–d), data are expressed as mean ± SD and statistical significance was calculated using an unpaired One-way ANOVA (a-c) or Student's *t* test (d). In (l, m), data are expressed as median values and statistical significance was calculated using the Mann Whitney test. In (e–f, i), scale bar = 200 μm. In (h, n), scale bar = 50 μm.

Figure 5

CD3 CC3 DAPI Ki67 αSMA CK19   100 μm

Figure 5. *Dpy30*-KO tumors display higher T cell infiltration, activation, and heterogeneity.

a-b. UMAP plot shows bi-dimensional sample (a) and cell type (b) distribution of control (sgCTR) or *Dpy30*-KO (sgDpy30) KPC tumors after filtering and quality control calculated using Seurat. **c.** Cell fraction of T cells, NK cells, tumor cells and fibroblasts calculated using Seurat in sgCTR and sgDpy30 KPC tumors. **d.** Percentage of CD8+ T cells, CD4+ conventional T cells, CD4+/Foxp3+ T regulatory cells, CD3+ NK, and NK

cells in sgCTR or sgDpy30 KPC tumors grown in immune-competent C57BL/6 mice. Data

were analyzed using flow cytometry. **e.** Percentage of CD8+/PD-1+ T cells in sgCTR or

sgDpy30 KPC tumors grown in C57BL/6 mice and expressing the cytotoxic markers

Gzmb, Ifng, Tnf as analyzed by flow cytometry. **f.** Violin plots showing the percentage of

barcode changes in sgCTR and sgDpy30 KPC injected in NSG (blue) or C57/BL6 (pink)

background. **g.** Multiplexed immunofluorescence staining of CD3 (green), cleaved

caspase 3 (CC3, red), Ki67 (yellow), αSMA (cyan), cytokeratin 19 (CK19, magenta) and

DAPI (blue) in sgCTR or sgDpy30 KPC cells grown in C57BL/6 mice and collected after

10, 15 or 25 days after cell injection. **h-j.** Bar plot showing the percentage of CD3/Ki67+

cells (h), CC3+ cells (i) or Ki67+ cells (j) in the samples described in g. In (c-e, h-j) each

dot represents a biological replicate and data are presented as mean $\pm$ SD and statistical

significance was calculated using Student's *t* test (c-e) or one-way ANOVA (h-j).

Figure 6

Figure 6. DPY30 expression correlates with tumor stage, poor outcome, and resistance to ICB.

a. Schematic representation of the in vivo experiment: control (sgCTR) or *Dpy30*-KO (sgDpy30) KPC cells were injected subcutaneously into C57BL/6 mice. After 5 days mice were randomized and treated starting from day 7 with monoclonal antibodies against

CD8 (α-CD8) or PD-1 (α-PD-1) or isotype control three times per week, for two weeks.

b-c. Graphs reporting the tumor volumes (b) or Kaplan-Meier survival curves (c) of sgCTR

or sgDpy30 KPC tumors as described in a. **d.** DPY30 immunohistochemical (IHC) staining

in normal pancreas and PDAC samples included in the tissue microarray (TMA). **e.**

Kaplan–Meier survival curve of patients with PDAC included in the TMA described in (d)

and divided according to low (IHC score 0 or 1), medium (IHC score 2), or high (IHC score

3) DPY30 expression. **f.** Kaplan–Meier survival curve of patients in the TCGA PDAC

cohort and divided according to low or high *DPY30* expression*.* **g.** Volcano plot

representing GSEA in high-purity TCGA PDAC samples and divided according to high--

or low *DPY30* expression. **h.** Kaplan–Meier progression-free survival curve of patients

included in the phase 2 PRINCE PDAC trial and divided according to low or high *DPY30*

expression*.* In (c, e, f, h) statistical significance and Hazard Ratio (HR) were calculated

using the Log-rank test. In (b) data are presented as mean $\pm$ SD and statistical

significance was calculated using one-way ANOVA.

Extended Data Figure 1

1739

Extended Data Figure 1. DPY30 is dispensable for cell proliferation.

a-d. Western Blot analysis of PATC53 cells harboring shRNAs against WDR5, RBBP5, ASH2L, DPY30 or control (respectively shWDR5, shRBBP5, shASH2L, shDPY30 or shCTR). **e.** Growth curve analysis of PATC53 cells described in a-d. **f.** Western Blot analysis of control (sgCTR) or DPY30-KO PATC53 cells. **g.** Western Blot analysis of KPC cells silenced for Dpy30 (shDPY30) or control (shCTR). **h.** Growth curve of KPC cells described in g. **i.** Representation (left) and quantification of a clonogenic assay of KPC cells, as described in g. **j.** Western Blot analysis of sgCTR or sgDpy30 KP$^{-/-}$C cells. **k.** Growth curve of KPC cells described in j. **l.** Representation (left) and quantification of a clonogenic assay of KP$^{-/-}$C cells, as described in j. **m.** Western Blot analysis of shCTR or shDpy30 KP$^{-/-}$C cells. **n.** Growth curve of KP$^{-/-}$C cells described in m. **o.** Representation (left) and quantification (right) of a clonogenic assay of KP$^{-/-}$C cells, as described in m. **p.** Western Blot analysis of shCTR or shDPY30 PATC66 cells. **q.** Volume (left) and weight (right) of tumors from shCTR or shDPY30 PATC66 cells injected subcutaneously into the flank of NSG mice. **r.** Histopathological analysis of tumors described in (q). Hematoxylin and eosin (H&E) staining (left), Ki67 IHC staining (middle) and Ki67 quantification (right) are shown. In (a-e, g-l, m-r) cells were treated or not with doxycycline to induce shRNA expression. In western blot analyses vinculin or actin were used as loading controls. In e, h,i, k, l, n, o, q, r data are expressed as mean ±SD. Each dot represents a biological replicate. Statistical significance was calculated using one-way ANOVA followed by Tukey's multiple comparison tests (e, h, i, n, o) or using Student's t test (k, l, q, r).

Extended Data Figure 2

Extended Data Figure 2. WDR5 loss, but not DPY30, broadly affect H3K4

methylation patterns.

a. Dot plots of BrdU incorporation assay of PATC53 cells silenced for WDR5 or

DPY30 (respectively shWDR5 and shDPY30) or control (shCTR). Cells were treated with

doxycycline for 48 hrs to induce shRNA expression. Data are expressed as mean ±SD,

each dot represents a biological replicate. Statistical significance was calculated using

one-way ANOVA. **b.** Western Blot analysis of whole lysate (upper panel) or histone

lysates (lower panel) in PATC53 cells, as described in a. **c.** Plots showing normalized

reads intensity (left) and the quantification (right) of ATACseq signals from three biological

replicates of PATC53 cells as described in a-b. **d.** Plots showing normalized reads

intensity (left) and the quantification (right) of H3K27ac signals from three biological

replicates of PATC53 cells in a-b. **e.** Plots showing normalized reads intensity (left) and

the quantification (right) of H3K4me3 signals from three biological replicates of PATC53

cells as described in a-b. **f.** Western Blot analysis of whole lysate (upper) or histone

lysates (lower) in KPC cells silenced/knock out for Dpy30 (respectively shDpy30 and

sgDpy30) or control (shCTR or sgCTR). Cells were treated with doxycycline or vehicle for

72 hrs to induce shRNA expression. **g.** Plot representing the top regulated pathways

resulting from genes having H3K4me3 peak loss in shWDR5 PATC53 cells compared to

controls (shCTR). In c-d, data in the violin plots are presented as median value.

Extended Data Figure 3

Extended Data Figure 3. DPY30 loss moderately upregulates inflammatory
pathways.

a. Heatmap showing Spearman correlation between samples and biological
replicates for bulk RNA sequencing (RNAseq) of PATC53 cells silenced for WDR5
(shWDR5) or DPY30 (shDPY30) or control (shCTR). **b.** Volcano plot representing the
GSEA results from bulk RNAseq data in PATC53 cells shWDR5 versus shCTR. **c.** Impact
of shCTR, shDPY30 or shWDR5 in PATC53 cells on *FEN1*, *GINS* and *RFC3* analyzed
by chromatin accessibility (ATAC), histone H3 methylation (H3K4me3), histone H3
acetylation (H3K27ac) and mRNA expression (RNA). **d.** Graphs reporting the mRNA
expression of *FEN1*, *GINS* and *RFC3* analyzed by quantitative PCR in shCTR, shDPY30
and shWDR5 PATC53 cells. **e.** Volcano plot representing GSEA from bulk RNAseq data
in PATC53 cells shDPY30 versus shCTR. **f.** Impact of shCTR, shDPY30 or shWDR5 in
PATC53 cells on *IFNL1*, *IRF-AS1* and *TRIM10* analyzed by ATAC, H3K4me3, H3K27ac
and RNA expression. **g.** Graphs reporting the mRNA expression of *IFNL1*, *IRF-AS1* and
TRIM10 analyzed as described in d. In b-c, top upregulated pathways are indicated. In d
and g, data are expressed as mean ±SD, each dot represents a biological replicate, and
statistical significance was calculated using one-way ANOVA.

Extended Data Figure 4

Extended Data Figure 4. The WRAD core interacts with the replisome machinery in a non-canonical manner.

a-c. Dot plots reporting pathway enrichment analyses of DPY30 interactors in human PATC66 (a), PATC53 (b), or mouse KPC (c) cells. Pathway enrichment is calculated using STRING online tool and Reactome module. **d.** Venn diagram showing the DPY30 common interactors among PATC53 (PDX2), PATC66 (PDX1), and KPC cells. **e.** Western Blot analysis of 293T cells co-overexpressing HA-tagged mouse DPY30 and EGFP-tagged mouse MCM members (MCM2-7). Control IgG and input were included. **f.** Contour plot showing the gating strategy for sorting PATC53 cells expressing FUCCI according to cell cycle phases. **g.** Western blot analysis of PATC53 cells sorted as in f. **h-i.** Bar plots showing the fold change enrichment of the epigenetic interactors (h) or replisome interactors (i) in DPY30 immunoprecipitates analyzed by mass spectrometry. Each dot represents a biological replicates and data are expressed as mean ± SD. **j.** Representative PLA staining detecting WDR5 interaction with WDHD1 and MCM6 in PATC66 cells. **k-m.** Western Blot analysis of the immunoprecipitation assay for endogenous WDR5 (k), Ash2L (l) and RbBP5 (m) in PATC53 cells. Control IgG and input were included. **n.** Western blot analysis of PATC53 cells expressing DPY30 wild type (WT) or mutant (L69D) isoforms. **o.** Western blot analysis of PATC53 cells expressing WDR5 wild type (WT) or mutant (L240K, V268E) isoforms. In (n, o), control IgG and input are shown, and H3 was used as a loading control.

Extended Data Figure 5

Extended Data Figure 5. DPY30 loss leads to DNA replication stress and DNA

damage.

a. Western blot analysis of proteins bound to nascent DNA retrieved using iPOND

or input control in KPC cells. Cells were treated as follows: control (negative control, no

biotin in the click reaction), EdU pulse (EdU treatment for 2, 5 or 10min), chase (EdU

treatment for 10 min, followed by a 60 min thymidine chase). Histone H3 was used as a

loading control. **b.** Western blot analysis of iPOND or input control in KPC cells, control

(sgCTR) or *Dpy30*-KO (sgDpy30). Cells were treated as described in a. Histone H3 was

used as a loading control. **c.** Schematic representation of the re-replication sequencing

(re-repli-seq) in KPC sg/ctr or sgDpy30 cells. **d-e.** Plots showing the genomic regions and

the ratio (score) of incorporated CldU and IdU in KPC sgCTR and sgDpy30 cells. **f.** Left,

panels show immunofluorescence images of KPC and KP$^{-/-}$C cells silenced for Dpy30

(shDpy30) or control (shCTR), treated with doxycycline for 96 hrs to induce shRNA

expression. In green DPY30, in red pS139 H2AX (γH2AX), in blue nuclei (DAPI). Right,

graphs report the percentage of γH2AX+ in KPC cells. **g.** Left, panels show

immunofluorescence staining of KPC and KP$^{-/-}$C cells, as described in f, stained for pS4-

8 RPA (green), γH2AX (red) and nuclei (blue). Right, graphs report the percentage of

pS4-8 RPA+ in each cell group. **h.** Immunofluorescence staining of sgDpy30 or sgCTR

KPC tumors grown in NSG (upper) or C57BL/6 (lower) mice and then stained for Ki67

(green), γH2AX (red), and nuclei (blue). In (f-g), data are expressed as mean ± SD and

each dot represents a different analyzed field. Statistical significance was calculated

using one-way ANOVA (f, g).

Extended Data Figure 6

Extended Data Figure 6. DPY30 loss leads to DNA damage and accelerated S-to-G2/M transition.

a. Immunohistochemical staining of γH2AX (left) and quantification (right) in control (sgCTR) or DPY30-KO (sgDPY30) PATC53 tumors grown in NSG (upper) or nude (lower) mice. **b.** Immunohistochemical staining of γH2AX (left) and quantification (right) in control (shCTR) or DPY30-silenced (shDPY30) PATC66 tumors grown in NSG mice. **c-d.** Volcano plots representing the GSEA results from bulk RNAseq data in KPC (c) or KP$^{-/-}$C (d) cells sgDpy30 versus sgCTR. **e.** Histograms showing DNA content in sgCTR or sgDpy30 KPC cells. Cells were collected in exponential growth (async) or synchronized (t0) in S phase using a double thymidine block and then released in complete medium for the indicated time. **f.** Bar graph showing cell cycle distribution in KPC cells as described in e. **g.** Dot plots (left) and quantification (right) of the BrdU incorporation assay in KPC cells described as in e. BrdU (APC) mean intensity is shown (A.U. = arbitrary unit). **h-i.** Western Blot analysis of sgCTR or sgDpy30 KPC (h) or KP$^{-/-}$C (i) cells, as described in e. Vinculin was used as loading control. **j.** Western Blot analysis of sgCTR or sgDpy30 KPC cells synchronized in S or G2/M phase. Vinculin was used as loading control. In (a, b, f, g), data are expressed as mean ±SD, each dot represents a biological replicate. Statistical significance was calculated using one-way student t test (a-b, g) or two-way ANOVA test (f).

Extended Data Figure 7

Extended Data Figure 7. DPY30 loss increases chromosomal instability in

PDAC.

a. Western Blot analysis of control (sgCTR) or *DPY30*-KO (sgDPY30) RPE-1 cells

treated with vehicle or MLN4924 for 24 hrs. **b.** Dot plots (left) and cell percentage (right)

of BrdU incorporation in RPE-1 cells as described in a. **c.** Western Blot analysis of sgCTR

and sgDpy30 KPC cells treated with vehicle or MLN4924 for 24 hrs. **d.** Histograms of

DNA content (left) and quantification of G2/M and aneuploid (>4N) cells (right) in KPC

cells, as described in c. **e. f.** Bar graphs reporting the mean percentage of aberrant mitosis

in control (shCTR) or Dpy30-silenced (shDpy30) KPC and KP$^{-/-}$C cells *in vitro*. Cells were

treated with doxycycline (dox) for 96 hrs to induce shRNA expression. **g.** Left, hematoxylin

and eosin (H&E) staining of mitotic sgCTR and sgDPY30 PATC53 cells grown in NSG

mice. Yellow arrows indicate chromosomal laggings. Right, bar graph reporting the

percentage of aberrant mitosis in PATC53 cells. **h.** Left, images of metaphasic spread in

shCTR or shDPY30 PATC66 cells treated with dox for 48 hrs to induce shRNA

expression. **i.** Left, images of metaphasic spread in sgCTR and sgDPY30 PATC53 cells.

Right, graph reporting the percentage of normal (yellow) or aberrant (grey) mitosis. In (a,

c), vinculin was used as loading control. In (h-i), red arrows indicate chromosome fusions

or breaks. In (b, d-g), data are expressed as mean ±SD, dots indicate biological replicates

and statistical significance was calculated using one-way ANOVA (b, e, f, g), two-way

ANOVA (d) or Chi-square test (h, i).

Extended Data Figure 8. In pancreatic ductal carcinoma, DPY30 loss has a
context-dependent phenotype.

a-c. Tumor volume of control (sgCTR) and Dpy30-KO (sgDpy30) KP$^{-/-}$C cells
injected subcutaneously into the flank of NSG (a), nude (b), or C57BL/6 (c) mice. **d.**
Weight (mg) of tumors described in a-c. **e.** Left, T2 Magnetic Resonance Imaging (MRI)
scans of sgCTR and sgDpy30 KP$^{-/-}$C tumors grown orthotopically in the pancreas of
C57BL/6 mice. Right, Kaplan-Meier survival curve of tumor-bearing mice. **f-h.** Tumor
volume (f, g) and weight (h) of control (shCTR) or Dpy30-silenced (shDpy30) KP$^{-/-}$C cells
injected subcutaneously into the flank of NSG (f) and C57BL/6 (g) mice. **i.** Hematoxylin
and eosin (H&E) staining of KP$^{-/-}$C tumors as described in f-h. **j.** Left, T2 Magnetic
Resonance Imaging (MRI) scans of shCTR or shDpy30 KP$^{-/-}$C tumors grown
orthotopically in the pancreas of C57BL/6 mice. Right, Kaplan-Meier survival curve of
tumor-bearing mice. **k-l.** Immunofluorescence staining of shCTR and shDpy30 KPC
tumors grown in NSG (k) or C57BL/6 (l) mice and then stained for αSMA, pS10-H3, and
nuclei. **m-n.** Volcano plots representing GSEA analyses from bulk RNAseq data in
sgDpy30 versus sgCTR KP$^{-/-}$C tumors grown in NSG (k) or C57BL/6 (l) mice. **o.**
Immunofluorescence staining of pS366 STING (green) and nuclei (blue) in sgCTR and
sgDpy30 KPC cells, in vitro. **p.** Masson's trichrome staining of sgCTR and sgDpy30 KPC
tumors grown in C57BL/6 mice. In (a-d, f-h, o), data are expressed as mean value ±SD.
Dots represent biological replicates. In (e, j), statistical significance and Hazard Ratio
(HR) were calculated using Log-rank test. Statistical significance was calculated using
Student t test (a-d) or one-way ANOVA (f-h, o). Scale bar = 200 μm (i, p), 100 μm (k, l),
50 μm (o).

Extended Data Figure 9

Extended Data Figure 9. Dpy30 knockout tumors display higher T cell infiltration and activation.

a. Bubble plot showing expression levels of genes selected for cluster annotation in single-cell RNA sequencing data from control (sgCTR) or Dpy30-KO (sgDpy30) KPC tumors grown in C57BL/6 mice. Size of dots represents the percentage of cells expressing the gene; color scale shows the average expression level. **b.** Bar plot showing the percentage of cell distribution in the different clusters (cell type) calculated using Seurat in sgCTR or sgDpy30 KPC tumors (n=3 per condition). **c-g.** Contour plots showing the flow cytometry gating strategy for detecting the NK or CD3+ NK cells (c), CD4+ conventional or Treg cells (d), CD8+ T cells (e), CD8/PD-1+ cells expressing Granzyme B (Gzmb, f), Interferon gamma (Infg, g) or Tumor Necrosis Factor (tnf, g). **h.** Violin plot showing the expression of T cell exhaustion marker (Tox) or cytotoxic markers *(Gzmb, Ifng, Nkg7)* in the T cell of sgCTR or sgDpy30 KPC tumors, calculated using Seurat. **i.** Bar plot showing the percentage of single nucleotide variants (SNVs) and indels in sgCTR or sgDpy30 KP-/-C tumors grown in NSG or C57BL/6 mice (n=3 or 4 tumors per group). Data are expressed as median (h) or mean value ±S.D. (i). Statistical significance was calculated using Student's t test (h) or a two-way ANOVA (i).

Extended Data Figure 10

1928 **Extended Data Figure 10. DPY30 correlates with tumor stage and predicts**

1929 **response to ICB treatment.**

1930   **a-c.** Kaplan–Meier survival curves of TCGA PDAC patients divided according to

1931 low- or high-expression of *WDR5* (a), *RBBP5* (b) and *ASH2L* (c). **d.** DPY30

1932 immunohistochemical staining in normal pancreas, pancreatic metaplasia and

1933 intraepithelial neoplasia (PanIN) and PDAC samples from KPC mouse model. **e.** Volcano

1934 plot representing GSEA in high-purity PDAC PDXs samples and represented as low

1935 versus high *DPY30* expression. **f.** Kaplan–Meier overall survival curve of patients in the

1936 PRINCE trial cohort divided according to low- or high-expression of *DPY30.* **h.** Violin plot

1937 showing the median expression of *DPY30* in the patience included in the analysis in f and

1938   Fig. 6h. In (a-c, f), statistical significance and Hazard Ratio (HR) were calculated using

1939   the Log-rank test.