**Vitamin E supplementation prevents ferroptosis in round spermatids of aged mice**

Jasper Germeraad, Takako Kikkawa, Noriko Osumi*

Department of Developmental Neuroscience, Tohoku University Graduate School of Medicine,

2-1, Seiryo-Machi, Aoba-ku, Sendai, Miyagi 980-8575, Japan

Running title: Ferroptotic cell death in the testes

Keywords**:** Spermatogenesis, cell death, ferroptosis

*Corresponding author

Noriko Osumi, PhD

Professor, Department of Developmental Neuroscience,

Tohoku University Graduate School of Medicine,

2-1 Seiryo-machi, Aoba-ku, Sendai 980-8575, Japan

Tel.: +81 22 717 8201. Fax: +81 22 717 8205.

E-mail address: noriko.osumi.c7@tohoku.ac.jp (N. Osumi)

Grant Sponsors: JSPS KAKENHI (#17K19372, 22K19322), The Canon Foundation

**Abstract**

Germ cell depletion in the aged testes has traditionally been attributed to removal by apoptosis.

This study aimed to determine whether ferroptosis, an alternative form of cell death driven by

iron-dependent lipid peroxidation, also contributes to germ cell loss in the lipid-rich

environment of the testis. Here, we demonstrate that pre-meiotic cells are eliminated via

apoptosis, whereas post-meiotic round spermatids (RSs) are mainly removed through

ferroptosis. Surprisingly, we detected a greater abundance of Y-chromosome-bearing RSs (Y-

RSs) than X-carrying RSs (X-RSs) in the aged testis, implying that X-RSs might be more prone

to ferroptosis. Young mice fed a vitamin E (VE) deficient diet recapitulated age-related

phenotypes, while VE supplementation prevented ferroptosis and promoted the survival of X-

RSs in aged mice. Overall, this study reveals that aging causes ferroptosis in RSs, specifically

impacting X-RSs, which can be prevented by VE supplementation, effectively reversing age-

induced deterioration and contributing to healthy testicular aging.

**Introduction**

In recent decades, a global demographic shift towards aging populations has resulted in delayed marriages and parenthood[1,2], placing increasing pressure on healthcare systems. These trends have contributed to higher risks of fertility challenges, pregnancy complications, and neurodevelopmental disorders in offspring. While the reproductive lifespan of females is biologically limited, male fertility, although maintained throughout life, experiences a gradual decline with age[3]. Numerous studies have documented age-related reductions in semen volume, sperm concentration and motility, as well as increased DNA fragmentation in sperm[4,5,6], which collectively contribute to decreased reproductive potential. As a result, preserving male reproductive health during aging has become increasingly important for couples seeking to conceive later in life.

Sperm cells, produced through spermatogenesis, are essential for sexual reproduction, as they carry either the X or Y chromosome, which determines the sex of the offspring. Spermatogenesis occurs within the seminiferous tubules of the testis and takes approximately 75 days in humans and 35 days in mice[7]. The process starts with spermatogonial stem cells that undergo asymmetric division to form spermatocytes. These spermatocytes then enter meiosis, producing haploid round spermatids (RSs), which elongate and condense their genetic material, ultimately developing into mature spermatozoa. While various studies have documented age-related decline in murine spermatogenesis[8,9,10], the full implications of these changes on reproductive health remain insufficiently understood.

Germ cell depletion is commonly described in the aged testis, typically attributed to apoptotic elimination triggered by oxidative stress[11,12]. However, we found a study reporting that spermatocytes and RSs may respond differentially to oxidative stress, particularly to lipid peroxides *in vitro*; while spermatocytes undergo apoptosis, RSs seem to die via ferroptosis[13]. Ferroptosis is a regulated form of cell death, distinct from apoptosis, characterized by iron-dependent oxidation of membrane phospholipids[14,15] and the failure of Glutathione peroxidase 4 (Gpx4) to neutralize lipid peroxides[16]. Notably, Gpx4 was identified as indispensable for preventing germ cell death already before the discovery of ferroptosis[17,18]. These findings suggest a previously undiscovered role of ferroptosis in the aging testis *in vivo*.

Vitamin E (VE), a group of fat-soluble micronutrients comprising of four tocopherols and four tocotrienols, is well-known for its antioxidant properties and its ability to prevent ferroptosis by inhibiting lipid peroxidation *in vitro*[19,20]. VE has also been shown to positively impact the

male reproductive system and fertility across various species *in vivo*[21,22]. Additionally, VE

deficiency (VED) has been reported in mice to cause alterations in the testis and epidydimis

that closely resemble aging[23]. These findings suggest a potential link between ferroptosis and

germ cell death during aging, which may be modulated by VE.

In this study, we demonstrate that premeiotic cells primarily undergo apoptosis, while RSs

predominantly initiate an alternative form of cell death, exhibiting key features of ferroptosis

as mice age. Suprisingly, we noticed an alteration in sex chromosome ratio among RSs of aged

mice, with a greater proportion of Y-chromosome bearing RSs (Y-RSs), likely due to increased

susceptibility of X-bearing RSs (X-RSs) to ferroptosis. Furthermore, VED was found to induce

ferroptosis and eventually alter the Y/X ratio, while VE supplementation reversed several

aging-related changes including improving RSs survival and restoring the YX ratio. Our

findings suggest that VE may hold clinical potential in mitigating aging-related testicular

decline by preserving RS viability.

**Results**

**Round spermatids predominantly undergo a form of cell death distinct from apoptosis**

We began by confirming aging-related phenotypic alterations in the testis of 18-month old mice.

As we expected, aged mice exhibited an increase in body weight (Fig. 1a) and a corresponding

decrease in testis weight (Fig. 1b), resulting in a lower gonadosomatic index[24] (testis

weight/body weight, GSI, Fig. 1c). Next, we assessed spermatogenesis within the seminiferous

tubules (Fig. 1d). In young mice, the tubules appeared morphologically intact, with a dense,

well-organized cell arrangement from the periphery to the lumen (Fig. 1e, upper panels). In

contrast, aged mice showed significant depletion of cells in both the peripheral and luminal

areas (Fig. 1e, lower panels). A detailed immunohistochemical germ cell analysis revealed a

marked reduction in the number of Type A spermatogonial cells (Type A Spg) in aged mice

compared to young mice (Fig. 1f), which extended to later stages of spermatogenesis, including

Type B spermatogonia (Type B Spg, Fig. 1g), meiotic pachytene spermatocytes (PSs, Fig. 1h),

and haploid RSs (Fig. 1i). Additionally, a decrease in the number of somatic Sertoli cells (SCs)

was observed in aged mice (Fig. 1j).

We next examined differential cell death mechanisms in young and aged testes by double

labeling with active caspase-3 (Casp3) and terminal deoxynucleotidyl transferase dUTP nick

end labelling (TUNEL). Casp3 is a marker specific for apoptosis, while TUNEL detects DNA

fragmentation associated with various forms of cell death, including ferroptosis[25]. Our analysis
clearly showed a higher percentage of tubules containing TUNEL-positive (TUNEL+) cells in
aged mice compared to young mice (32% in young vs 46% in aged, Figure 2a). We then
calculated the cell death index (CDI), defined as the average number of dying cells per tubule
(TUNEL+ cells/tubule number). The CDI was significantly elevated in aged mice (Fig. 2b).

We further characterized cell death in pre-meiotic cells, including spermatogonia and PSs,
which are located in the peripheral region of the seminiferous tubules (between the dashed and
solid lines), and in post-meiotic RSs situated in the luminal area (inside the solid line) (Fig. 2c).
Cell death within the tubules exhibited heterogeneity; some tubules contained only
Casp3+/TUNEL+ cells, indicating apoptotis in the peripheral area (closed arrowheads, Fig. 2d,
d′), while others displayed a mixture of Casp3+/TUNEL+ apoptotic cells and Casp3-/TUNEL+
non-apoptotic cells (open arrowheads, Fig. 2e, e′). In other cases, only Casp3-/TUNEL+ cells
were observed in both the peripheral and luminal areas (Fig. 2f, f′, g, g′). Interestingly, both
Casp3+/TUNEL+ and Casp3-/TUNEL+ cells were found in the peripheral area, but in the
luminal area, most dying cells were Casp3-/TUNEL+ (Fig. 2g, g′), with only few
Casp3+/TUNEL+ cells (Extended data 1).

We quantified these labeled cells in both regions of young and aged testes. In the aged testis,
there was a significant increase in Casp3+/TUNEL+ cells in the peripheral area compared to
the young testis (Fig. 2h), while the number of Casp3-/TUNEL+ cells remained unchanged
(Fig. 2h). In the luminal area, the number of Casp3+/TUNEL+ RSs was consistently low, with
no significant difference between age groups (Fig. 2i). However, aged mice showed a notable
increase in Casp3-/TUNEL+ RSs in aged mice (Fig. 2i). These findings suggest that aging
primarily induced apoptosis in pre-meiotic cells, whereas non-apoptotic cell death
predominated in post-meiotic RSs. This reveals a novel aspect of testicular degeneration due
to aging, highlighting a shift from apoptosis to non-apoptotic cell death in RSs.

**126 Round spermatids in aged mice exhibit hallmarks of ferroptosis**
We hypothesized that aging-related oxidative stress may trigger ferroptosis in RSs *in vivo*.
Therefore, we aimed to characterize ferroptosis in RSs at the molecular level by isolating RSs
from whole testes (Fig. 3a). Using a modified density gradient method[26], we collected fractions
that were microscopically assessed for round spermatid (RS) enrichment using PNA staining.
Fractions with >80% RSs purity were deemed sufficient for the following analyses (Fig. 3b).
With these RS-enriched fractions, we examined key molecular markers of ferroptosis. We

133 hypothesized that aging would result in a reduced levels of antioxidant enzyme Gpx4, along

134 with increased lipid peroxidation of polyunsaturated fatty acids (PUFAs) mediated by long-

135 chain-fatty-acid-CoA ligase 4 (*Ascl4*), and arachidonate 15-lipoxygenase (*Alox15*) (Fig. 3c).

136 We first histologically confirmed the expression of Gpx4 in the cytoplasm of RSs and in the

137 tail region of sperm, while weak expression was observed in spermatocytes. However, no

138 obvious difference between young and aged testes was observed (Fig. 3d). Gpx4 exists in three

139 isoforms: nuclear (nGpx4), mitochondrial (mGpx4), and cytosolic (cGpx4). Previous studies

140 have reported that while both nGpx4 and mGpx4 are indispensible for spermatogenesis, cGpx4

141 plays a crucial role in protecting germ cells from cell death[17,27]. We therefore sought to

142 determine which isoforms are affected by aging. Western blot analysis using an anti-Gpx4

143 antibody that recognizes both mGpx4 and cGpx4 isoforms, revealed no significant difference

144 in Gpx4 protein levels (approximately 17 kDa) between young and aged RS-enriched fractions

145 (Fig. 3e). To further assess isoform-specific expression at the mRNA level, we conducted

146 quantitative PCR (qPCR). In aged RSs, we found that the expression levels of both *nGpx4* and

147 *mGpx4* remained unchanged (Fig. 3f, g). However, *cGpx4* expression was significantly reduced

148 in aged RSs (Fig. 3h).

149 Next we examined lipid peroxidation in RSs. Quantification of *Acsl4* and *Alox15* expression in

150 RS-enriched fractions from young and aged testes revealed a significant increase in aged RSs

151 (Fig. 3i, j). Since Acsl4 and Alox15 promote PUFA oxidation, leading to the production of

152 aldehydes like 4-Hydroxynonenal (4HNE), a key marker of lipid peroxidation, we quantified

153 4HNE levels by Western blot analysis. This revealed a significant increase in the 4HNE levels

154 in RS-enriched fractions from the aged testes (Fig. 3k). These findings indicate that aged RSs

155 accumulate lipid peroxides, as evidenced by the upregulation of *Acsl4* and *Alox15*, along with

156 increased 4HNE levels. The concurrent reduction in *cGpx4* expression suggests a diminished

157 capacity to neutralize reactive oxygen species (ROS), which may promote ferroptosis as a

158 distinct cell death pathway in RSs, contrasting to the apoptotic mechanism observed in

159 spermatogonia and spermatocytes.

160 We then investigated potential differences in cell death vulnerability between X-RSs and Y-

161 RSs. During spermatogenesis, X-RSs and Y-RSs are theoretically produced in equal propotions

162 through meisosis. However, using fluorescent *in situ* hybridization (FISH), we unexpectedly

163 discovered that the sex chromosome ratio in the aged testis was skewed towards Y-RSs

164 compared to the young testis (young: 1.02 vs aged: 1.11; *P* <0.001, Fig. 3l). This finding

165 suggests that aging may selectively impact the survival of X-RSs. Given the established

relationship between reduced *cGpx4* levels, lipid peroxidation and ferroptosis, it is plausible
that X-RSs are more susceptible to ferroptosis. This increased vulnerability could lead to the
observed skewed Y/X ratio in the aged testis.

**Vitamin E ensures RSs survival and restores the Y/X ratio alteration**

Given the observed susceptibility of RSs to oxidative stress and ferroptosis, we next
investigated whether varying dietary levels of VE, a potent antioxidant, could mitigate these
age-related changes in RS survival and restore the skewed Y/X ratio. To simulate the effect of
aging, we fed young mice either a control diet (YC) or VE-deficient diet (YVE-), while to
counteract aging effects, we provided aged mice either a control diet (AC) or VE-supplemented
diet (AVE+) (Fig. 4a).

We first assessed whether VE- or VE+ diets would result in systemic health abnormalities by
measuring bodyweight. While body weight naturally increased over time across all groups, it
remained unaffected by dietary changes (Extended Data Fig. 2a, b). Similarly, testis weight
and GSI showed no significant alterations under these dietary conditions (Extended Data Fig.
2c, d). However, the progression of spermatogenesis was influenced by VE modulation (Fig.
4b, Extended Data Fig. 2e). Cell-count analyses revealed an age-related decline in Type A and
B Spg in both AC and AVE+ groups (Extended Data Fig. 2f, g). PSs showed a moderate
sensitivity to VE levels, exhibiting a slight but non-significant decrease in the YVE- group and
a minor increase in the AVE+ group (Extended Data Fig. 2h). In contrast, RSs were highly
responsive to VE modulation, demonstrating significant reductions in the YVE- group and,
conversely, partial recovery in the AVE+ group (Fig. 4b, c). These findings suggest that VE
plays a crucial role in modulating spermatogenesis, particularly by mitigating age-related
declines in RSs and enhancing their recovery under VE+ conditions.

Analyzing cell death under the influence of VE (Fig. 4d, Extended Data Fig. i), we found that
the testes of YVE- and AC mice had the highest percentage of tubules containing TUNEL+
cells, while the percentage was reduced in AVE+ mice (YC: 28%, YVE-: 40%, AC: 41%,
AVE: 30%, Fig. 4e). Similar trends were observed for the CDI (Fig. 4f). In the peripheral region
of the testis, neither VE- nor VE+ significantly affected the number of Casp3+/TUNEL+ cells
(Fig. 4g) or Casp3-/TUNEL+ cells (Extended Data Fig. 2i, j). However, in the lumen, RSs were
notably impacted; YVE- led an increase in Casp3-/TUNEL+ RSs, while AVE+ reduced the
number of these cells (Fig. 4h). Casp3+/TUNEL+ RSs remained rare and unchanged across all
groups (Extended Data Fig. 2i, k). These observations imply that while VE modulation has

minimal effects on apoptosis in peripheral testis cells, it significantly influences RS survival,

with VE- increasing and VE+ reducing cell death.

We next assessed whether VE modulation affected the molecular hallmarks of ferroptosis in

RSs. No significant changes were observed in Gpx4 protein levels across the groups (Fig. 4i)

nor in mRNA levels for *nGpx4*, or *mGpx4* (Fig. 4j, k). However, *cGpx4*, showed a slight, yet

non-significant reduction in YVE- group, while no changes in *cGpx4* mRNA levels were

observed in AVE+ compared to AC (Fig. 4l). In terms of lipid peroxidation markers, there was

a marginal, but non-significant, increase in *Acsl4* mRNA levels in YVE-, with no changes

observed in AVE+ (Fig. 4m). *Alox1*5 expression remained unaffected in both YVE- and AVE+

groups (Fig. 4n). Notably, however, 4HNE levels were significantly altered; they were elevated

in YVE- group and dramatically reduced in AVE+ compared with AC (Fig. 4o). These results

indicate that while VE modulation does not significantly affect a key ferroptosis-related

enzyme like Gpx4, it has a pronounced impact on lipid peroxidation, as evidenced by the

marked changes in 4HNE levels, suggesting a protective role of VE against oxidative damage

in RSs.

Lastly, we evaluated the effect of VE on the Y/X ratio. As expected, the Y chromosome became

more dominant in the RSs of the YVE- group (YC: 1.02 vs YVE-: 1.07; *P*=0.023). On the

contrary, the AVE+ group show a marked decrease in the Y/X ratio compared to AC, resulting

in a ratio similar to the young control condition (AC: 1.10 vs AVE+: 1.01, *P*= <0.001, Fig. 4p).

Taken together, these data suggest that VE- accelerates testicular aging, as evidenced by

increased cell death, oxidative stress, and Y-chromosome dominance, while VE+ in aged mice

partially reversed these aging effects by improving RS survival, reducing lipid peroxidation,

and restoring the Y/X ratio, potentially by lowering the risk of ferroptosis in X-RSs.

222
223
**Discussion**

In humans, aging is associated with significant changes in the testes, including reductions in

testicular volume and weight[28,29]. Similar aging effects have been reported in rodent models by

us and others[10,30]. The observed reduction in testicular weight is largely attributed to the loss

of both germ cells and Sertoli cells, resulting from from impaired spermatogenesis[31,32,33]. In

aged mice, we observed a decline in the number of Type A Spg and Type B Spg, PSs, RSs, and

SCs, all of which likely contribute to the noted loss in testicular weight.

In this study, we focused on aging-induced cell death during spermatogenesis. Previous studies

have indentified aging-related apoptosis in the murine testis[34,35], consistent with findings in

other tissues[36,37]. Our analysis of young and aged testes revealed that different germ cell types

initiate distinct forms of cell death. Specifically, our results suggest that pre-meiotic male germ

cells predominantly undergo apoptosis, whereas post-meiotic RSs display a preference for

ferroptosis. This distinction aligns with findings from a previous *in vitro* study[13]. Furthermore,

research from our laboratory, investigating miRNA expression in sperm from aged mice, has

revealed a significant upregulation of miRNAs that target both apoptosis and ferroptosis[38]. This

suggests that cells at different stages of development posess the necessary machinery to initiate

distinct cell death pathways in response to alterations in the aging testicular environment.

A previous ultrastructural and histological study using electron microscopy, toluidine blue and

acridine orange (dyes to label different types of cell death), has suggested that RSs rarely initate

apoptosis in normal, healthy testes[39]. In our study, we observed many RSs positive for TUNEL

but negative for Casp3. This suggests that RSs in aged mice appear to favor a non-apoptotic

cell death pathway. We indeed identified a combination of molecular hallmarks of ferroptosis

in RSs of aged mice, indicating that this might be the predominant cell death modality in these

cells. Specifically, we observed increased mRNA levels of *Acsl4* and *Alox15*, along with

reduced *cGpx4*, leading to elevated lipid peroxidation, as evidenced by higher 4HNE levels.

These lines of evidence collectively align with the characteristics of ferroptosis observed in

somatic cell lines[16,40], implying that aging may sensitize RSs to this form of cell death. This

heightened vulnerability may be partly explained by an increased incorporation of arachidonic

acid-derived PUFAs into membranes of RSs compared to other germ cells[41], making them

more susceptible to aging-related oxidative stress.

Mammalian Y-chromosomes are essential for sex determination in offspring and are expected

to be present in a 1:1 ratio to X-chromosomes in haploid RSs. Surprisingly, we detected a

higher number of Y chromosomes than X chromosomes in the aged RS population, shifting

the ratio (young: Y/X 1:02, *n=13,303, N=6* vs aged: Y/X: 1.11, *n=13,008, N=6)*. To our

knowledge, this is the first study that investigated the skewed sex chromosome ratio in aged

RSs. A previous study examined the Y/X ratio in RSs in young (8-week old) mice and found

no difference between Y and X chromosome numbers (Y/X: 1.02, *n=2,029*)[42], which aligns

with our observations in young (12-week old) mice. Other studies have focused on the Y/X

ratio of mature sperm in healthy men, with mixed results: some reported an excess of X

chromosomes[43], others found no difference[44], while another observed a predominance of Y

chromosomes[45]. Whether the aging-altered Y/X ratio in RSs persists in mature sperm remains
to be investigated.

We present compelling evidence that X-RSs are more vulnerable to aging-related
environmental changes than Y-RSs, suggesting that X-RSs may selectively be eliminated. The
reason why X-RSs preferentially die, potentially via ferroptosis, remains unclear. One possible
explanation is that X-RSs are more susceptible to lipid peroxidation because *Acsl4*, which we
found to be upregulated in aged RSs, is located on the X chromosome[46]. In addition, during
spermatogenesis, round spermatids remain physically bound through cytoplasmic bridges,
facilitating the sharing of gene products to maintain transcriptional diploidy. However, some
gene transcripts have been observed to not be equally shared, suggesting the potential for allelic
bias in gene expression[47]. Upon reanalyzing their data[47], we found that several genes involved
in ferroptosis, including *Lpcat3* and *Dhr7*, are only partially shared, with an allelic bias
favoring X-RSs. These findings imply that lipid peroxidation levels might naturally be higher
in X-RSs, and with the age-related reduction of Gpx4, X-RSs might be more susceptible to
ferroptosis in response to oxidative stress.

Our observations in VE- recapitulated the characteristics of aged testes, a finding consistent
with a previous report that studied the influence of VE- in mice testes at the morphological
level[23]. In contrast, VE+ mice showed partial restoration of certain aging parameters,
suggesting that lipid-related oxidative stress may play a key role in modulating aging-related
changes. Specifically, our findings suggest that VE modulation primarily affects RSs, as shown
by changes in their cell number in both VE- and VE+ groups, while other cell types were
largely unaffected. In line, a previous study showed that rats fed VE- diet from postnatal day
10 (P10) and sacrificed at P42 (before completion of the first spermatogenic wave), halted
spermatogenesis at the RS stage, while other germ cells remained present[48]. We observed that
spermatogenesis continued beyond the RS stage, likely because our VE dietary intervention
was initiated in sexually mature mice and the feeding period (10 weeks, covering two cycles
of spermatogenesis) was not sufficient to induce complete arrest. Nonetheless, these lines of
evidence underscore the critical role of VE in survival of RSs.

We found that VE modulation may influence the initiation of ferroptosis independently of
*cGpx4, Acsl4* and *Alox15* function, as evidenced by its impact on 4HNE levels in both VED
and VE+ conditions. Other studies have shown that Gpx4 and VE cooperatively protect
regulatory T cells and hematopoietic stem cells from ferroptosis[49,50], suggesting that interaction
between Gpx4 and VE may be cell-type or tissue-specific. Regarding the vulnerability of

ferroptosis in RSs, VE plays a critical role in supporting the survival of RSs, particularly X-
RSs, by preventing the initiation of ferroptosis through neutralization of lipid peroxyl radicals
in aged mice.

Overall, our data suggest that VE supplementation may help prevent aging-related changes in
the testis, with notable benefits for RSs. It is reported that VE supplementation can potentially
enhance the success of a *in vitro* fertilization technique by injecting RSs for men with complete
meiotic arrest[51]. Furthermore, it has recently been reported that overall sperm count worldwide
decreased in men from 1973 to 2018, even after accounting for confounders, including age[52,53].
As such, VE supplementation might be beneficial not only for aged men but also for young
men. However, our study has a few limitations. We do not present direct evidence that
ferroptosis induces the skewed Y/X ratio in RSs, nor do we have data on the long-term effects
of VE supplementation in clinical applications, including optimal concentrations or ideal
delivery methods. Additionally, it remains unclear whether the altered Y/X ratio persists in
mature sperm or if VE supplementation can influence live birth ratios in aged mice. These
questions merit further investigation.

**Methods**
**Animals**

C57BL/6J mice were obtained from Charles River laboratories and were bred and maintained in the Animal Experiment Facility at the Tohoku University Graduate School of Medicine. Mice were sacrificed at 3 months old (Young) or 18 months old (Aged). All animals were housed in standard cages in a temperature- and humidity-controlled room on a 12-hour light/dark cycle (light on at 8 am) and had access to food and water *ad libitum*. All experimental procedures were approved by the Ethics Committee for Animal Experiments in Tohoku University (#2023-MED03701) and the animals were treated according to the National Institutes of Health (NIH) guide for the care and use of laboratory animals.

**Testis collection**

Animals were sacrificed by short sedation with isoflurane (MSD Animal health) and cervical dislocation. Testes were isolated from the peritoneal cavity and separated from the surrounding fat tissue and epididymis. Testis samples were fixed in 4% paraformaldehyde in phosphate buffered saline (PBS, 4% PFA, Nacalai-Tesque) for 2 hours at 4°C, cut in half, and fixed for an additional 18 hours. Cryopreservation was executed as follows: the testes samples were immersed, sequentially, in 10% sucrose dissolved in PBS for 1 hour and 20% sucrose in PBS overnight, both at 4°C. After cryopreservation, samples were frozen in Tissue-Tek optimal cutting compound (OCT, Sakura Finetek) on a metal pedestal using liquid nitrogen and stored in -80°C until further use. Cryosections (10μm) were obtained using a cryostat (Leica CM3050 S) and used for histological analyses.

**Immunohistochemistry (IHC)**

Testicular sections on slide glasses (Matsunami) were immunostained using antibodies targeting anti-Plzf (1:500, sc28319, Santa Cruz), anti-Scp3 (1:500, ab97672, Abcam), anti-Kit (1:500, D13A2, Cell Signalling technologies), anti-Sox9 (1:900, 5535, EMD Millipore), and anti-Gpx4 (1:1000, ab125066, Abcam). Sections were washed using Tris-buffered saline containing 0.1% Tween20 (TBST). Antigen retrieval was performed (except for anti-cKit), for 10 minutes in 0.01 M citrate buffer (pH 6.0) at 95°C. Afterwards, slides were washed 3 times for 10 min-each with TBST, and subsequently blocked with 3% Bovine serum albumin (BSA) and 0.3% Triton X-100 in PBS for 1 hour at room temperature (RT). The slides were then incubated with primary antibodies diluted in blocking solution overnight at 4°C, washed again with TBST and incubated for 1 hour at RT with 4', 6-diamino-2-phenylindole (DAPI, 1:1000,

Sigma) as well as secondary antibodies (1:500, Anti-mouse alexa 488 or anti-rabbit Cy3,

Jackson ImmunoResearch) or peanut agglutinin (PNA-Cy5, 1:2000, Vector Laboratories)

when staining the acrosome for round spermatid identification. The sections were washed again

with TBST and mounted with Prolong Glass Antifade Mountant (Thermofisher). All images

were obtained using a confocal laser-scanning microscope (LSM800, Carl Zeiss), and

subsequently processed in Fiji (ImageJ).

**Fluorescent *in situ* hybridization**

FISH was performed as described previously[54]. In short, slides with frozen sections of the testis

were subjected to antigen retrieval in Histo-VT One (Nacalai-Tesque) for 30 minutes at 90˚C

and denatured using 50% formamid/2x saline-sodium citrate buffer (SSC), followed by

dehydration using 70% and 100% ethanol. A custom mixture of X (Conjugated with Cy3) and

Y (conjugated with FITC) chromosome probes (Chromosome Science Labo) was applied

subsequently. The slides were covered with hybridization cover (Hybrislip, Thermofisher) and

incubated at 80˚C for 10 minutes. Hybridization occurred overnight in a humid chamber at

37˚C. IHC was performed to enhance probe signals. Blocking was performed using 5%

Blocking One solution (Nacalai-Tesque) for 1 hour at RT. Primary antibodies (anti-FITC,

1:2000, A889, Invitrogen and Anti-Cy3, 1:2000, sc-166894, Santa Cruz) were diluted in 5%

Blocking One solution and incubated 1 hour at RT. After washing with 4x SSC, secondary

antibodies (1:2000, Anti-rabbit alexa 488 and anti-mouse Cy3, Jackson ImmunoResearch)

together with HOECHST-44432 (1:5000, Thermofisher) were applied and incubated for 1 hour

at RT. Slides were mounted with Prolong Glass Antifade Mountant (Thermofisher).. All

images were obtained using a fluorescence microscope (Keyence BZ-X series) and processed

in Fiji. RSs were identified according to nuclear morphology and location within the tubule.

Hybdridization efficiency was determined by microscopic evaluation and images were deemed

appropriate for analysis when in more than 95% of cells, a sex chromosome could be assigned.

**Cell death detection**

Apoptotic or non-apoptotic cell death was identified by double labeling for terminal

deoxynucleotidyl transferase dUTP nick end labelling (TUNEL, Merck Millipore) and

subsequently for an antibody against active-caspase-3 (1:500, 559565, BD Biosciences).

TUNEL labeling was performed first and according to the manufacturers' instructions. In short,

slides with testis sections were washed 1x 5 min in PBS and subjected to antigen retrieval by

microwaving 5 minutes in 0.01M citrate buffer. TdT mixture was applied and incubated at

37˚C for one hour followed by incubation with an anti-digoxigenin antibody conjugated with

fluorescein at RT for 30 min. Next, active-caspase-3 IHC was performed. Slides with testicular

sections were covered in blocking solution (3% BSA and 0.3% Triton X-100 in PBS) at RT for

one hour followed by primary antibody incubation at RT for one hour. Slides were washed

three times for five min in PBS and incubated with a secondary antibody (1:500, Anti-rabbit-

Cy3, Jacksonimmunoresearch) and HOECHST-44432 (1:5000) for one hour at RT. The slides

were washed again in PBS and mounted with Prolong Glass Antifade Mountant (Thermofisher).

and finally analysed with a confocal laser-scanning microscope (LSM800, Carl Zeiss). Using

Fiji, male germ line cells in the periphery or in the region towards the lumen of the seminiferous

tissue were independently quantified.

**Round spermatid isolation**

Round spermatid isolation was performed according to a previously published method[26].

Briefly, fresh testicular tissue was harvested from young or aged male mice followed by

removal of the epididymis and tunica albuginea. A single cell suspension was obtained via

serial digestion with Collagenase IV (1mg/ml, Sigma) and Trypsin (0.6 mg/ml, Worthington)

/ DNAse I (>3.2 ku/ml, Sigma) dissolved in 1x Krebs bicarbonate buffer (containing 1.2 mM

$KH_2PO_4$, 0.35 mM MgSO4, 1.7 mM $CaCl_2$, 4.7 mM KCl & 2.5 mM $NaHCO_3$) . The cell

suspension was loaded on a 1-5% BSA / Krebs buffer gradient and allowed to sediment for 1.5

hour on ice at 1 unit gravity. Twenty-eight fractions of 1 mL were collected and analysed for

purity using PNA (1:2000, Rhodamine, Vector Laboratories) and DAPI (1:1000, Sigma). Only

fractions #4-7 were routinely enriched with RSs (>80%). Therefore, in subsequent isolations,

fractions #4-7 were collected and pooled. The fractions were washed with Krebs buffer to

remove excess BSA and immediately used for RNA and protein isolation.

**Simultaneous RNA and Protein isolation**

The simultaneous isolation of RNA and Protein was carried out using the AllPrep

DNA/RNA/Protein mini kit (Qiagen) according to the manufacturers protocol with minor

modifications. All reagents and spin columns are supplied by the AllPrep mini kit (except for

Sodium dodecyl sulfate (SDS) buffer). In short, tissue was lysed in RLT buffer containing 1%

β-mercapto-ethanol (Sigma). The lysate was applied to the DNA spin column to remove DNA.

The flow-through was subsequently mixed with 100% ethanol and transferred to the RNA

binding column spin column. After centrifugation, flow-through was kept aside for protein

isolation while RNA was bound to the column. The column was subjected to washes with RW1

and RPE buffers, after which RNA was eluted using RNAse free water and assessed for

concentration and purity using the Nanodrop (Thermofisher). Isolated RNA was stored at -

80˚C until further used for complementary DNA (cDNA) synthesis. To the flow-through

containing protein, buffer APP was added and incubated for 10 minutes at RT. The sample was

pelleted, and 70% ethanol was added. The pellet was air dried and then vortexed for 1 minute

in SDS buffer containing 4% SDS, 20% sucrose, 10 mM Tris-HCL pH 6.8 and 1x

ethylenediaminetetraacetic acid (EDTA) free protease inhibitor (Roche) followed by

incubation at 37˚C for 30 minutes. The protein sample was vortexed for 5 minutes and was

next pelleted via centrifugation. Supernatant was transferred to a new tube and protein

concentration was determined using the bicinchoninic acid assay (BCA, Nacalai-Tesque).

Protein samples were stored in -80˚C until used for western blot.

**cDNA synthesis and quantitative polymerase chain reaction (qPCR)**

RNA isolated from pooled round spermatid fractions (1µg) was reverse transcribed into

cDNA using the SuperScipt III™ First-Strand Synthesis System for RT-PCR (18080-400,

Invitrogen). cDNA was subsequently used for qPCR using Taq Pro Universal SYBR qPCR

Master Mix (Vazyme). Primer sequences used in this study can be found in Supplementary

Table 1. For *Gpx4,* 3 primer sets were used to quantify the isoforms with a set targeting

*nGpx4* and *mGpx4* individually and a set that targeted both *mGpx4* and *cGpx4.* Cytosolic

*Gpx4* was quantified by subtracting Δct values of mitochondrial *Gpx4* from cytosolic +

mitochondrial *Gpx4*. All Values were normalized against *ActB* as a reference gene using the

$2^{-\Delta\Delta ct}$ method.

**Western blot**

Protein samples (5 µg) were separated by SDS-PAGE at 150V for 50 minutes (12%, TGX™

FastCast™ Acrylamide Kit; Bio-Rad). Primary antibodies targeting Gpx4 (1:1000, ab125066,

Abcam), 4-hydroxynonenal (4HNE, 1:1000, ab46545, Abcam), ActB (1:2500, sc-47778, Santa

Cruz), and Gapdh (1:4000, ab8245, Abcam) were used. Proteins were transferred onto 0.2 µm

polyvinylidene difluoride membranes (PDVF) via turboblot transfer (Trans-Blot Turbo Mini

PVDF Transfer Packs, Bio-rad), at 2.5A, 25V, 3 min. The membranes were then blocked in

10% Li-cor Intercept TBS blocking buffer (Licor) for 1 hour (Gpx4 & ActB) or in 5% blocking

One (Nacalai-Tesque) for 30 min (4HNE & Gapdh), and incubated with primary antibodies

overnight at 4 ˚C. The membrane was washed with Tris-buffered saline with 0.1% Tween-20

(TBST) and incubated with a secondary antibody (Gpx4 1:10000 or 4HNE 1:10000, W4011,

Promega and Gapdh 1:10000 or ActB 1:5000, W4021, Promega), diluted in 10% TBS blocking

buffer (Licor) or 5% blocking One (Nacalai-Tesque) for 1 hour at RT. The membranes were

then washed with TBST and visualized using the Bio-rad ChemiDoc Touch Gel Imaging

system and quantified relative to ActB for Gpx4 and Gapdh for 4HNE using Fiji.

**Vitamin E feeding**

For this experiment, young and aged C57BL/6J mice were purchased from Jackson laboratories

(JAX, Japan) as age-matched mice were available at the time of purchase. Differences in

bodyweight and testis were assessed and accounted for. Diets were custom-made and ordered

from Funabashi Farm, Japan. Young mice were fed control (containing 135 mg/kg VE), or

vitamin E deficient diet (containing >0.1 mg/kg VE). Aged mice were maintained on control,

VE-, or VE+ diet (containing 500 mg/kg VE). Detailed ingredients are listed in Supplementary

Table 2. All mice were raised on these diets for 10 weeks (70 days), covering two full cycles

of spermatogenesis (35 days)[7]. Young and aged mice were sacrificed at three and 18 months

of age, respectively. Testes were extracted as described with minor modifications: one testis

was cut in half, and one half was fixed using 4% PFA, cryopreserved and sectioned (10µm)

using a Leica 3050S cryostat as described previously. RSs were isolated from the remaining

testicular tissue (one and a half), after which RNA and protein were isolated simultaneously

from RS fractions. Testicular sections and RNA/protein were prepared for histology, qPCR

and Western blot experiments as described above.

**Statistical analysis**

Statistical information for individual experiments is shown in corresponding figure legends.

Values are always presented as mean ± standard deviation. Data were first analysed for

normality by Shapiro-wilk test. Statistical comparisons between groups were then analysed for

significance by Student's *t*-test, Mann-Whitney U test, One-way ANOVA, followed by

Tukey's multiple corrections, Kruskal-Wallis test followed by Dunn's multiple corrections, or

Mixed-Model (Fixed effects: diet, week and bodyweight, Random effects: individual mice,

using repeated measures). Results were considered significant at $P < 0.05$. Statistical analyses

were conducted using GraphPad Prism 10 (GraphPad Software).

**Declaration of interest**

The authors declare no conflict of interest.

**Acknowledgements**

We are grateful for the financial support from Japanese Society for the Promotion of Science

Kakenhi grant number #17K19372 and #22K19322, The Canon Foundation awarded to N.O,

and The establishment of university fellowships towards the creation of science technology

innovation, grant number JPMJFS2102, JPMJSP2114 awarded to J.G. We thank Ms. Sayaka

Makino for animal care and all other members of the Developmental Neuroscience laboratory

at Tohoku University for their useful comments.

**Author contributions**

Study design: J.G, TK, N.O; Data collection: J.G, T.K ; Data analysis: J.G; Drafted the

manuscript: J.G; Edited the manuscript: N.O, T.K; Funding acquisition: N.O, J.G.

**References**

1. Raymo, J. M. The second demographic transition in Japan: a review of the evidence. *China Popul Dev Stud* **6**, 267–287 (2022).

2. Schmidt, L., Sobotka, T., Bentzen, J. G. & Nyboe Andersen, A. Demographic and medical consequences of the postponement of parenthood. *Hum Reprod Update* **18**, 29–43 (2012).

3. Martins da Silva, S. & Anderson, R. A. Reproductive axis ageing and fertility in men. *Rev Endocr Metab Disord* **23**, 1109–1121 (2022).

4. Johnson, S. L., Dunleavy, J., Gemmell, N. J. & Nakagawa, S. Consistent age-dependent declines in human semen quality: A systematic review and meta-analysis. *Ageing Res Rev* **19**, 22–33 (2015).

5. Verón, G. L. *et al.* Impact of age, clinical conditions, and lifestyle on routine semen parameters and sperm kinematics. *Fertil Steril* **110**, 68-75.e4 (2018).

6. Evenson, D. P., Djira, G., Kasperson, K. & Christianson, J. Relationships between the age of 25,445 men attending infertility clinics and sperm chromatin structure assay (SCSA®) defined sperm DNA and chromatin integrity. *Fertil Steril* **114**, 311–320 (2020).

7. Griswold, M. D. Spermatogenesis: The Commitment to Meiosis. *Physiol Rev* **96**, 1–17 (2016).

8. Gosden, R. G., Richardson, D. W., Brown, N. & Davidson, D. W. Structure and gametogenic potential of seminiferous tubules in ageing mice. *Reproduction* **64**, 127–133 (1982).

9. Wright, W. W., Fiore, C. & Zirkin, B. R. The effect of aging on the seminiferous epithelium of the brown Norway rat. *J Androl* **14**, 110–7 (1993).

10. Silva, J. V. *et al.* Effects of Age and Lifelong Moderate-Intensity Exercise Training on Rats' Testicular Function. *Int J Mol Sci* **23**, 11619 (2022).

11. Asadi, A., Ghahremani, R., Abdolmaleki, A. & Rajaei, F. Role of sperm apoptosis and oxidative stress in male infertility: A narrative review. *International Journal of Reproductive BioMedicine (IJRM)* (2021) doi:10.18502/ijrm.v19i6.9371.

12. Gunes, S., Hekim, G. N. T., Arslan, M. A. & Asci, R. Effects of aging on the male reproductive system. *J Assist Reprod Genet* **33**, 441–454 (2016).

13. Bromfield, E. G. *et al.* Differential cell death decisions in the testis: evidence for an exclusive window of ferroptosis in round spermatids. *MHR: Basic science of reproductive medicine* **25**, 241–256 (2019).

14.    Halliwell, B. & Gutteridge, J. M. C. *Free Radicals in Biology and Medicine*. (Oxford
University Press, 2015). doi:10.1093/acprof:oso/9780198717478.001.0001.

15.    Gao, M., Monian, P., Quadri, N., Ramasamy, R. & Jiang, X. Glutaminolysis and
Transferrin Regulate Ferroptosis. *Mol Cell* **59**, 298–308 (2015).

16.    Dixon, S. J. *et al.* Ferroptosis: An Iron-Dependent Form of Nonapoptotic Cell Death.
*Cell* **149**, 1060–1072 (2012).

17.    Schneider, M. *et al.* Mitochondrial glutathione peroxidase 4 disruption causes male
infertility. *The FASEB Journal* **23**, 3233–3242 (2009).

18.    Imai, H. *et al.* Depletion of Selenoprotein GPx4 in Spermatocytes Causes Male
Infertility in Mice. *Journal of Biological Chemistry* **284**, 32522–32532 (2009).

19.    Friedmann Angeli, J. P. *et al.* Inactivation of the ferroptosis regulator Gpx4 triggers
acute renal failure in mice. *Nat Cell Biol* **16**, 1180–1191 (2014).

20.    Agmon, E., Solon, J., Bassereau, P. & Stockwell, B. R. Modeling the effects of lipid
peroxidation during ferroptosis on membrane properties. *Sci Rep* **8**, 5155 (2018).

21.    Muhammad, Z. Effects of dietary vitamin E on male reproductive system. *Asian Pacific*
*Journal of Reproduction* 145–150 (2017) doi:10.12980/apjr.6.20170401.

22.    Zhou, X., Shi, H., Zhu, S., Wang, H. & Sun, S. Effects of vitamin E and vitamin C on
male infertility: a meta-analysis. *Int Urol Nephrol* **54**, 1793–1805 (2022).

23.    Saito, H., Hara, K., Kitajima, S. & Tanemura, K. Effect of vitamin E deficiency on
spermatogenesis in mice and its similarity to aging. *Reproductive Toxicology* **98**, 225–
232 (2020).

24.    Şen, E., Tunali, Y. & Erkan, M. Testicular development of male mice offsprings
exposed to acrylamide and alcohol during the gestation and lactation period. *Hum Exp*
*Toxicol* **34**, 401–414 (2015).

25.    Ueta, T. *et al.* Glutathione Peroxidase 4 Is Required for Maturation of Photoreceptor
Cells. *Journal of Biological Chemistry* **287**, 7675–7682 (2012).

26.    Da Ros, M., Lehtiniemi, T., Olotu, O., Meikar, O. & Kotaja, N. Enrichment of
Pachytene Spermatocytes and Spermatids from Mouse Testes Using Standard
Laboratory Equipment. *Journal of Visualized Experiments* (2019) doi:10.3791/60271.

27.    Conrad, M. *et al.* The Nuclear Form of Phospholipid Hydroperoxide Glutathione
Peroxidase Is a Protein Thiol Peroxidase Contributing to Sperm Chromatin Stability.
*Mol Cell Biol* **25**, 7637–7644 (2005).

28.    Paniagua, R., Martín, A., Nistal, M. & Amat, P. Testicular involution in elderly men:
565            comparison of histologic quantitative studies with hormone patterns. *Fertil Steril* **47**,
566            671–679 (1987).

567    29.    Lasiene, K. *et al.* Age-related morphological peculiarities of human testes. *Folia*
568            *Morphol (Warsz)* **80**, 122–126 (2021).

569    30.    Endo, T. *et al.* Multiple ageing effects on testicular/epididymal germ cells lead to
570            decreased male fertility in mice. *Commun Biol* **7**, 16 (2024).

571    31.    Jiang, H. *et al.* Quantitative histological analysis and ultrastructure of the aging human
572            testis. *Int Urol Nephrol* **46**, 879–885 (2014).

573    32.    Petersen, P. M., Seierøe, K. & Pakkenberg, B. The total number of Leydig and Sertoli
574            cells in the testes of men across various age groups – a stereological study. *J Anat* **226**,
575            175–179 (2015).

576    33.    Xia, Y., Zhu, W.-J., Hao, S.-F., Liang, W.-B. & Li, J. Stereological analysis of age-
577            related changes of testicular peritubular cells in men. *Arch Gerontol Geriatr* **55**, 116–
578            119 (2012).

579    34.    Barnes, C. J., Covington, B. W. & Lee, M. Effect of Aging and Dietary Restriction on
580            Rat Testicular Germ Cell Apoptosis. *J Gerontol A Biol Sci Med Sci* **54**, B199–B204
581            (1999).

582    35.    Morales, E. *et al.* Effect of ageing on the proliferation and apoptosis of testicular germ
583            cells in the Syrian hamster Mesocricetus auratus. *Reprod Fertil Dev* **15**, 89 (2003).

584    36.    Kajstura, J. *et al.* Necrotic and apoptotic myocyte cell death in the aging heart of Fischer
585            344 rats. *American Journal of Physiology-Heart and Circulatory Physiology* **271**,
586            H1215–H1228 (1996).

587    37.    Martin, D. S. D. *et al.* Apoptotic Changes in the Aged Brain Are Triggered by
588            Interleukin-1β-induced Activation of p38 and Reversed by Treatment with
589            Eicosapentaenoic Acid. *Journal of Biological Chemistry* **277**, 34239–34246 (2002).

590    38.    Miyahara, K., Tatehana, M., Kikkawa, T. & Osumi, N. Investigating the impact of
591            paternal aging on murine sperm miRNA profiles and their potential link to autism
592            spectrum disorder. *Sci Rep* **13**, 20608 (2023).

593    39.    Blanco-Rodríguez, J. & Martínez-García, C. Spontaneous germ cell death in the testis
594            of the adult rat takes the form of apoptosis: re-evaluation of cell types that exhibit the
595            ability to die during spermatogenesis. *Cell Prolif* **29**, 13–31 (1996).

596    40.    Yuan, H., Li, X., Zhang, X., Kang, R. & Tang, D. Identification of ACSL4 as a
597            biomarker and contributor of ferroptosis. *Biochem Biophys Res Commun* **478**, 1338–
598            1343 (2016).

599    41.    Oresti, G. M. *et al.* Differentiation-related changes in lipid classes with long-chain and
600            very long-chain polyenoic fatty acids in rat spermatogenic cells. *J Lipid Res* **51**, 2909–
601            2921 (2010).

602    42.    Otaka, K., Hiradate, Y., Kobayashi, N., Shirakata, Y. & Tanemura, K. Distribution of
603            the sex chromosome during mouse spermatogenesis in testis tissue sections. *J Reprod
604            Dev* **61**, 375–81 (2015).

605    43.    Martin, R. H. *et al.* The chromosome constitution of 1000 human spermatozoa. *Hum
606            Genet* **63**, 305–309 (1983).

607    44.    van Kooij, R. J. & van Oost, B. A. Determination of sex ratio of spermatozoa with a
608            deoxyribonucleic acid-probe and quinacrine staining: a comparison. *Fertil Steril* **58**,
609            384–386 (1992).

610    45.    Quinlivan, W. L. G., Preciado, K., Long, T. L. & Sullivan, H. Separation of human X
611            and Y spermatozoa by albumin gradients and Sephadex chromatography. *Fertil Steril*
612            **37**, 104–107 (1982).

613    46.    Watkins, P. A., Maiguel, D., Jia, Z. & Pevsner, J. Evidence for 26 distinct acyl-
614            coenzyme A synthetase genes in the human genome. *J Lipid Res* **48**, 2736–2750 (2007).

615    47.    Bhutani, K. *et al.* Widespread haploid-biased gene expression enables sperm-level
616            natural selection. *Science* **371**, (2021).

617    48.    Bensoussan, K, Morales, C. R. & Hermo, L. Vitamin E Deficiency Causes Incomplete
618            Spermatogenesis and Affects the Structural Differentiation of Epithelial Cells of the
619            Epididymis in the Rat. *J Androl* **19**, 266–288 (1998).

620    49.    Luo, S. *et al.* Vitamin E and GPX4 cooperatively protect treg cells from ferroptosis and
621            alleviate intestinal inflammatory damage in necrotizing enterocolitis. *Redox Biol* **75**,
622            103303 (2024).

623    50.    Hu, Q. *et al.* GPX4 and vitamin E cooperatively protect hematopoietic stem and
624            progenitor cells from lipid peroxidation and ferroptosis. *Cell Death Dis* 12, 706 (2021).

625    51.    Round Spermatid Nucleus Injection (ROSNI). *Fertility and Sterility* **90** (2008).

626    52.    Levine, H. *et al.* Temporal trends in sperm count: a systematic review and meta-
627            regression analysis. *Hum Reprod Update* **23**, 646–659 (2017).

628    53.    Levine, H. *et al.* Temporal trends in sperm count: a systematic review and meta-
629           regression analysis of samples collected globally in the 20th and 21st centuries. *Hum*
630           *Reprod Update* **29**, 157–176 (2023).

631    54.    Tatehana, M., Kimura, R., Mochizuki, K., Inada, H. & Osumi, N. Comprehensive
632           histochemical profiles of histone modification in male germline cells during meiosis
633           and spermiogenesis: Comparison of young and aged testes in mice. *PLoS One* **15**,
634           e0230930 (2020).

**Figure legends**

**Fig. 1: Age-related alterations in mouse testis germ cell populations**

**a**, Bodyweight (g) in young and aged male mice. **b**, Testis weight (mg) in young and aged males. **c**, Gonadosomatic index (testis weight/bodyweight) in young and aged males. **d**, Schematic showing the process of spermatogenesis, highlighting key differentiation stages, undifferentiated spermatogonia (Type A Spg), differentiating spermagotogonia (Type B Spg), pachytene spermatocytes (PSs), round spermatids (RSs) and Sertoli cells (SCs). **e**, Representative images of seminiferous tubules from young (top row) and aged (bottom row) mice, showing cells positive for Plzf (Type A Spg marker), Kit (Type B Spg marker), Scp3 (meiotic PSs marker), PNA (acrosome RSs marker) and Sox9 (SCs marker). Arrowheads indicate Kit positive cells. Images are merged with DAPI. Scale bar, 50 μm. **f-j**, Quantification of Plzf positive Type A Spg (**f**), Kit positive Type B Spg (**g**), Scp3 positive PSs (**h**), PNA positive RSs (**i**) and Sox9 positive SCs (**j**). Quantification in (**f**) is based on 20 images per mouse, whereas (**g-j**) are based on 10 images per mouse, normalized to tubule circumference. Data are presented as mean ± SD. ***P* <0.01, ****P* <0.001 determined by Two-tailed Student's *t*-test or Mann-Whitney U test. Sample size: *N* = 6 mice per group (young, 3-months; aged 18-months). Each dot represents one mouse.

**Fig. 2: Distinct cell death decisions in pre- and post-meiotic germ cells**

**a**, Percentage of seminiferous tubules containing cells TUNEL+ cells relative to the total number of tubules within a section. **b**, Quantification of cell death index (number of dead cells/total tubules). **c**, Image of a seminiferous tubule from a young mouse, indicating the peripheral area (in between the dashed and solid line) and the luminal area (within the solid line) areas. **d-g′**, Representative images of tubules in young (top row) and aged (bottom row) mice stained for cell death with Casp3 and TUNEL, showing Casp3+/TUNEL+ cells (closed arrowheads) in the periphery (**d, d′**), tubules with both Casp3+/TUNEL+ and Casp3-/TUNEL+ cells (open arrowheads) in the periphery (**e, e′**), tubules with only Casp3-/TUNEL+ cells in the periphery (**f, f′**) and tubules with Casp3-/TUNEL+ in both the periphery and lumen (**g, g′**). Images are merged with Hoechst. Scale bar, 50 μm. **h, i**, Quantification of Casp3+/TUNEL+ and Casp3-/TUNEL+ cells in (**h**) the periphery and (**i**) luminal areas, normalized to 100 tubules. Data are presented as mean ± SD in all panels, ns = not significant, ***P* <0.01, ****P* <0.001 as determined by two-tailed Student's *t*-test or Mann-Whitney U test. Sample size: *N* = 6 mice per group (young, 3-months; aged 18-months). Each dot represents one mouse.

**Fig. 3: Round spermatids show hallmarks of ferroptosis**

**a**, Schematic outlining the procedure of obtaining RNA and protein from round spermatid

(RS)- enriched fractions in young and aged mice using a modified density gradient method. **b**,

RS enrichment analysis. Fraction 4-7 consistently showed >80% enrichment, confirmed by

PNA (acrosomal) staining. Scale bar, 40 μm. **c**, Simplified diagram illustrating ferroptosis,

which occurs in response to excessive iron-dependent lipid peroxidation mediated by Acsl4

and Alox15 (red box), coupled with a loss of Gpx4 antioxidant function (green box). **d**, Gpx4

protein localization in young (top panel) and aged (bottom panel) mice, merged with DAPI.

Scale bar, 40 μm. **e**, Western blot images showing Gpx4 protein levels relative to ActB (top),

with quantitative analysis (bottom). **f-j**, Quantification of the mRNA levels of the *nGpx4*

isoform (**f**), *mGpx4* isoform (**g**), *cGpx4* isoform (**h**), *Acsl4* (**i**) and *Alox15* (**j**), in young and aged

RSs showing fold change relative to *ActB*. **k**, Western blot images of 4HNE levels relative to

Gapdh (top), with quantitative analysis (bottom). **l**, FISH analysis of RSs with either X (red,

open arrowhead) or Y (green, closed arrowhead) signal (left), and quantification of the Y/X

ratio (Y-chromosomes/X-chromosomes, young: Y/X 1.02, *n*=13,303 vs aged: Y/X: 1.11,

*n*=13,008, right), merged with Hoechst. Scale bar, 5 μm. Data are presented as mean ± SD in

all panels, ns = not significant, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$ as determined by Two-tailed

Student's *t*-test or Mann-Whitney U test. Sample size: $N = 6$ mice per group (young, 3-months;

aged, 18-months). Each dot represents one mouse.

**Fig. 4: Vitamin E modulation induces or rescues age-related changes in spermatogenesis**

**a**, Study design, young mice were fed either a control diet (YC) or a vitamin E-deficient diet

(YVE-) and aged mice were fed a control (AC) or VE-supplemented diet (AVE+) for 70 days.

**b,c**, Representative images of tubules containing PNA positive RSs (**b**), quantified from 10

images per mouse (**c**), Scale bar, 5 μm. **d**, Representative images of tubules showing

Casp3+/TUNEL+ (closed arrowheads) and Casp3-/TUNEL+ (open arrowheads) cells in the

periphery, and Casp3-/TUNEL+ cells in the lumen. Scale bar, 50 μm. **e**, Percentage of

seminiferous tubules containing cells TUNEL+ cells relative to the total number of tubules

within a section. **f**, Quantification of cell death index (number of dead cells/total tubules). **g,h**,

Quantification of Casp3+/TUNEL+ cells in the peripheral area (**g**), and Casp3-/TUNEL+ in

the luminal area (**h**). **i**, Western blot images of Gpx4 and ActB (top) and quantitative analysis

(bottom). **j-n**, Quantification of the mRNA levels of the *nGpx4* isoform (**j**), *mGpx4* isoform

(**k**), *cGpx4* isoform (**l**), *Ascl4* (**m**) and *Alox15* (**n**) relative to *ActB*. **o,** Western blot images of

4HNE relative to Gapdh (top) and quantitative analysis (bottom). **p,** FISH analysis of RSs with

X (Red, open arrowhead) or Y (Green, closed arrowhead) signals (left), and quantification of

the Y/X ratio (Y chromosomes/X chromosomes, YC: 1.02, $n$ = 13,100, YVE-: 1.07, $n$= 13,300,

AC: 1.10, $n$= 13,022, AVE+: 1.01 $n$= 12,018). Scale bar, 5 µm merged with Hoechst. Data are

presented as mean ± SD in all panels. Groups that do not share a common letter are statistically

different as determined by ANOVA with *post-hoc* Tukey HSD or Kruskal-Wallis with *post-*

*hoc* Dunn's test. Sample size: $N$=6 for panels **c, e-i, o,** and **p**; $N$=9 (YC), $N$=8 (YVE-), $N$=7

(AC), and $N$=9 (AVE+) for panels **j-n**. Each dot represents one mouse.

**Extended data Fig. 1: Casp3+/TUNEL+ is observed in round spermatids**

Representative images of tubules containing Casp3+/TUNEL+ round spermatids (closed

arrowheads) in young (top panel) and aged (bottom panel) mice. Scale bar, 50 µm.

**Extended data Fig. 2: Vitamin E modulation induces or rescues spermatogenesis**

**impairment**

**a,b,** Bodyweight (g) of young control (YC) and VE deficient (YVE-) mice (**a**) and aged control

(AC) and VE supplemented (AVE+) mice (**b**). ***$P$<0.001 = significant fixed effect of week

on bodyweight, n.s = non significant fixed effect of diet on bodyweight (Mixed-model). **c,**

Testis weight (mg) of YC, YVE-, AC, and AVE+ mice. **d**, Gonadosomatic index (testis weight

/bodyweight) of YC, YVE-, AC, and AVE+ mice. **e**, Representative images of seminiferous

tubules in YC, YVE-, AC and AVE+ mice, containing germ cells positive for Plzf (Type A

Spg, top row), Kit (Type B Spg, middle row), and Scp3 (meiotic PSs, bottom row). Scale bar,

50 µm. **f-h,** Quantification of Plzf positive Type A Spg (**f**), Kit positive Type B Spg (**g**), Scp3

positive PSs (**h**). Quantification in (**f**) is based on 20 images per mouse, and in (**g**) and (**h**) is

based on 10 images per mouse, normalized to the circumference of each tubule. **i,** A variety of

tubules in YC, YVE-, AC, AVE+ mice, showing heterogeneity of cell death in the peripheral

and luminal areas, Casp3+/TUNEL+ cells (closed arrowheads) and Casp3-/TUNEL+ (open

arrowheads). Scale bar, 50 µm. **j,k,** Quantification of Casp3-/TUNEL+ cells in the peripheral

area (**j**), and Casp3+/TUNEL+ in the luminal area (**k**). Data is presented as mean ± SD in all

panels, groups that do not share a common letter are statistically different, determined by

ANOVA with *post-hoc* Tukey HSD or Kruskal-Wallis with *post-hoc* Dunn's test. Each dot

represents one mouse. Sample size: $N$=9 (YC), $N$=8 (YVE-), $N$=7 (AC), and $N$=9 (AVE+) for

panels **a-d**. $N$=6 for panels **f-h, j, k**; Each dot represents one mouse.

Fig. 1

Casp3/TUNEL/Hoechst

Fig. 2

Fig. 3

Fig. 4

Casp3/TUNEL/Hoechst

Young

Aged

Extended data Fig. 1

Extended data Fig. 2