Ferroptotic vulnerability of post-meiotic germ cells skews sex chromosome ratios in

aged testis

Jasper K. Germeraad[1], Takako Kikkawa[1]*, Junya Ito[2,3], Leon S.Y. Giesselink[4], Wilfred T.V.

Germeraad[5], Bernhard Henkelmann[3], Maceler Aldrovandi[3], Kenshiro Hara[6], Kentaro

Tanemura[6], Kiyotaka Nakagawa[2], Eikan Mishima[3,7], Marcus Conrad[3,8], Noriko Osumi[1]*

[1]Department of Developmental Neuroscience, Tohoku University Graduate School of Medicine, Sendai,

Japan

[2]Laboratory of Food Function Analysis, Graduate School of Agricultural Science, Tohoku University,

Sendai, Japan

[3]Institute of Metabolism and Cell Death, Molecular Targets and Therapeutics Center, Helmholtz

Munich, Neuherberg, Germany

[4]Centre for Health Protection, National Institute for Public Health and the Environment (RIVM),

Bilthoven, The Netherlands

[5]Department of Internal Medicine, Division of Haematology, Maastricht University Medical Center,

Maastricht, The Netherlands

[6]Laboratory of Animal Reproduction and Development, Graduate School of Agricultural Science,

Tohoku University, Sendai, Japan

[7]Environmental Medicine and Molecular Toxicology, Graduate School of Medicine, Tohoku University,

Sendai, Japan

[8]Translational Redox Biology, Natural School of Sciences, Technical University of Munich, Garching,

Germany

Running title: Non-apoptotic cell death in aged testes

Keywords**:** Spermatogenesis, regulated cell death, lipid peroxidation, vitamin E

*Corresponding author

Noriko Osumi, PhD

Professor, Department of Developmental Neuroscience,

Tohoku University Graduate School of Medicine,

2-1 Seiryo-machi, Aoba-ku, Sendai 980-8575, Japan

Tel.: +81 22 717 8201. Fax: +81 22 717 8205.

E-mail address: noriko.osumi.c7@tohoku.ac.jp (N. Osumi)

Takako Kikkawa, PhD

Assistant Professor, Department of Developmental Neuroscience,

Tohoku University Graduate School of Medicine,

2-1 Seiryo-machi, Aoba-ku, Sendai 980-8575, Japan

Tel.: +81 22 717 8203. Fax: +81 22 717 8205.

E-mail address: takako.kikkawa.c4@tohoku.ac.jp (T. Kikkawa)

Grant Sponsors: JSPS KAKENHI (#17K19372, 22K19322, 25K22536), The Canon

Foundation, JST FOREST Program (JPMJFR2213), European Union's Horizon 2020 research

and innovation program (GA 884754).

Abstract

Age-related testicular germ cell depletion has been predominantly attributed to apoptosis;

however, the contribution of alternative cell death modalities remains unclear. Using young

and aged mice, we demonstrate compartment-specific cell-death signatures in the testis.

Apoptotic cell increased in peripheral germ layers, whereas the luminal round spermatid (RS)

layer mainly accumulated non-apoptotic events sharing features with ferroptosis, a lipid-

peroxidation dependent cell death. Lipidomic analysis revealed age-related phospholipid

remodelling in whole testis, while RS-enriched fractions showed increased lipid peroxidation.

Interestingly, aging shifted the balance of sex-chromosome–bearing RS toward Y-bearing cells,

and this bias persisted in mature sperm. Dietary vitamin E modulated these phenotypes

bidirectionally: vitamin E deficiency in young mice increased RS lipid peroxidation and

reproduced the Y-skew, whereas supplementation in aged mice lowered RS lipid peroxidation,

decreased non-apoptotic events, and restored the skewed RS sex-chromosome ratio. These

findings define an age-related RS vulnerability linking redox stress to haploid spermatogenic

output.

Introduction

Introduction

Advanced paternal age is associated with reduced semen volume, lower sperm concentration

and motility, and increased sperm DNA fragmentation[1–3]. Although male fertility can be

maintained throughout life, reproductive capacity progressively declines because of age-

dependent deterioration of testicular function[4]. Sperm are produced through spermatogenesis,

a strictly controlled process within the seminiferous tubules in which spermatogonial stem cells

generate differentiating spermatogonia, meiotic spermatocytes and post-meiotic haploid round

spermatids (RSs) that subsequently elongate into spermatozoa[5]. Each round spermatid (RS)

carries either a Y- or an X-chromosome. Therefore, selective survival or loss at post meiotic

stages has the potential to bias sex-chromosome transmission by altering the relative

representation of Y- and X-bearing gametes. While age-related impairment of murine

spermatogenesis is well documented[6–8], the mechanisms driving germ-cell loss during

testicular aging remain insufficiently understood, particularly at post-meiotic stages[9].

Germ-cell loss in aging testes has frequently been linked to oxidative stress-induced

apoptosis[10,11]. However, oxidative stress can activate multiple cell death modalities[12], and

susceptibility is likely to vary along the germ-cell differentiation axis. Previous reports showed

that pre-meiotic germ and somatic cells predominantly engage apoptosis[13], whereas recent *in*

vitro studies suggests that spermatocytes and RSs respond differently to oxidative insults,

including lipid peroxides[14]. RS membranes are enriched in polyunsaturated phospholipids and

undergo extensive membrane remodelling, features predicted to increase sensitivity to lipid

peroxidation[15,16]. Consistent with this notion, elevated lipid peroxidation represents a recurring

biochemical signature of reproductive aging in testes[17].

Lipid peroxidation can trigger ferroptosis, a non-apoptotic form of cell death driven by iron-

mediated phospholipid peroxidation[18]. Ferroptosis susceptibility is shaped by three core pillars:

antioxidant defence mechanisms, membrane phospholipid composition and control of labile

iron[19,20]. For instance, the selenoenzyme glutathione peroxidase 4 (GPX4) detoxifies

potentially harmful lipid hydroperoxides, and impaired GPX4 functions confers cellular

vulnerability to ferroptosis[21,22]. Conversely, long-chain fatty acid CoA ligase 4 (ACSL4)

promotes incorporation of poly-unsaturated fatty acids (PUFAs) into membrane phospholipids,

generating highly peroxidizable substrates that sensitize cells to ferroptosis[23]. Iron-handling

pathways further modify this process; transferrin receptor 1 (TFR1) mediated iron uptake and

heme oxygenase-1 (HO-1) dependent heme degradation can elevate iron levels, amplifying

ferroptotic sensitivity[24].

Ferroptosis is suppressed by lipophilic antioxidants, among which vitamin E is one of the most

potent natural antioxidants that limits membrane lipid peroxidation and exerts anti-ferroptotic

activity[25]. Vitamin E deficiency induces spermatogenic defects resembling features of

reproductive aging[26] and can preferentially reduce post meiotic germ-cell populations, despite

differences in dietary regimens and species[27]. These observations suggest that disruption of

lipid redox homeostasis may be sufficient to activate aging-relevant vulnerability pathways,

with RSs representing a particularly susceptible stage where lipid peroxidation could

selectively influence the survival of X- versus Y-bearing haploid cells.

Here, we investigate how aging alters spermatogenic output and cell death modalities,

including apoptosis and ferroptosis, across the seminiferous epithelium. We characterize age-

related lipid peroxidation and related defence pathways in whole testis and in RS-enriched

fractions and quantify sex-chromosome ratios in both RSs and mature sperm. Additionally, by

bidirectionally manipulating dietary vitamin E, through deficiency in young mice and

supplementation in aged mice, we test whether aging-associated RS vulnerability and sex-

chromosome ratio shifts are causally linked to the dysregulation of the lipid peroxidation-

ferroptosis axis.

Results

Aging impairs spermatogenesis and promotes non-apoptotic cell death in RSs

By 18 months of age, C57BL/6J males exhibit measurable testicular aging[28]. To investigate aging-related gonadal changes, we therefore compared young (3-month-old) and aged (18-month-old) male mice. Aged mice displayed increased body weight (Fig. 1a) and decreased testis weight (Fig. 1b), resulting in a lower gonadosomatic index (testis weight/body weight, Fig. 1c). To assess spermatogenic output in aged mice, we focused on stage VII-VIII seminiferous tubules, which contain undifferentiated spermatogonia, differentiating spermatogonia, meiotic pachytene spermatocytes, post-meiotic RSs and somatic Sertoli cells (spermatogenesis support cells), thus representing key stages of spermatogenesis (Fig. 1d). Testis from aged mice showed reduced cellular density in germ and somatic cells of the seminiferous epithelium (Fig. 1e). Quantitative analysis showed a reduction in undifferentiated spermatogonia in aged compared with young testes (Fig. 1f), which extended to differentiating spermatogonia, pachytene spermatocytes, RSs, and was also observed in somatic SCs (Fig. 1g–j).

Next, to investigate germ cell type–specific cell death associated with aging, we evaluated cell death using terminal deoxynucleotidyl transferase dUTP nick end labelling (TUNEL) across seminiferous tubules, irrespective of stage. Although TUNEL staining was classically used as a marker of apoptosis, it is also known to detect cells undergoing other regulated cell death modalities[19]. A higher proportion of seminiferous tubules in aged mice contained TUNEL-positive dead cells compared to those in young mice (Extended data Fig. 1a, b).

To identify apoptotic cells, we performed double labelling with TUNEL (a general cell death marker) and immunostaining against cleaved-caspase3 (CASP3), a specific marker of apoptosis. We quantified CASP3$^+$/TUNEL$^+$ cells (apoptotic) and CASP3$^-$/TUNEL$^+$ cells (dead cells) separately within the peripheral compartment, containing spermatogonia, spermatocytes and Sertoli cells, and the luminal layer, which is enriched in RSs (Fig. 1k). In the peripheral compartment, aging mice showed a significantly larger number of CASP3$^+$/TUNEL$^+$ apoptotic cells (Fig. 1l), whereas the number of CASP3$^-$/TUNEL$^+$ dead cells remained unchanged (Fig. 1m). In contrast, within the luminal layer, CASP3$^+$/TUNEL$^+$ apoptotic RSs were rare and did not differ between age groups (Fig. 1n, Extended Data Fig. 1c), while CASP3$^-$/TUNEL$^+$ cells prevailed and were more abundant in aged testes (Fig. 1o). Although cleaved CASP3 can be transient, the marked enrichment of CASP3$^-$/TUNEL$^+$ events specifically in the luminal RS

149   layer supports a compartment-specific shift toward TUNEL[+] cell-death events without

150   detectable CASP3 activation in aging testes.

Aging enhances lipid peroxidation in RSs

Increased lipid peroxidation is a feature of testicular aging and capable of inducing cell death

in aging rodent models[29–31]. To define age-related changes in phospholipid composition and

oxidation status, we profiled testicular phospholipids by mass-spectrometry-based lipidomic

analysis. Consistent with previous studies[32,33], phosphatidylcholine (PC) and

phosphatidylethanolamine (PE) constituted the predominant phospholipid classes detected in

mouse testis (Extended Data Fig. 2a, b). Age-related changes in individual PC and PE species

were modest overall, although aged testes showed an increase in PC plasmalogen species

(Extended Data Fig. 2c), but not PE plasmalogen species (Extended Data Fig. 2d). The increase

in PC plasmalogen species may reflect age-dependent remodelling of ether lipids, which are

implicated in antioxidant defence under oxidative stress conditions[34].

Given that RSs are enriched in polyunsaturated phospholipid species[15], this may suggest

heightened vulnerability to oxidative stress and lipid peroxidation. We isolated RS-enriched

fractions from young and aged testes using a density-gradient (Fig. 2a) and confirmed >80%

RS purity by peanut agglutinin (PNA) staining and nuclear morphology (Extended Data Fig.

3). Immunoblot analysis of RS-enriched fractions revealed increased levels of 4-

hydroxynonenal (4HNE)-modified proteins in aged compared with young samples, indicating

enhanced lipid peroxidation in aged RSs (Fig. 2b). In line, gene expression levels of *Alox15*,

encoding a PUFA-oxidizing lipoxygenase, was elevated in aged RSs (Fig. 2c). Because

susceptibility to lipid peroxidation is influenced by how PUFAs are incorporated into

membrane phospholipids[23], we examined gene expression levels of key enzymes involving

PUFA loading and elongation. In aged RSs, expression of *Acsl4,* encoding an acyl-CoA

synthetase that facilitates arachidonic/adrenic acid (AA/AdA)-PUFA enrichment in

phospholipids*,* was increased, whereas other ACSL family genes such as *Acsl6* and the PUFA-

elongating enzymes *Elovl2* and *Elovl5* were not altered with age (Fig. 2d, e, Extended Data Fig.

2e, f). Together, these data are consistent with a selective shift toward AA/AdA-type PUFA

phospholipid remodelling rather than enhanced docosahexaenoic acid (DHA) loading in aged

RSs, supporting increased susceptibility of AA/AdA-containing phospholipids to oxidation.

Aging reduces cGpx4 expression and remodels iron handling in RSs

To investigate the mechanism underlying increased lipid peroxidation in aged testes, we

assessed antioxidant defence and iron-handling pathways in RSs. GPX4 is a central regulator

of lipid peroxide detoxification and the prime regulator of ferroptosis. Immunostaining

confirmed GPX4 expression in the RS cytoplasm and in the sperm tail (Extended Data Fig. 4a).

In testis, GPX4 is expressed as three isoforms: nuclear (nGPX4), mitochondrial (mGPX4) and

cytosolic (cGPX4) isoforms, of which cGPX4 has been implicated as the critical isoform

protecting germ cell from cell death[35–38]. We therefore quantified *Gpx4* mRNA transcripts in

RSs from young and aged mice, using primer sets targeting the nuclear (*nGpx4*) transcript and

the mitochondrial-targeted (*mGpx4*) transcript, together with a primer set amplifying both

mitochondrial and cytosolic (*cGpx4*) transcripts[39]. Using this strategy, we estimated the *cGpx4*

fraction as the difference between the shared signal and the mitochondrial-specific signal.

While *nGpx4* and *mGpx4* were unchanged with age, *cGpx4* expression was reduced in aged

RSs (Fig. 2f–h). Total GPX4 protein levels, detected with an antibody recognizing all isoforms,

did not differ between young and aged RSs fractions (Fig. 2i). Given the predominance of

mGPX4 in RSs[36], these findings suggest that age-associated downregulation of *cGpx4* occurs

without a detectable change in bulk RS GPX4 protein levels.

Nuclear erythroid factor 2 (NRF2) controls a broad antioxidant and cytoprotective program[40]

(Fig. 2j). Among NRF2 target genes, superoxide dismutase-*1* (*Sod1*) and catalase (*Cat*)

mRNAs did not change in RSs (Fig. 2k–m), whereas *HO-1* was selectively induced in aged

RSs (Fig. 2n), consistent with increased heme catabolism. Because HO-1 can elevate the labile

iron pool, we assessed iron handling and found increased *Tfr1* mRNA and increased ferritin

heavy chain (FTH1), with no change in ferritin light chain (FTL) and nuclear receptor co-

activator 4 (NCOA4) protein in aged RSs (Fig. 2o–s; Extended Data Fig. 4b, c). Together, these

changes support an age-associated shift toward a more pro-oxidant iron environment in RSs

(Fig. 2t), which would be expected to sensitize these cells to iron-dependent lipid damage.

Aging skews the Y/X sex chromosome ratio in RSs and mature sperm

Since *Acsl4* is encoded by the X-chromosome and is upregulated in aged RSs (Fig. 2f), we

asked whether aging is associated with a bias in the relative abundance of X- and Y-bearing

RSs. To distinguish X- and Y-bearing RSs, we performed fluorescent *in situ* hybridization

(FISH), targeting the X- and Y- chromosome with specific probes. We found that the ~1:1 ratio

was maintained in young testes but was significantly skewed toward Y-bearing RSs in aged

testes (young: mean Y/X ratio 1.02 [$n = 6,282 \pm 100$ Y-RSs, $n = 6,188 \pm 113$ X-RSs]; aged: mean

Y/X ratio 1.11 [$n = 6,694\pm85$ Y-RSs, $n = 6,056\pm98$ X-RSs]; Fig. 3a,b). To determine whether

this bias persisted in mature sperm, we quantified X- and Y-bearing sperm using fluorescent

activated cell sorting (FACS) using Hoechst-33342 to visualize the DNA content difference

between Y- and X- bearing RSs (Fig. 3c, d). Consistent with the findings in RSs, the Y/X ratio

was similarly shifted toward Y-bearing sperm in aged males compared with young controls

(young: mean Y/X 1.02 [$n = 3,382,147 \pm 140,662$ Y-sperm vs $n = 3,293,224 \pm 126,522$ X-

sperm]; aged: mean Y/X 1.09 [$n = 3,294,779 \pm 156,878$ Y-sperm; $n = 3,032,412 \pm 145,748$ X-

sperm], Fig. 3e,f). These results indicate that aging is associated with a bias toward Y-bearing

RSs, and this sex chromosome bias is maintained in mature sperm.

Vitamin E depleted diet recapitulated aging-associated phenotypes

Given the elevated lipid peroxidation observed in aged testis, we asked whether a lipid

peroxidation-prone condition induced by vitamin E-depleted diet may cause aging-like

testicular phenotypes in young mice[26]. Young animals were fed either a vitamin E-deficient

diet (VE(−), <0.1mg/kg vitamin E) or control diet (135mg/kg vitamin E) for two cycles of

spermatogenesis[5] (70 days, Fig. 4a). Although body weight, testis weight and gonadosomatic

index were not affected by vitamin E-deficient diet (Fig. 4b–d), VE(−) testes showed impaired

spermatogenic output (Fig. 4e). Quantitative analysis revealed that numbers of undifferentiated

and differentiating spermatogonia remained unchanged, whereas pachytene spermatocytes,

RSs and Sertoli cells were reduced in VE(−) testes compared with controls (Fig. 4f–j),

indicating impaired spermatogenic progression.

Consistent with the reduction in germ and somatic cell numbers, VE(−) testes contained more

TUNEL$^+$ dead cells than control testes (Extended Data Fig. 5a, b). When cell death was

analysed separately in the peripheral compartment and the luminal RS layer (Fig. 4k), vitamin

E deficiency significantly increased CASP3$^+$/TUNEL$^+$ apoptotic cells in the peripheral regions

without altering CASP3$^-$/TUNEL$^+$ cells (Fig. 4k, l). By contrast, in the luminal compartment,

vitamin E-deficient diet only induced a trend toward higher CASP3$^-$/TUNEL$^+$ non-apoptotic

RSs, but CASP3$^+$/TUNEL$^+$ RSs remained rare and unchanged (Fig. 4l, m; Extended Data Fig.

5d). Together, vitamin E deficiency preferentially elevates apoptosis-associated death events

in the peripheral compartment, while its effect on RS-layer CASP3$^-$/TUNEL$^+$ events is weaker

under these conditions.

In RSs isolated from VE(−) testes, expression of lipid-metabolic and stress-response genes was

significantly altered. *Acsl4, Hmox1, Ftl* and *Tfr1* transcripts were upregulated, whereas all three

Gpx4 isoforms were reduced (Fig. 4n). Consistent with enhanced lipid peroxidation, the levels

of 4HNE-modified proteins were elevated in VE(−) RSs (Fig. 4o). At the protein level, FTH1

was decreased, whereas total GPX4, FTL and NCOA4 were unaffected (Fig. 4p-t).

We further examined whether feeding mice with a vitamin E-deficient diet altered the sex

chromosome composition of RSs. In control testes, Y- and X-bearing RSs were similar in

number, whereas VE(−) testes exhibited a significant enrichment of Y-bearing RSs (control:

mean Y/X ratio 1.02 [$n = 6,390\pm125$ Y-RSs vs $n = 6,274\pm127$], X-RSs; VE(−): Y/X ratio 1.07

[$n = 6,548\pm222$ Y-RSs vs $n = 6,186\pm212$ X-RSs]; Fig. 4u). Overall, vitamin E deficiency in

young mice induced an aging-like RS stress state, with increased lipid peroxidation and broad

remodelling of lipid/iron-defence transcripts in RS fractions. Notably, despite limited induction

of RS-layer CASP3^{-}/TUNEL^{+} events, vitamin E deficiency produced a significant shift in

haploid output, enriching Y-bearing RSs, thereby recapitulating the sex-chromosome skew

observed in aged testes.

Vitamin E supplementation reduced aging-associated testicular phenotypes

Given that vitamin E deficiency in young mice recapitulated multiple aging-like testicular

phenotypes, we next asked whether vitamin E supplementation could counteract these changes

in aged mice. To this end, aged animals were fed either a vitamin E-supplemented diet (VE(+))

or control diet according to the regimen shown in Fig. 5a. Body weight, testis weight and

gonadosomatic index did not differ between VE(+) and control groups (Fig. 5a–d), indicating

that vitamin E supplementation does not affect gross somatic growth or testis size in aged mice

during this feeding regimen. Despite unchanged testis size, VE(+) improved spermatogenic

output in aged mice (Fig. 5e). While the numbers of undifferentiated and differentiating

spermatogonia were not altered, pachytene spermatocyte, RS and Sertoli cell numbers were

significantly increased in VE(+) compared with control testes (Fig. 5f–j). Consistent with this

improvement, VE(+) testes contained fewer TUNEL^{+} cells than control testes (Extended Data

Fig. 5a, c). In the peripheral compartment, vitamin E supplementation significantly lowered

the Casp3^{+}/TUNEL^{+} apoptotic population without affecting CASP3^{-}/TUNEL^{+} cells (Fig. 5k,

l). In contrast, within the luminal RS layer, VE(+) selectively reduced non-apoptotic

CASP3^{-}/TUNEL^{+} RSs, whereas CASP3^{+}/TUNEL^{+} apoptotic RSs remained infrequent and

unchanged across groups (Fig. 5k, m, Extended Data Fig. 5d). These results indicate that

vitamin E attenuates apoptosis in the periphery and CASP3$^-$/TUNEL$^+$ non-apoptotic events in

the luminal layer, consistent with reduced oxidative burden.

We next assessed the transcriptional changes in RSs from aged mice. In VE(+) RSs, *Alox15*

mRNA was significantly reduced, whereas expression of other lipid/oxidized lipid-metabolic

genes including, *Gpx4*, NRF2-target genes and iron-related transcripts were unchanged relative

to the aged controls (Fig. 5n). Nonetheless, 4HNE-modified protein levels were significantly

reduced in VE(+) RSs (Fig. 5o). Among iron-handling proteins, in RSs from VE(+) mice,

FTH1 and FTL protein levels were significantly reduced, whereas NCOA4 was increased (Fig.

5q–t), a pattern consistent with enhanced ferritin turnover, suggesting mobilization of ferritin-

bound iron; however, whether this increases the labile iron pool or is coupled to iron

utilization/export remains to be determined.

VE(+) also exerted an opposite effect on the RS sex chromosome composition compared to

VE(−). In aged control testes, Y-bearing RSs outnumbered X-bearing RSs, whereas, in contrast,

VE(+) restored the Y- and X-bearing RSs ratio to a near 1:1 ratio (control: mean Y/X ratio 1.10

[$n = 6,527\pm135$ Y-RSs vs $n = 5,950\pm156$ X-RSs]; VE(+): Y/X ratio 1.01 [$n = 5,772\pm156$ Y-

RSs vs $n = 5,702\pm149$ X-RSs]; Fig. 5u). Taken together, the vitamin E deficiency and

supplementation experiments show that in vitamin E deficient conditions, higher lipid

peroxidation is associated with enrichment of Y-bearing RSs, whereas vitamin E

supplementation in aged mice lowers lipid peroxidation and restores the RS sex ratio toward

~1:1. These findings are consistent with preferential vulnerability of X-bearing RSs under

conditions of elevated lipid peroxidation.

Discussion

Aging in mammals is accompanied by testicular atrophy and progressive decline in spermatogenic capacity[28,41,42]. Here, we identified post-meiotic RSs as a particularly age-sensitive stage in which lipid peroxidation increases, coinciding with a compartment-specific shift in death signatures and an altered haploid sex-chromosome output. Dietary modulation of vitamin E reversed these phenotypes, indicating that membrane redox buffering can influence spermatogenic efficiency and the relative abundance of Y- and X-bearing haploid cells.

Aging increased germ-cell death, but not uniformly across the seminiferous epithelium. Peripheral layers showed an apoptotic signature consistent with prior work [43,44], whereas the luminal RS layer accumulated predominantly non-apoptotic TUNEL$^+$ events. This stage specificity may reflect intrinsic vulnerabilities of RSs, whose membranes are enriched in peroxidation-prone PUFAs[15,45], potentially pushing RSs closer to a lipid-damage threshold. RS-enriched fractions indeed displayed characteristics indicative of enhanced lipid peroxidation and reduced antioxidant capacity. The convergence of lipidomic oxidation signatures, altered antioxidant and iron-handling pathways, and bidirectional modulation by vitamin E supports a death modality that shares features with ferroptosis[18,23,46].

A striking consequence of this RS vulnerability is a shift in haploid sex-chromosome composition. We observed a skew toward Y-bearing RSs in aged testes, indicating that the sex-chromosome ratio can deviate from ~1:1 and persists in mature sperm. Because the skew is already present at the RS stage, it is unlikely to arise solely from selective processes during epididymal transit, although upstream meiotic drive and downstream quality-control mechanisms could still change the final output[47,48]. Round spermatids share gene transcripts across cytoplasmic bridges[49], however, incomplete sharing of gene products across intercellular bridges provides a plausible route for divergence between X- and Y-bearing RSs, and geno-informativity analyses have identified incompletely shared gene transcripts relevant to lipid peroxidation pathways[50]. Interestingly, independent *in vitro* studies report an X-skew among surviving mature sperm following acute oxidative stress[51]. Although the direction of this bias differs from our observed RS-stage Y-skew, the two observations are not directly comparable given differences in developmental stage, context, and stress regime. Nonetheless, our dietary vitamin E data indicate that membrane redox buffering can influence RS haploid output, shifting the Y/X ratio toward or away from parity, consistent with the broader principle

that redox conditions can bias sex-chromosome representation at the gamete level[52], potentially

via differential vulnerability of X- and Y-bearing cells.

This study has several limitations. First, although the lipid peroxidation profile and vitamin E

sensitivity in RSs support a lipid damage state, definitive pathway assignment *in vivo* is

constrained by the essential role of GPX4 in spermatogenesis, which limits straightforward

genetic testing of this axis. Establishing causality will therefore require complementary

approaches, such as perturbation of more genetically tractable suppressor pathways. Second,

the basis of the aging-associated Y/X skew remains unknown because RS-enriched fractions

were not sex-resolved and upstream or downstream selection may also contribute. Finally, we

did not test whether vitamin E modifies the Y/X ratio in offspring sex ratio in aged males,

which will require controlled mating studies.

Overall, our findings position RSs as a key stage at which aging-associated lipid redox stress

may bias haploid sex-chromosome ratios (Extended Data Fig. 6). Defining how lipid

peroxidation, antioxidant capacity and Y- versus X-linked vulnerabilities are coordinated *in*

vivo will be important for understanding how male reproductive aging shapes sex-chromosome

transmission.

Methods

Animals

Male C57BL/6J mice were maintained in the Animal Experiment Facility at the Tohoku

University Graduate School of Medicine. All animals were housed in standard cages in a

temperature- and humidity-controlled room on a 12-hour light/dark cycle (light on at 8 am) and

had access to food and water ad libitum. All experimental procedures were approved by the

Ethics Committee for Animal Experiments in Tohoku University (#2023-MED03701), and the

animals were treated according to the National Institutes of Health (NIH) guide for the care

and use of laboratory animals. For age-comparison experiments, C57BL/6J mice were obtained

from Charles River Laboratories and were bred and maintained in-house. Mice were sacrificed

at 3 months old (Young) or 18 months old (Aged).

Vitamin E feeding

A separate cohort of male 1-month and 16-month-old C57BL/6J mice was purchased from

Jackson Laboratories (JAX, Japan). Vitamin E diets were custom-made and ordered from

Funabashi Farm, Japan. 1-month-old animals were fed control (containing 135 mg/kg vitamin

E) or vitamin E-deficient diet (VE(−), containing <0.1 mg/kg vitamin E), and 16-month-old

animals were maintained on control or vitamin E-supplemented diet (VE(+), containing 500

mg/kg vitamin E). Detailed ingredients are listed in Extended Data Table 1. All mice were

raised on these diets for 10 weeks (70 days), covering two full cycles of spermatogenesis (35

days)[7], after which mice were sacrificed at 3- and 18-months of age, respectively.

Testis collection

Animals were sacrificed by sedation with isoflurane (MSD Animal health) and cervical

dislocation. Testes were isolated, separated from the surrounding fat tissue and epididymis and

weighed together. Samples were then fixed in 4% paraformaldehyde in phosphate buffered

saline (PBS, 4% PFA, Nacalai-Tesque) for 2 hours at 4°C, cut in half, and fixed for an

additional 18 hours. Cryoprotection was executed as follows: the testes samples were immersed,

sequentially, in 10% sucrose dissolved in PBS for 1 hour and 20% sucrose in PBS overnight,

both at 4˚C. After cryoprotection, samples were frozen in Tissue-Tek optimal cutting

compound (OCT, Sakura Finetek) on a metal pedestal using liquid nitrogen and stored in -80°C

until further use. Cryosections (10μm) were obtained using a cryostat (Leica CM3050 S) and

used for histological analyses.

Immunohistochemistry (IHC)

Testicular sections on slide glasses (Matsunami) were stained using a previously established

method[53] with antibodies targeting anti-PLZF (1:500, sc28319, Santa Cruz), anti-SCP3 (1:500,

ab97672, Abcam), anti-KIT (1:500, D13A2, Cell Signalling Technologies), anti-SOX9 (1:900,

5535, EMD Millipore), and anti-GPX4 (1:1,000, ab125066, Abcam). Sections were washed

using Tris-buffered saline containing 0.1% Tween20 (TBST). Antigen retrieval was performed

(except for anti-Kit) for 10 minutes in 0.01 M citrate buffer (pH 6.0) at 95°C. Afterwards, slides

were washed with TBST and blocked with 3% bovine serum albumin (BSA) and 0.3% Triton

X-100 in PBS for 1 hour at room temperature (RT). The slides were then incubated with

primary antibodies diluted in blocking solution overnight at 4°C, washed with TBST, and

incubated for 1 hour at RT with 4′,6-diamidino-2-phenylindole (DAPI, 1:1,000, Sigma) as well

as either a Cy3- or Alexa488-conjugated secondary antibody (1:500, Jackson

ImmunoResearch) or peanut agglutinin (PNA-Cy5, 1:2,000, Vector Laboratories) when

staining the acrosome for round spermatid identification. Sections were washed again with

TBST and mounted with Prolong Glass Antifade Mountant (Thermofisher). All images were

obtained using a confocal laser-scanning microscope (LSM800, Carl Zeiss) and processed in

Fiji (ImageJ).

Cell death detection

Apoptotic or non-apoptotic cell death was identified by double labelling for terminal

deoxynucleotidyl transferase dUTP nick end labelling (TUNEL, Merck Millipore) and an

antibody against active-caspase-3 (1:500, BD Biosciences). TUNEL labelling was performed

first and according to the manufacturer's instructions. In short, slides with testis sections were

washed in PBS and subjected to antigen retrieval by microwaving for 5 minutes in 0.01 M

citrate buffer. TdT mixture was applied and incubated at 37°C for 1 hour, followed by

incubation with an anti-digoxigenin antibody conjugated with fluorescein at RT for 30 minutes.

Next, active-caspase-3 IHC was performed as described above with a slight modification:

instead of DAPI, Hoechst-44432 (1:5,000) was used. Using Fiji, male germ line cells in the

peripheral region of the seminiferous tissue and round spermatids in the luminal region were

independently quantified.

Fluorescent *in situ* hybridization (FISH)

FISH was performed according to a previously established protocol[54]. Slides with frozen

sections of the testis were subjected to antigen retrieval in Histo-VT One (Nacalai-Tesque) for

30 minutes at 90˚C and denatured using 50% formamide/2× saline-sodium citrate buffer (SSC),

followed by dehydration using 70% and 100% ethanol. A custom mixture of X (conjugated

with Cy3) and Y (conjugated with FITC) chromosome probes (Chromosome Science Labo)

was applied to the slides, covered with a hybridization cover (Hybrislip, Thermofisher) and

incubated at 80˚C for 10 minutes. Further hybridization occurred overnight in a humid chamber

at 37˚C. To enhance probe signals, IHC was performed. Samples were blocked using 5%

Blocking One solution (Nacalai-Tesque) for 1 hour at RT. Primary antibodies (anti-FITC,

1:2,000, A889, Invitrogen, and anti-Cy3, 1:2,000, sc-166894, Santa Cruz) were diluted in 5%

Blocking One solution and incubated for 1 hour at RT. After washing with 4× SSC, Cy3- or

Alexa488-conjugated secondary antibodies (1:2,000, Jackson ImmunoResearch) together with

Hoechst-44432 (1:5,000, Thermofisher) were applied and incubated for 1 hour at RT. Slides

were mounted with Prolong Glass Antifade Mountant (Thermofisher). All images were

obtained using a fluorescence microscope (Keyence BZ-X series) and processed in Fiji. RSs

were identified according to nuclear morphology and location within the tubule. Hybridization

efficiency was determined by microscopic evaluation and images were deemed appropriate for

analysis when, in more than 95% of cells, a sex chromosome could be assigned.

Chemicals

Butylated hydroxytoluene (BHT, Cat#W218405-1KG-K), ammonium acetate for LC-MS

LiChropur™ (Cat#5.33004) and dichloromethane (Cat#270997) were from Sigma-Aldrich.

EquiSPLASH™ (Cat#330731) and UltimateSPLASH® ONE Mass Spec Standard

(Cat#330820) were from Avanti Polar Lipids Inc. Methyl-tert-butyl ether (MTBE, Cat#34875),

ammonium formate (NH4HCO2, Cat#55674) and formic acid (Cat#94318) were from

Honeywell. Water (Cat#455), methanol (Cat#1428), acetonitrile (Cat#2697) and 2-propanol

(Cat#1178) were from CHEMSOLUTE®.

Lipid extraction from whole testis for lipidomic analysis

Lipid extraction from testes was performed using the methyl-tert-butyl ether (MTBE) method[55].

In short, approximately 100 mg of tissue was weighed into Precellys Lysing Kit tubes

(P000912-LYSK1-A, Bertin Technologies) and 1 mL methanol was added. Samples were kept

at 4˚C and homogenized using a Precellys® 24 homogenizer (5,500 rpm, 3 times for 30 seconds,

with a pause of 20 seconds between runs). Next, 500 μL of homogenate were transferred to a

10 mL screw-top glass vial (Chromacol 10-SV, Thermo) and spiked with internal standards

(50 μL EquiSPLASH™ diluted 1:100 and 50 μL UltimateSPLASH® ONE Mass Spec

Standard diluted 1:10, Avanti). Methanol (500 µL) and MTBE (3,330 µL) were added, and

samples were vortexed using an Ohaus Multi-Tube Vortex Mixer (2,500 rpm for 2 minutes,

pulsed), followed by incubation at 4˚C for 20 minutes. A second vortexing step was performed

(2,500 rpm for 2 minutes, pulsed) and samples were incubated at 4˚C for an additional 20

minutes. Ice-cold water (830 µL) was added, samples were vortexed (2,500 rpm for 3 minutes,

pulsed), and then centrifuged at 526 × g for 10 minutes at 4˚C. The upper phase was transferred

to a new screw-top vial. To the lower phase, MTBE (1,290 µL), methanol (388 µL), and water

(323 µL) were added for re-extraction, followed by vortexing (2,500 rpm for 3 minutes, pulsed)

and centrifugation at 526 × g for 10 minutes at 4˚C. Upper phases were combined and

evaporated to dryness using a RapidVap (Labconco) with the following settings: 45% speed,

35˚C, 250 mbar for 30 minutes; 45% speed, 35˚C, 150 mbar for 20 minutes; and 45% speed,

40˚C, 50 mbar for 50 minutes. Dried extracts were resuspended in 200 µL 2-propanol and

transferred to an amber 300 µL fixed insert vial (Chromacol 03-FIRV(A), Thermo) for LC–

MS/MS analysis. A pooled sample was prepared by combining 7.5 µL of each extract. All

solvents were supplemented with 0.01% (w/v) butylated hydroxytoluene (BHT) and cooled on

ice prior to lipid extraction.

Lipidomic analysis

Lipidomics was performed using hydrophilic interaction chromatography (HILIC) coupled to

triple quadrupole mass spectrometry. HILIC was carried out on a SCIEX Triple Quad™ 7500

system equipped with a Luna® NH2 column (100 × 2 mm; 3 µm, 100 Å, Phenomenex). Lipids

were separated by gradient elution using solvent A (acetonitrile/dichloromethane, 93:7, v/v)

and solvent B (acetonitrile/water, 50:50, v/v), both containing 5 mM ammonium acetate.

Solvent B was adjusted to pH 8.2 with NH3 (15%). Separation was performed at 35˚C using

the following gradient: 0 to 2 minutes, 0% B isocratic (flow rate 0.2 mL/min); 2 to 11 minutes,

0% to 40% B (flow rate 0.5 mL/min); 11 to 11.5 minutes, 40% to 70% B (flow rate 0.5

mL/min); 11.5 to 12.5 minutes, 70% to 100% B (flow rate 0.5 mL/min); 12.5 to 15 minutes,

100% B isocratic (flow rate 0.5 mL/min); 15 to 15.1 minutes, 100% to 0% B, followed by re-

equilibration at 0% B for 2.5 minutes. The triple quadrupole mass spectrometer was operated

with the following settings: TEM 500˚C, GS1 45, GS2 70, CUR 45, CAD 9, IS −3,500 V. Data

were acquired in multiple reaction monitoring (MRM) mode using a published assay (SCIEX

technical note RUO-MKT-02-8477-C). For quantification, the area under the curve for each

precursor mass to fragment mass transition was integrated using SCIEX OS software and

normalized to appropriate internal standards from UltimateSPLASH® ONE (Avanti),

including PC(17:0_22:4(d5)), PE(17:0_18:1(d5)), LPC(17:0(d5)), and LPE(17:0(d5)). Isotopic

correction was performed using LICAR[56].

Round spermatid (RS) isolation

Isolation was carried out according to a previously established protocol[57]. Briefly, Fresh

testicular tissue was harvested from young or aged male mice followed by removal of the

epididymis and tunica albuginea. A single cell suspension was obtained via serial digestion

with Collagenase IV (1mg/ml, Sigma) and Trypsin (0.6 mg/ml, Worthington) / DNAse I (>3.2

kU/ml, Sigma) dissolved in Krebs bicarbonate buffer. The cell suspension was loaded on a 1-

5% BSA / Krebs buffer gradient and allowed to sediment for 1.5 hour on ice at 1 unit gravity.

Twenty-eight fractions of 1 mL were collected and analyzed for purity using PNA (1:2,000,

Rhodamine, Vector Laboratories) and DAPI (1:1,000, Sigma). Only fractions #4-7 were

routinely enriched with RSs (>80%, Extended Data Fig. 3). Therefore, in subsequent isolations,

fractions #4-7 were collected and pooled. The fractions were washed with Krebs buffer to

remove excess BSA and immediately used for RNA and protein isolation.

Simultaneous RNA and Protein isolation

RNA and protein were simultaneously isolated using the AllPrep DNA/RNA/Protein mini kit

(Qiagen) according to the manufacturer's protocol, with one minor modification: the protein

fraction was air dried and then vortexed for 1 minute in SDS buffer (containing 4% SDS, 20%

sucrose, 10 mM Tris-HCl pH 6.8 and 1× ethylenediaminetetraacetic acid (EDTA)-free protease

inhibitor (Roche), followed by incubation at 37°C for 30 minutes. The protein sample was

vortexed and pelleted via centrifugation. Supernatant was transferred to a new tube and protein

concentration was determined using the bicinchoninic acid assay (BCA, Nacalai-Tesque).

RNA was assessed for concentration and purity using the Nanodrop (Thermofisher). Isolated

RNA and protein were stored at -80°C until further use.

cDNA synthesis and quantitative polymerase chain reaction (qPCR)

RNA isolated from pooled round spermatid fractions (1 µg) was reverse transcribed into cDNA

using the SuperScript III™ First-Strand Synthesis System for RT-PCR (Invitrogen). cDNA (2

µl, 1:10 diluted) was subsequently used for qPCR using Taq Pro Universal SYBR qPCR Master

Mix (Vazyme). Primer sequences used in this study can be found in Extended Data Table 2.

Assay specificity was verified by single-peak melt curves and a single band of the expected

size on agarose gel. For *Gpx4* transcript analysis, three qPCR assays were used[39]: a primer set

specific for the *nGpx4* transcript, a primer set specific for the mitochondrial-targeted *Gpx4*

transcript (*mGpx4*), and a primer set amplifying a region shared by mitochondrial and cytosolic

transcripts (*mGpx4+cGpx4*). Ct values were normalized to *ActB* (ΔCt). To estimate the

cytosolic component, expression values were converted to linear scale ($2^{-\Delta Ct}$), and cytosolic

expression was calculated as $\boldsymbol{E_c = E_{m+c} - E_m}$, where $\boldsymbol{E_{m+c}}$ is the signal from the assay

amplifying the shared mitochondrial+cytosolic transcripts and E_m is the mitochondrial-specific

assay. Group comparisons were performed on log2-transformed expression values.

Western blot

Protein samples (5 μg) were separated by SDS-PAGE on 12% TGX™ FastCast™ acrylamide

gels (Bio-Rad) at 150 V for 50 minutes and transferred to 0.2 μm polyvinylidene difluoride

(PVDF) membranes (Trans-Blot Turbo Mini PVDF Transfer Packs; Bio-Rad) using the Trans-

Blot Turbo system (2.5 A, 25 V, 3 minutes). Membranes were blocked in 10% Intercept TBS

blocking buffer (Licor; GPX4, ActB), 5% Blocking One (Nacalai-Tesque; 4HNE, GAPDH),

or 5% skimmed milk in TBST (FTH1, FTL and NCOA4) and incubated overnight at 4˚C with

primary antibodies against GPX4 (1:1,000, ab125066, Abcam), 4HNE, (1:1,000, ab46545,

Abcam), ACTB (1:2,500, sc-47778, Santa Cruz), GAPDH (1:4,000, ab8245, Abcam), FTH1

(1:1,000, ab183781, abcam) FTL (1:1,000, ab69090, abcam) and NCOA4 (1:1,000, ab314553,

abcam). After washing with Tris-buffered saline containing 0.1% Tween-20 (TBST),

membranes were incubated for 1 hour at room temperature with HRP-conjugated secondary

antibodies: for GPX4 and ACTB, W4011 (1:10,000) and W4021 (1:5,000) in 10% Intercept;

for 4HNE and GAPDH, both 1:10,000 in 5% Blocking One; and for FTH1, FTL and NCOA4,

both 1:10,000 in 'Can Get Signal?' (Toyobo). Membranes were washed again in TBST, and

signals were visualized using a ChemiDoc Touch Imaging System (Bio-Rad). Band intensities

were quantified in Fiji and normalized to ACTB or GAPDH as specified.

Flow cytometric analysis of mature sperm

The caudal epididymis was excised from two testes, incised 4 times, and transferred to a dish

containing 2 mL prewarmed (37°C) phosphate buffered saline (PBS) with 2% FBS. Viable

mature sperm were allowed to swim out into the medium for 10 minutes in a 37˚C, 5.0% CO_2

incubator. The sperm suspension was filtered through a 40 μm PET cell strainer and stained

with Hoechst (5 μg/mL) for 45 minutes at 35°C in the dark, mixing every 15 minutes. Samples

were kept on ice until analysis, which proceeded within 1 hour. Directly before analysis, 2 mL

cell suspensions were filtered through a 35 μm strainer cap attached to a 5 mL Polystyrene

Round Bottom FACS tube (Falcon). Cells were subjected to flow cytometry using an

LSRFortessa cell analyzer (BD Biosciences). Hoechst was excited using the 355 nm UV laser

and detected using a 450/50 bandpass filter. Forward Scatter (FSC) and Side Scatter (SSC)

were detected using a 488 nm laser and a 488/10 bandpass filter. Flow rate was set to achieve

2,500-3,500 events/second. A minimum of 100,000 events within a specified gate (Fig 3d)

were recorded using the BD FACSDiva Software (BD Biosciences). Data analysis was done

using FlowJo software v10.10.0 (BD Life Sciences).

Statistical analysis

Statistical information for individual experiments is shown in the corresponding figure

legends. Values are presented as mean ± standard deviation. Data were first analysed for

normality using the Shapiro–Wilk test. Depending on distribution and variance, statistical

comparisons between groups were performed using Student's *t*-test, Welch's *t*-test, or the

Mann–Whitney *U* test. Longitudinal effects were analysed using two-way ANOVA (fixed

effects: diet, week and bodyweight; random effect: individual mouse, repeated measures).

Results were considered significant at $P < 0.05$. Statistical analyses were conducted using

GraphPad Prism 10 (GraphPad Software).

Declaration of interest

The authors declare no conflict of interest.

Acknowledgements

We thank the Japanese Society for the Promotion of Science KAKENHI grant number #17K19372, #22K19322, and #25K22536, The Canon Foundation awarded to N.O., the establishment of university fellowships towards the creation of science technology innovation, grant number JPMJFS2102, JPMJSP2114 awarded to J.G., and JST FOREST Program (JPMJFR2213) awarded to E.M. for financial support. M.C. received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement no. GA 884754). We thank Sayaka Makino and Asuka Takehara for animal care and all other members of the Developmental Neuroscience laboratory at Tohoku University for their useful comments. We also thank Prof. Junken Aoki (Graduate School of Pharmaceutical Sciences, The University of Tokyo) for advice on lipidomic analysis and Dr. Ibuki Kusumoto (Laboratory of Food Function Analysis, Tohoku University) for technical support.

Author contributions

Study design: J.G., T.K., W.G., E.M., N.O.; Data collection: J.G., T.K., J.I., L.G., B.H.; Data analysis: J.G.; J.I., B.H., M.A.; Drafted the manuscript: J.G.; Edited the manuscript: T.K., W.G., E.M., M.C., N.O.; Funding acquisition: J.G., E.M., M.C., N.O.

Competing interests

M.C. is co-founder and shareholder of ROSCUE Therapeutics GmbH. The other authors declare no competing interests.

References

1. Johnson, S. L., Dunleavy, J., Gemmell, N. J. & Nakagawa, S. Consistent age-dependent declines in human semen quality: A systematic review and meta-analysis. *Ageing Res. Rev.* **19**, 22–33 (2015).

2. Verón, G. L. *et al.* Impact of age, clinical conditions, and lifestyle on routine semen parameters and sperm kinematics. *Fertil. Steril.* **110**, 68-75.e4 (2018).

3. Evenson, D. P., Djira, G., Kasperson, K. & Christianson, J. Relationships between the age of 25,445 men attending infertility clinics and sperm chromatin structure assay (SCSA®) defined sperm DNA and chromatin integrity. *Fertil. Steril.* **114**, 311–320 (2020).

4. Martins da Silva, S. & Anderson, R. A. Reproductive axis ageing and fertility in men. *Rev. Endocr. Metab. Disord.* **23**, 1109–1121 (2022).

5. Griswold, M. D. Spermatogenesis: The Commitment to Meiosis. *Physiol. Rev.* **96**, 1–17 (2016).

6. Gosden, R. G., Richardson, D. W., Brown, N. & Davidson, D. W. Structure and gametogenic potential of seminiferous tubules in ageing mice. *Reproduction* **64**, 127–133 (1982).

7. Wright, W. W., Fiore, C. & Zirkin, B. R. The effect of aging on the seminiferous epithelium of the brown Norway rat. *J. Androl.* **14**, 110–7 (1993).

8. Silva, J. V. *et al.* Effects of Age and Lifelong Moderate-Intensity Exercise Training on Rats' Testicular Function. *Int. J. Mol. Sci.* **23**, 11619 (2022).

9. Dong, S. *et al.* Testicular aging, male fertility and beyond. *Front. Endocrinol. (Lausanne).* **13**, (2022).

10. Nguyen-Powanda, P. & Robaire, B. Oxidative Stress and Reproductive Function in the Aging Male. *Biology (Basel).* **9**, 282 (2020).

11. Gunes, S., Hekim, G. N. T., Arslan, M. A. & Asci, R. Effects of aging on the male reproductive system. *J. Assist. Reprod. Genet.* **33**, 441–454 (2016).

12. Villalpando-Rodriguez, G. E. & Gibson, S. B. Reactive Oxygen Species (ROS) Regulates Different Types of Cell Death by Acting as a Rheostat. *Oxid. Med. Cell. Longev.* **2021**, 9912436 (2021).

13. Blanco-Rodríguez, J. & Martínez-García, C. Spontaneous germ cell death in the testis of the adult rat takes the form of apoptosis: re-evaluation of cell types that exhibit the ability to die during spermatogenesis. *Cell Prolif.* **29**, 13–31 (1996).

14. Bromfield, E. G. *et al.* Differential cell death decisions in the testis: evidence for an

exclusive window of ferroptosis in round spermatids. *MHR: Basic science of*

reproductive medicine **25**, 241–256 (2019).

15. Oresti, G. M. *et al.* Differentiation-related changes in lipid classes with long-chain and

very long-chain polyenoic fatty acids in rat spermatogenic cells. *J. Lipid Res.* **51**,

2909–2921 (2010).

16. Li, Z., Lange, M., Dixon, S. J. & Olzmann, J. A. Lipid Quality Control and

Ferroptosis: From Concept to Mechanism. *Annu. Rev. Biochem.* **93**, 499–528 (2024).

17. Matzkin, M. E., Calandra, R. S., Rossi, S. P., Bartke, A. & Frungieri, M. B. Hallmarks

of Testicular Aging: The Challenge of Anti-Inflammatory and Antioxidant Therapies

Using Natural and/or Pharmacological Compounds to Improve the Physiopathological

Status of the Aged Male Gonad. *Cells* **10**, (2021).

18. Dixon, S. J. *et al.* Ferroptosis: An Iron-Dependent Form of Nonapoptotic Cell Death.

Cell **149**, 1060–1072 (2012).

19. Mishima, E. *et al.* Recommendations for robust and reproducible research on

ferroptosis. *Nat. Rev. Mol. Cell Biol.* **26**, 615–630 (2025).

20. Dixon, S. J. & Olzmann, J. A. The cell biology of ferroptosis. *Nat. Rev. Mol. Cell Biol.*

25, 424–442 (2024).

21. Friedmann Angeli, J. P. *et al.* Inactivation of the ferroptosis regulator Gpx4 triggers

acute renal failure in mice. *Nat. Cell Biol.* **16**, 1180–1191 (2014).

22. Yang, W. S. *et al.* Regulation of Ferroptotic Cancer Cell Death by GPX4. *Cell* **156**,

317–331 (2014).

23. Doll, S. *et al.* ACSL4 dictates ferroptosis sensitivity by shaping cellular lipid

composition. *Nat. Chem. Biol.* **13**, 91–98 (2017).

24. Galy, B., Conrad, M. & Muckenthaler, M. Mechanisms controlling cellular and

systemic iron homeostasis. *Nat. Rev. Mol. Cell Biol.* **25**, 133–155 (2024).

25. Conrad, M. & Pratt, D. A. The chemical basis of ferroptosis. *Nat. Chem. Biol.* **15**,

1137–1147 (2019).

26. Saito, H., Hara, K., Kitajima, S. & Tanemura, K. Effect of vitamin E deficiency on

spermatogenesis in mice and its similarity to aging. *Reproductive Toxicology* **98**, 225–

232 (2020).

27. Bensoussan, K., Morales, C. R. & Hermo, L. Vitamin E deficiency causes incomplete

spermatogenesis and affects the structural differentiation of epithelial cells of the

epididymis in the rat. *J. Androl.* **19**, 266–88 (1998).

28. Endo, T. *et al.* Multiple ageing effects on testicular/epididymal germ cells lead to
decreased male fertility in mice. *Commun. Biol.* **7**, 16 (2024).

29. Chen, Y.-F. *et al.* Cisd2 deficiency drives premature aging and causes mitochondria-
mediated defects in mice. *Genes Dev.* **23**, 1183–1194 (2009).

30. Luo, L. *et al.* Aging and the Brown Norway Rat Leydig Cell Antioxidant Defense
System. *J. Androl.* **27**, 240–247 (2006).

31. Liao, C.-H. *et al.* Optimizing a Male Reproductive Aging Mouse Model by d-
Galactose Injection. *Int. J. Mol. Sci.* **17**, 98 (2016).

32. Kimura, A. K. & Kim, H.-Y. Phosphatidylserine synthase 2: high efficiency for
synthesizing phosphatidylserine containing docosahexaenoic acid. *J. Lipid Res.* **54**,
214–222 (2013).

33. Sulc, A. *et al.* MALDI Imaging Mass Spectrometry Reveals Lipid Alterations in
Physiological and Sertoli Cell-Only Syndrome Human Testicular Tissue Sections. *Int.*
J. Mol. Sci. **25**, (2024).

34. Braverman, N. E. & Moser, A. B. Functions of plasmalogen lipids in health and
disease. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1822**,
1442–1452 (2012).

35. Imai, H. *et al.* Depletion of Selenoprotein GPx4 in Spermatocytes Causes Male
Infertility in Mice. *Journal of Biological Chemistry* **284**, 32522–32532 (2009).

36. Schneider, M. *et al.* Mitochondrial glutathione peroxidase 4 disruption causes male
infertility. *The FASEB Journal* **23**, 3233–3242 (2009).

37. Conrad, M. *et al.* The Nuclear Form of Phospholipid Hydroperoxide Glutathione
Peroxidase Is a Protein Thiol Peroxidase Contributing to Sperm Chromatin Stability.
Mol. Cell. Biol. **25**, 7637–7644 (2005).

38. Mishima, E. *et al.* DHODH inhibitors sensitize to ferroptosis by FSP1 inhibition.
Nature **619**, E9–E18 (2023).

39. Borchert, A. *et al.* The Role of Phospholipid Hydroperoxide Glutathione Peroxidase
Isoforms in Murine Embryogenesis. *Journal of Biological Chemistry* **281**, 19655–
19664 (2006).

40. Ngo, V. & Duennwald, M. L. Nrf2 and Oxidative Stress: A General Overview of
Mechanisms and Implications in Human Disease. *Antioxidants (Basel).* **11**, (2022).

41. Lasiene, K. *et al.* Age-related morphological peculiarities of human testes. *Folia*
Morphol. (Warsz). **80**, 122–126 (2021).

42.   Jiang, H. *et al.* Quantitative histological analysis and ultrastructure of the aging human
testis. *Int. Urol. Nephrol.* **46**, 879–885 (2014).

43.   Barnes, C. J., Covington, B. W. & Lee, M. Effect of Aging and Dietary Restriction on
Rat Testicular Germ Cell Apoptosis. *J. Gerontol. A Biol. Sci. Med. Sci.* **54**, B199–
B204 (1999).

44.   Morales, E. *et al.* Effect of ageing on the proliferation and apoptosis of testicular germ
cells in the Syrian hamster Mesocricetus auratus. *Reprod. Fertil. Dev.* **15**, 1–10 (2003).

45.   Mortensen, M. S., Ruiz, J. & Watts, J. L. Polyunsaturated Fatty Acids Drive Lipid
Peroxidation during Ferroptosis. *Cells* **12**, (2023).

46.   Kagan, V. E. *et al.* Oxidized arachidonic and adrenic PEs navigate cells to ferroptosis.
Nat. Chem. Biol. **13**, 81–90 (2017).

47.   Sutovsky, P. *et al.* A putative, ubiquitin-dependent mechanism for the recognition and
elimination of defective spermatozoa in the mammalian epididymis. *J. Cell Sci.* **114**,
1665–1675 (2001).

48.   Cocquet, J. *et al.* A Genetic Basis for a Postmeiotic X Versus Y Chromosome
Intragenomic Conflict in the Mouse. *PLoS Genet.* **8**, e1002900 (2012).

49.   Braun, R. E., Behringer, R. R., Peschon, J. J., Brinster, R. L. & Palmiter, R. D.
Genetically haploid spermatids are phenotypically diploid. *Nature* **337**, 373–376
(1989).

50.   Bhutani, K. *et al.* Widespread haploid-biased gene expression enables sperm-level
natural selection. *Science (1979).* **371**, (2021).

51.   You, Y.-A. *et al.* Sex chromosome-dependent differential viability of human
spermatozoa during prolonged incubation. *Human Reproduction* **32**, 1183–1191
(2017).

52.   Rahman, M. S. & Pang, M.-G. New Biological Insights on X and Y Chromosome-
Bearing Spermatozoa. *Front. Cell Dev. Biol.* **7**, 388 (2019).

53.   Kimura, R., Yoshizaki, K. & Osumi, N. Dynamic expression patterns of Pax6 during
spermatogenesis in the mouse. *J. Anat.* **227**, 1–9 (2015).

54.   Tatehana, M., Kimura, R., Mochizuki, K., Inada, H. & Osumi, N. Comprehensive
histochemical profiles of histone modification in male germline cells during meiosis
and spermiogenesis: Comparison of young and aged testes in mice. *PLoS One* **15**,
e0230930 (2020).

55.   Matyash, V., Liebisch, G., Kurzchalia, T. V., Shevchenko, A. & Schwudke, D. Lipid

738   extraction by methyl-tert-butyl ether for high-throughput lipidomics. *J. Lipid Res.* **49**,

739   1137–1146 (2008).

740   56.   Gao, L. *et al.* LICAR: An Application for Isotopic Correction of Targeted Lipidomic

741   Data Acquired with Class-Based Chromatographic Separations Using Multiple

742   Reaction Monitoring. *Anal. Chem.* **93**, 3163–3171 (2021).

743   57.   Da Ros, M., Lehtiniemi, T., Olotu, O., Meikar, O. & Kotaja, N. Enrichment of

744   Pachytene Spermatocytes and Spermatids from Mouse Testes Using Standard

745   Laboratory Equipment. *Journal of Visualized Experiments*

746   https://doi.org/10.3791/60271 (2019) doi:10.3791/60271.

Fig. 1: Age-related alterations in mouse testis germ cell populations.

a–c, Body weight (**a**), combined testes weight (**b**) and gonadosomatic index (combined testes

weight/body weight; **c**) in young (3 months) and aged (18 months) males. **d**, Schematic of

spermatogenesis highlighting undifferentiated and differentiating spermatogonia, pachytene

spermatocytes, round spermatids (RSs) and spermatozoa. **e**, Representative stage VII–VIII

seminiferous tubules stained for PLZF (undifferentiated spermatogonia), KIT (differentiating

spermatogonia), SCP3 (pachytene spermatocytes), PNA (RSs) and SOX9 (Sertoli cells); nuclei

counterstained with DAPI. Scale bar, 50 μm. f–j, Quantification of PLZF$^+$ (**f**), KIT$^+$ (**g**), SCP3$^+$

(**h**), PNA$^+$ (**i**) and SOX9$^+$ (**j**) cells, normalized to tubule circumference (**f**: 20 images per mouse;

g–j: 10 images per mouse). **k**, Representative tubules showing CASP3$^+$/TUNEL$^+$ and

CASP3$^-$/TUNEL$^+$ events in peripheral and luminal compartments; nuclei counterstained with

Hoechst. Scale bar, 50 μm. **l–o**, Quantification of CASP3$^+$/TUNEL$^+$ and CASP3$^-$/TUNEL$^+$

cells in the periphery (**l**, **m**) and lumen (**n**, **o**), normalized to 100 tubules. Data are mean ± s.d.;

dots represent mice. Statistics: two-tailed Student's *t*-test (**a–c**, **f**, **h**, **i**, **l**, **n**, **o**) or Mann–Whitney

U test (**g**, **j**, **m**), exact *P* values are given. $N = 6$ mice per group. Illustration in **d** was created in

BioRender, Germeraad, J. (2026) https://BioRender.com/8iim1z1

Fig. 2: Increased lipid peroxidation in aged whole testis and round spermatids.

a, Workflow for isolation of RS-enriched fractions and extraction of RNA/protein using a

modified density-gradient approach. **b**, Representative immunoblots of 4HNE adducts and

quantification, normalized to GAPDH, black triangle indicates quantified band. **c–h**, RS

mRNA expression of *Alox15* (**c**), *Acsl4* (**d**), *Acsl6* (**e**) and *Gpx4* isoforms (*nGpx4* (**f**), *mGpx4*

(**g**), *cGpx4* (**h**)), shown as fold change relative to *ActB*. **i**, Representative immunoblots and

quantification of GPX4 protein, normalized to ACTB. **j**, Schematic outlining NRF2 activation

and initiating its target gene expression. Oxidative stress activates NRF2 ,via KEAP1 release,

to induce antioxidant/HO-1 programs, while iron uptake (TFR1), storage (FTH1/FTL), and

ferritinophagy (NCOA4) regulate the labile iron pool (LIP) **k–o**, RS mRNA expression of *Nrf2*

(**k**), *Sod1* (**l**), *Cat* (**m**), *HO-1* (**n**) and *Tfr1* (**o**), relative to *ActB*. **p–s**, Representative

immunoblots (**p**) and quantification of FTH1 (**q**), FTL (**r**) and NCOA4 (**s**). **t**, Summary

schematic of aging-associated changes across lipid peroxidation, redox defence and iron

metabolism (red, increased; blue, decreased; gray boxes, protein; white boxes, mRNA). Data

are mean ± s.d.; dots represent mice. Statistics: Welch's *t*-test (**a**); two-tailed Student's *t*-test

(**b–e**, **g–i**, **k**, **l**, **n**, **q–s**); Mann–Whitney *U* test (**f**, **m**, **o**), exact P values are given. $N = 6$ mice

per group.

Fig. 3: Aging alters the sex chromosome ratio in round spermatids and mature sperm.

a, **b**, FISH in RSs showing Y-chromosomes (green; closed arrowhead) and X-chromosomes

(red; open arrowhead) signals with Hoechst counterstain (**a**) and quantification of Y/X ratio

(Young: n = 6,282±100 Y-RSs, n = 6,188±113 X-RSs; mean Y/X ratio 1.02; Aged: n =

6,694±85 Y-RSs, n = 6,056±98 X-RSs; mean Y/X ratio 1.11) (**b**). Scale bar, 5 μm. **c**, Sperm

collection and DNA staining workflow: cauda epididymides were excised and sperm was

collected via swim-out (37 °C, 10 min) followed by Hoechst 33342 staining (45 min) before

flow cytometry. **d**, Gating strategy to resolve X- and Y-bearing sperm: debris exclusion (FSC-

A vs SSC-A), singlet selection (FSC-H vs FSC-W and Hoechst Blue vs SSC-A), and

histogram-based separation by Hoechst Blue intensity. **e**, **f**, Representative Hoechst Blue

distributions (young vs aged and overlay normalized to mode) (**e**) and Y/X ratio quantification

(young: n = 3,382,147 ± 140,662 Y-sperm vs n = 3,293,224 ± 126,522 X-sperm; mean Y/X

1.02; aged: n = 3,294,779 ± 156,878 Y-sperm; n = 3,032,412 ± 145,748 X-sperm; mean Y/X

1.09) (**f**). Data are mean ± s.d.; dots represent mice. Statistics: two-tailed Student's t-test (**b**, **f**).

Exact P values are shown. N = 6 mice per group. Illustration in **c** was created in BioRender,

Germeraad, J. (2026) https://BioRender.com/j1osdbj

Fig. 4: Vitamin E deficient feeding recapitulates age-associated testis phenotypes.

a, Experimental design: young males (1 month) fed control or vitamin E–deficient diet (VE(−))

for 70 days. **b**, body weight was recorded weekly for 10 weeks in the same cohort of mice. **c**,

d, combined testes weight (**c**) and gonadosomatic index (combined testes weight/body weight;

d). **e**, Representative stage VII–VIII seminiferous tubules stained for PLZF, KIT, SCP3, PNA

and SOX9: counterstained with DAPI. **f–j**, Quantification of PLZF⁺ (**f**), KIT⁺ (**g**), SCP3⁺ (**h**),

PNA⁺ (**i**) and SOX9⁺ (**j**) cells, normalized to tubule circumference (**f**: 20 images per mouse; **g**–

j: 10 images per mouse). **k**, Representative tubules showing CASP3⁺/TUNEL⁺ cells in the

periphery and CASP3⁻/TUNEL⁺ cells in the lumen. **l**, **m**, Quantification of CASP3⁺/TUNEL⁺

and CASP3⁻/TUNEL⁺ cells in peripheral (**l**) and luminal compartments (**m**). **n**, qPCR of fatty-

acid metabolism, GPX4 defence, NRF2-response and iron-homeostasis transcripts, relative to

ActB. **o**, **p**, Representative immunoblots and quantification of 4HNE (**o**) and GPX4 (**p**),

normalized to GAPDH, black triangle indicates quantified band. **q–t**, Representative

immunoblots (**q**) and quantification of FTH1 (**r**), FTL (**s**) and NCOA4 (**t**). **u**, FISH in RSs with

X (red) or Y (green) signals and Y/X ratio (control: n = 6,390±125 Y-RSs vs n = 6,274±127

X-RSs; mean Y/X ratio 1.02; VE(−): n = 6,548±222 Y-RSs vs n = 6,186±212 X-RSs; mean

Y/X ratio 1.09). Scale bars: **e**, 50 µm (overview) and 5 µm (inset); **k**, 50 µm; **u**, 5 µm. Data are

mean ± s.d.; dots represent mice except for **b** where it represents the mean of each group.

Statistics: two-way repeated measures-ANOVA with Geisser–Greenhouse correction (**b**); two-

tailed Student's *t*-test (**c, d, f–i, m–p, r–t, u**); Mann–Whitney *U* test (**j, l, n**), exact *P* values are

given. Sample sizes: **b–d**, *N* = 9 per group; all other panels, *N* = 6 per group.

Fig. 5: Vitamin E supplemented feeding rescues age-associated phenotypes within the

testis.

a, Experimental design: aged (16-months) males fed control or vitamin E–supplemented diet

(VE+) for 70 days. **b**, body weight was recorded weekly for 10 weeks in the same cohort of

mice. **c, d**, combined testes weight (**c**) and gonadosomatic index (combined testes weight/body

weight; **d**). **e**, Representative stage VII–VIII seminiferous tubules stained for PLZF, KIT, SCP3,

PNA and SOX9: nuclei counterstained with DAPI. **f–j**, Quantification of PLZF$^+$ (**f**), KIT$^+$ (**g**),

SCP3$^+$ (**h**), PNA$^+$ (**i**) and SOX9$^+$ (**j**) cells, normalized to tubule circumference (**f**: 20 images

per mouse; **g–j**: 10 images per mouse). **k**, Representative tubules showing CASP3$^+$/TUNEL$^+$

peripheral and CASP3$^-$/TUNEL$^+$ luminal events. **l,m**, Quantification of CASP3$^+$/TUNEL$^+$ and

CASP3$^-$/TUNEL$^+$ cells in peripheral (**l**) and luminal compartments (**m**). **n**, qPCR of fatty-acid

metabolism, GPX4 defence, NRF2-response and iron-homeostasis transcripts, relative to ActB.

o, **p**, Representative immunoblots and quantification of 4HNE (**o**) and GPX4 (**p**), normalized

to GAPDH, black triangle indicates quantified band. **q–t**, Representative immunoblots (**q**) and

quantification of FTH1 (**r**), FTL (**s**) and NCOA4 (**t**). **u**, FISH in RSs with X (red) or Y (green)

signals and Y/X ratio (control: *n* = 6,527±135 Y-RSs vs *n* = 5,950±156 X-RSs; Y/X ratio mean

1.10; VE(+): *n* = 5,772±156 Y-RSs vs *n* = 5,702±149 X-RSs; mean Y/X ratio 1.01). Scale bars:

e, 50 µm (overview) and **5** µm (inset); **k**, 50 µm; **u**, 5 µm. Data are mean ± s.d.; dots represent

mice except for **b** where it represents the mean of each group. Statistics: two-way repeated

measures-ANOVA with Geisser–Greenhouse correction (**b**); two-tailed Student's *t*-test (**c, f, g,**

i, l, m, n, p, r, s, u); Welch's *t*-test (**o**); Mann–Whitney *U* test (**d, h, j, t**), exact *P* values are

given. Sample sizes: **b–d**, *N* = 9 per group; all other panels, *N* = 6 per group.

Extended data Fig. 1: Age-associated cell death is increased in seminiferous tubules

a, Representative images of tubules containing TUNEL$^+$ cells in young and aged mice.

Asterisks indicate tubules containing TUNEL$^+$ cells. Scale bar: 200 µm. **b**, Percentage of

seminiferous tubules containing cells TUNEL$^+$ cells relative to the total number of tubules

within a section. **c**, Representative images of CASP3$^+$/TUNEL$^+$ round spermatids in young and

aged testis sections. Scale bar: 50μm. Data is presented as mean ± s.d. Significance was

determined by two-tailed Student's *t*-test (**b**). Exact *P* value is given. Sample size: *N* = 6 in

each group. Each dot represents one mouse.

Extended data Fig. 2: Aging shifts phosphatidylcholine composition toward plasmalogen

PC

a-d, Heatmaps showing relative abundances of individual phospholipid species in young and

aged whole testis: diacyl phosphatidylcholine (PC; **a**), diacyl phosphatidylethanolamine (PE;

b), plasmalogen phosphatidylcholine (PCp; **c**) and plasmalogen phosphatidylethanolamine

(PEp; **d**). Columns represent biological replicates and rows represent lipid species. Values are

scaled to show relative differences across samples. **e, f,** mRNA level quantification analysis of

Elovl2 (**e**) and *Elovl5* (**f**) expression in young and aged round spermatids. Data is presented as

mean ± s.d. Significance was determined by two-tailed Student's *t*-test (**e, f**). Sample size: *N* =

6 in each group. Each dot represents one mouse.

Extended data Fig. 3: Enrichment profile of round spermatids across fractions.

a, Representative image of the enriched cell suspension (fraction #3) stained with peanut

agglutinin (PNA) to label the acrosome and DAPI to label nuclei. Scale bar: 20μm. **b,**

Representative cellular composition of fractions showing the percentage of round spermatids

versus other germ-cell populations (including spermatogonia and other cells). Each bar

represents one fraction; stacked colors indicate the relative contribution of round spermatids

(orange) and other cells (green).

Extended data Fig. 4: GPX4 localization and ferritin expression in testicular aging

a, Representative images of GPX4 localization in seminiferous tubules from young and aged

testes, with DAPI counterstain. Scale bar: 50μm. **b, c,** mRNA level quantification of iron-

handling transcripts in round spermatids from young and aged mice: *Fth1* (**b**) and *Ftl* (**c**),

relative to *ActB*. Data are presented as mean ± s.d. Significance was determined by two-tailed

Student's *t*-test (**b, c**). Exact *P* values are given. Sample size: *N* = 6 in each group. Each dot

represents one mouse.

Extended data Fig. 5: Dietary vitamin E modulates cell death in young and aged testes

a, Representative TUNEL staining in testis sections from young control (3-months), vitamin

E–deficient (VE(−)), aged (18-months) control and aged vitamin E–supplemented (VE(+))

mice; asterisks mark tubules containing TUNEL$^+$ cells. Scale bar: 200 µm. **b, c,** Quantification

of the percentage of tubules containing TUNEL$^+$ cells in young mice (control vs VE(−); **b**) and

aged mice (control vs VE(+); **c**). **d,** Representative images of CASP3$^+$/TUNEL$^+$ round

spermatids in young and aged testis sections. Scale bars: 50µm. Data are presented as mean ±

s.d. Significance was determined by two-tailed Student's *t*-test (**b, c**). Exact *P* values are given.

Sample size: *N* = 6 in each group. Each dot represents one mouse.

Extended Data Fig. 6: Age-linked round spermatid loss skews Y/X output.

Graphical summary of the model supported by this study, linking aging-associated lipid redox

stress to altered haploid sex-chromosome output. Along the male germline lineage, cell death

is enriched in pre-meiotic and meiotic stages, whereas post-meiotic round spermatids (RSs)

exhibit heightened lipid peroxidation and ferroptosis-like vulnerability. In young mice, lipid

peroxidation is low and X- and Y-bearing RSs are maintained, resulting in a balanced Y/X

output. In aged mice, lipid peroxidation is increased and RS loss is associated with a skewed

Y/X chromosome ratio. Vitamin E deficiency phenocopies the aging-associated increase in

lipid peroxidation and RS, whereas vitamin E supplementation reduces lipid peroxidation and

restores Y/X balance. Arrows denote the direction of change; the lightning symbol indicates

RS loss. Illustration was created in BioRender. Germeraad, J. (2026)

https://BioRender.com/k04h4kg

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

a

b

c

Extended data 1

Extended data 2

a

b

Extended data 3

a

b

ns

Fth1 fold change relative to *ActB*

Young   Aged

c

P = 0.026

Ftl fold change relative to *ActB*

Young   Aged

Extended data 4

Extended data 5

Age-linked round spermatid loss skews Y/X output

Extended data 6