The population structure of invasive *Lantana camara* is shaped by its mating

system

P. Praveen[1*], Rajesh Gopal[2], Uma Ramakrishnan[1*]

[1] National Centre for Biological Sciences - TIFR, GKVK Campus, Bellary Road,

Bangalore 560065, India

[2] Global Tiger Forum – East of Kailash, New Delhi, India

*Corresponding authors: praveenprakash@ncbs.res.in, uramakri@ncbs.res.in

24 Keywords: Invasive species, Invasion-genomics, self-fertilization, population genetic

25 simulations, India, *Lantana camara*

Abstract

Over the last century, invasive species have emerged as an important driver of global

biodiversity loss. Many invasive species have low genetic diversity in the invaded

habitats, owing to the demographic bottleneck during introduction. *Lantana camara* is

one of the hundred most problematic invasive species globally. Despite its ecological

importance in many countries, our understanding of the genetic diversity patterns of

this plant remains poor. Previous studies hypothesize that invasive *L. camara* is a

species complex with a hybrid origin, but this remains untested. We investigated the

population genetic patterns of this invasive species by sampling 359 plants that

represented a spectrum of flower colour variants across 36 locations, spanning most

of the biogeographic regions across India. Analyses of the population structure using

19,008 SNPs revealed that *L. camara* in India exhibits a strong genetic structure.

Interestingly, the structuring pattern does not exhibit a strong correlation with

geography. In the structure analysis, individuals with similar flower colours clustered

together regardless of their location of origin. The genetic distance between most of

the individuals was low, indicating the absence of multiple species. A high inbreeding

coefficient and a low proportion of heterozygous sites observed suggested that the

strong structure could be due to self-fertilization. Thus we infer that *L. camara* exists

as homozygous inbred lines formed by self-fertilization and that these inbred lines

could be associated with distinct flower colours. Together, this would explain the

correlation between flower colour and genetic structure, and the lack of geographic

structure. These results refute the argument that *L. camara* is a species complex and

emphasize the importance of the mating system in shaping the patterns of diversity in

this invasive species. Our findings highlight a hitherto unknown role for mating

systems in invasive species, furthering our understanding of evolution in invasive

species.

1. Introduction

Invasive species are one of the major causes of ongoing biodiversity loss and

extinctions in the Anthropocene (Linders et al., 2019; Mollot et al., 2017).

Understanding the ecological and evolutionary underpinnings of invasion success is

critical to controlling the emergence of new invasive species. Evolutionary and

population genetic theory suggests that among other factors, genetic variation is

critical to demographic success and range expansion (Ørsted et al., 2019).

Populations that have undergone a demographic bottleneck are expected to have

low genetic variation, which can negatively impact their ability to survive and adapt

(Olazcuaga et al., 2023).

Invasive species typically undergo demographic bottlenecks during

colonization (Alves et al., 2022). Low propagule pressure (the number of individuals

introduced) leads to low effective population size. Consequently, invasive populations

are established under conditions of limited genetic variation (Dlugosch & Parker,

2008; Simberloff, 2009). Despite the theoretical expectation of low genetic variation,

invasive populations paradoxically demonstrate the ability to adapt, thrive and

reproduce in large numbers in these novel habitats. This phenomenon is called the

genetic paradox of biological invasions(Estoup et al., 2016). While extensively

debated, some studies support the presence of such low genetic variation (Alves et

al., 2022) while others contradict it (Bastien Lavergne & Molofsky, 2007; Zhang et al.,

2021). The inherent complexities of human-mediated introduction like the choice of

source population, number of introductions, artificial hybridizations etc. along with

evolutionary forces make the patterns of genetic variation in invasive populations

complicated (Roman & Darling, 2007). Several factors such as propagule pressure,

hybridizations, gene flow, mating system etc. play an important role in determining

the amount of genetic variation in invasive populations (Jaspers et al., 2021).

The introduction of many individuals either through single or multiple events

can bring high variation into the novel environment (Bastien Lavergne & Molofsky,

2007). The source populations from which these individuals originate can influence

the patterns of genetic diversity (Ryan et al., 2019) in new populations. Gene flow

between multiple invasive populations can homogenize genetic variation (Vallejo-

Marín et al., 2021). Intraspecific hybridizations between genetically distinct individuals

can lead to the formation of new genotypes with novel allelic combinations, while

hybridization with native species can aid in rapid adaptation (Blair & Hufbauer, 2010;

Mesgarana et al., 2016). In plant populations, mating systems play a role in

determining the pattern of genetic diversity (Hamrick et al., 1992; Ness et al., 2010;

Schoen & Brown, 1991). Theoretically, cross-fertilizing species have high

heterozygosity, and hence high genetic variation (Koelling et al., 2011). Conversely,

with self-fertilization heterozygosity decreases by 50% every generation (Hedrick

2011). This leads to high genome-wide homozygosity and individuals exist as

homozygous inbred lines (Guo et al., 2008; Pérez De La Vega & Garda, 1997). In

self-fertilizing species, neutral diversity is low due to low effective population size

(Foxe et al., 2009; Guo et al., 2008; Koelling et al., 2011), and it results in high

genetic drift. The combination of low gene flow, high drift and high homozygosity

leads to a strong genetic structure in self-fertilizing species compared to cross-

fertilizing species (Edwards et al., 2019; Koelling et al., 2011; Mairal et al., 2023).

While mating systems play an important role in determining genetic diversity, to the

best of our knowledge, their role in invasive species has not been specifically

investigated.

Here we present the genetic diversity patterns of the globally invasive plant

species, *Lantana camara* (Lantana hereafter) in India. Lantana is a perennial shrub

native to Central and South America. During the colonial era, this species was

introduced into many countries across the globe, like India, Australia and South

Africa as a garden plant (Kannan et al., 2013). Currently, it is one of the most

important invasive species in India with a significant impact on the native

ecosystems. On average, Lantana threatens almost 44 per cent of the forest areas in

India (Mungi et al., 2020). In the Bandipur tiger reserve in southern India, Lantana

occupies at least 40 per cent of the land area (R et al., 2016). Unlike many invasive

species, Lantana exhibits extensive phenotypic variation in India (Negi, Sharma, &

Vishvakarma, 2019), with the most striking being the variation in flower colour. By

1900, approximately 16 species of the genus Lantana were reported in India,

potentially suggesting multiple introductions. The taxonomy of the genus Lantana is

complicated due to the presence of polyphyletic groups (Lu-Irving et al., 2021). In the

taxonomy, many flower colour variants of Lantana are classified as distinct species

(Sanders, 2006). However, genetic validation of this classification, focusing on

different flower colour variants has not been done. It is possible that active

hybridization to produce new hybrids took place in botanical gardens (Goyal &

Sharma, 2015a; Urban et al., 2011). Previous studies have hypothesized that

invasive Lantana is a species complex, potentially formed from such hybridization

events (Goyal & Sharma, 2015b; Sanders, 2006). Variation in Lantana flower colour

lends support to the species complex hypothesis. As a result, the invasive Lantana is

sometimes referred to as *Lantana camara* sensu lato (Goyal & Sharma, 2015c).

Detailed genetic studies are important in checking this hypothesis and characterising

different species of this genus.

Despite its status as one of the most pervasive invasive species globally, our

understanding of its biology and genetics remains limited. It is further complicated by

uncertainties regarding its species identity. Investigating the genetic diversity patterns

of Lantana presents an opportunity to improve our understanding regarding the

uncertainties about the species identity, as well as the genetic paradox of biological

invasions. This can help in improving our understanding of the taxonomy of Lantana

itself. A previous population genetic study using microsatellite markers suggests that

Lantana populations in India are genetically homogeneous (Ray & Quader, 2014).

However, as discussed earlier, its ecological and demographic success is very high

along with the high phenotypic variation. We use thousands of genome-wide Single

Nucleotide Polymorphism (SNP) markers and genetic simulations to investigate the

patterns of genetic diversity, spatial genetic structure and the factors shaping these

patterns in Lantana populations in India. We use India as an example of a large,

ecologically diverse landmass where Lantana has been highly successful. More

specifically, we investigate i) the genetic diversity and population structure of Lantana

populations in India, and ii) the evolutionary forces shaping the genetic diversity

patterns of Lantana.

2. Material and Methods

2.1 Study system

   Lantana camara is considered to be one of the hundred most invasive species in the world (IUCN 2000). It was introduced across the globe as a garden plant because of its strikingly attractive flowers. The earliest recorded presence of Lantana in India comes from the British botanical garden in Kolkata in 1807 (Kannan et al., 2013). Currently, it is widely distributed in India occupying almost every terrestrial habitat (Mungi et al., 2020). In India, Lantana exhibits extensive phenotypic variation with the most remarkable being the variation in the flower colour (Negi, Sharma, Vishvakarma, et al., 2019). Lantana produces umbellate inflorescence with younger flowers in the centre and older ones in the periphery. The colour of the flowers may change post-pollination, thus a single inflorescence can have flowers of different colours (Negi, Sharma, & Vishvakarma, 2019). Yellow-pink (yellow young flowers in the centre of the inflorescence and older pink flowers in the periphery), White-pink and Orange (flowers maintain more or less the same colour) flower colour types are commonly observed in India (Fig. 1b). Flower colours like reddish-pink, yellow-orange and pink (Fig. 1b) are relatively uncommon. A single location can have multiple of these flower colour variants growing together next to each other.

Figure 1 Sampling locations (a) of Lantana across different biogeographic regions in India Flower colour variation (b) of Lantana. i- White-pink, ii- Yellow-pink, iii- orange, iv- Yellow-pink-b, v - Yellow-pin-c, vi - Reddish-pink, vii- pink, viii- yellow-orange, ix - yellow-orange-b, x – Yellow-dark-pink a, xi - Yellow-dark-pink b

2.2 Sampling locations

Lantana plant samples were collected between the years 2020 and 2023. Plants were collected from 36 locations across the country, representing various bioclimatic regions (Fig. 1a, Supplementary Table 1). Wild-growing plant saplings were collected and planted in the greenhouse at the National Center for Biological Sciences, Bengaluru. From each location, a few plants that represented different flower colour types were selected randomly for the genomic analysis. We received silica gel-dried leaf samples from four locations which were also included in the study. Whenever possible, we have noted the flower colour during collection, or in the

greenhouse—however, the exact flower colour of 59 plants that died before flowering

is unknown.

2.3 DNA extraction and sequencing library preparation

Two to three fresh leaves were collected from each plant for DNA extraction.

Leaves were crushed using liquid nitrogen in a mortar and pestle. A Qiagen DNeasy

plant pro kit was used for the extractions, following the manufacturer's instructions.

DNA was quantified using a Qubit fluorometer. 199 µl Qubit buffer and 1 µl dye were

mixed and 199 µl of this mix was used with 1µl DNA to check the concentrations.

Extracted DNA was stored at -20 degrees Celsius. Double-digest Restriction-site-

associated DNA (ddRAD) libraries were prepared using the modified protocol by

Tyagi et al 2024. Briefly, genomic DNA was digested using two restriction enzymes;

SphI a six-base pair and MlucI a four-base pair cutter. Adapters specifically designed

for SphI and MlucI are ligated to the cut ends. Unbound adapters were removed

using magnetic bead-based cleaning, where 0.5X beads were used to remove

fragments shorter than 50 base pairs. Individual specific index combinations were

ligated to each sample using PCR and amplification was carried out. The PCR

products were dual size selected for a target library size of 200 to 500 bp using

magnetic beads. Library quality was evaluated using a TapeStation automated

electrophoresis system. All the samples were pooled to get a 2nM library and the

final library was sequenced in HiSeq2500 and NovaSeq6000 platforms.

2.4 Variant calling and filtering

Raw reads were demultiplexed and barcodes were trimmed using a custom

script. Raw reads were quality-controlled using Trimmonatic - v.0.33.0 (Bolger et al.,

2014). Reads with an average quality lower than 30 and reads shorter than 30 bp

were excluded. Trimmed reads were aligned to a draft reference genome of *L.*

camara (Joshi et al, 2022) using BWA-MEM (Li & Durbin, 2009) with default settings.

We used Freebayes (Garrison & Marth, 2012) to call variants using the default

settings, which were filtered using VCFtools (Danecek et al., 2011) to retain only

high-quality SNPs. The filtering was done in the following order: biallelic SNPs were

retained, SNPs with a minimum depth of 10 and a maximum depth of 500 were

retained. Sites with a minimum genotyping quality of 30 and a minor allele count of at

least three were kept. Individuals with more than 90% missing data were removed.

SNPs that are not in Hardy-Weinberg equilibrium were removed. Finally, SNPs

present in at least 90 per cent of the individuals were retained. The filtered VCF was

used for all further analyses.

2.5 Population structure and genetic diversity estimations

We used a Bayesian clustering approach (ADMIXTURE) and a multivariate

approach (Principal Component Analysis) to understand the genetic structuring of

Lantana. Principal Component Analysis (PCA) was carried out using PLINK (Purcell

et al., 2007). Genetic population structuring was investigated using the model-based

clustering algorithm ADMIXTURE (Alexander et al., 2009), with K values ranging

from 2 to 17. We used Evanno's delta K method (Evanno et al., 2005) implemented

in the web server CLUMPAK (Kopelman et al., 2015) to find the optimum K.

CLUMPAK was used to plot the admixture proportions as well. The structure plot was

rearranged for easy visualization. Inbreeding coefficients and heterozygosity were

estimated using the R package Adegenet (v2.1.10) (Jombart, 2008). F_{ST} was

estimated using Weir and Cockerham's method implemented in VCFtools (Danecek

et al., 2011). Adegenet (v2.1.10) was used to test for isolation by distance. Nei's

genetic distance was calculated using the StAMPP package (Pembleton et al., 2013)

in R. A phylogenetic network was generated using the r package StAMPP and the

SplitsTree4 (Huson, 1997).

2.6 Genetic simulations

We used SLiM (Haller & Messer, 2019), a forward-in-time, individual-based

simulation program to understand the importance of mating systems on the pattern of

genetic diversity in invasive plant species. Populations analogous to both native and

invasive populations were simulated with different mating systems. All the individuals

in the simulations were diploid hermaphrodites, with 10 chromosomes each with

1,00,000 bps. The mutation rate and the recombination rate were assumed to be 10^{-8}

and 10^{-7} respectively (Igor Kovalchuk, 2000), which is the general mutation and

recombination rate for plants. Generations were assumed to be non-overlapping.

Scenarios with different mating systems having varying levels of self-fertilization (zero

- complete crossing, 25, 50, 75 -mixed-mating and 100- complete selfing) were

simulated. A schematic representation of the demographic changes during the

simulation is shown in Figure 2. The simulation started with a single population of 100

genetically identical individuals, which grew exponentially (growth rate=1.05) until

they reached a carrying capacity of 10,000 individuals. This carrying capacity was

decided, to reduce the computational complexity. In the burn-in phase, the individuals

evolve under these conditions for 20,000 years/generations. In the 20,000[th] year, two

new populations were created by individuals migrating from the initial simulated

population (assumed to be the native population). The number of founding individuals

(in this case propagule pressure) was either 10, 100 or 1000. These populations

grew exponentially to a carrying capacity of 10,000 with the same growth rate. The

population evolved till the year 20,200 and thus 200 generations after invasion is

simulated. The population level heterozygosity, nucleotide diversity and F_{ST} between

populations were measured at the end of the 20th and 200th generations. For

randomly selected 50 individuals from each population, VCF files with SNP

information were created at the 20,200th generation.

Figure 2 Schematic of the SLiM simulations. In generation one, a single population is
formed with 100 individuals. It exponentially grows and reaches a carrying capacity of
10,000. At generation 20,000 two new invasive populations are formed, founded by
10 individuals each. Further, all three populations evolve for 200 more years.

3 Results

3.1 Invasive Lantana populations in India show strong genetic structure

Using ddRAD sequencing, we obtained 19,008 SNP markers for 359 Lantana

individuals from different regions across India. This SNP dataset was utilised to

assess the genetic differentiation in Lantana through three different analyses:

Structure, Principal component analyses and fixation index. Structure analyses using

the ADMIXTURE package revealed strong genetic structure in the Indian Lantana

populations (Fig. 3a). The optimum number of Hardy-Weinberg populations was

between four and eleven (Supplementary Figure 1). At K=4, which had the highest

support for the optimum number of clusters, the geographic origin did not dictate the

genetic structuring. Despite belonging to the same geographic location, individuals

were assigned across different genetic clusters. At K=3, nine individuals from North-

Eastern India clustered together and maintained a distinctive identity at higher K as

well. Many individuals were admixed. K=8 and 11 had the second and third highest

log likelihood values for the optimum K.

The principal component analysis results corroborated the findings from the

admixture analyses (Fig. 3b and c). In the PCA encompassing all 359 samples

(Fig.3b), PC 1 explained 44.8% of the variation and PC 2 explained 6.4% of the total

variation. Nine samples from North-eastern India formed a distinct cluster separate

from the rest of the samples in the PC1 axis (Fig. 3b). To explore the structuring

pattern within the remaining samples, a separate analysis excluding these nine

individuals was carried out (Fig. 3c). In this analysis, PC 1 explained 16.2% of the

variation and PC 2 explained 11% of the variation. In the PCA, samples were

coloured based on the geographic location from south to north of India (brown to

green respectively). Individuals from geographically distant locations were clustering

together while those from the same geographic area were assigned to different

clusters in many locations, reinforcing the absence of a strong geography-based

pattern. A flower colour-based clustering pattern was evident in the PCA with

samples colour-coded according to the flower colour (Supplementary Figure 2).

The isolation by distance analysis using the Mantel test (Fig. 3d) revealed a weak

correlation between genetic distance and geographic distance (Mantel r = 0.039, p-

value = 0.075), indicating that geography is not the major factor influencing the

structuring pattern. The fixation index (F_{ST}) between most of the population was low

(Supplementary Table 2). Arunachal Pradesh (in north-eastern India) samples

showed high F_{ST} with many populations (0.46 to 0.54) (Supplementary Table 2).

Figure 3 Population structure of Lantana in India. a) Genetic ancestry assignment
based on the programme ADMIXTURE. Ancestry fractions assuming four, eight and
eleven ancestral populations are given. Clusters are arranged based on the flower
colour variants b) PCA of all 359 samples c) PCA of all samples except nine
northeast Indian samples. Individuals are coloured based on their geographic
locations from north to south of India d) Isolation by distance showing a weak
correlation between genetic distance and geographic distance

3.2 Genetic population structuring in Lantana is associated with flower colour

In the Admixture analysis, the clusters closely aligned with distinct flower colour

variants of Lantana. At K=4, yellow-pink and some of the orange-flowered Lantana

clustered together, while a separate cluster comprised pink and some of the white-

pink individuals (Fig. 3a) . A third cluster was formed by dark-pink-coloured plants

from Ooty (Southern India) and North-east India. Nine individuals from northeast

India formed a fourth cluster. Other flower colour variants were admixed.

Interestingly, at K=11, a pronounced flower colour-based clustering pattern emerged,

with distinct clusters formed by yellow-pink, pink, reddish-pink, dark pink and yellow-

orange individuals, while both orange and white-pink Lantana each formed two

separate clusters. In the PCA, the clustering pattern often mirrored the flower colour

variants (Supplementary Figure 2). Yellow-pink colour variant formed a tight cluster.

One of the orange colour variants also showed a similar trend. In the phylogenetic

network generated using SplitsTree, (Fig. 4a) most of the individuals were grouped

according to their flower colour. The most common flower colour in the country,

yellow-pink, formed a single cluster in admixture analysis and in the SplitsTree

network. Most of the yellow-pink and orange individuals across the country had short

branch lengths in the phylogenetic network.

High Inbreeding and low genetic diversity characterize Lantana populations

Nearly all the individuals exhibited a very low proportion of heterozygous sites, with

less than 10% heterozygous sites (Fig. 4c). However, the nine individuals from North-

East India showed the highest proportion of heterozygous sites (>0.3). The

inbreeding coefficient of many of the individuals was extremely high, suggesting high

levels of self-fertilization (Fig. 4d). The Nei's genetic distance between most of the

Lantana individuals across the country was very low indicating the absence of

genetically distinct lineages (Fig. 4b). Even with low genetic distance, there is often

strong differentiation within a location. The genetic distance was high only for the

nine individuals from the northeast of India with the rest of the samples. This was

reflected in the phylogenetic network, where the nine northeast individuals formed a

distinct group with long branch lengths (Fig. 4a).

Figure 4 Signatures of high inbreeding due to self-fertilization in Lantana a)
Phylogenetic network using SplitsTree4, lineages are coloured based on the flower
colour b) Nei's genetic distance between all the individuals, nine northeast India
samples are genetically most distinct c) Proportion of heterozygous sites per
individual d) Inbreeding coefficient

3.3 Genetic diversity in invasive populations is influenced by the mating system.

To understand the influence of the mating system on genetic diversity in invasive

plant species, we have carried out an individual-based genetic simulation using SLiM.

These simulations confirmed that the mating system strongly impacts standing

genetic variation within a species. Simulations of cross-fertilisation resulted in higher

heterozygosity and nucleotide diversity than those of self-fertilization (Fig. 5 a and b).

This was consistent across various propagule pressures (10, 100 and 1000)

simulated. The heterozygosity and nucleotide diversity were low in the simulated

invasive populations compared to the simulated native population (Fig. 5 a and b). An

increase in the propagule pressure did not substantially increase the genetic

diversity. Under the same propagule pressure, simulations of cross-fertilization gave

rise to invasive populations with high genetic diversity compared to scenarios with

self-fertilization. The variance in heterozygosity and nucleotide diversity was high in

the self-fertilizing simulation scenarios.

Mating systems had an impact on the differentiation of simulated native and invasive

populations. F_{ST} between simulated native and invasive populations revealed that the

F_{ST} value was lowest under complete cross-fertilization (Fig. 5c). The differentiation

increased as the percentage of self-fertilization increased. Under simulations of

complete self-fertilization, the variance was high and F_{ST} ranged from approximately

0.03 to 0.69. Whereas, in complete cross-fertilization, it ranged from 0.07 to 0.2.

Similarly, in simulations of cross-fertilization, the differentiation between simulated

invasive populations was low compared to self-fertilisation scenarios (Fig. 5d). The

differentiation increased as the allowed selfing percentage increased. F_{ST} ranged

from 0.05 to 0.98 in self-fertilizing simulation scenarios. Within each category of

selfing percentage, an increase in the propagule pressure reduced the differentiation.

Figure 5 Individual-based genetic simulation results showing putative patterns of
genetic diversity in invasive plants under different assumed mating systems and
propagule pressure. Heterozygosity (a) and nucleotide diversity (b) under different
mating systems in 'native' and 'invasive' populations founded by 10, 100 and 1000
individuals c) F_{ST} between 'native' and 'invasive populations' under different
simulated mating systems and propagule pressure. d) F_{ST} between two simulated
'invasive populations' under different mating systems and propagule pressure.

Self-fertilization leads to strong structuring within the simulated populations.

PCA using SNPs from individuals of all three populations revealed a strong genetic

structure in self-fertilization scenarios (Supplementary Figure 3). Whereas, in cross-

fertilization scenarios, there was a weak structure (Supplementary Figure 3). Under

selfing, individuals from the same population were assigned to distinct genetic

clusters, and within a cluster, individuals from multiple populations were present.

4 Discussion

Biological invasions pose a serious global challenge, with millions of dollars spent

annually on their control (IPBES, 2023). Yet, it is often difficult to control the

emergence of new invasive species and their spread due to the lack of

understanding of the invasion process and the factors contributing to invasion

success. Critical to understanding invasion success is an understanding of standing

genetic variation and population differentiation. We used genomic techniques to

understand patterns of genetic diversity and differentiation in invasive Lantana

populations in India. Our findings revealed that Lantana populations in India are

genetically structured, but the structuring pattern was not strongly correlated with the

geography. In the Admixture analysis, the genetic structure aligned with flower colour

variation. Notably, many individuals exhibited a high inbreeding coefficient and a very

low proportion of heterozygous sites. Genetic distance and phylogenetic network

indicated that lineages are relatively similar. We suggest that the strong genetic

structure despite the low genetic distance is because of the increased homozygosity

due to self-fertilization. Through genetic simulations, we show that the mating system

can impact genetic diversity and the structure of simulated, putatively invasive

species.

4.1 Population structure

Our investigation into the genetic structure of Lantana using both Admixture and

principal component analysis revealed a strong population structure. Individuals from

the same geographic location were assigned to different genetic clusters, contrary to

a previous study using microsatellite markers which suggested genetic homogeneity

in Lantana populations in India (Ray & Quader, 2014). SNP markers are numerous in

the genome, and as a result, have higher statistical power than microsatellite markers

(Zimmerman et al., 2020). We have used 19,008 SNP markers and thus we expected

population structure to be better resolved.

In PCA, admixture, and phylogenetic network nine individuals from the northeast of

India were genetically distinct compared to the rest of the samples. These samples

exhibited high heterozygosity compared to other plants, which could explain their

genetic distinctiveness. These plants also had noticeable phenotypic differences,

such as a lack of thorns and dense trichomes on leaves. Further detailed analyses of

these plants are required to better understand the basis of such a large genetic

difference.

Many invasive species show strong genetic structure in the invaded ranges (Wang &

Chen, 2017) due to multiple introductions, followed by the lack of gene flow between

populations (Bastien Lavergne & Molofsky, 2007). However, in many such studies, a

geography-based population structuring pattern is observed (Le Cam et al., 2020).

Our genetic results corroborate the expectation of multiple introductions as

suggested by the historical records on Lantana (Kannan et al., 2013). As expected

with multiple introductions, we find multiple genetic clusters (across populations) and

phenotypic (varied flower colours) diversity in invasive populations of Lantana in

India. However, our findings are unique because, despite having multiple genetic

clusters, we do not find geographic structuring.

4.2 Self-pollination leading to high homozygosity in Lantana

Most of the Lantana individuals in India show a high inbreeding coefficient and a low proportion of heterozygous sites. In plants, high inbreeding coefficient and low heterozygosity are associated with self-fertilization (Hedrick 2011). These results indicate that Lantana is predominantly self-fertilizing in India and exists as homozygous inbred lines. This is further supported by Structure and PCA results. In many natural populations of plants, the type of mating system is shown to influence the patterns of genetic diversity (Bomblies et al., 2010). Mating systems in plants range from uniparental reproduction or self-fertilization to obligate cross-fertilization or outcrossing (Charlesworth, 2006). Approximately 10-15% of seed plants predominantly self-fertilize (Goodwillie et al., 2005). Many species exhibit mixed mating systems, including a combination of both selfing and outcrossing (Whitehead et al., 2018). Mating systems influence the effective population size and thus play a crucial role in shaping genetic diversity within a species (Laenen et al., 2018).

In its native range, Lantana primarily relies on cross-fertilization for reproduction (Barrows, 1976). However, in the introduced range, Lantana is self-compatible (Goulson & Derwent, 2004; Ram & Mathur, 1984). During colonization and invasion, plant species can shift mating systems from cross-fertilization to self-fertilization (Petanidou et al., 2012; Willi et al., 2022). A detailed study of mating systems in native and invaded ranges is required to confirm such a shift in the case of Lantana. Self-fertilization is shown to increase the genome-wide homozygosity and inbreeding coefficient (Barragan et al., 2024). The strong genetic structure even with low genetic differences can be attributed to the high homozygosity of the inbred lines formed

through selfing (Bomblies et al., 2010). The presence of multiple genetic groups in a

single location is mainly due to the presence of such inbred lines. This is supported

by studies like Bhattacharya et al., 2022. Selfing acts as a barrier to gene flow

between these inbred lines, thereby maintaining their genetic identity. However, the

genetic distance between these lines suggests that they are not genetically very

different. The presence of admixed individuals in many populations indicates the

occurrence of cross-fertilization as well. These crossing events can help in

generating new recombinant inbred lines. Future research including field-based

studies on mating systems in Lantana can help better understand the extent and

frequency of these various mating strategies.

4.3 Association of flower colour with genetic structure

Our study revealed an association between genetic clustering and flower colour

variation. Different flower colour variants were assigned to distinct genetic clusters

despite being from the same geographic location. This indicates the lack of gene flow

between flower colour variants which grow physically close to each other in many

locations. This correlation between flower colour and genetic structuring underscores

the influence of the mating system. Here, self-fertilization acts as a barrier to gene

flow and different inbred lines are associated with different flower colours. Thus, we

think that the identity of these lines is maintained by self-fertilization. Historically,

Lantana species were primarily defined mainly based on the flower colours and many

such variants were considered as different species (Sanders, 2006). Our study

suggests that these are within species variation and the overall phenotypic similarity

within a flower colour variant is maintained by self-fertilization. We think that the

correlation between flower colour and genetic structure is caused by the formation of

inbred lines. Further genetic analysis is required to understand the genetic basis of

these flower colour variations.

4.4 Is Lantana a species complex?

The findings from the molecular analysis of Lantana challenge the notion of invasive

Lantana being a species complex. While genetic structuring was evident, it

predominantly arose from the prevalence of highly homozygous inbred lines rather

than from the presence of genetically different species. Most lines exhibited low

genetic distance between each other, indicating the high genetic similarity between

flower colour variants. The high phenotypic variation observed could be the result of

the introduction of multiple inbred lines to the subcontinent or the emergence of novel

inbred lines through hybridization post-introduction. These variations could be within

species phenotypic variations typical to many plant species. Through citizen science

platforms like iNaturalist, it is evident that Lantana in its native range also exhibits

similar flower colour variation. However, genomic analysis of related species of

Lantana can help in understanding the chances of any past introgressions.

5 **Conclusions**

Studying genetic diversity patterns in invasive species is crucial for understanding

their evolution. We explored genetic diversity patterns of the global invasive species

L. camara. Our results reveal that lantana populations in India are strongly structured

with genetic differences between flower colour variants. The low genetic distance

observed between most plants negates the chances of lantana being a species

complex. The presence of strong genetic structure even in the absence of high

genetic distance is attributed to self-fertilization and the formation of putative inbred

lines. This is supported by the low heterozygosity and high inbreeding coefficient. In

predominantly self-fertilizing species, individuals exist as inbred lines, which restricts

gene flow and allows these inbred lines to maintain their genetic identity. We suggest

that the correlation between flower colour and genetic structure is due to the

association of specific flower colours to specific inbred lines.

Uniquely, our study highlights the significance of the mating system in shaping the

genetic diversity pattern in invasive plant species. Our results also highlight the

importance of multiple introductions and the introduction of genetically different

variants in contributing to high genetic structuring within invasive populations. Further

research on other invasive species will reveal how general these patterns might be.

6   Acknowledgements

We are thankful to Keval Palya, Mamta M., and Rajat Rastogi for their help with

sampling and laboratory experiments. We are thankful to Divyashree Rana, Kritagnya

Vadar, Nishma Dahal, Rachana Rao, Tikily Tayeng, Tista Ghosh and Vinay Sagar for

aiding in the sample collection. We thank Mayuresh Gangal, Divyashree Rana,

Abhinav Tyagi, Anubhab Khan, Vinay Sagar and Arpitha Jayanth for their valuable

suggestions. We thank the forest departments of Karnataka (No: PCCF (WL) /E2/CR-

52/2019-20), Tamil Nadu (No: WL(A)/52852/2019) and Madhya Pradesh (MP permit

no:835, dated: 30-01-2020) for providing necessary permits. PP was supported by

NCBS/TIFR (Department of Atomic Energy). This work was supported under DBT

project No-BT/PR29251/FCB/125/18/2018. The NCBS data cluster used is supported

under project 12- R&D-TFR-5.04-0900, Department of Atomic Energy, Government

of India.

7 **Author contributions**

Conceptualization: P.P., U. R, and R. G., Data collection: P.P., Laboratory work: P.

P., Data analysis: P.P, Project administration: P.P., U.R., and R. G., Funding

acquisition: U. R, and R. G., Writing original draft: P.P., Writing – Review and editing:

U.R., Supervision: U.R.

8 **Data availability statement**

Accession numbers for the raw ddRAD sequencing data and the GitHub link for

analysis codes will be provided upon acceptance of the manuscript.

9 **References**

Adwait G. Joshi, P. P. U. R. R. S. (2022). Draft genome sequence of an invasive
plant Lantana camara L. Bioinformation, 18(9), 739–741.
https://doi.org/10.6026/97320630018739

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of
ancestry in unrelated individuals. Genome Research, 19(9), 1655–1664.
https://doi.org/10.1101/gr.094052.109

Alves, J. M., Carneiro, M., Day, J. P., Welch, J. J., Duckworth, J. A., Cox, T. E.,
Letnic, M., Strive, T., Ferrand, N., & Jiggins, F. M. (2022). A single introduction of
wild rabbits triggered the biological invasion of Australia.
https://doi.org/10.1073/pnas

Barragan, A. C., Collenberg, M., Schwab, R., Kersten, S., Kerstens, M. H. L.,
Požárová, D., Bezrukov, I., Bemm, F., Kolár, F., & Weigel, D. (2024). Deleterious
phenotypes in wild Arabidopsis arenosa populations are common and linked to
runs of homozygosity. G3: Genes, Genomes, Genetics, 14(3).
https://doi.org/10.1093/g3journal/jkad290

Barrows, E. M. (1976). Nectar Robbing and Pollination of Lantana camara
(Verbenaceae) (Vol. 8, Issue 2).

Bastien Lavergne, S., & Molofsky, J. (2007). Increased genetic variation and
evolutionary potential drive the success of an invasive grass.
www.pnas.org/cgi/content/full/
Bhattacharya, S., Hernández, F., Alves, M. F., Machado, R. M., Sun, Y. Y., Wang, M.
R., Zhang, C. Bin, & Hao, J. H. (2022). Genetic diversity and population structure
of invasive and native populations of Erigeron canadensis L. Journal of Plant
Ecology, 15(4), 864–876. https://doi.org/10.1093/jpe/rtac016
Blair, A. C., & Hufbauer, R. A. (2010). Hybridization and invasion: One of North
America's most devastating invasive plants shows evidence for a history of
interspecific hybridization. Evolutionary Applications, 3(1), 40–51.
https://doi.org/10.1111/j.1752-4571.2009.00097.x
Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for
Illumina sequence data. Bioinformatics, 30(15), 2114–2120.
https://doi.org/10.1093/bioinformatics/btu170
Bomblies, K., Yant, L., Laitinen, R. A., Kim, S. T., Hollister, J. D., Warthmann, N.,
Fitz, J., & Weigel, D. (2010). Local-scale patterns of genetic variability,
outcrossing, and spatial structure in natural stands of Arabidopsis thaliana. PLoS
Genetics, 6(3). https://doi.org/10.1371/journal.pgen.1000890
Charlesworth, D. (2006). Evolution of Plant Breeding Systems. In Current Biology
(Vol. 16, Issue 17). https://doi.org/10.1016/j.cub.2006.07.068
Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A.,
Handsaker, R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., & Durbin,
R. (2011). The variant call format and VCFtools. Bioinformatics, 27(15), 2156–
2158. https://doi.org/10.1093/bioinformatics/btr330
Dlugosch, K. M., & Parker, I. M. (2008). Founding events in species invasions:
Genetic variation, adaptive evolution, and the role of multiple introductions. In
Molecular Ecology (Vol. 17, Issue 1, pp. 431–449).
https://doi.org/10.1111/j.1365-294X.2007.03538.x
Edwards, C. E., Albrecht, M. A., Bassüner, B., & Yatskievych, G. A. (2019).
Population genetic analysis reveals a predominantly selfing mating system and
strong genetic structuring in a naturally fragmented, threatened plant.
Conservation Genetics, 20(6), 1437–1448. https://doi.org/10.1007/s10592-019-
01226-9
Estoup, A., Ravign, V., Hufbauer, R., Vitalis, R., Gautier, M., & Facon, B. (2016). Is
There a Genetic Paradox of Biological Invasion ?
https://doi.org/10.1146/annurev-ecolsys-121415
Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of
individuals using the software STRUCTURE: A simulation study. Molecular
Ecology, 14(8), 2611–2620. https://doi.org/10.1111/j.1365-294X.2005.02553.x
Foxe, J. P., Slotte, T., Stahl, E. A., Neuffer, B., Hurka, H., & Wright, S. I. (2009).
Recent speciation associated with the evolution of selfing in Capsella. In PNAS
(Vol. 106).
Garrison, E., & Marth, G. (2012). Haplotype-based variant detection from short-read
sequencing. http://arxiv.org/abs/1207.3907

Goodwillie, C., Kalisz, S., & Eckert, C. G. (2005). The evolutionary enigma of mixed mating systems in plants: Occurrence, theoretical explanations, and empirical evidence. In Annual Review of Ecology, Evolution, and Systematics (Vol. 36, pp. 47–79). https://doi.org/10.1146/annurev.ecolsys.36.091704.175539

Goulson, D., & Derwent, L. C. (2004). Synergistic interactions between an exotic honeybee and an exotic weed: Pollination of Lantana camara in Australia. Weed Research, 44(3), 195–202. https://doi.org/10.1111/j.1365-3180.2004.00391.x

Goyal, N., & Sharma, G. P. (2015a). Lantana camara L. (sensu lato): An enigmatic complex. NeoBiota, 25, 15–26. https://doi.org/10.3897/neobiota.25.8205

Goyal, N., & Sharma, G. P. (2015b). NeoBiota Lantana camara L . (sensu lato): an enigmatic complex. April. https://doi.org/10.3897/neobiota.25.8205

Goyal, N., & Sharma, G. P. (2015c). NeoBiota Lantana camara L . (sensu lato): an enigmatic complex. April. https://doi.org/10.3897/neobiota.25.8205

Guo, Y.-L., Bechsgaard, J. S., Slotte, T., Neuffer, B., Lascoux, M., Weigel, D., & Schierup, M. H. (2008). Recent speciation of Capsella rubella from Capsella grandiflora, associated with loss of self-incompatibility and an extreme bottleneck. www.pnas.orgcgidoi10.1073pnas.0808012106

Haller, B. C., & Messer, P. W. (2019). SLiM 3: Forward Genetic Simulations Beyond the Wright-Fisher Model. Molecular Biology and Evolution, 36(3), 632–637. https://doi.org/10.1093/molbev/msy228

Hamrick, J. L., Godt, J. W., & Sherman-Broyles, S. L. (1992). Factors influencing levels of genetic diversity in woody plant species. In New Forests (Vol. 6).

Huson, D. H. (1997). SplitsTree: analyzing and visualizing evolutionary data.

Igor Kovalchuk, O. K. and B. H. (2000). mutation.

IPBES. (2023). The thematic assessment report on INVASIVE ALIEN SPECIES AND THEIR CONTROL. https://doi.org/10.5281/zenodo.7430682

Jaspers et al. (2021). contributed new reagents/analytic tools. 118. https://doi.org/10.1073/pnas.2116211118/-/DCSupplemental

Jombart, T. (2008). Adegenet: A R package for the multivariate analysis of genetic markers. Bioinformatics, 24(11), 1403–1405. https://doi.org/10.1093/bioinformatics/btn129

Kannan, R., Shackleton, C. M., & Shaanker, R. U. (2013). Reconstructing the history of introduction and spread of the invasive species , Lantana , at three spatial scales in India. 1287–1302. https://doi.org/10.1007/s10530-012-0365-z

Koelling, V. A., Hamrick, J. L., & Mauricio, R. (2011). Genetic diversity and structure in two species of Leavenworthia with self-incompatible and self-compatible populations. Heredity, 106(2), 310–318. https://doi.org/10.1038/hdy.2010.59

Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., & Mayrose, I. (2015). Clumpak: A program for identifying clustering modes and packaging population structure inferences across K. Molecular Ecology Resources, 15(5), 1179–1191. https://doi.org/10.1111/1755-0998.12387

Laenen, B., Tedder, A., Nowak, M. D., Toräng, P., Wunder, J., Wötzel, S., Steige, K. A., Kourmpetis, Y., Odong, T., Drouzas, A. D., Bink, M. C. A. M., Ågren, J., Coupland, G., & Slotte, T. (2018). Demography and mating system shape the

genome-wide impact of purifying selection in Arabis alpina. Proceedings of the
National Academy of Sciences of the United States of America, 115(4), 816–
821. https://doi.org/10.1073/pnas.1707492115

Le Cam, S., Daguin-Thiébaut, C., Bouchemousse, S., Engelen, A. H., Mieszkowska,
N., & Viard, F. (2020). A genome-wide investigation of the worldwide invader
Sargassum muticum shows high success albeit (almost) no genetic diversity.
Evolutionary Applications, 13(3), 500–514. https://doi.org/10.1111/eva.12837

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-
Wheeler transform. Bioinformatics, 25(14), 1754–1760.
https://doi.org/10.1093/bioinformatics/btp324

Linders, T. E. W., Schaffner, U., Eschen, R., Abebe, A., Choge, S. K., Nigatu, L.,
Mbaabu, P. R., Shiferaw, H., & Allan, E. (2019). Direct and indirect effects of
invasive species: Biodiversity loss is a major mechanism by which an invasive
tree affects ecosystem functioning. Journal of Ecology, 107(6), 2660–2672.
https://doi.org/10.1111/1365-2745.13268

Lu-Irving, P., Bedoya, A. M., Salimena, F. R. G., dos Santos Silva, T. R., Viccini, L.
F., Bitencourt, C., Thode, V. A., Cardoso, P. H., O'Leary, N., & Olmstead, R. G.
(2021). Phylogeny of Lantana, Lippia, and related genera (Lantaneae:
Verbenaceae). American Journal of Botany, 108(8), 1354–1373.
https://doi.org/10.1002/ajb2.1708

Mairal, M., García-Verdugo, C., Le Roux, J. J., Chau, J. H., van Vuuren, B. J., Hui,
C., Münzbergová, Z., Chown, S. L., & Shaw, J. D. (2023). Multiple introductions,
polyploidy and mixed reproductive strategies are linked to genetic diversity and
structure in the most widespread invasive plant across Southern Ocean
archipelagos. Molecular Ecology, 32(4), 756–771.
https://doi.org/10.1111/mec.16809

Mesgarana, M. B., Lewisb, M. A., Adesc, P. K., Donohued, K., Ohadia, S., Lia, C., &
Cousensa, R. D. (2016). Hybridization can facilitate species invasions, even
without enhancing local adaptation. Proceedings of the National Academy of
Sciences of the United States of America, 113(36), 10210–10214.
https://doi.org/10.1073/pnas.1605626113

Mollot, G., Pantel, J. H., & Romanuk, T. N. (2017). The Effects of Invasive Species
on the Decline in Species Richness: A Global Meta-Analysis. In Advances in
Ecological Research (Vol. 56, pp. 61–83). Academic Press Inc.
https://doi.org/10.1016/bs.aecr.2016.10.002

Mungi, N. A., Qureshi, Q., & Jhala, Y. V. (2020). Expanding niche and degrading
forests: Key to the successful global invasion of Lantana camara (sensu lato).
Global Ecology and Conservation, 23.
https://doi.org/10.1016/j.gecco.2020.e01080

Negi, G. C. S., Sharma, S., & Vishvakarma, S. C. R. (2019). Ecology and Use of
Lantana camara in India. 109–130.

Negi, Sharma, S., Vishvakarma, S. C. R., & Samant, S. S. (2019). Ecology and Use
of Lantana camara in India. Botanical Review, 85(2), 109–130.
https://doi.org/10.1007/s12229-019-09209-8

Ness, R. W., Wright, S. I., & Barrett, S. C. H. (2010). Mating-system variation, demographic history and patterns of nucleotide diversity in the tristylous plant Eichhornia paniculata. Genetics, 184(2), 381–392. https://doi.org/10.1534/genetics.109.110130

Olazcuaga, L., Lincke, B., DeLacey, S., Durkee, L. F., Melbourne, B. A., & Hufbauer, R. A. (2023). Population demographic history and evolutionary rescue: Influence of a bottleneck event. Evolutionary Applications, 16(8), 1483–1495. https://doi.org/10.1111/eva.13581

Ørsted, M., Hoffmann, A. A., Sverrisdóttir, E., Nielsen, K. L., & Kristensen, T. N. (2019). Genomic variation predicts adaptive evolutionary responses better than population bottleneck history. PLoS Genetics, 15(6). https://doi.org/10.1371/journal.pgen.1008205

Pembleton, L. W., Cogan, N. O. I., & Forster, J. W. (2013). StAMPP: An R package for calculation of genetic differentiation and structure of mixed-ploidy level populations. Molecular Ecology Resources, 13(5), 946–952. https://doi.org/10.1111/1755-0998.12129

Pérez De La Vega, M., & Garda, P. (1997). Genetic structure of self-pollinating species: the case of wild A vena.

Petanidou, T., Godfree, R. C., Song, D. S., Kantsa, A., Dupont, Y. L., & Waser, N. M. (2012). Self-compatibility and plant invasiveness: Comparing species in native and invasive ranges. Perspectives in Plant Ecology, Evolution and Systematics, 14(1), 3–12. https://doi.org/10.1016/j.ppees.2011.08.003

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J., Sklar, P., De Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: A tool set for whole-genome association and population-based linkage analyses. American Journal of Human Genetics, 81(3), 559–575. https://doi.org/10.1086/519795

R, F. M., Vardhan, V., Madan, R., Kannan, S., Arunkumar, B., & Prakash, G. (2016). Mapping the Distribution of Lantana Camara in the Bandipur Tiger Reserve , Karnataka Draft Report submitted to Karnataka Forest Department March 2016. March.

Ram, H. Y. M., & Mathur, G. (1984). Flower Colour Changes in Lantana camara. 35(160), 1656–1662.

Ray, A., & Quader, S. (2014). Genetic diversity and population structure of Lantana camara in India indicates multiple introductions and gene flow. 16, 651–658. https://doi.org/10.1111/plb.12087

Roman, J., & Darling, J. A. (2007). Paradox lost: genetic diversity and the success of aquatic invasions. In Trends in Ecology and Evolution (Vol. 22, Issue 9, pp. 454–464). https://doi.org/10.1016/j.tree.2007.07.002

Ryan, S. F., Lombaert, E., Espeset, A., Vila, R., Talavera, G., Dinca, V., Doellman, M. M., Renshaw, M. A., Eng, M. W., Hornett, E. A., Li, Y., Pfrender, M. E., & Shoemaker, D. W. (2019). Global invasion history of the agricultural pest butterfly Pieris rapae revealed with genomics and citizen science. Proceedings

of the National Academy of Sciences of the United States of America, 116(40), 20015–20024. https://doi.org/10.1073/pnas.1907492116

Sanders, R. W. (2006). TAXONOMY OF LANTANA SECT. LANTANA (VERBENACEAE): I. CORRECT APPLICATION OF LANTANA CAMARA AND ASSOCIATED NAMES. In SIDA (Vol. 22, Issue 1).

Schoen, D. J., & Brown, A. H. D. (1991). Intraspecific variation in population gene diversity and effective population size correlates with the mating system in plants (inbreeding species/genetic conservation/neutral model/maximnum likelihood/isozyme). In Proc. Natl. Acad. Sci. USA (Vol. 88). https://www.pnas.org

Simberloff, D. (2009). The role of propagule pressure in biological invasions. Annual Review of Ecology, Evolution, and Systematics, 40, 81–102. https://doi.org/10.1146/annurev.ecolsys.110308.120304

Urban, A. J., Simelane, D. O., Retief, E., Heystek, F., Williams, H. E., & Madire, L. G. (2011). The invasive ' Lantana camara L .' hybrid complex (Verbenaceae): a review of research into its identity and biological control in South Africa. 19(2), 315–348.

Vallejo-Marín, M., Friedman, J., Twyford, A. D., Lepais, O., Ickert-Bond, S. M., Streisfeld, M. A., Yant, L., van Kleunen, M., Rotter, M. C., & Puzey, J. R. (2021). Population genomic and historical analysis suggests a global invasion by bridgehead processes in Mimulus guttatus. Communications Biology, 4(1). https://doi.org/10.1038/s42003-021-01795-x

Wang, G., & Chen, Z. (2017). Strong population genetic structure of an invasive species , Rhynchophorus ferrugineus (Olivier), in southern China. February, 10770–10781. https://doi.org/10.1002/ece3.3599

Whitehead, M. R., Lanfear, R., Mitchell, R. J., & Karron, J. D. (2018). Plant mating systems often vary widely among populations. In Frontiers in Ecology and Evolution (Vol. 6, Issue APR). Frontiers Media S. A. https://doi.org/10.3389/fevo.2018.00038

Willi, Y., Lucek, K., Bachmann, O., & Walden, N. (2022). Recent speciation associated with range expansion and a shift to self-fertilization in North American Arabidopsis. Nature Communications, 13(1). https://doi.org/10.1038/s41467-022-35368-1

Zhang, L., Hua, Y., & Wei, S. (2021). High genetic diversity of an invasive alien species: Comparison between fur-farmed and feral american mink (neovison vison) in China. Animals, 11(2), 1–11. https://doi.org/10.3390/ani11020472

Zimmerman, S. J., Aldridge, C. L., & Oyler-Mccance, S. J. (2020). An empirical comparison of population genetic analyses using microsatellite and SNP data for a species of conservation concern. BMC Genomics, 21(1). https://doi.org/10.1186/s12864-020-06783-9