Title: An unexplored diversity for adaptation of germination to high temperatures in
***Brassica* species**

Authors:

Tiret, M.[1*], Wagner, M.-H.[2], Gay, L.[3], Chenel, E.[1], Dupont, A.[2], Falentin, C.[1], Gavory, F.[4], Labadie,
K.[5], Ducournau, S.[2], Chèvre, A.-M.[1]

Institutions:

[1] IGEPP, INRAE, Institut Agro, Université de Rennes, 35650 Le Rheu, France
[2] GEVES, Station Nationale d'Essais de Semences, 49071 Beaucouzé, France
[3] UMR AGAP Institut, Université de Montpellier, CIRAD, INRAE, Institut Agro, Montpellier,
France
[4] Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Univ Evry, Université
Paris-Saclay, Evry, France
[5] Genoscope, Institut François Jacob, CEA, Université Paris-Saclay, Evry, 91057, France

[*] Corresponding author: Mathieu Tiret (mathieu.tiret@inrae.fr)

Abstract: (186 words)
Main Text: 5,692 words.

Elevated temperatures inhibit the germination of a concerning number of crop species. One strategy
to mitigate the impact of warming temperatures is to identify and introgress adaptive genes into
elite germplasm. Diversity must be sought in wild populations, coupled with an understanding of
the complex pattern of adaptation across a broad range of landscapes. By investigating the
landraces, wild, and feral populations of Algeria, Italy, France, Slovenia, Spain, and Tunisia, we
assessed the response of germination to temperature increase in an unexplored diversity of 112
accessions of *B. rapa* and 60 of *B. oleracea*. Our results show that *B. rapa* accessions exhibited
tolerance to the temperature range tested, with an increase in speed and uniformity of germination
time, as well as an increase in germination rate for the landraces as temperature increased.
Conversely, *B. oleracea* accessions displayed a lower germination rate and speed as temperature
increased. The southern wild populations of *B. oleracea* were identified as heat tolerant, in relation
to their warmer climate of origin. These findings highlight the complex interplay between
domestication, feralization, and current agronomic practices in shaping germination characteristics
in *Brassica* species.

keywords: germination, heat tolerance, *Brassica oleracea, Brassica rapa,* landscape genomics.

Introduction

Environmental factors, including temperature, precipitation patterns, increased atmospheric CO_2 concentrations, and extreme weather events, will undergo significant deviations from their current states as a direct consequence of climate change (IPCC, 2013; IPCC, 2021). Agriculture is among the most endangered sector (Wheeler & Von Braun, 2013; Arunanondchai et al., 2018; Raza et al., 2019), across many major regions (Rosenzweig et al., 2014; Thornton et al., 2014), particularly due to warming temperatures often leading to reduced yields, altered crop quality, and increased susceptibility to environmental variations (Lobell & Field, 2007; DaMatta et al., 2010; Zhao et al., 2017; Chaudhry et al., 2022). On many major crops, the detrimental effects of warming temperatures are evident in the disruption of developmental stages and phenological cycles (Angadi et al., 2000; Barnabás et al., 2008; Chmielewski et al., 2004; Hatfield & Prueger, 2015).

Seed vigor is particularly vulnerable to the environmental changes induced by climate change (Hampton et al., 2016; Castillo-Lorenzo et al., 2019). Seed vigor encompasses seed longevity, germination speed, seedling growth, and early stress tolerance (Pollock & Roos, 1972; Reed et al., 2022). It is a key determinant of a plant's early growth and competitive ability, and is essential for ensuring successful establishment (TeKrony & Egli, 1991; Finch-Savage & Bassel, 2016; Ebone et al., 2020). The processes involved are influenced by a variety of environmental factors, either experienced during seed development (i.e., the maternal effect; Wolf & Wade, 2009; Penfield & MacGregor, 2017), storage (Taylor, 2020), or on the seedbed (Lamichhane et al., 2018). Elevated temperatures are known to impair seed fertility, delay germination, and lower seed vigor in various crop species (Gareca et al., 2012; Cochrane et al., 2015; Tribouillois et al., 2016; Reed et al., 2022).

The susceptibility of seed vigor to environmental factors has often been ascribed to evolutionary responses to specific geographical conditions (Clauss & Venable, 2000; Galloway, 2005; Franks et al., 2014; Dürr et al., 2015; Becklin et al., 2016). The adaptation to the local environment may be widespread, as plants that evolved in warmer climates often possess the ability to thrive under high-temperature conditions (Cochrane et al., 2015; El-Keblawy et al., 2018). On the other hand, the recent ongoing climate change may require a shift in phenotypes to remain locally adapted; but whether rapid genetic change is sufficient to keep pace with climate change remains a topic of debate (Jump & Peñuelas, 2005; Jump et al., 2008). The relationship between climate of origin and heat tolerance is complex and it is not always straightforward to identify a pattern of local adaptation due to phenotypic plasticity and the interplay of various environmental and genetic factors (Jump & Peñuelas, 2005; Nicotra et al., 2010). These studies underscore the intricate nature of a 'climate-smart agriculture' strategy to mitigate the impact of climate change, including, among other approaches, assisted migration (Pareek et al., 2020; Sloat et al., 2020). Indeed, the inadvertent consequence of assisted migration may be the disruption of the local adaptation patterns of crops, potentially leading to a reduction in yield or fitness.

The *Brassica* genus is notable for its extensive morphological and genetic diversity, which has made it a global economic resource, particularly in the production of edible root crops, vegetables, and oilseeds (Warwick & Black, 1991; Rakow, 2004; Cheng et al., 2016). Despite this diversity, cultivated lines exhibit limited genetic variation, particularly those of oilseed rape (*Brassica napus*

L.), which constrains their adaptive potential (Allender & King, 2010). Establishment failure has
been identified as a recurring problem for oilseed rape, and despite extensive research, a solution
remains elusive (Nelson et al., 2022). As in many plant genera, the utilization of plant genetic
resources in the *Brassica* genus has been considered a strategy to mitigate climate change (Salgotra
& Chauhan, 2023). Wild populations and crop wild relatives represent an untapped gene pool,
adapted to a wide range of habitats and potentially more resilient to climate change than cultivated
crops due to their higher genetic diversity (Castillo-Lorenzo et al., 2019; Renzi et al., 2022). They
offer valuable traits for improving stress tolerance and adaptability to changing environmental
conditions, making them prime candidates for breeding programs (Tanksley & McCouch, 1997;
Zamir, 2001; Nevo et al., 2012; McCouch et al., 2013). For instance, crop wild relatives of the
Brassica genus have been screened recently for heat and water stress resistance (Castillo-Lorenzo et
al., 2019). Nevertheless, a large-scale sampling strategy across a broad range of climatic conditions
remains to be conducted.
100
The objective of this study was to assess, with high-throughput imaging phenotyping, the
germination abilities under heat stress conditions of an unexplored diversity panel of wild
populations and landraces comprising 112 accessions of *Brassica rapa* L. and 60 accessions of
Brassica oleracea L., the two progenitors of *B. napus*, and provide insights for future targets of
breeding aiming to mitigate the effect of climate change. The results indicate that while *B. rapa*
landraces display an ability to germinate at high temperature, *B. oleracea* accessions generally do
not. Notably, wild populations of *B. oleracea* from southern regions show a significant degree of
heat tolerance, which is likely due to the warmer climate of origin, making them a promising group
of candidates for consideration in improving the adaptation of landraces to the warming climate.
110

Materials & Methods

Plant material. The material, which is a subsample of the plant collection presented in Falentin et al. (2024), comprises 117 accessions of *B. rapa* and 66 of *B. oleracea*. Among them, 55 accessions of *B. rapa* and 50 of *B. oleracea* are landraces, and 62 accessions *B. rapa* and 16 of *B. oleracea* accessions are populations sampled in the field (in areas that are not part of a cultivated plot). For simplicity, we will refer to these populations as 'wild' in the following, knowing that it may reflect different origins (true wild or feral populations; see Mittel et al., 2020; McAlvay et al., 2021; Saban et al., 2023). Seeds were collected in Algeria, France, Italy, Slovenia, Spain, and Tunisia. For wild populations, seeds were collected from a maximum of 30 mother plants per sampling site; for landraces, they were collected from local farmers. To circumvent any potential maternal effect, the collection was then multiplied over two consecutive years (2020/2021 and 2021/2022) at a single location (Le Rheu, France, 48°06'22"N 1°47'25"W). Seedlings were grown for seed production under cages covered with pollen-proof netting: two cages per population of *B. rapa* with five plants each , and three cages per population of *B. oleracea* with two plants each. The production of the multiplication is what we will refer to as the "accession", which is a heterogeneous pool representing the descent of 6 to 10 mother plants (Table 1). Hereafter, we refer to the accessions' affiliation with wild populations or cultivated landraces as their 'type'. Twenty-eight accessions of *B. rapa* and 50 of *B. oleracea* were harvested in the summer of 2021, and 104 accessions of *B. rapa* and 15 of *B. oleracea* were harvested in the summer of 2022 (15 accessions of *B. rapa* overlapped between the two seasons). Seed lots were stored at 4°C and 10 % relative humidity in the dark until phenotyping. Seed lots were then split for different analyses and measurements (germination in controlled conditions and emergence in field conditions).

Phenotyping – seed germination. We phenotyped the accessions for germination time (GT) and rate (GR) under two treatments in controlled conditions: at 20°C following the ISTA reference temperature for both species (ISTA, 2024), and at 35°C for *B. rapa* and 30°C for *B. oleracea* to measure heat tolerance. The maximum temperatures were determined based on tests on commercial germplasm, on what was perceived as a stress (germination rate drops).

Germination was monitored using the high-throughput Plant Imaging PHENOTIC platform at the National Seed Testing Station at Angers, France (GEVES, Angers, France). Measurements were performed on 16 trials of eight days on Jacobsen germinators, from 10/15/2021 to 06/06/2023. Seeds were imbibed in the dark for 168 h at a water potential of -0.4 Mpa. During the experiment, the germinators maintained a constant temperature of 20°C or 30/35 ± 5°C. The germination process was automatically assessed via the image analysis method described in Demilly et al. (2014), with images captured every two to three hours throughout the experiment. When necessary, the data collection was terminated after seven days to minimize the occurrence of false positive. At the conclusion of the experiment, if germination occurred within the first 168 hours, the time to the start of germination (in hours), namely GT, was recorded; otherwise, it was declared as non-germinated.

In each trial, a randomized incomplete block design was implemented, with blocks of 25 seeds of a single accession (one batch; see details in Wagner et al., 2011). All accessions were replicated 100

times (four batches) per treatment (low or high temperature), with the exception of four accessions of *B. rapa* replicated 95 times at 35°C, all due to material availability. Additionally, to investigate the effect of the seed multiplication year, supplementary seeds of *B. rapa* were tested at 20°C, leading to 200 seeds for 26 accessions and 125 seeds for 32 accessions. For *B. rapa*, only 79 accessions (out of 117) were tested at 35°C given that the germination rate was already low at 20°C: temperature increase would not have broken dormancy, which is the major reason of low germination at 20°C for Algerian and Italian wild accessions. For *B. oleracea*, all 66 accessions were tested at both temperatures. In total, 25,080 seeds of *B. rapa* (16,700 at 20°C and 8,380 at 35°C) and 12,725 of *B. oleracea* (6,025 at 20°C and 6,700 at 30°C) were phenotyped.

Phenotyping – seed emergence. To confirm that the germination ranking is maintained under field conditions, we conducted an experiment in a common garden setting. The emergence rate, defined as the ratio of cotyledon-stage seedlings to the total number of seeds sown, was measured in the field in 2022 on the same lots of seeds that were used in the laboratory. For the *B. rapa* accessions, the seeds were sown in a random incomplete block design on September 10, 2022 at Le Rheu (France, 48°06'33"N 1°47'11"W), in three repetitions of 120 blocks (including a control replicated three times in each repetition, the "Purple Top Milan"), with 150 seeds per plot of 1 m x 1.5 m. For the *B. oleracea* accessions, the seeds were sown in small pots within a greenhouse between September 19, 2022 and September 22, 2022, at Ploudaniel, France (48°30'06"N 4°19'32"W), with 140 seeds per accession. The emergence rate was measured on a block on October 6, 2022 for *B. rapa* and on October 3, 2022 for *B. oleracea*.

Sequence data and bioinformatics. For each accession, a pool of 30 seedlings (for landrace accessions, 30 random seeds; for wild accessions, at least one seed per mother plant) was used to generate DNA bulks with a standardized sampling per plant (see details in Maillet et al., 2023). Libraries were constructed from the DNA pools (250 ng) using the NEBNext DNA Sample Prep modules (New England Biolabs) with 'on beads' protocol. DNA libraries were sequenced using Illumina NovaSeq 6000 technology (short reads, paired-end, 150 bp; Illumina Inc., San Diego, CA) with a target coverage of 50X. The sequences of the Illumina adapters and primers used during library construction were removed from the whole reads. Low quality nucleotides with quality value < 20 were removed from both ends. The longest sequence without adapters and low quality bases was kept. Sequences between the second unknown nucleotide (N) and the end of the read were also trimmed. Reads shorter than 30 nucleotides after trimming were discarded. These trimming steps were achieved using fastx_clean (http://www.genoscope.cns.fr/fa), an internal software based on the FASTX library (http://hannonlab.cshl.edu/fast). The reads and their mates that mapped onto run quality control sequences (Enterobacteria phage PhiX174 genome) were removed using SOAP aligner (Li et al., 2008). The cleaned reads were mapped on the *B. rapa* 'C1.3' var. *rapifera* genome for *B. rapa* (which corresponds to the A subgenome of the *B. napus* 'RCC-S0'; Maillet et al., 2023) and *B. oleracea* 'HDEM' ssp. *captitata* genome for *B. oleracea* (Belser et al., 2018), using the default parameters of the mem program from the bwa v. 0.7.17 software (Li & Durbin 2009; Li, 2013). Read alignments with a mapping quality Phred score below 20 or PCR duplicates were then removed using the view (option '-q 20') and 'markdup' programs from the SAMtools v. 1.14 (Danecek et al., 2021) software suite. Then, we completed the mapping by a step of realignment around indels for *B. oleracea*, as implemented in GATK (McKenna et al., 2010). Subsequently,

198 variant calling was conducted using BCFtools v. 1.11 (Danecek et al., 2021) with the options '-q 20
199 -Q 20', which set the minimum mapping quality for an alignment and the minimum base quality for
200 a base to 20.
201
The resulting VCF file was filtered using BCFtools to retain only biallelic SNPs, and remove SNPs
situated less than 5 bp away from indels. Only accessions with an average depth higher than 20
were retained (removing five accessions of *B. rapa* and six of *B. oleracea*). Then, SNPs were
retained if they exhibited an average depth between its 5th and 95th percentiles, a mapping quality
exceeding 40, and mapped on chromosome scaffolds. SNPs were also removed when both allelic
variants had a depth below 10. The VCF file was then pruned for linkage disequilibrium, with the
prune program from the BCFtools v 1.11 suite program, with the option '-l 0.2 -w 1000'. The
variant filtering resulted in 106,064 SNPs for the 112 accessions of *B. rapa*, and 80,054 SNPs for
the 60 accessions of *B. oleracea*.
211
Population genetic analyses. The genomic relationship matrix (G) was calculated using GCTA
v1.94.1 (Yang et al., 2011) with the default parameters, and subsequently transformed into a
dissimilarity matrix (1-G*), where the correlation matrix (G*) was derived from G using the
'cov2cor' function of the R package 'stats'. The dissimilarity matrix was analyzed using a
multidimensional scaling (MDS) decomposition with the 'cmdscale_lanczos' function of the R
package 'refund' (Goldsmith et al., 2024). The subpopulation membership was calculated using the
ADMIXTURE program v. 1.22 (Alexander et al., 2009), with the default parameters and a cross-
validation (CV) error rate used to determine the likelihood of the K parameter.
220
221 **Statistical analyses**. All analyses were performed with R v 4.4.1 (R Core Team, 2024). The
222 genomic BLUP (gBLUP) of germination time (GT) for the accessions was calculated using a mixed
223 linear model (one for each species) according to the following model:
224
225 $GT_{ijkl} = \mu_1 + temperature + type_i + batch_j + year_k + conservation_l + G_i + e_{ijkl}$,   (1)
226
where 'GT_{ijkl}' is the germination time of the i-th accession, in the j-th batch, of the k-th seed lot, of
the l-th seed. The factor 'μ_1' is the intercept, 'temperature' the fixed effect of the heat treatment,
'type' the fixed effect of the type of the accession (wild or landrace), 'batch' the fixed effect of the
batch in the germination experiment, 'year' the fixed effect capturing the environmental conditions
of the year of multiplication (2020/2021 or 2021/2022), and 'conservation' the fixed effect of the
number of days the seeds had been conserved before phenotyping (controlling for the detrimental
effect of seed aging). 'G' is the random polygenic effect following a normal distribution of mean 0
and variance matrix $\sigma_G^2 G$, where σ_G^2 is the additive genetic variance, and 'e' is the residual, also
following a normal distribution of mean 0 and variance matrix $\sigma_e^2 I$, where σ_e^2 is the residual
variance. In accordance with the recommendations set forth by Crawley (2007), we also fitted the
model to subdivided datasets: separating wild populations and landraces (two models denoted 1b),
and separating wild and landraces for each temperature (four models denoted 1c). The model was
fitted using the function 'remlf90' of the 'breedR' package (Muñoz & Sanchez, 2024), with the EM
method, and one step of the AI method to estimate the standard error.
241

For the proportion of seeds that germinated, hereafter called the germination rate (GR), it was first corrected for residual factors, a process analogous to the approximated method implemented in GRAMMAR as a pre-treatment for binary variables for computing gBLUP (Aulchenko et al., 2007):

$$GR_{ijkl} = \mu_2 + batch_j + year_k + conservation_l + e_{ijkl}, \tag{2}$$

where 'GR_{ijkl}' is the indicator (0 or 1) of whether the seed germinated before 168 h, 'μ_2' is the intercept, with the indices and the other factors defined as above. Model (2) was fitted to a binomial distribution using the function 'glm' of the 'stats' package. The residues of the fitted model (2) (in the latent scale), thereafter called 'GRA', were then fitted with the following mixed linear model:

$$GRA_{ijkl} = \mu_3 + temperature + type_i + G_i + e_{ijkl}, \tag{3}$$

where 'GRA_{ijkl}' is the adjusted germination rate, 'μ_3' is the intercept, with indices and factors defined as above. Once more, the model was fitted to subdivided datasets: separating wild and landraces (two models denoted 3b), and separating wild and landraces for each temperature (four models denoted 3c). All estimations of GR were conducted on the latent scale. The decision to employ two successive models ensures reliable convergence of the estimation process, which is not currently guaranteed by any available software implementing a (Bayesian) all-in-one GLM approach for GBLUP, including tools like the 'BGLR' package (Pérez & de Los Campos, 2022) that offer robust methods but require separate model specification. The effect of temperature was tested using a Z-test, which provides a reliable approximation of the Student's t-test since the sample size is large, and the effect of the type was tested with a Tukey's HSD test. Heritability, defined as $\sigma_G^2/(\sigma_G^2 + \sigma_e^2)$, was estimated with models (1c) and (3c). For each species, the gBLUP from the models (1c) and (3c) of the different temperatures were compared with the following model:

$$W_i = \mu_4 + C_i + e_i \tag{4}$$

where 'W_i' is the gBLUP at warm temperature, C_i the gBLUP at cold temperature, 'μ_4' is the intercept, with indices and factors defined as above. The GPS coordinate of the sampling site was used to extract the bioclimatic variables from WorldClim v. 2.1 (climate data for 1970-2000; Fick & Hijmans, 2017) using 'geodata' (Hijmans et al., 2024) and 'terra' (Hijmans, 2024) packages, with an accuracy of ten minutes of degree. The Pearson's correlation between the latitude and the 'bio1' bioclimatic variable (annual mean temperature) was very high (-0.89) and we therefore used the latitude as a proxy of bioclimatic variables in the following analyses. For each species, we studied the link between the latitudes of the sampling sites and heat tolerance H_i, defined as W_i-C_i for the i-th accession, with the following model:

$$H_i = \mu_5 + C_i + l_i + e_i \tag{5}$$

where 'H_i' is the heat tolerance of the i-th accession, 'l_i' is the latitude at the sampling site, 'μ_5' is the intercept, with indices and factors defined as above. 'C_i' was added as an explanatory factor to control for the baseline effect.

Finally, we tested the relationship between the gBLUP of GR from model (3c) and the emergence rate by fitting the following model for each species:

$$E_i = \mu_6 + (\mu_3 + C_i) + e_i \tag{6}$$

where 'E_i' is the emergence rate, 'μ_3' is the intercept of the model (3c) for cold temperatures of the corresponding species, 'μ_6' is the intercept, with indices and factors defined as above.

Results

1. The type of the accession was a reliable predictor of its germination time and rate

The genetic divergence among the accessions was mainly explained by the wild-landrace axis (Fig. 1A and 1D). In *B. rapa*, Algerian and Italian wild accessions exhibited stronger divergence from the landraces and wild French accessions. In both MDS and admixture analyses (best K = 3 from CV), French accessions clustered with landraces. The differentiation between wild and landraces was carried by the first axis of the MDS, which explained 77.88% of the variance. In *B. oleracea*, however, this distinction was not as clear as for *B. rapa*: MDS and admixture analyses (best K = 4 from CV) partially differentiated wild and landraces (48.07% in the first axis). The wild accessions were closest to *B. oleracea* ssp. *acephala* (Fig. S1)*.

Similar to the genetic divergence between wild and landrace accessions, there was a clear distinction regarding the germination pattern (Table 2). The type of accession (wild or landrace) was the most discriminating factor: germination time (GT) was significantly longer for wild accessions than for landraces in the case of *B. rapa* (Tukey's test, q = 12.80, p < 0.001; model 1) and of *B. oleracea* (q = 7.98, p < 0.001; model 1), with a slowing effect of 25.92h and 21.82h, respectively. Similarly, germination rate (GR) was significantly lower for wild accessions in both species (q = -10.80 and -5.67, p < 0.001 and < 0.001 for *B. rapa* and *B. oleracea* respectively; model 3). *B. oleracea* accessions were slower than *B. rapa* to germinate, regardless of the type or condition (Student's t-test, t > 19.56, p < 0.001), except for wild accessions at 20°C (t = 0.61, p = 0.540). The heritability of GT and GR was intermediate (Table S1), with an average of 12.25% for GR (ranging from 5 to 20%) and of 21.25% for GT (ranging from 13 to 34%). As expected, the additive variances are systematically larger in wild accessions than in landraces (Table S2), in particular for GR. The genome-wide association studies (GWAS) on germination time and rate across temperatures did not identify any major genes (Fig. S2-S9; see section A of Supplementary Material).

2. Elevated temperature altered germination of *B. oleracea* accessions, but not of *B. rapa* accessions

In *B. rapa*, GR was impacted both by the year of production and the conservation time: significantly higher for those produced in 2021/2022 (Z-test, z = 31.37, p < 0.001; model 2), and significantly lower for those conserved for a long period of time (z = -68.88, p < 0.001; model 2). Similarly, in *B. oleracea*, GR was affected by both the year of production (z = -5.28, p < 0.001; model 2; with higher GR for 2020/2021) and conservation time (z = -11.69, p < 0.001; model 2). Models (1) and (3) account for these effects.

Accessions of *B. rapa* germinated faster when temperature increased, for both wild and landrace accessions (z < -7.36, p < 0.001; model 1b; Fig. 1B), with a reduction of 0.64 h/°C and 1.60 h/°C respectively. Temperature significantly increased GR for landrace accessions (z = 20.28, p < 0.001; model 1c) and significantly decreased it on wild accessions (z = -4.37, p < 0.001; model 1c; Fig. 1C). In contrast, in *B. oleracea* (Fig. 1E), wild accessions germinated significantly slower when temperature increased (z = 9.56, p < 0.001; model 3b), adding 1.31 h/°C, but not for landrace accessions (z = -0.83, p = 0.405). In terms of GR, a slight but significant impact was observed in

both wild and landrace accessions: an increase for landraces (z = 2.62, p = 0.009; model 3b) and a decrease for wild accessions (z = 2.54, p = 0.01; model 3b).

3. Ranking was maintained across temperatures for germination time, but not for germination rate

To test the hypothesis that there is little genotype x environment effect for germination time and rate, i.e. that the ranking of populations was conserved at the two temperatures tested, the low temperature phenotype was connected to high temperature phenotype for both species (Fig. 2; model 4). For GT, the relationship was significant for both species (Z-test, $|z| > 3.95$, $p < 0.001$). The adjusted R^2 were of 0.86 and 0.51 for *B. rapa* and *B. oleracea* respectively. For GR, a significant link was observed for all accessions ($|z| > 2.77$, $p < 0.007$), except for wild accessions of *B. oleracea* (z = 0.71, p = 0.482). The adjusted R^2 were of 0.76 and 0.08 for *B. rapa* and *B. oleracea* respectively.

4. The geographic provenance explained heat tolerance only for *B. oleracea* accessions

The sampling site latitude, which was closely correlated with the annual mean temperature, only impacted the heat tolerance of *B. oleracea* accessions (Fig. 3). We tested this hypothesis in a manner analogous to Finlay-Wilkinson regression (Finlay & Wilkinson, 1963) with model (5). For *B. oleracea* accessions, southern accessions were more tolerant, as latitude had a significant effect on heat tolerance (Z-test, $z < -2.60$, $p < 0.01$) with an adjusted R^2 of 0.50. The effect of latitude was particularly pronounced among wild accessions (the effect of the latitude estimated at -0.017 for wild accessions, and -0.011 for landraces), where Spanish populations exhibited complete tolerance compared to French accessions, with only one exception. In contrast, a slight but significant effect was observed for *B. rapa* wild accessions (z = -2.80, p = 0.006, with the effect of the latitude estimated at -0.002), but no significant effect for *B. rapa* landraces (z = -0.46, p = 0.647, with the effect of the latitude estimated at -0.009), albeit with an adjusted R^2 of 0.29.

5. Germination was correlated with emergence only in *B. rapa* wild accessions

The genetic value of germination rate of *B. rapa* accessions measured in controlled conditions proved to be a reliable predictor of the emergence rate in the field for *B. rapa* accessions (Fig. 4; model 6), whereas it was not the case for *B. oleracea* accessions. For *B. rapa,* both the wild accessions and landraces showed a significant positive relationship (z > 10.56, p < 0.001), with an adjusted R^2 of 0.79. For *B. oleracea*, the relationship was not significant ($|z| = 1.80$, $p > 0.078$), with an adjusted R^2 of 0.02.

Discussion

The extensive sampling of Mediterranean landscapes revealed a strong pattern of differentiation between the wild populations and the landraces, manifesting in both genetics and germination patterns. The elevated temperatures had a significant inhibitory effect on the *B. oleracea* accessions, whereas the *B. rapa* accessions demonstrated resilience and were able to thrive despite the temperature increase. Comparing the gBLUP at both temperatures, results showed that an accession with superior germination capacity at 20°C also exhibited a robust capacity at warmer temperatures (30/35°C), indicating minimal GxE. Nevertheless, the geographic provenance explained heat tolerance for *B. oleracea* accessions, wherein southern accessions demonstrated superior resilience to heat stress. Finally, the germination rate measured in laboratory settings proved to be a reliable predictor of emergence in the field only for *B. rapa*, suggesting other factors involved for *B. oleracea* emergence.

1. Different stories explains the different patterns of germination
The critical factor influencing germination appears to be the 'type' of the accessions, i.e., whether they were sampled in the wild or a cultivated landrace, regardless of the genetic clustering (e.g., French wild accessions of *B. rapa* cluster with landraces). It is important to note that a maternal effect is excluded as an explanatory factor, since all populations were multiplied for one generation in a common environment. As an illustration, germination processes of French *B. rapa* 'wild' populations are closest to that of wild populations, with lower germination rates (GR) and higher germination times (GT), while being genetically closer to landraces. The literature suggests that these French populations may be feral (McAlvay et al., 2021). If this is the case, our results show that they tend to revert back to patterns akin to wild populations for germination. Similarly, the literature also suggests that all wild accessions of *B. oleracea* are more likely to be feral (Maggioni et al., 2020; Mittell et al., 2020; Cai et al., 2022; Saban et al., 2023), yet these populations displayed differences in GR and GT between 'types' of populations, albeit with weaker effects. This phenomenon suggests that, regardless of the ancestry of the populations, the environment is less predictable in the wild than in cultivated fields, so that bet-hedging strategies (Bewley, 1997) based on variable seed dormancy capacities are highly advantageous. This adaptation is evident in *B. rapa* French populations, where dormancy – a trait typical of wild plants – was observed with high frequency (data not shown). In landraces, we can assume that such bet-hedging strategies have been selected against to favour homogeneity in fields. This is consistent with the remarkably low variance observed for GR in landraces, which marks both the domestication bottleneck and the exhaustion of genetic variance by selection. Reverting to pre-domestication traits is common, especially for seed dormancy (Ayal & Levy, 2005; Zhou et al., 2021), although in our case it should be treated with caution due to the very low variance remaining in landraces.

The different level of domestication between *B. rapa* and *B. oleracea* may also explain the lower differences between 'types' of *B. oleracea*. Either germination rate of *B. oleracea* has always been inherently high, and the selection pressure in cultivated lines was weak, or the observed patterns are the result of a shared selective pressure, a domestication bottleneck that led to the fixation of a high germination rate. In regard to our findings, the weak selection hypothesis seems unlikely: the preliminary investigation revealed no evidence of dormancy in *B. oleracea* accessions (data not

shown), one of the first sign of domestication (Fullet & Allaby, 2009), and germination traits of cultivated lines of *B. napus* have been reported to be under strong selection (Laurençon et al., 2024). Futhermore, seed vigor is more closely correlated with vegetative growth than with performance at full reproductive maturity (Tekrony & Egli, 1991), which should result in a stronger selection for seed vigor in cultivated lines of *B. oleracea* that are less often exploited for their seeds.

Interestingly, our findings indicate that the germination response to temperature is correlated with sampling site latitude (i.e., annual mean temperature) only for *B. oleracea*. The range of temperatures tested is a stress for both species, with different intensity depending on the accession, which is reflected in the different levels of heat tolerance among the accessions. We expected accessions at lower latitudes to display higher adaptation to heat stress, resulting in higher heat tolerance, as they have been exposed to this selective pressure, and this is the case for *B. oleracea*, but lesser for *B. rapa*. The lower nucleotide diversity of *B. oleracea* than *B. rapa* (Saban et al., 2023) could indicate that stronger selection has acted on the former, perhaps imputable to a more pronounced domestication process or lower population sizes. In this study, the adaptation of landraces and feral populations to the local environment may be a signature of selection accounting for the environmental profiling, i.e., envirotyping (Xu, 2016; Colouer et al., 2019). In contrast, the lack of such a correlation in *B. rapa*, though a small effect on wild populations, suggests a strong selection of germination traits that are advantageous under a wide range of controlled agricultural environments, thereby reducing the influence of local climatic factors on germination (Purugganan & Fuller, 2009).

2. Genetic improvement and assisted migration for heat tolerance would be beneficial
Our findings show that there is genetic variance for germination traits, which suggests that those traits in these *Brassica* landraces could be selected and introgressed into commercial germplasms, which generally have an optimal germination temperature lower than 30-35°C (ISTA, 2024). The genotype x environment interaction was found to be minimal and there was significant genetic variances for GR and GT, which indicate that breeding can be conducted and worth promoting, given the unequal utilization of landraces across different countries (e.g., more common in Algeria than in France). For instance, *B. napus* seedlings show inhibited establishment at high temperatures (Zhang et al., 2015), indicating a potential benefit from integrating heat-tolerant traits from *B. rapa*. The resilience of *B. rapa* landraces suggests that they will not suffer from future temperature rises and could be useful to improve commercial germplasms (Cochrane et al., 2015; Castillo-Lorenzo et al., 2019). Additionally, seed priming techniques could be explored to further enhance the germination performance of *B. rapa* under varying environmental conditions (Paparella et al., 2015). In contrast, *B. oleracea* landraces experience heat as a stressor, significantly affecting their germination. However, wild southern accessions, particularly those from Spain, show higher heat tolerance and could serve as sources for assisted migration into French 'wild' populations. Indeed, assisted migration, grounded in evolutionary knowledge, can enhance the strategy to cope with rapid environmental changes (Hoffmann & Sgrò, 2011; Walck et al., 2011). The Spanish wild accessions could also prove useful for introgression into elite *B. oleracea* germplasms. Optimizing their use in breeding program is an active field of research; nowadays, pre-breeding and bridging populations act as a buffer to limit the linkage drag from wild materials and to maintain the level of performance of elite crop lines (Cowling et al., 2009; Sanchez et al., 2023).

The introgression of these traits is crucial for *B. napus* that could benefit from the germinative vigor
of *B. rapa* landraces, especially if it also enhances emergence rates and crop field performance
(Smith et al., 2000; Ghassemi-Golezani et al., 2010). Although our genome-wide association studies
(GWAS) did not detect any major genes – consistent with the polygenic nature of germination traits
(Hatzig et al., 2015; Bettey et al., 2000) – there is potential for improvement through phenomic and
genomic selection (Laurençon et al., 2024). Nonetheless, the study has identified valuable sources
of genetic diversity within *Brassica* populations, thereby highlighting the potential of expanding
sampling efforts within these populations to circumvent the effect of population structure and
increase the power of GWAS analyses (Tibbs Cortes et al., 2021). This would enable the
identification of QTL underlying adaptation (Basnet et al., 2015), particularly those conferring
resilience to temperature variation and improve the understanding of the genetic basis of adaptive
traits (Huang & Han, 2014).

Conclusion
The germination time and rate of an unexplored diversity of 25,080 seeds of *Brassica rapa* and
12,725 seeds of *Brassica oleracea* were compared at two temperature conditions. The results
demonstrate that temperature elevation does not constitute a stressor for *B. rapa* accessions, which
maintained high germination rates and short germination times at high temperatures, especially
landraces. In contrast, accessions of *B. oleracea* exhibited clear signs of stress when exposed to
elevated temperatures, which in turn leads to a notable decline in their germination performance. In
the context of global warming, wild populations of *B. oleracea* could benefit from assisted
migration in order to adapt to the anticipated temperature increases. This can be achieved by
incorporating traits from southern wild populations that have demonstrated high heat tolerance,
likely due to their adaptation to higher annual mean temperatures. The integration of these heat-
tolerant traits into commercial germplasms would enhance the resilience of *B. oleracea*, ensuring its
productivity and sustainability.

Funding

The investigation was supported by H2020 Prima funding (grant no. 1425), entitled "BrasExplor: Wide exploration of genetic diversity in *Brassica* species for sustainable crop production"; by INRAE through the TSARA initiative (Transforming food systems and agriculture through a partnership research with Africa), which fosters Franco-Algerian collaborations; and by the France Génomique National infrastructure, funded as part of the «Investissements d'Avenir» program managed by the Agence Nationale pour la Recherche (contract ANR-10-INBS-09).

Acknowledgement

We thank the Genetic Resource Centers BrACySol (https://www6.rennes.inrae.fr/igepp_eng/About-IGEPP/Platforms/BrACySol), especially V. Richer, S. Théréné, and S. Doré; and the Agricultural Institute of Slovenia (https://www.kis.si/en/) for providing seeds from different landraces. We thank the GenOuest bioinformatics platform (https://www.genouest.org/), as well as the Plant imaging platform PHENOTIC in Angers (https://phenotic.hub.inrae.fr/). We also extend our appreciation to the staff who provided care for the plant material, particularly L. Charlon, J-P. Constantin, and F. Letertre; P. Glory and P. Vallée for field experiments; and to M. Gautier and M. Delmond for the SNP calling.

Contributions

MHW, CF, DS, and AMC designed the study; FG and KL generated Illumina data; MHW, AD performed experiments and contributed data; MT, MHW, EC, and LG analyzed the data; MT, LG, and MHW drafted the manuscript. All authors read and approved the final version of the manuscript.

Conflict of interest statement

The authors declare that there is no conflict of interest.

Data availability

The short-read resequencing data that support the findings are available in NCBI's sequence read archive (SRA) under the project PRJNA1174687. The VCF file and the raw phenotype dataset will be freely available with a DOI upon acceptance. Seeds of the accessions are available at the following Genetic Resource Centers: BrACySol for French accessions, de Los Campos seed genebank (ESP003) for Spanish accessions, and Agricultural Institute of Slovenia for Slovenian accessions. The unique accession numbers are detailed in Table S3. For inquiries regarding Algerian, Italian, and Tunisian accessions, please refer to the contact specified in Table S3.

Figure legends & Tables

Figure 1. Genetic clusters and germination of Brassica wild and landrace accessions. Wild accessions are in orange and blue; landraces are in green. **(A)** MDS of the genotypes of *B. rapa*: the first axis explains 77.58 %, and the second 12.11 %. Admixture plot in the bottom with the corresponding color (light green is for French landraces, colored post-analysis). **(B)** Germination time (in hours) of *B. rapa*: 20°C in blue, 35°C in red. **(C)** Germination rate (in percent) of *B. rapa*: 20°C in blue, 35°C in red. **(D)** MDS of the genotypes of *B. oleracea*: the first axis explains 48.07 %, 23.23 %. Admixture plot in the bottom. **(E)** Germination time (in hours) of *B. oleracea*: 20°C in blue, 30°C in red. **(F)** Germination rate (in percent) of *B. oleracea*: 20°C in blue, 30°C in red.

Figure 2. gBLUP of germination traits at 20°C as a function of the gBLUP of germination traits at 30/35°C. Wild accessions in orange; landraces in green. **(A)** BLUP of germination time of *B. rapa*. **(B)** BLUP of germination rate of *B. rapa*. **(C)** BLUP of germination time of *B. oleracea*. **(D)** BLUP of germination rate of *B. oleracea*.

Figure 3. Heat tolerance (defined as the difference between the gBLUP at 30/35°C and that at 20°C) as a function of latitude. Wild accessions in orange; landraces in green. Provenance: Algeria: full circle; France: full triangle; Italy: full square; Slovenia: cross; Spain: crossed square; Tunisia: star. **(A)** Heat tolerance of *B. rapa*. **(B)** Heat tolerance of *B. oleracea*.

Figure 4. Emergence rate in field as a function of the gBLUP of germination rate in controlled conditions at 20°C. **(A)** Emergence rate of *B. rapa*. **(B)** Emergence rate of *B. oleracea*.

	B. rapa		B. oleracea	
	Wild	Landrace	Wild	Landrace
Algeria	28	22		6
France	16	21	8	26
Italy	17	5		4
Slovenia		3		11
Spain			8	
Tunisia		5		3

Table 1. The accessions by types (wild or landrace) and countries (Algeria, France, Italy, Slovenia, Spain, and Tunisia), out of the 117 accessions of *B. rapa* and 66 accessions of *B. oleracea*.

	T°	B. rapa		B. oleracea	
		Wild	Landrace	Wild	Landrace
GR	20°C	0.39 ± 0.49	0.97 ± 0.16	0.88 ± 0.32	0.96 ± 0.20
	30/35°C	0.67 ± 0.47	0.97 ± 0.18	0.84 ± 0.37	0.96 ± 0.20
GT	20°C	74.0 ± 41.9	42.6 ± 23.9	74.5 ± 23.6	59.7 ± 24.0
	30/35°C	56.6 ± 40.4	27.0 ± 18.7	85.9 ± 40.0	57.8 ± 33.7

Table 2. Average germination rate (before 7 days) and germination time (in hours), and standard deviations, of *B. rapa* and *B. oleracea* in controlled conditions (20°C and 35°C for *B. rapa*, 20°C and 30°C for *B. oleracea*).

References

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome research*, *19*(9), 1655-1664.

Allender, C. J., & King, G. J. (2010). Origins of the amphiploid species Brassica napus L. investigated by chloroplast and nuclear molecular markers. *BMC plant biology*, *10*, 1-9.

Angadi, S. V., Cutforth, H. W., Miller, P. R., McConkey, B. G., Entz, M. H., Brandt, S. A., & Volkmar, K. M. (2000). Response of three Brassica species to high temperature stress during reproductive growth. *Canadian Journal of plant science*, *80*(4), 693-701.

Arunanondchai, P., Fei, C., Fisher, A., McCarl, B. A., Wang, W., & Yang, Y. (2018). How does climate change affect agriculture?. In *The Routledge handbook of agricultural economics* (pp. 191-210). Routledge.

Aulchenko, Y. S., De Koning, D. J., & Haley, C. (2007). GRAMMAR: a fast and simple method for genome-wide pedigree-based quantitative trait loci association analysis. *Genetics*, *177*(1), 577-585.

Ayal, S., & Levy, A. A. (2005). Wheat domestication and dedomestication-What are the odds?.

Barnabás, B., Jäger, K., & Fehér, A. (2008). The effect of drought and heat stress on reproductive processes in cereals. *Plant, cell & environment*, *31*(1), 11-38.

Basnet, R. K., Duwal, A., Tiwari, D. N., Xiao, D., Monakhos, S., Bucher, J., ... & Maliepaard, C. (2015). Quantitative trait locus analysis of seed germination and seedling vigor in Brassica rapa reveals QTL hotspots and epistatic interactions. *Frontiers in plant science*, *6*, 1032.

Becklin, K. M., Anderson, J. T., Gerhart, L. M., Wadgymar, S. M., Wessinger, C. A., & Ward, J. K. (2016). Examining plant physiological responses to climate change through an evolutionary lens. *Plant physiology*, *172*(2), 635-649.

Belser, C., Istace, B., Denis, E., Dubarry, M., Baurens, F. C., Falentin, C., ... & Aury, J. M. (2018). Chromosome-scale assemblies of plant genomes using nanopore long reads and optical maps. *Nature plants*, *4*(11), 879-887.

Bettey, M., Finch-Savage, W. E., King, G. J., & Lynn, J. R. (2000). Quantitative genetic analysis of seed vigour and pre-emergence seedling growth traits in Brassica oleracea. *The New Phytologist*, *148*(2), 277-286.

Bewley, J. D. (1997). Seed germination and dormancy. *The plant cell*, *9*(7), 1055.

Cai, C., Bucher, J., Bakker, F. T., & Bonnema, G. (2022). Evidence for two domestication lineages supporting a middle-eastern origin for Brassica oleracea crops from diversified kale populations. *Horticulture research, 9*, uhac033.

Castillo-Lorenzo, E., Finch-Savage, W. E., Seal, C. E., & Pritchard, H. W. (2019). Adaptive significance of functional germination traits in crop wild relatives of Brassica. *Agricultural and Forest Meteorology*, *264*, 343-350.

Chaudhry, S., & Sidhu, G. P. S. (2022). Climate change regulated abiotic stress mechanisms in plants: A comprehensive review. *Plant Cell Reports*, *41*(1), 1-31.

Cheng, F., Sun, R., Hou, X., Zheng, H., Zhang, F., Zhang, Y., ... & Wang, X. (2016). Subgenome parallel selection is associated with morphotype diversification and convergent crop domestication in Brassica rapa and Brassica oleracea. *Nature genetics*, *48*(10), 1218-1224.

Chmielewski, F. M., Müller, A., & Bruns, E. (2004). Climate changes and trends in phenology of fruit trees and field crops in Germany, 1961–2000. *Agricultural and Forest Meteorology*, *121*(1-2), 69-78.

Clauss, M. J., & Venable, D. L. (2000). Seed germination in desert annuals: an empirical test of adaptive bet hedging. *The American Naturalist, 155*(2), 168-186.

Cochrane, J. A., Hoyle, G. L., Yates, C. J., Wood, J., & Nicotra, A. B. (2015). Climate warming delays and decreases seedling emergence in a Mediterranean ecosystem. *Oikos, 124*(2), 150-160.

Corlouer, E., Gauffreteau, A., Bouchet, A. S., Bissuel-Bélaygue, C., Nesi, N., & Laperche, A. (2019). Envirotypes based on seed yield limiting factors allow to tackle G× E interactions. *Agronomy*, *9*(12), 798.

Cowling, W. A., Buirchell, B. J., & Falk, D. E. (2009). A model for incorporating novel alleles from the primary gene pool into elite crop breeding programs while reselecting major genes for domestication or adaptation. *Crop and Pasture Science, 60*(10), 1009-1015.

Crawley, M. J. (2007). Statistical modelling. In *The R book* (pp. 323–386). Chichester, UK: John Wiley & Sons.

DaMatta, F. M., Grandis, A., Arenque, B. C., & Buckeridge, M. S. (2010). Impacts of climate changes on crop physiology and food quality. *Food Research International, 43*(7), 1814-1823.

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., ... & Li, H. (2021). Twelve years of SAMtools and BCFtools. *Gigascience, 10*(2), giab008.

Demilly, D., Ducournau, S., Wagner, M. H., & Dürr, C. (2014). Digital imaging of seed germination. *Plant image analysis: fundamentals and applications, 1*, 147-164.

Dürr, C., Dickie, J. B., Yang, X. Y., & Pritchard, H. W. (2015). Ranges of critical temperature and
water potential values for the germination of species worldwide: contribution to a seed trait
database. *Agricultural and forest meteorology*, *200*, 222-232.

Ebone, L. A., Caverzan, A., Tagliari, A., Chiomento, J. L. T., Silveira, D. C., & Chavarria, G.
(2020). Soybean seed vigor: Uniformity and growth as key factors to improve yield. *Agronomy*,
10(4), 545.

El-Keblawy, A., Al-Shamsi, N., & Mosa, K. (2018). Effect of maternal habitat, temperature and
light on germination and salt tolerance of Suaeda vermiculata, a habitat-indifferent halophyte of arid
Arabian deserts. *Seed Science Research*, *28*(2), 140-147.

Falentin, C., Hadj-Arab, H., Aissiou, F., Bartoli, C., Bazan, G., Boudet, M., ... & Chèvre, A.-M.
(2024). Combined cytogenetic and molecular methods for taxonomic verification and description of
Brassica populations deriving from different origins. *Genetic Resources*, *5*(9), 61–71.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for
global land areas. *International journal of climatology*, *37*(12), 4302-4315.

Finch-Savage, W. E., & Bassel, G. W. (2016). Seed vigour and crop establishment: extending
performance beyond adaptation. *Journal of experimental botany*, *67*(3), 567-591.

Finlay, K. W., & Wilkinson, G. N. (1963). The analysis of adaptation in a plant-breeding
programme. *Australian journal of agricultural research*, *14*(6), 742-754.

Franks, S. J., Weber, J. J., & Aitken, S. N. (2014). Evolutionary and plastic responses to climate
change in terrestrial plant populations. *Evolutionary applications*, *7*(1), 123-139.

Fuller, D. Q., & Allaby, R. (2009). Seed dispersal and crop domestication: shattering, germination
and seasonality in evolution under cultivation. *Annual plant reviews volume 38: fruit development*
and seed dispersal, *38*, 238-295.

Galloway, L. F. (2005). Maternal effects provide phenotypic adaptation to local environmental
conditions. *New phytologist*, *166*(1), 93-100.

Gareca, E. E., Vandelook, F., Fernández, M., Hermy, M., & Honnay, O. (2012). Seed germination,
hydrothermal time models and the effects of global warming on a threatened high Andean tree
species. *Seed Science Research*, *22*(4), 287-298.

Ghassemi-Golezani, K., Bakhshy, J., Yaeghoob, R. A. E. Y., & Hossainzadeh-Mahootchy, A. (2010).
Seed vigor and field performance of winter oilseed rape (Brassica napus L.) cultivars. *Notulae*
Botanicae Horti Agrobotanici Cluj-Napoca, *38*(3), 146-150.

Goldsmith, J., Scheipl, F., Huang, L., Wrobel, J., Di, C., Gellar, J., ... & Cui E (2024). refund:
Regression with Functional Data. R package version 0.1-35,
https://CRAN.R-project.org/package=refund

Hampton, J. G., Conner, A. J., Boelt, B., Chastain, T. G., & Rolston, P. (2016). Climate change:
seed production and options for adaptation. *Agriculture, 6*(3), 33.

Hatfield, J. L., & Prueger, J. H. (2015). Temperature extremes: Effect on plant growth and
development. *Weather and climate extremes, 10*, 4-10.

Hatzig, S. V., Frisch, M., Breuer, F., Nesi, N., Ducournau, S., Wagner, M. H., ... & Snowdon, R. J.
(2015). Genome-wide association mapping unravels the genetic control of seed germination and
vigor in Brassica napus. *Frontiers in plant science, 6*, 221.

Hijmans, R. J., Barbosa, M., Ghosh, A., & Mandel, A. (2024). geodata: Download Geographic
Data. R package version 0.6-2, https://CRAN.R-project.org/package=geodata.

Hijmans, R. (2024). terra: Spatial Data Analysis. R package version 1.7-78, https://CRAN.R-
project.org/package=terra.

Hoffmann, A. A., & Sgrò, C. M. (2011). Climate change and evolutionary adaptation. *Nature,
470*(7335), 479-485.

Huang, X., & Han, B. (2014). Natural variations and genome-wide association studies in crop
plants. *Annual review of plant biology, 65*(1), 531-551.

IPCC. (2013). Climate Change 2013: The Physical Science Basis. Contribution of Working Group I
to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (T. F. Stocker, D.
Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, & P. M.
Midgley, Eds.). Cambridge University Press.

IPCC. (2021). Climate Change 2021: The Physical Science Basis. Contribution of Working Group I
to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change (V. Masson-
Delmotte, P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I.
Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O.
Yelekçi, R. Yu, & B. Zhou, Eds.). Cambridge University Press.

ISTA. (2024). International rules for seed testing: International Seed Testing Association (ISTA).
Wallisellen, Switzerland.
Jump, A. S., & Peñuelas, J. (2005). Running to stand still: adaptation and the response of plants to
rapid climate change. *Ecology letters, 8*(9), 1010-1020.
735

736 Jump, A. S., Peñuelas, J., Rico, L., Ramallo, E., Estiarte, M., Martinez-izquierdo, J. A., & Lloret, F.
737 (2008). Simulated climate change provokes rapid genetic change in the Mediterranean shrub
738 Fumana thymifolia. *Global Change Biology*, *14*(3), 637-643.
739
740 Lamichhane, J. R., Debaeke, P., Steinberg, C., You, M. P., Barbetti, M. J., & Aubertot, J. N. (2018).
741 Abiotic and biotic factors affecting crop seed germination and seedling emergence: a conceptual
742 framework. *Plant and soil*, *432*, 1-28.
743
744 Laurençon, M., Legrix, J., Wagner, M. H., Demilly, D., Baron, C., Rolland, S., ... & Nesi, N. (2024).
745 Genomic and phenomic predictions help capture low-effect alleles promoting seed germination in
746 oilseed rape in addition to QTL analyses. *Theoretical and Applied Genetics*, *137*(7), 1-16.
747
748 Li, R., Li, Y., Kristiansen, K., & Wang, J. (2008). SOAP: short oligonucleotide alignment program.
749 *Bioinformatics*, *24*(5), 713-714.
750
751 Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
752 *arXiv preprint arXiv:1303.3997*.
753
754 Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler
755 transform. *bioinformatics*, *25*(14), 1754-1760.
756
757 Lobell, D. B., & Field, C. B. (2007). Global scale climate–crop yield relationships and the impacts
758 of recent warming. *Environmental research letters*, *2*(1), 014002.
759
760 Maggioni, L., von Bothmer, R., Poulsen, G., & Härnström Aloisi, K. (2020). Survey and genetic
761 diversity of wild Brassica oleracea L. germplasm on the Atlantic coast of France. *Genetic Resources*
762 *and Crop Evolution*, *67*(7), 1853-1866.
763
764 Maillet, L., Norest, M., Kautsky, A., Geraci, A., Oddo, E., Troia, A., ... & Bartoli, C. (2023). Plant
765 genetic bases explaining microbiota diversity shed light into a novel holobiont generalist gene
766 theory. *bioRxiv*, 2023-12.
767
768 McAlvay, A. C., Ragsdale, A. P., Mabry, M. E., Qi, X., Bird, K. A., Velasco, P., ... & Emshwiller, E.
769 (2021). Brassica rapa domestication: untangling wild and feral forms and convergence of crop
770 morphotypes. *Molecular biology and evolution*, *38*(8), 3358-3372.
771
772 McCouch, S., Baute, G. J., Bradeen, J., Bramel, P., Bretting, P. K., Buckler, E., ... & Zamir, D.
773 (2013). Feeding the future. *Nature*, *499*(7456), 23-24.
774
775 McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., ... & DePristo,
776 M. A. (2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
777 generation DNA sequencing data. *Genome research*, *20*(9), 1297-1303.
778

Mittell, E. A., Cobbold, C. A., Ijaz, U. Z., Kilbride, E. A., Moore, K. A., & Mable, B. K. (2020).
Feral populations of Brassica oleracea along Atlantic coasts in western Europe. *Ecology and*
evolution, *10*(20), 11810-11825.

Muñoz, F., & Sanchez, L. (2024). breedR: Statistical Methods for Forest Genetic Resources
Analysts. R package version 0.12-5, commit 92779f422f72823ce6db28e805592a9f1be78b27,
https://github.com/famuvie/breedR

Nelson, M. N., Nesi, N., Barrero, J. M., Fletcher, A. L., Greaves, I. K., Hughes, T., ... & Kirkegaard,
J. A. (2022). Strategies to improve field establishment of canola: a review. *Advances in Agronomy*,
175, 133-177.

Nevo, E., Fu, Y. B., Pavlicek, T., Khalifa, S., Tavasi, M., & Beiles, A. (2012). Evolution of wild
cereals during 28 years of global warming in Israel. *Proceedings of the National Academy of*
Sciences, *109*(9), 3412-3415.

Nicotra, A. B., Atkin, O. K., Bonser, S. P., Davidson, A. M., Finnegan, E. J., Mathesius, U., ... &
van Kleunen, M. (2010). Plant phenotypic plasticity in a changing climate. *Trends in plant science*,
15(12), 684-692.

Paparella, S., Araújo, S. S., Rossi, G., Wijayasinghe, M. A. L. A. K. A., Carbonera, D., &
Balestrazzi, A. (2015). Seed priming: state of the art and new perspectives. *Plant cell reports*, *34*,
1281-1293.

Pareek, A., Dhankher, O. P., & Foyer, C. H. (2020). Mitigating the impact of climate change on
plant productivity and ecosystem sustainability. *Journal of Experimental Botany*, *71*(2), 451-456.

Penfield, S., & MacGregor, D. R. (2017). Effects of environmental variation during seed production
on seed dormancy and germination. *Journal of experimental botany*, *68*(4), 819-825.

Pérez-Rodríguez, P., & de Los Campos, G. (2022). Multitrait Bayesian shrinkage and variable
selection models with the BGLR-R package. *Genetics*, *222*(1), iyac112.

Purugganan, M. D., & Fuller, D. Q. (2009). The nature of selection during plant domestication.
Nature, *457*(7231), 843-848.

Pollock, B. M., & Roos, E. E. (1972). Seed and seedling vigor. *Seed Biology, I. Importance,*
development and germination, 314-387.

R Core Team (2024). R: A Language and Environment for Statistical Computing. R Foundation for
Statistical Computing, Vienna, Austria. https://www.R-project.org/

Rakow, G. (2004). Species origin and economic importance of Brassica. In *Brassica* (pp. 3-11).
Berlin, Heidelberg: Springer Berlin Heidelberg.

Raza, A., Razzaq, A., Mehmood, S. S., Zou, X., Zhang, X., Lv, Y., & Xu, J. (2019). Impact of climate change on crops adaptation and strategies to tackle its outcome: A review. *Plants, 8*(2), 34.

Reed, R. C., Bradford, K. J., & Khanday, I. (2022). Seed germination and vigor: ensuring crop sustainability in a changing climate. *Heredity, 128*(6), 450-459.

Renzi, J. P., Coyne, C. J., Berger, J., von Wettberg, E., Nelson, M., Ureta, S., ... & Brus, J. (2022). How could the use of crop wild relatives in breeding increase the adaptation of crops to marginal environments?. *Frontiers in Plant Science, 13*, 886162.

Rosenzweig, C., Elliott, J., Deryng, D., Ruane, A. C., Müller, C., Arneth, A., ... & Jones, J. W. (2014). Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison. *Proceedings of the national academy of sciences, 111*(9), 3268-3273.

Saban, J. M., Romero, A. J., Ezard, T. H., & Chapman, M. A. (2023). Extensive crop–wild hybridization during Brassica evolution and selection during the domestication and diversification of Brassica crops. *Genetics, 223*(4), iyad027.

Salgotra, R. K., & Chauhan, B. S. (2023). Genetic diversity, conservation, and utilization of plant genetic resources. *Genes, 14*(1), 174.

Sanchez, D., Sadoun, S. B., Mary-Huard, T., Allier, A., Moreau, L., & Charcosset, A. (2023). Improving the use of plant genetic resources to sustain breeding programs' efficiency. *Proceedings of the National Academy of Sciences, 120*(14), e2205780119.

Sloat, L. L., Davis, S. J., Gerber, J. S., Moore, F. C., Ray, D. K., West, P. C., & Mueller, N. D. (2020). Climate adaptation by crop migration. *Nature communications, 11*(1), 1243.

Smith, S. E., Riley, E., Tiss, J. L., & Fendenheim, D. M. (2000). Geographical variation in predictive seedling emergence in a perennial desert grass. *Journal of Ecology, 88*(1), 139-149.

Tanksley, S. D., & McCouch, S. R. (1997). Seed banks and molecular maps: unlocking genetic potential from the wild. *science, 277*(5329), 1063-1066.

Taylor, A. G. (2020). Seed storage, germination, quality, and enhancements. In *The physiology of vegetable crops* (pp. 1-30). Wallingford UK: CABI.

TeKrony, D. M., & Egli, D. B. (1991). Relationship of seed vigor to crop yield: a review. *Crop science, 31*(3), 816-822.

Thornton, P. K., Ericksen, P. J., Herrero, M., & Challinor, A. J. (2014). Climate variability and vulnerability to climate change: a review. *Global change biology, 20*(11), 3313-3328.

Tibbs Cortes, L., Zhang, Z., & Yu, J. (2021). Status and prospects of genome-wide association studies in plants. *The plant genome, 14*(1), e20077.

Tribouillois, H., Dürr, C., Demilly, D., Wagner, M. H., & Justes, E. (2016). Determination of germination response to temperature and water potential for a wide range of cover crop species and related functional groups. *PloS one, 11*(8), e0161185.

Wagner, M., Demilly, D., Ducournau, S., Dürr, C., & Léchappé, J. (2011). Computer vision for monitoring seed germination from dry state to young seedlings. *Seed Test, 142*, 49-51.

Walck, J. L., Hidayati, S. N., Dixon, K. W., Thompson, K. E. N., & Poschlod, P. (2011). Climate change and plant regeneration from seed. *Global change biology, 17*(6), 2145-2161.

Warwick, S. I., & Black, L. D. (1991). Molecular systematics of Brassica and allied genera (subtribe Brassicinae, Brassiceae)—chloroplast genome and cytodeme congruence. *Theoretical and Applied Genetics, 82*, 81-92.

Wheeler, T., & Von Braun, J. (2013). Climate change impacts on global food security. *Science, 341*(6145), 508-513.

Wolf, J. B., & Wade, M. J. (2009). What are maternal effects (and what are they not)?. *Philosophical Transactions of the Royal Society B: Biological Sciences, 364*(1520), 1107-1115.

Xu, Y. (2016). Envirotyping for deciphering environmental impacts on crop plants. *Theoretical and Applied Genetics, 129*, 653-673.

Yang, J., Lee, S. H., Goddard, M. E., & Visscher, P. M. (2011). GCTA: a tool for genome-wide complex trait analysis. *The American Journal of Human Genetics, 88*(1), 76-82.

Zamir, D. (2001). Improving plant breeding with exotic genetic libraries. *Nature reviews genetics, 2*(12), 983-989.

Zhang, J., Jiang, F., Yang, P., Li, J., Yan, G., & Hu, L. (2015). Responses of canola (Brassica napus L.) cultivars under contrasting temperature regimes during early seedling growth stage as revealed by multiple physiological criteria. *Acta Physiologiae Plantarum, 37*, 1-10.

Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D. B., Huang, Y., ... & Asseng, S. (2017). Temperature increase reduces global yields of major crops in four independent estimates. *Proceedings of the National Academy of sciences, 114*(35), 9326-9331.

Zhou, C., Feng, Y., Li, G., Wang, M., Jian, J., Wang, Y., ... & Yang, J. (2021). The new is old: Novel germination strategy evolved from standing genetic variation in weedy rice. *Frontiers in Plant Science, 12*, 699464.