Chemotherapy resistance due to epithelial-to-mesenchymal transition is caused by abnormal lipid metabolic balance

Atsushi Matsumoto[1]†, Akihito Inoko[2,3]†, Waki Hosoda[4], Takahiro Kojima[5], Koji Ohnishi[2], Junichi Ikenouchi[6] *

Affiliations

1. Department of Biology, Faculty of Sciences, Kyushu University, Fukuoka, 819-0395, Japan

2. Department of Pathology, Aichi Medical University School of Medicine, Nagakute, 480-1195, Japan

3. Division of Cancer Epidemiology and Prevention, Aichi Cancer Center Research Institute, Nagoya, 464-8681, Japan

4. Department of Pathology and Molecular Diagnostics, Aichi Cancer Center Hospital, Nagoya, 464-8681, Japan

5. Department of Urology, Aichi Cancer Center Hospital, Nagoya, 464-8681, Japan

6. Department of Biochemistry, Kyushu University Graduate School of Medical Sciences, Fukuoka, 812-8582, Japan.

†. These authors contributed equally to this work.

*Corresponding author. Junichi Ikenouchi M.D., Ph.D.

Tel.: +81-92-802-4292; Fax: +81-92-802-4292

Email: ikenouchi.junichi.033@m.kyushu-u.ac.jp

Abstract

Invasive cancer is defined by the loss of epithelial cell traits resulting from the ectopic expression of epithelial-mesenchymal transition (EMT)-related transcription factors such as Snail. Although EMT is known to impart chemoresistance to cancer cells, the precise molecular mechanisms remain elusive. We found that Snail expression confers chemoresistance by upregulating the cholesterol efflux pump ABCA1 as a countermeasure to the excess of cytotoxic free cholesterol relative to its major

interaction partner in cellular membranes, sphingomyelin. This imbalance is introduced

by the transcriptional repression of enzymes involved in the biosynthesis of

sphingomyelin by Snail. Inhibiting esterification of cholesterol, which renders it inert,

selectively suppresses growth of a xenograft model of Snail-positive kidney cancer. Our

findings offer a new perspective on lipid-targeting strategies for invasive cancer

therapy.

Introduction

Metastatic cancer, characterized by the spread of tumor cells to distant organs, remains

a major challenge in cancer treatment due to its high mortality rate. Metastasis involves

a series of biological events, such as acquisition of cancer stem cell characteristics,

tumor-microenvironment interactions, alterations in cell metabolism and resistance to

chemotherapy, that are mediated by epithelial-mesenchymal transition (EMT) (Fischer

et al, 2015; Lambert & Weinberg, 2021; Nieto *et al*, 2016; Zheng *et al*, 2015).

Transcription factors driving EMT (EMT-TFs), including Snail, Slug and Twist, that are

ectopically expressed in invasive cancers were initially thought to promote invasion and

metastasis by suppressing the expression of adhesion molecules in cancer cells.

However, since E-cadherin-mediated adhesion is essential for cell survival after

metastasis (Padmanaban *et al*, 2019), invasive cancer cells do not undergo complete

EMT but rather sustain a hybrid state with both epithelial and mesenchymal (hybrid

E/M) characteristics. In a mouse model of pancreatic carcinoma and breast cancers

where key EMT-TFs were deleted, the suppression of EMT did not alter the appearance

of systemic dissemination or metastasis but rather contributed to increased sensitivity to

anticancer drugs (Fischer *et al.*, 2015; Zheng *et al.*, 2015). Therefore, in-depth

understanding of the molecular mechanisms of resistance to anti-cancer drugs conferred

by EMT-TFs is an essential prerequisite for devising effective therapeutic interventions

(Davis *et al*, 2014). It was recently reported that expression of a Rho family GTPase,

RHOJ, is upregulated in cancer cells undergoing EMT and that its activation enhances

the DNA damage response, which enables tumor cells to efficiently repair

chemotherapy-induced DNA damage (Debaugnies *et al*, 2023). Here, we investigated

EMT-dependent acquisition of resistance to more common anticancer drugs that do not

directly cause DNA damage, such as those targeting the ERK signaling pathway, and

found that expression of EMT-TFs causes metabolic alterations of sphingomyelin,

resulting in ectopic induction of drug efflux transporter expression and conferring

chemotherapy resistance to cancer cells.

Results

Snail-induced ABCA1 expression confers chemoresistance to hybrid E/M cells

Renal cell carcinoma (RCC) is notorious for the lack of sensitivity to

chemotherapy (Makhov *et al*, 2018), but the underlying mechanisms are not elucidated.

Nitidine chloride (NC) holds promise as a drug treatment against RCC as it was shown

to have a pro-apoptotic effect on RCC and inhibit tumor growth by suppressing the

ERK signaling pathway in a xenograft model (Cui *et al*, 2020; Fang *et al*, 2014).

However, in human lung adenocarcinoma cells, NC efficacy was inversely correlated

with the expression level of the drug transporter ABCA1: cancer cells with

downregulated ABCA1 were more sensitive to NC treatment (Iwasaki *et al*, 2010).

Therefore, we wondered whether ABCA1 could also confer NC resistance in RCC. We

treated five different RCC cell lines with a combination of NC and an ABCA1 inhibitor

cyclosporin A (CsA). Anti-proliferative effect of NC was significantly enhanced in the

presence of CsA for three lines among five, Caki-1, 786-O and A498 (Fig. 1A). These

cells consistently expressed higher levels of ABCA1 compared to the non-responsive

cells (Fig. 1B), suggesting the acquisition of ABCA1-mediated drug resistance. To

investigate clinical relevance of ABCA1 upregulation in cancers, we analyzed ABCA1

expression in cancer using the UCSC Xena platform. Comparative ABCA1 expression

data of both normal tissue and primary tumors were available for 24 cancer subtypes in

the GDC TCGA database. In addition to glioblastoma and head and neck cancer, all

three types of renal cancers (clear cell carcinoma (ccRCC), chromophobe carcinoma

and papillary cell carcinoma) exhibited significant ABCA1 upregulation in tumor (Fig.

1C and Fig. S1), indicating that ABCA1 upregulation frequently occurs in clinical renal

cancers.

Regulatory mechanisms of ABCA1 expression in cancer cells is largely

unknown. Interestingly, ABCA1 expression was positively correlated with expression

of the EMT-TF Snail in the RCC cell lines (Fig. 1B). To examine whether the

enrichment of ABCA1 protein in malignant cancer cells is detectable in clinical cases,

we conducted immunohistochemical analysis on surgical specimens taken from patients

with primary ccRCC. The findings showed that ABCA1 was upregulated at the lesion

site of high-grade ccRCC (Fig. 1D). This clinical data aligns with our cellular

experimental results, as Snail expression is known to be upregulated in high-grade

ccRCC. Upon detailed comparison among grades, ABCA1 signal was more prominent

in the patients with the high-grade classification (Fig. S2), suggesting that ABCA1 could serve as a marker for high-grade ccRCC. Xena analysis also revealed significantly higher expression of Snail in ccRCC, also supporting the idea that ABCA1 upregulation is correlated with exogenous Snail expression (Fig. S1).

In order to clarify whether Snail expression alone can confer resistance to anti-cancer drugs in epithelial cells, Snail was ectopically expressed in normal epithelial cells and we investigated changes in resistance to anti-cancer drugs. Overexpression of Snail in a mouse mammary gland epithelial cell line EpH4 abolishes transcription of various genes involved in cell-cell adhesion including E-cadherin (Fig. 1E) (Batlle *et al*, 2000; Cano *et al*, 2000; Ikenouchi *et al*, 2003). EpH4-Snail cells exhibited coexistence of the mesenchyme-specific intermediate filament Vimentin and the epithelium-specific Cytokeratin-18 (Fig. 1F), which is a typical feature of a hybrid E/M state of cancer cells. The hybrid E/M EpH4-Snail cells were significantly more resistant to NC than parental EpH4 cells (Fig. 1G). Consistent with the results of kidney cancer cell lines, this resistance was cancelled by CsA (Fig. 1G), and upregulation of ABCA1 expression was also observed in EpH4-Snail cells (Fig. 1H). We also examined another model with a different cell line. In MDCK II, a canine kidney epithelial cell line, TGF-β treatment induced EMT under the expression of the oncogenic KRAS-G12V mutant (Arner *et al*, 2019). The upregulation of ABCA1 expression was also observed in this transient hybrid E/M model (Fig. 1H), suggesting that similar mechanisms to those in EpH4-Snail are also at play in kidney cell EMT. Notably, induction of ABCA1 expression requires a longer period of treatment with TGF-β (day 8) than Snail induction (day 3), indicating that ABCA1 is upregulated in response to cellular alteration by Snail but not induced directly by TGF-β signaling. Furthermore, in comparison between human

esophageal cancer cell lines, Snail-positive TE-8 cells exhibited higher expression of ABCA1 than Snail-negative TE-15 (Fig. 1I). These results suggest that the induction of ABCA1 by ectopic expression of Snail occurs regardless of cellular background.

Intracellular cholesterol accumulation drives ABCA1 expression in hybrid E/M cells

Then how does Snail induce ABCA1 expression, since as a transcriptional repressor it is incapable of doing so directly? Upregulation of ABCA1 was not observed in either E-cadherin or α-catenin KO cells (Fig. 1I), indicating that ABCA1 induction is driven as a specific response to a fundamental change in cellular state induced by Snail overexpression, not an ancillary effect of disrupting either cell-cell adhesion or apical-basal polarity. Generally, ABCA1 is a transporter for the efflux of cholesterol out of the cell when there is an excess of cholesterol in the cell and ABCA1 upregulation in response to excess loading of cholesterol is mediated by sterol-activated nuclear receptors, LXRs (Ignatova *et al*, 2013). Aside from the cholesterol-LXRs regulatory axis, it was recently reported that ABCA1 expression in epithelial cells is regulated via FOXO3a and c-Myc (Frechin *et al*, 2015). Activation of FAK/PI3K/Akt signaling depending on cell crowding state downregulates ABCA1 expression via phosphorylation of FOXO3a, a positive transcriptional regulator of ABCA1 (Frechin *et al.*, 2015). We found that phosphorylation level of FOXO3a at S453 is lower in EpH4-Snail cells than in wild-type EpH4 cells (Fig. S3A). However, since downregulation of Akt signaling by treatment with inhibitors of its activators GSK2334470 (PDK1), LY294002 (PI3K) and Wortmannin (PI3K) did not induce ABCA1 expression in wild-

type EpH4 cells (Fig. S3A), FOXO3a would not be responsible for ABCA1 expression

in EpH4-Snail cells. Although ABCA1 expression is also downregulated by c-Myc in

breast cancer, no remarkable difference of c-Myc expression was observed between

EpH4 and EpH4-Snail cells, indicating that c-Myc is not likely involved in ABCA1

expression in EpH4-Snail cells (Fig. S3B). Therefore, we next pursued in detail the

possibility that LXRs induce ABCA1 in EpH4-Snail cells. Immunofluorescence

microscopy revealed that LXRs are highly accumulated in the nuclei of EpH4-Snail

cells, suggesting that ABCA1 upregulation could be mediated by LXRs (Fig. 2A).

Consistently, treatment with an LXR inhibitor GSK2033 abolished ABCA1 expression

to an undetectable level in EpH4-Snail cells (Fig. 2B). Since transcriptional activity of

LXRs are regulated by cellular cholesterol level, we examined whether ABCA1 is

expressed in response to high cholesterol accumulation in the cells. Inhibition of

cholesterol biosynthesis by simvastatin, an inhibitor of the rate-limiting enzyme of

cholesterol biosynthesis 3-hydroxy-3-methylglutaryl-coenzyme A reductase,

remarkably decreased ABCA1 expression in EpH4-Snail cells but residual ABCA1 was

still detected (Fig. 2C, normal FBS). In contrast, treatment with simvastatin in the

medium supplemented with lipid-free FBS completely abolished ABCA1 expression

(Fig. 2C, lipid-free FBS). Together with the finding that incubation in lipid-free FBS

medium alone did not markedly decrease ABCA1 expression, these results suggest that

cholesterol biosynthesis is the primary source of excess accumulation of cholesterol in

EpH4-Snail cells, but cholesterol uptake from media also significantly contribute to the

accumulation as a secondary source.

These findings imply dysregulation of cholesterol metabolism in EpH4-Snail.

Filipin staining revealed that cholesterol was prominently accumulated in intracellular

structures in EpH4-Snail cells, whereas it localized primarily in the plasma membrane

of wild-type EpH4 cells (Fig. 2D). Since excess cholesterol can be stored in lipid

droplets, we stained cells with Lipi-Green to visualize lipid droplets. We found that

lipid droplets were significantly enlarged in EpH4-Snail (Figs. 2, E and F).   We next

compared the total cholesterol levels in EpH4 cells and EpH4-Snail cells; the cellular

amount of cholesterol slightly increased in EpH4-Snail cells, although the difference

was not significant (Fig. 2G). These results suggest that the accumulation of cholesterol

in lipid droplets in EpH4-Snail cells was not due to an increase in the amount of

cholesterol, but to an abnormality in the subcellular distribution of cholesterol.

Collectively, dysregulation of intracellular distribution of cholesterol activates LXR to

induce ABCA1 expression in hybrid E/M cells.

Downregulation of VLCFA-SM synthesis induces Chol/SM imbalance in

hybrid E/M cells

A large proportion (30-50%) of cellular cholesterol is associated with

sphingomyelin (Das *et al*, 2014). Therefore, we next examined the possibility that

sphingomyelin levels are altered in EpH4-Snail cells. We found that the sphingomyelin

content relative to total phospholipids decreased in EpH4-Snail cells (Fig. 3A). To

investigate the SM profile in detail, we next analyzed the fatty acid composition of SM

using LC-MS. In EpH4-Snail cells, the fractions of VLCFA-SMs (22:1, 22:0 and 24:1)

were significantly decreased whereas those of LCFA-SMs (14:0, 16:1 and 16:0) were

rather increased (Fig. 3B), resulting in a prominent decrease in the total VLCFA-SM

fraction (Figs. 3, C and D). These results suggest that decrease in VLCFA-SMs are

responsible for decrease in total SM content in EpH4-Snail cells.

Fatty acid composition of SM is mainly modulated by substrate specificity of

two families of enzymes, Elovls and CerSs, that respectively catalyze the rate-limiting

step of the fatty acid elongation cycle and the N-acylation of sphingoid bases to produce

ceramides (Fig. 3E) (Levy & Futerman, 2010; Sassa & Kihara, 2014). Therefore, we

hypothesized that transcriptional repression of genes from these families causes the

decrease in VLCFA-SM content in EpH4-Snail cells. RT-PCR analyses identified

ELOVL7 and CERS3 as genes which are completely repressed specifically in EpH4-

Snail cells (Fig. 3F). Consistent with our hypothesis, Elovl7 and CerS3 control VLCFA-

SM production (Fig. 3E). Collectively, exogenous expression of Snail represses

transcription of VLCFA-SM-producing enzymes and results in a decrease in the amount

of VLCFA-SM. Consequently, the ratio of cholesterol to SM (Chol/SM ratio) in the

cells significantly increased in EpH4-Snail cells (Fig. 3H), suggesting that a substantial

amount of cholesterol exist in a sphingomyelin-unbound form in EpH4-Snail.

Since ABCA1 expression is not aligned with either the loss of epithelial state (E-

cadherin and α-catenin KO) or a non-epithelial basal state (fibroblasts), upregulation of

ABCA1 is characteristic to the hybrid E/M cells expressing ectopic Snail (Fig. 3I). To

investigate the mechanism underlying this difference in ABCA1 expression, we again

compared Chol/SM ratio in EpH4-Snail and normal fibroblast cell lines L-929 and

NIH3T3. Chol/SM ratio of these fibroblasts were slightly but not significantly higher

than that of EpH4 (Fig. 3I). The cholesterol content in NIH-3T3 cells is significantly

higher than in EpH4 cells (Fig. S2). Consequently, normal fibroblast, would synthesize

more SMs to counteract the cholesterol load imposed by the decrease in VLCFA-SMs

and stabilize the Chol/SM ratio. By contrast, hybrid E/M cells are unable to respond

appropriately to the harmful imbalance between cholesterol and SM due to the

incomplete activation of the mesenchymal transcriptome.

To examine whether the above findings apply to human cancer cells, we used

two esophageal squamous cell carcinoma cell lines, TE-15 and TE-8, that exhibit

epithelial and hybrid-E/M phenotype, respectively (Fig. 3J) (Nishihira *et al*, 1993). As

expected, expressions of ABCA1 and Snail were cooperatively upregulated in TE-8

cells (Fig. 3J). Consistent with our findings in the EpH4-Snail system, Chol/SM ratio

was significantly higher in TE-8 cells than in TE-15 cells (Fig. 3K). Together with the

results of mouse cell lines, normal epithelial and fibroblast cells maintain Chol/SM ratio

around 1.0-1.5, and ABCA1 expression as a response to the excess Chol is induced

when Chol/SM ratio exceed 1.5 (Fig. 3H). Notably, the difference in either cholesterol

or SM contents between TE-15 and TE-8 cells, and cholesterol content between EpH4

and EpH4-Snail is not significant when each are compared alone (Fig. 2D and Fig. S4),

indicating the importance of considering their functional cooperativity, i.e. to analyze

the Chol/SM ratio.

Inhibitor of cholesterol esterification selectively inhibits growth of Snail-positive kidney cancer cells

Finally, we examined the possibility of targeting the imbalance of Chol/SM ratio

in Snail-positive cancer cells to selectively inhibit their growth. As discussed above,

hybrid E/M cells of high Chol/SM ratio must either export cholesterol out of cells or

isolate it within cells in lipid droplets as cholesteryl esters to avoid the cytotoxic effects

of free cholesterol. Thus, inhibiting the means of clearing excess cholesterol could

specifically target cells in the hybrid E/M state for self-extermination. Cholesterol efflux

from cells is mainly dependent on ABCA1, and esterification of cholesterol is catalyzed

by acyl-CoA:cholesterol acyl transferases (ACATs) (Fig. 4A) (Song *et al*, 2021).

According to our findings above, hybrid E/M cells activate both cholesterol efflux and

isolation system to handle excess cholesterol. Therefore, we examined the effect of

ABCA1 inhibitor CsA and ACAT inhibitor TMP-153 on the EpH4-Snail model and

renal carcinoma cells. While CsA non-selectively inhibited growth of all cell lines

tested (Figs. 4B and 4D), TMP-153 selectively inhibited growth of Snail-positive cells

(EpH4-Snail, 786-O, A498 and Caki-1; Figs. 4C and 4E). Thus, sequestration of

cholesterol via esterification by ACATs is the primary mechanism for handling excess

cholesterol to avoid cell death in these cells. Notably, TMP-153 treatment alone did not

provide this selectivity between esophageal cancer cell lines, but combination of TMP-

153 and cyclosporin A selectively inhibited growth of the Snail-positive TE-8 cells

(Figs. S5, A-C). Collectively, these results suggest that inhibition of cholesterol efflux

and cholesterol esterification selectively inhibit growth of cells stably in a Snail-induced

hybrid E/M state.

We also investigated the potential of ACAT inhibitors as anti-tumor agents in

vivo using xenograft models. We successfully transplanted the 786-O line into nude

mice among Snail-positive ccRCC cell lines. Using this system, we found that TMP-

153 treatment significantly reduced 786-O tumor volume in vivo (Figs. 4, F-H).

Moreover, the combination of TMP-153 and cyclosporin A also inhibited the growth of

TE-8 tumor xenografts (Figs. S5D-F). Together with the in vitro assays (Figs. 4, B-J

and Figs. 5, A-C), we propose that cholesterol clearance systems, particularly ACATs,

can be used as molecular targets to selectively inhibit the growth of Snail-positive

cancers.

Discussion

In recent years, it has been revealed that the expression of EMT-TFs, such as

Snail, induces cancer cells to acquire mesenchymal characteristics while retaining

epithelial traits, transforming them into so-called hybrid E/M cells. Hybrid E/M cells

have been demonstrated to exhibit the highest plasticity and malignancy (Aiello *et al*,

2018; Pastushenko *et al*, 2018). Moreover, Snail expression contributes to cancer

stemness and resistance to chemotherapy (Lambert & Weinberg, 2021). To date,

molecular mechanisms connecting to EMT-TFs to chemoresistance remain to be

established. Here we identify lipid metabolic imbalance-mediated ABCA1 expression

as a mechanism of chemoresistance induced by Snail. We found reduced VLCFA-SM

biosynthesis in Snail-positive cells with hybrid E/M phenotypes, resulting in high

Chol/SM ratio. This should be detrimental in theory since it increases exposure to the

toxic effect of free cholesterol but cancer cells deploy countermeasures: upregulating

ABCA1 to move cholesterol out of cells and sequestering esterified cholesterol in the

form of lipid droplets. Notably, it has been also reported that not only ABCA1 (Iwasaki

et al., 2010; Wang *et al*, 2021) but also enhanced lipid droplet formation triggers

chemoresistance to various drugs (Cotte *et al*, 2018; Hultsch *et al*, 2018; Schlaepfer *et*

al, 2012). Therefore, dietary or pharmacological control of cholesterol and also

sphingolipids would be a new strategy to overcome resistance to chemotherapy.

The aberrant activation of EMT-TFs is implicated in multiple crucial steps in

cancer progression, prompting efforts to explore therapeutic approaches aimed at

suppressing EMT-TF functions in cancer cells (Davis *et al.*, 2014). However, as EMT-

TFs are predominantly located in the nucleus, no effective treatment methods, including

antibody drugs, have been established thus far. Therefore, instead of directly targeting

EMT-TFs, focusing on the metabolic network vulnerabilities specific to hybrid E/M

cells may prove to be an effective therapeutic strategy. In the present study, we found

increased sensitivity of Snail-positive cells to toxicity of free-cholesterol accumulation

induced by the ACAT inhibitor TMP-153. Notably, kidney cancer growth could be

suppressed by inhibiting cholesterol esterification alone while esophageal cancer

required the additional downregulation of ABCA1, indicating varying dependence on

free cholesterol clearance strategies among cancers of different primary origin. Other

cancers might employ alternative pathways to evade cytotoxic effects of excess

cholesterol and identifying these traits will pave the way for the development of novel

strategies in targeted molecular therapeutics against hybrid E/M cancers. An intriguing

corollary here is the potential utility of ABCA1 expression as a diagnostic marker to

identify cancers that would be sensitive to targeting the Chol/SM imbalance.

Although various metabolic changes in cancer cells focusing on energy sources

including sugars, amino acids and fatty acids have been intensely studied (Pavlova *et al*,

2022), alteration of sphingolipid homeostasis driven by EMT-TFs remains largely

unknown. We found both quantitative and qualitative alteration of sphingomyelin

profile: decrease in the total SM amount and the VLCFA-SM percentage. This

alteration is attributed to transcriptional repression of enzymes for biosynthesis of

VLCFA-SMs, CerS3 and Elovl7. Genes whose transcription is repressed by Snail

include those encoding proteins directly involved in epithelial adhesive structures, such as E-cadherin and Claudins. Hence, it is not surprising that VLCFA-SMs play important roles cooperatively with the adhesion molecules. Indeed, cholesterol accumulation is required for tight junction formation, and VLCFA-SMs are enriched in the junctional region (Shigetomi *et al*, 2018; Shigetomi *et al*, 2023). Studies focusing on the VLCFA-synthetic enzymes would provide a new layer of knowledge about mechanisms underlying the establishment of epithelial structures and functions. Such studies would also contribute to understanding how dysfunctions in lipid metabolisms promote cancer malignancy and tissue fibrosis as a consequence of EMT. The regulation of cell adhesion by fine-tuning of lipid composition should be reversible and therefore would explain the two conflicting events: the invasion and metastasis process and the viability process at the metastatic site.

Materials and methods

Cells and reagents

Cells were grown in DMEM supplemented with 10% FBS under 5% CO_2 at 37°C. Mouse Snail-expressing EpH4 cells (EpH4-Snail)(Ikenouchi *et al.*, 2003), E-cadherin KO and α-catenin KO EpH4 cells (Shigetomi *et al.*, 2018) were generated previously. EpH4-Snail cells were maintained in medium supplemented with 500 μg/ml G418 (Wako Pure Chemical). For KRAS G12V expression in MDCK II cells, pLVET-HA-K-RasG12V-IRES-GFP (Addgene Plasmid #107140) was transduced using the lentiviral system, and GFP-positive cells were collected using a cell sorter Sony SH-800. Nitidine chloride was purchased from Selleck Biotech, cyclosporin A from Wako, and TMP-153

from Cayman Chemical, filipin complex from *Streptomyces filipinensis* from Sigma-

Aldrich, Lipi-Green from Dojindo Laboratories.

The following primary antibodies were used for immunoblotting,

immunofluorescence microscopy and immunohistochemistry: Rabbit anti-ABCA1 mAb

(96292S, IB) and rabbit anti-Snail mAb (3879S, IB) from Cell Signaling Technology;

Rabbit anti-ABCA1 pAb (NB400-105, IHC) from Novus Biologicals; Rat anti-E-

cadherin mAb (ECCD-2, IB) from TaKaRa; Goat anti-LXR alpha + LXR beta pAb

(ab24362, IF) from Abcam; Mouse anti-α-tubulin (12G10, IB) produced in house.

The following secondary antibodies were used: HRP-conjugated anti-rat IgG

(HAF005) and anti-mouse IgG (HAF007) from R&D Systems; HRP-conjugated anti-

rabbit IgG (4030-05) from Southern Biotech; Cy3-conjugated anti-goat IgG (705-165-

147) from Jackson Immunoresearch Laboratories.

RT-PCR

Total RNAs were prepared with RNeasy Mini Kit (Qiagen) and the

retrotranscribed with Superscript III First-Strand Synthesis System (Invitrogen). RT-

PCR was then performed using KOD FX (TOYOBO) following manufacturer's

instructions. Primer sequences used are shown in Table S1.

Lipid analysis

Total lipids were extracted from cultured cells by Bligh-Dyer's method, dissolved

in methanol/chloroform = 1/1 (v/v). Phospholipid content was determined by phosphate

assay. SM and Chol content were determined using Sphingomyelin assay kit STA-601

(Cell Biolabs) and Amplex Red Cholesterol Assay Kit (Invitrogen), respectively,

according to the manufacturer's instructions. Fatty acid profile of SM was determined

with LC-MS as described previously[32]. Briefly, electrospray ionization mass

spectrometry analysis was performed on a 6420 triple-quadrupole liquid

chromatography–mass spectrometer (Agilent Technologies) equipped with an HPLC

system and an auto sampler (Infinity 1260; Agilent Technologies). The extracted lipids

were directly subjected to electrospray ionization mass spectrometry analysis. The

mobile phase composition was acetonitrile/methanol/water = 18:11:1 (0.1% ammonium

formate). The flow rate was 4 μl/min. The mass range of the instrument was set at 650–

950 *m/z*. *m/z* profiles of SM were extracted according to total ion chromatogram

patterns, and all peaks corresponding to $[M+H]^+$ ions of d18:1-even chain length fatty

acid-SMs were subjected to analyses.

Fluorescence microscopy

To visualize subcellular localization of cholesterol, cells cultured in glass-bottom

dishes were fixed with 4% paraformaldehyde for 15 min at room temperature and

stained with 50 μg/ml filipin prepared in PBS for 30 min. To visualize lipid droplets,

live cells were washed once with HBSS and stained with Lipi-Green (Dojindo) for 20

min at 37°C. Cells were washed twice with HBSS and observed on a heating stage at

37°C. For immunofluorescence microscopy, cells cultured on coverslips were fixed

with 4% paraformaldehyde prepared in PBS for 15 min at RT and permeabilized with

0.1% Triton X-100 prepared in PBS. Fixed cells were blocked with 1% BSA prepared

in PBS for 1 h at RT. Cells were incubated with primary antibodies for 1 h at RT and

secondary antibodies for 1 h at RT. Antibodies were prepared in the blocking solution.

Fixed cells were observed at RT. All observation was performed with a confocal

microscope (Carl Zeiss LSM900) equipped with Plan-APO (63×/1.40 NA, oil

immersion) objective. Images were acquired using Carl Zeiss Zen 3.4 software. Images

were analyzed using ImageJ/Fiji.

Immunoblotting

Whole cells were lysed with SDS sample buffer and samples were resolved by

SDS-PAGE and transferred to nitrocellulose membranes. After blocking with 5% skim-

milk prepared in TBS-T, membranes were incubated with primary antibody for 1 h at

RT or overnight at 4°C. Membranes were then washed and incubated with HRP-

conjugated secondary antibody for 1 h at RT. Chemiluminescence signal was detected

using a LAS4000mini imaging system (Fujifilm) and images were analyzed using

ImageJ/Fiji.

Growth inhibition assay

Cells cultured in 96-well microtiter plates (2×10^3 cells/well, 48 h) were subjected

to treatments with nitidine chloride, cyclosporin A, TMP-153 or their combination

dissolved in DMSO for 24 h at 37°C. Final DMSO concentrations were less than 0.25%.

After treatment, metabolically active cell number was determined using Cell Counting

Kit-8 (Dojindo) according to manufacturer's instructions. Briefly, 10 μl of CCK-8

solution was directly added to each well containing 100 μl culture medium and cells

were incubated 1 h at 37°C. Reaction was stopped by adding 10 μl of 1% sodium

dodecyl sulfate prepared in MQ and plates were put at 4°C to avoid additional reaction.

Absorbance of solution at 450 nm were measured using a spectrophotometer (Thermo

GENESYS 10S UV-Vis). Data were expressed as relative to 0.25% DMSO control.

Immunohistochemistry

Formalin-fixed and paraffin-embedded surgical specimens were cut into 3 μm

sections. These sections were immersed in antigen retrieval solution (pH 9.0) (Nichirei

Bioscience Inc.) and heated in a pressure chamber. After that, the sections were

incubated with the primary antibodies (rabbit anti-ABCA1 pAb, NB400-105) and later

incubated with HRP-labeled goat anti-rabbit secondary antibodies (Nichirei Bioscience

Inc.). Immunoreactivity was visualized by using 3–3 Diaminobenzidine Dab Substrate

Kits (Nichirei Bioscience Inc.). Informed consent for experimental use of the samples

was obtained from the patients in accordance with the hospital's ethical guidelines.

Tumor xenograft model

Four-week-old female athymic nude mice were purchased from Japan SLC and

housed in a specific pathogen-free facility. The animal experiments were approved by

the ethics committee of Aichi Medical University and performed according to the

established guidelines.

To initiate xenografting of 786-O, 2×10^6 cells in 200 μL DMEM were

subcutaneously injected into the back of mice by using a 27 G needle. After 7 days,

when the inoculated tumor became visible, the mice were randomly divided into

treatment and control groups, and treatment was initiated (day 0). TMP-153 (50 mg/kg

body weight) or vehicle (0.5% hydroxypropyl methylcellulose with 0.1% polysorbate

80) was intraperitoneally administered to each mouse on days 0, 4, 7, 11, 14, and 18 to

each mouse. The tumor volume was measured every week and calculated using the

modified ellipsoid formula ($1/2 \times$ length $\times$ width^2). The xenografting of TE-8 was

performed with some modifications. For this, 1×10^7 cells were injected. A combination

of TMP-153 and cyclosporin A (both at 10 mg/kg body weight) or vehicle was

intraperitoneally administered to each mouse on days 0 and 4. Finally, the tumor

volume was measured 36 days after the first administration.

Statistical analysis

Microsoft Excel for Microsoft 365 MSO 2306 and GraphPad Prism v8.4.1 were

used for analyses and displays of quantitative data. Data are expressed as mean with

each point or error bars representing standard deviations (SD). Unpaired two-tailed

Student's *t*-test, Dunnett's multiple comparison test and Welch's *t*-test are performed

using Prism. Unless otherwise noted, *n* denotes biological replicates.

Acknowledgments

We thank all members of the Ikenouchi laboratory for helpful discussions, Takana

Motoyoshi (K.I. Stainer) for immunohistochemistry, Teruaki Fujishita and Masahiro

Aoki for technical advice on xenograft experiments, Minako Suzuki for technical

assistance and Kenji Kasai for pathological advice. This work was supported by

AMED-FORCE (21444781) (J.I.), JSPS KAKENHI [JP22H02618 (J.I.), JP23K18141

(J.I.) and 22KJ2374 (A.M.)], JST-FOREST (JPMJFR204L) (J.I.), and the Bioscience

Research Grant from Takeda Science Foundation (J.I.).

Author contributions

A.M. performed most of experiments and analyzed the data. A.I. designed and

performed in vivo experiments with clinical specimens and tumor xenografts. W.H.

made pathological diagnoses and provided clinical specimens. T.K. performed surgery

and managed patient information. K.O. performed pathological analysis. A.M. and J.I.

designed research and wrote the paper.

Declaration of interests

There are no competing interests to declare.

References

Aiello NM, Maddipati R, Norgard RJ, Balli D, Li J, Yuan S, Yamazoe T, Black T, Sahmoud A, Furth EE *et al* (2018) EMT Subtype Influences Epithelial Plasticity and Mode of Cell Migration. *Dev Cell* 45: 681-695.e684

Arner EN, Du W, Brekken RA (2019) Behind the Wheel of Epithelial Plasticity in KRAS-Driven Cancers. *Frontiers in oncology* 9: 1049

Batlle E, Sancho E, Francí C, Domínguez D, Monfar M, Baulida J, García De Herreros A (2000) The transcription factor snail is a repressor of E-cadherin gene expression in epithelial tumour cells. *Nat Cell Biol* 2: 84-89

Cano A, Pérez-Moreno MA, Rodrigo I, Locascio A, Blanco MJ, del Barrio MG, Portillo F, Nieto MA (2000) The transcription factor snail controls epithelial-mesenchymal transitions by repressing E-cadherin expression. *Nat Cell Biol* 2: 76-83

Cotte AK, Aires V, Fredon M, Limagne E, Derangère V, Thibaudin M, Humblin E, Scagliarini A, de Barros JP, Hillon P *et al* (2018) Lysophosphatidylcholine acyltransferase 2-mediated lipid droplet production supports colorectal cancer chemoresistance. *Nat Commun* 9: 322

Cui Y, Wu L, Cao R, Xu H, Xia J, Wang ZP, Ma J (2020) Antitumor functions and mechanisms of nitidine chloride in human cancers. *Journal of Cancer* 11: 1250-1256

Das A, Brown MS, Anderson DD, Goldstein JL, Radhakrishnan A (2014) Three pools of plasma membrane cholesterol and their relation to cholesterol homeostasis. *Elife* 3

Davis FM, Stewart TA, Thompson EW, Monteith GR (2014) Targeting EMT in cancer: opportunities for pharmacological intervention. *Trends in pharmacological sciences* 35: 479-488

Debaugnies M, Rodríguez-Acebes S, Blondeau J, Parent MA, Zocco M, Song Y, de Maertelaer V, Moers V, Latil M, Dubois C *et al* (2023) RHOJ controls EMT-associated resistance to chemotherapy. *Nature* 616: 168-175

Fang Z, Tang Y, Jiao W, Xing Z, Guo Z, Wang W, Xu Z, Liu Z (2014) Nitidine chloride induces apoptosis and inhibits tumor cell proliferation via suppressing ERK signaling pathway in renal cancer. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association* 66: 210-216

Fischer KR, Durrans A, Lee S, Sheng J, Li F, Wong ST, Choi H, El Rayes T, Ryu S, Troeger J *et al* (2015) Epithelial-to-mesenchymal transition is not required for lung metastasis but contributes to chemoresistance. *Nature* 527: 472-476

Frechin M, Stoeger T, Daetwyler S, Gehin C, Battich N, Damm EM, Stergiou L, Riezman H, Pelkmans L (2015) Cell-intrinsic adaptation of lipid composition to local crowding drives social behaviour. *Nature* 523: 88-91

Hultsch S, Kankainen M, Paavolainen L, Kovanen RM, Ikonen E, Kangaspeska S, Pietiäinen V, Kallioniemi O (2018) Association of tamoxifen resistance and lipid reprogramming in breast cancer. *BMC cancer* 18: 850

Ignatova ID, Angdisen J, Moran E, Schulman IG (2013) Differential regulation of gene expression by LXRs in response to macrophage cholesterol loading. *Molecular endocrinology (Baltimore, Md)* 27: 1036-1047

Ikenouchi J, Matsuda M, Furuse M, Tsukita S (2003) Regulation of tight junctions during the epithelium-mesenchyme transition: direct repression of the gene expression of claudins/occludin by Snail. *J Cell Sci* 116: 1959-1967

Iwasaki H, Okabe T, Takara K, Yoshida Y, Hanashiro K, Oku H (2010) Down-regulation of lipids transporter ABCA1 increases the cytotoxicity of nitidine. *Cancer chemotherapy and pharmacology* 66: 953-959

Lambert AW, Weinberg RA (2021) Linking EMT programmes to normal and neoplastic epithelial stem cells. *Nat Rev Cancer* 21: 325-338

Levy M, Futerman AH (2010) Mammalian ceramide synthases. *IUBMB life* 62: 347-356

Makhov P, Joshi S, Ghatalia P, Kutikov A, Uzzo RG, Kolenko VM (2018) Resistance to Systemic Therapies in Clear Cell Renal Cell Carcinoma: Mechanisms and Management Strategies. *Molecular cancer therapeutics* 17: 1355-1364

Nieto MA, Huang RY, Jackson RA, Thiery JP (2016) EMT: 2016. *Cell* 166: 21-45

Nishihira T, Hashimoto Y, Katayama M, Mori S, Kuroki T (1993) Molecular and

cellular features of esophageal cancer cells. *Journal of cancer research and clinical*

oncology 119: 441-449

Padmanaban V, Krol I, Suhail Y, Szczerba BM, Aceto N, Bader JS, Ewald AJ (2019) E-

cadherin is required for metastasis in multiple models of breast cancer. *Nature* 573:

439-444

Pastushenko I, Brisebarre A, Sifrim A, Fioramonti M, Revenco T, Boumahdi S, Van

Keymeulen A, Brown D, Moers V, Lemaire S *et al* (2018) Identification of the tumour

transition states occurring during EMT. *Nature* 556: 463-468

Pavlova NN, Zhu J, Thompson CB (2022) The hallmarks of cancer metabolism: Still

emerging. *Cell metabolism* 34: 355-377

Sassa T, Kihara A (2014) Metabolism of very long-chain Fatty acids: genes and

pathophysiology. *Biomolecules & therapeutics* 22: 83-92

Schlaepfer IR, Hitz CA, Gijón MA, Bergman BC, Eckel RH, Jacobsen BM (2012)

Progestin modulates the lipid profile and sensitivity of breast cancer cells to docetaxel.

Molecular and cellular endocrinology 363: 111-121

Shigetomi K, Ono Y, Inai T, Ikenouchi J (2018) Adherens junctions influence tight

junction formation via changes in membrane lipid composition. *J Cell Biol* 217: 2373-

2381

Shigetomi K, Ono Y, Matsuzawa K, Ikenouchi J (2023) Cholesterol-rich domain

formation mediated by ZO proteins is essential for tight junction formation. *Proc Natl*

Acad Sci U S A 120: e2217561120

Song Y, Liu J, Zhao K, Gao L, Zhao J (2021) Cholesterol-induced toxicity: An

integrated view of the role of cholesterol in multiple diseases. *Cell metabolism* 33:

1911-1925

Wang W, Lokman NA, Noye TM, Macpherson AM, Oehler MK, Ricciardelli C (2021)

ABCA1 is associated with the development of acquired chemotherapy resistance and

predicts poor ovarian cancer outcome. *Cancer drug resistance (Alhambra, Calif)* 4:

485-502

Yang H, Chen K, Zhang X, Wang L, Li C, Tao H, Wang L, Li Q (2010) Vitamin A

deficiency results in dysregulation of lipid efflux pathway in rat kidney. *Pediatric*

nephrology (Berlin, Germany) 25: 1435-1444

Zheng X, Carstens JL, Kim J, Scheible M, Kaye J, Sugimoto H, Wu CC, LeBleu VS,

Kalluri R (2015) Epithelial-to-mesenchymal transition is dispensable for metastasis but

induces chemoresistance in pancreatic cancer. *Nature* 527: 525-530

Figure legends

Figure 1

Ectopic expression of Snail confers epithelial cells nitidine chloride

resistance via induction of ABCA1 expression

(**A**) Enhancement of growth inhibitory effect of nitidine chloride (NC) on human renal

carcinoma cells by co-treatment with an ABCA1 inhibitor cyclosporin A (CsA). Cells

were grown with or without 20 μM NC and 10 μM CsA for 24 h, and cell number was

determined with CCK-8. Fold change in cell number by NC was shown as relative to

controls without NC. Results are shown as mean of at least four independent

experiments ± SD (Tukey-Kramer's multiple comparison test: **, $p<0.01$; ***,

$p<0.001$; ****, $p<0.0001$).

(**B**) Immunoblot analysis of whole cell lysate of human renal carcinoma cell lines.

(**C**) Comparison of ABCA1 expression between normal tissue and primary tumor of

indicated subtypes of renal cancers analyzed using UCSC Xena. Bars indicate means

(Welch's t-test: ***, $p<0.001$; ****, $p<0.0001$).

(**D**) Immunohistochemistry of a surgically extracted renal tissue from a patient with

Fuhrman grade 3 primary ccRCC, indicating upregulation of ABCA1 in the lesion site.

The ABCA1 staining was verified by the accumulation of known signals in renal

tubules and glomeruli in normal tissue (Yang *et al*, 2010). Scale bars, 250 μm (left), 50

μm (right).

(**E**) Phase contrast images of EpH4 wild-type cells (top) and Snail-overexpression cells

(bottom, EpH4-Snail). Scale bar, 50 μm.

(**F**) Immunofluorescence images of EpH4 and EpH4-Snail cells. Scale bars, 20 μm.

(**G**) Acquisition of NC resistance by exogenous expression of Snail in EpH4 cells and

enhancement of NC effect on EpH4-Snail cells by co-treatment with CsA. Shown are

means of three independent experiments ± SD (Tukey-Kramer's multiple comparison

test: *, $p<0.05$; ****, $p<0.0001$).

(**H-J**) Immunoblot analyses of whole cell lysates of (H) EpH4, EpH4-Snail, E-cadherin

KO and α-catenin KO EpH4 cells, (I) MDCK II cells expressing KRAS G12V treated

with 5 μg/ml TGFβ for 0, 3 and 8 days and (J) human esophageal carcinoma cell lines

TE-15 and TE-8. Phase contrast images are also shown on top (J). Scale bars, 50 μm.

Figure 2

Alteration of cellular cholesterol distribution induces ABCA1 expression

in hybrid E/M cells

(**A**) Immunofluorescence images of cells stained with the LXRα+β antibody. Scale

bars, 10 μm.

(**B**) Immunoblot analyses of whole cell lysates of EpH4-Snail cells treated with 1 μM

GSK-2033 for 24 h.

(**C**) Immunoblot analyses of whole cell lysates of EpH4-Snail cells treated with

indicated concentration of simvastatin or 0.1% DMSO ('0' μM) in DMEM

supplemented with 10% normal or lipid-free fetal bovine serum (FBS) for 48 h.

(**D**) Subcellular distribution of Chol stained with filipin. Scale bar, 10 μm.

(**E**) Live cell images of lipid droplets (LDs) visualized by Lipi-Green dye.

(**F**) Quantification of LD signal per cell. The Lipi-Green signals of all optical sections

were summed and divided by the number of cells counted using DAPI images. Shown

are means of three different fields of view ±SD.

(**G**) Comparison of cholesterol (Chol) content relative to phospholipid concentration.

Shown are means of three biological replicates ± SDs. (F,G) *p*-value of Student's *t*-test

are shown.

Figure 3

Effects of Snail-induced EMT on sphingomyelin profile

(**A**) Comparison of sphingomyelin (SM) content relative to phospholipid content.

(**B**) Comparison of cholesterol (Chol)/SM ratio between EpH4 and EpH4-Snail cells.

(**C**) Fatty acid composition of SM determined using LC-MS. All peaks corresponding to

d18:1-even fatty acid-SMs were included in the analysis and each peak value is

expressed as a percentage. n.d., peak not detected.

(**D**) Pie charts of SM chain length profile from c. SMs were classified into long-chain

fatty acid (LCFA, orange) (≤20) and very long-chain fatty acid (VLCFA, purple) (>20)

SMs.

(**E**) Chain length specificity of Elovls and CerSs responsible for fatty acid profile of

sphingolipids reported (Levy & Futerman, 2010; Sassa & Kihara, 2014). Elovls catalyze

a rate-limiting step of fatty acid elongation cycle, and CerSs catalyze *N*-acylation of

sphingoid bases to produce ceramides.

(**F**) Comparison of transcription levels of *ELOVL*s and *CERS*s. Notable decreases in

expressions of *ELOVL7* and *CERS3* in EpH4-Snail cells are indicated by asterisks.

(**G**) Immunoblot analyses of whole cell lysates of mouse epithelial (MTD-1A, CSG)

and fibroblast (L-929, NIH3T3) cells.

(**H**) Comparison of Chol/SM ratio of EpH4, EpH4-Snail and normal fibroblasts.

(**I**) Comparison of Chol/SM ratio between TE-15 and TE-8.

(**J**) Scatter plot of Chol content against SM content used in B, H and I. Dashed lines

indicate Chol/SM = 1.0 (blue) and 1.5 (red). The region of Chol/SM > 1.5 is shown with

red background.

(**K**) Possible mechanism of LD enlargement induction through decrease in SM. In

EpH4, a certain amount of Chol is sequestered through interaction with SM. In EpH4-

Snail, SM-unbound form of Chol is increased, detected by cellular cholesterol sensors,

internalized and stored in LDs. In A, B, C, H and I, data are means of three independent

experiments ± SDs (Student's *t*-test (A,B,C,I) or Dunnett's test (H): n.s., $p \geq 0.05$; *, *p*

< 0.05; **, $p < 0.01$; ***, $p < 0.001$).

Figure 4

Selective growth inhibition of Snail-positive cells by an ACAT inhibitor

TMP-153

(**A**) Schematic image of molecular mechanisms for handling excess cholesterol to avoid

cell death. Cells eliminate excess cholesterol through efflux mediated by transporters

including ABCA1 or isolation into lipid droplets (LD) via esterification of cholesterol

by acyl-CoA: cholesterol acyltransferases (ACATs) to avoid cell death. Cyclosporin A

(CsA) and TMP-153 inhibits ABCA1 and ACATs, respectively.

(**B-E**)**,** Effects of treatments with cyclosporin A (B,D) or TMP-153 (C,E) to cellular

growth of EpH4 and EpH4-Snail (B,C) or human kidney cancer cell lines (D,E). Cells

were treated with drugs for 24 h and relative cell number to 0.1% DMSO control was

determined by CCK-8. Results are shown as mean of at least three independent

experiments ± SD.

(**F-H**) TMP-153 inhibits the growth of tumor xenografts composed of Snail-positive

renal cancer 786-O cells. Experiments were performed as described in Materials and

Methods. In brief, 50 mg/kg TMP-153 was administered intraperitoneally to nude mice

bearing 786-O tumor xenografts. After 21 days of the first administration, the overall

appearance (F) and the excised tumors (G) of control and treated mice were compared.

(H) Tumor growth curve in the xenograft model. Results are shown as mean of four

biological replicates ± SD (Student's t-test: *, $p < 0.05$; **, $p < 0.01$).

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary Figure 1

Cancer types that exhibit tumor specific upregulation of ABCA1 transcription

(**A-C**) Comparison of mRNA transcription of ABCA1 (A,B) and Snail (C) in different cancer types.

Expression data from GDC TCGA database were analyzed using USCS Xena platform. Significant

differences (***, $p < 0.001$) by Welch's t-tests are shown.

Supplementary Figure 2

Comparison of ABCA1 expression in different progression grades of kidney cancer cases

Immunohistochemistry of surgically extracted renal tissues from patients with Fuhrman grade 13 primary ccRCC, indicating elevation of ABCA1 signal at the lesion site of higher-grade ccRCC. All samples were collected from independent patients. The ABCA1 staining was verified by the accumulation of known signals in renal tubules and glomeruli in normal tissues (32). Scale bars, 100 μm.

Supplementary Figure 3

Contribution of transcription factors involved in ABCA1 transcription in epithelial cells

(**A**) Immunoblot analysis of whole cell lysates of EpH4-Snail (without treatment) and EpH4 cells treated with inhibitors of PI3K-Akt pathway (5 μM GSK2334470: PDK1, 50 μM LY294002: PI3K, or 1 μM Wortmannin: PI3K for 24 h).

(**B**) Immunoblot analysis of whole cell lysates of EpH4 and EpH4-Snail cells.

Supplementary Figure 4

Comparison of SM and cholesterol content

(**A,B**) Comparison among EpH4, EpH4-Snail, and fibroblasts.

(**C,D**) Comparison between esophageal cancer cell lines TE-15 (Snail-negative) and TE-8 (Snail-positive). n = 3, ** $p < 0.01$, * $p < 0.05$, n.s., $p \geq 0.05$ by Tukey's multiple comparison test (A,B) and Student's t-test (C,D).

Supplementary Figure 5

Selective growth inhibition of a Snail positive esophageal cancer cell line by combined treatment of cyclosporin A and TMP-153

(**A-C**) Effects of treatments with cyclosporin A (**a**), TMP-153 (**b**) or combination of 50 μM cyclosporin A and indicated concentration of TMP-153 (**c**) to cellular growth of human esophageal cancer cell lines TE-15 and TE-8. Cells were treated with drugs for 24 h and relative cell number to 0.1% DMSO control was determined by CCK-8. $n \geq 3$, error bars, SDs. Significant differences between TE-15 and TE-8 by unpaired one-sided Student's t-test are shown (*, $p<0.05$).

(**D-F**), Combined treatment of cyclosporin and TMP-153 inhibits the growth of TE-8 tumor xenografts. Experiments were performed as described in Materials and Methods. In brief, 10 mg/kg cyclosporin A and 10 mg/kg TMP-153 were simultaneously administered intraperitoneally to nude mice bearing TE-8 tumor xenografts. After 36 days of the first administration, the overall appearance (D), the excised tumors (E), and the tumor size (F) of control and treated mice were compared. $n = 5$, error bars, SDs. Significant differences by unpaired one-sided Student's t-test are shown (*, $p < 0.05$).

Table S1 Primer sequences used in this study

Gene		Primer sequence	
CerS isozymes			
mouse	CerS1	forward	5'-CTGTACATTGTGGCTTTCGC-3'
		reverse	5'-GGCTGTCTGAGCTTCCAGAG-3'
mouse	CerS2	forward	5'-CCCTGCTCTTTCTCGTCATTC-3'
		reverse	5'-CGTGACAAAAGGTCTACCTCC-3'
mouse	CerS3	forward	5'-TTTCAAGCATTCCACAAGCAAAC-3'
		reverse	5'-ATCGCCAGCAAGACTCCT-3'
mouse	CerS4	forward	5'-GGCTGTGCGAATTGTCTT-3'
		reverse	5'-CGGGTTGGGCTTTATCTTT-3'
mouse	CerS5	forward	5'-AGTAAGGTTATTGGCTCTGTGTT-3'
		reverse	5'-GCTCAGAAGTGACCAGACAAT-3'
mouse	CerS6	forward	5'-CCGGAGTGGACAAAGCAAGAT-3'
		reverse	5'-GCCGTTCTTACCTCTCTGGAAGATG-3'
Elovl isozymes			
mouse	Elovl1 variant1	forward	5'-TGGAGGCTGTTGTGAACTTG-3'
		reverse	5'-ATGAAGCCTCGAAGTTGGAA-3'
mouse	Elovl1 variant2	forward	5'-GATCCCTTTG AACCCTTCAC-3'
		reverse	5'-TCAGAAACCGCATGCTTCAT-3'
mouse	Elovl2	forward	5'-GATCCCTTTGAACCCTTCAC-3'
		reverse	5'-GAGAATCTCGTGGTCCAAACA-3'
mouse	Elovl3	forward	5'-CGGGTTAAAAATGGACCTGA-3'
		reverse	5'-CCAACAACGATGAGCAACAG-3'
mouse	Elovl4	forward	5'-ACCGTGGAGTTCTATCGCTG-3'
		reverse	5'-TGCTTATGCTTATCGTTGGC-3'
mouse	Elovl5	forward	5'-CTTGCACATCCTCCTGCTC-3'
		reverse	5'-CTGAGTGACGCATCGAAATG-3'
mouse	Elovl6	forward	5'-CCAATGGATGCAGGAAAACT-3'
		reverse	5'-AACTTGGCTCGCTTGTTCAT-3'
mouse	Elovl7	forward	5'-GGAAAATGGCGTTCAGTGAT-3'
		reverse	5'-CATAGAGGCCCAGGATGATG-3'