Chemotherapy resistance due to epithelial-to-mesenchymal transition is caused by abnormal lipid metabolic balance

Atsushi Matsumoto[1]†, Akihito Inoko[2,3]†, Takuya Tanaka[4], Gen-Ichi Konishi[4], Waki Hosoda[5], Takahiro Kojima[6], Koji Ohnishi[2], Junichi Ikenouchi[1] *

Affiliations

1. Department of Biochemistry, Kyushu University Graduate School of Medical Sciences, Fukuoka, 812-8582, Japan.

2. Department of Pathology, Aichi Medical University School of Medicine, Nagakute, 480-1195, Japan

3. Division of Cancer Epidemiology and Prevention, Aichi Cancer Center Research Institute, Nagoya, 464-8681, Japan

4. Department of Chemical Science and Engineering, Institute of Science Tokyo, Meguro-ku, Tokyo 152-8552, Japan.

5. Department of Pathology and Molecular Diagnostics, Aichi Cancer Center Hospital, Nagoya, 464-8681, Japan

6. Department of Urology, Aichi Cancer Center Hospital, Nagoya, 464-8681, Japan

†. These authors contributed equally to this work.

*Corresponding author. Junichi Ikenouchi M.D., Ph.D.

Tel.: +81-92-802-4292; Fax: +81-92-802-4292

Email: ikenouchi.junichi.033@m.kyushu-u.ac.jp

Abstract

Invasive cancer is defined by the loss of epithelial cell traits resulting from the ectopic expression of epithelial-mesenchymal transition (EMT)-related transcription factors such as Snail. Although EMT is known to impart chemoresistance to cancer cells, the precise molecular mechanisms remain elusive. We found that Snail expression confers chemoresistance by upregulating the cholesterol efflux pump ABCA1 as a

countermeasure to the excess of cytotoxic free cholesterol relative to its major

interaction partner in cellular membranes, sphingomyelin. This imbalance is introduced

by the transcriptional repression of enzymes involved in the biosynthesis of

sphingomyelin by Snail. Inhibiting esterification of cholesterol, which renders it inert,

selectively suppresses growth of a xenograft model of Snail-positive kidney cancer. Our

findings offer a new perspective on lipid-targeting strategies for invasive cancer therapy.

Introduction

Metastatic cancer, characterized by the spread of tumor cells to distant organs, remains

a major challenge in cancer treatment due to its high mortality rate. Metastasis involves

a series of biological events, such as acquisition of cancer stem cell characteristics,

tumor-microenvironment interactions, alterations in cell metabolism and resistance to

chemotherapy, that are mediated by epithelial-mesenchymal transition (EMT) (1-4).

Transcription factors driving EMT (EMT-TFs), including Snail, Slug and Twist, that are

ectopically expressed in invasive cancers were initially thought to promote invasion and

metastasis by suppressing the expression of adhesion molecules in cancer cells. However,

since E-cadherin-mediated adhesion is essential for cell survival after metastasis (5),

invasive cancer cells do not undergo complete EMT but rather sustain a hybrid state with

both epithelial and mesenchymal (hybrid E/M) characteristics. In a mouse model of

pancreatic carcinoma and breast cancers where key EMT-TFs were deleted, the

suppression of EMT did not alter the appearance of systemic dissemination or metastasis

but rather contributed to increased sensitivity to anticancer drugs (3, 4). Therefore,

in-depth understanding of the molecular mechanisms of resistance to anti-cancer drugs

conferred by EMT-TFs is an essential prerequisite for devising effective therapeutic

interventions (6). It was recently reported that expression of a Rho family GTPase,

RHOJ, is upregulated in cancer cells undergoing EMT and that its activation enhances the

DNA damage response, which enables tumor cells to efficiently repair

chemotherapy-induced DNA damage (7). Here, we investigated EMT-dependent

acquisition of resistance to anticancer agents that do not directly cause DNA damage,

such as those targeting the ERK signaling pathway and found that expression of

EMT-TFs causes metabolic alterations of sphingomyelin, resulting in ectopic induction

of drug efflux transporter expression and conferring chemotherapy resistance to cancer

cells.

Results

Snail-induced ABCA1 expression confers chemoresistance to hybrid E/M cells

Renal cell carcinoma (RCC) is notorious for the lack of sensitivity to

chemotherapy (8), but the underlying mechanisms are not elucidated. Nitidine chloride

(NC) holds promise as a drug treatment against RCC as it was shown to have a

pro-apoptotic effect on RCC and inhibit tumor growth by suppressing the ERK

signaling pathway in a xenograft model (9, 10). However, in human lung

adenocarcinoma cells, NC efficacy was inversely correlated with the expression level of

the drug transporter ABCA1: cancer cells with downregulated ABCA1 were more

sensitive to NC treatment (11). Therefore, we wondered whether ABCA1 could also

confer NC resistance in RCC. We treated five different RCC cell lines with a

combination of NC and an ABCA1 inhibitor cyclosporin A (CsA). Anti-proliferative

effect of NC was significantly enhanced in the presence of CsA for three lines among

five, Caki-1, 786-O and A498 (Figure 1A). These cells consistently expressed higher

levels of ABCA1 compared to the non-responsive cells (Figure 1B and C), suggesting

the acquisition of ABCA1-mediated drug resistance. Notably, ABCA1 expression was

positively correlated with expression of the EMT-TF Snail in the RCC cell lines (Figure

1D and E). To investigate clinical relevance of ABCA1 upregulation in cancers, we

analyzed ABCA1 expression in cancer using the UCSC Xena platform. Comparative

ABCA1 expression data of both normal tissue and primary tumors were available for 24

cancer subtypes in the GDC TCGA database. In addition to glioblastoma and head and

neck cancer, all three types of renal cancers (clear cell carcinoma (ccRCC),

chromophobe carcinoma and papillary cell carcinoma) exhibited significant ABCA1

upregulation in tumor (Figure 1C and Figure 1—figure supplement 1A and B),

indicating that ABCA1 upregulation frequently occurs in clinical renal cancers.

To examine whether the enrichment of ABCA1 protein in malignant cancer cells

is detectable in clinical cases, we conducted immunohistochemical analysis on surgical

specimens taken from patients with primary ccRCC. The findings showed that ABCA1

was upregulated at the lesion site of high-grade ccRCC (Figure 1G). This clinical data

aligns with our cellular experimental results, as Snail expression is known to be

upregulated in high-grade ccRCC. Upon detailed comparison among grades, ABCA1

signal was more prominent in the patients with the high-grade classification (Figure 1H

and Figure 1—figure supplement 2), suggesting that ABCA1 could serve as a marker for

high-grade ccRCC. Xena analysis also revealed significantly higher expression of Snail

in ccRCC, also supporting the idea that ABCA1 upregulation is correlated with

exogenous Snail expression (Figure 1—figure supplement 1C).

Our results suggest the correlation between chemoresistance and Snail

expression level. In order to clarify whether Snail expression alone can confer resistance

to anti-cancer drugs in epithelial cells, Snail was ectopically expressed in normal

epithelial cells and we investigated changes in resistance to anti-cancer drugs.

Overexpression of Snail in a mouse mammary gland epithelial cell line EpH4 abolishes

transcription of various genes involved in cell-cell adhesion including E-cadherin (Figure

1I) (12-14). EpH4-Snail cells exhibited coexistence of the mesenchyme-specific

intermediate filament Vimentin and the epithelium-specific Cytokeratin-18 (Figure 1J),

which is a typical feature of a hybrid E/M state of cancer cells. The hybrid E/M

EpH4-Snail cells were significantly more resistant to NC than parental EpH4 cells

(Figure 1K). Consistent with the results of kidney cancer cell lines, this resistance was

cancelled by CsA (Figure 1K), and upregulation of ABCA1 expression was also

observed in EpH4-Snail cells (Figure 1L and M). Consistent with the resistance to the

antitumor agent, ABCA1 mainly localized plasma membrane in EpH4-Snail cells

(Figure 1M). We also examined another model with a different cell line. In MDCK II, a

canine kidney epithelial cell line, TGF-β treatment induced EMT under the expression of

the oncogenic KRAS-G12V mutant (15). The upregulation of ABCA1 expression was

also observed in this transient hybrid E/M model (Figure 1N), suggesting that similar

mechanisms to those in EpH4-Snail are also at play in kidney cell EMT. Notably,

induction of ABCA1 expression requires a longer period of treatment with TGF-β (day 8)

than Snail induction (day 3), indicating that ABCA1 is upregulated in response to cellular

alteration by Snail but not induced directly by TGF-β signaling. Furthermore, in

comparison between human esophageal cancer cell lines (21), Snail-positive TE-8 cells

exhibited higher expression of ABCA1 than Snail-negative TE-15 (Figure 1O). These

results suggest that the induction of ABCA1 by ectopic expression of Snail occurs

regardless of cellular background.

Intracellular cholesterol accumulation drives ABCA1 expression in hybrid E/M cells

Then how does Snail induce ABCA1 expression, since as a transcriptional repressor it is

incapable of doing so directly? Upregulation of ABCA1 was not observed in either

E-cadherin or α-catenin KO cells (Figure 1L), indicating that ABCA1 induction is driven

as a specific response to a fundamental change in cellular state induced by Snail

overexpression, not an ancillary effect of disrupting either cell-cell adhesion or

apical-basal polarity. Generally, ABCA1 is a transporter for the efflux of cholesterol out

of the cell when there is an excess of cholesterol in the cell and ABCA1 upregulation in

response to excess loading of cholesterol is mediated by sterol-activated nuclear

receptors, LXRs (16). Aside from the cholesterol-LXRs regulatory axis, it was recently

reported that ABCA1 expression in epithelial cells is regulated via FOXO3a and c-Myc

(17). Activation of FAK/PI3K/Akt signaling depending on cell crowding state

downregulates ABCA1 expression via phosphorylation of FOXO3a, a positive

transcriptional regulator of ABCA1 (17). We found that phosphorylation level of

FOXO3a at S453 tended to be lower in EpH4-Snail cells than in wild-type EpH4 cells

(Figure 2—figure supplement 1A). However, since downregulation of Akt signaling by

treatment with inhibitors of its activators GSK2334470 (PDK1), LY294002 (PI3K) and

Wortmannin (PI3K) did not induce ABCA1 expression in wild-type EpH4 cells (Figure

2—figure supplement 1A), FOXO3a would not be responsible for ABCA1 expression in

EpH4-Snail cells. Although ABCA1 expression is also downregulated by c-Myc in breast

cancer, no remarkable difference of c-Myc expression was observed between EpH4 and

EpH4-Snail cells, indicating that c-Myc is not likely involved in ABCA1 expression in

EpH4-Snail cells (Figure 2—figure supplement 1B). Therefore, we next pursued in detail

the possibility that LXRs induce ABCA1 in EpH4-Snail cells. Immunofluorescence

microscopy revealed that LXRs are highly accumulated in the nuclei of EpH4-Snail cells,

suggesting that ABCA1 upregulation could be mediated by LXRs (Figure 2A).

Consistently, treatment with an LXR inhibitor GSK2033 abolished ABCA1 expression to

an undetectable level in EpH4-Snail cells (Figure 2B). GSK 2033 treatment also

canceled NC resistance (Figure 2C), suggesting that cholesterol efflux system activated

by LXR is important for acquisition of chemoresistance. However, EpH4-snail cells did

not show enhanced sensitivity to NC treatment even when ABCA1 was depleted by

knockout (Figure 2—figure supplement 2). Given that CsA inhibits other isoforms of

ABC transporters (18), other LXR-regulated efflux pumps may also contribute to

chemoresistance.

Since transcriptional activity of LXRs are regulated by cellular cholesterol level,

we examined whether ABCA1 is expressed in response to high cholesterol accumulation

in the cells. Inhibition of cholesterol biosynthesis by simvastatin, an inhibitor of the

rate-limiting enzyme of cholesterol biosynthesis 3-hydroxy-3-methylglutaryl-coenzyme

A reductase, remarkably decreased ABCA1 expression in EpH4-Snail cells but residual

ABCA1 was still detected (Figure 2D, normal FBS). In contrast, treatment with

simvastatin in the medium supplemented with lipid-free FBS completely abolished

ABCA1 expression (Figure 2D, lipid-free FBS). Together with the finding that

incubation in lipid-free FBS medium alone did not markedly decrease ABCA1

expression, these results suggest that cholesterol biosynthesis is the primary source of

excess accumulation of cholesterol in EpH4-Snail cells, but cholesterol uptake from

media also significantly contribute to the accumulation as a secondary source.

These findings imply dysregulation of cholesterol metabolism in EpH4-Snail. Filipin staining revealed that cholesterol was prominently accumulated in intracellular structures in EpH4-Snail cells, whereas it localized primarily in the plasma membrane of wild-type EpH4 cells (Figure 2E). Since excess cholesterol can be stored in lipid droplets, we stained cells with Lipi-Green to visualize lipid droplets and found that lipid droplets were significantly enlarged in EpH4-Snail (Figure 2F and G). Consistently, TopFluor cholesterol, a fluorescent cholesterol analogue, was localized to lipid droplets in EpH4-Snail cells (Figure 2H). On the other hand, the intracellular filipin signals did not co-localize with the lipid droplet protein ADRP (perilipin-2) but co-localize with the lysosomal protein LAMP1 instead (Figure 2I). Given that filipin stains unesterified cholesterol, internalized cholesterol accumulates in lysosomes in its unesterified form and in lipid droplets in its esterified form in EpH4-Snail cells.

We next compared the total cholesterol levels in EpH4 cells and EpH4-Snail cells; the cellular amount of cholesterol slightly increased in EpH4-Snail cells, but the difference was modest and not significant (Figure 2J). Therefore, the internalization of cholesterol in EpH4-Snail cells is not simply due to an increase in the amount of cholesterol, but would be due to an abnormality in the subcellular distribution of cholesterol. A large proportion (30-50%) of cellular cholesterol is associated with sphingomyelin, and depletion of sphingomyelin induces internalization of cholesterol (19). To investigate contents of cholesterol and sphingomyelin in the plasma membrane, we utilized the lipid-binding protein probes domain 4 of perfringolysin O (cholesterol,

D4) and lysenin (sphingomyelin). Intriguingly, cholesterol content in the plasma

membrane was not affected by EMT, but sphingomyelin content decreased in

EpH4-Snail cells (Figure 2K and L). Collectively, the increase in a

sphingomyelin-unbound form of cholesterol in the plasma membrane results in the

internalization of cholesterol and activation of LXR to induce ABCA1 expression in

hybrid E/M cells.

The strong interaction between sphingolipids and cholesterol contributes to the

formation of tightly packed and less fluid plasma membrane (20). Alteration of this

physical property of plasma membrane frequently occurs in cancer cells and is thought

to affect malignancy (21). Therefore, we next addressed the impact of sphingomyelin

decrease on the lipid packing of plasma membrane using FπCM-SO$_3$, a recently

developed plasma membrane-specific solvatochromic probe (22). Solvatochromic dyes

respond to the polarity of their environment, exhibiting a red shift in more polar

solvents. In lipid bilayers, these dyes similarly sense differences in lipid packing (order)

and display a red shift in more fluid, disordered membranes (Figure 2—figure

supplement 3A). Thus, variations in lipid order can be visualized as changes in the ratio

of signal intensities of the red to blue channels (Figure 2—figure supplement 3B). The

negatively charged and alkylated modifications enable FπCM-SO$_3$ to stably reside in

the plasma membrane of living cells. Ratiometric imaging using FπCM-SO$_3$ revealed

that the plasma membrane of EpH4-Snail cells was more disordered (Figure 2M–O).

While plasma membrane fluidity is usually attributed to the cholesterol content (23), our

results suggest that ectopic expression of Snail induces fluidization of plasma

membrane, not by altering the surface cholesterol level but by decreasing

sphingomyelin content.

Downregulation of VLCFA-SM synthesis induces Chol/SM imbalance in hybrid E/M cells

The present findings suggest that the cause of chemoresistance may trace back to alterations of the balance of sphingomyelin and cholesterol. Therefore, we next investigate how sphingomyelin metabolism is affected by ectopic expression of Snail. We found that the total sphingomyelin content relative to total phospholipids decreased in EpH4-Snail cells (Figure 3A). Consequently, the ratio of cholesterol to SM (Chol/SM ratio) in the cells significantly increased in EpH4-Snail cells (Figure 3B). To investigate the SM profile in detail, we next analyzed the fatty acid composition of SM using LC-MS. In EpH4-Snail cells, the fractions of VLCFA-SMs (22:1, 22:0 and 24:1) were significantly decreased whereas those of LCFA-SMs (14:0, 16:1 and 16:0) were rather increased (Figure 3C), resulting in a prominent decrease in the total VLCFA-SM fraction (Figure 3D). These results suggest that decrease in VLCFA-SMs are responsible for decrease in total SM content in EpH4-Snail cells. On the other hand, the decrease in the cell surface content (Figure 2L) was more prominent than that in total SM (Figure 3A). Given that VLCFA-SMs are suggested to be differently trafficked in recycling from endosome to plasma membrane (24), their reduction may lead to decreased plasma membrane sphingomyelin content by altering its intracellular distribution.

Fatty acid composition of SM is mainly modulated by substrate specificity of two families of enzymes, Elovls and CerSs, that respectively catalyze the rate-limiting step of the fatty acid elongation cycle and the N-acylation of sphingoid bases to produce ceramides (Figure 3E) (25, 26). Therefore, we hypothesized that transcriptional

repression of genes from these families causes the decrease in VLCFA-SM content in

EpH4-Snail cells. RT-PCR analyses identified ELOVL7 and CERS3 as genes which are

repressed specifically in EpH4-Snail cells (Figure 3F and G). Consistent with our

hypothesis, Elovl7 and CerS3 control VLCFA-SM production (Figure 3E). Collectively,

ectopic expression of Snail represses transcription of VLCFA-SM-producing enzymes

and results in a decrease in the amount of VLCFA-SM. This downregulation of

biosynthesis pathway is responsible for the induction of ABCA1 expression, because

supplementation of VLCFA ceramide decreased the ABCA1 expression in EpH4-Snail

cells (Figure 3H). Together with the results of the cholesterol depletion experiment

(Figure 2D), the high Chol/SM ratio due to downregulation of VLCFA-SM biosynthesis

accounts for the elevated ABCA1 expression in EpH4-Snail cells.

Since ABCA1 expression is not aligned with either the loss of epithelial state

(E-cadherin and α-catenin KO; Figure 1L) or a non-epithelial basal state (fibroblasts),

upregulation of ABCA1 is characteristic to the hybrid E/M cells expressing ectopic Snail

(Figure 3I). To investigate the mechanism underlying this difference in ABCA1

expression, we again compared Chol/SM ratio in EpH4-Snail and normal fibroblast cell

lines L-929 and NIH3T3. Chol/SM ratio of these fibroblasts were slightly but not

significantly higher than that of EpH4 (Figure 3J). The cholesterol content in NIH-3T3

cells is significantly higher than in EpH4 cells (Figure3—figure supplement 1).

Consequently, normal fibroblast would synthesize more SMs to counteract the

cholesterol load imposed by the decrease in VLCFA-SMs and stabilize the Chol/SM

ratio. By contrast, hybrid E/M cells are unable to respond appropriately to the harmful

imbalance between cholesterol and SM due to the incomplete activation of the

mesenchymal transcriptome.

To examine whether the above findings apply to human cancer cells, we

investigate the Chol/SM ratio of TE-15 and TE-8 (see also Figure 1O) (27). Consistent

with our findings in the EpH4-Snail system, Chol/SM ratio was significantly higher in

TE-8 cells than in TE-15 cells (Figure 3K). Together with the results of mouse cell lines,

normal epithelial and fibroblast cells maintain Chol/SM ratio around 1.0-1.5, and

ABCA1 expression as a response to the excess Chol is induced when Chol/SM ratio

exceed 1.5 (Figure 3L). Notably, the difference in either cholesterol or SM contents

between TE-15 and TE-8 cells, and cholesterol content between EpH4 and EpH4-Snail is

not significant when each are compared alone (Figure 2D and Figure 3—figure

supplement 1), indicating the importance of considering their functional cooperativity,

i.e. to analyze the Chol/SM ratio. Collectively, the cause of chemoresistance can be

traced back to the downregulation of VLCFA-SM biosynthesis (Figure 3M). A decrease

in sphingomyelin but not in cholesterol induces plasma membrane fluidization, which

leads to internalization of cholesterol. The internalized, SM-unbound cholesterol

activates LXRs, which in turn elevate ABCA1 expression and thereby confer

chemoresistance to cells.

Inhibitor of cholesterol esterification selectively inhibits growth of

Snail-positive kidney cancer cells

Finally, we examined the possibility of targeting the imbalance of Chol/SM ratio

in Snail-positive cancer cells to selectively inhibit their growth. As discussed above,

hybrid E/M cells of high Chol/SM ratio must either export cholesterol out of cells or

isolate it within cells in lipid droplets as cholesteryl esters to avoid the cytotoxic effects of

free cholesterol. Thus, inhibiting the means of clearing excess cholesterol could

specifically target cells in the hybrid E/M state for self-extermination. Cholesterol efflux

from cells is mainly dependent on ABCA1, and esterification of cholesterol is catalyzed

by acyl-CoA:cholesterol acyl transferases (ACATs) (Figure 4A) (28). Among the two

ACAT isoforms in humans, SOAT1 is widely expressed, including in the kidney (29).

Indeed, all RCC cell lines tested express SOAT1 (Figure 4B), although the expression

levels did not correlate with those of Snail in the respective cell lines (Figure 1B).

Therefore, we examined the effect of ABCA1 inhibitor CsA and ACAT inhibitor

TMP-153 on the EpH4-Snail model and renal carcinoma cells. While CsA

non-selectively inhibited growth of all cell lines tested (Figure 4C and E), TMP-153

selectively inhibited growth of Snail-positive cells (EpH4-Snail, 786-O, A498 and

Caki-1; Figure 4D and 4F). Thus, sequestration of cholesterol via esterification by

ACATs is the primary mechanism for handling excess cholesterol to avoid cell death in

these cells. Notably, TMP-153 treatment alone did not provide this selectivity between

esophageal cancer cell lines, but combination of TMP-153 and CsA selectively inhibited

growth of the Snail-positive TE-8 cells (Figure 4—figure supplement 1 A-C).

Collectively, these results suggest that inhibition of cholesterol efflux and cholesterol

esterification selectively inhibit growth of cells stably in a Snail-induced hybrid E/M

state.

We also investigated the potential of ACAT inhibitors as anti-tumor agents in

vivo using xenograft models. We successfully transplanted the 786-O line into nude mice

among Snail-positive ccRCC cell lines. Using this system, we found that TMP-153

treatment significantly reduced 786-O tumor volume in vivo (Figure 4G-I). Moreover,

the combination of TMP-153 and CsA also inhibited the growth of TE-8 tumor

1 xenografts (Figure 4—figure supplement 1D-F). Together with the in vitro assays

2 (Figure 4C-F and Figure 4—figure supplement 1 A-C), we propose that cholesterol

3 clearance systems, particularly ACATs, can be used as molecular targets to selectively

4 inhibit the growth of Snail-positive cancers.

Discussion

In recent years, it has been revealed that the expression of EMT-TFs, such as

Snail, induces cancer cells to acquire mesenchymal characteristics while retaining

epithelial traits, transforming them into so-called hybrid E/M cells. Hybrid E/M cells

have been demonstrated to exhibit the highest plasticity and malignancy (30, 31).

Moreover, Snail expression contributes to cancer stemness and resistance to

chemotherapy (2). To date, molecular mechanisms connecting to EMT-TFs to

chemoresistance remain to be established. Here we identify lipid metabolic

imbalance-mediated ABCA1 expression as a mechanism of chemoresistance induced by

Snail. We found reduced VLCFA-SM biosynthesis in Snail-positive cells with hybrid

E/M phenotypes, resulting in high Chol/SM ratio. This should be detrimental in theory

since it increases exposure to the toxic effect of free cholesterol but cancer cells deploy

countermeasures: upregulating ABCA1 to move cholesterol out of cells and sequestering

esterified cholesterol in the form of lipid droplets. Notably, it has been also reported that

not only ABCA1 (11, 32) but also enhanced lipid droplet formation triggers

chemoresistance to various drugs (33-35). Therefore, dietary or pharmacological control

of cholesterol and also sphingolipids would be a new strategy to overcome resistance to

chemotherapy.

The aberrant activation of EMT-TFs is implicated in multiple crucial steps in cancer progression, prompting efforts to explore therapeutic approaches aimed at suppressing EMT-TF functions in cancer cells (6). However, as EMT-TFs are predominantly located in the nucleus, no effective treatment methods, including antibody drugs, have been established thus far. Therefore, instead of directly targeting EMT-TFs, focusing on the metabolic network vulnerabilities specific to hybrid E/M cells may prove to be an effective therapeutic strategy. In the present study, we found increased sensitivity of Snail-positive cells to toxicity of free-cholesterol accumulation induced by the ACAT inhibitor TMP-153. Notably, kidney cancer growth could be suppressed by inhibiting cholesterol esterification alone while esophageal cancer required the additional downregulation of ABCA1, indicating varying dependence on free cholesterol clearance strategies among cancers of different primary origin. Other cancers might employ alternative pathways to evade cytotoxic effects of excess cholesterol and identifying these traits will pave the way for the development of novel strategies in targeted molecular therapeutics against hybrid E/M cancers. An intriguing corollary here is the potential utility of ABCA1 expression as a diagnostic marker to identify cancers that would be sensitive to targeting the Chol/SM imbalance.

Although various metabolic changes in cancer cells focusing on energy sources including sugars, amino acids and fatty acids have been intensely studied (36), alteration of sphingolipid homeostasis driven by EMT-TFs remains largely unknown. We found both quantitative and qualitative alteration of sphingomyelin profile: decrease in the total SM amount and the VLCFA-SM percentage. This alteration is attributed to transcriptional repression of enzymes for biosynthesis of VLCFA-SMs, CerS3 and Elovl7. Genes whose transcription is repressed by Snail include those encoding proteins

directly involved in epithelial adhesive structures, such as E-cadherin and Claudins.

Hence, it is not surprising that VLCFA-SMs play important roles cooperatively with the

adhesion molecules. Indeed, cholesterol accumulation is required for tight junction

formation, and VLCFA-SMs are enriched in the junctional region (37, 38). Studies

focusing on the VLCFA-synthetic enzymes would provide a new layer of knowledge

about mechanisms underlying the establishment of epithelial structures and functions.

Such studies would also contribute to understanding how dysfunctions in lipid

metabolisms promote cancer malignancy and tissue fibrosis as a consequence of EMT.

The regulation of cell adhesion by fine-tuning of lipid composition should be reversible

and therefore would explain the two conflicting events: the invasion and metastasis

process and the viability process at the metastatic site.

Materials and methods

Cells and reagents

Cells were grown in DMEM supplemented with 10% FBS under 5% CO_2 at 37°C.

Mouse Snail-expressing EpH4 cells (EpH4-Snail)(14), E-cadherin KO and α-catenin

KO EpH4 cells (37) were generated previously. EpH4-Snail cells were maintained in

medium supplemented with 500 μg/ml G418 (Wako Pure Chemical). EpH4 cells were

kindly provided by Dr. Reichmann (Institut Suisse de Recherches, Lausanne,

Switzerland), and amplification of mouse-specific genomic sequences was confirmed.

HEK293, 786-O, and A498 cells were obtained from the American Type Culture

Collection (ATCC); Caki-1 and KMRC-3 cells were obtained from the Japanese

Collection of Research Bioresources (JCRB) Cell Bank; TE-8 and TE-15 cells were

obtained from the Tohoku University Cell Resource Center for Biomedical Research;

and TUHR14TKB cells were obtained from the RIKEN BioResource Research Center

(BRC). Human cell lines were authenticated by short tandem repeat (STR) profiling

using the GenePrint10 System (Promega). All cell lines were routinely tested and

confirmed to be negative for mycoplasma contamination.

pLVET-HA-K-RasG12V-IRES-GFP (Addgene #107140), LAMP1-mRFP-FLAG

(#34611), pLentiPGK DEST H2B-iRFP670 (#90237) were transduced using the

lentivirus produced in HEK293 cells, and fluorescence-positive cells were collected

using a cell sorter Sony SH-800 as needed. pEGFP-C1-ADRP (Addgene #78161) was

transfected to cells using Lipofectamine 3000 (Thermo Fischer). Nitidine chloride was

purchased from Selleck Biotech, cyclosporin A from Wako, and TMP-153 from

Cayman Chemical, filipin complex from *Streptomyces filipinensis* and GSK2033 from

Sigma-Aldrich, Lipi-Green from Dojindo Laboratories. Ceramide (d18:1/22:0; C22:0

ceramide) from Avanti Polar Lipids.

The following primary antibodies were used for immunoblotting (IB),

immunofluorescence microscopy (IF) and immunohistochemistry (IHC): Rabbit

anti-ABCA1 mAb (96292S, IB, IF), rabbit anti-Snail mAb (3879S, IB), rabbit anti-Akt

mAb (4691S, IB), rabbit anti-phospho-Akt (T308) mAb (2965S, IB), rabbit

anti-phospho-FoxO3a (S253) pAb (9466S, IB) from Cell Signaling Technology; Rabbit

anti-ABCA1 pAb (NB400-105, IHC) from Novus Biologicals; Rat anti-E-cadherin

mAb (ECCD-2, IB, IF) from TaKaRa; Mouse anti-cytokeratin 18 mAb (LS-C84878, IF)

from LS Bio; Goat anti-LXR alpha + LXR beta pAb (ab24362, IF) from Abcam; Mouse

anti-α-tubulin mAb (12G10, IB) produced in house.

The following secondary antibodies were used: HRP-conjugated anti-rat IgG

(HAF005) and anti-mouse IgG (HAF007) from R&D Systems; HRP-conjugated

anti-rabbit IgG (4030-05) from Southern Biotech; Cy3-conjugated anti-goat IgG

(705-165-147) from Jackson Immunoresearch Laboratories.

RT-PCR

Total RNAs were prepared with RNeasy Mini Kit (Qiagen) and the

retrotranscribed with Superscript III First-Strand Synthesis System (Invitrogen).

RT-PCR was then performed using KOD FX (TOYOBO) following manufacturer's

instructions. Primer sequences used are shown in Table S1.

Lipid analysis

Total lipids were extracted from cultured cells by Bligh-Dyer's method, dissolved

in methanol/chloroform = 1/1 (v/v). Phospholipid content was determined by phosphate

assay. SM and Chol content were determined using Sphingomyelin assay kit STA-601

(Cell Biolabs) and Amplex Red Cholesterol Assay Kit (Invitrogen), respectively,

according to the manufacturer's instructions. Fatty acid profile of SM was determined

with LC-MS as described previously (39). Briefly, electrospray ionization mass

spectrometry analysis was performed on a 6420 triple-quadrupole liquid

chromatography–mass spectrometer (Agilent Technologies) equipped with an HPLC

system and an auto sampler (Infinity 1260; Agilent Technologies). The extracted lipids

were directly subjected to electrospray ionization mass spectrometry analysis. The

mobile phase composition was acetonitrile/methanol/water = 18:11:1 (0.1% ammonium

formate). The flow rate was 4 µl/min. The mass range of the instrument was set at

650–950 *m/z*. *m/z* profiles of SM were extracted according to total ion chromatogram

patterns, and all peaks corresponding to $[M+H]^+$ ions of d18:1-even chain length fatty

acid-SMs were subjected to analyses.

Fluorescence microscopy

To visualize subcellular localization of cholesterol, cells cultured on glass-bottom

dishes were fixed with 4% paraformaldehyde for 15 min at room temperature and

stained with 50 μg/ml filipin prepared in PBS for 30 min. To visualize lipid droplets,

live cells were washed once with HBSS and stained with Lipi-Green (Dojindo) for 20

min at 37°C. Cells were washed twice with HBSS and observed on a heating stage at

37°C. For immunofluorescence microscopy, cells cultured on coverslips were fixed

with 4% paraformaldehyde or 2% formalin prepared in PBS for 15 min at RT and

permeabilized with 0.1% Triton X-100 prepared in PBS. Fixed cells were blocked with

1% BSA prepared in PBS for 1 h at RT. Cells were incubated with primary antibodies

for 1 h at RT and secondary antibodies for 1 h at RT. Antibodies were prepared in the

blocking solution. For detection of ABCA1, cells were fixed with MeOH for 20 min at

-30°C and antibodies were prepared in Can Get Signal immunostain Immunoreaction

Enhancer Solution B (TOYOBO). Fixed cells were observed at RT. For imaging of

plasma membrane cholesterol and sphingomyelin, cells cultured on glass-bottom dishes

were washed with HBSS three times, incubated with 1x working solution of CellMask

Green plasma membrane stain (Molecular Probes) in HBSS, subsequently incubated

with 5 μg recombinant scarlet-tagged D4 or 10 μg RFP-tagged lysenin in 100 μl HBSS

for 30 min, washed three times with HBSS and observed at 37°C. For ratiometric

imaging using FπCM-SO$_3$ (22), cells cultured on glass-bottom dishes were incubated

with 1 mM FπCM-SO$_3$ in HBSS for 10 min at 37°C and observed at 37°C without

washout. The dye was excited with a 405 nm laser, and 400–500 nm (Blue) and

500–700 nm (Red) signals were split with variable dichroic mirror and simultaneously

detected with two photomultipliers. The raw images were gaussian blurred, ratio and

mean intensity of two channels was calculated, and then data are expressed as a hue

(ratio) – brightness (intensity) image (see Figure 2—figure supplement 3). All

observation was performed with a confocal microscope (Carl Zeiss LSM900) equipped

with Plan-APO (63×/1.40 NA, oil immersion) objective. Images were acquired using

Carl Zeiss Zen 3.4 software. Images were analyzed using ImageJ/Fiji.

Immunoblotting

Whole cells were lysed with SDS sample buffer and samples were resolved by

SDS-PAGE and transferred to nitrocellulose membranes. After blocking with 5%

skim-milk prepared in TBS-T, membranes were incubated with primary antibody for 1

h at RT or overnight at 4°C. Membranes were then washed and incubated with

HRP-conjugated secondary antibody for 1 h at RT. Chemiluminescence signal was

detected using a LAS4000mini imaging system (Fujifilm) and images were analyzed

using ImageJ/Fiji. Protein levels were calculated relative to the signal intensity of

α-tubulin as a loading control.

Ceramide supplementation to culture medium

C22:0 ceramide was supplemented to culture medium as described previously (40).

The ceramide was dissolved in dodecane/ethanol = 2/98 solution at 0.63 mM (stock

solution). The stock solution (or vehicle control) was added to DMEM+10% FBS

medium at a final solvent concentration of 0.4%, resulting in a final ceramide

1　concentration of 2.5 µM. The prepared medium was used immediately. Cells were

2　cultured in this medium for 24 h and then subjected to further analysis.

Growth inhibition assay

Cells cultured in 96-well microtiter plates (2×10^3 cells/well, 48 h) were subjected

to treatments with nitidine chloride, GSK2033, CsA, TMP-153 or their combination

dissolved in DMSO for 24 h at 37°C. In experiments involving GSK2033, the

compound was added 24 h prior to treatment with the other drugs. Final DMSO

concentrations were less than 0.25%. After treatment, metabolically active cell number

was determined using Cell Counting Kit-8 (Dojindo) according to manufacturer's

instructions. Briefly, 10 µl of CCK-8 solution was directly added to each well

containing 100 µl culture medium and cells were incubated 1 h at 37°C. Reaction was

stopped by adding 10 µl of 1% sodium dodecyl sulfate prepared in MQ and plates were

put at 4°C to avoid additional reaction. Absorbance of solution at 450 nm were

measured using a spectrophotometer (Thermo GENESYS 10S UV-Vis). Data were

expressed as relative to 0.25% DMSO control.

Immunohistochemistry

Formalin-fixed and paraffin-embedded surgical specimens were cut into 3 µm

sections. These sections were immersed in antigen retrieval solution (pH 9.0) (Nichirei

Bioscience Inc.) and heated in a pressure chamber. After that, the sections were

incubated with the primary antibodies (rabbit anti-ABCA1 pAb, NB400-105) and later

incubated with HRP-labeled goat anti-rabbit secondary antibodies (Nichirei Bioscience

Inc.). Immunoreactivity was visualized by using 3–3 Diaminobenzidine Dab Substrate

Kits (Nichirei Bioscience Inc.). Informed consent for experimental use of the samples was obtained from all patients in accordance with the ethical guidelines of Aichi Cancer Center, and the study was approved by the Institutional Review Board (Approval No. 2021-0-037). DAB signals were extracted from RGB images by color deconvolution, inverted, and quantified as the mean intensity of pixels exceeding a defined threshold using ImageJ/Fiji.

Tumor xenograft model

Four-week-old female athymic nude mice were purchased from Japan SLC and housed in a specific pathogen-free facility. All animal experiments were conducted in accordance with the Guidelines for Animal Experiments of Aichi Medical University and approved by the Institutional Animal Care and Use Committee (Approval No. 2024-58).

To initiate xenografting of 786-O, 2×10^6 cells in 200 μL DMEM were subcutaneously injected into the back of mice by using a 27 G needle. After 7 days, when the inoculated tumor became visible, the mice were randomly divided into treatment and control groups, and treatment was initiated (day 0). TMP-153 (50 mg/kg body weight) or vehicle (0.5% hydroxypropyl methylcellulose with 0.1% polysorbate 80) was intraperitoneally administered to each mouse on days 0, 4, 7, 11, 14, and 18 to each mouse. The tumor volume was measured every week and calculated using the modified ellipsoid formula ($1/2 \times$ length $\times$ width^2). The xenografting of TE-8 was performed with some modifications. For this, 1×10^7 cells were injected. A combination of TMP-153 and CsA (both at 10 mg/kg body weight) or vehicle was

intraperitoneally administered to each mouse on days 0 and 4. Finally, the tumor

volume was measured 36 days after the first administration.

Statistical analysis

Microsoft Excel for Microsoft 365 MSO 2306, GraphPad Prism v8.4.1 and Python

version 3.12.7 were used for analyses and displays of quantitative data. Data are

expressed as mean with each point or error bars representing standard deviations (SD).

Significant difference was indicated as follows: *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$;

****, $p<0.0001$.

Data Availability

All data generated or analysed during this study are included in the manuscript and

supporting files.

Acknowledgments

We thank all members of the Ikenouchi laboratory for helpful discussions, Takana

Motoyoshi (K.I. Stainer) for immunohistochemistry, Teruaki Fujishita and Masahiro

Aoki for technical advice on xenograft experiments, Minako Suzuki for technical

assistance and Kenji Kasai for pathological advice. This work was supported by

AMED-FORCE (21444781) (J.I.), JSPS KAKENHI [JP25H00994 (J.I.), JP25H01325

Author contributions

Declaration of interests

There are no competing interests to declare.

References

1.   Nieto MA, Huang RY, Jackson RA, & Thiery JP (2016) EMT: 2016. *Cell*
       166(1):21-45.
2.   Lambert AW & Weinberg RA (2021) Linking EMT programmes to normal and
       neoplastic epithelial stem cells. *Nat Rev Cancer* 21(5):325-338.
3.   Zheng X*, et al.* (2015) Epithelial-to-mesenchymal transition is dispensable for
       metastasis but induces chemoresistance in pancreatic cancer. *Nature*
       527(7579):525-530.
4.   Fischer KR*, et al.* (2015) Epithelial-to-mesenchymal transition is not required for
       lung metastasis but contributes to chemoresistance. *Nature* 527(7579):472-476.
5.   Padmanaban V*, et al.* (2019) E-cadherin is required for metastasis in multiple
       models of breast cancer. *Nature* 573(7774):439-444.

6. Davis FM, Stewart TA, Thompson EW, & Monteith GR (2014) Targeting EMT in cancer: opportunities for pharmacological intervention. *Trends in pharmacological sciences* 35(9):479-488.

7. Debaugnies M*, et al.* (2023) RHOJ controls EMT-associated resistance to chemotherapy. *Nature* 616(7955):168-175.

8. Makhov P*, et al.* (2018) Resistance to Systemic Therapies in Clear Cell Renal Cell Carcinoma: Mechanisms and Management Strategies. *Molecular cancer therapeutics* 17(7):1355-1364.

9. Cui Y*, et al.* (2020) Antitumor functions and mechanisms of nitidine chloride in human cancers. *Journal of Cancer* 11(5):1250-1256.

10. Fang Z*, et al.* (2014) Nitidine chloride induces apoptosis and inhibits tumor cell proliferation via suppressing ERK signaling pathway in renal cancer. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association* 66:210-216.

11. Iwasaki H*, et al.* (2010) Down-regulation of lipids transporter ABCA1 increases the cytotoxicity of nitidine. *Cancer chemotherapy and pharmacology* 66(5):953-959.

12. Cano A*, et al.* (2000) The transcription factor snail controls epithelial-mesenchymal transitions by repressing E-cadherin expression. *Nat Cell Biol* 2(2):76-83.

13. Batlle E*, et al.* (2000) The transcription factor snail is a repressor of E-cadherin gene expression in epithelial tumour cells. *Nat Cell Biol* 2(2):84-89.

14. Ikenouchi J, Matsuda M, Furuse M, & Tsukita S (2003) Regulation of tight junctions during the epithelium-mesenchyme transition: direct repression of the gene expression of claudins/occludin by Snail. *J Cell Sci* 116(Pt 10):1959-1967.

15. Arner EN, Du W, & Brekken RA (2019) Behind the Wheel of Epithelial Plasticity in KRAS-Driven Cancers. *Frontiers in oncology* 9:1049.

16. Ignatova ID, Angdisen J, Moran E, & Schulman IG (2013) Differential regulation of gene expression by LXRs in response to macrophage cholesterol loading. *Molecular endocrinology (Baltimore, Md.)* 27(7):1036-1047.

17. Frechin M*, et al.* (2015) Cell-intrinsic adaptation of lipid composition to local crowding drives social behaviour. *Nature* 523(7558):88-91.

18. Cserepes J*, et al.* (2004) Functional expression and characterization of the human ABCG1 and ABCG4 proteins: indications for heterodimerization. *Biochem Biophys Res Commun* 320(3):860-867.

19. Das A, Brown MS, Anderson DD, Goldstein JL, & Radhakrishnan A (2014) Three pools of plasma membrane cholesterol and their relation to cholesterol homeostasis. *Elife* 3.

20. van Meer G, Voelker DR, & Feigenson GW (2008) Membrane lipids: where they are and how they behave. *Nat Rev Mol Cell Biol* 9(2):112-124.

21. Szlasa W, Zendran I, Zalesińska A, Tarek M, & Kulbacka J (2020) Lipid composition of the cancer cell membrane. *Journal of bioenergetics and biomembranes* 52(5):321-342.

22. Tanaka T, *et al.* (2024) Fluorescent Solvatochromic Probes for Long-Term Imaging of Lipid Order in Living Cells. *Advanced science (Weinheim, Baden-Wurttemberg, Germany)* 11(17):e2309721.

23. Subczynski WK, Pasenkiewicz-Gierula M, Widomska J, Mainali L, & Raguz M (2017) High Cholesterol/Low Cholesterol: Effects in Biological Membranes: A Review. *Cell biochemistry and biophysics* 75(3-4):369-385.

24. Koivusalo M, Jansen M, Somerharju P, & Ikonen E (2007) Endocytic trafficking of sphingomyelin depends on its acyl chain length. *Mol Biol Cell* 18(12):5113-5123.

25. Levy M & Futerman AH (2010) Mammalian ceramide synthases. *IUBMB life* 62(5):347-356.

26. Sassa T & Kihara A (2014) Metabolism of very long-chain Fatty acids: genes and pathophysiology. *Biomolecules & therapeutics* 22(2):83-92.

27. Nishihira T, Hashimoto Y, Katayama M, Mori S, & Kuroki T (1993) Molecular and cellular features of esophageal cancer cells. *Journal of cancer research and clinical oncology* 119(8):441-449.

28. Song Y, Liu J, Zhao K, Gao L, & Zhao J (2021) Cholesterol-induced toxicity: An integrated view of the role of cholesterol in multiple diseases. *Cell metabolism* 33(10):1911-1925.

29. Lee O, Chang CC, Lee W, & Chang TY (1998) Immunodepletion experiments suggest that acyl-coenzyme A:cholesterol acyltransferase-1 (ACAT-1) protein plays a major catalytic role in adult human liver, adrenal gland, macrophages, and kidney, but not in intestines. *J Lipid Res* 39(8):1722-1727.

30. Aiello NM, *et al.* (2018) EMT Subtype Influences Epithelial Plasticity and Mode of Cell Migration. *Dev Cell* 45(6):681-695.e684.

31. Pastushenko I, *et al.* (2018) Identification of the tumour transition states occurring during EMT. *Nature* 556(7702):463-468.

32. Wang W, *et al.* (2021) ABCA1 is associated with the development of acquired chemotherapy resistance and predicts poor ovarian cancer outcome. *Cancer drug resistance (Alhambra, Calif.)* 4(2):485-502.

33. Schlaepfer IR, *et al.* (2012) Progestin modulates the lipid profile and sensitivity of breast cancer cells to docetaxel. *Molecular and cellular endocrinology* 363(1-2):111-121.

34. Cotte AK, *et al.* (2018) Lysophosphatidylcholine acyltransferase 2-mediated lipid droplet production supports colorectal cancer chemoresistance. *Nat Commun* 9(1):322.

35. Hultsch S, *et al.* (2018) Association of tamoxifen resistance and lipid reprogramming in breast cancer. *BMC cancer* 18(1):850.

36. Pavlova NN, Zhu J, & Thompson CB (2022) The hallmarks of cancer metabolism: Still emerging. *Cell metabolism* 34(3):355-377.

37. Shigetomi K, Ono Y, Inai T, & Ikenouchi J (2018) Adherens junctions influence tight junction formation via changes in membrane lipid composition. *J Cell Biol* 217(7):2373-2381.

38. Shigetomi K, Ono Y, Matsuzawa K, & Ikenouchi J (2023) Cholesterol-rich domain formation mediated by ZO proteins is essential for tight junction formation. *Proc Natl Acad Sci U S A* 120(8):e2217561120.

39. Ikenouchi J, *et al.* (2012) Lipid polarity is maintained in absence of tight junctions. *J Biol Chem* 287(12):9525-9533.

40. McNally BD, *et al.* (2022) Long-chain ceramides are cell non-autonomous signals linking lipotoxicity to endoplasmic reticulum stress in skeletal muscle. *Nat Commun* 13(1):1748.

41. Yang H, *et al.* (2010) Vitamin A deficiency results in dysregulation of lipid efflux pathway in rat kidney. *Pediatric nephrology (Berlin, Germany)* 25(8):1435-1444.

Figure legends

Figure 1

Ectopic expression of Snail confers epithelial cells nitidine chloride resistance via induction of ABCA1 expression

(**A**) Enhancement of growth inhibitory effect of nitidine chloride (NC) on human renal carcinoma cells by co-treatment with an ABCA1 inhibitor CsA. Cells were grown with or without 20 μM NC and 10 μM CsA for 24 h, and cell number was determined with CCK-8. Fold change in cell number by NC was shown as relative to controls without NC. Results are shown as mean of at least four independent experiments ± SD (Tukey-Kramer's multiple comparison test).

(**B-E**) Immunoblot analysis of whole cell lysate of human renal carcinoma cell lines. A representative result from three independent experiments is shown (B). Quantitative analyses of Snail (C) and ABCA1 (D) expression are presented. A scatter plot illustrating the correlation between ABCA1 and Snail expression levels is also shown (E). Quantitative data are shown as mean of at least three independent experiments ± SD (Tukey-Kramer's multiple comparison test).

(**F**) Comparison of ABCA1 expression between normal tissue and primary tumor of indicated subtypes of renal cancers analyzed using UCSC Xena. Bars indicate means (Welch's *t*-test).

(**G**) Immunohistochemistry of a surgically extracted renal tissue from a patient with Fuhrman grade 3 primary ccRCC, indicating upregulation of ABCA1 in the lesion site. The ABCA1 staining was verified by the accumulation of known signals in renal tubules and glomeruli in normal tissue (41). Scale bars, 250 μm (left), 50 μm (right).

(**H**) Quantification of ABCA1 signal from surgically extracted renal tissues from three patients for each Furhman grade (Tukey-Kramer's test; see also Figure1—figure supplement 2).

(**I**) Phase contrast images of EpH4 wild-type cells (top) and Snail-overexpression cells

(bottom, EpH4-Snail). Scale bars, 50 µm.

(**J**) Immunofluorescence images of EpH4 and EpH4-Snail cells. The cells were fixed

Scale bars, 20 µm.

(**K**) Acquisition of NC resistance by exogenous expression of Snail in EpH4 cells and

enhancement of 20 µM NC effect on EpH4-Snail cells by co-treatment with 10 µM CsA.

Data are presented as in Figure 1A (Tukey-Kramer's multiple comparison test).

(**L**) Immunoblot analyses of whole cell lysates of EpH4, EpH4-Snail, E-cadherin KO

and α-catenin KO EpH4 cells.

(**M**) Immunofluorescence images of EpH4 and EpH4-Snail cells. Cells were fixed with

MeOH. Scale bars, 20 µm.

(**N,O**) Immunoblot analyses of whole cell lysates of MDCK II cells (N) expressing

KRAS G12V treated with 5 µg/ml TGFβ for 0, 3 and 8 days and human esophageal

carcinoma cell lines TE-15 and TE-8 (O). Graphs on bottom shows analyses from three

biological replicates (Tukey's multiple comparison test (N) and Student's *t*-test (O)).

Phase contrast images are also shown on top (O). Scale bars, 50 µm.

Figure 1—Figure supplement 1

Cancer types that exhibit tumor specific upregulation of ABCA1

transcription

(**A-C**) Comparison of mRNA transcription of ABCA1 (A,B) and Snail (C) in different

cancer types. Expression data from GDC TCGA database were analyzed using USCS

Xena platform (Welch's *t*-test).

 Figure 1—Figure supplement 2

 Comparison of ABCA1 expression in different progression grades of

 kidney cancer cases

Immunohistochemistry of surgically extracted renal tissues from patients with Fuhrman

grade 13 primary ccRCC, indicating elevation of ABCA1 signal at the lesion site of

higher-grade ccRCC. All samples were collected from independent patients. The

ABCA1 staining was verified by the accumulation of known signals in renal tubules and

glomeruli in normal tissues (41). Scale bars, 100 μm.

Figure 2

Alteration of cellular cholesterol distribution induces ABCA1 expression in

hybrid E/M cells

(**A**) Immunofluorescence images of cells stained with the LXRα+β antibody. Scale bars,

10 μm. Cells were fixed with 4% PFA. Comparison of the abundance of LXR is also

shown (bottom). LXR signal intensity overlapped with DAPI signals was determined

for three independent fields of view (Student's t-test).

(**B**) Immunoblot analyses of whole cell lysates of EpH4-Snail cells treated with 1 μM

GSK-2033 for 24 h ($n = 4$, Student's t-test).

(**C**) Enhancement of growth inhibitory effect of nitidine chloride (NC) by co-treatment

with an LXR inhibitor GSK2033. Cells were treated with or without 1 μM GSK2033 for

24 h, then cultured with or without 20 μM NC and 1 μM GSK2033 for 24 h, and cell

number was determined with CCK-8. Fold change in cell number by NC was shown as

relative to controls without NC (Student's t-test).

(**D**) Immunoblot analyses of whole cell lysates of EpH4-Snail cells treated with

indicated concentration of simvastatin or 0.1% DMSO ('0' μM) in DMEM

supplemented with 10% normal or lipid-free fetal bovine serum (FBS) for 48 h ($n = 3$,

Tukey-Kramer's test).

(**E**) Subcellular distribution of cholesterol stained with filipin. Scale bar, 10 μm.

Intracellular signal to surface signal ratio was determined for three independent fields of

view (Student's t-test).

(**F**) Live cell images of lipid droplets (LDs) visualized by Lipi-Green dye.

(**G**) Quantification of LD signal per cell. The Lipi-Green signals of all optical sections

were summed and divided by the number of cells counted using DAPI images. Shown

are means of three different fields of view ±SD.

(**H**) Subcellular distribution of Topfluor-(TF-)cholesterol. Living cells were incubated

with 5 μg/ml TF-cholesterol for 1 h, then stained with Lipi-Deepred and Nuc Blue.

Scale bar, 20 μm.

(**I**) Observation of filipin in EpH4-Snail cells expressing either EGFP-ADRP (lipid

droplets) or mRFP-LAMP1 (lysosome). Scale bar, 20 μm.

(**J**) Comparison of cholesterol content relative to phospholipid (Student's *t*-test).

(**K,L**) Comparison of cell surface cholesterol (D4) and sphingomyelin (lysenin).

Wild-type EpH4 cells expressing H2B-iRFP and EpH4-Snail cells were co-cultured and

stained with recombinant lipid-binding proteins and CellMask Green (K).

Quantification of fluorescent signals of lipid binding probes (L). Signal intensities per

cell, relative to those of wild-type cells, were quantified and analyzed (Student's *t*-test).

(**M−O**) Observation of lipid order of plasma membrane using FπCM-SO₃. See also

Figure 2—figure supplement 2. Wild-type EpH4 cells expressing H2B-iRFP and

EpH4-Snail cells were co-cultured and stained with 1 μM FπCM-SO₃ for 10 min. The

hue (Red/Blue ratio) – brightness (intensity) images of FπCM-SO₃ are shown with the

confocal image of H2B-iRFP (magenta) (M). Normalized histogram of Red/Blue ratio

values within FπCM-SO₃ positive pixels (N). Comparison of mean Red/Blue ratio

values (P) (Student's *t*-test).

Figure 2—Figure supplement 1

Contribution of transcription factors involved in ABCA1 transcription in

epithelial cells

(**A**) Immunoblot analysis of whole cell lysates of EpH4-Snail (without treatment) and

EpH4 cells treated with inhibitors of PI3K-Akt pathway (5 μM GSK2334470: PDK1,

50 μM LY294002: PI3K, or 1 μM Wortmannin: PI3K for 24 h). Comparison of protein

levels of ABCA1 and phospho-FoxO3a is also shown (Tukey-Kramer's test).

(**B**) Immunoblot analysis of whole cell lysates of EpH4 and EpH4-Snail cells (Student's

t-test). Quantitative analysis of c-myc expression is also shown.

Figure 2—Figure supplement 2

Effect of ABCA1 knockout on nitidine chloride resistance of EpH4-Snail

cells

(**A**) Immunoblot analysis of whole cell lysates of wild-type and two lines of ABCA1

KO EpH4-Snail cells (#1 and #2).

(**B**) Effect of ABCA1 knockout on growth inhibitory effect of nitidine chloride. Cells

were grown with or without 20 μM NC for 24 h, and cell number was determined with

CCK-8. Fold change in cell number by NC was shown as relative to controls without

NC (Tukey-Kramer's test).

Figure 2—Figure supplement 3

Visualization and analyses of lipid order using FπCM-SO₃

(**A**) Schematic illustrating the alteration of fluorescence spectra of solvatochromic dyes

in lipid bilayers with different lipid compositions. A high content of sphingolipids and

cholesterol induces an ordered acyl chain orientation. The dyes exhibit red-shifted

fluorescence in disordered membranes and blue-shifted fluorescence in ordered

membranes.

(B) Schematic showing lipid-order visualization and analysis using FπCM-SO₃. Living

cells cultured on glass-bottom dishes were stained with 1 μM FπCM-SO₃ for 10 min.

The dye was excited with a 405 nm laser, and emission signals of 400–500 nm (Blue)

and 500–700 nm (Red) were separated using a variable dichroic mirror and

simultaneously detected by two photomultipliers. The raw images were

Gaussian-blurred, Red/Blue ratio and mean intensity of both channels were calculated.

For visualization of lipid order, a ratio-intensity image was generated in which the hue

represents the Red/Blue ratio and the brightness corresponds to the signal intensity

(bottom left). For quantitative comparison, a threshold was set based on the

fluorescence intensity of FπCM-SO₃, and histograms of the Red/Blue ratio were

generated from positive pixels (bottom center). For statistical assessment, mean

Red/Blue ratio values of positive pixels were obtained and compared (bottom right).

Figure 3

Effects of Snail-induced EMT on sphingomyelin profile

(**A**) Comparison of sphingomyelin (SM) content relative to phospholipid content.

(**B**) Comparison of cholesterol (Chol)/SM ratio between EpH4 and EpH4-Snail cells.

(**C**) Fatty acid composition of SM determined using LC-MS. All peaks corresponding to d18:1-even fatty acid-SMs were included in the analysis and each peak value is expressed as a percentage. n.d., peak not detected.

(**D**) Pie charts of SM chain length profile from c. SMs were classified into long-chain fatty acid (LCFA, orange) ($\leq$20) and very long-chain fatty acid (VLCFA, purple) (>20) SMs.

(**E**) Chain length specificity of Elovls and CerSs responsible for fatty acid profile of sphingolipids reported (25, 26). Elovls catalyze a rate-limiting step of fatty acid elongation cycle, and CerSs catalyze *N*-acylation of sphingoid bases to produce ceramides.

(**F**) Comparison of transcription levels of *ELOVL*s and *CERS*s. Notable decreases in expressions of *ELOVL7* and *CERS3* in EpH4-Snail cells are indicated by asterisks.

(**G**) Quantification of transcription levels of *ELOVL7* and *CERS3* relative to that of *ZO-1*, whose expression does not change between EpH4 and EpH4-Snail (14).

(Student's *t*-test).

(**H**) Effect of supplementation of C22:0 ceramide (Cer22:0) on ABCA1 expression. EpH4-Snail cells were treated with 10 μM ceramide or 0.4% vehicle (dodecane/ethanol = 2/98) for 24 h, then whole cell lysates were analyzed by immunoblot.

(**I**) Immunoblot analyses of whole cell lysates of mouse epithelial (MTD-1A, CSG) and fibroblast (L-929, NIH3T3) cells.

(**J**) Comparison of Chol/SM ratio of EpH4, EpH4-Snail and normal fibroblasts (Dunnett's test).

(**K**) Comparison of Chol/SM ratio between TE-15 and TE-8 (Student's *t*-test).

(**L**) Scatter plot of Chol content against SM content used in B, H and I. Dashed lines indicate Chol/SM = 1.0 (blue) and 1.5 (red). The region of Chol/SM > 1.5 is shown with red background.

(**M**) Possible mechanism of induction of ABCA1 expression and lipid droplet enlargement through decrease in VLCFA-SM. In EpH4 (epithelial) cells, a certain amount of choleseterol is sequestered through interaction with SM, organizing ordered plasma membrane (PM). In EpH4-Snail (hybrid E/M) cells, the lower level of VLCFA-SM biosynthesis leads to a decrease in plasma membrane SM content, resulting in membrane disorder. The increase in SM-unbound form of cholesterol in PM induces its internalization and activation of cellular cholesterol sensors, resulting in to lipid droplet enlargement and enhanced ABCA1 expression.

Figure 3—Figure supplement 1

Comparison of SM and cholesterol content

(**A,B**) Comparison among EpH4, EpH4-Snail, and fibroblasts.

(**C,D**) Comparison between esophageal cancer cell lines TE-15 (Snail-negative) and

TE-8 (Snail-positive). $n = 3$, ** $p < 0.01$, * $p < 0.05$, n.s., $p \geq 0.05$ by Tukey's multiple

comparison test (A,B) and Student's t-test (C,D).

Figure 4

Selective growth inhibition of Snail-positive cells by an ACAT inhibitor

TMP-153

(**A**) Schematic illustrating the molecular mechanisms for handling excess cholesterol to

avoid cell death. Cells eliminate excess cholesterol through efflux mediated by

transporters including ABCA1 or isolation into lipid droplets (LD) via esterification of

cholesterol by acyl-CoA: cholesterol acyltransferases (ACATs) to avoid cell death. CsA

and TMP-153 inhibit ABCA1 and ACATs, respectively.

(**B**) Immunoblot analysis of whole cell lysates of human kidney cell lines. Quantitative

analysis of SOAT1 expression is also shown.

(**C-F**), Effects of treatments with CsA (C,E) or TMP-153 (D,F) to cellular growth of

EpH4 and EpH4-Snail (C,D) or human kidney cancer cell lines (E,F). Cells were treated

with drugs for 24 h and relative cell number to 0.1% DMSO control was determined by

CCK-8. Results are shown as mean of at least three independent experiments ± SD.

(**G-I**) TMP-153 inhibits the growth of tumor xenografts composed of Snail-positive

renal cancer 786-O cells. Experiments were performed as described in Materials and

Methods. In brief, 50 mg/kg TMP-153 was administered intraperitoneally to nude mice

bearing 786-O tumor xenografts. After 21 days of the first administration, the overall

appearance (G) and the excised tumors (H) of control and treated mice were compared.

(I) Tumor growth curve in the xenograft model. Results are shown as mean of four

biological replicates ± SD (Student's *t*-test).

Figure 4—Figure supplement 1

Selective growth inhibition of a Snail positive esophageal cancer cell line

by combined treatment of CsA and TMP-153

(**A-C**) Effects of treatments with CsA (**a**), TMP-153 (**b**) or combination of 50 μM CsA

and indicated concentration of TMP-153 (**c**) to cellular growth of human esophageal

cancer cell lines TE-15 and TE-8. Cells were treated with drugs for 24 h and relative

cell number to 0.1% DMSO control was determined by CCK-8. n ≥ 3, error bars, SDs.

(one-sided Student's *t*-test).

(**D-F**)**,** Combined treatment of CsA and TMP-153 inhibits the growth of TE-8 tumor

xenografts. Experiments were performed as described in Materials and Methods. In

brief, 10 mg/kg CsA and 10 mg/kg TMP-153 were simultaneously administered

intraperitoneally to nude mice bearing TE-8 tumor xenografts. After 36 days of the first

administration, the overall appearance (D), the excised tumors (E), and the tumor size

(F) of control and treated mice were compared. *n* = 5, error bars, SDs (Student's *t*-test).

Figure 1

Figure 2

Figure 3

Figure 4