1 **Wag31, a membrane tether, is crucial for lipid homeostasis in mycobacteria**

2 Yogita Kapoor[1,2], Himani Khurana[3#], Debatri Dutta[1#], Arnab Chakraborty[3], Anshu Priya[2,4], Archana

3 Singh[2,4], Siddhesh S. Kamat[3], Neeraj Dhar,[5,6,] Thomas J. Pucadyil[3], Vinay Kumar Nandicoori[1,2,7]

    1. CSIR-Centre for Cellular and Molecular Biology (CSIR-CCMB), Hyderabad, 500007
    2. AcSIR, Ghaziabad-201002
    3. Indian Institute of Science Education and Research, Pune, 411008
    4. CSIR-Institute of Genomics and Integrative Biology (CSIR-IGIB), New Delhi, 110025
    5. Vaccine and Infectious Disease Organisation, University of Saskatchewan, Canada, S7N 5E3
    6. Department of Biochemistry, Microbiology and Immunology, University of Saskatchewan, Canada S7N 5E5
    7. National Institute of Immunology, New Delhi 110067
    #   contributed equally to this work

    Address correspondence to vinaykn@ccmb.res.in, vinaykn@nii.ac.in

Keywords: Wag31, intracellular lipid inclusions (ILIs), lipid homeostasis, membrane-tether

Abstract

The mycobacterial cytoskeletal protein Wag31 is necessary for maintaining cell shape and directing
cellular growth and elongation. Wag31 has a characteristic N-terminal DivIVA-domain and a C-terminal
coiled-coil domain. While the role of Wag31 in polar elongation is known, there is limited mechanistic
insight on how it orchestrates growth and elongation. In this report, we delineate roles of the N- and C-
terminal domains of Wag31 using genetics, state-of-the-art multi-omics, biochemical, and imaging
approaches. We show that Wag31 predominantly interacts with several membrane-associated proteins
involved in lipid metabolism, cell wall synthesis and division. Native levels of Wag31 are critical for the
maintenance and distribution of membrane lipids. Both depletion and overexpression of Wag31
perturbs lipid homeostasis, leading to the formation of intracellular lipid inclusions (ILIs). Protein-lipid
crosslinking and imaging studies reveal that purified Wag31 can bind and effectively tether Cardiolipin
(CL)-containing liposomes. We further show that the tethering activity lies in the DivIVA-domain
containing N-terminal of Wag31 while the C-terminal mediates protein-protein interactions of Wag31.
Despite retaining its ability to interact with partner proteins, the DivIVA domain-deleted Wag31 mutant
shows defects in liposome tethering *in vitro* and non-polar localization of CL *in vivo*, which eventually
causes lethality. Our study suggests that membrane tethering 'licenses' Wag31 to form scaffolds that
help orchestrate protein-lipid and protein-protein interactions necessary for mycobacterial growth and
survival.

Introduction

Cell division is a critical event in the life of a bacterium as it is the prime mode of bacterial growth and reproduction. Hence, the process of their division is tightly knitted. The key process in bacterial cell division, i.e., assembly of the FtsZ ring at the mid-cell (1), is tightly regulated by the Min and Nucleoid occlusion systems (2-6). Mycobacterium, which comprises clinically relevant pathogens-*Mycobacterium tuberculosis (Mtb)*, *Mycobacterium leprae*, and *Mycobacterium abscessus* lacks both the regulators of FtsZ ring placement (7). Instead, they encode for a DivIVA protein, usually encoded by Firmicutes and other Actinobacteria (8-11). It is necessary that the bacterial shape, crucial for normal cellular physiology, is maintained during cell division. Rod-shaped bacteria such as *Escherichia coli* and *Bacillus subtilis* encode MreB and multiple MreB-like proteins that sustain the rod morphology of cells (12). *Mycobacterial spp.* lack MreB homologs (13, 14). The DivIVA protein of *Mtb* called Wag31 is known to regulate both cellular growth and morphology (10, 15), which makes it an exciting candidate to investigate.

DivIVA proteins were discovered three decades ago during the study of cell division mutants of *B. subtilis* (16). DivIVA proteins are rich in coiled-coil domains (17-21), which enable them to form higher oligomeric structures and, hence, scaffolds, making them suitable adaptor proteins (22, 23). They are known to play diverse roles in various species (24), e.g., they play dual roles of septum positioning and chromosome partitioning in *B. subtilis* (9, 25); cell-separation/resolution post-cell-division in *Listeria monocytogenes* (26, 27); pole formation and maturation in *Streptococcus spp.* (Fadda, 2007) In *Mycobacterium spp.*, Wag31 is an essential protein that senses negative membrane curvature found at poles and septum and localizes there (28). It plays roles in cellular elongation and septation in concert with other divisome and elongasome components (10, 15, 29). It is specifically found in higher amounts at the old pole, where it directs polar elongation by recruiting the acyl CoA carboxylase (ACCase) complex involved in fatty acid, mycolic acid, and methyl-branched chain lipid precursor synthesis (28, 30-32). It localizes to the septum after cytokinesis is over, perhaps for septal synthesis via its interactions with FtsI (32-34), though the process remains largely unknown. Consequently, loss of Wag31 leads to cell division defects- loss of polar elongation, which renders the cells 'round' in shape that occurs due to affected peptidoglycan (PG) synthesis (10, 15, 28). Though Wag31 participates in coordinating PG metabolism, the mechanism or the protein network by which it does so remains unknown.

Wag31 has also been associated with the maintenance of the intracellular membrane domain (IMD), a specialized compartment for lipid synthesis, localized predominantly at the old pole in consonance with pole elongation and spottily all around the plasma membrane where it may function to repair membrane defects (35-37). The loss of Wag31 leads to the delocalization of IMD and its protein markers, such as MurG and GlfT2, but it is not understood whether Wag31 plays a direct role in membrane-partitioning or it is just a consequence of localized membrane synthesis (34, 37). The role of Wag31in PG and lipid synthesis that drives cell division remains unclear. Despite numerous studies, the details of how Wag31 orchestrates mega-complexes at the pole or partitions the membrane into two to govern cellular elongation remain elusive.

Here, we endeavoured to address the lacunae in the functionality of Wag31 by asking the

following questions- i) what are the ultrastructural changes that occur in mycobacteria due to the

absence of Wag31, ii) do Wag31 deficient cells undergo transcriptional rewiring to adjust to cell division

defects, iii) is the change in cellular morphology from 'rod-to-round' accompanied by altered lipid levels,

iv) what is the interactome of Wag31, v) what are the functions of its N and C-terminal domains and

finally vi) what is its relationship with the mycobacterial membrane. The present study shows that

Wag31 levels are critical for maintaining lipid homeostasis in mycobacteria. Our results demonstrate

that the N- and C-terminals of Wag31 have distinct functions. The N-terminal of Wag31 binds and

tethers the membrane together while the C-terminal interacts with various proteins. Together, the

domains help Wag31 to orchestrate membrane-localized processes essential for maintaining normal

cellular morphology and physiology.

 Results

Loss of Wag31 leads to the formation of intracellular lipid inclusions (ILIs)

In the pursuit of bridging the knowledge gap existing in the functionality of Wag31 (Fig. 1a), we
generated a *Wag31* conditional mutant in *Msm*, by integrating a copy of *wag31* under tetracycline (ATc-
tet off) regulation at the L5 locus and replacing the native copy with a hygromycin cassette (Fig. S1a).
The mutant, *Δwag31*, thus generated, was confirmed by PCRs (Fig. S1b) and Western blotting (Fig.
1b). The lack of Wag31 led to a drop in the turbidity of culture (Fig. 1c). Analysis of bacillary survival *in*
vitro by enumerating CFUs 12h post-ATc addition, revealed a 2 log fold difference in the survival of
Δwag31 (Fig. 1d). Given the essential role of Wag31 in causing polar growth of the cell, we investigated
the effect of depletion of Wag31 on bacterial cellular morphology using electron microscopy (EM).
Scanning EM (SEM) suggested that the absence of Wag31 caused a transition in the morphology of
the cells from rod to round, as previously documented (10, 15, 38). Apart from round morphology (~36
%), we also observed a rod-to-round transformation stage in which cells were bulged at one pole with
the other pole intact (Fig. 1e-g). To obtain detailed insights, we performed transmission EM (TEM),
which divulges information on the internal structure of cells. We discovered spherical, electron-lucent
bodies (indicated by an arrowhead in Fig.1h) in the cytosol of all the samples. While *Msm* and *Δwag31*-
ATc harbored a few (~2-3) electron-lucent bodies, their numbers were drastically higher in the case of
Δwag31+ATc (Fig. 1h). The cytosol of *Δwag31*+ATc was filled with multiple variable-sized bodies which
resembled Lipid bodies that *Mycobacterium tuberculosis* (*Mtb)* accumulates inside the host (39, 40). As
the term 'lipid bodies/lipid droplets' is used specifically for the lipids acquired by *Mtb* from the host cell,
we referred to them as intracellular lipid inclusions (ILIs). We categorized the cells based on the number
of ILIs they contained. Their distribution remained reasonably similar within *Msm* and *Δwag31*-ATc cells
for all the classes (Fig. 1i). Whereas *Δwag31* treated with ATc had as high as 85% cells harboring more
than 5 ILIs and very few cells with less than 5 ILIs (Fig. 1i). Our data, therefore, reports a novel
observation of lipid accumulation when Wag31 is depleted from mycobacteria.

Wag31 levels influence lipid homeostasis

Next, we set out to analyze if the ILIs are indeed lipidic in nature by staining cells with BODIPY
(493/503), a dye that preferentially partitions into neutral lipid droplets (41, 42). We stained either
untreated or ATc-treated *Msm* and *Δwag31* with BODIPY and visualized them on a microscope. ATc-
treated *Δwag31* cells displayed the highest level of BODIPY fluorescence, indicating the maximum
abundance of lipid bodies among the three strains (Fig. 2a, c). Further, we examined whether the
appearance of ILIs is associated just with the loss of Wag31 or is a consequence of deviations from the
native expression level of Wag31. Towards this, we overexpressed Wag31 episomally using an
isovaleronitrile (IVN) inducible system (*Msm::wag31*) (Fig. S2a) and stained the cells with BODIPY.
Unlike Wag31 depletion, which resulted in spherical cells (Fig. 1e), Wag31 overexpression resulted in
the bulging of cells from one pole (Fig. 2b). Additionally, the overexpression of Wag31 also led to lipid
aggregates in the cell with a few larger 'donut-like' aggregates in the bulged pole (Fig 2b,d).

Lipid accumulation in both cases led us to inspect whether lipid metabolism is closely tied to
Wag31 expression levels in mycobacteria. To investigate that possibility, total cellular lipids were
extracted from *Msm,* Wag31 depletion, and overexpression conditions and subjected to semi-

quantitative LC-MS analysis (illustrated in Fig. 2e). Lipidomic analysis of *Δwag31*+ATc revealed an imbalance in the levels of neutral lipids, phospholipids, and free fatty acids in comparison to *Msm*-ATc (Fig. 2f-g). *Δwag31*+ATc cells had a significant accumulation of unsaturated diacylglycerols (~3-5 fold) and a very long chain containing (>C36) phosphatidylethanolamines (~5-32 fold). This was accompanied by increased levels of different chain lengths (C18-C24) of free fatty acids (~1.7-4 fold) and reduced levels of cardiolipins C70:0 & C70:1 (~3 fold). Wag31 overexpression, on the other hand, led to a 2-fold increase in the level of different species of alpha-mycolic acids (C74-C78) as compared to *Msm*+IVN (Fig. 2h), while other tested lipids remained unchanged (Fig. S2). Altogether, the results presented above demonstrate that Wag31 levels are crucial for maintaining lipid homeostasis in mycobacteria, the depletion and overexpression of which impact lipid metabolism differently.

Molecular interactions of Wag31 with membrane-associated proteins govern cellular homeostasis.

We reasoned that Wag31, an adaptor protein, might maintain lipid homeostasis by interacting with proteins involved in these pathways. To examine the protein interaction network of Wag31, we performed an interactome analysis by utilizing *Δwag31*, which expresses FLAG-Wag31 integrated at the L5 locus, and used FLAG-GFP as the control to differentiate from non-specific interactions. FLAG-Wag31 and FLAG-GFP from biological triplicates were immunoprecipitated using whole cell lysates from the log phase cells and subjected to mass spectrometry analysis (Fig. S3a & Table S2). This study identified both direct and indirect protein-partners of Wag31. The proteins identified were categorized based on their functions. It was found that Wag31 interacts with proteins involved in cell-wall elongation and division, lipid metabolism, intermediary metabolism and respiration and information pathways (translation, replication) (Fig. 3a, Fig. S4). Interactions of Wag31 with AccA3 and AccD5, subunits of the ACCase complex, and Rne or RNaseE (Msm4626), a membrane-associated protein, have been reported previously (28, 43). Subunits of ACCase complex and Rne were among the top 5 hits in our MS/MS data, validating our approach (Fig. 3a, marked with asterisk). The classification of the top 20 hits is represented in Fig. 3a. The highest number of protein interactors, expectedly, fell in the cell wall and cell processes category (Fig. 3a). For the present study, we focussed on the interactors found in the cell wall elongation and cell division and lipid metabolism functional categories to probe the cause behind the lipid dysbiosis phenotype.

MurG, involved in PG metabolism (36, 44-46), and SepIVA, involved in regulating septation and the spatial regulator of MurG (47) also surfaced in the interactome of Wag31. Msm1813, or acyl CoA carboxylase (ACAD), crucial for nascent fatty acid precursor synthesis, was also enriched in the interactome. Other lipid metabolism interactors of Wag31 included acyl-CoA dehydrogenase (Msm2080), involved in lipid catabolism; GpsA, involved in phospholipid biosynthesis; and Msm0372, involved in fatty acid biosynthesis. To validate the high-throughput data, we chose MurG, SepIVA and Msm2092 and performed pull-down assays with His-tagged Wag31. MmpL4 and MmpS5 which didn't appear in the interactome database were used as negative controls in this experiment. His-tagged Wag31 (Fig. 3b) was incubated with *Msm* lysates expressing 3X-FLAG tagged versions of the interactors and non-interactors mentioned above (Fig. 3c) and pulled-down with the help of cobalt beads followed by a western blot analysis. Results obtained confirmed the interaction of Wag31 with

MurG, SepIVA and Msm2092, but not with the MmpL4 and MmpS5, thus validating our data (Fig.
3d).

Since Wag31 is a spatiotemporal regulator of other proteins, we were intrigued to discover the
localisation of its interacting partners. We used the data curated by Zhu et al. (48) and determined that
Wag31 majorly interacts with membrane-associated proteins with a few exceptions (Fig. 3e, Fig. S3b).
The mycobacterial plasma membrane has been recently shown to have a cell wall-associated
compartment called PM-CW and a cell wall-free compartment called PM_f or IMD, which is the site for
lipid synthesis. Based on the available data (36, 48, 49), we found that Wag31 interacts more with IMD-
proteins than with PM-CW proteins (Fig. 3f.) In accordance with assigned roles (36), PM-CW residing
proteins in the interactome are involved in cell wall and cell processes and intermediary metabolism
and respiration. Similarly, IMD-homing proteins clustered in the lipid metabolism and cell wall synthesis
category. These results suggest that Wag31 maintains lipid homeostasis and cellular morphology
primarily through its interactions with the membrane proteins participating in such pathways.

Wag31 binds and tethers cardiolipin-containing membranes

Wag31 is a cytoplasmic protein that recognizes negative curvature regions in the cell, i.e., poles
and septum (28). It also forms higher-order structural assemblies at the poles (21, 50). However, the
precise mechanism by which Wag31 assemblies localize to the poles remains elusive. The predominant
interactions of Wag31 with membrane-associated proteins (Fig. 3) were intriguing and led us to
investigate if it can directly bind to membranes. We reasoned that such membrane binding activity might
render it capable of acting as a scaffold that anchors and organizes other proteins, thereby modulating
their functions. To test this, we performed a lipid crosslinking-based PLiMAP assay (Fig. 4a). Purified
Wag31 was incubated with ~100 nm extruded liposomes containing each of the abundant classes of
bacterial lipids phosphatidylethanolamine (PE), phosphatidylglycerol (PG), or cardiolipin (CL).
Liposomes contained trace amounts of a reactive fluorescent lipid (RFL), which crosslinks with
membrane-bound proteins upon photoactivation. These assays revealed that Wag31 preferentially
binds CL- but not PG- or PE-containing liposomes (Fig. 4b). Both CL and PG are anionic lipids, and the
preferential binding of Wag31 to CL indicates that the binding cannot be attributed to a charge-based
interaction. Remarkably, the sensitive fluorescence-based detection revealed the presence of SDS-
resistant higher-order oligomers of membrane-bound Wag31 (Fig. 4b).

Wag31 is a homolog of the DivIVA protein from *B. subtilis.* The latter has been shown to bind
to the membrane via the N-terminal DivIVA domain (N-DivIVA$_{BS}$) (51). However, the functional
significance of the N- & C- terminal domains of mycobacterial Wag31 has not been dissected.
Considering that the Wag31 N-terminal region shares 42% sequence identity with N-DivIVA$_{BS}$ (50), we
sought to inspect the effect of disruption of a particular residue (K20) in the N-terminus and the deletion
of the entire N-terminus, encompassing the DivIVA domain (1-60 amino acids) on membrane binding
(Fig. S5a). The K20 residue lies within a positively charged patch, which likely constitutes a membrane-
associating surface in the intertwined loop region of Wag31(50). Moreover, mutating the K20 residue
has been shown to decrease the cell length and elongation rate by 18% and 22%, respectively (34).
The K20A mutation showed a marginal defect in binding to CL-liposomes (Fig. 4c-d), indicating that
K20 alone is insufficient for membrane binding. However, Wag31$_{K20A}$ showed a marked defect in

forming SDS-resistant higher-order oligomers, as seen in the binding assays. Interestingly, deletion of the entire N-terminal domain caused a significant defect in both membrane binding and forming SDS-resistant higher-order oligomers (Fig. 4c-d). Surprisingly, binding was not entirely abolished, suggesting that the C-terminus of Wag31 may also contribute to membrane binding. Taken together, our results indicate that Wag31 preferentially binds CL-containing membranes and that regions other than the N-terminal domain contribute to membrane binding. The latter feature distinguishes Wag31 from DivIVA$_{BS}$, and Wag31 likely contains membrane binding sites located at both the N- and C-terminal regions, which might lead to differences in the manner in which it interacts with the membrane and partner proteins compared to DivIVA$_{BS}$.

To further confirm membrane binding, we imaged 100 nm-extruded CL-containing liposomes with a membrane dye DiI in the presence of Wag31. We used a GFP-tagged Wag31 mixed with an untagged Wag31 at a 1:10 molar ratio for these experiments. In the absence of Wag31-GFP, liposomes appeared as faint but distinct fluorescent puncta (Fig. 4e, first panel). Surprisingly, in reactions containing Wag31-GFP, we observed liposomes clustered into bright fluorescent puncta that were positive for Wag31-GFP (Fig. 4e, remaining panels). The Wag31 tetramer has been modelled as a long rod-shaped protein filament (21, 50) and such clustering of liposomes falls in congruence with the model. It indicates that these long filaments of Wag31 act to tether liposomes. We then systematically evaluated this effect by testing liposomes of different compositions and Wag31 mutants. Again, in the absence of Wag31, both PE- and CL-containing liposomes appeared as faint but distinct fluorescent puncta (Fig. 4f). The addition of Wag31 caused clustering of CL-containing liposomes into bright foci (Fig. 4f), which is quantified by a particle intensity analysis (Fig. 4g). This clustering was caused by membrane binding of Wag31 because PE-containing liposomes that do not bind Wag31 (Fig. 4b) showed no clustering (Fig. 4f,g). As expected, Wag31$_{\Delta N}$ was defective in membrane binding (Fig. 4c,d) and showed a severe tethering capacity defect (Fig. 4f,g). Surprisingly, Wag31$_{K20A}$, which showed a marginal defect in membrane binding (Fig. 4c,d), showed a marked defect in tethering capacity (Fig. 4f,g). Together, these results suggest that membrane-binding and membrane-tethering are non-overlapping attributes in the structure of Wag31, and the N-terminal DivIVA-domain is indispensable for membrane-tethering activity.

Tethering is crucial for the survival of mycobacteria

Results presented in Fig.4c indicate that Wag31 binds strongly to CL-containing membranes. CL is a negatively charged inverted cone-shaped lipid that has been shown to be concentrated at the sites of negative curvature in the bacterial cell, i.e., pole and septum (52, 53). Wag31 has been demonstrated previously to localize at sites of negative curvature (28). Their similar localization pattern, coupled with Wag31's affinity to bind to CL, led us to examine the effect of this binding *in vivo*. We hypothesized that modulations in Wag31 levels might alter CL distribution in the cell.

To examine this, we utilized a fluorophore 10-N-Nonyl-acridine orange (NAO) that is widely used to examine the localization of CL in prokaryotes and in the mitochondria of eukaryotes (52, 54, 55). NAO binds to anionic phospholipids with an affinity constant of 7×10^4 M^{-1} and to CL with an affinity constant of 2×10^6 M^{-1} (56). We stained *Msm and* Wag31 depleted and overexpressing cells with NAO and imaged them using a fluorescence microscope. While the localization pattern of NAO was polar in

Msm+ATc and *Δwag31*-ATc, there was a redistribution of red fluorescence to the cytosol and along the
perimeter of the cell in *Δwag31*+ATc (Fig.5a-b). We quantified the distribution of fluorescence using a
line profile which yielded peaks of intensity at the poles and a basal level of fluorescence along the
lateral cell body for both *Msm*+ATc and *Δwag31*-ATc (Fig.5b). Wag31 overexpression also led to a
change in the distribution of NAO from the poles and septum to 'donut-like' fluorescent foci in the cytosol
(Fig.5c-d). As the affinity of NAO for CL is more than other anionic phospholipids and our result (shown
in Fig. 4b) indicate that Wag31 prefers to bind CL over PG (another negatively charged phospholipid),
the change in the distribution of NAO fluorescence implies that the loss or overexpression of Wag31
causes the mislocalization of CL in mycobacterium.

Subsequently, we investigated the impact of tethering-deficient Wag31 mutants on cell survival.
Wag31 has an N-terminal domain joined to the C-terminal via a linker region. The N-terminal domain
contains a positively charged patch in the intertwined region (the region between two helices in the N-
terminus) and membrane binding has been attributed to this region (50). This patch is lined with
conserved arginine and lysine residues, K15, K20, and R21 (50) (Fig. 5e). As our *in vitro* results suggest
the importance of K20 and the DivIVA-domain in tethering, we performed complementation assays with
the tethering-defective mutants to examine cell survival and morphology. *Δwag31* was complemented
with an integrative copy of either Wag31 or Wag31$_{K20A}$ or Wag31$_{ΔN.}$ *Δwag31*+ATc showed a two log-
fold difference in survival and a 40 % decrease in the percentage of rod-shaped cells compared with
Δwag31-ATc (Fig. 5f-g). Complementation with Wag31 and Wag31$_{K20A}$ rescued the loss of cell viability
and restored the rod shape of the cells (Fig. 5f-g), whereas Wag31$_{ΔN}$ behaved similar to *Δwag31*+ATc.
It neither restored the rod shape of the cells nor reversed the survival defect (Fig. 5f-g). These results
highlight the importance of the N-terminal DivIVA-domain for sustaining mycobacterial morphology and
survival.

Tethering is correlated to lipid levels and CL localization in mycobacteria
As described above, Wag31-mediated membrane tethering is an absolute for the survival of
mycobacteria. Perturbations in lipid homeostasis and mislocalization of CL due to altered Wag31 levels
hint at a correlation between its ability to function as a homotypic tether and maintain lipid composition
and localization. To derive the mechanism behind it, we performed fluorescence imaging experiments
with native and tethering defective mutants of Wag31. To investigate a role of tethering in lipid
homeostasis in mycobacteria, we stained wildtype and complementation strains with BODIPY and
quantified the fluorescence intensities. We observed comparable staining intensity in *Msm* +ATc and,
Δwag31 -ATc (Fig. 6a-b). *Δwag31*+ATc expectedly had the highest BODIPY staining intensity,
indicating higher than normal lipid levels (Fig. 6a-b). Complementation with Wag31 fully restored the
lipid levels to normal, whereas complementation with Wag31$_{K20A}$ couldn't restore the lipid levels
completely (Fig. 6a-b). Wag31$_{ΔN}$ exhibited higher lipid levels than *Δwag31*-ATc (Fig. 6a-b). These
results highlight that the maintenance of lipid homeostasis by Wag31 is a consequence of its tethering
activity.

Next, we examined the dependence of CL localisation on tethering by staining *Msm* +ATc,
Δwag31 -ATc*, Δwag31*+ATc*, Δwag31::wag31*+ATc*, Δwag31::wag31$_{K20A}$*+ATc*,* and *Δwag31:: wag31$_{ΔN}$*
+ATc with NAO. NAO staining showed a polar and septal (in dividing cells) localization pattern in the

case of *Msm* and *Δwag31*-ATc, whereas it stained the cytosol and perimeter of the Wag31 depleted cells (Fig. 6c-d). Complementation with Wag31 restored NAO localization completely, whereas Wag31$_{K20A}$ restored it partially, which could be attributed to its limited tethering activity (Fig. 6c-d). Importantly, we observed a mis-localized pattern of NAO with Wag31$_{\Delta N}$ similar to Wag31 depletion (Fig. 6c-d). These results inform of the dependence of CL localization on membrane tethering activity of Wag31. Taken together, the data suggests the importance of the N-terminal region of Wag31 in regulating lipid homeostasis and localization.

The N and C-terminal domains of Wag31 have distinct functions.

Wag31 is a 272 amino-acids long protein that contains an N- and a C-terminal domain that are joined by a linker region (Fig. 7a). In this report, we established a role for the N-terminal region of Wag31 that comprises DivIVA-domain in membrane-tethering (Fig. 4). We also identified the protein-interaction network of Wag31 with the help of MS/MS analysis (Fig. 3 & Fig. S4). To investigate which domain of Wag31 promotes its interactions with other proteins, we generated *Msm* strains expressing either N-terminal (Wag31$_{\Delta N}$) or C-terminal truncated (Wag31$_{\Delta C}$) Wag31 (Fig. 7a) and performed a series of His pull-down experiments with both the mutants. To investigate the protein-binding ability of the domain truncation mutants, we tested them against novel interactors of Wag31 viz. MurG, SepIVA, and Msm2092 from our interactome database (Fig.3a) and AccA3, a known interactor of Wag31. MmpS5 served as a non-interactor negative control in the experiment.

We performed His pull-down assays utilizing *E. coli* lysates expressing either Wag31 or Wag31$_{\Delta C}$ or Wag31$_{\Delta N}$ (Fig. 7b, e). These lysates were mixed and incubated with *Msm* lysates expressing 3x-FLAG tagged versions of the above-mentioned proteins (Fig.7c, f) in separate reactions and subsequently pulled-down with Cobalt beads. The western blot analysis revealed that all the tested interactors bound to Wag31 (Fig. 7d, g). Expectedly, MmpS5, being a non-interactor of Wag31 did not bind to it (Fig. 7d). As shown in Fig. 7d, none of the interactors bound to Wag31$_{\Delta C}$, indicating that the presence of N-terminal DivIVA-domain is not sufficient to enable Wag31 to interact with other proteins. Interestingly, interactions with all the proteins were retained in the case of Wag31$_{\Delta N}$ (Fig. 7g), thus suggesting that Wag31 interacts with other proteins via its C-terminal. Taken together, the results suggest that the N-terminal region of Wag31 is involved in membrane-tethering through interactions with CL, and the C-terminal region is involved in modulating interactions with other membrane-associated proteins. Thus N-terminal and C-terminal of Wag31 have distinct molecular functions that together are crucial for maintaining cell shape, lipid homeostasis and survival of mycobacteria.

Discussion.

Wag31 modulates mycobacterial cell division by acting as an adaptor protein that recruits the elongation machinery to the old pole for unipolar growth of cells (10, 15, 28). In the absence of Wag31, rod-shaped cells advance into a spheroplast-like stage, which is thought to arise from dysregulated PG metabolism (10, 11, 28). Recently, Wag31 has been implicated in maintaining the lipid-rich IMD compartment (37). But the molecular mechanisms by which Wag31 manages these functions has remained unclear. In this study, we discover an additional function of Wag31 as a membrane tether,

which likely reconciles many of the previously ascribed functions of this protein into a unified model. To
gain insights into defects associated with a change in cell shape, we imaged Wag31-deficient cells and
found a dramatic accumulation of lipids in the form of ILIs. Interestingly, both *Msm* and *ΔWag31*-ATc
showed the presence of ILIs, but in much less abundance, indicating that ILIs are a normal occurrence
that may function to regulate cellular functions (Fig. 1g-h).

The loss of Wag31 leads to the accumulation of neutral lipids while its overexpression increase
the levels of alpha-mycolic acids (α-MA), implying that the maintenance of native Wag31 levels is critical
for the proper balance of lipid composition (Fig. 2f-h). The lipid classes that get impacted by the
depletion of Wag31 vs overexpression are different. Wag31 is an adaptor protein that interacts with
proteins of the ACCase complex (28, 30) that synthesize fatty acid precursors and regulate their activity
(34). The varied response on lipid homeostasis could be attributed to a change in the stoichiometry of
these interactions of Wag31. While Wag31 depletion would prevent such interactions from occurring
and might affect lipid synthesis that directly depends on Wag31-protein partner interactions, its
overexpression would lead to promiscuous interactions and a change in the stoichiometry of native
interactions that would ultimately modulate lipid synthesis pathways.

A protein's interactome is akin to its functional fingerprint. Here, we identify 116 novel
interacting partners of Wag31 through mass spectrometry and validate three of these interactions,
namely with MurG, SepIVA and Msm2092, through pull-down studies (Fig. 3 & 5i). Strikingly, the bulk
of Wag31 interactors are membrane proteins, which might be attributed to the formation of Wag31
oligomers on the membrane. Our results identify MurG, SepIVA, which regulates MurG (47), the D-
aminopeptidase Msm2092 and AmiB, both of which are PG remodellers, as Wag31 interactors.
Together, these results contribute towards our understanding of how Wag31 affects PG synthesis (Fig
3a,d). AccA3 is an interactor of Wag31 (28, 34) and catalyses the first committed step in the synthesis
of long-chain fatty acids by forming malonyl-CoA. In addition, Long-chain specific acyl dehydrogenase
(LCAD), which we report here to be another interactor of Wag31, catalyses the first step in the β-
oxidation of fatty acids, which might explain the accumulation of long chain PEs and long-chain fatty
acids.

As mentioned above, the overexpression of Wag31 leads to an upregulation in α-MA levels
(Fig. 2h). The localization of the α-MA transporter MmpL3 depends on polar PG levels (57, 58). MurG
is an essential enzyme for PG synthesis (59) and we report here that it strongly interacts with Wag31
(Fig. 3a,d). It is tempting to hypothesize that homeostatic control on lipid composition arises from the
ability of Wag31 to modulate the activity of enzymes involved in lipid metabolism and PG synthesis.
That both depletion and overexpression of Wag31 affects lipid composition is likely also a manifestation
of its ability to function as a homotypic tether, where the tethering ability is inherently sensitive to the
tether concentration.

Our results are the first to establish a membrane tethering function of Wag31. DivIVA proteins
possess an N-terminal membrane-binding domain and a C-terminal coiled-coil domain (50, 60). Our
data show that disruption of the N-terminal domain (Wag31$_{\Delta N}$) markedly reduces membrane binding
and completely abolishes membrane tethering (Fig. 4f, g). The N-terminal domain binds CL while the
C-terminal coiled-coil domain facilitates multimerization, which together could explain the molecular

basis for Wag31 to function in homotypic tethering of CL-containing membranes. Results from testing
mutants defective in membrane tethering indicate that this function is important for the proper
localisation of CL and eventually the survival of mycobacteria (Fig. 5f & 6c-d). Thus, alterations in the
native levels of Wag31 results in a displacement of CL from the poles (Fig. 5a-d). However, given
NAO's ability to bind to anionic phospholipids (albeit with lower affinity), a smaller degree of
NAO redistribution intensity might be coming indirectly from other anionic phospholipids displaced
from the membrane due to the loss of membrane integrity or cell shape changes caused by Wag31
depletion.

While, complementation with Wag31 reverses the phenotypic changes observed upon Wag31
depletion, complementation with the membrane tethering-defective mutant (Wag31$_{\Delta N}$) fails to do so
(Fig. 5f-g & 6a-d). These results not only highlight the dependence of CL localization on membrane-
tethering but also underscores the importance of tethering for maintaining cellular morphology,
physiology and survival. CL micro-domains at the poles (52, 53) have been shown to be an important
hub for protein localisation and activity. Several studies have implicated CL in modulating the activities
of membrane-resident proteins involved in electron-transport, protein translocation, bacterial two-
component system etc. (61-64). Our work delineates the roles of the N- and C-terminal domains of
Wag31. The N-terminal DivIVA-domain facilitates lipid binding and membrane tethering, and the C-
terminal coiled-coil domain engages in protein-protein interactions and facilitates the multimerization of
Wag31. This is evident from findings that deletion of the N-terminal DivIVA-domain affects lipid binding
and tethering but not its ability to bind interacting partners (Fig. 4c-d, f-g & 7c). Based on our results,
we propose a model in which the N-terminal of Wag31 binds and tethers CL ensuring its proper
localization while the C-terminal is free to interact and recruit proteins to the membrane to facilitate cell
growth and division (Fig. 8). We reason that the loss of cell-shape and viability in the absence of
tethering occurs due to mis-localization of CL which in turn has a negative impact on membrane-
localized processes at a global scale.

393
394

Materials and Methods

All the illustrations were made using BioRender. List of constructs and strains used in the study are

given in Table S1. Mass spectrometry samples were processed and performed by Valerian Chem.

Bacterial strains and culturing

Middlebrook 7H9 medium (BD Biosciences) or 7H11 agar (BD Biosciences) was used to cultivate

mycobacterial strains. 7H9 medium was supplemented with 10% ADC (albumin, dextrose, NaCl, and

catalase) and 0.2% glycerol (Sigma) and, 0.05% Tween-80 (Sigma) or 7H11 agar (BD Biosciences)

with 10% OADC (oleic acid added to ADC) and 0.2% glycerol were added to the media. Antibiotic

concentrations used: Hygromycin (Hyg) was used at 100 µg/mL, Kanamycin (Kan) at 25 µg/mL, and

Apramycin (Apra) at 30 µg/mL for *Msm*. *E. coli* was cultured in Miller Luria Bertani Broth (Himedia) or

Miller Luria Bertani Agar. Antibiotic concentrations used: Hyg was used at 150 µg/mL, Kan at 50 µg/mL,

Apra at 30 µg/mL, Ampicillin (Amp) at 100 µg/mL for *E. coli*.

Generation of *wag31* conditional mutant and *in vitro* Growth kinetics

Recombineering method, as described in (65), was used to generate a Wag31 conditional gene

replacement mutant. First, wag31 was cloned under a tet regulatable promoter in pST-KiT$_{off}$ (kan res;

(66)), which turns off transcription upon the addition of anhydrotetracycline. pST-wag31 construct thus

generated was electroporated into a recombineering proficient strain, Msm::pJV53. An allelic exchange

substrate was generated by cloning 675 bp from upstream (Left hand sequence, LHS) and 700 bp from

downstream (Right hand sequence, RHS) of *wag31* (including ~150 bp at both ends) with *hyg*[res]

cassette and *oriE+cosλ* and digested with SnaBI to release the donor LHS-hyg[res]-RHS. The linear donor

was electroporated into Msm::wag31 expressing gp60 and gp61 recombinases. The colonies were

selected on Hyg and ten colonies were screened with multiple primer pairs to confirm recombination at

the native locus. Confirmed recombinants were passaged multiple times to cure them of pJV53 to

generate conditional mutant Δwag31.To perform *in vitro* growth kinetics, secondary cultures of *Msm*

and *Δwag31* were seeded at a density of A_{600}~ 0.05 and either left untreated or treated with 100 ng/mL

ATc for 12 h. At 0 and 12 h, appropriate serial dilutions were plated on 7H11 agar plate and incubated

at 37°C for 2-3 days, and CFUs were enumerated. CFUs were plotted using GraphPad Prism 9 and

analysed by performing Two-way ANOVA (0.0021 (**), 0.0002 (***)).

Western blot analysis

Msm, Msm::wag31 and *Δwag31* transformants grown in the presence or absence of IVN/ ATc for 12 h

and harvested by centrifugation at 4000 rpm for 10 min at 4ºC. The pellets were resuspended in 1:3

(pellet weight: buffer volume) Phosphate Buffer Saline containing 5% glycerol and 1 mM protease

inhibitor- PMSF. The resuspension was transferred into pre-equilibrated bead-beating tubes filled to

one-fourth with 0.1mm Zirconia beads. Pre-equilibration was performed by washing the beads with

autoclaved MilliQ followed by lysis buffer. The tubes were placed in a bead-beater and subjected to 6-

8 cycles of bead-beating -1 min on and 2 min rest on ice. To obtain WCL, the tubes were centrifuged

at 13000 rpm for 10 min at 4°C, and the supernatant was collected in a microcentrifuge tube. The

concentration of protein in each sample was estimated by Pierce BCA protein Assay kit (Thermo Fischer

Scientific), and 40 µg WCLs were resolved on 10-12% SDS-PAGE gels at constant current and

transferred onto Nitrocellulose (NC) membrane. This was followed by blocking the membrane with 5%

Bovine Serum Albumin (BSA) for an hour at RT and subsequently incubating it with the desired primary

antibody for 2 h at RT or overnight at 4°C. The primary antibodies used were either anti-RpoA (raised

in rabbit), anti-Wag31 (raised in rabbit), anti-FLAG (raised in mice, Sigma), anti-His (raised in rabbit,

CST). The membrane was washed thrice for 10 min each with PBST (PBS + 0.5% Tween-20) to remove

any non-specific binding. Next, the membrane was probed with appropriate secondary antibody (anti-

rabbit, Invitrogen or anti-mice, Invitrogen) for an hour at RT, washed thrice for 10 min each with PBST,

developed using ECL reagent (Millipore) and visualized on BioRad ChemiDoc Touch Gel imaging

system.

Sample preparation for SEM and TEM

Msm and *Δwag31* were set up at A_{600}~0.05 and grown overnight in either the presence of

absence of 100 ng/ml ATc. Post 12 h, the cells were harvested by centrifugation at 4000 rpm for 10 min

at RT. Cells equivalent to A_{600} ~2 were taken further for sample preparation and washed thrice with 5

mL sterilized PBS. The cells were fixed with 5 mL of fixative (4% Paraformaldehyde (Sigma), 2.5%

Glutaraldehyde (EMS), 0.1M Na-Cacodylate) for 4 h at RT protected from light. Cells were tap-mixed

every 30 min to ensure that they remained in suspension. This was followed by washing the cells twice

for 10 min with 0.1 M Na-Cacodylate buffer. Subsequently, cells were stained with 3 mL of 1% Osmium

Tetraoxide (Os_4O_2) protected from light for a maximum of 1.5 h with tap-mixing every 30 min. Os_4O_2

was removed from the samples by centrifugation at 4000 rpm at 10 min at RT and discarded as per

safety guidelines. The cells were then subjected to serial dehydration at RT with increasing

concentrations of ethanol from 25% to 100%. Briefly, the cells were incubated for 5 min with 25%

ethanol, harvested, and incubated for 7 min with 50% ethanol. This was followed by serially incubating

the cells for 15, 20, and 30 min for 75%, 95%, and 100% (thrice) ethanol, respectively. Cells were then

taken forward for critical-point drying with the help of 100 μl Hexamethyldisilazine (HMDS). Meanwhile,

stubs were prepared by placing carbon tape and coverslip on them. 10 μl of the dried sample was

spotted on the coverslip and evenly spread. Next, the stub was coated with 0.8 mm Gold and stored in

a desiccator or imaged at 20,000X using FEI Nova NanoSEM 450. For TEM, cells were grown and

harvested as described above for SEM. For TEM sample preparation, 50 OD cells were taken, fixed,

stained, and gradually dehydrated with ethanol, as explained above. The cells were then infiltrated with

Epon 812 resin. This was followed by cutting the resin into several 62 nm thick ultra-thin sections using

an ultramicrotome (Leica EM UC7). The sections were then stained with Uranyl acetate and lead citrate

and examined on a 120 kV transmission electron microscope (L120C, Talos, Thermo Scientific) at

17,500 magnification.

BODIPY labelling, imaging and analysis

1 mL cells of *Msm* and *Δwag31* either untreated or treated with 100 ng/mL ATc for 12 h were stained

with 20 μg/mL BODIPY (Invitrogen). For examining lipids in Wag31 overexpression strain, *Msm* and

Msm::wag31 were treated with 5 μM IVN for 12 h and 1 mL cells of both strains were stained with 2.5

μg BODIPY. Staining was performed for 30 min protected from light in an incubator shaker followed by

washing with sterilized Phosphate buffer saline (PBS). For Msm and Δwag31, ~10 μl of cells were spotted

on a 1% agar pad (prepared in 7H9) and visualized using a 100X/1.4 NA oil immersion objective on a

Zeiss, LSM880 at 488 nm/510 nm (Excitation: Emission). Background correction in the collected images

followed by Corrected Total Cell Fluorescence analysis was performed using Fiji (2) and plotted using Graph Pad Prism 9. For examining lipids in BODIPY-stained Msm and Msm::wag31, 2 μl of cells were spotted on a 1% agar pad (prepared in 7H9) and visualized using a 100X/1.32 NA oil immersion objective on a Leica THUNDER Imaging systems at 475nm:519nm (Excitation: Emission). About 20 Z-stack images, each spaced 0.1 μM apart, were acquired using a scientific CMOS K8 camera (Leica microsystems). Background correction and Thunder image processing was performed on the images using the Las-X software module (Leica microsystems). Processed images were then analysed for CTCF using Fiji (2) and plotted using Graph Pad prism 9.

Sample preparation for Lipidomics and analysis

Secondary cultures of *Msm* and *Δwag31* were seeded at a density of A600~ 0.05 and either left untreated or treated with 100 ng/mL ATc for 12 h until A600~ 0.8-1.0. Post 12 h incubation with ATc, Cells corresponding to A_{600}~20 were harvested and processed for lipid extraction. Bacterial pellets were resuspended in 1 ml sterilized PBS and transferred to a 3 ml glass vial. 2:1 Chloroform::Methanol mixture was added to it and vortexed vigorously for 2 min. Chloroform contained 1 nm 15:0 FFA to be used as the internal standard for normalization. The samples were centrifuged at 1500 rpm for 5 min at RT to separate the organic layer. The latter was transferred to a new glass vial, and 2 ml Chloroform (containing 2% Formic acid) was added to the old vial to extract phospholipids from the sample. The samples were centrifuged at 1500 rpm for 5 min at RT to extract the phospholipid-containing organic layer. Both organic layers were pooled and dried under a stream of Nitrogen gas at RT. The dried lipid extracts were re-solubilized in 200 μL of 2:1 CHCl3: CH$_3$OH, and 10 μL was injected into an Agilent 6545 QTOF (quadrupole-time-of-flight) instrument for semi-quantitative analysis using high-resolution auto MS/MS methods and chromatography techniques. A Gemini 5U C-18 column (Phenomenex) coupled with a Gemini guard column (Phenomenex, 4x3 mm, Phenomenex security cartridge) was used for LC separation. The solvents used for negative ion mode were buffer A: 95:5 H2O:CH$_3$OH + 0.1% ammonium hydroxide and buffer B: 60:35:5 Isopropanol: CH$_3$OH:H2O + 0.1% ammonium hydroxide. The 0.1% ammonium hydroxide in each buffer was replaced by 0.1% Formic acid + 10 mM Ammonium Formate for positive ion mode runs. All LC/MS runs were for 60 min, starting with 0.3 mL/min 100% buffer A for 5 minutes, 0.5 mL/min linear gradient to 100% buffer B over 40 minutes, 0.5 mL/min 100% buffer B for 10 minutes, and equilibration with 0.5 mL/min 100% buffer A for 5 minutes. The following settings were used for the ESI-MS analysis: drying gas and sheath gas temperature: 320°C, drying gas and sheath gas flow rate: 10 L/min, fragment or voltage: 150 V, capillary voltage: 4000 V, nebulizer (ion source gas) pressure: 45 psi and nozzle voltage: 1000 V. For analysis, a lipid library was employed in the form of a Personal Compound Database Library (PCDL), and the peaks were validated based on relative retention times and fragments obtained. This library was selectively curated from the MycoMass database and the LIPID MAPS Structure Database (LMSD). All robustly detected lipid species were quantified by normalizing areas under the curve (AUC) to the AUC of the relevant internal standard added and by normalizing to the total cell number. Log2 fold changes were then plotted for each lipid compared to the Msm-ATc samples.

Immunoprecipitation (IP) and MS/MS analysis

250 mL secondary cultures of *Msm*::*gfp* and *Δwag31* were inoculated at an A600~ 0.05 and

incubated in a shaker incubator until A600~ 0.8-1.0. Cells were collected and lysed using bead-beating to make Whole cell lysate (WCL) and passed through a 0.2 µM filter. 3 mg lysate from each group was immunoprecipitated overnight with anti-FLAG® M2 magnetic beads (Millipore), washed thrice, and heated in 4X Laemmli Buffer. One-fifth of the IP supernatant was resolved on a 10% gel, transferred onto a Nitrocellulose membrane, and probed with anti-FLAG Antibody (Sigma). Lysates from confirmed IPs (biological quadruplets) were taken further for mass spectrometry sample preparation and analysis. Sample preparation and, mass spectrometry analysis were outsourced to VProteomics, New Delhi. Briefly, the IP samples were reduced with 5 mM TCEP, followed by alkylation with 50 mM iodoacetamide. The samples were digested with 1µg Trypsin for 16 h at 37 °C. Digests were cleaned using a C18 silica cartridge and resuspended in a buffer containing 2% acetonitrile and 0.1% formic acid. 1 µg of peptide was used for analysis in an Easy-nlc-1000 system coupled to an Orbitrap Exploris mass spectrometer. The gradients were run for 110 min. MS1 spectra were acquired in the Orbitrap (Max IT = 60 ms, AGQ target = 300%; RF Lens = 70%; R=60K, mass range = 375−1500; Profile data). Dynamic exclusion was employed for 30 seconds, excluding all charge states for a given precursor. MS2 spectra were collected for the top 20 peptides. MS2 (Max IT=60ms, R= 15K, AGC target 100%). Proteome Discoverer (v2.5) was used to analyse the RAW data against the Uniprot Msm database. The precursor and fragment mass limitations for the dual Sequest and Amanda searches were established at 10 ppm and 0.02 Da, respectively. The false discovery rate for proteins and peptide spectrum matches was adjusted to 0.01 FDR. For identification of interactors unique to Wag31, we removed all the proteins obtained in the three replicates of *Msm::gfp* from the *Δwag31* datasets. In the corrected datasets, proteins common to all the three replicates with minimum unique peptides #2 were considered interactors. To identify top hits, PSM cut-off was set to 18 and unique peptides to 5.

Protein expression and purification

BL21(DE3) cells transformed with pET-wag31, or pET-wag31K20A or pET-wag31$_{\Delta N}$ were cultured at 37ºC until the A600 ~0.6. 0.1 mM IPTG was added to the culture to induce protein expression, followed by incubating the culture at 18ºC for 12 h. Cells were pelleted and stored at -40 ºC. The bacterial pellet was resuspended in a lysis buffer (20 mM HEPES pH 7.4, 150 mM NaCl, and 20 mM imidazole with 1 mM phenylmethylsulphonyl fluoride (PMSF)) and lysed by sonication in an ice-water bath. Lysate was spun at 30,000 x g for 20 min, and the supernatant obtained was incubated with TALON® Metal Affinity Resin (Takara Bio) 4 ºC for 30 min. The resin was extensively washed with lysis buffer, and the bound protein was eluted with 20 mM HEPES pH 7.4, 150 mM NaCl, and 250 mM imidazole. The elution was dialyzed against 20 mM HEPES pH 7.4 150 mM NaCl and stored at 4 ºC for the duration of the experiments. Aggregates were removed by centrifugation at 100,000 g. Concentration of the proteins was quantified by measuring A280 using the molar extinction coefficient predicted by the Expasy ProtParam tool.

Liposome preparation

1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE) and1′,3′-bis[1,2-dioleoyl-sn-glycero-3-phospho]-glycerol (sodium salt), CL were obtained from Avanti Polar Lipids. 1,1'-dilinoleyl-3,3,3',3'-tetramethylindocarbocyanine, 4-chlorobenzenesulfonate (FAST DiI™), and 3,3'-dilinoleyloxacarbocyanine perchlorate (FAST DiO™)

were obtained from Invitrogen. The UV-activable, diazirine-containing reactive fluorescent lipid probe BODIPY-diazirine phosphatidylethanolamine (BDPE) was prepared as described earlier (1, 2). For liposome preparation, lipids were aliquoted from chloroform stocks at desired ratios in a clean glass tube and dried under a high vacuum for an hour to a thin film. The dried lipids were used at a final concentration of 1 mM (deionised water was added to them). After hydrating at 50 °C for 30 minutes, lipids were vigorously vortexed and then extruded through polycarbonate filters with pore sizes of 100 nm (Whatman). For PLiMAP assays, liposomes were CL, DOPG, or DOPE with 69 mol% DOPC and 1 mol% BDPE. Liposomes used in tethering assays were made similarly with 30 mol% CL or DOPE and 1 mol% of the fluorescent lipid DiI or DiO) in the DOPC background.

Proximity-Based Labelling of Membrane-Associated Proteins (PLiMAP)

For carrying out the PLiMAP assay, Wag31 and its mutants were purified and desired liposomes were prepared as described in supplementary. PLiMAP was carried out and analysed as described (67). Briefly, 2 µM Wag31 or its mutants were incubated with liposomes containing 30 mol% CL, DOPG, or DOPE (200 µM total lipid) in a final volume of 30 µL. The reaction was incubated for 30 min in the dark at room temperature followed by exposure to a 1 min long pulse of 365 nm UV light (UVP crosslinker CL-1000L) at an intensity of 200mJ cm^{-2}. The reaction was mixed with Laemmli sample buffer, boiled, and equal volumes for all reactions were loaded onto a 10% polyacrylamide gel and resolved using SDS-PAGE. iBright1500 (Thermo Fischer Scientific) or Amersham Typhoon (GE) were used for imaging gels. Gels were first imaged for BODIPY fluorescence and later fixed and stained with Coomassie Brilliant Blue.

Tethering assays, fluorescence imaging, and image analysis

1 µM of Wag31 and its mutants were mixed with 10 µM of 100 nm extruded liposomes in Eppendorf. The mixture was then transferred to a LabTek chamber and imaged using 100x 1.4 NA oil-immersion objective on an Olympus IX73 inverted microscope connected to an LED light source (CoolLED) and an Evolve 512 EMCCD camera (Photometrics). Image acquisition was controlled by µManager, and images were analysed using Fiji. For single liposome intensity analysis, all images from the same condition from two independent experiments were combined into one montage. The montage was duplicated and thresholded using Fiji's Otsu algorithm to generate a binary mask. The mask was overlaid onto the original montage to obtain the maximum intensity of liposomes.

NAO labelling, imaging and analysis

Msm, *Msm::wag31* were treated with 5 µM IVN for 12 h and *Msm* and *Δwag31* were either left untreated or treated with 100 ng/mL ATc for 12 h. For complementation experiment, 1 mL of each strain was stained with 2 µM NAO for 1 h protected from light in an incubator shaker and washed thrice with sterilised 1X PBS. 2 µl of cells were spotted on a 1% agar pad (prepared in 7H9) and visualized using a 100X/1.32 NA oil immersion objective on a Leica THUNDER Imaging systems at 475nm:535nm (Excitation: Emission). About 20 Z-stack images, each spaced 0.1 µM apart, were acquired using a scientific CMOS K8 camera (Leica microsystems). For CL specific labelling, collection was also performed at 642nm.

Background correction and Thunder image processing was performed on the images using the Las-X software module (Leica microsystems). Processed images were then analysed using Fiji (68)and

plotted using Graph Pad prism 9. Briefly, N=100 cells across two independent experiments were examined for their profile of CL distribution by normalizing the cell length and assigning pole with brighter NAO fluorescence intensity as 0 and the other pole as 100. The fluorescence intensity was normalised to the highest fluorescent value for each cell. Corresponding fluorescence intensity value for all cells for every point between 0 to 100 were calculated from the LOOKUP function of excel. The values representing CL fluorescence across 100 cells were averaged out and plotted against normalised cell length in GraphPad prism 9. 0th order smoothening with 4 neighbours on each size was performed on the curves obtained.

Complementation experiment

Secondary cultures of *Msm, Δwag31, Δwag31::wag31, Δwag31::wag31$_{K20A}$,* and *Δwag31::wag31$_{ΔN}$* were set up at A$_{600}$ ~0.05. For each strain, two types of cultures were set up in triplicates- untreated and treated with 100ng/mL ATc and grown for 12 h. At 0 and 12 h, appropriate serial dilutions were plated on 7H11 agar plates and incubated at 37$^{°}$C for 2-3 days. CFUs were enumerated and plotted using GraphPad Prism 9. Statistical analysis was performed using two-way RM ANOVA followed by *Tukey's* multiple comparison test, α=0.01, GP: 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), <0.0001 (****). For BODIPY- and NAO- labelling, the strains were grown as described above. At 12 h, 1 mL cultures of *Msm*+ATc, *Δwag31, Δwag31* +ATc, *Δwag31::wag31*+ATc*, Δwag31::wag31$_{K20A}$* +ATc*,* and *Δwag31::wag31$_{ΔN}$* +ATc were taken and labelled with either BODIPY or NAO. The labelling, imaging and analysis for both fluorophores was performed as described in the respective sections above.

His pulldown assay.

BL21 (DE3) cells transformed with no plasmid, or pET-wag31 or pET-wag31$_{ΔN}$ or pET-wag31$_{ΔC}$ were cultured and cell lysates were prepared as described for PLiMAP and tethering assays. Wag31 or Wag31$_{ΔN}$ was assessed by resolving the samples on 10% SDS-PAGE followed by Coomassie staining whereas Wag31$_{ΔC}$ was resolved on Tris-Tricine PAGE as described elsewhere (69). The interacting partners, namely *murG, sepIVA, msm2092* and *accA3* and the non-interactor *mmpS5* were amplified from Msm genomic DNA using gene specific primers harbouring NdeI-HindIII sites and amplicons were digested and cloned into the corresponding sites in pNit-3X-FLAG shuttle vector. The constructs thus generated were electroporated in *Msm* to generate *Msm::murG, Msm::SepIVA, Msm::msm2092, Msm::accA3* and *Msm::mmpS5.* The cultures grown till A600 ~0.8 were used for fresh inoculation at A$_{600}$ of 0.05 in 7H9 media containing 0.5 μM IVN. Cultures were grown for 12 h and lysates were prepared by bead-beating and 40 μg were resolved on 10% gels and probed with α-FLAG antibody.

500 μg of *E. coli* lysates expressing Wag31 or mutant were mixed with 300 μg *Msm* lysates expressing Wag31 interactors in a total reaction volume of 700 μL and the samples were twirled at 4^{o}C for 2 h. 50 μL cobalt beads/ sample (GoldBio) were washed twice with lysis buffer and incubated with 5% BSA solution for 2 h at 4^{o}C. Subsequently, beads were washed with lysis buffer and incubated with samples for 2 h (twirled at 4^{o}C). The pull downs were washed thrice in lysis buffer containing 1%

Triton X-100 and finally resuspended in 2X-SDS sample buffer. Pulldown samples were resolved on

10-12% SDS-PAGE and probed with α-His and α-FLAG antibodies.

Figure Legends:

Fig.1| Loss of Wag31 leads to the formation of Intracellular Lipid Inclusions (ILIs).

(a.) Illustration depicting the known role of Wag31 in directing polar elongation and maintaining rod-

shaped morphology of cells. **(b.)** Whole Cell Lysates **(**WCL) were prepared from *Msm* and *Δwag31*

(Table S1) either untreated or treated with 100 ng/mL ATc for 12 h, and 40 µg lysate was resolved on

a 12% gel, transferred on Nitrocellulose membrane, and probed with either α-Wag31 or α-rpoA antibody

(loading control), respectively. **(c.)** Cultures showing growth of *Msm* and *Δwag31* either untreated or

treated with 100 ng/mL ATc for 12 h. **(d.)** *Msm+*ATc, *Δwag31*-ATc, and *Δwag31+*ATc were tracked for

bacillary survival at 0 and 12 h by enumerating CFUs. Appropriate serial dilutions were plated on 7H11

plates (without antibiotic or ATc), and bar graphs demonstrating bacillary survival (CFU $\log_{10}$/mL ±

standard deviations (SD, represented by error bars) were plotted at indicated time points. Statistical

significance was calculated using Two-way ANOVA (0.0021 (**), 0.0002 (***). The experiment was

performed twice independently, each performed in triplicates. **(e.)** SEM micrographs showing *Msm*,

Δwag31-ATc, and *Δwag31+*ATc at 12 h post-treatment. **(f.)** Cells depicting rod, bulged or round

morphology. The yellow arrowhead indicates bulged pole of the cell. **(g.)** 200 cells from each group

were checked for either rod, bulged or round morphology, and the percentage of cells with these

morphologies in all the three strains was plotted in a bar graph. **(h.)** TEM micrographs showing *Msm*,

Δwag31-ATc, and *Δwag31+*ATc at 12 h post-treatment. **(i.)** 250 cells, each from *Msm*, *Δwag31*-ATc

and 185 cells from *Δwag31+*ATc, were divided into three classes (0-2, 2-5, and >5 ILIs) based on the

number of ILIs and the percentage of each class in every strain was plotted in a bar graph.

Fig.2| Wag31 levels perturb lipid homeostasis

Representative micrographs showing BODIPY staining in **(a.)** *Msm+*ATc, *Δwag31*-ATc and,

*Δwag31+*ATc samples imaged 12 h post-ATc treatment **(b.)** *Msm+* IVN, *Msm*::*wag31* + IVN samples

imaged 12 h post 5 µM IVN treatment. **(c.)** Corrected total cell fluorescence was calculated in individual

bacterial cells from *Msm+*ATc, *Δwag31*-ATc, and *Δwag31+*ATc strains (N=48) and plotted in a graph.

Horizontal bar represents median CTCF. (Statistical analysis was performed using one-way ANOVA

followed by Brown-Forsythe test (Tukey's multiple comparison test; 0.0021 (**), <0.0001 (****)). **(d.)**

Corrected total cell fluorescence was calculated for 42 individual cells from *Msm+*IVN, and

Msm::*wag31+*IVN strains and plotted in a graph (Statistical analysis was performed using Welch's t-

test 0.0021 (**), **e.)** Schematic depicting the workflow of Lipidomics (LC-MS). **(f-g.)** Heatmap depicting

$\log_2$ fold change (FC) in indicated lipid classes of *Msm* +ATc, *Δwag31*-ATc, and *Δwag31+ATc* as

compared to *Msm*-ATc. The experiment was performed with six biological replicates, and the

abundances of identified lipid classes were normalized to *Msm*-ATc to yield relative abundances of

lipids in other strains. Blue and red colours indicate upregulated and downregulated lipid classes; color

intensity variation changes with $\log_2$ FC. **(h.)** Bar graph (Mean ± SEM) depicting the abundance of α-

MA in *Msm* and *Msm::wag31* treated with 5 µM IVN for 12 h. Values on the Y-axis represent area under

the curve (AUC) normalised to an internal standard (FFA 15:0). X-axis represent various species (chain

length variations) of α-MA. (Statistical analysis was performed using two-way ANOVA followed by a

Tukey's multiple comparison's test; <0.0001 (****)).

Fig.3| Molecular interactions of Wag31 with membrane-associated proteins govern cellular homeostasis.

(a.) Illustration showing the distribution of the top 20 interactors of Wag31 identified from MS/MS analysis from three independent immunoprecipitations. Interactors were classified on the basis of their functional category based on data available in Mycobrowser (70). MSMEG numbers or the identity of interactors are marked in smaller circles. **(b.)** Coomassie Brilliant blue (CBB) staining of plain *E.coli* lysate and *E.coli* lysate expressing His-Wag31. *E. coli* lysate prepared from BL21-DE3 strain was used as the control. **(c.)** Representative western blots demonstrating expression of 3XFLAG tagged-MurG, -SepIVA and -Msm2092. WCLs were prepared from 0.5 μM IVN induced *Msm::murG*, *Msm::sepIVA*, *Msm::msm2092* and 40 μg of each was resolved on a 10 % SDS-PAGE, transferred onto a nitrocellulose membrane and probed with α-FLAG and α-RpoA (loading control). **(d.)** Representative western blots showing interaction of MurG, SepIVA, and, Msm2092 with Wag31. *Msm* and *E. coli* lysates represented in (b & c) incubated together and pulled down with Cobalt beads were resolved on a 10-12 % SDS-PAGE, transferred onto nitrocellulose membrane and probed with α-His to detect the pulldown and α-FLAG to detect the interaction. The experiment was performed independently twice.

(e.) Graphic representation (adapted from (48)) shows the distribution of the top 20 interactors in different compartments of a mycobacterial cell: membrane, cytoplasm, or core (involved in DNA replication and recombination). Interactors from each compartment are marked inside the respective domain. **(f.)** Pie-chart shows the distribution of interactors belonging to the IMD, PM-CW, or membrane proteins whose specific localization is not determined. Percentage of proteins (top 20 hits) belonging to each membrane-compartment is mentioned.

Fig.4| Wag31 binds and tethers mycobacterial membrane.

(a.) Schematic showing the workflow of the PLiMAP assay. **(b.)** Coomassie Brilliant blue (CBB) staining of His-Wag31 and in-gel fluorescence (FI) from a representative PLiMAP experiment performed using 2 μM Wag31 and hundred-fold excess of 30 mol% of either 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE-), and1′,3′-bis[1,2-dioleoyl-sn-glycero-3-phospho]-glycerol (DOPG-), or CL-containing liposomes. **(c.)** CBB and FL gels from a representative PLiMAP experiment performed with 2 μM Wag31, Wag31$_{K20A,}$ or Wag31$_{\Delta N}$ either in the presence or absence of 30 mol% CL liposomes. Note the presence of SDS-resistant higher-order polymers in gel indicated by the red curly bracket. **(d.)** Quantitation of PLiMAP data from (c.) showing a difference in binding propensies of His-Wag31, His-Wag31$_{K20A}$ and His-Wag31$_{\Delta N}$. Data represent the mean ± SD of three independent experiments. Statistical analysis was performed using a 2-tailed unpaired student's t test in Graph Pad Prism 9. Wag31 vs Wag31$_{K20A}$ showed no significant difference (ns, p value = 0.5667) whereas Wag31 vs Wag31$_{\Delta N}$ (p value = 0.0081, **) and Wag31$_{K20A}$ vs Wag31$_{\Delta N}$ (p value = 0.0002, ***) showed significant difference in binding **(e.)** Representative micrographs from binding and tethering assay showing CL-Dil-liposomes (~100 nm) before (first panel) and after incubation (second panel) with 1 μM His-Wag31-GFP. Membrane and protein fluorescence are rendered in red and green, respectively. The experiment was performed independently twice. **(f.)** Representative micrographs from tethering experiments performed with His-Wag31, His-Wag31$_{K20A,}$ or His-Wag31$_{\Delta N}$ proteins. The left panels indicate PE

liposomes only (upper panel) and PE liposomes incubated with 1 μM Wag31 (lower panel), the middle

panels indicate CL liposomes (upper) or CL liposomes incubated with 1 μM Wag31 (lower panel), the

right panel indicates CL liposomes either incubated with His-Wag31$_{K20A}$ (upper) and His-Wag31$_{\Delta N}$

(lower). For visualization, PE- and CL-containing liposomes were doped with trace amounts of 3,3'-

dilinoleyloxacarbocyanine perchlorate (FAST DiO™) and 4-chlorobenzenesulfonate (FAST DiI™),

respectively. **(g.)** Data from particle analysis showing the maximum intensity of liposomes for all the

reactions from two independent experiments.

Fig.5| Tethering is crucial for the survival of mycobacteria

(a.) Top panel: Representative Fluorescent micrographs showing NAO intensity across *Msm*+ATc,

Δwag31-ATc and, *Δwag31*+ATc samples imaged at 12 h post-ATc treatment. Bright red foci of polar

CL are indicated with yellow arrowheads and septal CL foci with white arrowheads. Bottom panel:

shows respective phase images. **(b.)** Line plot depicting CL distribution along the length of the cells

across *Msm*+ATc, *Δwag31*-ATc and, *Δwag31*+ATc. N=100 cells across two independent biological

replicates were analysed using Fiji and their normalised fluorescence intensities were plotted against

cell length; fluorescence emitted from 0-20 % & 80-100 % of cell length was considered polar. **(c.)**

Representative fluorescent micrographs showing NAO staining in *Msm*+IVN, *Msm*::*wag31*+IVN

samples imaged at 12 h post 5 μM IVN treatment. Bright red foci of polar CL are indicated with yellow

arrowheads, and septal CL foci with white arrowheads. The panels below show corresponding phase

images. **(d.)** Line plot depicting CL distribution along the length of the cells in *Msm*+IVN,

Msm::*wag31*+IVN. N=100 cells across two independent biological replicates were analysed using Fiji.

The y-axis represents the normalized fluorescence intensities, and the x-axis represents cell length

expressed as a percentage of total cell length. Fluorescence peaks in 0-20 % & 80-100 % of cell length

were considered polar. **(e.)** Structure of Wag31 (MSMEG_4217) from AlphaFold protein structure

database (71). The structure represents the N-terminal, linker, and C-terminal domains in Wag31. The

N-terminal contains a short helix, followed by an intertwined loop harbouring positively charged amino

acid K20 (highlighted in red) and the DivIVA domain. The N-terminus is linked to the C-terminus via a

linker. **(f.)** *Msm*+ATc, *Δwag31*, *Δwag31::wag31*, *Δwag31::wag31$_{K20A}$*, *Δwag31::wag31$_{\Delta N}$* either

untreated or treated with ATc were scored for bacillary survival post 12 h ATc addition by enumerating

CFUs. Appropriate serial dilutions were plated on 7H11 plates (without antibiotic or ATc), and bar

graphs demonstrating bacillary survival (CFU log$_{10}$/mL ± standard deviations (SD, represented by error

bars) were plotted and statistical significance was calculated using two-way RM ANOVA followed by

Tukey's multiple comparison test, α=0.01, GP: 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), <0.0001

(****). The experiment was performed twice independently, each with triplicates. **(g.)** Strains described

in **(f.)** were scored for their morphology-rod, bulged or round at 12 h post ATc treatment. 308 cells of

each strain across two independent experiments were analysed.

Fig.6| Tethering is correlated with lipid levels and CL localization in mycobacteria

(a.) Top panel: Representative fluorescent micrographs showing BODIPY staining across *Δwag31-ATc*

and Msm, Δwag31, Δwag31::wag31, Δwag31::wag31$_{K20A}$, Δwag31::wag31$_{\Delta N}$ imaged at 12 h post ATc

addition. Bottom panel: corresponding phase images. **(b.)** Corrected total cell fluorescence (CTCF) of
BODIPY-stained cells was calculated from N=100 cells across two independent experiments and
analysed using Fiji and plotted in a graph. The horizontal bar represents the median CTCF. (Statistical
analysis was performed using one-way ANOVA followed by Brown-Forsythe test (Tukey's multiple
comparison test; 0.1234 (ns), 0.0332 (*), <0.0001 (****)). **(c.)** Fluorescent micrographs showing NAO
intensity of strains as mentioned above. Bottom panel: corresponding phase images. **(d.)** Line plot
analysis of strains described in **(c.)** depicting CL localisation. N=100 cells across two independent
experiments were analysed using Fiji and their normalised fluorescence intensities were plotted against
cell length.
766
Fig.7| The N and C-terminal domains of Wag31 have distinct functions.
a.) Illustration depicting the domain architecture of Wag31, Wag31$_{\Delta N}$ and Wag31$_{\Delta C}$. **(b.)** Coomassie
Brilliant blue (CBB) staining of plain *E. coli* lysate and *E. coli* lysate expressing His-Wag31$_{\Delta C}$. *E. coli* lysate
prepared from BL21-DE3 strain was used as the control. **(c.)** Representative western blots
demonstrating expression of 3XFLAG tagged-MmpS5, -MurG, -SepIVA, -Msm2092 and, -AccA3.
WCLs were prepared from 0.5 µM IVN induced *Msm::mmpS5*, *Msm::murG*, *Msm::sepIVA,*
Msm::msm2092, Msm::accA3 and 40 µg of each was resolved a 10 % SDS-PAGE, transferred onto a
nitrocellulose membrane and probed with α-FLAG and α-RpoA (loading control). **(d.)** Representative
western blots showing interaction of MurG, SepIVA, Msm2092 and, Acca3 with Wag31 and Wag31$_{\Delta c.}$
The non-interactor MmpS5 was used as a negative control. *E. coli* and *Msm* lysates represented in (b &
c) incubated together and pulled down with Cobalt beads were resolved on Tris Tricine gel as described
elsewhere (69), transferred onto nitrocellulose membrane and probed with α-His to detect the pulldown
and α-FLAG to detect the interaction. The experiment was performed independently twice. **(e.)**
Coomassie Brilliant blue (CBB) staining of *E. coli* lysates expressing His-Wag31$_{\Delta N}$. *E. coli* lysate prepared
from BL21-DE3 strain was used as the control. **(f.)** Representative western blots demonstrating
expression of 3XFLAG tagged-MurG, -SepIVA, -Msm2092 and, -AccA3. WCLs were prepared from
0.5 µM IVN induced *Msm::murG*, *Msm::sepIVA, Msm::msm2092, Msm::accA3* and 40 µg of each was
resolved a 10 % SDS-PAGE, transferred onto a nitrocellulose membrane and probed with α-FLAG and
α-RpoA (loading control). (g.) Representative western blots showing interaction of MurG, SepIVA,
Msm2092 and, Acca3 with Wag31 and Wag31$_{\Delta N.}$ *E. coli* and *Msm* lysates represented in (e & f) incubated
together and pulled down with Cobalt beads were resolved on 10% SDS-PAGE, transferred onto
nitrocellulose membrane and probed with α-His to detect the pulldown and α-FLAG to detect the
interaction. The experiment was performed independently twice.
790
Fig.8| Model for Wag31 mediated tethering of the membrane
a.) Model of Wag31 showing the N- and C-terminals. N-terminal houses the DivIVA-domain that is
involved in membrane-tethering and the C-terminal is the protein-binding domain that facilitates
interactions of Wag31 with other proteins. **(b.)** Left: Illustration showing Wag31 scaffolds bound to
cardiolipin in the membrane via N-terminal, tethering it and creating CL-microdomains at the poles.
The C-terminal of Wag31 bound to protein-partners, localize them to the membrane, facilitating several

membrane-centric processes such as PG and lipid synthesis, that ensure mycobacterial survival. The right panel shows the time-dependent consequences of Wag31 depletion i.e. early and late stage. The cells are bulged at one pole and start accumulating ILIs in the early stage and as the time progresses, the effect of Wag31 depletion becomes severe, rendering the cells round and full of ILIs. In the absence of Wag31, CL-microdomains become delocalised due to the loss of tethering, pulling away several membrane-associated proteins causing non-polar growth and loss of curvature.

Figure S1| Generation of Wag31 knockdown

(a.) Schematic showing the *wag31* locus in *Msm* and *Δwag31* and illustrating the strategy utilized to replace *wag31* locus with hygromycin resistance cassette. Primers used for knockout confirmation are indicated as F1, R1, F2, R2, F3 and R3. **(b.)** 1% agarose gels showing confirmatory PCRs performed with different primer sets. The first PCR, or the control PCR, was done using SepIVA (738 bp) primers showing amplicon at ~750bp. It indicates an equal amount of gDNA isolated from *Msm* and *Δwag31* and subsequently used for all PCRs. The next panel shows amplicons from the PCR performed with the F1-R1 pair, which bind to *wag31* locus (~819 bp). They yield an 819 bp amplicon in the case of *Msm* and 819 bp and an additional ~1.5 kb amplicon in the case of *Δwag31.* The third and fourth panels show amplicons amplified using the F2-R2 set (expected amplicon ~1.2 Kb) and F3-R3 set (expected amplicon ~0.9 Kb), which are expected only in the case of *Δwag31* confirming legitimate recombination at the native locus.

Fig.S2| Wag31 levels perturb lipid homeostasis

(a.) Whole Cell Lysates **(**WCL) were prepared from *Msm* and *Msm::wag31* (Table S1) treated with 5 μM IVN for 12 h, and 40 μg lysate was resolved on a 12% gel, transferred on Nitrocellulose membrane, and probed with either α-FLAG or α-rpoA antibody (loading control), respectively. **(b-g)** Bar graphs (Mean ± SEM) depicting the abundance of various lipids classes **(b.)** TAG, **(c.)** DAG, **(d.)** FFA, **(e.)** PE, **(f.)** PI, **(g.)** CL in *Msm* and *Msm::wag31* treated with 5 μM IVN for 12 h. Values on the Y-axis represent area under the curve (AUC) normalised to an internal standard. X-axis represent various species (chain length variations) of the above mentioned lipid classes. The experiment was performed with six biological replicates, and the abundances of identified lipid classes were normalized to *Msm*-IVN to yield relative abundances of lipids in *Msm::wag31* +IVN.

Figure S3| *Molecular interactions of wag31 with membrane proteins govern cellular homeostasis.*

(a.) Western blots showing FLAG-GFP (left panel) and FLAG-Wag31 (right panel) immunoprecipitations. 3 mg lysate from each group was processed for overnight FLAG immunoprecipitations, resolved on a 10% gel, and probed with α-Flag antibody. HC and LC are short for Heavy chain and light chain, respectively. **(b.)** Western blot showing expression of His-Wag31 used for the His pulldown experiments. **(c.)** Representative western blots demonstrating expression of 3XFLAG tagged-MmpL4, -MmpS5 and -Msm2092. WCLs were prepared from 0.5 μM IVN induced *Msm::mmpL4*, *Msm::mmpS5, Msm::msm2092,* and 40 μg of each was resolved on a 10 % SDS-PAGE,

transferred onto a nitrocellulose membrane and probed with α-FLAG and α-RpoA (loading control). **(d.)**
Representative western blots showing the absence of interaction of Wag31 with both MmpL4 and
MmpS5. Msm2092, a Wag31 interactor was used as a positive control. *E. coli* and *Msm* lysates
represented in (b & c) incubated together and pulled down with Cobalt beads were resolved on a 10-
12 % SDS-PAGE, transferred onto nitrocellulose membrane and probed with α-His to detect the
pulldown and α-FLAG to detect the interaction. The experiment was performed independently twice. **(e.)**
Classification of cellular processes that occur in different cellular compartments of *Msm,* as shown in
(48)*.*
844
845 **Figure S4| *Molecular interactions wag31 with membrane proteins govern cellular homeostasis.***
846 The list of Wag31 interactors is classified according to functional categories based on the data available
847 in Mycobrowser. The colour of the textbox indicates localization of the proteins: Pink- membrane, blue-
848 cytoplasm, and core- green.
849
850 **Table S1**
851 Plasmids and strains used in the study.
852 **Table S2.** It depicts unique interacting partners found for Wag31.
853

Acknowledgments
We thank Dharani Kumar, SEST, University of Hyderabad for helping with imaging of SEM stubs; CCMB
for access to its Genomics, Advanced Microscopy, and Electron Microscopy facilities; VIDO, USask
for providing infrastructure to carry out some part of the work; Asis Kumar Khuntia for the generation of
pET28b-*wag31*$_{K20A}$ and pET28b-*wag31*$_N$. YK was supported by a Research Associateship from NIH
(Grant 6044-SC23-26). YK and ND acknowledge MITACS for the Globalink Research Award
(FR118569) which supported the work carried out at VIDO. ND also acknowledges support from
Canadian Institutes of Health Research (CIHR, funding reference number 185715). VIDO receives
operational funding from the Government of Saskatchewan through Innovation Saskatchewan and the
Ministry of Agriculture and from the Canada Foundation for Innovation through the Major Science
Initiatives Fund. DD thanks DBT for Senior Project Associateship (Grant no.:
BT/PR13522/COE/34/27/2015). Research in this publication was supported by SERB grant
(CRG/2018/001294), Govt. of India, and JC Bose Award (JCB/2019/000015), Govt. of India to VKN.
Work in the SSK laboratory was supported by the Swarna Jayanti Fellowship to SSK by SERB, Govt.
of India (Grant: SB/SJF/2021-22/01). AC was supported by a Senior research fellowship from CSIR,
India. HK thanks the Indian Institute of Science Education and Research, Pune, for a graduate
fellowship. TP thanks the Howard Hughes Medical Institute for funding support.
871

References

1.   E. F. Bi, J. Lutkenhaus, FtsZ ring structure associated with division in Escherichia coli. *Nature* **354**, 161-
164 (1991).

2.   P. A. de Boer, R. E. Crossley, A. R. Hand, L. I. Rothfield, The MinD protein is a membrane ATPase
required for the correct placement of the Escherichia coli division site. *Embo j* **10**, 4371-4380 (1991).

3.   Z. Hu, A. Mukherjee, S. Pichoff, J. Lutkenhaus, The MinC component of the division site selection system
in Escherichia coli interacts with FtsZ to prevent polymerization. *Proc Natl Acad Sci U S A* **96**, 14819-
14824 (1999).

4.   D. M. Raskin, P. A. de Boer, The MinE ring: an FtsZ-independent cell structure required for selection of
the correct division site in E. coli. *Cell* **91**, 685-694 (1997).

5.   H. Meinhardt, P. A. de Boer, Pattern formation in Escherichia coli: a model for the pole-to-pole oscillations
of Min proteins and the localization of the division site. *Proc Natl Acad Sci U S A* **98**, 14202-14207 (2001).

6.   C. L. Woldringh *et al.*, Role of the nucleoid in the toporegulation of division. *Res Microbiol* **141**, 39-49
(1990).

7.   M. M. Logsdon, B. B. Aldridge, Stable Regulation of Cell Cycle Events in Mycobacteria: Insights From
Inherently Heterogeneous Bacterial Populations. *Front Microbiol* **9**, 514 (2018).

8.   J. H. Cha, G. C. Stewart, The divIVA minicell locus of Bacillus subtilis. *J Bacteriol* **179**, 1671-1683 (1997).

9.   D. H. Edwards, J. Errington, The Bacillus subtilis DivIVA protein targets to the division septum and controls
the site specificity of cell division. *Mol Microbiol* **24**, 905-915 (1997).

10. C. M. Kang, S. Nyayapathy, J. Y. Lee, J. W. Suh, R. N. Husson, Wag31, a homologue of the cell division
protein DivIVA, regulates growth, morphology and polar cell wall synthesis in mycobacteria. *Microbiology*
(Reading) **154**, 725-735 (2008).

11. C. Jani *et al.*, Regulation of polar peptidoglycan biosynthesis by Wag31 phosphorylation in mycobacteria.
BMC Microbiol **10**, 327 (2010).

12. M. Letek *et al.*, DivIVA is required for polar growth in the MreB-lacking rod-shaped actinomycete
Corynebacterium glutamicum. *J Bacteriol* **190**, 3283-3292 (2008).

13. N. R. Thanky, D. B. Young, B. D. Robertson, Unusual features of the cell cycle in mycobacteria: polar-
restricted growth and the snapping-model of cell division. *Tuberculosis (Edinb)* **87**, 231-236 (2007).

14. K. J. Kieser, E. J. Rubin, How sisters grow apart: mycobacterial growth and division. *Nat Rev Microbiol*
12, 550-562 (2014).

15. L. Nguyen *et al.*, Antigen 84, an effector of pleiomorphism in Mycobacterium smegmatis. *J Bacteriol* **189**,
7896-7910 (2007).

16. L. R. Hammond, M. L. White, P. J. Eswara, ¡vIVA la DivIVA! *J Bacteriol* **201** (2019).

17. H. Stahlberg *et al.*, Oligomeric structure of the Bacillus subtilis cell division protein DivIVA determined by
transmission electron microscopy. *Mol Microbiol* **52**, 1281-1290 (2004).

18. K. N. Muchová *et al.*, Oligomerization of the Bacillus subtilis division protein DivIVA. *Microbiology*
(Reading) **148**, 807-813 (2002).

19. M. Letek *et al.*, DivIVA uses an N-terminal conserved region and two coiled-coil domains to localize and
sustain the polar growth in Corynebacterium glutamicum. *FEMS Microbiol Lett* **297**, 110-116 (2009).

20. M. D. Rigden *et al.*, Identification of the coiled-coil domains of Enterococcus faecalis DivIVA that mediate
oligomerization and their importance for biological function. *J Biochem* **144**, 63-76 (2008).

21. K. Choukate *et al.*, Higher order assembling of the mycobacterial polar growth factor DivIVA/Wag31. *J*
Struct Biol **209**, 107429 (2020).

22. K. Flärdh, Essential role of DivIVA in polar growth and morphogenesis in Streptomyces coelicolor A3(2).
Mol Microbiol **49**, 1523-1536 (2003).

23. D. Fadda *et al.*, Streptococcus pneumoniae DivIVA: localization and interactions in a MinCD-free context.
J Bacteriol **189**, 1288-1298 (2007).

24. K. G. Kaval, S. Halbedel, Architecturally the same, but playing a different game: the diverse species-
specific roles of DivIVA proteins. *Virulence* **3**, 406-407 (2012).

25. H. B. Thomaides, M. Freeman, M. El Karoui, J. Errington, Division site selection protein DivIVA of Bacillus
subtilis has a second distinct function in chromosome segregation during sporulation. *Genes Dev* **15**,
1662-1673 (2001).

26. K. G. Kaval, S. Hauf, J. Rismondo, B. Hahn, S. Halbedel, Genetic Dissection of DivIVA Functions in
Listeria monocytogenes. *J Bacteriol* **199** (2017).

27.   K. G. Kaval, J. Rismondo, S. Halbedel, A function of DivIVA in Listeria monocytogenes division site
927   selection. *Mol Microbiol* **94**, 637-654 (2014).

928   28.   X. Meniche *et al.*, Subpolar addition of new cell wall is directed by DivIVA in mycobacteria. *Proc Natl Acad
929   Sci U S A* **111**, E3243-3251 (2014).

930   29.   E. S. Melzer, C. E. Sein, J. J. Chambers, M. S. Siegrist, DivIVA concentrates mycobacterial cell envelope
931   assembly for initiation and stabilization of polar growth. *Cytoskeleton (Hoboken)* **75**, 498-507 (2018).

932   30.   W. X. Xu *et al.*, The Wag31 protein interacts with AccA3 and coordinates cell wall lipid permeability and
933   lipophilic drug resistance in Mycobacterium smegmatis. *Biochem Biophys Res Commun* **448**, 255-260
934   (2014).

935   31.   M. T. M. Hannebelle *et al.*, A biphasic growth model for cell pole elongation in mycobacteria. *Nat Commun*
936   **11**, 452 (2020).

937   32.   I. Santi, N. Dhar, D. Bousbaine, Y. Wakamoto, J. D. McKinney, Single-cell dynamics of the chromosome
938   replication and cell division cycles in mycobacteria. *Nat Commun* **4**, 2470 (2013).

939   33.   P. Mukherjee *et al.*, Novel role of Wag31 in protection of mycobacteria under oxidative stress. *Mol
940   Microbiol* **73**, 103-119 (2009).

941   34.   N. Habibi Arejan *et al.*, Polar protein Wag31 both activates and inhibits cell wall metabolism at the poles
942   and septum. *Front Microbiol* **13**, 1085918 (2022).

943   35.   Y. S. Morita *et al.*, Compartmentalization of lipid biosynthesis in mycobacteria. *J Biol Chem* **280**, 21645-
944   21652 (2005).

945   36.   J. M. Hayashi *et al.*, Spatially distinct and metabolically active membrane domain in mycobacteria. *Proc
946   Natl Acad Sci U S A* **113**, 5400-5405 (2016).

947   37.   A. García-Heredia *et al.*, Membrane-partitioned cell wall synthesis in mycobacteria. *Elife* **10** (2021).

948   38.   V. Singh *et al.*, Identification of aminopyrimidine-sulfonamides as potent modulators of Wag31-mediated
949   cell elongation in mycobacteria. *Mol Microbiol* **103**, 13-25 (2017).

950   39.   J. Daniel, H. Maamar, C. Deb, T. D. Sirakova, P. E. Kolattukudy, Mycobacterium tuberculosis uses host
951   triacylglycerol to accumulate lipid droplets and acquires a dormancy-like phenotype in lipid-loaded
952   macrophages. *PLoS Pathog* **7**, e1002093 (2011).

953   40.   C. Barisch, P. Paschke, M. Hagedorn, M. Maniak, T. Soldati, Lipid droplet dynamics at early stages of
954   Mycobacterium marinum infection in Dictyostelium. *Cell Microbiol* **17**, 1332-1349 (2015).

955   41.   R. Bartz *et al.*, Lipidomics reveals that adiposomes store ether lipids and mediate phospholipid traffic. *J
956   Lipid Res* **48**, 837-847 (2007).

957   42.   K. Tauchi-Sato, S. Ozeki, T. Houjou, R. Taguchi, T. Fujimoto, The surface of lipid droplets is a
958   phospholipid monolayer with a unique Fatty Acid composition. *J Biol Chem* **277**, 44507-44512 (2002).

959   43.   A. Griego, T. Douché, Q. G. Gianetto, M. Matondo, G. Manina, RNase E and HupB dynamics foster
960   mycobacterial cell homeostasis and fitness. *iScience* **25**, 104233 (2022).

961   44.   R. Priyadarshini, M. A. de Pedro, K. D. Young, Role of peptidoglycan amidases in the development and
962   morphology of the division septum in Escherichia coli. *J Bacteriol* **189**, 5334-5347 (2007).

963   45.   C. Heidrich *et al.*, Involvement of N-acetylmuramyl-L-alanine amidases in cell separation and antibiotic-
964   induced autolysis of Escherichia coli. *Mol Microbiol* **41**, 167-178 (2001).

965   46.   D. Mengin-Lecreulx, L. Texier, M. Rousseau, J. van Heijenoort, The murG gene of Escherichia coli codes
966   for the UDP-N-acetylglucosamine: N-acetylmuramyl-(pentapeptide) pyrophosphoryl-undecaprenol N-
967   acetylglucosamine transferase involved in the membrane steps of peptidoglycan synthesis. *J Bacteriol*
968   **173**, 4625-4636 (1991).

969   47.   A. H. Freeman *et al.*, Arginine methylation sites on SepIVA help balance elongation and septation in
970   Mycobacterium smegmatis. *Mol Microbiol* **119**, 208-223 (2023).

971   48.   J. Zhu *et al.*, Spatiotemporal localization of proteins in mycobacteria. *Cell Rep* **37**, 110154 (2021).

972   49.   Z. He, J. De Buck, Cell wall proteome analysis of Mycobacterium smegmatis strain MC2 155. *BMC
973   Microbiol* **10**, 121 (2010).

974   50.   K. Choukate, B. Chaudhuri, Structural basis of self-assembly in the lipid-binding domain of mycobacterial
975   polar growth factor Wag31. *IUCrJ* **7**, 767-776 (2020).

976   51.   M. A. Oliva *et al.*, Features critical for membrane binding revealed by DivIVA crystal structure. *Embo j* **29**,
977   1988-2001 (2010).

978   52.   L. D. Renner, D. B. Weibel, Cardiolipin microdomains localize to negatively curved regions of Escherichia
979   coli membranes. *Proc Natl Acad Sci U S A* **108**, 6264-6269 (2011).

53. P. M. Oliver *et al.*, Localization of anionic phospholipids in Escherichia coli cells. *J Bacteriol* **196**, 3386-
3398 (2014).

54. M. I. Garcia Fernandez, D. Ceccarelli, U. Muscatello, Use of the fluorescent dye 10-N-nonyl acridine
orange in quantitative and location assays of cardiolipin: a study on different experimental models. *Anal*
Biochem **328**, 174-180 (2004).

55. E. Mileykovskaya, W. Dowhan, Visualization of phospholipid domains in Escherichia coli by using the
cardiolipin-specific fluorescent dye 10-N-nonyl acridine orange. *J Bacteriol* **182**, 1172-1175 (2000).

56. J. M. Petit, A. Maftah, M. H. Ratinaud, R. Julien, 10N-nonyl acridine orange interacts with cardiolipin and
allows the quantification of this phospholipid in isolated mitochondria. *Eur J Biochem* **209**, 267-273 (1992).

57. C. Carel *et al.*, Mycobacterium tuberculosis proteins involved in mycolic acid synthesis and transport
localize dynamically to the old growing pole and septum. *PLoS One* **9**, e97148 (2014).

58. N. H. Arejan, D. R. Czapski, J. A. Buonomo, C. C. Boutte, MmpL3, Wag31 and PlrA are involved in
coordinating polar growth with peptidoglycan metabolism and nutrient availability. *bioRxiv*
10.1101/2024.04.29.591792 (2024).

59. G. P. Salmond, J. F. Lutkenhaus, W. D. Donachie, Identification of new genes in a cell envelope-cell
division gene cluster of Escherichia coli: cell envelope gene murG. *J Bacteriol* **144**, 438-440 (1980).

60. R. Lenarcic *et al.*, Localisation of DivIVA by targeting to negatively curved membranes. *Embo j* **28**, 2272-
2282 (2009).

61. G. Carranza *et al.*, Cardiolipin plays an essential role in the formation of intracellular membranes in
Escherichia coli. *Biochim Biophys Acta Biomembr* **1859**, 1124-1132 (2017).

62. V. A. Gold *et al.*, The action of cardiolipin on the bacterial translocon. *Proc Natl Acad Sci U S A* **107**,
10044-10049 (2010).

63. W. S. Yeo, S. Dyzenhaus, V. J. Torres, S. R. Brinsmade, T. Bae, Regulation of Bacterial Two-Component
Systems by Cardiolipin. *Infect Immun* **91**, e0004623 (2023).

64. S. P. Ouellette *et al.*, Localized cardiolipin synthesis is required for the assembly of MreB during the
polarized cell division of Chlamydia trachomatis. *PLoS Pathog* **18**, e1010836 (2022).

65. J. C. van Kessel, G. F. Hatfull, Recombineering in Mycobacterium tuberculosis. *Nat Methods* **4**, 147-152
(2007).

66. A. Parikh *et al.*, Development of a new generation of vectors for gene expression, gene replacement, and
protein-protein interaction studies in mycobacteria. *Appl Environ Microbiol* **79**, 1718-1729 (2013).

67. G. P. Jose, S. Gopan, S. Bhattacharyya, T. J. Pucadyil, A facile, sensitive and quantitative membrane-
binding assay for proteins. *Traffic* **21**, 297-305 (2020).

68. J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682
(2012).

69. H. Schägger, Tricine-SDS-PAGE. *Nat Protoc* **1**, 16-22 (2006).

70. A. Kapopoulou, J. M. Lew, S. T. Cole, The MycoBrowser portal: a comprehensive and manually annotated
resource for mycobacterial genomes. *Tuberculosis (Edinb)* **91**, 8-13 (2011).

71. K. Tunyasuvunakool *et al.*, Highly accurate protein structure prediction for the human proteome. *Nature*
596, 590-596 (2021).

1019

Figure 1

a.

New Pole · Septum · Old Pole · ACC complex · New cell wall deposition · Depletion of Wag31 · Septum

cell wall precursors · Wag31 · Septum

b.

Msm - ATc · *Msm* + ATc · Δ*wag31* - ATc · Δ*wag31* + ATc

43 — α-Wag31

54 — 43 — α-rpoA

c.

Msm - ATc · *Msm* + ATc · Δ*wag31* - ATc · Δ*wag31* + ATc

d.

Msm + ATc · Δ*wag31* - ATc · Δ*wag31* + ATc

CFU log$_{10}$/ml

0 h · 12 h

***   **

e.

Msm · Δ*wag31* - ATc · Δ*wag31* + ATc

5 μm

f.

Rod · Bulged · Round

g.

Percentage of cells

Rod · Bulged · Round

Msm · Δ*wag31* - ATc · Δ*wag31* + ATc

i.

Percentage of lipid bodies

Lipid bodies 0-2 · Lipid bodies 3-5 · Lipid bodies >5

Msm · Δ*wag31* - ATc · Δ*wag31* + ATc

h.

Msm · Δ*wag31* - ATc · Δ*wag31* + ATc

500 nm

Figure 2

Figure 3

a.

His-Wag31/His-Wag31mut + Liposomes (+ 1% RFL) → RFL labelled liposome membrane → +UV → RFL cross-linked protein → PAGE → -UV +UV

b.

His-Wag31 — Coomassie
His-Wag31 — Fluorescence

Liposome: - PE PG CL

kDa: 250, 150, 100, 75, 50, 37, 25, 20

c.

CL-liposomes / no liposomes

Wag31ΔN
Wag31K20A
Wag31

kDa: 150, 100, 75, 50, 37, 25, 20

Coomassie Fluorescence Coomassie Fluorescence

d.

Fluorescence Intensity (x 10³)

Wag31, Wag31K20A, Wag31ΔN

** / ***

e.

CL-DiI-Liposomes - Wag31
CL-DiI-Liposomes + Wag31
CL-DiI-Liposomes + Wag31
Merged
10μm

f.

PE Liposomes / CL Liposomes

10μm

His-Wag31K20A
His-Wag31
His-Wag31
His-Wag31ΔN

g.

Particle Intensity (x 10³)

PE Liposomes: -, Wag31
CL Liposomes: -, Wag31, Wag31K20A, Wag31ΔN

Figure 5

Figure 6

Figure7

Figure 8

a.

b.

Figure S1

a.

b.

Figure S3

a.

Msm::pST-GFP
Heavy chain
FLAG-GFP
Light Chain
α-FLAG

Δwag31::pST-Wag31
Heavy chain
Flag-Wag31
Light chain
α-FLAG

b.

Control / His-Wag31
α-His
α-RpoA

c.

Msm / MmpL4
Msm / MmpS5
Msm / Msm2092
α-FLAG
α-RpoA

d.

MmpL4 / Msm2092
His-Wag31 + +
Control + +

MmpS5 / Msm2092
His-Wag31 + +
Control + +

α-FLAG
α-His

e.

Membrane:
- Energy production and conversion
- Cell cycle control, cell divison and chromosome partitioning
- Cell wall membrane and envelope biogenesis
- Intracellular trafficking, secretion, and vesicular transport

Core:
- Translation, ribosomal structure and biogenesis
- Transcription
- Replication, recombination and repair

Cytoplasm:
- Amino acid transport and metabolism

Figure S4

Cell wall & cell processes

- MSMEG_0402/mps2: linear gramicidin synthetase subunit D
- MSMEG_6099: conserved transmembrane protein
- MSMEG_4227: murG: undecaprenyldiphospho-muramoylpentapeptide beta-N-acetylglucosaminyltransferase
- MSMEG_2092: D-aminopeptidase
- MSMEG_6863: class II aldolase/adducin domain protein
- MSMEG_2416: SepIVA: DivIVA-domain containing protein
- MSMEG_1046: ABC-type molybdenum transport system
- MSMEG_0386: NAD dependent epimerase/dehydratase family protein
- MSMEG_3654: ATPase SecA2
- MSMEG_4958: diaminopimelate decarboxylase
- MSMEG_0919: heparin-binding hemagglutinin
- MSMEG_1954: ABC1 family protein
- MSMEG_2326: ABC-type molybdenum transport system

Lipid metabolism

- MSMEG_1813: propionyl-CoA carboxylase beta chain
- MSMEG_0372: oxidoreductase, short chain dehydrogenase/ reductase family protein
- MSMEG_2393: NAD-dependent glycerol-3-phosphate dehydrogenase
- MSMEG_2080: putative acyl-CoA dehydrogenase
- MSMEG_0912: 3-hydroxybutyryl-CoA dehydrogenase
- MSMEG_2648: Sfp-type phosphopantetheinyl transferase
- MSMEG_0406: fadE5: acyl-coA-dehydrogenase
- MSMEG_5199: putative acyl-CoA dehydrogenase
- MSMEG_4709: enoyl-CoA hydratase
- MSMEG_5197: long-chain specific acyl-CoA dehydrogenase
- MSMEG_6512: acyl-CoA dehydrogenase domain protein
- MSMEG_6511: acyl-CoA dehydrogenase domain protein
- MSMEG_6208: palmitoyl-CoA hydrolase
- MSMEG_5198: carnitinyl-CoA dehydratase

Replication, recombination & repair

- MSMEG_3596: ATPase- DNA replication
- MSMEG_2174: superfamily protein I DNA or RNA helicase
- MSMEG_1061: phosphohydrolase
- MSMEG_2768: OB-fold nucleic acid binding domain protein
- MSMEG_4450: alpha/beta hydrolase
- MSMEG_1930: DEAD/DEAH box helicase
- MSMEG_5156: base excision DNA repair protein, HhH-GPD family protein

Translation, ribosome structure & biogenesis

- MSMEG_4626: rne: ribonuclease, Rne/Rng family protein
- MSMEG_2931: thrS: threonyl-tRNA synthetase
- MSMEG_3003: aspS: aspartyl-tRNA synthetase
- MSMEG_4959: argS: arginyl-tRNA synthetase
- MSMEG_3777: pheT: phenylalanyl-tRNA synthetase, beta subunit

Regulatory proteins

- MSMEG_6219: moxR2: ATPase family protein associated with various cellular activities (AAA)
- MSMEG_4659: GntR-family protein transcriptional regulator
- MSMEG_3363: regulatory protein, TetR
- MSMEG_0035: fhaA: FHA domain protein

Unknown

- MSMEG_3785: PfkB-family protein carbohydrate kinase
- MSMEG_6783: integral membrane protein
- MSMEG_4652: D-oliose 4-ketoreductase
- MSMEG_1679: AmiB

Miscellaneous

- MSMEG_6514: trehalose synthase-fused probable maltokinase

Intermediary metabolism & respiration

- MSMEG_4915: alpha-glucan phosphorylase family protein
- MSMEG_0502: glucosidase
- MSMEG_6459: ferredoxin-dependent glutamate synthase 1
- MSMEG_3138: trx: thioredoxin
- MSMEG_0594: iron-sulfur cluster binding protein
- MSMEG_5427:prs: ribose-phosphate pyrophosphokinase
- MSMEG_2611: pyridine nucleotide-disulphide oxidoreductase family protein
- MSMEG_5937: bphl-2: 4-hydroxy-2-oxovalerate aldolase
- MSMEG_3051: gmk: guanylate kinase
- MSMEG_3044: pyrC: dihydroorotase
- MSMEG_1037: adhc1: alcohol dehydrogenase, zinc-containing
- MSMEG_4294: glnA: glutamine synthetase, type I
- MSMEG_3632: gnd: 6-phosphogluconate dehydrogenase, decarboxylating
- MSMEG_3100: opcA: OpcA protein: Putative OXPP cycle protein OpcA
- MSMEG_3894: prcA: proteasome alpha subunit
- MSMEG_2379: leuB: 3-isopropylmalate dehydrogenase
- MSMEG_6874: aldehyde dehydrogenase
- MSMEG_6114: ppa: inorganic pyrophosphatase
- MSMEG_1073: oxidoreductase, short-chain dehydrogenase/reductase family protein
- MSMEG_1825: rmlD: dTDP-4-dehydrorhamnose reductase
- MSMEG_3746: pyrG: CTP synthase
- MSMEG_4272: HesB/YadR/YfhF family protein
- MSMEG_3451: short-chain dehydrogenase/reductase, NAD(P)H-binding
- MSMEG_5592: Luciferase like monooxygenase
- MSMEG_0220; monoglyceride lipase
- MSMEG_3880: nitrilase/cyanide hydratase and apolipoprotein N-acyltransferase
- MSMEG_5903: oxidoreductase, short chain dehydrogenase/reductase family protein
- MSMEG_0759: purA: CTP synthase
- MSMEG_6598: ribH: 6,7-dimethyl-8-ribityllumazine synthase
- MSMEG_6351: histidinol-phosphate aminotransferase
- MSMEG_4528: cysH: phosphoadenosine phosphosulfate reductase
- MSMEG_4276: ilvE: branched-chain amino acid aminotransferase
- MSMEG_1670: sdhA: succinate dehydrogenase, flavoprotein subunit
- MSMEG_0539: transcriptional regulator, Crp/Fnr family protein
- MSMEG_4258: trpD: anthranilate phosphoribosyltransferase
- MSMEG_2357: cysteine desulfurase
- MSMEG_3207: hisB: imidazoleglycerol-phosphate dehydratase
- cobyrinic Acid a,c-diamide synthase
- alpha oxoglutarate ferredoxin oxidoreductase, beta subunit
- short chain dehydrogenase
- fructose-1,6-bisphosphatase, class II
- MSMEG_2617: cobB: cobyrinic Acid a,c-diamide synthase
- MSMEG_4645: orB: alpha oxoglutarate ferredoxin oxidoreductase, beta subunit
- MSMEG_3619: short chain dehydrogenase
- MSMEG_5239: glpX: fructose-1,6-bisphosphatase, class II
- MSMEG_3071: ribE: riboflavin synthase, alpha subunit
- MSMEG_0793: thiG: thiazole biosynthesis protein
- MSMEG_0956: hemB: delta-aminolevulinic acid dehydratase
- MSMEG_3219: indole-3-glycerol phosphate synthase
- MSMEG_0239: O-acetylhomoserine/O-acetylserine sulfhydrylase
- MSMEG_5515: purH: bifunctional purine biosynthesis protein
- MSMEG_6256: asd: aspartate-semialdehyde dehydrogenase

Amino acid transport and metabolism

- MSMEG_5211: aminotransferase class-III

Conserved hypotheticals

- MSMEG_2436; MSMEG_0897
- MSMEG_6941; MSMEG_3309
- MSMEG_2481; MSMEG_2434
- MSMEG_3148; MSMEG_6901
- MSMEG_6499; MSMEG_1340
- MSMEG_1514; MSMEG_5691
- MSMEG_3942; MSMEG_4282
- MSMEG_1049; MSMEG_3024
- MSMEG_3244; MSMEG_0240
- MSMEG_2685; MSMEG_4193
- MSMEG_2329

Membrane

Cytoplasm

Core