1 **Wag31, a membrane tether, is crucial for lipid homeostasis in mycobacteria**

2 Yogita Kapoor[1,2], Himani Khurana[3], Arnab Chakraborty[3], Debatri Dutta[1], Anshu Priya[2,4], Archana

3 Singh[2,4], Siddhesh S. Kamat[3], Neeraj Dhar,[5,6,] Thomas J. Pucadyil[3], Vinay Kumar Nandicoori[1,2,7]

1. CSIR-Centre for Cellular and Molecular Biology (CSIR-CCMB), Hyderabad, 500007
2. AcSIR, Ghaziabad-201002
3. Indian Institute of Science Education and Research, Pune, 411008
4. CSIR-Institute of Genomics and Integrative Biology (CSIR-IGIB), New Delhi, 110025
5. Vaccine and Infectious Disease Organisation, University of Saskatchewan, Canada, S7N 5E3
6. Department of Biochemistry, Microbiology and Immunology, University of Saskatchewan, Canada S7N 5E5
7. National Institute of Immunology, New Delhi 110067

Address correspondence to vinaykn@ccmb.res.in, vinaykn@nii.ac.in

Keywords: Wag31, intracellular lipid inclusions (ILIs), lipid homeostasis, membrane-tether

Abstract

The mycobacterial cytoskeletal protein Wag31 is necessary for maintaining cell shape and
directing cellular growth and elongation. Wag31 has a characteristic N-terminal DivIVA-domain and a
C-terminal coiled-coil domain. While the role of Wag31 in polar elongation is known, there is limited
mechanistic insight on how it orchestrates growth and elongation. In this report, we delineate roles of
the N- and C-terminal domains of Wag31 using genetics, state-of-the-art multi-omics, biochemical, and
imaging approaches. We show that Wag31 predominantly interacts with several membrane-associated
proteins involved in lipid metabolism, cell wall synthesis and division. Native levels of Wag31 are critical
for the maintenance and distribution of membrane lipids. Both depletion and overexpression of Wag31
perturbs lipid homeostasis, leading to the formation of intracellular lipid inclusions (ILIs). Protein-lipid
crosslinking and imaging studies reveal that purified Wag31 can bind and effectively tether Cardiolipin
(CL)-containing liposomes. Despite retaining its ability to interact with partner proteins, the DivIVA
domain-deleted Wag31 mutant shows defects in liposome tethering *in vitro* and non-polar localization
of CL *in vivo*, which eventually causes lethality. Our study suggests that membrane tethering 'licenses'
Wag31 to form scaffolds that help orchestrate protein-lipid and protein-protein interactions necessary
for mycobacterial growth and survival.

Introduction

Cell division is a critical event in the life of a bacterium as it is the prime mode of bacterial growth and reproduction. Hence, the process of their division is tightly knitted. The key process in bacterial cell division, i.e., assembly of the FtsZ ring at the mid-cell (1), is tightly regulated by the Min and Nucleoid occlusion systems (2-6). Mycobacterium, which comprises clinically relevant pathogens- *Mycobacterium tuberculosis (Mtb)*, *Mycobacterium leprae,* and *Mycobacterium abscessus* lacks both the regulators of FtsZ ring placement (7). Instead, they encode for a DivIVA protein, usually encoded by Firmicutes and other Actinobacteria (8-11). It is necessary that the bacterial shape, crucial for normal cellular physiology, is maintained during cell division. Rod-shaped bacteria such as *Escherichia coli* and *Bacillus subtilis* encode MreB and multiple MreB-like proteins that sustain the rod morphology of cells (12). *Mycobacterial spp.* lack MreB homologs (13, 14). The DivIVA protein of *Mtb* called Wag31 is known to regulate both cellular growth and morphology (10, 15), which makes it an exciting candidate to investigate.

DivIVA proteins were discovered three decades ago during the study of cell division mutants of *B. subtilis* (16). DivIVA proteins are rich in coiled-coil domains (17-21), which enable them to form higher oligomeric structures and, hence, scaffolds, making them suitable adaptor proteins (22, 23). They are known to play diverse roles in various species (24), e.g., they play dual roles of septum positioning and chromosome partitioning in *B. subtilis* (9, 25); cell-separation/resolution post-cell-division in *Listeria monocytogenes* (26, 27); pole formation and maturation in *Streptococcus spp.* (Fadda, 2007) In *Mycobacterium spp.*, Wag31 is an essential protein that senses negative membrane curvature found at poles and septum and localizes there (28). It plays roles in cellular elongation and septation in concert with other divisome and elongasome components (10, 15, 29). It is specifically found in higher amounts at the old pole, where it directs polar elongation by recruiting the acyl CoA carboxylase (ACCase) complex involved in fatty acid, mycolic acid, and methyl-branched chain lipid precursor synthesis (28, 30-32). It localizes to the septum after cytokinesis is over, perhaps for septal synthesis via its interactions with FtsI (32-34), though the process remains largely unknown. Consequently, loss of Wag31 leads to cell division defects- loss of polar elongation, which renders the cells 'round' in shape that occurs due to affected peptidoglycan (PG) synthesis (10, 15, 28). Though Wag31 participates in coordinating PG metabolism, the mechanism or the protein network by which it does so remains unknown.

Wag31 has also been associated with the maintenance of the intracellular membrane domain (IMD), a specialized compartment for lipid synthesis, localized predominantly at the old pole in consonance with pole elongation and spottily all around the plasma membrane where it may function to repair membrane defects (35-37). The loss of Wag31 leads to the delocalization of IMD and its protein markers, such as MurG and GlfT2, but it is not understood whether Wag31 plays a direct role in membrane-partitioning or it is just a consequence of localized membrane synthesis (34, 37). The role of Wag31in PG and lipid synthesis that drives cell division remains unclear. Despite numerous studies, the details of how Wag31 orchestrates mega-complexes at the pole or partitions the membrane into two to govern cellular elongation remain elusive.

Here, we endeavoured to address the lacunae in the functionality of Wag31 by asking the following questions- i) what are the ultrastructural changes that occur in mycobacteria due to the absence of Wag31, ii) do Wag31 deficient cells undergo transcriptional rewiring to adjust to cell division defects, iii) is the change in cellular morphology from 'rod-to-round' accompanied by altered lipid levels, iv) what is the interactome of Wag31, v) what are the functions of its N and C-terminal domains and finally vi) what is its relationship with the mycobacterial membrane. The present study shows that Wag31 levels are critical for maintaining lipid homeostasis in mycobacteria. Our results demonstrate that Wag31 binds and tethers the membrane together to orchestrate membrane-localized processes essential for maintaining normal cellular morphology and physiology.

  Results

Loss of Wag31 leads to the formation of intracellular lipid inclusions (ILIs)

In the pursuit of bridging the knowledge gap existing in the functionality of Wag31 (Fig. 1a), we

generated a *Wag31* conditional mutant in *Msm*, by integrating a copy of *wag31* under tetracycline (ATc-

tet off) regulation at the L5 locus and replacing the native copy with a hygromycin cassette (Fig. S1a).

The mutant, *Δwag31*, thus generated, was confirmed by PCRs (Fig. S1b) and Western blotting (Fig.

1b). The lack of Wag31 led to a drop in the turbidity of culture (Fig. 1c). Analysis of bacillary survival *in*

vitro by enumerating CFUs 12h post-ATc addition, revealed a 2 log fold difference in the survival of

Δwag31 (Fig. 1d). Given the essential role of Wag31 in causing polar growth of the cell, we investigated

the effect of depletion of Wag31 on bacterial cellular morphology using electron microscopy (EM).

Scanning EM (SEM) suggested that the absence of Wag31 caused a transition in the morphology of

the cells from rod to round, as previously documented (10, 15, 38). Apart from round morphology (~36

%), we also observed a rod-to-round transformation stage in which cells were bulged at one pole with

the other pole intact (Fig. 1e-g). To obtain detailed insights, we performed transmission EM (TEM),

which divulges information on the internal structure of cells. We discovered spherical, electron-lucent

bodies (indicated by an arrowhead in Fig.1h) in the cytosol of all the samples. While *Msm* and *Δwag31*-

ATc harbored a few (~2-3) electron-lucent bodies, their numbers were drastically higher in the case of

Δwag31+ATc (Fig. 1h). The cytosol of *Δwag31*+ATc was filled with multiple variable-sized bodies which

resembled Lipid bodies that *Mycobacterium tuberculosis* (*Mtb)* accumulates inside the host (39, 40). As

the term 'lipid bodies/lipid droplets' is used specifically for the lipids acquired by *Mtb* from the host cell,

we referred to them as intracellular lipid inclusions (ILIs). We categorized the cells based on the number

of ILIs they contained. Their distribution remained reasonably similar within *Msm* and *Δwag31*-ATc cells

for all the classes (Fig. 1i). Whereas *Δwag31* treated with ATc had as high as 85% cells harboring more

than 5 ILIs and very few cells with less than 5 ILIs (Fig. 1i). Our data, therefore, reports a novel

observation of lipid accumulation when Wag31 is depleted from mycobacteria.

Wag31 levels influence lipid homeostasis

Next, we set out to analyze if the ILIs are indeed lipidic in nature by staining cells with BODIPY

(493/503), a dye that preferentially partitions into neutral lipid droplets (41, 42). We stained either

untreated or ATc-treated *Msm* and *Δwag31* with BODIPY and visualized them on a microscope. ATc-

treated *Δwag31* cells displayed the highest level of BODIPY fluorescence, indicating the maximum

abundance of lipid bodies among the three strains (Fig. 2a, c). Further, we examined whether the

appearance of ILIs is associated just with the loss of Wag31 or is a consequence of deviations from the

native expression level of Wag31. Towards this, we overexpressed Wag31 episomally using an

isovaleronitrile (IVN) inducible system (*Msm::wag31*) (Fig. S2a) and stained the cells with BODIPY.

Unlike Wag31 depletion, which resulted in spherical cells (Fig. 1e), Wag31 overexpression resulted in

the bulging of cells from one pole (Fig. 2b). Additionally, the overexpression of Wag31 also led to lipid

aggregates in the cell with a few larger 'donut-like' aggregates in the bulged pole (Fig 2b,d).

Lipid accumulation in both cases led us to inspect whether lipid metabolism is closely tied to

Wag31 expression levels in mycobacteria. To investigate that possibility, total cellular lipids were

extracted from *Msm,* Wag31 depletion, and overexpression conditions and subjected to semi-

quantitative LC-MS analysis (illustrated in Fig. 2e). Lipidomic analysis of *Δwag31*+ATc revealed an
imbalance in the levels of neutral lipids, phospholipids, and free fatty acids in comparison to *Msm*-ATc
(Fig. 2f-g). *Δwag31*+ATc cells had a significant accumulation of unsaturated diacylglycerols (~3-5 fold)
and a very long chain containing (>C36) phosphatidylethanolamines (~5-32 fold). This was
accompanied by increased levels of different chain lengths (C18-C24) of free fatty acids (~1.7-4 fold)
and reduced levels of cardiolipins C70:0 & C70:1 (~3 fold). Wag31 overexpression, on the other hand,
led to a 2-fold increase in the level of different species of alpha-mycolic acids (C74-C78) as compared
to *Msm*+IVN (Fig. 2h), while other tested lipids remained unchanged (Fig. S2). Altogether, the results
presented above demonstrate that Wag31 levels are crucial for maintaining lipid homeostasis in
mycobacteria, the depletion and overexpression of which impact lipid metabolism differently.

Molecular interactions of Wag31 with membrane-associated proteins govern cellular
homeostasis.

We reasoned that Wag31, an adaptor protein, might maintain lipid homeostasis by interacting
with proteins involved in these pathways. To examine the protein interaction network of Wag31, we
performed an interactome analysis by utilizing *Δwag31*, which expresses FLAG-Wag31 integrated at
the L5 locus, and used FLAG-GFP as the control to differentiate from non-specific interactions. FLAG-
Wag31 and FLAG-GFP from biological triplicates were immunoprecipitated using whole cell lysates
from the log phase cells and subjected to mass spectrometry analysis (Fig. S3a & Table S2). This study
identified both direct and indirect protein-partners of Wag31. The proteins identified were categorized
based on their functions. It was found that Wag31 interacts with proteins involved in cell-wall elongation
and division, lipid metabolism, intermediary metabolism and respiration and information pathways
(translation, replication) (Fig. 3a, Fig. S4). Interactions of Wag31 with AccA3 and AccD5, subunits of
the ACCase complex, and Rne or RNaseE (Msm4626), a membrane-associated protein, have been
reported previously (28, 43). Subunits of ACCase complex and Rne were among the top 5 hits in our
MS/MS data, validating our approach (Fig. 3a, marked with asterisk). The classification of the top 20
hits is represented in Fig. 3a. The highest number of protein interactors, expectedly, fell in the cell wall
and cell processes category (Fig. 3a). For the present study, we focussed on the interactors found in
the cell wall elongation and cell division and lipid metabolism functional categories to probe the cause
behind the lipid dysbiosis phenotype.

MurG, involved in PG metabolism (36, 44-46), and SepIVA, involved in regulating septation
and the spatial regulator of MurG (47) also surfaced in the interactome of Wag31. Msm1813, or acyl
CoA carboxylase (ACAD), crucial for nascent fatty acid precursor synthesis, was also enriched in the
interactome. Other lipid metabolism interactors of Wag31 included acyl-CoA dehydrogenase
(Msm2080), involved in lipid catabolism; GpsA, involved in phospholipid biosynthesis; and Msm0372,
involved in fatty acid biosynthesis. To validate the high-throughput data, we chose MurG, SepIVA and
Msm2092 and performed pull-down assays with His-tagged Wag31. Briefly, His-tagged Wag31 (Fig.
3b) was incubated with *Msm* lysates expressing 3X-FLAG tagged versions of the tested proteins (Fig.
3c) and pulled-down with the help of cobalt beads followed by a western blot analysis. Results obtained
confirmed the interaction of Wag31 with MurG, SepIVA and Msm2092, thus validating our data (Fig.
3d).

Since Wag31 is a spatiotemporal regulator of other proteins, we were intrigued to discover the
localisation of its interacting partners. We used the data curated by Zhu et al. (48) and determined that
Wag31 majorly interacts with membrane-associated proteins with a few exceptions (Fig. 3e, Fig. S3b).
The mycobacterial plasma membrane has been recently shown to have a cell wall-associated
compartment called PM-CW and a cell wall-free compartment called PM_f or IMD, which is the site for
lipid synthesis. Based on the available data (36, 48, 49), we found that Wag31 interacts more with IMD-
proteins than with PM-CW proteins (Fig. 3f.) In accordance with assigned roles (36), PM-CW residing
proteins in the interactome are involved in cell wall and cell processes and intermediary metabolism
and respiration. Similarly, IMD-homing proteins clustered in the lipid metabolism and cell wall synthesis
category. These results suggest that Wag31 maintains lipid homeostasis and cellular morphology
primarily through its interactions with the membrane proteins participating in such pathways.

Wag31 binds and tethers cardiolipin-containing membranes

Wag31 is a cytoplasmic protein that recognizes negative curvature regions in the cell, i.e., poles
and septum (28). It also forms higher-order structural assemblies at the poles (21, 50). However, the
precise mechanism by which Wag31 assemblies localize to the poles remains elusive. The predominant
interactions of Wag31 with membrane-associated proteins (Fig 3) were intriguing and led us to
investigate if it can directly bind to membranes. We reasoned that such membrane binding activity might
render it capable of acting as a scaffold that anchors and organizes other proteins, thereby modulating
their functions. To test this, we performed a lipid crosslinking-based PLiMAP assay (Fig. 4a). Purified
Wag31 was incubated with ~100 nm extruded liposomes containing each of the abundant classes of
bacterial lipids phosphatidylethanolamine (PE), phosphatidylglycerol (PG), or cardiolipin (CL).
Liposomes contained trace amounts of a reactive fluorescent lipid (RFL), which crosslinks with
membrane-bound proteins upon photoactivation. These assays revealed that Wag31 preferentially
binds CL- but not PG- or PE-containing liposomes (Fig. 4b). Both CL and PG are anionic lipids, and the
preferential binding of Wag31 to CL indicates that the binding cannot be attributed to a charge-based
interaction. Remarkably, the sensitive fluorescence-based detection revealed the presence of SDS-
resistant higher-order oligomers of membrane-bound Wag31 (Fig. 4b).

Wag31 is a homolog of the DivIVA protein from *B. subtilis.* The latter has been shown to bind
to the membrane via the N-terminal DivIVA domain (N-DivIVA$_{BS}$) and localize other cell division proteins
through its interactions with the C-terminal region (51, 52). However, the functional significance of the
N- & C- terminal domains of mycobacterial Wag31 has not been dissected. Considering that the Wag31
N-terminal region shares 42% sequence identity with N-DivIVA$_{BS}$ (50), we sought to inspect the effect
of disruption of a particular residue (K20) in the N-terminus and the deletion of the entire N-terminus,
encompassing the DivIVA domain (1-60 amino acids) on membrane binding (Fig. S5a). The K20
residue lies within a positively charged patch, which likely constitutes a membrane-associating surface
in the intertwined loop region of Wag31(50). Moreover, mutating the K20 residue has been shown to
decrease the cell length and elongation rate by 18% and 22%, respectively (34). The K20A mutation
showed a marginal defect in binding to CL-liposomes (Fig. 4c-d), indicating that K20 alone is insufficient
for membrane binding. However, Wag31$_{K20A}$ showed a marked defect in forming SDS-resistant higher-
order oligomers, as seen in the binding assays. Interestingly, deletion of the entire N-terminal domain

caused a significant defect in both membrane binding and forming SDS-resistant higher-order
oligomers (Fig. 4c-d). Surprisingly, binding was not entirely abolished, suggesting that the C-terminus
of Wag31 may also contribute to membrane binding. Taken together, our results indicate that Wag31
preferentially binds CL-containing membranes and that regions other than the N-terminal domain
contribute to membrane binding. The latter feature distinguishes Wag31 from DivIVA$_{BS}$, and Wag31
likely contains membrane binding sites located at both the N- and C-terminal regions, which might lead
to differences in the manner in which it interacts with the membrane and partner proteins compared to
DivIVA$_{BS}$.

To further confirm membrane binding, we imaged 100 nm-extruded CL-containing liposomes
with a membrane dye DiI in the presence of Wag31. We used a GFP-tagged Wag31 mixed with an
untagged Wag31 at a 1:10 molar ratio for these experiments. In the absence of Wag31, liposomes
appeared as faint but distinct fluorescent puncta (Fig. 4e). Surprisingly, in reactions containing Wag31,
we observed liposomes clustered into bright fluorescent puncta that were positive for GFP-Wag31 (Fig.
4e). The Wag31 tetramer has been modelled as a long rod-shaped protein filament (21, 50) and such
clustering of liposomes falls in congruence with the model. It indicates that these long filaments of
Wag31 act to tether liposomes. We then systematically evaluated this effect by testing liposomes of
different compositions and Wag31 mutants. Again, in the absence of Wag31, both PE- and CL-
containing liposomes appeared as faint but distinct fluorescent puncta (Fig. 4f). The addition of Wag31
caused clustering of CL-containing liposomes into bright foci (Fig. 4f), which is quantified by a particle
intensity analysis (Fig. 4g). This clustering was caused by membrane binding of Wag31 because PE-
containing liposomes that do not bind Wag31 (Fig. 4b) showed no clustering (Fig. 4f,g). As expected,
Wag31$_{\Delta 1-60}$ was defective in membrane binding (Fig. 4c,d) and showed a severe tethering capacity
defect (Fig. 4f,g). Surprisingly, Wag31$_{K20A}$, which showed a marginal defect in membrane binding (Fig.
4c,d), showed a marked defect in tethering capacity (Fig. 4f,g). Together, these results suggest that
membrane-binding and membrane-tethering are non-overlapping attributes in the structure of Wag31,
and the N-terminal DivIVA-domain is indispensable for membrane-tethering activity.

Tethering is crucial for the survival of mycobacteria

Results presented in Fig.4c indicate that Wag31 binds strongly to CL-containing membranes.
CL is a negatively charged inverted cone-shaped lipid that has been shown to be concentrated at the
sites of negative curvature in the bacterial cell, i.e., pole and septum (53, 54). Wag31 has been
demonstrated previously to localize at sites of negative curvature (28). Their similar localization pattern,
coupled with Wag31's affinity to bind to CL, led us to examine the effect of this binding *in vivo*. We
hypothesized that modulations in Wag31 levels might alter CL distribution in the cell. To examine this,
we utilized a fluorophore 10-N-Nonyl-acridine orange (NAO) that binds to anionic phospholipids (54,
55). We stained *Msm and* Wag31 depleted and overexpressing cells with NAO and imaged them using
a fluorescence microscope. While the localization pattern of NAO was polar in *Msm*+ATc and *Δwag31*-
ATc, there was a redistribution of red fluorescence to the cytosol and along the perimeter of the cell in
Δwag31+ATc (Fig.5a-b). We quantified the distribution of fluorescence using a line profile which yielded
peaks of intensity at the poles and a basal level of fluorescence along the lateral cell body for both
Msm+ATc and *Δwag31*-ATc (Fig.5b). Wag31 overexpression also led to a change in the distribution of

NAO from the poles and septum to 'donut-like' fluorescent foci in the cytosol (Fig.5c-d). Since our *in vitro* results indicate that Wag31 prefers to bind CL over PG (Fig. 4b), the change in the distribution of NAO fluorescence implies that the loss or overexpression of Wag31 causes the mislocalization of CL in mycobacterium.

Subsequently, we investigated the impact of tethering-deficient Wag31 mutants on cell survival. Wag31 has an N-terminal domain joined to the C-terminal via a linker region. The N-terminal domain contains a positively charged patch in the intertwined region (the region between two helices in the N-terminus) and membrane binding has been attributed to this region (50). This patch is lined with conserved arginine and lysine residues, K15, K20, and R21 (50) (Fig. 5e). As our *in vitro* results suggest the importance of K20 and the DivIVA-domain in tethering, we performed complementation assays with the tethering-defective mutants to examine cell survival and morphology. *Δwag31* was complemented with an integrative copy of either Wag31 or Wag31$_{K20A}$ or Wag31$_{\Delta1-60}$. *Δwag31*+ATc showed a two log-fold difference in survival and a 40 % decrease in the percentage of rod-shaped cells compared with *Δwag31*-ATc (Fig. 5f-g). Complementation with Wag31 and Wag31$_{K20A}$ rescued the loss of cell viability and restored the rod shape of the cells (Fig. 5f-g), whereas Wag31$_{\Delta1-60}$ behaved similar to *Δwag31*+ATc. It neither restored the rod shape of the cells nor reversed the survival defect (Fig. 5f-g). These results highlight the importance of tethering for sustaining mycobacterial morphology and survival.

Tethering is correlated to lipid levels and CL localization in mycobacteria

As described above, Wag31-mediated membrane tethering is an absolute for the survival of mycobacteria. Perturbations in lipid homeostasis and mislocalization of CL due to altered Wag31 levels hint at a correlation between its ability to function as a homotypic tether and maintain lipid composition and localization. To derive the mechanism behind it, we performed fluorescence imaging experiments with native and tethering defective mutants of Wag31. To investigate a role of tethering in lipid homeostasis in mycobacteria, we stained wildtype and complementation strains with BODIPY and quantified the fluorescence intensities. We observed comparable staining intensity in *Msm* +ATc and, *Δwag31* -ATc (Fig. 6a-b). *Δwag31*+ATc expectedly had the highest BODIPY staining intensity, indicating higher than normal lipid levels (Fig. 6a-b). Complementation with Wag31 fully restored the lipid levels to normal, whereas complementation with Wag31$_{K20A}$ couldn't restore the lipid levels completely (Fig. 6a-b). Wag31$_{\Delta1-60}$ exhibited higher lipid levels than *Δwag31*-ATc (Fig. 6a-b). These results highlight that the maintenance of lipid homeostasis by Wag31 is a consequence of its tethering activity.

Next, we examined the dependence of CL localisation on tethering by staining *Msm* +ATc, *Δwag31* -ATc, *Δwag31*+ATc, *Δwag31::wag31*+ATc, *Δwag31::wag31$_{K20A}$*+ATc, and *Δwag31:: wag31$_{\Delta1-60}$* +ATc with NAO. NAO staining showed a polar and septal (in dividing cells) localization pattern in the case of *Msm* and *Δwag31*-ATc, whereas it stained the cytosol and perimeter of the Wag31 depleted cells (Fig. 6c-d). Complementation with Wag31 restored NAO localization completely, whereas Wag31$_{K20A}$ restored it partially, which could be attributed to its limited tethering activity (Fig. 6c-d). Importantly, we observed a mis-localized pattern of NAO with Wag31$_{\Delta1-60}$ similar to Wag31 depletion (Fig. 6c-d). These results inform of the dependence of CL localization on membrane tethering activity

of Wag31. Taken together, the data suggests that Wag31 regulates lipid homeostasis and localisation
by the virtue of its tethering activity.

The N and C-terminal domains of Wag31 have distinct functions.

From the results presented above, we established a role for the N-terminal region of Wag31
that comprises DivIVA-domain in membrane-tethering (Fig. 4). We also identified the protein-interaction
network of Wag31 with the help of MS/MS analysis (Fig. 3 & Fig. S4). This implored us to investigate
the functionality of the C-terminal region of Wag31. We hypothesized that it might be responsible for
mediating interactions of Wag31 with other proteins. We performed a pull-down assay utilizing *E. coli*
lysates expressing either Wag31 or Wag31$_{\Delta1-60}$ (Fig. 7a). We chose novel interactors of Wag31 viz.
MurG, SepIVA, and Msm2092 from our interactome database (Fig.3a) and a known interactor of
Wag31, AccA3. The *Msm* lysates expressing 3x-FLAG tagged versions of the above-mentioned
proteins (Fig.7b) were incubated with *E. coli* lysate or *E. coli* lysates expressing either Wag31 or
Wag31$_{\Delta1-60,}$ followed by pulldown with Cobalt beads. The western blot analysis revealed that all the
tested interactors bound to Wag31 (Fig. 7c). Interestingly, interactions with all the proteins were retained
in the case of Wag31$_{\Delta1-60}$ as well, thus indicating that Wag31 interacts with other proteins, largely via
its C-terminal. Taken together, the results suggest that the N-terminal region of Wag31 is involved in
membrane-tethering through interactions with CL, and the C-terminal region is involved in modulating
interactions with other membrane-associated proteins. Thus N-terminal and C-terminal of Wag31 have
distinct molecular functions that together are crucial for maintaining cell shape, lipid homeostasis and
survival of mycobacteria.

300
301

302   ***Discussion.***

303   Wag31 modulates mycobacterial cell division by acting as an adaptor protein that recruits the

elongation machinery to the old pole for unipolar growth of cells (10, 15, 28). In the absence of Wag31,

rod-shaped cells advance into a spheroplast-like stage, which is thought to arise from dysregulated PG

metabolism (10, 11, 28). Recently, Wag31 has been implicated in maintaining the lipid-rich IMD

compartment (37). But the molecular mechanisms by which Wag31 manages these functions has

remained unclear. In this study, we discover an additional function of Wag31 as a membrane tether,

which likely reconciles many of the previously ascribed functions of this protein into a unified model. To

gain insights into defects associated with a change in cell shape, we imaged Wag31-deficient cells and

found a dramatic accumulation of lipids in the form of ILIs. Interestingly, both *Msm* and *ΔWag31*-ATc

showed the presence of ILIs, but in much less abundance, indicating that ILIs are a normal occurrence

that may function to regulate cellular functions (Fig. 1g-h).

314   The loss of Wag31 leads to the accumulation of neutral lipids while its overexpression increase

the levels of alpha-mycolic acid (α-MA), implying that the maintenance of native Wag31 levels is critical

for the proper balance of lipid composition (Fig. 2f-h). A protein's interactome is akin to its functional

fingerprint. Here, we identify 116 novel interacting partners of Wag31 through mass spectrometry and

validate three of these interactions, namely with MurG, SepIVA and Msm2092, through pull-down

studies (Fig. 3 & 5i). Strikingly, the bulk of Wag31 interactors are membrane proteins, which might be

attributed to the formation of Wag31 oligomers on the membrane. Previous studies have reported that

the depletion of Wag31 delocalize MurG, but if this was caused by loss of direct interaction or was a

consequence of change in cell shape was not known (34, 37), which our results clarify. Furthermore,

our results identify SepIVA, which regulates MurG (47), the D-aminopeptidase Msm2092 and AmiB,

both of which are PG remodellers, as Wag31 interactors. Together, these results contribute towards

our understanding of how Wag31 affects PG synthesis (Fig 3a,d). AccA3 is an interactor of Wag31 (28)

(34) and catalyses the first committed step in the synthesis of long-chain fatty acids by forming malonyl-

CoA. In addition, Long-chain specific acyl dehydrogenase (LCAD), which we report here to be another

interactor of Wag31, catalyses the first step in the β-oxidation of fatty acids, which might explain the

accumulation of long chain PEs and long-chain fatty acids.

330   As mentioned above, the overexpression of Wag31 leads to an upregulation in α-MA levels

(Fig. 2h). The localization of the α-MA transporter MmpL3 depends on polar PG levels (56, 57). MurG

is an essential enzyme for PG synthesis (58) and we report here that it strongly interacts with Wag31

(Fig. 3a,d). It is tempting to hypothesize that homeostatic control on lipid composition arises from the

ability of Wag31 to modulate the activity of enzymes involved in lipid metabolism and PG synthesis.

That both depletion and overexpression of Wag31 affects lipid composition is likely also a manifestation

of its ability to function as a homotypic tether, where the tethering ability is inherently sensitive to the

tether concentration.

338   Our results are the first to establish a membrane tethering function of Wag31. DivIVA proteins

possess an N-terminal membrane-binding domain and a C-terminal coiled-coil domain (50, 51). Our

data show that disruption of the N-terminal domain (Wag31$_{\Delta 1-60}$) markedly reduces membrane binding

and completely abolishes membrane tethering (Fig. 4f, g). The N-terminal domain binds CL while the

C-terminal coiled-coil domain facilitates multimerization, which together could explain the molecular
basis for Wag31 to function in homotypic tethering of CL-containing membranes. Results from testing
mutants defective in membrane tethering indicate that this function is important for the proper
localisation of CL and eventually the survival of mycobacteria (Fig. 5f & 6c-d). Thus, alterations in the
native levels of Wag31 results in a displacement of CL from the poles (Fig. 5a-d). However, given
NAO's ability to bind to anionic phospholipids, the observed staining could also typify a
redistribution of anionic phospholipids in the cytosol and membrane. In the absence of Wag31
binding to the poles (due to gene silencing or deletion of the DivIVA-domain), the multiprotein
scaffold required for cell envelope biosynthesis and cell elongation is not assembled properly,
leading to changes in lipid and PG metabolism and consequently to cells failing to form poles
where CL would normally concentrate. The failure of a Wag31-centered multiprotein scaffold
formation likely cause of this phenotype is further supported by the wag31 overexpression study,
which shows similar changes in NAO staining. By affecting the stoichiometry of Wag31 relative to
other interacting protein partners, Wag31 overexpression negatively impacts the formation and
activity of the multiprotein scaffold.

While, complementation with Wag31 reverses the phenotypic changes observed upon Wag31
depletion, complementation with the membrane tethering mutant (Wag31$_{\Delta 1-60}$) fails to do so (Fig. 5f-g
& 6a-d). These results not only highlight the dependence of CL localization on membrane-tethering but
also underscore the importance of tethering for maintaining cellular morphology, physiology and
survival. CL micro-domains at the poles (53, 54) have been shown to be an important hub for protein
localisation and activity. Several studies have implicated CL in modulating the activities of membrane-
resident proteins involved in electron-transport, protein translocation, bacterial two-component system
etc. (59-62).

Our work delineates the roles of the N- and C-terminal domains of Wag31. The N-terminal
DivIVA-domain facilitates lipid binding and membrane tethering, and the C-terminal coiled-coil domain
engages in protein-protein interactions and facilitates the multimerization of Wag31. This is evident from
findings that deletion of the N-terminal DivIVA-domain affects lipid binding and tethering but not its
ability to bind interacting partners (Fig. 4c-d, f-g & 7c). Based on our results, we propose a model in
which the N-terminal of Wag31 binds and tethers CL ensuring its proper localization while the C-terminal
is free to interact and recruit proteins to the membrane to facilitate cell growth and division (Fig. 8). We
reason that the loss of cell-shape and viability in the absence of tethering occurs due to mis-localization
of CL which in turn has a negative impact on membrane-localized processes at a global scale.
374
375

Materials and Methods

All the illustrations were made using BioRender. List of constructs and strains used in the study are given in Table S1. Mass spectrometry samples were processed and performed by Valerian Chem.

Bacterial strains and culturing

Middlebrook 7H9 medium (BD Biosciences) or 7H11 agar (BD Biosciences) was used to cultivate mycobacterial strains. 7H9 medium was supplemented with 10% ADC (albumin, dextrose, NaCl, and catalase) and 0.2% glycerol (Sigma) and, 0.05% Tween-80 (Sigma) or 7H11 agar (BD Biosciences) with 10% OADC (oleic acid added to ADC) and 0.2% glycerol were added to the media.

Generation of *wag31* conditional mutant and *in vitro* Growth kinetics

Recombineering method, as described in (63), was used to generate a Wag31 conditional gene replacement mutant. Detailed methodology is provided in the supplementary text. To perform *in vitro* growth kinetics, secondary cultures of *Msm* and *Δwag31* were seeded at a density of A_{600}~ 0.05 and either left untreated or treated with 100 ng/mL ATc for 12 h. At 0 and 12 h, appropriate serial dilutions were plated on 7H11 agar plate and incubated at 37°C for 2-3 days, and CFUs were enumerated. CFUs were plotted using GraphPad Prism 9 and analysed by performing Two-way ANOVA (0.0021 (**), 0.0002 (***).

Western blot analysis

Msm, Msm::wag31 and *Δwag31* transformants grown in the presence or absence of IVN/ ATc for 12 h were resuspended in Phosphate Buffer Saline containing 5% glycerol and 1mM protease inhibitor-phenymethylsulfonylfluoride, PMSF (Sigma) and whole cell lysates (WCL) were prepared with the help of bead-beating. Concentration of protein in each sample was estimated by Pierce BCA protein Assay kit (Thermo Fischer Scientific) and 40 μg WCLs were resolved on 10-12% SDS-PAGE gels, transferred onto nitrocellulose membrane, probed with desired antibodies and developed using ECL reagent (Millipore).

Sample preparation for SEM and TEM

Msm and *Δwag31* were incubated in the absence of the presence of ATc for 12 h, and the samples were prepared for both SEM and TEM as described previously in (64). For SEM, the samples were imaged at 20,000X using FEI Nova NanoSEM 450. TEM sections were examined on a 120 kV transmission electron microscope (L120C, Talos, Thermo Scientific) at 17,500 magnification.

BODIPY labelling, imaging and analysis

1 mL cells of *Msm* and *Δwag31* either untreated or treated with 100 ng/mL ATc for 12 h were stained with 20 µg/mL BODIPY (Invitrogen). For examining lipids in Wag31 overexpression strain, *Msm* and *Msm::wag31* were treated with 5 μM IVN for 12 h and 1 mL cells of both strains were stained with 2.5 μg BODIPY. Staining was performed for 30 min protected from light in an incubator shaker followed by washing with sterilized Phosphate buffer saline (PBS).

Sample preparation for Lipidomics and analysis

Secondary cultures of *Msm* and *Δwag31* were seeded at a density of A_{600}~ 0.05 and either left untreated or treated with 100 ng/mL ATc for 12 h until A_{600}~ 0.8-1.0. Post 12 h incubation with ATc, cells corresponding to A_{600}~20 were processed for lipid extraction as described (64). Detailed methodology is provided in supplementary text.

Immunoprecipitation (IP) and MS/MS analysis

250 mL secondary cultures of *Msm::gfp* and *Δwag31* were inoculated at an A_{600}~ 0.05 and incubated in a shaker incubator until A_{600}~ 0.8-1.0. Cells were collected and lysed using bead-beating to make Whole cell lysate (WCL) and passed through a 0.2 µM filter. 3 mg lysate from each group was immunoprecipitated overnight with anti-FLAG® M2 magnetic beads (Millipore), washed thrice, and heated in 4X Laemmli Buffer. One-fifth of the IP supernatant was resolved on a 10% gel, transferred onto a Nitrocellulose membrane, and probed with anti-FLAG Antibody (Sigma). Lysates from confirmed IPs (biological quadruplets) were taken further for mass spectrometry sample preparation and analysis as described in supplementary text.

Proximity-Based Labelling of Membrane-Associated Proteins (PLiMAP)

For carrying out the PLiMAP assay, Wag31 and its mutants were purified and desired liposomes were prepared as described in supplementary. PLiMAP was carried out and analysed as described (65). Briefly, 2 µM Wag31 or its mutants were incubated with liposomes containing 30 mol% CL, DOPG, or DOPE (200 µM total lipid) in a final volume of 30µL. The reaction was incubated for 30 min in the dark at room temperature followed by exposure to a 1 min long pulse of 365 nm UV light (UVP crosslinker CL-1000L) at an intensity of 200mJ cm^{-2}. The reaction was mixed with Laemmli sample buffer, boiled, and equal volumes for all reactions were loaded onto a 10% polyacrylamide gel and resolved using SDS-PAGE. iBright1500 (Thermo Fischer Scientific) or Amersham Typhoon (GE) were used for imaging gels. Gels were first imaged for BODIPY fluorescence and later fixed and stained with Coomassie Brilliant Blue.

Tethering assays, fluorescence imaging, and image analysis

1 µM of Wag31 and its mutants were mixed with 10 µM of 100 nm extruded liposomes in Eppendorf. The mixture was then transferred to a LabTek chamber and imaged using 100x 1.4 NA oil-immersion objective on an Olympus IX73 inverted microscope connected to an LED light source (CoolLED) and an Evolve 512 EMCCD camera (Photometrics). Image acquisition was controlled by µManager, and images were analysed using Fiji. For single liposome intensity analysis, all images from the same condition from two independent experiments were combined into one montage. The montage was duplicated and thresholded using Fiji's Otsu algorithm to generate a binary mask. The mask was overlaid onto the original montage to obtain the maximum intensity of liposomes.

NAO labelling, imaging and analysis

Msm, *Msm::wag31* were treated with 5 µM IVN for 12 h and *Msm* and *Δwag31* were either left untreated or treated with 100 ng/mL ATc for 12 h. For complementation experiment, 1 mL of each strain was stained with 2 µM NAO for 1 h protected from light in an incubator shaker and washed thrice with sterilised 1X PBS. 2 µl of cells were spotted on a 1% agar pad (prepared in 7H9) and visualized using a 100X/1.32 NA oil immersion objective on a Leica THUNDER Imaging systems at 475nm:535nm (Excitation: Emission). About 20 Z-stack images, each spaced 0.1 µM apart, were acquired using a scientific CMOS K8 camera (Leica microsystems). For CL specific labelling, collection was also performed at 642nm.

Complementation experiment

Secondary cultures of *Msm, Δwag31, Δwag31::wag31, Δwag31::wag31$_{K20A}$,* and *Δwag31::wag31$_{Δ1-60}$*

were set up at A$_{600}$ ~0.05. For each strain, two types of cultures were set up in triplicates- untreated and

treated with 100ng/mL ATc and grown for 12 h. At 0 and 12 h, appropriate serial dilutions were plated

on 7H11 agar plates and incubated at 37°C for 2-3 days. CFUs were enumerated and plotted using

GraphPad Prism 9. Statistical analysis was performed using two-way RM ANOVA followed by *Tukey's*

multiple comparison test, α=0.01, GP: 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), <0.0001 (****).

For BODIPY- and NAO- labelling, the strains were grown as described above. At 12 h, 1 mL cultures

of *Msm*+ATc, *Δwag31, Δwag31* +ATc, *Δwag31::wag31*+ATc *, Δwag31::wag31$_{K20A}$* +ATc*,* and

Δwag31::wag31$_{Δ1-60}$ +ATc were taken and labelled with either BODIPY or NAO. The labelling, imaging

and analysis for both fluorophores was performed as described in the respective sections above.

His pulldown assay.

E.coli and *Msm* lysate preparation for the assay is described in supplementary. Briefly, 500 μg of *E. coli*

lysates expressing Wag31 or mutant were mixed with 300 μg *Msm* lysates expressing Wag31

interactors in a total reaction volume of 700 μL and the samples were twirled at 4°C for 2 h. 50 μL

cobalt beads/ sample (GoldBio) were washed twice with lysis buffer and incubated with 5% BSA solution

for 2 h at 4°C. Subsequently, beads were washed with lysis buffer and incubated with samples for 2 h

(twirled at 4°C). The pull downs were washed thrice in lysis buffer containing 1% Triton X-100 and

finally resuspended in 2X-SDS sample buffer. Pulldown samples were resolved on 10-12% SDS-PAGE

and probed with α-His and α-FLAG antibodies.

Figure Legends:

Fig.1| Loss of Wag31 leads to the formation of Intracellular Lipid Inclusions (ILIs).

(a.) Illustration depicting the known role of Wag31 in directing polar elongation and maintaining rod-

shaped morphology of cells. **(b.)** Whole Cell Lysates **(**WCL) were prepared from *Msm* and *Δwag31*

(Table S1) either untreated or treated with 100 ng/mL ATc for 12 h, and 40 μg lysate was resolved on

a 12% gel, transferred on Nitrocellulose membrane, and probed with either α-Wag31 or α-rpoA antibody

(loading control), respectively. **(c.)** Cultures showing growth of *Msm* and *Δwag31* either untreated or

treated with 100 ng/mL ATc for 12 h. **(d.)** *Msm*+ATc, *Δwag31*-ATc, and *Δwag31*+ATc were tracked for

bacillary survival at 0 and 12 h by enumerating CFUs. Appropriate serial dilutions were plated on 7H11

plates (without antibiotic or ATc), and bar graphs demonstrating bacillary survival (CFU log$_{10}$/mL ±

standard deviations (SD, represented by error bars) were plotted at indicated time points. Statistical

significance was calculated using Two-way ANOVA (0.0021 (**), 0.0002 (***). The experiment was

performed twice independently, each performed in triplicates. **(e.)** SEM micrographs showing *Msm*,

Δwag31-ATc, and *Δwag31*+ATc at 12 h post-treatment. **(f.)** Cells depicting rod, bulged or round

morphology. **(g.)** 200 cells from each group were checked for either rod, bulged or round morphology,

and the percentage of each morphology category was plotted in a bar graph. **(h.)** TEM micrographs

showing *Msm*, *Δwag31*-ATc, and *Δwag31*+ATc at 12 h post-treatment. **(i.)** 250 cells, each from *Msm*,

Δwag31-ATc and 185 cells from *Δwag31*+ATc, were divided into three classes (0-2, 2-5, and >5 ILIs)

based on the number of ILIs and the percentage of each class in every strain was plotted in a bar graph.

Fig.2| Wag31 levels perturb lipid homeostasis

Representative micrographs showing BODIPY staining in **(a.)** *Msm*+ATc, *Δwag31*-ATc and,
Δwag31+ATc samples imaged 12 h post-ATc treatment **(b.)** *Msm*+ IVN, *Msm*::*wag31* + IVN samples
imaged 12 h post 5 μM IVN treatment. **(c.)** Corrected total cell fluorescence was calculated in individual
bacterial cells from *Msm*+ATc, *Δwag31*-ATc, and *Δwag31*+ATc strains (N=48) and plotted in a graph.
Horizontal bar represents median CTCF. (Statistical analysis was performed using one-way ANOVA
followed by Brown-Forsythe test (Tukey's multiple comparison test; 0.0021 (**), <0.0001 (****)). **(d.)**
Corrected total cell fluorescence was calculated for 42 individual cells from *Msm*+IVN, and
Msm::*wag31*+IVN strains and plotted in a graph (Statistical analysis was performed using Welch's t-
test 0.0021 (**), **e.)** Schematic depicting the workflow of Lipidomics (LC-MS). **(f-g.)** Heatmap depicting
$\log_2$ fold change (FC) in indicated lipid classes of *Msm* +ATc, *Δwag31*-ATc, and *Δwag31*+*ATc* as
compared to *Msm*-ATc. The experiment was performed with six biological replicates, and the
abundances of identified lipid classes were normalized to *Msm*-ATc to yield relative abundances of
lipids in other strains. Blue and red colours indicate upregulated and downregulated lipid classes; color
intensity variation changes with $\log_2$ FC. **(h.)** Bar graph (Mean ± SEM) depicting the abundance of α-
MA in *Msm* and *Msm::wag31* treated with 5 μM IVN for 12 h. Values on the Y-axis represent area under
the curve (AUC) normalised to an internal standard (FFA 15:0). X-axis represent various species (chain
length variations) of α-MA. (Statistical analysis was performed using two-way ANOVA followed by a
Tukey's multiple comparison's test; <0.0001 (****)).

Fig.3| Molecular interactions of Wag31 with membrane-associated proteins govern cellular
homeostasis.

(a.) Illustration showing the distribution of the top 20 interactors of Wag31 identified from MS/MS
analysis from three independent immunoprecipitations. Interactors were classified on the basis of their
functional category based on data available in Mycobrowser (66). MSMEG numbers or the identity of
interactors are marked in smaller circles. **(b.)** Coomassie Brilliant blue (CBB) staining of *E.coli* lysates
expressing either His-Wag31. *E. coli* lysate prepared from BL21-DE3 strain was used as the control.
(c.) Representative western blots demonstrating expression of 3XFLAG tagged-MurG, -SepIVA, -
Msm2092 and, -AccA3. WCLs were prepared from 0.5 μM IVN induced *Msm::murG*, *Msm::sepIVA*,
Msm::msm2092 and 40 μg of each was resolved on a 10 % SDS-PAGE, transferred onto a
nitrocellulose membrane and probed with α-FLAG and α-RpoA (loading control). **(d.)** Representative
western blots showing interaction of MurG, SepIVA, and, Msm2092 with Wag31. *Msm* and *E. coli* lysates
represented in (b & c) incubated together and pulled down with Cobalt beads were resolved on a 10-
12 % SDS-PAGE, transferred onto nitrocellulose membrane and probed with α-His to detect the
pulldown and α-FLAG to detect the interaction. The experiment was performed independently twice.
(e.) Graphic representation (adapted from (48)) shows the distribution of the top 20 interactors in
different compartments of a mycobacterial cell: membrane, cytoplasm, or core (involved in DNA
replication and recombination). Interactors from each compartment are marked inside the respective
domain. **(f.)** Pie-chart shows the distribution of interactors belonging to the IMD, PM-CW, or membrane
proteins whose specific localization is not determined. Percentage of proteins (top 25 hits) belonging to
each membrane-compartment is mentioned.

Fig.4| Wag31 binds and tethers mycobacterial membrane.

(a.) Schematic showing the workflow of the PLiMAP assay. **(b.)** Coomassie Brilliant blue (CBB) staining of His-Wag31 and in-gel fluorescence (FI) from a representative PLiMAP experiment performed using 2 µM Wag31 and hundred-fold excess of 30 mol% of either 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE-), and1′,3′-bis[1,2-dioleoyl-sn-glycero-3-phospho]-glycerol (DOPG-), or CL-containing liposomes. **(c.)** CBB and FL gels from a representative PLiMAP experiment performed with 2 µM Wag31, $Wag31_{K20A,}$ or $Wag31_{\Delta1-60}$ either in the presence or absence of 30 mol% CL liposomes. Note the presence of SDS-resistant higher-order polymers in gel indicated by the red curly bracket. **(d.)** Quantitation of PLiMAP data from (c.) showing a difference in binding propensities of His-Wag31, $His-Wag31_{K20A}$ and $His-Wag31_{\Delta1-60}$. Data represent the mean ± SD of three independent experiments. Statistical analysis was performed using a 2-tailed unpaired student's t test in Graph Pad Prism 9. Wag31 vs $Wag31_{K20A}$ showed no significant difference (ns, p value = 0.5667) whereas Wag31 vs $Wag31_{\Delta1-60}$(p value = 0.0081, **) and $Wag31_{K20A}$ vs $Wag31_{\Delta1-60}$ (p value = 0.0002, ***) showed significant difference in binding **(e.)** Representative micrographs from binding and tethering assay showing CL-DiI-liposomes (~100 nm) before and after incubation with 1 µM His-Wag31-GFP. Membrane and protein fluorescence are rendered in red and green, respectively. The experiment was performed independently twice. **(f.)** Representative micrographs from tethering experiments performed with His-Wag31, $His-Wag31_{K20A,}$ or $His-Wag31_{\Delta1-60}$ proteins. The left panels indicate PE liposomes only (upper panel) and PE liposomes incubated with 1 µM Wag31 (lower panel), the middle panels indicate CL liposomes (upper) or CL liposomes incubated with 1 µM Wag31 (lower panel), the right panel indicates CL liposomes either incubated with $His-Wag31_{K20A}$ (upper) and $His-Wag31_{\Delta1-60}$ (lower). For visualization, PE- and CL-containing liposomes were doped with trace amounts of 3,3'-dilinoleyloxacarbocyanine perchlorate (FAST DiO™) and 4-chlorobenzenesulfonate (FAST DiI™), respectively**. (g.)** Data from particle analysis showing the maximum intensity of liposomes for all the reactions from two independent experiments.

Fig.5| Tethering is crucial for the survival of mycobacteria

(a.) Top panel: Representative Fluorescent micrographs showing NAO intensity across *Msm*+ATc, *Δwag31*-ATc and, *Δwag31*+ATc samples imaged at 12 h post-ATc treatment. Bright red foci of polar CL are indicated with yellow arrowheads and septal CL foci with white arrowheads. Bottom panel: shows respective phase images. **(b.)** Line plot depicting CL distribution along the length of the cells across *Msm*+ATc, *Δwag31*-ATc and, *Δwag31*+ATc. N=100 cells across two independent biological replicates were analysed using Fiji and their normalised fluorescence intensities were plotted against cell length; fluorescence emitted from 0-20 % & 80-100 % of cell length was considered polar. **(c.)** Representative fluorescent micrographs showing NAO staining in *Msm*+IVN, *Msm*::*wag31*+IVN samples imaged at 12 h post 5 µM IVN treatment. Bright red foci of polar CL are indicated with yellow arrowheads, and septal CL foci with white arrowheads. The panels below show corresponding phase images. **(d.)** Line plot depicting CL distribution along the length of the cells in *Msm*+IVN, *Msm*::*wag31*+IVN. N=100 cells across two independent biological replicates were analysed using Fiji. The y-axis represents the normalized fluorescence intensities, and the x-axis represents cell length expressed as a percentage of total cell length. Fluorescence peaks in 0-20 % & 80-100 % of cell length

were considered polar. **(e.)** Structure of Wag31 (MSMEG_4217) from AlphaFold protein structure

database (67). The structure represents the N-terminal, linker, and C-terminal domains in Wag31. The

N-terminal contains a short helix, followed by an intertwined loop harbouring positively charged amino

acid K20 (highlighted in red) and the DivIVA domain. The N-terminus is linked to the C-terminus via a

linker. **(f.)** *Msm+ATc, Δwag31, Δwag31::wag31, Δwag31::wag31$_{K20A}$, Δwag31::wag31$_{Δ1-60}$* either

untreated or treated with ATc were scored for bacillary survival post 12 h ATc addition by enumerating

CFUs. Appropriate serial dilutions were plated on 7H11 plates (without antibiotic or ATc), and bar

graphs demonstrating bacillary survival (CFU log$_{10}$/mL ± standard deviations (SD, represented by error

bars) were plotted and statistical significance was calculated using two-way RM ANOVA followed by

Tukey's multiple comparison test, α=0.01, GP: 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), <0.0001

(****). The experiment was performed twice independently, each with triplicates. **(g.)** Strains described

in **(f.)** were scored for their morphology-rod, bulged or round at 12 h post ATc treatment. 308 cells of

each strain across two independent experiments were analysed.

Fig.6| Tethering is correlated with lipid levels and CL localization in mycobacteria

(a.) Top panel: Representative fluorescent micrographs showing BODIPY staining across *Δwag31-ATc*

and Msm, Δwag31, Δwag31::wag31, Δwag31::wag31$_{K20A}$, Δwag31::wag31$_{Δ1-60}$ imaged at 12 h post ATc

addition. Bottom panel: corresponding phase images. **(b.)** Corrected total cell fluorescence (CTCF) of

BODIPY-stained cells was calculated from N=100 cells across two independent experiments and

analysed using Fiji and plotted in a graph. The horizontal bar represents the median CTCF. (Statistical

analysis was performed using one-way ANOVA followed by Brown-Forsythe test (Tukey's multiple

comparison test; 0.1234 (ns), 0.0332 (*), <0.0001 (****)). **(c.)** Fluorescent micrographs showing NAO

intensity of strains as mentioned above. Bottom panel: corresponding phase images. **(d.)** Line plot

analysis of strains described in **(c.)** depicting CL localisation. N=100 cells across two independent

experiments were analysed using Fiji and their normalised fluorescence intensities were plotted against

cell length.

Fig.7| The N and C-terminal domains of Wag31 have distinct functions.

(a.) Coomassie Brilliant blue (CBB) staining of *E.coli* lysates expressing either His-Wag31. *E. coli* lysate

prepared from BL21-DE3 strain was used as the control. **(b.)** Representative western blots

demonstrating expression of 3XFLAG tagged-MurG, -SepIVA, -Msm2092 and, -AccA3. WCLs were

prepared from 0.5 μM IVN induced *Msm::murG, Msm::sepIVA, Msm::msm2092, Msm::accA3* and 40

μg of each was resolved on a 10 % SDS-PAGE, transferred onto a nitrocellulose membrane and probed

with α-FLAG and α-RpoA (loading control). **(i.)** Representative western blots showing interaction of

MurG, SepIVA, Msm2092 and, Acca3 with Wag31 and Wag31$_{Δ1-60}$. *Msm* and *E. coli* lysates represented

in (g & h) incubated together and pulled down with Cobalt beads were resolved on a 10-12 % SDS-

PAGE, transferred onto nitrocellulose membrane and probed with α-His to detect the pulldown and α-

FLAG to detect the interaction. The experiment was performed independently twice.

Fig.8| Model for Wag31 mediated tethering of the membrane

(a.) Model of Wag31 showing the N- and C-terminals. N-terminal houses the DivIVA-domain that is

involved in membrane-binding and the C-terminal binds to various interactors of Wag31. **(b.)** Left:

Illustration showing Wag31 scaffolds bound to cardiolipin in the membrane via N-terminal, tethering it

and creating CL-microdomains at the poles. The C-terminal of Wag31 bound to protein-partners, localizing them to the membrane, facilitating several membrane-centric processes such as PG and lipid synthesis, that ensure mycobacterial survival. The right panel shows the time-dependent consequences of Wag31 depletion i.e. early and late phase. The cells are bulged at one pole and start accumulating ILIs in the early phase and as time progresses, the effect of Wag31 depletion becomes severe rendering the cells round and full of ILIs. In the absence of Wag31, CL-microdomains become delocalised due to the loss of tethering, pulling away several membrane-associated proteins causing non-polar growth and loss of curvature. **(b.)** A zoomed-in micrograph of the left panel of (a.) highlighting Wag31-mediated cellular activities that occur in IMD and PM-CW. The proteins shown in the front in dark solid colours are interactors of Wag31; light-coloured proteins in the background are other membrane proteins. *From left*: AccA3 & AccD5 (subunits of Acyl CoA complex) showing cell wall lipid-precursor synthesis; MurG is involved in Lipid II (precursor of PG) synthesis, SepIVA is a spatio-temporal regulator of MurG. Thioredoxin is an antioxidant critical for maintaining cellular redox balance, ICS (Iron-sulfur cluster) and SDH (Succinate Dehydrogenase), or respiratory complex II, are involved in bacterial respiration. GPD (NAD-dependent Glycerol 3-phosphate Dehydrogenase) engaged in *de novo* phospholipid biosynthesis. ABC transporters are involved in transmembrane transport (grey spheres represent solutes).

Figure S1| Generation of Wag31 knockdown

(a.) Schematic showing the *wag31* locus in *Msm* and *Δwag31* and illustrating the strategy utilized to replace *wag31* locus with hygromycin resistance cassette. Primers used for knockout confirmation are indicated as F1, R1, F2, R2, F3 and R3. **(b.)** 1% agarose gels showing confirmatory PCRs performed with different primer sets. The first PCR, or the control PCR, was done using SepIVA (738 bp) primers showing amplicon at ~750bp. It indicates an equal amount of gDNA isolated from *Msm* and *Δwag31* and subsequently used for all PCRs. The next panel shows amplicons from the PCR performed with the F1-R1 pair, which bind to *wag31* locus (~819 bp). They yield an 819 bp amplicon in the case of *Msm* and 819 bp and an additional ~1.5 kb amplicon in the case of *Δwag31.* The third and fourth panels show amplicons amplified using the F2-R2 set and F3-R3 set, which are expected only in the case of *Δwag31* confirming legitimate recombination at the native locus.

Fig.S2| Wag31 levels perturb lipid homeostasis

(a.) Whole Cell Lysates **(**WCL) were prepared from *Msm* and *Msm::wag31* (Table S1) treated with 5 μM IVN for 12 h, and 40 μg lysate was resolved on a 12% gel, transferred on Nitrocellulose membrane, and probed with either α-FLAG or α-rpoA antibody (loading control), respectively. **(b-g)** Bar graphs (Mean ± SEM) depicting the abundance of various lipids classes **(b.)** TAG, **(c.)** DAG, **(d.)** FFA, **(e.)** PE, **(f.)** PI, **(g.)** CL in *Msm* and *Msm::wag31* treated with 5 μM IVN for 12 h. Values on the Y-axis represent area under the curve (AUC) normalised to an internal standard. X-axis represent various species (chain length variations) of the above mentioned lipid classes. The experiment was performed with six biological replicates, and the abundances of identified lipid classes were normalized to *Msm*-IVN to yield relative abundances of lipids in *Msm::wag31* +IVN.

Figure S3| *Molecular interactions of wag31 with membrane proteins govern cellular homeostasis.*

(a.) Western blots showing FLAG-GFP (left panel) and FLAG-Wag31 (right panel) immunoprecipitations. 3 mg lysate from each group was processed for overnight FLAG immunoprecipitations, resolved on a 10% gel, and probed with α-Flag antibody. HC and LC are short for Heavy chain and light chain, respectively. **(b.)** Classification of cellular processes that occur in different cellular compartments of *Msm,* as shown in (48).

Figure S4| *Molecular interactions wag31 with membrane proteins govern cellular homeostasis.* The list of Wag31 interactors is classified according to functional categories based on the data available in Mycobrowser. The colour of the textbox indicates localization of the proteins: Pink- membrane, blue- cytoplasm, and core- green.

Table S1
Plasmids and strains used in the study.
Table S2. It depicts unique interacting partners found for Wag31.

Acknowledgments
We thank Dharani Kumar, SEST, University of Hyderabad for helping with imaging of SEM stubs; CCMB for access to its Genomics, Advanced Microscopy, and Electron Microscopy facilities; VIDO, USask for providing infrastructure to carry out some part of the work; Asis Kumar Khuntia for the generation of pET28b-*wag31*$_{K20A}$ and pET28b-*wag31*$_{1-60}$. YK was supported by a Research Associateship from NIH (Grant 6044-SC23-26). YK and ND acknowledge MITACS for the Globalink Research Award (FR118569) which supported the work carried out at VIDO. ND also acknowledges support from Canadian Institutes of Health Research (CIHR, funding reference number 185715). VIDO receives operational funding from the Government of Saskatchewan through Innovation Saskatchewan and the Ministry of Agriculture and from the Canada Foundation for Innovation through the Major Science Initiatives Fund. Research in this publication was supported by SERB grant (CRG/2018/001294), Govt. of India, and JC Bose Award (JCB/2019/000015), Govt. of India to VKN. Work in the SSK laboratory was supported by the Swarna Jayanti Fellowship to SSK by SERB, Govt. of India (Grant: SB/SJF/2021-22/01). AC was supported by a Senior research fellowship from CSIR, India. HK thanks the Indian Institute of Science Education and Research, Pune, for a graduate fellowship. TP thanks the Howard Hughes Medical Institute for funding support. We thank Dr. Christopher Sassetti for his critical comments.

References

1.   E. F. Bi, J. Lutkenhaus, FtsZ ring structure associated with division in Escherichia coli. *Nature* **354**, 161-
164 (1991).
2.   P. A. de Boer, R. E. Crossley, A. R. Hand, L. I. Rothfield, The MinD protein is a membrane ATPase
required for the correct placement of the Escherichia coli division site. *Embo j* **10**, 4371-4380 (1991).
3.   Z. Hu, A. Mukherjee, S. Pichoff, J. Lutkenhaus, The MinC component of the division site selection system
in Escherichia coli interacts with FtsZ to prevent polymerization. *Proc Natl Acad Sci U S A* **96**, 14819-
14824 (1999).
4.   D. M. Raskin, P. A. de Boer, The MinE ring: an FtsZ-independent cell structure required for selection of
the correct division site in E. coli. *Cell* **91**, 685-694 (1997).
5.   H. Meinhardt, P. A. de Boer, Pattern formation in Escherichia coli: a model for the pole-to-pole oscillations
of Min proteins and the localization of the division site. *Proc Natl Acad Sci U S A* **98**, 14202-14207 (2001).
6.   C. L. Woldringh *et al.*, Role of the nucleoid in the toporegulation of division. *Res Microbiol* **141**, 39-49
(1990).
7.   M. M. Logsdon, B. B. Aldridge, Stable Regulation of Cell Cycle Events in Mycobacteria: Insights From
Inherently Heterogeneous Bacterial Populations. *Front Microbiol* **9**, 514 (2018).
8.   J. H. Cha, G. C. Stewart, The divIVA minicell locus of Bacillus subtilis. *J Bacteriol* **179**, 1671-1683 (1997).
9.   D. H. Edwards, J. Errington, The Bacillus subtilis DivIVA protein targets to the division septum and controls
the site specificity of cell division. *Mol Microbiol* **24**, 905-915 (1997).
10.   C. M. Kang, S. Nyayapathy, J. Y. Lee, J. W. Suh, R. N. Husson, Wag31, a homologue of the cell division
protein DivIVA, regulates growth, morphology and polar cell wall synthesis in mycobacteria. *Microbiology*
(Reading) **154**, 725-735 (2008).
11.   C. Jani *et al.*, Regulation of polar peptidoglycan biosynthesis by Wag31 phosphorylation in mycobacteria.
BMC Microbiol **10**, 327 (2010).
12.   M. Letek *et al.*, DivIVA is required for polar growth in the MreB-lacking rod-shaped actinomycete
Corynebacterium glutamicum. *J Bacteriol* **190**, 3283-3292 (2008).
13.   N. R. Thanky, D. B. Young, B. D. Robertson, Unusual features of the cell cycle in mycobacteria: polar-
restricted growth and the snapping-model of cell division. *Tuberculosis (Edinb)* **87**, 231-236 (2007).
14.   K. J. Kieser, E. J. Rubin, How sisters grow apart: mycobacterial growth and division. *Nat Rev Microbiol*
12, 550-562 (2014).
15.   L. Nguyen *et al.*, Antigen 84, an effector of pleiomorphism in Mycobacterium smegmatis. *J Bacteriol* **189**,
7896-7910 (2007).
16.   L. R. Hammond, M. L. White, P. J. Eswara, ¡vIVA la DivIVA! *J Bacteriol* **201** (2019).
17.   H. Stahlberg *et al.*, Oligomeric structure of the Bacillus subtilis cell division protein DivIVA determined by
transmission electron microscopy. *Mol Microbiol* **52**, 1281-1290 (2004).
18.   K. N. Muchová *et al.*, Oligomerization of the Bacillus subtilis division protein DivIVA. *Microbiology*
(Reading) **148**, 807-813 (2002).
19.   M. Letek *et al.*, DivIVA uses an N-terminal conserved region and two coiled-coil domains to localize and
sustain the polar growth in Corynebacterium glutamicum. *FEMS Microbiol Lett* **297**, 110-116 (2009).
20.   M. D. Rigden *et al.*, Identification of the coiled-coil domains of Enterococcus faecalis DivIVA that mediate
oligomerization and their importance for biological function. *J Biochem* **144**, 63-76 (2008).
21.   K. Choukate *et al.*, Higher order assembling of the mycobacterial polar growth factor DivIVA/Wag31. *J*
Struct Biol **209**, 107429 (2020).
22.   K. Flärdh, Essential role of DivIVA in polar growth and morphogenesis in Streptomyces coelicolor A3(2).
Mol Microbiol **49**, 1523-1536 (2003).
23.   D. Fadda *et al.*, Streptococcus pneumoniae DivIVA: localization and interactions in a MinCD-free context.
J Bacteriol **189**, 1288-1298 (2007).
24.   K. G. Kaval, S. Halbedel, Architecturally the same, but playing a different game: the diverse species-
specific roles of DivIVA proteins. *Virulence* **3**, 406-407 (2012).
25.   H. B. Thomaides, M. Freeman, M. El Karoui, J. Errington, Division site selection protein DivIVA of Bacillus
subtilis has a second distinct function in chromosome segregation during sporulation. *Genes Dev* **15**,
1662-1673 (2001).
26.   K. G. Kaval, S. Hauf, J. Rismondo, B. Hahn, S. Halbedel, Genetic Dissection of DivIVA Functions in
Listeria monocytogenes. *J Bacteriol* **199** (2017).
27.   K. G. Kaval, J. Rismondo, S. Halbedel, A function of DivIVA in Listeria monocytogenes division site
selection. *Mol Microbiol* **94**, 637-654 (2014).
28.   X. Meniche *et al.*, Subpolar addition of new cell wall is directed by DivIVA in mycobacteria. *Proc Natl Acad*
Sci U S A **111**, E3243-3251 (2014).
29.   E. S. Melzer, C. E. Sein, J. J. Chambers, M. S. Siegrist, DivIVA concentrates mycobacterial cell envelope
assembly for initiation and stabilization of polar growth. *Cytoskeleton (Hoboken)* **75**, 498-507 (2018).
30.   W. X. Xu *et al.*, The Wag31 protein interacts with AccA3 and coordinates cell wall lipid permeability and
lipophilic drug resistance in Mycobacterium smegmatis. *Biochem Biophys Res Commun* **448**, 255-260
(2014).
31.   M. T. M. Hannebelle *et al.*, A biphasic growth model for cell pole elongation in mycobacteria. *Nat Commun*
11, 452 (2020).

32. I. Santi, N. Dhar, D. Bousbaine, Y. Wakamoto, J. D. McKinney, Single-cell dynamics of the chromosome
replication and cell division cycles in mycobacteria. *Nat Commun* **4**, 2470 (2013).
33. P. Mukherjee *et al.*, Novel role of Wag31 in protection of mycobacteria under oxidative stress. *Mol*
Microbiol **73**, 103-119 (2009).
34. N. Habibi Arejan *et al.*, Polar protein Wag31 both activates and inhibits cell wall metabolism at the poles
and septum. *Front Microbiol* **13**, 1085918 (2022).
35. Y. S. Morita *et al.*, Compartmentalization of lipid biosynthesis in mycobacteria. *J Biol Chem* **280**, 21645-
21652 (2005).
36. J. M. Hayashi *et al.*, Spatially distinct and metabolically active membrane domain in mycobacteria. *Proc*
Natl Acad Sci U S A **113**, 5400-5405 (2016).
37. A. García-Heredia *et al.*, Membrane-partitioned cell wall synthesis in mycobacteria. *Elife* **10** (2021).
38. V. Singh *et al.*, Identification of aminopyrimidine-sulfonamides as potent modulators of Wag31-mediated
cell elongation in mycobacteria. *Mol Microbiol* **103**, 13-25 (2017).
39. J. Daniel, H. Maamar, C. Deb, T. D. Sirakova, P. E. Kolattukudy, Mycobacterium tuberculosis uses host
triacylglycerol to accumulate lipid droplets and acquires a dormancy-like phenotype in lipid-loaded
macrophages. *PLoS Pathog* **7**, e1002093 (2011).
40. C. Barisch, P. Paschke, M. Hagedorn, M. Maniak, T. Soldati, Lipid droplet dynamics at early stages of
Mycobacterium marinum infection in Dictyostelium. *Cell Microbiol* **17**, 1332-1349 (2015).
41. R. Bartz *et al.*, Lipidomics reveals that adiposomes store ether lipids and mediate phospholipid traffic. *J*
Lipid Res **48**, 837-847 (2007).
42. K. Tauchi-Sato, S. Ozeki, T. Houjou, R. Taguchi, T. Fujimoto, The surface of lipid droplets is a
phospholipid monolayer with a unique Fatty Acid composition. *J Biol Chem* **277**, 44507-44512 (2002).
43. A. Griego, T. Douché, Q. G. Gianetto, M. Matondo, G. Manina, RNase E and HupB dynamics foster
mycobacterial cell homeostasis and fitness. *iScience* **25**, 104233 (2022).
44. R. Priyadarshini, M. A. de Pedro, K. D. Young, Role of peptidoglycan amidases in the development and
morphology of the division septum in Escherichia coli. *J Bacteriol* **189**, 5334-5347 (2007).
45. C. Heidrich *et al.*, Involvement of N-acetylmuramyl-L-alanine amidases in cell separation and antibiotic-
induced autolysis of Escherichia coli. *Mol Microbiol* **41**, 167-178 (2001).
46. D. Mengin-Lecreulx, L. Texier, M. Rousseau, J. van Heijenoort, The murG gene of Escherichia coli codes
for the UDP-N-acetylglucosamine: N-acetylmuramyl-(pentapeptide) pyrophosphoryl-undecaprenol N-
acetylglucosamine transferase involved in the membrane steps of peptidoglycan synthesis. *J Bacteriol*
173, 4625-4636 (1991).
47. A. H. Freeman *et al.*, Arginine methylation sites on SepIVA help balance elongation and septation in
Mycobacterium smegmatis. *Mol Microbiol* **119**, 208-223 (2023).
48. J. Zhu *et al.*, Spatiotemporal localization of proteins in mycobacteria. *Cell Rep* **37**, 110154 (2021).
49. Z. He, J. De Buck, Cell wall proteome analysis of Mycobacterium smegmatis strain MC2 155. *BMC*
Microbiol **10**, 121 (2010).
50. K. Choukate, B. Chaudhuri, Structural basis of self-assembly in the lipid-binding domain of mycobacterial
polar growth factor Wag31. *IUCrJ* **7**, 767-776 (2020).
51. R. Lenarcic *et al.*, Localisation of DivIVA by targeting to negatively curved membranes. *Embo j* **28**, 2272-
2282 (2009).
52. M. A. Oliva *et al.*, Features critical for membrane binding revealed by DivIVA crystal structure. *Embo j* **29**,
1988-2001 (2010).
53. L. D. Renner, D. B. Weibel, Cardiolipin microdomains localize to negatively curved regions of Escherichia
coli membranes. *Proc Natl Acad Sci U S A* **108**, 6264-6269 (2011).
54. P. M. Oliver *et al.*, Localization of anionic phospholipids in Escherichia coli cells. *J Bacteriol* **196**, 3386-
3398 (2014).
55. E. Mileykovskaya, W. Dowhan, Visualization of phospholipid domains in Escherichia coli by using the
cardiolipin-specific fluorescent dye 10-N-nonyl acridine orange. *J Bacteriol* **182**, 1172-1175 (2000).
56. C. Carel *et al.*, Mycobacterium tuberculosis proteins involved in mycolic acid synthesis and transport
localize dynamically to the old growing pole and septum. *PLoS One* **9**, e97148 (2014).
57. N. H. Arejan, D. R. Czapski, J. A. Buonomo, C. C. Boutte, MmpL3, Wag31 and PlrA are involved in
coordinating polar growth with peptidoglycan metabolism and nutrient availability. *bioRxiv*
10.1101/2024.04.29.591792 (2024).
58. G. P. Salmond, J. F. Lutkenhaus, W. D. Donachie, Identification of new genes in a cell envelope-cell
division gene cluster of Escherichia coli: cell envelope gene murG. *J Bacteriol* **144**, 438-440 (1980).
59. G. Carranza *et al.*, Cardiolipin plays an essential role in the formation of intracellular membranes in
Escherichia coli. *Biochim Biophys Acta Biomembr* **1859**, 1124-1132 (2017).
60. V. A. Gold *et al.*, The action of cardiolipin on the bacterial translocon. *Proc Natl Acad Sci U S A* **107**,
10044-10049 (2010).
61. W. S. Yeo, S. Dyzenhaus, V. J. Torres, S. R. Brinsmade, T. Bae, Regulation of Bacterial Two-Component
Systems by Cardiolipin. *Infect Immun* **91**, e0004623 (2023).
62. S. P. Ouellette *et al.*, Localized cardiolipin synthesis is required for the assembly of MreB during the
polarized cell division of Chlamydia trachomatis. *PLoS Pathog* **18**, e1010836 (2022).
63. J. C. van Kessel, G. F. Hatfull, Recombineering in Mycobacterium tuberculosis. *Nat Methods* **4**, 147-152
(2007).

64.   S. Kumar *et al.*, Mycobacterium tuberculosis Transcription Factor EmbR Regulates the Expression of Key
817   Virulence Factors That Aid in Ex Vivo and In Vivo Survival. *mBio* **13**, e0383621 (2022).
818   65.   G. P. Jose, S. Gopan, S. Bhattacharyya, T. J. Pucadyil, A facile, sensitive and quantitative membrane-
819   binding assay for proteins. *Traffic* **21**, 297-305 (2020).
820   66.   A. Kapopoulou, J. M. Lew, S. T. Cole, The MycoBrowser portal: a comprehensive and manually annotated
821   resource for mycobacterial genomes. *Tuberculosis (Edinb)* **91**, 8-13 (2011).
822   67.   K. Tunyasuvunakool *et al.*, Highly accurate protein structure prediction for the human proteome. *Nature*
823   **596**, 590-596 (2021).
824

Figure 1

a. New Pole | Septum | Old Pole | ACC complex | New cell wall deposition | Depletion of Wag31 | Septum
● cell wall precursors | Wag31 | Septum

b. *Msm* - ATc | *Msm* + ATc | Δ*wag31* - ATc | Δ*wag31* + ATc
43- α-Wag31
54- 43- α-rpoA

c. *Msm* - ATc | *Msm* + ATc | Δ*wag31* - ATc | Δ*wag31* + ATc

d.
Msm + ATc
Δ*wag31* - ATc
Δ*wag31* + ATc
CFU log₁₀/ml
*** **
0 h   12 h

e. *Msm* | Δ*wag31* - ATc | Δ*wag31* + ATc
5 µm

f. Rod | Bulged | Round

g. Percentage of cells | Rod | Bulged | Round
Msm | Δ*wag31* - ATc | Δ*wag31* + ATc

i. Percentage of lipid bodies | Lipid bodies 0-2 | Lipid bodies 3-5 | Lipid bodies >5
Msm | Δ*wag31* - ATc | Δ*wag31* + ATc

h. *Msm* | Δ*wag31* - ATc | Δ*wag31* + ATc
500 nm

Figure 2

a.
Msm + ATc *MsmΔwag31* - ATc

MsmΔwag31 + ATc

b.
Msm + IVN *Msm::wag31* + IVN

2 μm

2 μm

c.
Corrected total cell fluorescence

****
ns ****

Msm + ATc *MsmΔwag31* - ATc *MsmΔwag31* + ATc

d.
CTCF (x 10⁴ au)

**

Msm + IVN *Msm::wag31* + IVN

h.
Normalized AUC

Msm + IVN
Msm::wag31 + IVN

**** **** ****

C74 C76 C78
α-MA

e.
Biological triplicates → Extraction → Cellular Lipids

f.
Msm+ATc
ΔWag31-ATc
ΔWag31+ATc

32:0 34:0 36:0 | 32:1 34:1 34:2 | 32:0 32:1 34:0 34:1 35:0 35:1 36:0 36:1 37:0 37:1 38:0 | 16:0 18:0 20:0 20:1 18:1 18:2 20:4 22:0 22:1 24:0
DAG | PE | FFA

-4 -2 0 2 +4

g.
Msm+ATc
ΔWag31-ATc
ΔWag31+ATc

66:1 66:2 66:3 67:1 67:2 68:0 68:1 68:2 68:3 68:4 69:0 69:1 69:2 70:0 70:1 70:2
CL

Figure 3

a.

His-Wag31/His-Wag31mut + Liposomes (+ 1% RFL) → RFL labelled liposome membrane → +UV → RFL cross-linked protein → PAGE → -UV +UV

b.

His-Wag31 (Coomassie)
Liposome: - PE PG CL
His-Wag31 (Fluorescence)
Liposome: - PE PG CL

c.

CL-liposomes / no liposomes
Wag31Δ1-60 / Wag31K20A / Wag31
Coomassie / Fluorescence / Coomassie / Fluorescence

d.

Fluorescence Intensity (x 10³)
Wag31, Wag31K20A, Wag31Δ1-60
** ***

e.

CL-Dil-Liposomes / CL-Dil-Liposomes / Wag31-GFP / Merged / 10μM

f.

PE Liposomes / CL Liposomes
His-Wag31K20A / His-Wag31 / His-Wag31 / His-Wag31Δ1-60
10μM

g.

Particle Intensity (x 10³)
PE Liposomes: - , Wag31
CL Liposomes: - , Wag31 , Wag31K20A , Wag31Δ1-60

Figure 5

Figure 6

Figure7

Figure 8

a.

N-terminal

DivIVA-domain

Linker

C-terminal domain

Lipid binding domain

Protein binding domain

b.

Polar elongation

Wag31 depletion

PM-CW

IMD

Early stage

Late stage

c.

Intracellular Membrane domain

Plasma membrane-Cell Wall

Cardiolipin

SDH

e^-

ICS

e^-

AccD5

ABC transporters

AccA3

SepIVA

MurG

Wag31 Scaffold

Wag31 Scaffold

GPD

Lipid I

Lipid II

Thioredoxin

de novo
PL synthesis

Transmembrane
transport

Cell wall precursors synthesis

Lipid II synthesis

Redox balance

Respiration

Cardiolipin

Phospholipids

MurG

SepIVA

LipidI

LipidII

AccD5

AccA3

Lipid
precursors

C-terminal

N-terminal

Wag31 tetramer

Solutes

Delocalised protein-lipid
/protein-protein aggregatres

Intracellular Lipid
Inclusions

Figure S1

a.

b.

Figure S3

a.

b.

Membrane:
- Energy production and conversion
- Cell cycle control, cell divison and chromosome partitioning
- Cell wall membrane and envelope biogenesis
- Intracellular trafficking, secretion, and vesicular transport

Core:
- Translation, ribosomal structure and biogenesis
- Transcription
- Replication, recombination and repair

Cytoplasm:
- Amino acid transport and metabolism

Figure S4

Cell wall & cell processes

- MSMEG_0402/mps2: linear gramicidin synthetase subunit D
- MSMEG_6099: conserved transmembrane protein
- MSMEG_4227: murG: undecaprenyldiphospho-muramoylpentapeptide beta-N-acetylglucosaminyltransferase
- MSMEG_2092: D-aminopeptidase
- MSMEG_6863: class II aldolase/adducin domain protein
- MSMEG_2416: SepIVA: DivIVA-domain containing protein
- MSMEG_1046: ABC-type molybdenum transport system
- MSMEG_0386: NAD dependent epimerase/dehydratase family protein
- MSMEG_3654: ATPase SecA2
- MSMEG_4958: diaminopimelate decarboxylase
- MSMEG_0919: heparin-binding hemagglutinin
- MSMEG_1954: ABC1 family protein
- MSMEG_2326: ABC-type molybdenum transport system

Lipid metabolism

- MSMEG_1813: propionyl-CoA carboxylase beta chain
- MSMEG_0372: oxidoreductase, short chain dehydrogenase/ reductase family protein
- MSMEG_2393: NAD-dependent glycerol-3-phosphate dehydrogenase
- MSMEG_2080: putative acyl-CoA dehydrogenase
- MSMEG_0912: 3-hydroxybutyryl-CoA dehydrogenase
- MSMEG_2648: Sfp-type phosphopantetheinyl transferase
- MSMEG_0406: fadE5: acyl-coA-dehydrogenase
- MSMEG_5199: putative acyl-CoA dehydrogenase
- MSMEG_4709: enoyl-CoA hydratase
- MSMEG_5197: long-chain specific acyl-CoA dehydrogenase
- MSMEG_6512: acyl-CoA dehydrogenase domain protein
- MSMEG_6511: acyl-CoA dehydrogenase domain protein
- MSMEG_6208: palmitoyl-CoA hydrolase
- MSMEG_5198: carnitinyl-CoA dehydratase

Replication, recombination & repair

- MSMEG_3596: ATPase- DNA replication
- MSMEG_2174: superfamily protein I DNA or RNA helicase
- MSMEG_1061: phosphohydrolase
- MSMEG_2768: OB-fold nucleic acid binding domain protein
- MSMEG_4450: alpha/beta hydrolase
- MSMEG_1930: DEAD/DEAH box helicase
- MSMEG_5156: base excision DNA repair protein, HhH-GPD family protein

Translation, ribosome structure & biogenesis

- MSMEG_4626: rne: ribonuclease, Rne/Rng family protein
- MSMEG_2931: thrS: threonyl-tRNA synthetase
- MSMEG_3003: aspS: aspartyl-tRNA synthetase
- MSMEG_4959: argS: arginyl-tRNA synthetase
- MSMEG_3777: pheT: phenylalanyl-tRNA synthetase, beta subunit

Regulatory proteins

- MSMEG_6219: moxR2: ATPase family protein associated with various cellular activities (AAA)
- MSMEG_4659: GntR-family protein transcriptional regulator
- MSMEG_3363: regulatory protein, TetR
- MSMEG_0035: fhaA: FHA domain protein

Unknown

- MSMEG_3785: PfkB-family protein carbohydrate kinase
- MSMEG_6783: integral membrane protein
- MSMEG_4652: D-oliose 4-ketoreductase
- MSMEG_1679: AmiB

Miscellaneous

- MSMEG_6514: trehalose synthase-fused probable maltokinase

Intermediary metabolism & respiration

- MSMEG_4915: alpha-glucan phosphorylase family protein
- MSMEG_0502: glucosidase
- MSMEG_6459: ferredoxin-dependent glutamate synthase 1
- MSMEG_3138: trx: thioredoxin
- MSMEG_0594: iron-sulfur cluster binding protein
- MSMEG_5427:prs: ribose-phosphate pyrophosphokinase
- MSMEG_2611: pyridine nucleotide-disulphide oxidoreductase family protein
- MSMEG_5937: bphl-2: 4-hydroxy-2-oxovalerate aldolase
- MSMEG_3051: gmk: guanylate kinase
- MSMEG_3044: pyrC: dihydroorotase
- MSMEG_1037: adhc1: alcohol dehydrogenase, zinc-containing
- MSMEG_4294: glnA: glutamine synthetase, type I
- MSMEG_3632: gnd: 6-phosphogluconate dehydrogenase, decarboxylating
- MSMEG_3100: opcA: OpcA protein: Putative OXPP cycle protein OpcA
- MSMEG_3894: prcA: proteasome alpha subunit
- MSMEG_2379: leuB: 3-isopropylmalate dehydrogenase
- MSMEG_6874: aldehyde dehydrogenase
- MSMEG_6114: ppa: inorganic pyrophosphatase
- MSMEG_1073: oxidoreductase, short-chain dehydrogenase/reductase family protein
- MSMEG_1825: rmlD: dTDP-4-dehydrorhamnose reductase
- MSMEG_3746: pyrG: CTP synthase
- MSMEG_4272: HesB/YadR/YfhF family protein
- MSMEG_3451: short-chain dehydrogenase/reductase, NAD(P)H-binding
- MSMEG_5592: Luciferase like monooxygenase
- MSMEG_0220; monoglyceride lipase
- MSMEG_3880: nitrilase/cyanide hydratase and apolipoprotein N-acyltransferase
- MSMEG_5903: oxidoreductase, short chain dehydrogenase/reductase family protein
- MSMEG_0759: purA: CTP synthase
- MSMEG_6598: ribH: 6,7-dimethyl-8-ribityllumazine synthase
- MSMEG_6351: histidinol-phosphate aminotransferase
- MSMEG_4528: cysH: phosphoadenosine phosphosulfate reductase
- MSMEG_4276: ilvE: branched-chain amino acid aminotransferase
- MSMEG_1670: sdhA: succinate dehydrogenase, flavoprotein subunit
- MSMEG_0539: transcriptional regulator, Crp/Fnr family protein
- MSMEG_4258: trpD: anthranilate phosphoribosyltransferase
- MSMEG_2357: cysteine desulfurase
- MSMEG_3207: hisB: imidazoleglycerol-phosphate dehydratase
- cobyrinic Acid a,c-diamide synthase
- alpha oxoglutarate ferredoxin oxidoreductase, beta subunit
- short chain dehydrogenase
- fructose-1,6-bisphosphatase, class II
- MSMEG_2617: cobB: cobyrinic Acid a,c-diamide synthase
- MSMEG_4645: orB: alpha oxoglutarate ferredoxin oxidoreductase, beta subunit
- MSMEG_3619: short chain dehydrogenase
- MSMEG_5239: glpX: fructose-1,6-bisphosphatase, class II
- MSMEG_3071: ribE: riboflavin synthase, alpha subunit
- MSMEG_0793: thiG: thiazole biosynthesis protein
- MSMEG_0956: hemB: delta-aminolevulinic acid dehydratase
- MSMEG_3219: indole-3-glycerol phosphate synthase
- MSMEG_0239: O-acetylhomoserine/O-acetylserine sulfhydrylase
- MSMEG_5515: purH: bifunctional purine biosynthesis protein
- MSMEG_6256: asd: aspartate-semialdehyde dehydrogenase

Amino acid transport and metabolism

- MSMEG_5211: aminotransferase class-III

Conserved hypotheticals

- MSMEG_2436; MSMEG_0897
- MSMEG_6941; MSMEG_3309
- MSMEG_2481; MSMEG_2434
- MSMEG_3148; MSMEG_6901
- MSMEG_6499; MSMEG_1340
- MSMEG_1514; MSMEG_5691
- MSMEG_3942; MSMEG_4282
- MSMEG_1049; MSMEG_3024
- MSMEG_3244; MSMEG_0240
- MSMEG_2685; MSMEG_4193
- MSMEG_2329

Membrane

Cytoplasm

Core

FigureS5

a.

Supplementary material

Materials and methods

Bacterial culturing

Antibiotic concentrations used: Hygromycin (Hyg) was used at 100 µg/mL, Kanamycin (Kan) at 25 µg/mL, and Apramycin (Apra) at 30 µg/mL for *Msm. E. coli* was cultured in Miller Luria Bertani Broth (Himedia) or Miller Luria Bertani Agar. Antibiotic concentrations used: Hyg was used at 150 µg/mL, Kan at 50 µg/mL, Apra at 30 µg/mL, Ampicillin (Amp) at 100 µg/mL for *E. coli.*

Generation of *wag31* conditional mutant

First, *wag31* was cloned under a tet regulatable promoter in pST-KiT$_{off}$ (kan^{res} ; (1)), which turns of transcription upon the addition of anhydrotetracycline. pST-wag31 construct thus generated was electroporated into a recombineering proficient strain, *Msm::pJV53*. An allelic exchange substrate was generated by cloning 675 bp from upstream and 700 bp from downstream of *wag31* (including ~150 bp at both ends) with *hyg^{res}* cassette and *oriE*+cosλ and digested with SnaBI to release the donor LHS-*hyg^{res}*-RHS. The linear donor was electroporated into *Msm::wag31* expressing gp60 and gp61 recombinases. The colonies were selected on Hyg and ten colonies were screened with multiple primer pairs to confirm recombination at the native locus. Confirmed recombinants were passaged multiple times to cure them of pJV53 to generate conditional mutant *Δwag31*.

BODIPY imaging and analysis

For *Msm* and *Δwag31*, ~10 µl of cells were spotted on a 1% agar pad (prepared in 7H9) and visualized at 4X zoom through Confocal microscopy (LSM880) at 488 nm/510 nm (Excitation: Emission). Background correction in the collected images followed by Corrected Total Cell Fluorescence analysis was performed using Fiji (2) and plotted using Graph Pad Prism 9. For examining lipids in BODIPY-stained *Msm* and *Msm::wag31,* 2 µl of cells were spotted on a 1% agar pad (prepared in 7H9) and visualized using a 100X/1.32 NA oil immersion objective on a Leica THUNDER Imaging systems at 475nm:519nm (Excitation: Emission). About 20 Z-stack images, each spaced 0.1 µM apart, were acquired using a scientific CMOS K8 camera (Leica microsystems). Background correction and Thunder image processing was performed on the images using the Las-X software module (Leica microsystems). Processed images were then analysed for CTCF using Fiji (2) and plotted using Graph Pad prism 9.

Lipid extraction and analysis

The dried lipid extracts were re-solubilized in 200 µL of 2:1 CHCl$_3$: MeOH, and 10 µL was injected into an Agilent 6545 QTOF (quadrupole-time-of-flight) instrument for semi-quantitative analysis using high-resolution auto MS/MS methods and chromatography techniques. A Gemini 5U C-18 column (Phenomenex) coupled with a Gemini guard column (Phenomenex, 4x3 mm, Phenomenex security cartridge) was used for LC separation. The solvents used for negative ion mode were buffer A: 95:5 H$_2$O:MeOH + 0.1% ammonium hydroxide and buffer B: 60:35:5 Isopropanol:MeOH:H$_2$O + 0.1% ammonium hydroxide. The 0.1% ammonium hydroxide in each buffer was replaced by 0.1% Formic acid + 10 mM Ammonium Formate for positive ion mode runs. All LC/MS runs were for 60 min, starting with 0.3 mL/min 100% buffer A for 5 minutes, 0.5 mL/min linear gradient to 100% buffer B over 40 minutes, 0.5 mL/min 100% buffer B for 10 minutes, and equilibration with 0.5 mL/min 100% buffer A for

5 minutes. The following settings were used for the ESI-MS analysis: drying gas and sheath gas

temperature: 320°C, drying gas and sheath gas flow rate: 10 L/min, fragment or voltage: 150 V, capillary

voltage: 4000 V, nebulizer (ion source gas) pressure: 45 psi and nozzle voltage: 1000 V. For analysis,

a lipid library was employed in the form of a Personal Compound Database Library (PCDL), and the

peaks were validated based on relative retention times and fragments obtained. This library was

selectively curated from the MycoMass database and the LIPID MAPS Structure Database (LMSD). All

robustly detected lipid species were quantified by normalizing areas under the curve (AUC) to the AUC

of the relevant internal standard added and by normalizing to the total cell number. Log_2 fold changes

were then plotted for each lipid compared to the Msm-ATc samples.

Immunoprecipitation (IP) sample preparation and MS/MS analysis

Sample preparation and, mass spectrometry analysis were outsourced to VProteomics, New Delhi.

Briefly, the IP samples were reduced with 5 mM TCEP, followed by alkylation with 50 mM

iodoacetamide. The samples were digested with Trypsin (1:50, Trypsin/lysate ratio) for 16 h at 37 °C.

Digests were cleaned using a C18 silica cartridge and resuspended in a buffer containing 2%

acetonitrile and 0.1% formic acid. 1 μg of peptide was used for analysis in an Easy-nlc-1000 system

coupled to an Orbitrap Exploris mass spectrometer. The gradients were run for 110 min. MS1 spectra

were acquired in the Orbitrap (Max IT = 60 ms, AGQ target = 300%; RF Lens = 70%; R=60K, mass

range = 375−1500; Profile data). Dynamic exclusion was employed for 30 seconds, excluding all charge

states for a given precursor. MS2 spectra were collected for the top 20 peptides. MS2 (Max IT=60ms,

R= 15K, AGC target 100%). Proteome Discoverer (v2.5) was used to analyse the RAW data against

the Uniprot *Msm* database. The precursor and fragment mass limitations for the dual Sequest and

Amanda searches were established at 10 ppm and 0.02 Da, respectively. The false discovery rate for

proteins and peptide spectrum matches was adjusted to 0.01 FDR. Proteins found in at least three

replicates with minimum unique peptides #2 were considered interactors. To identify top hits, PSM cut-

off was set to 18 and unique peptides to 5.

Protein expression and purification

BL21(DE3) cells transformed with pET-wag31, or pET-wag31K20A or pET-wag31$_{Δ1-60}$ were cultured at

37ºC until the A_{600} ~0.6. 0.1 mM IPTG was added to the culture to induce protein expression, followed

by incubating the culture at 18ºC for 12 h. Cells were pelleted and stored at -40 ºC. The bacterial pellet

was resuspended in a lysis buffer (20 mM HEPES pH 7.4, 150 mM NaCl, and 20 mM imidazole with 1

mM phenylmethylsulphonyl fluoride (PMSF)) and lysed by sonication in an ice-water bath. Lysate was

spun at 30,000 g for 20 min, and the supernatant obtained was incubated with TALON® Metal Affinity

Resin (Takara Bio) 4 ºC for 30 min. The resin was extensively washed with lysis buffer, and the bound

protein was eluted with 20 mM HEPES pH 7.4, 150 mM NaCl, and 250 mM imidazole. The elution was

dialyzed against 20 mM HEPES pH 7.4 150 mM NaCl and stored at 4 ºC for the duration of the

experiments. Aggregates were removed by centrifugation at 100,000 g. Concentration of the proteins

was quantified by measuring A_{280} using the molar extinction coefficient predicted by the Expasy

ProtParam tool.

Liposome preparation

1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine
(DOPE) and1′,3′-bis[1,2-dioleoyl-sn-glycero-3-phospho]-glycerol (sodium salt), CL were obtained from
Avanti Polar Lipids. 1,1'-dilinoleyl-3,3,3',3'-tetramethylindocarbocyanine, 4-chlorobenzenesulfonate
(FAST DiI™), and 3,3'-dilinoleyloxacarbocyanine perchlorate (FAST DiO™) were obtained from
Invitrogen. The UV-activable, diazirine-containing reactive fluorescent lipid probe BODIPY-diazirine
phosphatidylethanolamine (BDPE) was prepared as described earlier (1, 2). For liposome preparation,
lipids were aliquoted from chloroform stocks at desired ratios in a clean glass tube and dried under a
high vacuum for an hour to a thin film. The dried lipids were used at a final concentration of 1 mM
(deionised water was added to them). After hydrating at 50 °C for 30 minutes, lipids were vigorously
vortexed and then extruded through polycarbonate filters with pore sizes of 100 nm (Whatman).. For
PLiMAP assays, liposomes were CL, DOPG, or DOPE with 69 mol% DOPC and 1 mol% BDPE.
Liposomes used in tethering assays were made similarly with 30 mol% CL or DOPE and 1 mol% of the
fluorescent lipid DiI or DiO) in the DOPC background.

NAO imaging analysis

Background correction and Thunder image processing was performed on the images using the Las-X
software module (Leica microsystems). Processed images were then analysed using Fiji (2) and plotted
using Graph Pad prism 9. Briefly, N=100 cells across two independent experiments were examined for
their profile of CL distribution by normalizing the cell length and assigning pole with brighter NAO
fluorescence intensity as 0 and the other pole as 100. The fluorescence intensity was normalised to the
highest fluorescent value for each cell. Corresponding fluorescence intensity value for all cells for every
point between 0 to 100 were calculated from the LOOKUP function of excel. The values representing
CL fluorescence across 100 cells were averaged out and plotted against normalised cell length in
GraphPad prism 9. 0^{th} order smoothening with 4 neighbours on each size was performed on the curves
obtained.

His pulldown assay- *E.coli* and *Msm* lysate preparation

BL21 (DE3) cells transformed with no plasmid, or pET-wag31 or pET-wag31$_{Δ1-60}$ were cultured as
described above and cell lysates were prepared. The expression of Wag31 or Wag31$_{Δ1-60}$ was assessed
by resolving the samples on 10% SDS-PAGE followed by Coomassie staining. The interacting partners,
namely *murG, sepIVA*, *msm2092* and *accA3* were amplified from *Msm* genomic DNA using gene
specific primers harbouring NdeI-HindIII sites and amplicons were digested and cloned into the
corresponding sites in pNit-3X-FLAG shuttle vector. The constructs thus generated were electroporated
in *Msm* to generate *Msm::murG*, *Msm::SepIVA*, *Msm::msm2092* and *Msm::accA3.* The cultures grown
till A$_{600}$ ~0.8 were used for fresh inoculation at A$_{600}$ of 0.05 in 7H9 media containing 0.5 μM IVN. Cultures
were grown for 12 h and lysates were prepared by bead-beating and 40 μg were resolved on 10% gels
and probed with α-FLAG antibody.

Table S1. Plasmids and strains used in the study
117

Plasmid	Description	Antibiotic resistance	Source
pJV53-Apra	pJV53 plasmid from Dr. Hatfull's lab wherein the *kan^r* antibiotic cassette was swapped with apra^r cassette.	apra	(3)
pST-KiT$_{off}$	L5 integrative pST-KiT plasmid harboring reverse tetR gene.	kan	(4)
pST-gfp	pST-KiT$_{off}$ construct harboring gfp between NdeI-HindIII sites.	Kan	This study
pST-*wag31*	pST-KiT$_{off}$ L5 integrative plasmid harboring FLAG- *wag31*-3X HA under the control of tetracycline induction	kan	This study
pET-*wag31*	pET28b harboring His-*wag31*-His	kan	This study
pET-*wag31$_{K20A}$*	pET28b harboring His-*wag31$_{K20A}$*-His	kan	This study
pET-*wag31$_{Δ1-60}$*	pET28b harboring His-*wag31$_{Δ1-60}$*-His	kan	This study
pET-*wag31-gfp*	pET28b harboring His-*wag31-gfp*-His	kan	This study
pSCG-*wag31*	Giles integrative plasmid harboring His-*wag31*-His under the control of wag31 native promoter	apra	This study
pSCG- *wag31$_{K20A}$*	Giles integrative plasmid harboring His-*wag31$_{K20A}$*-His under the control of wag31 native promoter	apra	This study
pSCG- *wag31$_{Δ1-60}$*	Giles integrative plasmid harboring His-*wag31$_{Δ1-60}$*-His under the control of wag31native promoter	apra	This study
pNiT-*wag31*	IVN inducible plasmid harboring 3X-FLAG-*wag31*	kan	This study
pNiT-*murG*	IVN inducible plasmid harboring 3X-FLAG-*murG*	kan	This study
pNiT-*sepIVA*	IVN inducible plasmid harboring 3X-FLAG-*sepIVA*	kan	This study
pNiT-*msm2092*	IVN inducible plasmid harboring 3X-FLAG-*msm2092*	kan	This study
pNiT-*accA3*	IVN inducible plasmid harboring 3X-FLAG-*accA3*	kan	This study

118
119

Strain	Description	Antibiotic resistance	Source
E. coli DH5α	*E. coli* strain used for cloning	none	Invitrogen
E. coli BL21DE3	*E. coli* strain used for protein expression	kan	Novagen
Msm	*M. smegmatis* mc^{2}155 strain	none	ATCC
Msm::pJV53	*Msm* strain electroporated with pJV53 plasmid that expresses recJ and	apra	This study
Msm::gfp	*Msm* strain electroporated with pST-KiT-gfp		
Msm::pS-wag31	*gp60* and *gp61* recombinase expressing Msm::pJV53 strain electroporated harboring integrative pST-KiT$_{off}$	apra, kan	This study
Δwag31	wag31 mutant in *Msm* in which the native *wag31* gene was replaced with hyg^r marker. The strain harbours integrative tet inducible pST-*wag31* at L5 site. Cured to remove pJV53 episomal copy.	hyg, kan	This study
Δwag31::wag31	*Δwag31* electroporated with pSCG-wag31	hyg, kan, apra	This study
Δwag31::wag31$_{K20A}$	*Δwag31* electroporated with pSCG-wag31$_{K20A}$	hyg, kan, apra	This study
Δwag31::wag31$_{\Delta1-60}$	*Δwag31* electroporated with pSCG-wag31$_{\Delta1-60}$	hyg, kan, apra	This study
Msm::wag31	*Msm* electroporated with pNiT-*wag31*	kan	This study
Msm::murG	*Msm* electroporated with pNiT-*murG*	kan	This study
Msm::sepIVA	*Msm* electroporated with pNiT-*sepIVA*	kan	This study
Msm::msm2092	*Msm* electroporated with pNiT-*msm2092*	kan	This study
Msm::accA3	*Msm* electroporated with pNiT-*accA3*	kan	This study

121
122

References

1. A. Parikh *et al.*, Development of a new generation of vectors for gene expression, gene replacement, and protein-protein interaction studies in mycobacteria. *Appl Environ Microbiol* **79**, 1718-1729 (2013).
2. J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012).
3. J. C. van Kessel, G. F. Hatfull, Recombineering in Mycobacterium tuberculosis. *Nat Methods* **4**, 147-152 (2007).
4. V. Soni *et al.*, Depletion of M. tuberculosis GlmU from Infected Murine Lungs Effects the Clearance of the Pathogen. *PLoS Pathog* **11**, e1005235 (2015).

Table S2: unique interacting partners found for Wag31

Accession	Gene ID	# PSMs	# Unique Peptides	Localisation (Based on PMID: 34965429, 27114527, 20412585)	Functional category
A0QYB1	MSMEG_3596	185	37	Unknown	replication, recombination & repair
A0QTE7	MSMEG_1813	104	16	IMD	lipid metabolism
Q3L891	MSMEG_0402	58	16	PM-CW	cell wall & cell processes
A0R1Y2	MSMEG_4915	61	16	PM-CW	Intermediary metabolism and respiration
A0R152	:MSMEG_4626	37	13	IMD	Translation, ribosomal structure & biogenisis
A0QWG2	MSMEG_2931	44	14	PM-CW	Translation, ribosomal structure & biogenisis
A0QPS6	MSMEG_0502	34	10	unknown	Intermediary metabolism and respiration
A0R684	MSMEG_6459	29	11	Cytosolic	Intermediary metabolism and respiration
A0R582	MSMEG_6099	41	10	PM-CW	Cell wall and cell processes
A0R016	MSMEG_4227	34	9	IMD	Cell wall and cell processes
A0QQW2	MSMEG_0897	20	7	unknown	conserved hypothetical
A0QU64	MSMEG_2092	41	7	unknown	Cell wall and cell processes
A0R7C9	MSMEG_6863	25	5	unknown	cell wall & cell processes
A0QX15	MSMEG_3138	28	4	Cytosolic	Intermediary metabolism and respiration
A0QPE7	MSMEG_0372	24	6	IMD	Lipid metabolism
A0QV38	MSMEG_2436	21	4	IMD	unknown
A0QUZ5	MSMEG_2393	24	6	Cytosolic	Lipid metabolism
A0QV18	MSMEG_2416	13	3	Cell wall	cell wall & cell processes
A0QRA5	EG_1046, MSMEG	12	4	Cell wall	Cell wall and cell processes
A0QUE0	MSMEG_2174	25	7	Cell wall	replication, recombination & repair
A0R2R8	MSMEG_5211	23	6	unknown	**Amino acid transport and metabolism**
A0R7J6	MSMEG_6941	18	5	IMD	conserved hypotheticals
A0QU53	MSMEG_2080	18	6	IMD	lipid metabolism
A0QQ17	MSMEG_0594	16	6	IMD	Intermediary metabolism and respiration
A0R3C8	MSMEG_5427	20	7	unknown	Intermediary metabolism and respiration
A0QVL0	MSMEG_2611	18	4	unknown	Intermediary metabolism and respiration
A0QXI0	MSMEG_3309	10	3	unknown	conserved hypothetical
A0QPG1	MSMEG_0386	7	3	unknown	Cell wall and cell processes
A0R4S6	MSMEG_5937	18	4	unknown	Intermediary metabolism and respiration
A0QV82	MSMEG_2481	11	4	unknown	conserved hypothetical
A0QWS9	MSMEG_3051	16	3	unknown	Intermediary metabolism and respiration
A0QWS2	MSMEG_3044	14	4	unknown	Intermediary metabolism and respiration
P0CH36	MSMEG_1037	16	4	unknown	Intermediary metabolism and respiration
A0R083	MSMEG_4294	10	3	unknown	Intermediary metabolism and respiration
A0QQX7	MSMEG_0912	14	4	unknown	lipid metabolism
A0QYE7	MSMEG_3632	10	3	PM-CW	Intermediary metabolism and respiration
A0QYG9	MSMEG_3654	13	4	PM-CW	cell wall & cell processes
A0QV36	MSMEG_2434	10	3	unknown	conserved hypotheticals
A0QVP7	MSMEG_2648	11	4	cell wall	lipid metabolism
A0QWX7	MSMEG_3100	14	4	unknown	Intermediary metabolism and respiration
Q3L887	MSMEG_0406	16	3	unknown	lipid metabolism
A0QX25	MSMEG_3148	15	4	Cell wall	conserved hypothetical
A0QZ46	MSMEG_3894	10	3	IMD	Intermediary metabolism and respiration
A0QUY3	MSMEG_2379	10	3	unknown	Intermediary metabolism and respiration
A0QRB9	MSMEG_1061	10	3	cell wall	replication, recombination & repair
A0R7D8	MSMEG_6874	13	3	unknown	Intermediary metabolism and respiration
Q9X5M1	MSMEG_4958	16	4	unknown	cell wall & cell processes
A0R7G3	MSMEG_6901	14	3	unknown	conserved hypothetical
A0R5K0	MSMEG_6219	12	1	PM-CW	regulatory protein
A0QW11	MSMEG_2768	19	3	unknown	anscription, replication, recombination & repair
A0R597	MSMEG_6114	10	3	unknown	Intermediary metabolism and respiration
A0QWN3	MSMEG_3003	10	2	unknown	Translation, ribosomal structure & biogenisis
A0QRD2	MSMEG_1073	14	4	IMD	Intermediary metabolism and respiration
A0R0N6	MSMEG_4450	10	3	IMD	replication, recombination & repair
A0QTF8	MSMEG_1825	10	4	unknown	Intermediary metabolism and respiration
A0R6C4	MSMEG_6499	10	2	unknown	conserved hypotheticals
A0QYQ7	MSMEG_3746	12	3	Cell wall	Intermediary metabolism and respiration
A0R184	MSMEG_4659	12	3	unknown	regulatory protein
A0R061	MSMEG_4272	10	2	cell wall	Intermediary metabolism and respiration

A0QXW7	MSMEG_3451	10	2	unknown	Intermediary metabolism and respiration
A0QYU0	MSMEG_3785	9	3	unknown	Unknown
A0R2Q7	MSMEG_5199	9	2	unknown	lipid metabolism
A0R1D1	MSMEG_4709	8	3	unknown	lipid metabolism
A0QTQ8	MSMEG_1930	8	3	IMD	replication, recombination & repair
A0R2Q5	MSMEG_5197	8	3	unknown	lipid metabolism
A0R3T9	MSMEG_5592	8	2	unknown	Intermediary metabolism and respiration
A0R6D9	MSMEG_6514	8	2	unknown	Virulence, detoxification, adaptation
A0QNZ7	AMSMEG_0220	8	2	IMD	Intermediary metabolism and respiration
A0QZ33	MSMEG_3880	8	2	unknown	Intermediary metabolism and respiration
A0R4P1	MSMEG_5903	8	2	unknown	Intermediary metabolism and respiration
A0QS40	MSMEG_1340	8	2	unknown	conserved hypotheticals
A0QQH7	MSMEG_0759	8	2	unknown	Intermediary metabolism and respiration
A0QSK8	MSMEG_1514	8	2	unknown	conserved hypotheticals
A0R6M2	MSMEG_6598	8	2	unknown	Intermediary metabolism and respiration
A0R436	MSMEG_5691	7	3	unknown	conserved hypothetical protein
A0R5X8	MSMEG_6351	7	3	unknown	Intermediary metabolism and respiration
A0QT17	MSMEG_1679	7	2	Cell wall	unknown
A0QXN2	MSMEG_3363	7	2	unknown	regulatory protein
A0R0W2	MSMEG_4528	7	2	unknown	Intermediary metabolism and respiration
A0R066	MSMEG_4276	7	2	unknown	Intermediary metabolism and respiration
A0QT08	MSMEG_1670	6	3	PM-CW	Intermediary metabolism and respiration
A0QPW2	MSMEG_0539	6	3	PM-CW	Intermediary metabolism and respiration
Q3I5Q7	MSMEG_0919	6	3	unknown	Cell wall and cell processes
A0QZ93	MSMEG_3942	6	3	unknown	conserved hypothetical protein
A0R6D7	MSMEG_6512	6	2	unknown	lipid metabolism
A0R071	MSMEG_4282	6	2	IMD	conserved hypothetical protein
A0R048	MSMEG_4258	6	2	PM-CW	Intermediary metabolism and respiration
A0R753	MSMEG_6783	6	2	unknown	Unknown
A0QRA8	EG_1049;MSMEG	6	2	unknown	Conserved hypotheticals
Q9X5M0	MSMEG_4959	6	2	PM-CW	Translation, ribosomal structure & biogenisis
A0QUW2	MSMEG_2357	6	2	unknown	Intermediary metabolism and respiration
A0QNG7	MSMEG_0035	6	2	PM-CW	regulatory protein
A0QX83	MSMEG_3207	6	2	unknown	Intermediary metabolism and respiration
A0QVL6	MSMEG_2617	6	2	unknown	Intermediary metabolism and respiration
A0R170	MSMEG_4645	6	2	unknown	Intermediary metabolism and respiration
A0QYD4	MSMEG_3619	6	2	PM-CW	Intermediary metabolism and respiration
A0R2U7	MSMEG_5239	6	2	unknown	Intermediary metabolism and respiration
A0R6D6	MSMEG_6511	6	2	unknown	lipid metabolism
A0QWQ3	MSMEG_3024	5	2	unknown	Conserved hypotheticals
A0R177	MSMEG_4652	5	2	unknown	Unknown
A0QXB8	MSMEG_3244	5	2	unknown	conserved hypothetical protein
A0QTT2	MSMEG_1954	4	2	PM-CW	Cell wall and cell processes
A0QP17	MSMEG_0240	4	2	unknown	conserved hypothetical protein
A0QYT3	MSMEG_3777	4	2	unknown	Translation, ribosomal structure & biogenisis
A0R2L4	MSMEG_5156	4	2	unknown	replication, recombination & repair
A0R5I9	MSMEG_6208	4	2	unknown	lipid metabolism
A0QWU9	MSMEG_3071	4	2	PM-CW	Intermediary metabolism and respiration
A0QQL0	MSMEG_0793	4	2	unknown	Intermediary metabolism and respiration
A0QR20	MSMEG_0956	4	2	unknown	Intermediary metabolism and respiration
A0QVT2	MSMEG_2685	4	2	unknown	Conserved hypotheticals
A0R2Q6	:MSMEG_5198	4	2	unknown	lipid metabolism
A0QX95	MSMEG_3219	4	2	Cell wall	Intermediary metabolism and respiration
A0QP16	MSMEG_0239	4	2	unknown	Intermediary metabolism and respiration
A0QZY4	MSMEG_4193	4	2	Cytosolic	Conserved hypotheticals
A0R3L4	MSMEG_5515	4	2	Cytosolic	Intermediary metabolism and respiration
A0R5N7	MSMEG_6256	3	2	PM-CW	Intermediary metabolism and respiration