Xenosiderophore transporter gene expression and clade-specific filamentation in *Candida auris* killifish infection

Hugh Gifford[a], Tina Bedekovic[a], Nicolas Helmstetter[a], Jack Gregory[a], Qinxi Ma[a], Alexandra C. Brand[a], Duncan Wilson[a], Johanna Rhodes[b,c], Mark Ramsdale[a], Tetsuhiro Kudoh[d,1], Rhys Farrer*[a,1]

Abstract

Candida auris is an emerging infectious agent and World Health Organisation (WHO) critical priority fungal pathogen. Rising drug resistance, massive nosocomial outbreaks and diagnostic challenges complicate clinical treatment, resulting in a patient mortality rate of ~45%. Surprisingly, gene expression profiles of *C. auris* have not yet been described during infection *in vivo*. To understand transcriptional responses during in-host infection, we developed a thermo-relevant fish embryo yolk-sac microinjection model (*Aphanius dispar*; Arabian killifish; AK) that mimics human body temperature. This allowed us to interrogate infection dynamics through dual host-pathogen RNA-seq at 24 and 48 h post injection (HPI) at 37 °C across the five major clades (I-V) of *C. auris*. Host gene expression following infection indicated features of heat shock, complement activation, and nutritional immunity, including haem oxygenase (*HMOX*) expression in response to clade IV infection. We identified an *in vivo* transcriptional signature across five clades of *C. auris* that was highly enriched for putative xenosiderophore transmembrane transporters. We describe this newly-discovered seventeen-member xenosiderophore transporter candidate (*XTC*) family in terms of individual gene expression patterns, and a sub-clade of five putative haem transport-related (*HTR*) genes, also up-regulated during infection. Only the basal clade V isolate formed filaments during infection, coinciding with typical and atypical regulators of

morphogenesis, including *UME6*, *HGC1*, and the novel adhesin *SCF1*. Clades I and IV

demonstrated increased virulence, coinciding with up-regulation of three *HTR* genes in

clade IV, and the mating-type locus (*MTL*) non-mating gene *PIKA* in both. Our study

suggests that *XTC* and *HTR* genes may play a critical role in *C. auris* virulence, making

excellent targets for further investigation and potential therapy.

Keywords: *Candida auris*, virulence, gene expression, transcriptome, pathogenicity,

Arabian killifish, *Aphanius dispar*, siderophore transport, mating-type locus,

filamentation

[a]MRC Center for Medical Mycology, University of Exeter, Geoffrey Pope Building,

Stocker Road, Exeter, United Kingdom, EX4 4QD

[b]Department of Medical Microbiology, Radboudumc, Geert Grooteplein Zuid 10,

Nijmegen, Netherlands, 6525 GA

[c]MRC Centre for Global Infectious Disease Analysis, Imperial College London South

Kensington Campus, London, United Kingdom, SW7 2AZ

[d]Biosciences, University of Exeter, Geoffrey Pope Building, Stocker Road, Exeter,

United Kingdom, EX4 4QD

[1]Co-Senior Author

*Corresponding Author - R.Farrer@exeter.ac.uk

Introduction

C. auris is an ascomycetous yeast in the *Metschnikowiaceae* family that has

rapidly transitioned from recent emergence to global public health threat[1] and World

Health Organisation (WHO) critical priority human fungal pathogen[2]. Despite the oldest

known historical isolate dating from 1996[3], the first formal species identification

occurred in 2008[4]. Within ten years, *C. auris* had caused nosocomial outbreaks around

the world, with high rates of azole antifungal drug resistance and a rising incidence of

pan-drug resistance[5]. These factors required extreme measures including Intensive

Care Unit (ICU) shutdowns and massive infection control costs[6–8]. Worryingly, *C. auris*

is now a leading cause of invasive candidiasis in several hospital settings globally[9,10].

Distinct clades of *C. auris*, including a sixth clade identified and described in

publications only in 2024[11,12], indicate the simultaneous emergence of multiple lineages

that are highly adapted for human infection[13,14]. The clinical impact, phenotypic traits

and simultaneous emergence of multiple clades of *C. auris* underscore the urgency to

understand its mechanisms of pathogenicity.

The factors governing in-host fitness and virulence during infection are poorly

understood. *C. auris* bloodstream infection is associated with a crude mortality of

~45%[15]. Virulence factors in other distantly related *Candida* species, such as

C. albicans, include hypha-mediated invasion[16], the invasin Als3[17], and the secreted

toxin, candidalysin[18]. In contrast, *C. auris* has been shown to reproduce as budding

yeasts in mammalian tissues[19,20]. Although *C. auris* possesses multiple adhesins

associated with an aggregation phenotype[21,22], there has been no evidence of toxin

synthesis[23,24]. *C. auris* evades host immune responses through cell surface masking[25],

secreted aspartyl protease digestion of host tissue[26], and intra-phagocytic survival with

cell lysis[27]. Clades I, III, and IV are more commonly associated with invasive disease

and nosocomial outbreaks, while clades II and V have been primarily reported in cases

of otomycosis and rarely in bloodstream infection[28]. Human mortality associated with

C. auris candidaemia is potentially higher with clades I and IV than clade III, but this

difference has not been shown to be statistically significant[29]. Clade and strain-specific

genotypic and phenotypic variation in these traits have been described and may

underlie differences in clinical presentation and pathogenesis.

Gene expression profiling *in vivo vs in vitro* conditions has become a standard

approach for assessing host and pathogen responses to infection. To date, no

experiment has described the gene expression of *C. auris* or its host during *in vivo*

infection, although studies using human whole blood *ex vivo*[30], co-incubation with

human dermal fibroblasts[31] or murine cell line-derived macrophages[32] have provided

useful insights. Additionally, host responses have been assessed in human peripheral

blood-derived monocytes[33] and murine bone marrow-derived macrophages[34], but not in

a living host. A dual RNA-seq study using the *ex vivo* whole blood model demonstrated

similar host responses to several *Candida* species such as *C. albicans*, but identified a

unique pattern of pathogen gene expression for *C. auris*, including up-regulation of

transporter genes such as orthologues of the *C. albicans* siderophore transporter

SIT1[30]. Interestingly, this family of predicted xenosiderophore transporter genes is

significantly expanded in several members of the *Metschnikowiaceae* clade compared

to other *Saccharomycetae*[23], raising questions as to the roles of siderophore

transporters in *C. auris* biology[35].

The dearth of studies profiling host-pathogen gene expression *in vivo* for *C. auris*

infection may be explained by multiple challenges. Most murine models, such as

BALB/c[19,36–38] and C57BL/6 mice[34], have only very rarely demonstrated lethality through

C. auris infection[33,39,40] without major immunosuppression with

cyclophosphamide/cortisone[41–52], monoclonal antibody neutrophil depletion[20,53], or other

immunodeficiency[21,54,55]. Additionally, mouse models are also expensive and associated

with concerns over environmental pathogen shedding[56]. Alternative host models such

as *Galleria mellonella*[36,46,57–63], *Drosophila melanogaster*[64] and

Caenorhabditis elegans[62] have been employed, but these may lack key elements of

mammalian-relevant immune systems, thus limiting their scope of application. In

patients, immune control of *C. auris* is thought to depend on myeloid cell activity *via* C-

type lectin receptors such as complement receptor 3 and mannose receptor MMR,

followed by interleukin (IL)-17 pathway activation[65,66]. Therefore, an animal model that

closely aligns with these mechanisms is required for human-relevant profiling of host-

pathogen interactions.

Though the environmental origins of *C. auris* have not been conclusively

demonstrated, a marine origin has been hypothesised and supported by successful

isolation of *C. auris* in coastal waters of both Indian and Pacific oceans[67,68]. Notably, a

close relative of *C. auris* in the *Metschnikowiaceae* clade, *C. haemulonii*, was first

isolated in 1961 from the gut of blue-striped grunt (genera name "*Haemulon*" for their

blood-red mouth interior)[69]. Therefore, a marine halotolerant model could shed light on a

possible evolutionarily adaptive niche. Fish larvae must survive hostile and varied ocean

environments, demonstrating very early innate immune responses such as antimicrobial

peptide production and the complement cascade[70,71]. Furthermore, adult teleost fish

possess functional IL-17 subtypes (A and F) that lead to T-helper-17 subset

differentiation and adaptive immunity in response to fungal infection[72]. The zebrafish

(*Danio rerio*) model has allowed hindbrain[25,73] and swim bladder[74] microinjection

modelling of *C. auris* infection, but rarely at temperatures above 30 °C, which zebrafish

are broadly intolerant of. Again, we are unaware of any in-host transcription studies

using the zebrafish model. Fungal transcriptional programmes can vary greatly across

temperatures and are critical for mammalian pathogenesis[75–77]. A thermotolerant teleost

fish embryo model such as the *Aphanius dispar*, also known as the Arabian killifish

(AK), which can acclimatise to temperatures up to and including 40 °C[78–81] offers a

balance between considerations of suitability for experimental manipulation, applicability

to human fungal infection at mammalian temperatures, and relevance to hypothetical

origins in an emerging pathogen.

In this paper, we present the first study exploring in-host virulence of *C. auris*

clades I-V in the AK yolk-sac microinjection model at 37 °C, with sufficient power to

detect significant differences in virulence. We supplemented our validation of the model

with histology, fungal burden recovery, and dual host-pathogen RNA-seq. By comparing

in vivo and *in vitro* expression profiles of *C. auris*, we detected a cross-clade expression

signature that was significantly enriched for siderophore transporter *SIT1* paralogues,

which mediate uptake of siderophores produced by other microbes (xenosiderophores).

We designated this expanded gene family as xenosiderophore transporter candidates

(*XTC*), which included a haem transport-related (*HTR*) sub-clade. We also identified

in vivo filamentation by clade V during infection, which was associated with orthologues

of key hypha-associated genes from *C. albicans*. Together, these findings substantially

enhance our knowledge of the processes underpinning *C. auris* infection.

Results

***C. auris* clades I-V are virulent in the AK microinjection model and clade V**

uniquely exhibits in-host filamentation

Microinjection of the AK yolk-sac with each of the five *C. auris* clades was lethal

at 37 °C (91.7-100% 7-day mortality for each clade) compared to sham injection (5.6%

mortality). Kaplan-Meier survival curves demonstrated a significantly higher lethality for

clade I (*vs* II, III or V, $p < 0.001$) and clade IV (*vs* II or III, $p < 0.04$) (**Fig. 1A**; **Table S1**).

Growth of the clade II strain in YPD-broth at 37 °C was noticeably slower over 24 h

compared with other clades (**Fig. S1A**). However, colony-forming units (CFU) recovered

post-infection from AK embryos were commensurate with similar in-host growth (**Fig.**

1B), with no differences identified between clades immediately post injection ($x = 673$

CFUs per infected embryo), 24 h post injection (HPI) ($x = 1.12 \times 10^5$) or 48 HPI ($x = 1.78$

$\times 10^5$), except between clade III and V at 48 HPI (x 4.48 $\times 10^5$ *vs* 1.56 $\times 10^4$, $p < 0.01$).

These results indicate that AK embryo yolk-sac microinjection is a suitable model for the

study of lethal *C. auris* infection at mammalian temperature.

Histological sectioning of the AK yolk-sac at 48 HPI revealed budding yeast

morphologies, except for the clade V strain, which demonstrated predominantly

filamentous forms (**Fig. 1C**). When grown in YPD-broth before microinjection, all

isolates exhibited yeast morphologies; clade V colonies recovered from AK yolk-sac

infection and grown on YPD-agar demonstrated rough (as opposed to smooth) colonies

in approx. 1-2% of plated colonies (**Fig. S1B**), which consisted of elongated yeast cells

and filamentous structures (> 10-50 µm) reminiscent of pseudohyphae (**Fig. S1C**).

Death was defined by cessation of heartbeat, which was followed by yolk-sac collapse

visible by timelapse microscopy (**Fig. 1D**). These findings informed the study design for

comparing *C. auris* gene expression between individual clades during *in vivo* and

in vitro growth to detect drivers of phenotypic and morphological variation relevant to in-

host virulence (**Fig. 1E**).

C. auris **triggered host iron-nutritional immunity, heat shock response, and**

complement activation

To determine whether microinjection interefered with host transcriptional

responses, we identified differentially expressed genes (DEGs) between sham injection

and no injection, defined as genes with significant ± >2 log-fold change in expression

after multiple testing. Though principle components analysis did not separate gene

expression profiles (**Fig. S1D**), correlation clustering separated sham injection and

negative controls without injection (**Fig. S1E**). However, there were only 30 DEGs for

sham injection compared to no injection (0.11% detected genes; **Fig. 2A**), with up-

regulation of zero and seven genes (at 24 and 48 HPI, respectively) and down-

regulation of four and nineteen genes (24 and 48 HPI, respectively). Four GO terms

were significantly enriched in the set of genes up-regulated at 48 HPI (relating to

sarcomere generation, organisation, M-band and muscle) with no enriched KEGG

pathways or PFAM domains, according to a falsed discovery rate (FDR) cut-off of

0.001. These GP terms were driven exclusively by three orthologues of murine *TITIN*

(log-fold change x = 7.59, gene ratio 0.16), a ~ 3-4 MDa cardiac muscle protein with a

putative role in haemodynamic stress tolerance and volume overload mechano-sensing

(**Fig. 2B**). Other up-regulated genes included vascular adhesion molecules and

hypothetical immunomodulator, dipeptidase 1 (*DPEP1*, **Table S2**). High survival of AK

embryos receiving sham injection and minor changes in gene expression (consistent

with physiological responses to exogenous fluid bolus) supported continued analysis of

host responses to infection.

To characterise host responses to *C. auris* during infection, we identified host

DEGs between infection by each clade and sham injection. Principle components

analysis (**Fig. S1D**) and correlation clustering (**Fig S1F**) did not cluster gene expression

profiles separately at either time-point (24 and 48 HPI). We identified 685 DEGs during

infection with any clade (2.49% potential genes), of which 183 were up-regulated and

508 were down-regulated in response to at least one clade. There were no DEGs

shared in common across host responses to all five clades, and only a small set of four

up-regulated genes and one down-regulated gene shared across host responses to four

clades of infection (**Table S2**). Up-regulated genes included Laminin subunit alpha-1

(*LAMA1*, not significant in response to clade III) and three genes that were not

significant in response to clade II, Glucose-6-phosphatase (*G6PC*), DN117939

(hypothetical protein) and Tryptophan aminotransferase protein 1 (*TAR1*);

immunoglobulin-like fibronectin type III domain-containing protein 1 (*IGFN1*) was down-

regulated. These results suggest AK larvae have a subtle and non-uniform response to

each clade of *C. auris* microinjection at these time-points, also reflecting levels of

virulence and stages of infection.

We identified clade-specific AK responses to infection. At 24 HPI, clade II elicited

the greatest host response, with up-regulation of 41 genes and down-regulation of 210

genes (**Fig. 2A**). However, at 48 HPI, clade IV elicited the greatest response (49 up-

regulated, 104 down-regulated). Heat shock proteins (*HSP30* and *HSP70*, but not

HSP90), complement cascade elements, pattern recognition receptors (PRRs),

inflammatory transcription regulators, and modulators of nutritional immunity were

differentially expressed with higher log-fold changes in clades I and IV, especially at

48 HPI (**Fig. 2C**). The most prominent gene relevant to pathogen recognition was

CARD14 in clade I and IV infection at 48 HPI (**Fig. 2D and 2E**). We found *NLRP3*

(inflammasome) expression during clade III infection at 24 HPI only. There was no

significant differential expression of several fungal immune response orthologues, such

as *IL-17*, for infection with any clade. Together, these findings suggest that heat shock

proteins, nutritional immunity, and the complement effector system are important

elements of the AK larval response to *C. auris* microinjection at 37 °C.

In order to highlight the roles of gene families in the host repsonse, we annotated

the AK transcriptome and identified enriched GO terms, KEGG pathways and PFAM

domains. Most significantly enriched gene features were identified from responses to

infection with the more virulent clades (I and IV) at 48 HPI (**Fig. S2**). The highest mean

log-fold changes for any enriched gene features were for clade IV infection, with haem

catabolism, haem binding, and haem oxygenase (*HMOX*) related GO terms, KEGG

pathway and PFAM domains. Two host orthologues of *HMOX* drove most of these

enrichments, such as haem oxidation (log-fold change x = 3.96, FDR < 0.0003, **Fig.**

2E). In terms of other genes involved in iron-nutritional immunity, the AK also up-

regulated Ferroxidase Hephaestin (*HPHL1*) in response to clade IV infection, but not

Hepcidin (*HEPC1*). The response to clade I infection included 15 enriched GO terms,

driven by up-regulation of two orthologues of Laminin subunit-α (*LAMA1*, *LAMA2*, **Fig.**

2D). No GO terms, KEGG pathways, or PFAM domains were enriched across

responses to all five clades. Only one GO term was enriched across responses to four

clades, which related to sarcomeric Z-disc in down-regulated genes at 48 HPI (II-V),

driven by genes involved in connective and muscular tissue, including *IGFN1* and

Calpain-3 (*CAN3*). The response to clade IV infection, in particular, indicates activation

of iron metabolism and sequestration, potentially comprising an iron-nutritional immunity

response to infection.

C. auris **xenosiderophore transporter expression was enriched across all five**

clades during in-host infection

To identify infection-specific *C. auris* gene expression, we compared the in-host

transcriptome to expression induced by growth in a standard rich laboratory media

(YPD). Principle components and sample correlation clustering from gene expression

data indicated a predominant separate clustering of *in vivo vs in vitro* conditions,

indicating that the gene expression profile of *C. auris* during infection is substantially

different to that when it is grown in nutrient-rich laboratory media (**Fig. S1G and S1H**).

We identified 960 DEGs between *in vivo* AK infection at 24 HPI or 48 HPI *versus in vitro*

culture in YPD over 24 h (499 up-regulated, 523 down-regulated, 17.7% of potential

genes, **Fig. 3A**). We identified few DEGs between the two time-points 24 HPI to 48 HPI,

with zero DEGs for clade IV and V (**Fig. 3B**). Numbers of up-regulated DEGs unique to

any single set were highest for individual clades, particularly in the filamentous clade V

at 24 HPI and in the more virulent clades I and IV at 48 HPI (**Fig. 3C-D**). We annotated

clade-specific reference genomes, and performed differential expression to unveil

enriched gene features across all five clades. Six GO terms, KEGG terms or PFAM

domains were enriched across in-host all five clades at 48 HPI, related to major

facilitator superfamily carbohydrate and siderophore-iron transmembrane transport (**Fig.**

S4). This highlighted the expected importance of up-regulation of genes involved in

sugar uptake *in vivo vs in vitro*, but also raised the unexpected possibility that up-

regulation of members of the *C. auris* expanded gene family of siderophore transporters

could be a general feature of in-host infection.

To detect a signature of *in vivo* gene expression across the species, we identified

the set of 31 DEGs that were significant for all five clades of *C. auris* (**Fig. 3E, Table**

S3). Thirteen DEGs common to all clades of in-host transcripion were transmembrane

transporters, including up-regulation of transporters for xenosiderophores (*SIT1_3921*,

SIT1_1487), sugar (*HGT2*, *HGT12*), ammonia (*ATO2*), and nicotinic acid (*TNA1*), and

down-regulated drug-resistance-related efflux pumps (*MDR2* and *DTR1*) and peptide

transporters (*TPO3*, *PTR22*). Several putative virulence factors such as lipase *LIP1* and

cell wall-related *KRE6* and *SOD6* were also down-regulated across all five clades. We

demonstrated prominent DEGs (such as down-regulated *ALS4* and *MDR1*) that were

significant for at least one clade by mean log-fold change expression volcano plots (**Fig.**

3F). We also noted that several members of the *SIT1* family were differentially

expressed by at least one clade, but were not significant across all five clades, including

five genes that were significantly up-regulated across four clades (*SIT1_1542*,

SIT1_1457, *SIT1_1458*, *SIT1_2110*, and *SIT1_4097*) and five that were up-regulated in

one (*SIT1_0002* and *SIT1_2465*), two (*SIT1_3908* and *SIT1_2465*) or three clades
(*SIT1_2241*). Two more family members were down-regulated across three clades
(*SIT1_1519* and *SIT1_1499*). To nuance these findings further, we noted that two up-
regulated genes were also down-regulated in one (*SIT1_3908*) or two clades
(*SIT1_2465*). The many up-regulated members of the *SIT1* expanded gene family
highlighted the potential importance of the wider family for in-host survival.

To examine orthologues across reference genomes for each clade (**Fig. S3**), we
identified 5,324 orthogroups across clades I-V with outliers *C. haemulonii* and
C. albicans containing 95.4% of all genes (n = 38,345 genes, **Fig. S3A-B**). The use of
clade-specific reference genomes identified higher numbers of DEGs (692 *vs* 499 up-
regulated and 714 *vs* 523 down-regulated genes) including genes within the accessory
genome (55 *vs* 40 up-regulated and 57 *vs* 45 down-regulated, **Fig. S3C**), which we
defined as orthogroups of genes not found in every clade. The core genome across all
three species and five clades of *C. auris* included 3619 single copy orthologues
(x = 66.9% *C. auris* genome per clade) and a further 869 single copy orthologues in
C. auris (x = 82.9% *C. auris* genome per clade), of which 133 were unique to *C. auris*.
The remaining core genome included 486-585 multiple copy orthologues shared across
all species and 110-121 in *C. auris*, of which 17-19 were present in each clade
(x = 643.2 genes per clade), bringing the total estimated core genome size to
5131.2/5413.2 genes per clade (x = 94.8%). A single copy orthologue tree confirmed
the basal status of clade V, and synteny plotting was consistent with limited
transversions/inversions within *C. auris* outwith clade II **Fig. S3A**. These findings are
consistent with a highly conserved core yeast genome across *Candida* species,
especially within the *Metschinikowiaceae* clade, which appears to contain a degree of
preserved genome structure.

We observed high correlation between logFC values obtained from clade-specific
and core (B8441) reference genomes (Pearson's >= 0.95, *p* <0.001 per comparison).
Among DEGs that had no orthologues in the clade I reference genome (**Fig. S3D**), the

highest number of clade-unique genes that were differentially expressed was identified
in clade IV; other DEGs included genes involved in cell wall, intracellular transport,
metabolism, transcription/translation and transmembrane transport (**Table S3**). In terms
of siderophore transporters, we additionally identified a clade IV unique gene associated
with a siderophore transport GO term that was differentially expressed at both time-
points of infection (CJJ09_005327). Overall, the *in vivo* expression profile across five
C. auris clades indicated a potential role for the expanded gene family of siderophore
transporters during *C. auris* in-host survival and pathogenicity, which was up-regulated
compared to growth in nutrient-rich laboratory media.
336
337 **More virulent *C. auris* clades I and IV up-regulated the non-mating *MTL* gene,**
338 ***PIKA*, and down-regulated *MDR1***
339
Given significant differences in virulence between clades, we examined gene
expression differences between clades I and IV (more virulent) and II and III (less
virulent), excluding clade V due to its major morphological differences. We identified
623 up-regulated and 779 down-regulated DEGs expressed in clades I and IV, of which
14 were up-regulated and 22 were down-regulated across all comparisons. In order to
explore potential drivers of virulence, we excluded DEGs that were significant only for
in vitro comparisons and examined the remaining set of 10 DEGs shared across all
in vivo comparisons (I *vs* II, I *vs* III, IV *vs* II, IV *vs* III, **Fig. 4A**, **Table S3**). Strikingly, three
of these genes were non-mating genes located at the mating-type locus (*MTL*),
including poly-A polymerase *PAP1*, oxysterol binding protein *OBPA* and
phosphatidylinositol 4-kinase *PIKA* (**Fig. 4B**). Within the set of 10 DEGs shared across
all comparisons, several genes outside the *MTL* were up-regulated, including vacuolar
protein sorting gene, *VPS70*, and filamentous growth regulator, *FGR14*. Additionally,
the drug efflux pump, *MDR1*, was down-regulated across all comparisons, supporting
the idea of a fitness trade-off between resistance and virulence.
355

Notably, clades I and IV contain the **a**-type *MTL*, whereas clades II, IV and V

contain the α-type *MTL*. That both *MTL* contain the non-mating genes is a conserved

feature across *Candida* species, in which the *MTLa* and *MTLα* genes occupy alternative

positions. Comparing transcript levels across *MTL* genes in all five clades, we observed

poorer detection of *MTLα* transcripts when aligned to the clade I reference genome, as

expected (**Fig. S5A**). Direct transcript-level testing, with correction for multiple testing to

minimise false positives, indicated that only up-regulation of *PIKA* and *PGA30* (48 HPI)

and down-regulation of *THI4* (24 HPI) and *MDR1* (at both time-points) were significantly

different across the set of 10 DEGs shared across all comparisons (**Fig. S5B**).

We used the clade I reference genome to annotate functional domains and

discover enriched gene families, revealing enrichment of up-regulated hypha-regulated

cell wall GPI-anchored proteins in more virulent clades (**Fig. S4**), driven by three pairs

of orthologues of *IFF4*, *HYR3* and *RBR3* (**Table S3**). Four secretory lipase *LIP1*

paralogues drove an enrichment of the related PFAM domain for clade I *vs* III at 24 HPI.

When comparing clade I and IV infection to clade II and 48 HPI, we also observed up-

regulated nutrient transporters including a 7 transmembrane domain putative ferric

reductase/iron importer, *CFL4*, as well as ferric reductase, *FRE3*, and siderophore

transporter orthologues *SIT1_1948*, *SIT1_2241* and *SIT1_4097*. Cell wall components,

non-mating *MTL* locus genes such as *PIKA*, and genes involved in iron and siderophore

transport are therefore potential modulators of strain-specific differences in virulence.

Typical and atypical hypha formation programmes in basal clade V during AK

yolk-sac infection

To explore drivers of filamentation by the clade V strain, we identified 1,046

DEGs between clade V and each of the other clades (I-IV), comprising 601 up-regulated

and 526 down-regulated genes across all comparisons. We then identified a set of 57

genes shared across all comparisons for *in vivo* conditions only, since we did observe

filamentation in nutrient-rich YPD. 22 genes were up-regulated at 24 HPI, 7 at 48 HPI,

and 18 at both time-points (**Fig. 4C**). The majority of up-regulated genes were metabolic

(n = 13), including two cytochrome P450 *ERG5* orthologues, though *ERG11* was down-

regulated. Transmembrane transporters for sugar (including two *HGT13* orthologues),

amino acids and siderophores (*SIT1_1487*) were also up-regulated (n = 10). Cell wall

genes up-regulated in filamentous clade V (n = 8) included the novel adhesin *SCF1*,

GPI-anchored genes *PGA31* and *IFF4*, and orthologues of classical *C. albicans* hyphal

adhesins *ALS3* and *RBR3*. We identified differentially expressed transcription factors

and related regulators, such as *HAP41* and the pheromone receptor *STE2*, which could

play a governing role on these responses. Given the minor differences in numbers of

DEGs identified when using clade-specific reference genomes, we also compared the

transcript levels for each gene in this set with annotations to identify a highly strict

shortlist of inter-clade difference. Using this method, the only significant differences

were for up-regulation of *MED7* and *SPS19* (at 24 HPI) and two *HGT13* orthologues

and *RNH70* (at both time-points, **Fig. S6**).

We also examined significant DEGs between clade V and at least one other

clade using mean-log fold change (**Fig. 4D**). Several genes that play critical roles in

C. albicans hypha formation were highly up-regulated, such as transcription factor

UME6 and hypha-specific G1 cyclin-dependent protein serine/threonine kinase *HGC1*.

Enrichment analysis indicated up-regulated gene sets involving CFEM domain,

cytochrome P450/oxidoreductase activity, hypha-regulated cell wall GPI-anchored

proteins and siderophore transport, particularly at 24 HPI and comparing clade V to

clade II (**Fig. 4D**). Significantly up-regulated iron-related transport genes in clade V

included *FRE3*, *CFL4*, *FRP1*, *SIT1_1519*, *SIT1_1487*, *SIT1_4097*, *SIT1_1542*, and

SIT1_1499. These findings are consistent with the preservation of a major filamentation

programme in *C. auris*, demonstrated in the basal clade V, with unique characteristics

such as expression of *SCF1* and the up-regulation of several members of the expanded

family of siderophore transmembrane transporters.

Xenosiderophore transporters are a *Metschnikowiaceae*-specific expanded gene family with varied expression patterns in *C. auris* infection

Given the prominence of *SIT1* orthologues across the transcriptomic analyses of *C. auris* during *in vivo* gene expression, we set out to characterise the expanded gene family across orthogroups, including outgroup species *C. haemulonii* and *C. albicans*. **Fig. 5A** shows the predicted structure of *Sit1_1542* (AlphaFold 3). Blastp searches identified proteins with sequence similarity across seven orthologue clusters, of which two main groupings of transporters with 14 transmembrane domains (TMRs) emerged (**Fig. 5B**). A larger branch was annotated with GO term, KEGG pathway and PFAM domains relating to siderophore import (including 12 genes in clade I), and a smaller branch was annotated with GO terms relating to haem transmembrane import (including 5 genes in clade I). The majority of orthologues retained 14 TMRs, and few had evidence of additional annotations except for the trichothecene mycotoxin efflux pump PFAM domain across multiple groupings. The orthologue *SIT1_1547* contained 12 TMRs and was up-regulated with *SIT1_1548* containing 2 TMRs, which was not identified as a *SIT1* orthologue in our pipeline but has previously been described as such[30,82].

Co-expression analysis, using an unweighted network (Pearson's correlation) with a cut-off of 0.85, demonstrated eight clusters featuring *SIT1* orthologues, one of which included five that were co-expressed (**Fig. 5C**). To understand how our *in vivo* expression of *C. auris* compared to other experimental conditions previously reported, we measured gene expression across publicly available RNA-seq datasets, identifying 35 datasets for which we were able to ascertain gene expression using our pipeline, but none of which included an in-host setting (**Table S4**). Gene expression patterns within the set of single copy orthologues clustered within both principal components analysis of experimental unit variance (**Fig. S3E**) and hierarchical clustering of $\log_{10}$ variance per gene (**Fig. 3F**). We used this data to construct an additional unweighted co-expression network as above for *SIT1* orthologues among single copy orthologue genes, identifying

two clusters only, featuring *SIT1_2110* and *SIT1_1519* (**Fig. 5D**). Given the difference in

co-expression patterns between the responses identified in the AK marine host across

this study, and a broader meta-analysis of available RNA-seq data, these results

highlight a potentially unique influence of a marine larval host environment on

transcription programmes and the need to explore in-host time series data with

weighted co-expression networks in the future.

To explore the evolutionary history of *SIT1*-related genes in fungi, we performed

Blastp searches for orthologues of *C. albicans SIT1* using an e-value cut-off of $1e^{-20}$

across 226 fungal species' reference genomes totalling 304 genomes. Separately, 28

Oomycota genomes were found to contain zero orthologues. We identified 1,689 hits,

which clustered into two branches: a smaller branch (n = 397) annotated for siderophore

transport including the 12 *C. auris* groupings which we have designated

"xenosiderophore transporter candidates" (*XTC*), and a larger branch (n = 1,292)

featuring genes predicted (by GO term) to encode haem transporters, including the 5

C. auris groupings, which we have additionally designated "haem transport-related"

(*HTR*, **Fig. 6A**). There were no putative haem transporters or members of the larger

branch for *C. albicans*, which only contained the single *SIT1* gene. We calculated the

mean number of orthogroups per genome per species (**Fig. 6B**). This showed the most

prominent expansion of the *XTC* branch in the *Metschnikowiaceae* clade, including

C. auris (12 orthologues) and even more so in *C. haemulonii*, with 15 orthologues. No

other fungal species or group of species contained so many. *XTC* orthologues were not

present in any members of the basal blasocladiomycota, chytridiomycota and

mucoromycota, or in several ascomycota (e.g. *Pichia kudriavzevii*,

Schizosaccharomyces pombe and *Pyricularia oryzae*) and basidiomycota (e.g. *Ustilago*

maydis). Common human fungal pathogens such as *Aspergillus fumigatus*, *C. albicans*

and *Nakaseomyces glabratus* contained only one *XTC* orthologue (**Fig. 6B**). The *HTR*

clade, by contrast, was more prominent across fungal phyla, with as many as nine in

Mucor species, 16 in *A. flavus* and 22 in *F. solani*. We therefore propose a naming of

these two expanded gene families as *XTC1-12* and *HTR1-5* in *C. auris*.

We note the absence of *XTC5* in clade V, *XTC10* in clade IV, and in addition to
these, *XTC4*, *XTC9* and *XTC11* in clade II (**Fig. 6C**). Interestingly, clade II, which
exhibits the greatest degree of *XTC* gene loss, is also considered to be the least virulent
strain. It is also noteworthy that the AK embryo expressed *HMOX* genes during clade IV
infection, while clade IV *C. auris* up-regulated *HTR1*, *HTR2* and *HTR5*. This suggests a
specific host-pathogen interaction axis in nutritional immunity related to haem
scavenging. No other clades of infection were associated with such iron-related
functional enrichment in the host, and no other clades expressed more than one *HTR*
gene (except *HTR1* and *HTR5* in clade III at 48 HPI). Clade II down-regulated *HTR1* in-
host. In summary, *XTC1*, *XTC2*, *XTC3*, *XTC4*, *XTC6*, *XTC7*, *XTC11* and *XTC12* were
up-regulated while *XTC8*, *XTC9* and *XTC12* were down-regulated *in vivo* in at least one
clade, while up-regulation of *XTC3* and *XTC7* was significant across all clades.
Additionally, the more virulent clades were associated with up-regulation of *XTC4,*
XTC5, and *XTC11*. The filamentous clade V up-regulated *XTC7* across comparisons
with all other clades, and up-regulated *XTC1, XTC4, XTC8* and *XTC9* compared to at
least one other clade.
491
492   **Discussion**
493
We have desribed a model for comparative functional genomics of *C. auris* in-
host survival using AK yolk-sac microinjections at a clinically relevant temperature. All
five clades demonstrated similar lethality, with a minor but significant increase in
pathogenicity for clades I and IV, consistent with clinical and laboratory findings[29]. In-
host growth was similar across all clades and high enough to indicate a large enough
in vivo fungal burden for robust bulk transcription assessment across the species.
Lower CFUs in clade V could be explained by increased aggregation of filamentous
cells. The contained nature of the yolk-sac provides a straightforward experimental unit
of host tissue without the challenges of pathogen RNA isolation such as locating foci of
infection in larger animals. Additional time-points at could be an informative future area

of study to develop fine-resolution strain-specific transcription networks. Thus, an AK

yolk-sac microinjection model has the potential to fill a substantial need for *in vivo* gene

expression studies in general.

We investigated the host transcriptional response to *C. auris* infection. The most

highly up-regulated gene family included two heme oxygenase (*HMOX*) genes. *HMOX*

genes have been shown to mask iron from invading pathogens as a feature of anti-

Candida nutritional immunity[83], so presumably have a similar function against *C. auris*.

This was particularly notable given *HMOX* genes were up-regulated in response to

clade IV infection, and the clade IV isolate had a corresponding significant up-regulation

of 3 haem transport-related *HTR* genes. In addition to nutritional immunity genes, we

found up-regulated heat shock and complement cascade genes during infection,

confirming the expected immune response to fungal infection[65,66]. Genes with complex

roles in immunity were up-regulated, such as *NLRP3*, for which *C. auris* may be less

activating than e.g. *C. albicans*[27,84]. It is notable that the yolk-sac model was not

associated with histological or transcriptional evidence of host neutrophil recruitment

and activation, a prominent feature of human blood infection modelling[30]. Reference

genome bias resulting from the available adult gill transcriptome assembly[85] may limit

detection of host gene expression features, so future long and short read assembly from

gene expression data across a range of larval stages could further understanding of AK

immunity.

We discovered a complex pattern of differential expression by *C. auris* during

infection compared with *in vitro* conditions. The lack of a major distinction across time-

points (24 *vs* 48 HPI) indicates that the transcriptional activity of *C. auris* in-host was

highly similar at these stages. Six functional annotation terms (GO, KEGG pathway and

PFAM domain) were significantly enriched among genes that were up-regulated across

all five clades, including sugar and siderophore transporters. The enrichment of sugar

transporters outside of nutrient-rich YPD-broth is unsurprising, such as High-affinity

Glucose Transporter *HGT12*, with evidence of redundancy in *C. albicans*[86–89], and

HGT2, part of an eight-gene "core filamentation response network" in *C. albicans*[90].

More surprising was the down-regulation of putative virulence factors such as Secreted

Aspartyl Protease 3 (*SAP3*), thought to be the most important SAP in *C. auris*, and

required for virulence in a mouse model[26], and Copper Superoxide Dismutase *SOD6*,

which has been shown to be down-regulated in iron starvation[91]. Multi-drug efflux

pumps such as *MDR2* (down-regulated *in vivo*) and *MDR1* (down-regulated *in vivo* and

in clades I and IV *vs* clades II and III) are understood to drive drug resistance through

multiple but unclear mechanisms in *C. auris*[92,93]; in-host down-regulation may indicate a

fitness trade-off for virulence *vs* drug resistance.

The mating type locus (*MTL*) genes have been hypothesised to contribute to

virulence in fungal species including *Cryptococcus neoformans*[94]. Given that *C. auris* is

assumed not to undergo meiotic recombination[95], there is a question of what role *MTL*-

associated genes play in pathogenicity and morphology. Two of the three non-mating

genes (*PAP1* and *PIKA*) in the *MTL* were up-regulated by more virulent strains (I and

IV) compared to the less virulent strains (II and III). Importantly, the two more virulent

clades (I and IV) contain *MTL**a*** rather than *MTLα* (clades II & III)[23], and clades V and VI

contain *MTL**a*** and *MTLα* respectively[12]. Thus, it is unclear if the up-regulation of non-

mating genes is a *MTL* specific transcriptional nuance, or a feature of more highly

virulent *C. auris* clades. The *MTL* also did not feature prominently in the filamentation

response observed in clade V, providing no evidence for an obvious role for the *MTL* in

such morphological changes.

Filamentation is an important pathogenicity trait in fungi, especially *Candida*

species, although its roles have been more poorly described in aquatic infections[96]. We

observed *in vivo* filamentation by a representative strain of *C. auris* clade V, the

basal/ancestral clade[97], accompanied by a genera-typical filamentous gene expression

signature. Specifically, we measured the up-regulation of *HGC1* and *UME6*, which are

two of the most important regulators of filamentation in *C. albicans*, and have recently

been described as drivers of *C. auris* filamentation and biofilm formation[98]. The

expression of adhesin *SCF1* in filamenting clade V is noteworthy because its cationic

adhesion mechanism has been postulated to be consistent with adhesion mechanisms

common to marine organisms including bivalves[21]. The generalist marine-isolated

pathogen *C. haemulonii* has also displayed filamentation competence at lower

temperatures[99], consistent with an ancestral aquatic state.

Human histopathology reports of *C. auris* infection are rarely published and have

not demonstrated classical in-host filamentation, to our knowledge. An *ex vivo* human

epithelial model using a clade I isolate did demonstrate filamentous *C. auris* at 37 °C

after injection into deeper dermis[100]. Clade III *C. auris* has also been observed to form

filamentous/hyphal forms in *Galleria mellonella* injection[60]. Fan and colleagues have

observed filamentous forms of *C. auris* across all four clades, even demonstrating a

higher virulence in the *G. mellonella* model[101], though the phenotype required passage

through a murine host with additional prolonged growth at 25 °C for at least 5 days

before and after injection[19]. Other stimuli have induced filamentation in clade I isolates,

such as hydroxyurea genotoxic stress[102] and Hsp90 inhibition[103], where several up-

regulated genes in *C. auris* filamentation (*SCF1*, *IFF4* and *PGA31*) matched those in

our study. Constitutive filamentous growth has also been demonstrated with a

transposon-mediated disruption of a long non-coding RNA[104], which sit adjacent to

chitin deacetylase *CDA2*, which was down-regulated at 24 HPI in this experiment during

filamentation. These findings indicate that hyphal formation is possible across *C. auris*

clades.

A total of twelve siderophore transporters were up-regulated during infection, and

four were down-regulated, with three family members up-regulated in more virulent

clades and five in the filamentous clade V. Siderophore synthesis may be unlikely in

C. auris, given the lack of evidence in other *Candida* species[105]. As a result, *C. albicans*

Sit1 - which is required for required for virulence[106] - is thought to be a xenosiderophore

transporter for iron-chelators produced by other microbes and specific to the

ferrioxamine type common in iron-poor ocean environments[107,108]. Siderophore

transporters can also act intracellularly for compartments such as the

mitochondrion[109,110], or act as efflux pumps for various molecules[111] including

trichothecene mycotoxins as identified by our functional annotations.

C. auris encodes a large expanded gene family of 17 siderophore transporters

that we named xenosiderophore transporter candidates (*XTC*) 1-12 and haem

transport-related (*HTR*) genes 1-5, the latter sharing sequence similarity with *S. pombe*

low-affinity haem transporter Str3[112]. In other *Candida* species such as *C. albicans*, iron

assimilation is understood to be central for in-host survival[113,114], where deprivation

triggers various pathogenic programmes such as alterations in cell surface beta-

glucan[115]. Haem is acquired by *C. albicans* through vesicular endocytosis and CFEM-

domain containing proteins[116,117] and does not contain *HTR* genes, suggesting a

possible loss of transmembrane haem transporters. The ubiquity of *HTR* across the

fungal kingdom and lack of *XTC* genes in the three basal fungal phyla suggest that

SIT1/HTC could have arisen from a duplication of an *HTR*-related gene in the Dikarya

(the ascomycota and basidiomycota) with subsequent independent loss of both *XTC*

and *HTC* genes in multiple fungal lineages. Our identification of up-regulated *XTC*

transport during infection corroborates the findings of up-regulated *XTC2A-B*

(*B9J08_1547-8*) in an *ex vivo* blood model of *C. auris* infection[30]. These findings

together suggest the importance of iron during infection, which could be targeted in the

clinic, e.g. via low-toxicity pyrvinium pamoate[118].

We show for the first time an *in vivo* transcriptional profile for the global public

health threat and critical priority WHO human fungal pathogen *C. auris*. Representative

strains from all five major clades of the recent simultaneous emergences highlighted

strain-specific differences in morphology, virulence and gene expression. This study

highlights the advantages and potential of a thermotolerant yolk-sac injection model,

adding to the debate around the putative marine origins of *C. auris*, and the emergence

of distinct clades. We demonstrate natural filamentation by clade V and virulence

advantages of clades I and IV, and comprehensively describe the associated gene

expression. We indicate the importance of the *XTC* and *HTR* gene clusters as

recommended targets for investigation of fungal biology, pathogenicity and treatment.

Our findings shed new light on the emergence and evolution of *C. auris* from a marine

yeast ancestor. Our analyses reveal new potential drug targets that enable *C. auris* to

survive in aquatic hosts, which may have a similar function during human infection, with

implications for patient care and public health control.

Methods

Ethical Statement: All animal experiments were performed under Project

Licence PP4402521 in compliance with University of Exeter Animal Welfare Ethical

Review Board regulations and MRC Center for Medical Mycology Project Review

Board, and are reported in line with "Animal Research: Reporting *in vivo* Experiments"

(ARRIVE 2.0) Guidelines[119].

***Aphanius dispar* Care, Monitoring, Husbandry and Procedures:** Wild-type

adult AK were kept at the Aquatic Resources Facility, Exeter, in two 28 °C incubation

tanks, with 12-hour day and night light cycles and artificial enrichment foliage. Embryos

were collected using three plastic netted collection chambers per tank with transfer of all

enrichment into tanks, and placed overnight from 16:00 to 09:15 in dark conditions.

Chambers were then drained, cleaned and strained with 35 parts per thousand artificial

seawater (ASW) to isolate all embryos. Subsequent ASW was sterile suction filtered *via*

0.2 µm pore Nalgene Rapid-Flow (Thermo Scientific, UK). Embryos were inspected for

viability in terms of an intact blastocyst without discolouration by stereo dissection

microscope (Optech, UK) and incubated at 30 °C (Memmert, Germany) for 72 h with

day and night cycles with daily ASW changes and removal of dead, unhealthy or

underdeveloped embryos, with a maximum of 30 embryos per standard size 100 mm x

15 mm petri dish (Greiner Bio-One, Austria). Excluded embryos were killed or disposed

of either by submission in Virkon solution (Sigma-Aldrich, UK) or freezing at -80 °C.

Experimental Procedures: Injection needles were prepared using borosilicate

glass capillaries (Harvard apparatus, USA) and a heated pulling system (heater level >

62, PC-10, Narishige, UK). We used tweezers to snap the tip to an estimated 1 mm

graticule-measured aperture of 10-20 μm, and sterilised needles before use with

253.7 nm UV light over 15 minutes (Environmental Validation Solutions, UK). Injection

bays were made with microwaved 1% agarose in 1 g (Sigma, USA) per 100 mL ASW

for up to 24 embryos in 100 mm petri dishes using custom-made molds in an inverted

60 mm petri dish. Embryos were randomly chosen from each dish and the sequence of

injection for each biological repeat was randomly allocated by computer. For

microinjection, *C. auris* yeasts were counted by haemocytometer and added to 25 μL

sterile filtered phenol red solution 0.5% (Sigma, UK) to a concentration of 0.5×10^8 per

mL in 100 μL. Injection needles were loaded with 5 μL injection solution using 20 μL

Microloader EpT.I.P.S. (Eppendorf, UK) immediately before injection of up to 30

embryos before re-loading. Injections were performed with 400-500 hPa injection

pressure and 7-40 hPa compensation pressure over 100-200 ms with the FemtoJet 4i

(Eppendorf UK) using a micromanipulator (World Precision Instruments, UK) and a

Microtec HM-3 stereo dissection microscope (Optech, UK). In order to deliver a dose of

approx. 500 yeasts in 10 nL solution, an injection bolus was administered to an

estimated diameter of 0.2 – 0.3 mm (10-15% of total embryo diameter) *via* a single

chorion puncture. Any embryos that failed to meet these criteria for injection were

excluded. Control *in vitro* inoculations were microinjected as above into 1 mL YPD in

1.5 mL Eppendorf tubes before incubation without shaking at 37 °C for 24 h

contemporaneously with embryos as below.

Study Design and Sample Sizes: We used an unblinded prospective virulence

study comparing six groups of AK embryo yolk-sac microinjection, including five

different clades of *C. auris*, and one control sham injection as the negative control. We

used >= 23 embryos per condition based on our power calculation of $\sigma = 0.25$,

significance = 0.05, power = 0.9, and δ effect size of 25% mortality (two-sided t-test).

After excluding embryos that were unhealthy or did not complete successful

microinjection, we included n = 10-13 embryos per group per experiment (total n = 34-

37 embryos per group). Each embryo was considered to be an experimental unit

(n = 70-72 embryos per experiment, total n = 213 embryos). For estimation of *C. auris*

in-host growth by homogenisation and plating of colony forming units, three embryos

per condition per time-point across two (0 and 24 HPI) or three (48 HPI) experiments

were used (total n = 126). For time lapse microscopy demonstration of *A. dispar* yolk-

sac collapse and death, representative image sets were chosen from a pilot experiment

using a reduced dosage (1/10) of *C. auris* (n = 4). For histological demonstration of

C. auris morphology, one embryo per condition at 24 HPI and 48 HPI each were used in

one experiment, and seven embryos per condition plus three embryos without injection

in a second experiment (n = 57). RNA was extracted from three embryos per condition

per time-point (0, 24 and 48 HPI, n = 54) plus three embryos per time-point without any

injection (n = 9, total n = 63) as a control for the impact of microinjection.

C. auris Strains and Growth: We selected five *C. auris* strains to represent

each major clade (**Table S1**). Culture stocks were kept in -80 °C in Yeast extract

Peptone Dextrose (YPD) broth (Sigma-Aldrich, UK) containing 25% glycerol

(ThermoFisher, UK) and plated onto YPD-agar (Sigma-Aldrich) for 48 h before

innoculation and overnight growth in 10 mL YPD broth at 30 °C in a shaking incubator at

200 rpm (Multitron, Infors HT, UK). Five mL from each overnight culture was then

centrifuged at 5000 g for 2 minutes before re-suspending and triple washing in 1 mL

autoclaved deionised water (DIW, Milli-Q, Merck, USA) to prevent aggregation[22] then

re-suspended in 200 µL DIW. Decontamination of surfaces was conducted with both

mopping and cleaning wipes in 70% Ethanol (Sigma-Aldrich) or 10% Chemgene HLD4L

(Medimark Scientific, UK) as per manufacturer's instructions. For growth curves,

C. auris strains were grown overnight in YPD in 4 replicates. Overnight cultures were

inoculated into YPD to a final OD 600 of 0.02 in a 96-well plate. OD was read over 25

hourly cycles at 37 °C with 300 s shaking at 2.5 mm amplitude on an Infinite 200 PRO

plate reader (Tecan, Switzerland).

Outcome measurement: individual embryos were kept in 1 mL sterile filtered

ASW per embryo in a 48 well flat bottom plate (Corning, USA) sealed with Parafilm "M"

(Bemis Company, USA) at 37 °C with passive humidification using a 500 mL open glass

beaker of water. Embryos were monitored for presence of heart rate at 24-hourly

intervals up to 7 days by stereo dissection microscope. Initial time of injection was

subtracted from each time-point to minimise bias between groups. Kaplan-Meier

survival curves were plotted with Log-Rank testing and Benjamini-Hochberg (BH)

correction[120] using survival v.3.5.8 and survminer v.0.4.9 packages in R v.4.4.0. Colony

forming units (CFUs) were calculated by homogenisation of embryos at 0, 24, and

48 HPI by autoclaved plastic micropestle in a 1.5 mL conical tube (Eppendorf, UK) in

100 µL 10,000 U/mL penicillin/streptomycin (P/S) solution (Gibco, UK). Dilutions were

made with P/S solution and DIW in 1/10-10,000 and counts were adjusted according to

dilution after plating onto YPD agar plates and read after 72 h incubation at 30 °C.

Statistical testing between CFU numbers was performed with Wilcoxon testing and BH

correction.

Imaging: For direct microscopy, AK were kept in 300 µL ASW containing 0.04%

Tricaine anaesthesia (Sigma-Aldrich, UK) in an optical adhesive and parafilm-sealed

96-well plate. These were then monitored for heart beat by 5 bright field 2X or 4X

images at pre-focused z slices taken every 2 h by Acquifer microscopy (Bruker,

Germany) over 72 h. All images were processed in ImageJ 1.53a (NIH, USA).

Sectioning and staining was performed by Microtechnical Services, Exeter, UK, with

fixation of embryos in 1 mL 10% neutral-buffered formalin (Sigma-Aldrich, UK) for >

24 h, followed by immersion in 4% phenol 10 mL glycerine BP made up to 100 mL with

distilled water[121], sectioned at 4 µm without orientation, and stained with Harris

Haematoxylin & 1% Alcoholic Eosin on Leica Autostainer. H&E stained histological

sections were imaged on an Olympus IX83 microscope coupled with an Olympus DP23

colour camera using an Olympus UPlanXApo x40/0.95 objective controlled by cellSens

software v.3.2. Brightfield microscopy was performed with an EVOS M5000 microscope

(ThermoFisher, UK). Microscopy of YPD-agar grown colonies was performed without

staining in DIW.

RNA Extraction and Sequencing: Total RNA was extracted using the

Monarch® Total RNA Miniprep Kit (NEB, USA). Embryos were flash frozen in liquid

nitrogen, re-suspended in DNA/RNA Protection Reagent and disrupted with an

autoclaved plastic micropestle. Bead beating was performed in a FastPrep 24

homogeniser (MP Biomedicals) with 0.5 mm zirconia/silica beads for 8 cycles of 35 s at

6 m/s plus 45 s on ice. Lysate was recovered and incubated for 5 min. at 55 °C with

proteinase K before addition of lysis buffer and RNA purification following

manufacturer's instructions. Total RNA samples were quantified using the Qubit 4.0

Fluorometer (Invitrogen, USA) and integrity was checked with the Tapestation 4200

(Agilent, USA). Library preparation and sequencing were performed by the Exeter

Sequencing Service. Samples were normalised and prepared using the NEBNext Ultra

II Directional RNA Library Prep - Poly(A) mRNA Magnetic Isolation Module following the

standard protocol. Libraries were sequenced on a Novaseq 6000 sequencer with 150bp

paired-end reads. Reads were trimmed with fastp[122] v.0.23.1 to remove reads < 75

bases and trim based from the 3' end with q-score < 22. Quality control was performed

with FastQC (https://github.com/s-andrews/FastQC) v.0.11.9 and MultiQC[123] v.1.6.

Differential Gene Expression: We aligned sequences (x = 52.9 m reads per

replicate, range 41.1-112.2 m) using the Trinity pipeline[124] v.2.15.1 to the *C. auris* clade

I B8441 reference genome (GCA_002759435.2)[23] downloaded from GenBank (mean

aligned reads in infected embryos 3.83 m, 0.81-13.1%). We further aligned clade-

specific reference genomes for each strain (**Table S1**), including clade II

(GCA_003013715.2)[4], III (GCA_002775015.1)[13], IV (GCA_008275145.1)[13] and V

(GCA_016809505.1)[97,125,126]. We aligned conditions containing *A. dispar* to the *de novo*

GJEY01 adult gill transcriptome assembly[85] downloaded from GenBank, for which we

detected expression over 27,481 potential genes. Alignment was performed with

Bowtie2[127] v.2.5.1, transcript estimation with RNA Seq by Expectation Maximisation

(RSEM)[128] v.1.3.3, differential expression with EdgeR[129] v.3.40.0, Samtools[130] v.1.17

and limma[131] v.3.54.0 using a false discovery rate (FDR) p-value cut-off of 0.001 and a

log fold change cut-off of +/-2.0. Principal components were calculated and plotted

using the prcomp function in Rv4.4.0 and ggfortify v.0.4.17 R package for fragments per

kilobase per million, and correlations plots from counts per million as per the default

output of the Trinity pipeline. Log fold change was plotted with ComplexHeatmap[132].

Annotation and Enrichment: Gene ontology terms[133,134] were assigned using

DIAMOND[135] v.2.1.8 *via* the DIAMOND2GO pipeline[136] for each of the *C. auris*

reference genomes. KEGG terms[137] were assigned using the BlastKOALA online

platform using the eukaryote database[138]. Protein FAMily (PFAM) domains[139] were

identified using the PFAM-A.hmm database

(https://ftp.ebi.ac.uk/pub/databases/Pfam/current_release/Pfam-A.full.gz) and HMMER

and HMM Scan[140] with a cut-off of $1e^{-5}$. Further domain-specific identification was

performed with SignalP 5.0[141], DeepTMHMM 1.0[142] and NetGPI 1.1[143]. Enrichment

analysis for DEG subsets was calculated using the Fisher's Exact test, adjusted using

the Benjamini-Hochberg method per set of up-regulated or down-regulated features in

each comparison as a conservative estimate of significant gene set enrichment, with a

false discovery rate cutoff of p=0.001. Annotation tables were checked with additional

information from available literature, supplemented by details and web-scraped

orthologue names from *Candida* Genome Database[82] and Blastp searches[144].

Annotations for the GJEY01 genome were also performed using Diamond2GO,

BlastKoala and HMMScan as above and checked against *A. dispar* annotations and

gene names obtained directly from the GJEY01 transcriptome makers[85]. For plotting,

lists of GO terms were identified (and for *A. dispar* enrichment, reduced by removal of

redundant terms) by REVIGO[145].

Orthologue Assignment: To identify accessory genes across *C. auris* clades,

we used the Synima pipeline[146] with Orthofinder v2.5.5[147] with additional reference

genomes for *C. albicans* (SC5314_GCA_000182965.3) and *C. haemulonii*

(B11899_ASM1933202v1). We also incorporated a very recent version of the *C. auris*

B8441 assembly (v3) to arrange contigs and illustrate re-arrangements, with broad

syntenic concordance; it contains 5 more genes than v2, for which 27 (0.5%) and 16

(0.3%) genes were unique to each genome, respectively. Multiple alignment of single

copy orthologues as defined by Orthofinder was performed with Muscle v5.1[148] and a

phylogeny was made with FastTree v2.1.11[149] using default settings.

RNA-Seq Meta-Analysis Variance: NCBI SRA was searched for

"*Candida auris*" on 16th April 2024 and downloaded as metadata *via* the "Send to: Run

Selector" function. Each project accession number for runs containing RNA was

searched against available published literature to identify publications, noting subject of

study, clades/strains, growth media, comparative conditions, temperature and time

scales used. Calculation of FPKM was performed and analysed as above using the set

of single copy orthologues to demonstrate variance via PCA with ggfortify and prcomp

packages in R and to calculate an unweighted co-expression network for *SIT1* genes

with and without the additional data from the RNA-Seq meta-analysis, using Pearson's

correlation with a correlation cut-off of 0.85.

SIT1 **Analysis:** The protein transcript for *C. albicans SIT1* was Blastp searched

against the draft pangenome database as above. We visualised the structure using

Alphafold v3[150] and ChimeraX v1.8[151]. Clade I-V *C. auris* and *C. haemulonii SIT1* genes

were aligned using Muscle as above before using RAXML-NG v1.2.2 with 1000

bootstraps and the LG4X model[152], which was visualised using ggtree v3.12.0 and

gheatmap. For panfungal *SIT1* analysis, we used the fungidb portal for Blastp using the

C. albicans SIT1 against all available fungal data (with no hits within oomyceta),

including 274 reference genomes representing 197 species and a cut-off of $1e^{-20}$.

Annotations for each protein transcript were performed as above to obtain GO terms,

KEGG pathways, PFAM domains, GPI-anchors, SignalP domains, and transmembrane

domains, while a phylogeny for *SIT1* similar sequences across all fungal references was

constructed as above following multiple alignment with Clustal Omega v1.2.4[153]

Data Availability: Raw reads have been deposited via NCBI/GEO *via* accession

GSE277854. All datasets and code required to make figures are available via Github

(https://github.com/hughgifford/Arabian_Killifish_C_auris_2024).

References

1.   Meis, J. F. & Chowdhary, A. *Candida auris*: A global fungal public health threat.

The Lancet Infectious Diseases **18**, 1298–1299 (2018).

2.   WHO. World Health Organisation (WHO) fungal priority pathogens list to guide

research, development and public health action. (2022)

doi:https://www.who.int/publications/i/item/9789240060241.

3.   Lee, W. G. *et al.* First three reported cases of nosocomial fungemia caused by

Candida auris. *Journal of Clinical Microbiology* **49**, 4 (2011).

4.   Satoh, K. *et al. Candida auris* sp. Nov., a novel ascomycetous yeast isolated

from the external ear canal of an inpatient in a Japanese hospital. *Microbiology and*

Immunology **53**, 41–44 (2009).

5.   Lyman, M. Notes from the field: Transmission of pan-resistant and echinocandin-

resistant *Candida auris* in health care facilities - Texas and the District of Columbia,

January-April 2021. *Morbidity and Mortality Weekly Report* **70**, (2021).

6.   Eyre, D. W. *et al.* A *Candida auris* outbreak and its control in an intensive care

setting. *New England Journal of Medicine* **379**, 1322–1331 (2018).

7.   Taori, S. K. *et al. Candida auris* outbreak: Mortality, interventions and cost of

sustaining control. *Journal of Infection* **79**, 601–611 (2019).

8.   Shuping, L. *et al.* High prevalence of *Candida auris* colonization during protracted
neonatal unit outbreak, South Africa. *Emerging Infectious Diseases* (2023)
doi:10.3201/eid2909.230393.

9.   Adam, R. D. *et al.* Analysis of *Candida auris* fungemia at a single facility in
Kenya. *International Journal of Infectious Diseases* **85**, 182–187 (2019).

10.   Prayag, P. S., Patwardhan, S., Panchakshari, S., Rajhans, P. A. & Prayag, A.
The dominance of *Candida auris*: A single-center experience of 79 episodes of
candidemia from Western India. *Indian Journal of Critical Care Medicine* **26**, 560–563
(2022).

11.   Khan, T. *et al.* Emergence of the novel sixth *Candida auris* Clade VI in
Bangladesh. *Microbiology Spectrum* **12**, e03540–23 (2024).

12.   Suphavilai, C. *et al.* Detection and characterisation of a sixth *Candida auris* clade
in Singapore: A genomic and phenotypic study. *Lancet Microbe* **0**, (2024).

13.   Lockhart, S. R. *et al.* Simultaneous emergence of multidrug-resistant *Candida*
auris on 3 continents confirmed by whole-genome sequencing and epidemiological
analyses. *Clinical Infectious Diseases* **64**, 134–140 (2017).

14.   Gifford, H., Rhodes, J. & Farrer, R. A. The diverse genomes of *Candida auris*.
The Lancet Microbe **0**, (2024).

15.   Chen, J. *et al.* Is the superbug fungus really so scary? A systematic review and
meta-analysis of global epidemiology and mortality of *Candida auris*. *BMC Infectious*
Diseases **20**, 827 (2020).

16.   Mayer, F. L., Wilson, D. & Hube, B. *Candida albicans* pathogenicity mechanisms.
Virulence **4**, 119–128 (2013).

17.   Phan, Q. T. *et al.* Als3 is a *Candida albicans* invasin that binds to cadherins and
induces endocytosis by host cells. *PLOS Biology* **5**, e64 (2007).

18.   Moyes, D. L. *et al.* Candidalysin is a fungal peptide toxin critical for mucosal

infection. *Nature* **532**, 64–68 (2016).

19.   Yue, H. *et al.* Filamentation in *Candida auris*, an emerging fungal pathogen of

humans: Passage through the mammalian body induces a heritable phenotypic switch.

Emerging Microbes & Infections **7**, 1–13 (2018).

20.   Chadwick, C., Jesus, M. D., Ginty, F. & Martinez, J. S. Pathobiology of *Candida*

auris infection analyzed by multiplexed imaging and single cell analysis. *PLOS ONE* **19**,

e0293011 (2024).

21.   Santana, D. J. *et al.* A *Candida auris*-specific adhesin, Scf1, governs surface

association, colonization, and virulence. *Science* **381**, 1461–1467 (2023).

22.   Pelletier, C., Shaw, S., Alsayegh, S., Brown, A. J. P. & Lorenz, A. *Candida auris*

undergoes adhesin-dependent and -independent cellular aggregation. *PLOS Pathogens*

20, e1012076 (2024).

23.   Muñoz, J. F. *et al.* Genomic insights into multidrug-resistance, mating and

virulence in *Candida auris* and related emerging species. *Nature Communications* **9**,

5346 (2018).

24.   Richardson, J. P. *et al.* Candidalysins are a new family of cytolytic fungal peptide

toxins. *mBio* **13**, e03510–21 (2022).

25.   Horton, M. V. *et al.* *Candida auris* cell wall mannosylation contributes to

neutrophil evasion through pathways divergent from *Candida albicans* and *Candida*

glabrata. *mSphere* **6**, e00406–21 (2021).

26.   Kim, J.-S., Lee, K.-T. & Bahn, Y.-S. Secreted aspartyl protease 3 regulated by

the Ras/cAMP/PKA pathway promotes the virulence of *Candida auris*. *Frontiers in*

Cellular and Infection Microbiology **13**, 1257897 (2023).

27.   Weerasinghe, H. *et al.* *Candida auris* uses metabolic strategies to escape and kill

macrophages while avoiding robust activation of the NLRP3 inflammasome response.

Cell Reports **42**, 112522 (2023).

28.   Byun, S. A. *et al.* Virulence traits and azole resistance in Korean *Candida auris*

Isolates. *Journal of Fungi* **9**, 979 (2023).

29.   Santana, D. J., Zhao, G. & O'Meara, T. R. The many faces of *Candida auris*:

Phenotypic and strain variation in an emerging pathogen. *PLOS Pathogens* **20**,

e1012011 (2024).

30.   Allert, S. *et al.* From environmental adaptation to host survival: Attributes that

mediate pathogenicity of *Candida auris*. *Virulence* **13**, 191–214 (2022).

31.   Fayed, B. *et al.* Transcriptome analysis of human dermal cells infected with

Candida auris identified unique pathogenesis/defensive mechanisms particularly

ferroptosis. *Mycopathologia* **189**, 65 (2024).

32.   Yang, B. *et al.* A correlative study of the genomic underpinning of virulence traits

and drug tolerance of *Candida auris*. *Infection and Immunity* **0**, e00103–24 (2024).

33.   Bruno, M. *et al.* Transcriptional and functional insights into the host immune

response against the emerging fungal pathogen *Candida auris*. *Nature Microbiology* **5**,

1516–1531 (2020).

34.   Wang, Y. *et al.* Innate immune responses against the fungal pathogen *Candida*

auris. *Nature Communications* **13**, 3553 (2022).

35.   Chybowska, A. D., Childers, D. S. & Farrer, R. A. Nine things genomics can tell

us about *Candida auris*. *Frontiers in Genetics* **11**, 351 (2020).

36.   Wang, X. *et al.* The first isolate of *Candida auris* in China: Clinical and biological

aspects. *Emerging Microbes & Infections* **7**, 1–9 (2018).

37.   Fan, S., Li, C., Bing, J., Huang, G. & Du, H. Discovery of the diploid form of the

emerging fungal pathogen *Candida auris*. *ACS Infectious Diseases* **6**, 2641–2646

(2020).

38.   Fan, S. *et al.* A biological and genomic comparison of a drug-resistant and a

drug-susceptible strain of *Candida auris* isolated from Beijing, China. *Virulence* **12**,

1388–1399 (2021).

39.   Wurster, S., Albert, N. D. & Kontoyiannis, D. P. *Candida auris* bloodstream

infection induces upregulation of the PD-1/PD-L1 immune checkpoint pathway in an

immunocompetent mouse model. *mSphere* **7**, e00817–21 (2022).

40.   Bing, J. *et al.* Rapid evolution of an adaptive multicellular morphology of *Candida

auris* during systemic infection. *Nature Communications* **15**, 2381 (2024).

41.   Ben-Ami, R. *et al.* Multidrug-resistant *Candida haemulonii* and *C. auris*, Tel Aviv,

Israel. *Emerging Infectious Diseases* **23**, (2017).

42.   Hager, C. L., Larkin, E. L., Long, L. A. & Ghannoum, M. A. Evaluation of the

efficacy of rezafungin, a novel echinocandin, in the treatment of disseminated *Candida

auris* infection using an immunocompromised mouse model. *Journal of Antimicrobial

Chemotherapy* **73**, 2085–2088 (2018).

43.   Zhao, M. *et al. In Vivo* Pharmacokinetics and pharmacodynamics of APX001

against *Candida* spp. In a neutropenic disseminated candidiasis mouse model.

Antimicrobial Agents and Chemotherapy **62**, e02542–17 (2018).

44.   Singh, S. *et al.* The NDV-3A vaccine protects mice from multidrug resistant

Candida auris infection. *PLOS Pathogens* **15**, e1007460 (2019).

45.   Zhang, F. *et al.* A marine microbiome antifungal targets urgent-threat drug-

resistant fungi. *Science* **370**, 974–978 (2020).

46.   Muñoz, J. E. *et al.* Pathogenicity levels of Colombian strains of *Candida auris*
and Brazilian strains of *Candida haemulonii* species complex in both murine and*Galleria*
mellonella experimental models. *Journal of Fungi* **6**, 104 (2020).

47.   Abe, M. *et al.* Potency of gastrointestinal colonization and virulence of *Candida*
auris in a murine endogenous candidiasis. *PLoS One* **15**, e0243223 (2020).

48.   Forgács, L. *et al.* Comparison of in vivo pathogenicity of four *Candida auris*
clades in a neutropenic bloodstream infection murine model. *Emerging Microbes &*
Infections **9**, 1160–1169 (2020).

49.   Nagy, F. *et al.* In vitro and in vivo interaction of caspofungin with isavuconazole
against *Candida auris* planktonic cells and biofilms. *Medical Mycology* **59**, 1015–1023
(2021).

50.   Forgács, L. *et al.* In vivo efficacy of amphotericin B against four *Candida auris*
Clades. *Journal of Fungi (Basel, Switzerland)* **8**, 499 (2022).

51.   Singh, S. *et al.* Protective efficacy of anti-Hyr1p monoclonal antibody against
systemic candidiasis due to multi-drug-resistant *Candida auris*. *Journal of Fungi* **9**, 103
(2023).

52.   Salama, E. A. *et al.* Lansoprazole interferes with fungal respiration and acts
synergistically with amphotericin B against multidrug-resistant *Candida auris*. *Emerging*
Microbes & Infections **13**, 2322649 (2024).

53.   Torres, S. R. *et al.* Impact of *Candida auris* infection in a neutropenic murine
model. *Antimicrobial Agents and Chemotherapy* **64**, e01625–19 (2020).

54.   Xin, H., Mohiuddin, F., Tran, J., Adams, A. & Eberle, K. Experimental mouse
models of disseminated *Candida auris* infection. *mSphere* **4**, e00339–19 (2019).

55.   Rosario-Colon, J., Eberle, K., Adams, A., Courville, E. & Xin, H. *Candida* cell-
surface-specific monoclonal antibodies protect mice against *Candida auris* invasive
infection. *International Journal of Molecular Sciences* **22**, 6162 (2021).

56.   Torres, S. R. *et al.* Assessment of environmental and occupational exposure

while working with multidrug resistant (MDR) fungus *Candida auris* in an animal facility.

Journal of Occupational and Environmental Hygiene **16**, 507–518 (2019).

57.   Borman, A. M. Of mice and men and larvae: *Galleria mellonella* to model the

early host-pathogen interactions after fungal infection. *Virulence* **9**, 9–12 (2017).

58.   Romera, D. *et al.* The *Galleria mellonella* infection model as a system to

investigate the virulence of *Candida auris* strains. *Pathogens and Disease* **78**, ftaa067

(2020).

59.   García-Carnero, L. C. *et al.* Early virulence predictors during the *Candida*

species–*Galleria mellonella* interaction. *Journal of Fungi* **6**, 152 (2020).

60.   Garcia-Bustos, V. *et al.* Characterization of the differential pathogenicity of

Candida auris in a *Galleria mellonella* infection model. *Microbiology Spectrum* **9**,

e00013–21 (2021).

61.   Carvajal, S. K. *et al.* Pathogenicity assessment of Colombian strains of *Candida*

auris in the *Galleria mellonella* invertebrate model. *Journal of Fungi* **7**, 401 (2021).

62.   Hernando-Ortiz, A. *et al.* Virulence of *Candida auris* from different clinical origins

in *Caenorhabditis elegans* and *Galleria mellonella* host models. *Virulence* **12**, 1063–

1075 (2021).

63.   Spettel, K. *et al. Candida auris* in Austria—what is new and what is different.

Journal of Fungi **9**, 129 (2023).

64.   Wurster, S. *et al. Drosophila melanogaster* as a model to study virulence and

azole treatment of the emerging pathogen *Candida auris*. *Journal of Antimicrobial*

Chemotherapy **74**, 1904–1910 (2019).

65.   Harpf, V., Rambach, G., Würzner, R., Lass-Flörl, C. & Speth, C. *Candida* and

complement: New aspects in an old battle. *Frontiers in Immunology* **11**, 1471 (2020).

66.   Pechacek, J. & Lionakis, M. S. Host defense mechanisms against *Candida auris*.
Expert Review of Anti-infective Therapy **21**, 1087–1096 (2023).

67.   Arora, P. *et al.* Environmental isolation of *Candida auris* from the coastal
wetlands of Andaman Islands, India. *mBio* **12**, e03181–20 (2021).

68.   Escandón, P. Novel environmental niches for *Candida auris*: Isolation from a
coastal habitat in Colombia. *Journal of Fungi* **8**, 748 (2022).

69.   Uden, V. & Kolipinski, M. C. *Torulopsis haemulonii nov. spec.*, A yeast from the
Atlantic Ocean. *Antonie Van Leeuwenhoek* **28**, 78–80 (1962).

70.   Bavia, L., Santiesteban-Lores, L. E., Carneiro, M. C. & Prodocimo, M. M.
Advances in the complement system of a teleost fish, *Oreochromis niloticus*. *Fish &*
Shellfish Immunology **123**, 61–74 (2022).

71.   Buchmann, K., Karami, A. M. & Duan, Y. The early ontogenetic development of
immune cells and organs in teleosts. *Fish & Shellfish Immunology* **146**, 109371 (2024).

72.   Tian, H. *et al.* Cytokine networks provide sufficient evidence for the differentiation
of CD4+ T cells in teleost fish. *Developmental & Comparative Immunology* **141**, 104627
(2023).

73.   Johnson, C. J., Davis, J. M., Huttenlocher, A., Kernien, J. F. & Nett, J. E.
Emerging fungal pathogen *Candida auris* evades neutrophil attack. *mBio* **9**, e01403–18
(2018).

74.   Pharkjaksu, S., Boonmee, N., Mitrpant, C. & Ngamskulrungroj, P.
Immunopathogenesis of Emerging *Candida auris* and *Candida haemulonii* Strains.
Journal of Fungi **7**, 725 (2021).

75.   Shapiro, R. S. & Cowen, L. E. Uncovering cellular circuitry controlling
temperature-dependent fungal morphogenesis. *Virulence* **3**, 400–404 (2012).

76.   Leach, M. D. *et al.* Hsf1 and Hsp90 orchestrate temperature-dependent global
transcriptional remodelling and chromatin architecture in *Candida albicans*. *Nature*
Communications **7**, 11704 (2016).

77.   Xiao, W. *et al.* Response and regulatory mechanisms of heat resistance in
pathogenic fungi. *Applied Microbiology and Biotechnology* **106**, 5415–5431 (2022).

78.   Akbarzadeh, A. & Leder, E. H. Acclimation of killifish to thermal extremes of hot
spring: Transcription of gonadal and liver heat shock genes. *Comparative Biochemistry*
and Physiology Part A: Molecular & Integrative Physiology **191**, 89–97 (2016).

79.   Hamied, A. *et al.* Identification and characterization of highly fluorescent pigment
cells in embryos of the Arabian killifish (*Aphanius Dispar*). *iScience* **23**, 101674 (2020).

80.   Alsakran, A. *et al.* Stage-by-stage exploration of normal embryonic development
in the Arabian killifish, Aphanius dispar. *Developmental Dynamics* **n/a**, (2024).

81.   Minhas, R. *et al.* The thermotolerant Arabian killifish, Aphanius dispar, as a novel
infection model for human fungal pathogens. 2024.10.08.617174 (2024)
doi:10.1101/2024.10.08.617174.

82.   Skrzypek, M. S. *et al.* The Candida Genome Database (CGD): Incorporation of
Assembly 22, systematic identifiers and visualization of high throughput sequencing
data. *Nucleic Acids Research* **45**, D592–D596 (2017).

83.   Potrykus, J. *et al.* Fungal iron availability during deep seated candidiasis is
defined by a complex interplay involving systemic and local events. *PLOS pathogens* **9**,
e1003676 (2013).

84.   Kasper, L. *et al.* The fungal peptide toxin Candidalysin activates the NLRP3
inflammasome and causes cytolysis in mononuclear phagocytes. *Nature*
Communications **9**, 4260 (2018).

85.   Bonzi, L. C. *et al.* The time course of molecular acclimation to seawater in a
euryhaline fish. *Scientific Reports* **11**, 18127 (2021).

86.　Varma, A. *et al.* Molecular cloning and functional characterisation of a glucose
transporter, CaHGT1, of Candida albicans. *FEMS microbiology letters* **182**, 15–21
(2000).

87.　Luo, L., Tong, X. & Farley, P. C. The *Candida albicans* gene HGT12
(Orf19.7094) encodes a hexose transporter. *FEMS Immunology & Medical Microbiology*
51, 14–17 (2007).

88.　Brown, V., Sexton, J. A. & Johnston, M. A glucose sensor in Candida albicans.
Eukaryotic Cell **5**, 1726–1737 (2006).

89.　Larcombe, D. E. *et al.* Glucose-enhanced oxidative stress resistance-A protective
anticipatory response that enhances the fitness of *Candida albicans* during systemic
infection. *PLOS Pathogens* **19**, e1011505 (2023).

90.　Martin, R. *et al.* A core filamentation response network in *Candida albicans* is
restricted to eight genes. *PLOS ONE* **8**, e58613 (2013).

91.　Schatzman, S. S. *et al.* Copper-only superoxide dismutase enzymes and iron
starvation stress in Candida fungal pathogens. *The Journal of Biological Chemistry* **295**,
570–583 (2020).

92.　Rybak, J. M. *et al.* Abrogation of Triazole Resistance upon Deletion of *CDR1* in a
Clinical Isolate of *Candida auris*. *Antimicrobial Agents and Chemotherapy* **63**, (2019).

93.　Li, J., Coste, A. T., Bachmann, D., Sanglard, D. & Lamoth, F. Deciphering the
Mrr1/Mdr1 Pathway in Azole Resistance of *Candida auris*. *Antimicrobial Agents and*
Chemotherapy **66**, e00067–22 (2022).

94.　Sun, S., Coelho, M. A., David-Palma, M., Priest, S. J. & Heitman, J. The
evolution of sexual reproduction and the Mating-type locus: Links to pathogenesis of
Cryptococcus human pathogenic fungi. *Annual Review of Genetics* **53**, 417–444 (2019).

95.   Wang, Y. & Xu, J. Population genomic analyses reveal evidence for limited

recombination in the superbug *Candida auris* in nature. *Computational and Structural*

Biotechnology Journal **20**, 3030–3040 (2022).

96.   Overy, D. P., Rämä, T., Oosterhuis, R., Walker, A. K. & Pang, K.-L. The

neglected marine fungi, *Sensu stricto*, and their isolation for natural products' discovery.

Marine Drugs **17**, 42 (2019).

97.   Chow, N. A. *et al.* Potential fifth clade of *Candida auris,* Iran, 2018. *Emerging*

Infectious Diseases **25**, 1780–1781 (2019).

98.   Louvet, M. *et al.* Ume6-dependent pathways of morphogenesis and biofilm

formation in *Candida auris*. *Microbiology Spectrum* **0**, e01531–24 (2024).

99.   Deng, Y., Li, S., Bing, J., Liao, W. & Tao, L. Phenotypic switching and

filamentation in *Candida haemulonii,* an emerging opportunistic pathogen of humans.

Microbiology Spectrum **9**, e00779–21 (2021).

100.   Seiser, S. *et al.* Native human and mouse skin infection models to study *Candida*

auris-host interactions. *Microbes and Infection* **26**, 105234 (2024).

101.   Fan, S. *et al.* Filamentous growth is a general feature of *Candida auris* clinical

isolates. *Medical Mycology* **59**, 734–740 (2021).

102.   Bravo Ruiz, G., Ross, Z. K., Gow, N. A. R. & Lorenz, A. Pseudohyphal growth of

the emerging pathogen *Candida auris* is triggered by genotoxic stress through the S

phase checkpoint. *mSphere* **5**, e00151–20 (2020).

103.   Kim, S. H. *et al.* Genetic analysis of *Candida auris* implicates Hsp90 in

morphogenesis and azole tolerance and Cdr1 in Azole Resistance. *mBio* **10**, e02529–

18 (2019).

104.   Gao, J. *et al.* LncRNA DINOR is a virulence factor and global regulator of stress

responses in *Candida auris*. *Nature Microbiology* **6**, 842–851 (2021).

105. Ismail, A., Bedell, G. W. & Lupan, D. M. Siderophore production by the
pathogenic yeast, *Candida albicans*. *Biochemical and Biophysical Research*
Communications **130**, 885–891 (1985).

106. Heymann, P. *et al.* The Siderophore iron transporter of *Candida albicans*
(Sit1p/Arn1p) mediates uptake of ferrichrome-type siderophores and Is required for
epithelial invasion. *Infection and Immunity* **70**, 5246–5255 (2002).

107. Boiteau, R. M. *et al.* Siderophore-based microbial adaptations to iron scarcity
across the eastern Pacific Ocean. *Proceedings of the National Academy of Sciences*
113, 14237–14242 (2016).

108. Park, J. *et al.* Siderophore production and utilization by marine bacteria in the
North Pacific Ocean. *Limnology and Oceanography* **68**, 1636–1653 (2023).

109. Renshaw, J. C. *et al.* Fungal siderophores: Structures, functions and
applications. *Mycological Research* **106**, 1123–1142 (2002).

110. Johnstone, T. C. & Nolan, E. M. Beyond iron: Non-classical biological functions
of bacterial siderophores. *Dalton Transactions* **44**, 6320–6339 (2015).

111. Nakajima, Y. *et al.* Effect of disrupting the trichothecene efflux pump encoded by
FgTri12 in the nivalenol chemotype of *Fusarium graminearum*. *The Journal of General*
and Applied Microbiology **61**, 93–96 (2015).

112. Xue, P. *et al.* Heme sensing and trafficking in fungi. *Fungal Biology Reviews* **43**,
100286 (2023).

113. Fourie, R., Kuloyo, O. O., Mochochoko, B. M., Albertyn, J. & Pohl, C. H. Iron at
the Centre of *Candida albicans* Interactions. *Frontiers in Cellular and Infection*
Microbiology **8**, 185 (2018).

114. Alselami, A. & Drummond, R. A. How metals fuel fungal virulence, yet promote
anti-fungal immunity. *Disease Models & Mechanisms* **16**, dmm050393 (2023).

115.   Pradhan, A. *et al.* Non-canonical signalling mediates changes in fungal cell wall
PAMPs that drive immune evasion. *Nature Communications* **10**, 5315 (2019).

116.   Weissman, Z. & Kornitzer, D. A family of *Candida* cell surface haem-binding
proteins involved in haemin and haemoglobin-iron utilization. *Molecular Microbiology* **53**,
1209–1220 (2004).

117.   Ding, C. *et al.* Conserved and divergent roles of Bcr1 and CFEM proteins in
Candida parapsilosis and Candida albicans. *PLOS ONE* **6**, e28151 (2011).

118.   Simm, C. *et al.* Disruption of iron homeostasis and mitochondrial metabolism are
promising targets to inhibit *Candida auris*. *Microbiology Spectrum* **10**, e00100–22
(2022).

119.   Sert, N. P. du *et al.* Reporting animal research: Explanation and elaboration for
the ARRIVE guidelines 2.0. *PLOS Biology* **18**, e3000411 (2020).

120.   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical and
powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B:*
Statistical Methodology **57**, 289–300 (1995).

121.   Bancroft, J. & Stevens, A. Theory and Practice of Histological Techniques.
(1977).

122.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. Fastp: An ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

123.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: Summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048
(2016).

124.   Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq
using the Trinity platform for reference generation and analysis. *Nature Protocols* **8**,
1494–1512 (2013).

125. Abastabar, M. *et al.* *Candida auris* otomycosis in Iran and review of recent
literature. *Mycoses* **62**, 101–105 (2019).

126. Muñoz, J. F. *et al.* Clade-specific chromosomal rearrangements and loss of
subtelomeric adhesins in *Candida auris*. *Genetics* **218**, iyab029 (2021).

127. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2.
Nature Methods **9**, 357–359 (2012).

128. Li, B. & Dewey, C. N. RSEM: Accurate transcript quantification from RNA-Seq
data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

129. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics **26**, 139–140 (2010).

130. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
25, 2078–2079 (2009).

131. Ritchie, M. E. *et al.* Limma powers differential expression analyses for RNA-
sequencing and microarray studies. *Nucleic Acids Research* **43**, e47 (2015).

132. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and
correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

133. Ashburner, M. *et al.* Gene Ontology: Tool for the unification of biology. *Nature*
Genetics **25**, 25–29 (2000).

134. The Gene Ontology Consortium *et al.* The Gene Ontology knowledgebase in
2023. *Genetics* **224**, iyad031 (2023).

135. Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life
scale using DIAMOND. *Nature Methods* **18**, 366–368 (2021).

136. Golden, C., Studholme, D. J. & Farrer, R. A. DIAMOND2GO: A rapid Gene
Ontology assignment and enrichment tool for functional genomics. 2024.08.19.608700
(2024) doi:10.1101/2024.08.19.608700.

137.   Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M. & Ishiguro-Watanabe, M.

KEGG for taxonomy-based analysis of pathways and genomes. *Nucleic Acids Research*

51, D587–D592 (2022).

138.   Kanehisa, M., Sato, Y. & Morishima, K. BlastKOALA and GhostKOALA: KEGG

tools for functional characterization of genome and metagenome sequences. *Journal of*

Molecular Biology **428**, 726–731 (2016).

139.   Bateman, A. *et al.* The Pfam protein families database. *Nucleic Acids Research*

32, D138–D141 (2004).

140.   Eddy, S. R. Accelerated Profile HMM Searches. *PLOS Computational Biology* **7**,

e1002195 (2011).

141.   Almagro Armenteros, J. J. *et al.* SignalP 5.0 improves signal peptide predictions

using deep neural networks. *Nature Biotechnology* **37**, 420–423 (2019).

142.   Hallgren, J. *et al.* DeepTMHMM predicts alpha and beta transmembrane proteins

using deep neural networks. (2022) doi:10.1101/2022.04.08.487609.

143.   Gíslason, M. H., Nielsen, H., Almagro Armenteros, J. J. & Johansen, A. R.

Prediction of GPI-anchored proteins with pointer neural networks. *Current Research in*

Biotechnology **3**, 6–13 (2021).

144.   Camacho, C. *et al.* BLAST+: Architecture and applications. *BMC Bioinformatics*

10, 421 (2009).

145.   Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO summarizes and

visualizes long lists of gene ontology terms. *PLOS ONE* **6**, e21800 (2011).

146.   Farrer, R. A. Synima: A Synteny imaging tool for annotated genome assemblies.

BMC Bioinformatics **18**, 507 (2017).

147.   Emms, D. M. & Kelly, S. OrthoFinder: Phylogenetic orthology inference for

comparative genomics. *Genome Biology* **20**, 238 (2019).

148.   Edgar, R. C. MUSCLE: Multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Research* **32**, 1792–1797 (2004).

149.   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – approximately maximum-
likelihood trees for large alignments. *PLOS ONE* **5**, e9490 (2010).

150.   Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493–500 (2024).

151.   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. *Protein Science* **30**, 70–82 (2021).

152.   Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-
analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

153.   Sievers, F. & Higgins, D. G. Clustal omega. *Current Protocols in Bioinformatics*
48, 3.13.1–3.13.16 (2014).

1217

1218   **End Matter**

1219

1220   **Acknowledgements**

1221

We acknowledge funding from the MRC Centre for Medical Mycology at the
University of Exeter (MR/N006364/2, MR/V033417/1), MRC Doctoral Training Grants
MR/P501955/2 and MR/W502649/1, Wellcome Trust Career Development Award
(215239/Z/19/Z) and Wellcome Trust fellowship (219551/Z/19/Z), National Centre for
the Replacement, Refinement and Reduction of Animals in Research (NC3Rs)
(NC/X001121/1), and the NIHR Exeter Biomedical Research Centre. TB also was
supported *via* Elizabeth Ballou and a Sir Henry Dale Fellowship jointly funded by the
Wellcome Trust and the Royal Society (211241/Z/18/Z) and the Lister Institute. The
views expressed are those of the authors and not necessarily those of the NIHR or the
Department of Health and Social Care. We thank staff in the Aquatic Resource Center

at the University of Exeter for husbandry of the fish and the support for developing

protocols for feeding, spawning, and raising new generations of this relatively new

model species, including Sam Worthington, Chloe Flint and Richard Silcox. We also

thank the Exeter Sequencing Service facility and support from Wellcome Trust

Institutional Strategic Support Fund (WT097835MF), Wellcome Trust Multi User

Equipment Awards (WT101650MA and 218247/Z/19/Z), Medical Research Council

Clinical Infrastructure Funding (MR/M008924/1) and BBSRC LOLA award

(BB/K003240/1). We also thank the University of Exeter High-Performance Computing

(HPC) facility funded by the UK MRC Clinical Research Infrastructure Initiative (award

number MR/M008924/1). We thank Tanmoy Chakraborty, Dhara Malavia and Neil Gow

for providing *C. auris* strains, Teigan Veale, Sumita Roy, Jonathan Ball and Darren

Thomson for assistance in microscopy and Chloe Pelletier for advice on aggregation.

We also thank Microtechnical Services, Exeter, UK, for assistance with histological

sectioning.

1246

1247 **Author Contributions**

1248

1249 HG: designed and conducted experiments, performed analysis, drafted and

1250 wrote the paper. NH, JG, QM, TB: optimised, designed or performed experiments. RF,

1251 TK, TB, MR, JR, DW, AB: designed experiments, obtained funding. All authors edited

1252 the manuscript.

1253

1254 **Competing Interests**

1255

1256 JR has received honoraria from Gilead for conference presentations. Other

1257 authors declare no competing interests.

1258

1259 **ORCID IDs**

1260

HG (https://orcid.org/0000-0003-2797-852X), TB (https://orcid.org/0000-0002-

5315-2759), NH (https://orcid.org/0000-0003-0260-023X), JG (https://orcid.org/0009-

0003-3803-9496), QM (https://orcid.org/0000-0001-9892-4997), AB

(https://orcid.org/0000-0002-7381-131X), DW (https://orcid.org/0000-0001-6617-0123),

JR (https://orcid.org/0000-0002-1338-7860), MR (https://orcid.org/0009-0009-0746-

4127), TK (https://orcid.org/0000-0002-5373-2737), RF (https://orcid.org/0000-0001-

8456-7458)

1268

Figures

Figure 1. Modelling C. auris infection in the Arabian killifish (A) _Survival curves for_

embryos injected with each C. auris clade or sham injection. Significance testing

between groups is given in **Table S1**; hours post injection (HPI). **(B)** Colony forming

units (CFUs) per clade from embryos homogenised after injection as an indicator of

*injection dose and in vivo growth. **(C)** H&E Staining of AK yolk-sac at 48 HPI for each*

clade or sham injection; morphology is indicated by blue arrows, with insets indicating

typical yeast (clade III) and filamentous morphology (clade V). The yolk-sac edge is

*oriented at the top. **(D)** Illustrative time-lapse microscopy of embryo death including*

yolk-sac collapse during infection (clade I infection) or survival (sham injection) using

*Acquifer live imaging. **(E)** Experimental scheme for differential gene expression analysis*

of RNA extracted from AK embryos injected at 3 days post fertilisation (DPF).

Figure 2. AK host responses to C. auris infection: (A) Number of differentially
expressed genes (DEGs) for each of two comparisons, sham injection vs no injection
(left panel), and C. auris injection vs sham injection (right panel), at either 24 or 48 h
post injection (HPI). (B) Volcano plot showing log-fold change and significance
according to -log^{10} false discovery rate (FDR) for sham injection vs no injection. (C)
Heatmap demonstrating log-fold change for differential expression between C. auris
injection and sham injection for groups of genes. (D-E) Volcano plots showing log-fold
change and significance for DEGs expressed during infection with the two more virulent
clades of C. auris at 48 HPI, clade I (D, DEGs in red) and clade IV (E, DEGs in green).

1293

***Figure 3: C. auris gene expression** in vivo vs in vitro **(A)** Total DEGs per clade for*

each comparison between 24 h post injection (HPI), 48 HPI, and growth in YPD alone.

***(B-D)** Upset plots for each comparison, indicating DEGs that were significant across all*

*five clades (far left) and for each individual clade (far right) and each set in between. **(E)***

Combined set of significant DEGs common to all five clades between in vivo infection

*and in vivo growth in YPD. Genes are coloured by manually annotated category **(Table***

S4). (F) Volcano plots demonstrate mean log-fold change for DEGs identified for any

clade. Genes with infinite false discovery rate (FDR) are indicated with an arrow. Genes

*included in the combined set are coloured by category as per panel **E**.*

Figure 4: Interclade comparisons for virulence and filamentation: (A): *The set of*

differentially expressed genes (DEGs) shared across all comparisons between clade I

and IV (more virulent) and clade II and III (less virulent), coloured by manually

annotated category, at either 24 or 48 h post infection (HPI). *(B)* Volcano plots

demonstrate mean log-fold change for each DEG found in any comparison; genes

included in the combined set are coloured as per *A. (C)* The set of DEGs shared across

all comparisons between filamenting clade V all other clades (I-IV) in vivo, coloured by

manually annotated category. *(D):* Volcano plots demonstrating mean log-fold change

for all comparisons between clade V and clades I-IV at 24 HPI and 48 HPI for each

DEG found in any comparison; genes included in the combined set are coloured as per

C.

1315

Figure 5. Xenosiderophore Transporter Analysis: (A) *AlphaFold 3 predicted protein*
folding for peptide sequences for a xenosiderophore transporter candidate, illustrating
the conserved 14-transmembrane domain similarity across the families. (B) *RAXML*
phylogeny using 1000 bootstraps and LG4X model of Muscle-aligned protein

sequences for orthologues of C. albicans Siderophore Transport 1 SIT1 gene. Scale bar

indicates number of substitutions per site. Orthofinder cluster assignment notes which

grouping each gene was assigned to by our Synima/Orthofinder pipeline. (C) Co-

expression network of associated gene expression involving siderophore transport

candidate genes. (D) Co-expression network across available RNA-seq data from meta-

*analysis of 35 studies (**Table S4**), demonstrating a different set of transcriptional*

networks indicated by external experimental data.

Figure 6. Panfungal Xenosiderophore Phylogeny: (A) *Phylogeny of Blastp hits for*

C. albicans SIT1 across fungal genomes available in FungiDB. Sequences were aligned

with Clustal Omega and FastTree, scale bar shows substitutions per site. Two main

branches emerged, a xenosiderophore transporter candidate branch (XTC) and a haem

transport-related branch (HTR), based mainly on the GO terms assigned (outer ring).

(B) *Bar chart depicting per-species mean Blastp hits for siderophore transporters.* ***(C)***

Heatmap of all DEGs across comparisons in this paper for each XTC and HTR gene,

assigned in order of contig in the B8441 v3 reference genome. White spaces represent

clades where each orthologue was not found in Orthofinder (XTC4, XTC9 and XTC11 in

clade II, XTC5 in II and V, XTC10 in II and IV, ***Fig. 5B***).

Supplementary Figures

1339

Supplementary Figure 1: *Profiles of C. auris growth, morphotype, morphology, and*

gene expression. ***(A)*** *Growth curves for five representative strains of C. auris over 24 h*

in vitro. ***(B)*** *Rough colonies demonstrated on YPD-agar after recovery from AK infection*

in clade V only (blue arrows). ***(C)*** *Filamentous forms (yellow arrows) isolated from rough*

colonies and imaged on light microscopy in distilled de-ionised water. ***(D)*** *Principle*

components analysis (PCA) of fragments per kilobase per million (FPKM) gene

expression for AK host transcription. ***(E)*** *Correlation matrix for individual replicates' log_2-*

transformed transcript counts per million for AK responses to sham injection and no

injection ***(F)*** *Correlation matrix as above for AK responses to C. auris infection and*

sham injection. ***(G)*** *PCA of FPKM gene expression for C. auris pathogen transcription.*

(H) *Correlation matrix as above for C. auris gene expression in vivo and in vivo across*

five clades.

1353

Supplementary Figure 2: A. dispar gene ontology, pathway and domain

enrichment during infection vs sham control: The GJEY01 A. dispar transcripts

were annotated for Gene Ontology (GO) terms using DIAMOND2GO, Kyoto

Encycopaedia of Genes and Genome pathways using BLASTKoala, and PFAM

domains using HMMScan. For each set of up-regulated or down-regulated genes in

each comparison between an injected C. auris clade and sham injection, Fisher's exact

test was used with Benjamini-Hochberg (BH) correction for multiple testing for each set,

with a false discovery rate (FDR) cut-off of 0.001. Mean log-fold change and ratio of

genes in that set was calculated for genes in each set possessing each feature.

Redundant GO terms were removed using Revigo.

Supplementary Figure 3. Contextualising differential gene expression across orthogroups and broder RNA-seq datasets. (A) Single copy orthologue phylogeny and Synima/Orthofinder synteny plot of reference genomes for *C. albicans, C. haemulonii* and the five clades of *C. auris* used in this study. Despite many orthologues, there is little synteny with *C. albicans*; prominent translocations and inversions are seen between *C. haemulonii* and *C. auris*, and for clade II. **(B)** Counts are shown for orthologues contained within and between species, whether single copy or multiple copy, with additional counts for accessory genomes. Clade V is the only *C. auris* clade with a single orthogroup, containing six genes, while clade IV has the highest number of apparently unique genes that were not assigned to any orthogroup. **(C)** Differential gene expression for *in vivo vs in vitro* comparisons, comparing numbers of DEGs identified when using a single core reference genome (B8441, clade I) or a clade-specific reference genome for each individual clade. **(D)** Differential expression of specific genes not present in core reference genome. **(E)** RNA-seq meta-analysis. 36 public datasets were identified for *C. auris* transcriptome expression and assessed for variance by PCA dimensionality reduction of normalised expression of single copy orthologues, indicating AK dataset clusters similarly to experiments using an *ex vivo* blood model and other experiments. **(F)** Hierarchical clustering of per-gene variance, calculated for all B8441 core genes for each study, demonstrating similar clustering and profile of within-experiment variance per gene.

1385

Supplementary Figure 4. *C. auris* Gene Set, Pathway and Domain Enrichment:

Transcripts from *C. auris* reference genomes for each clade were annotated for Gene

Ontology (GO) terms using DIAMOND2GO, Kyoto Encycopaedia of Genes and

Genome pathways using BLASTKoala, PFAM domains using HMMScan,

transemembrane domains with DeepTMHMM, GPI-anchored domains with NetGPI,

SignalP domains with SignalP5, and membership within the accessory genome using

Synima/Orthofinder. For each set of up-regulated or down-regulated genes in each

comparison, Fisher's exact test was used with Benjamini-Hochberg (BH) correction for

multiple testing for each set, with a false discovery rate (FDR) cut-off of 0.001. Mean

log-fold change and ratio of genes in that set was calculated for genes in each set

possessing each feature.

1397

Supplementary Figure 5. *C. auris* **Mating-Type Locus Gene Expression: (A)**
Transcripts (fragments per kilobase per million, FPKM) for orthogroups aligned to the
same reference genome (left-hand panels) and clade-specific reference genomes (right-
hand panels), indicating similar transcript levels between clades for e.g. *PIKA*, *OBPA*
and *PAP1* non-mating genes when aligned to clade-specific reference genomes. **(B)**
Comparing individual transcript levels for each orthogroup in more *vs* less virulent
clades *via* Wilcoxon signed-rank test with Bonferroni correction to minimise false
positives, demonstrating that only down-regulation of *THI4* (at 24 HPI), *MRD1* (at both
time-points), and up-regulation of *PGA30* and *PIKA* were significantly different.

1407   Significance levels are as follows: * p <= 0.05, ** <= 0.01, *** <= 0.001, **** <= 0.0001,

1408   ***** <= 0.00001.

 Supplementary Figure 6: (A-B) Comparing individual transcript levels for each

orthogroup in filamentous clade V vs all other clades via Wilcoxon signed-rank test with

Bonferroni correction to minimise false positives, demonstrating at 24 HPI (A) and

48 HPI (B), excluding any genes that were assigned to be 'hypothetical proteins' only,

up-regulation of two HGT13 orthologues and RNH70 were significantly different at both

time-points, with up-regulation of MED7 and SPS19 at 24 HPI. Significance levels are

as follows: Significance levels are as follows: * p <= 0.05, ** <= 0.01, *** <= 0.001, ****

<= 0.0001, ***** <= 0.00001.

Supplementary Tables

**Supplementary Table 1. Clade-Representative *Candida auris* Strains Used In This
Study:** Representative strain for each major clade of *C. auris* used in the study, with
notes on human disease source, country of discovery, and date of isolation. Mortality
statistics are given in terms of absolute numbers of embryos over biological triplicate
experiments. Additional significance testing is also noted for pairwise comparisons by
Log-Rank test with Benjamini-Hochberg (BH) correction.

**Supplementary Table 2: Differentially Expressed Genes in *A. dispar*
Microinjection:** All differentially expressed genes (DEGs) for *A. dispar* for (upper rows)
sham injection *vs* no injection and (lower rows) *C. auris* injection *vs* sham injection.
Gene IDs and descriptions follow annotations by Bonzi *et al.*[85]; log-fold change and -
$\log_{10}$(false discovery rate) are calculated across all clades as a mean value for *C. auris*
injection. Significant DEGs for each set of embryos injected with each clade are given
(1 = significant, 0 = not significant), and the total number of clades where DEGs where
significant is also given.

Supplementary Table 3. Differentially Expressed Genes in *C. auris* Infection: (A)
Combined set of differentially expressed genes (DEGs) expressed by five clades during

infection Each set of DEGs expressed *in vivo vs in vitro* for five clades was combined,

resulting in a set of genes that were specific to *C. auris* infection at these timepoints. **(B)**

Combined set of DEGs between more virulent and less virulent clades: Each set of

DEGs between clades I & IV (more virulent) and II & III (less virulent) was compared at

24 HPI and 48 HPI, resulting in a combined set expressed in infection. **(C)** Combined

DEG Sets: between filamentous clade V and all other clades: Each set of DEGs

between clade V and all each clades I-IV was compared at 24 HPI and 48 HPI, resulting

in a combined set expressed in infection, excluding those significant during growth in

YPD broth where clade V was not observed to filament. **(D)** Accessory genes not found

in clade I reference genome: DEGs without orthologues in the clade I reference were

identified and annotated.

1450

1451 **Supplementary Table 4. Publicly available RNA-seq Data on 16th April 2024:** A

1452 systematic re-analysis of RNA expression across NCBI-available datasets for *C. auris*

1453 gene expression, categorised by research theme for analysis (**Fig. S1D-E**).

1454