Xenosiderophore transporter gene expression and clade-specific filamentation in *Candida auris* killifish infection

Hugh Gifford[a], Tina Bedekovic[a], Nicolas Helmstetter[a], Jack Gregory[a], Qinxi Ma[a], Alexandra C. Brand[a], Duncan Wilson[a], Johanna Rhodes[b,c], Mark Ramsdale[a], Tetsuhiro Kudoh[d,1], Rhys A. Farrer*[a,1]

Abstract

Candida auris is a critical priority fungal pathogen (World Health Organization). Clinical management is challenging due to a high mortality rate, rapidly increasing antifungal resistance, and frequent nosocomial outbreaks. A critical bottleneck in understanding virulence is the lack of gene expression profiling models during infection. We developed a fish embryo yolk-sac microinjection model using *Aphanius dispar* (Arabian killifish; AK) at human body temperature. This enabled interrogation of infection dynamics *via* dual host-pathogen RNA-seq across five major clades of *C. auris* (I-V). Host responses included heat shock, complement activation, and nutritional immunity, notably haem oxygenase (*HMOX*) expression during clade IV infection. We identified a pathogen transcriptional signature across all five clades of *C. auris* strongly enriched for putative xenosiderophore transmembrane transporters. We describe this novel family and a sub-clade of five putative haem transport-related (*HTR*) genes. Only the basal clade V isolate formed filaments, associated with canonical and atypical regulators of morphogenesis. Clades I and IV demonstrated increased virulence, accompanied by up-regulation of three *HTR* genes in clade IV, and the non-mating mating-type locus (*MTL*) gene *PIKA* in both clades. Our study provides new insight into *C. auris* pathogenesis, highlighting species-wide *in vivo* up-regulation of *XTC* genes during host tissue infection.

Significance statement

Candida auris is an emerging human fungal pathogen and global public health threat, yet *in vivo* transcriptomic analysis of tissue infection has remained elusive. Using yolk-sac infection in Arabian killifish, we profiled gene expression across five major *C. auris* clades. We found that the basal clade V uniquely undergoes filamentation during infection, while all clades upregulate members of a large, expanded family of xenosiderophore transporter candidate genes. These findings highlight the important roles for iron acquisition and morphological switching in pathogenesis, revealing potential mechanisms of immune evasion and fungal persistence, and identifying candidate targets for antifungal therapy.

[a]MRC Center for Medical Mycology, University of Exeter, Geoffrey Pope Building, Stocker Road, Exeter,

United Kingdom, EX4 4QD

[b]Department of Medical Microbiology, Radboudumc, Geert Grooteplein Zuid 10, Nijmegen, Netherlands,

6525 GA

[c]MRC Centre for Global Infectious Disease Analysis, Imperial College London South Kensington Campus,

London, United Kingdom, SW7 2AZ

[d]Biosciences, University of Exeter, Geoffrey Pope Building, Stocker Road, Exeter, United Kingdom, EX4

4QD

[1]Co-Senior Author

Corresponding Author: Rhys Farrer

Email: R.Farrer@exeter.ac.uk

Author contributions: HG: designed and conducted experiments, performed analysis, drafted and wrote

the paper. NH, JG, QM, TB: optimised, designed or performed experiments. RF, TK, TB, MR, JR, DW,

AB: designed experiments, obtained funding. All authors edited the manuscript.

Competing interest statement: Authors declare no competing interests.

Classification: Biological sciences, Microbiology.

Pre-print: This article has been made available as a pre-print *via* the bioRxiv server.

Keywords: *Candida auris*, *Aphanius dispar*, dual host-pathogen RNA-seq, siderophore transport,

filamentation

Main text

Introduction

Candida auris is an ascomycetous yeast that has rapidly progressed from a recently identified

species to a World Health Organization (WHO) critical priority human fungal pathogen and global public

health threat[1,2]. The earliest known isolate was retrospectively identified from a bloodstream infection

dating to 1996[3], following its initial description as a novel species in 2008[4]. Over the past decade, *C. auris*

has caused nosocomial outbreaks globally, leading to intensive care unit (ICU) closures and substantial

infection control challenges and costs[5–7]. Alarmingly, *C. auris* is now a leading cause of invasive

candidiasis in several hospital settings globally[8,9]. *C. auris* bloodstream infections carry an associated

mortality rate of approximately 45%[10] and are often resistant to azole antifungals, with an increasing

number of pan-drug resistant isolates reported globally[11]. Phylogenetic analysis has revealed five major

clades of *C. auris*, and a minor sixth clade recently found[12,13], suggesting multiple populations of *C. auris*

have simultaneously emerged from unknown environmental reservoirs that have facilitated its rapid

adaptation to human infection, drug resistance, and healthcare-associated persistence[14,15]. Clade-specific

differences have been noted: clades I, III, and IV are more commonly associated with invasive disease,

whereas clades II and V have been linked to otomycosis[16]. Mortality from candidaemia may be higher

with clades I and IV than clade III, but this difference has not been shown to be statistically significant[17].

These differences in clinical presentation and virulence may reflect underlying genotypic variation across

clades.

Gene expression profiling has become a standard approach for assessing host and pathogen

responses to infection. However, no experiment has described the gene expression of *C. auris* during

in vivo infection of living tissue to date. Studies have assessed the pathogen transcriptome in an *in vivo*

murine catheter model[18], human whole blood *ex vivo*[19], co-incubation with human dermal fibroblasts[20] or

murine cell line-derived macrophages[21]. Additionally, host responses have been assessed in murine

peripheral skin[22], human peripheral blood-derived monocytes[23], and murine bone marrow-derived

macrophages[24]. *C. auris* has been shown to reproduce as budding yeasts in mammalian tissues[25,26] and

possess multiple adhesins associated with an aggregation phenotype[27,28]. There has been no evidence of

toxin synthesis unlike distantly related *C. albicans*[29,30]. *C. auris* digests of host tissue through secreted

aspartyl proteases[31] and evades host immune responses through cell surface masking[32] and intra-

phagocytic survival with cell lysis[33]. However, these studies have not interrogated *in vivo* in-host pathogen

gene expression, except for the dual RNA-seq study using the *ex vivo* whole blood model, for which it

was possible to compare gene expression profiles between *C. auris* and other *Candida* species[19].

The absence of *in vivo* tissue infection profiling for *C. auris* may be due to the lack of adequate

RNA recovery from current models. Most murine models, such as BALB/c[25,34–36] and C57BL/6 mice[24],

have rarely demonstrated lethality through *C. auris* infection[23,37,38] without major immunosuppression with

cyclophosphamide/cortisone[39–50], monoclonal antibody neutrophil depletion[26,51], or other

immunodeficiency[27,52,53]. Additionally, mouse models are also expensive and associated with concerns

over environmental pathogen shedding[54]. Alternative host models such as *Galleria mellonella*[34,44,55–61],

Drosophila melanogaster[62] and *Caenorhabditis elegans*[60] have been employed, but these may lack key

elements of mammalian-relevant immune systems, thus limiting their scope of application. In patients,

immune control of *C. auris* is thought to depend on myeloid cell activity *via* C-type lectin receptors such as

complement receptor 3 and mannose receptor MMR, followed by interleukin (IL-17) pathway

activation[63,64]. The zebrafish (*Danio rerio*) model has allowed hindbrain[32,65] and swim bladder[66]

microinjection modelling of *C. auris* infection, but rarely at temperatures above 30 °C, which zebrafish are

broadly intolerant of. Fungal transcriptional programmes can vary greatly across temperatures and are

critical for mammalian pathogenesis[67–69]. Despite these successes in profiling host responses, none of

these models have enabled the profiling of *C. auris* RNA during infection.

A marine model could also shed light not only on human infection but also on a possible

evolutionarily adaptive niche. The environmental origins of *C. auris* have not been conclusively

demonstrated; however, a marine origin has been hypothesised and supported by successful isolation of

C. auris in coastal waters of both Indian and Pacific oceans[70,71]. Notably, a close relative of *C. auris* in the

Metschnikowiaceae clade, *C. haemulonii*, was first isolated in 1961 from the gut of blue-striped grunt

(genera name "*Haemulon*" for their blood-red mouth interior)[72]. Fish larvae must survive hostile and varied

ocean environments, demonstrating very early innate immune responses such as antimicrobial peptide

production and the complement cascade[73,74]. Only in adulthood are teleost fish expected to possess

functional IL-17 subtypes (A and F) that lead to T-helper-17 subset differentiation and adaptive immunity

in response to fungal infection[75]. A thermotolerant teleost fish embryo model such as the *Aphanius dispar*,

also known as the Arabian killifish (AK), which can acclimatise to temperatures up to and including

40 °C[76–78], offers a balance between considerations such as suitability for experimental manipulation and

applicability to human fungal infection at mammalian temperatures[79,80]. Additionally, fish embryo models

align with the principles of replacement, refinement and reduction of animals in laboratory research and

offer an opportunity to develop novel animal models with ethical benefits over models such as adult

mice[81].

In this paper, we present the first study exploring in-host virulence of *C. auris* clades I-V in the AK

yolk-sac microinjection model at 37 °C. We supplemented our validation of the model with histology,

fungal burden recovery, and dual host-pathogen RNA-seq. By comparing *in vivo* and *in vitro* expression

profiles of *C. auris*, we detected a cross-clade expression signature that was significantly enriched for

siderophore transporter *SIT1* paralogues, which mediate uptake of siderophores produced by other

microbes (xenosiderophores). We designated this expanded gene family as xenosiderophore transporter

candidates (*XTC*), which included a haem transport-related (*HTR*) sub-clade. We also identified *in vivo*

filamentation by clade V during infection, which was associated with orthologues of key hypha-associated

genes from *C. albicans*. Together, these findings substantially enhance our knowledge of the processes

employed by *C. auris* during infection.

Results

AK yolk-sac microinjections with each of the five *C. auris* clades demonstrated lethality (defined

by cessation of heartbeat) at 37 °C (92-100% *vs* 6% for sham injections; **Figure 1A-B**). Earlier lethality

was significantly higher for clade I compared to clades II, III or V ($p < 0.001$) and for clade IV compared to

clades II or III ($p < 0.04$; **Figure 1B**, **Table S1**). To investigate morphological outcomes of infection, we

compared histological sections of infected yolk-sacs at 48 hours post infection (HPI). All clades exhibited

budding yeast morphologies both *in vitro* and *in vivo*, except for the clade V strain, which demonstrated

predominantly filamentous forms when grown *in vivo* (**Figure 1C**). Growth kinetics showed only minor

inter-clade variation. Clade II grew more slowly in YPD-broth at 37 °C (**Figure S1A**), although this did not

translate into lower CFU recovery from infected embryos (**Figure 1D**). On average (x), 673 CFUs were

recovered per infected embryo post injection, which increased 166-fold at 24 HPI ($x = 1.12 \times 10^5$) and by

an additional ~60% at 48 HPI ($x = 1.78 \times 10^5$). Filamentous clade V isolates formed rough (*vs* smooth)

colonies on YPD-agar in approximately 1-2% of cases (**Figure S1B**), composed of elongated yeast cells,

filamentous and pseudohyphal structures measuring >10-50 µm (**Figure S1C**). These results confirm that

AK yolk-sac microinjection supports active *C. auris* infection, leading to embryo mortality by 7 days post-

injection. This system therefore provides a robust and informative *in vivo* model for studying the

pathogenicity of human fungal pathogens such as *C. auris*.

To characterise host responses to *C. auris* during infection, we identified host differentially

expressed genes (DEGs) in AK yolk-sacs between infection by each clade and sham injections

(**Figure 2**). Across all infections, 685 host DEGs were detected (representing 2.49% of potential genes),

including 183 up-regulated and 508 down-regulated genes in response to at least one clade. Notably,

there were no DEGs common to infection by all five clades. Only five DEGs (four up-regulated genes and

one down-regulated gene) were shared across infections with four clades (**Table S2**). Up-regulated

genes included Laminin subunit alpha-1 (*LAMA1*, not significant in response to clade III) and three genes

that were not significant in response to clade II, Glucose-6-phosphatase (*G6PC*), DN117939 (hypothetical

protein) and Tryptophan aminotransferase protein 1 (*TAR1*); immunoglobulin-like fibronectin type III

domain-containing protein 1 (*IGFN1*) was down-regulated. These findings suggest host transcriptional

responses to *C. auris* are modest and clade-specific at these early time points, possibly reflecting

variation in virulence and stages of infection.

Although we did not observe a shared host transcriptional response across all *C. auris* clades, we

identified distinct, clade-specific host responses to infection. At 24 HPI, clade II elicited the largest host

response, with 41 up-regulated genes and 210 down-regulated genes (**Figure 3A**). Minimal differences

were detected between sham injection and uninjected controls, confirming the specificity of the infection-

induced response (**Figure 3B**, SI Appendix). By 48 HPI, clade IV induced the strongest transcriptional

response, with 49 up-regulated and 104 down-regulated host genes. Notably, we detected differential

expression several key immune-related genes across clades, with the most pronounced changes in

clades I and IV. These included heat shock proteins (*HSP30* and *HSP70*), complement cascade

elements, pattern recognition receptors (PRRs), inflammatory transcription regulators, and genes involved

in nutritional immunity (**Figure 3C**). Among genes associated with pathogen recognition, *CARD14* was

most prominently up-regulated during clade I and IV infection at 48 HPI (**Figure 3D-E**), while *NLRP3* (a
key inflammasome component) was expressed during clade III infection at 24 HPI.

Functional enrichment analysis of host DEGs revealed several significantly enriched Gene
Ontology (GO) terms (and additional annotations), particularly in response to the more virulent clades I
and IV at 48 HPI. Clade I infection was associated with 15 enriched GO terms, driven in part by up-
regulation of *LAMA1* and *LAMA2* (orthologues of Laminin subunit-α, **Figure 3D**). In clade IV-infected yolk-
sacs, enriched terms were related to haem catabolism, haem binding, and haem oxygenase (*HMOX*)
activity. Two *HMOX* orthologues were major contributors to this response, such as haem oxidation (log-
fold change x = 3.96, FDR < 0.0003, **Figure 3E**, **Figure S3**). Quantitative PCR confirmed increased
expression of *HMOX* genes in host; embryos infected by clade IV demonstrated late up-regulation at
48 HPI with a fold change of 3.26 and 9.21 for two *HMOX* genes DN109585 and DN112160 respectively
(compared to uninfected embryos; **Figure S2A**). Another host up-regulated gene from infection by clade
IV was ferroxidase hephaestin (*HPHL1*), which is also involved in iron nutritional immunity. Together,
these findings suggest that heat shock proteins, nutritional immunity (particularly iron), and the
complement effect system are important elements of the AK larval response to *C. auris* microinjection at
37° C.

We compared the in-host transcriptome to expression induced by growth in a standard rich
laboratory media (YPD). Principle components and sample correlation clustering from gene expression
data indicated a predominant separate clustering of *in vivo vs in vitro* conditions, indicating that the gene
expression profile of *C. auris* during infection is substantially different to that when it is grown in nutrient-
rich laboratory media (**Figure S1G and S1H**). We identified 960 DEGs between *in vivo* AK infection at
24 HPI or 48 HPI *vs in vitro* culture in YPD over 24 h (499 up-regulated, 523 down-regulated, 17.7% of
potential genes, **Figure 4A**). We identified few DEGs between the two time-points 24 HPI to 48 HPI, with
zero DEGs for clade IV and V between those time points (**Figure 5A**). The highest numbers of up-
regulated DEGs were identified in the filamentous clade V at 24 HPI and in the more virulent clades I and
IV at 48 HPI (**Figure 5B-C**). Six functional annotation terms were enriched across all in-host infections at
48 HPI, related to major superfacilitator family carbohydrate and siderophore-iron transmembrane
transport (**Figure 6**). Quantitative PCR confirmed increased expression of *SIT* genes that had been
significantly up-regulated by *C. auris* at 48 HPI (**Figure S2B**): during embryo infection, clade IV
demonstrated up-regulation with a fold change of 2.55 and 8.16 at 24 and 48 HPI respectively for *XTC3*
(B9J08_003921) compared to culture in YPD alone. Furthermore, *XTC7* (B9J08_001487) was up-
regulated with a fold change of 39.5 and 130.8 at 24 and 48 HPI respectively. Together, this analysis
reveals that during infection, which is a nutrient-limited environment, *C. auris* is responding by
significantly increasing expression of genes involved in sugar uptake, as well as members of the *C. auris*

expanded gene family of siderophore transporters.

Thirteen DEGs common to all clades of in-host transcription were transmembrane transporters,

including up-regulation of transporters for xenosiderophores (*SIT1* genes 3921 and 1487), sugar

transporters (*HGT2*, *HGT12*), an ammonia transporter (*ATO2*), and a nicotinic acid transporter (*TNA1*).

We also found down-regulation of several drug-resistance-related efflux pumps (*MDR2* and *DTR1*) and

peptide transporters (*TPO3*, *PTR22*, **Figure 4B**, **Table S3**). Surprisingly, several putative virulence

factors such as lipase *LIP1* and cell wall-related *KRE6* and *SOD6* were also down-regulated across all

five clades, as were *ALS4* and *MDR1* in at least one clade (**Figure 4C**). Several members of the *SIT1*

family were differentially expressed by <5 of the clades including 0002, 1457, 1458, 1542,2110, 2241,

2465, 3908 and 4097. Meanwhile, two *SIT1* genes were down-regulated across 3 of the clades (1519 and

1499). Several *SIT1* genes were upregulated in ≥1 clade and down-regulated in ≥1 clade (3908 and

2465). The strong signal of differential expression in the *SIT1* expanded gene family highlights the

potential importance during in-host survival.

The more virulent clades (I and IV) are **a**-type mating type locus (*MTL*), while the less virulent

clades (II, III and V) contain the **α**-type *MTL*. Surprisingly, we found differential expression in several non-

mating genes between more virulent and less virulent clades, including poly-A polymerase (*PAP1*),

oxysterol binding protein (*OBPA*) and phosphatidylinositol 4-kinase (*PIKA*, **Figure 7B**). Within the set of

10 DEGs shared across all comparisons, several genes outside the *MTL* were up-regulated, including

vacuolar protein sorting gene, *VPS70*, and filamentous growth regulator, *FGR14*. Additionally, the drug

efflux pump, *MDR1*, was down-regulated across all comparisons, supporting the idea of a fitness trade-off

between resistance and virulence.

To explore drivers of filamentation by the clade V strain, we identified 1,046 DEGs between clade

V and each of the other clades (I-IV), comprising 601 up-regulated and 526 down-regulated genes across

all comparisons. We identified a set of 57 genes shared across all comparisons for *in vivo* conditions only,

since we did not observe filamentation in nutrient-rich YPD. Twenty-two genes were up-regulated at

24 HPI, seven at 48 HPI, and 18 at both time-points (**Figure 7C**). Most of the up-regulated genes were

metabolic (*n* = 13), including two cytochrome P450 *ERG5* orthologues, though *ERG11* was down-

regulated. Transmembrane transporters for sugar (including two *HGT13* orthologues), amino acids and

siderophores (*SIT1* 1487) were also up-regulated (*n* = 10). Cell wall genes up-regulated in filamentous

clade V (*n* = 8) included the novel adhesin *SCF1*, GPI-anchored genes *PGA31* and *IFF4*, and orthologues

of classical *C. albicans* hyphal adhesins *ALS3* and *RBR3*. We identified differentially expressed

transcription factors and related regulators, such as *HAP41* and the pheromone receptor *STE2*, which

could play a governing role on these responses.

We also examined significant DEGs between clade V and at least one other clade using mean-
log fold change (**Figure 7D**). Several genes that play critical roles in *C. albicans* hypha formation were
highly up-regulated, such as transcription factor *UME6* and hypha-specific G1 cyclin-dependent protein
serine/threonine kinase *HGC1*. Enrichment analysis indicated up-regulated gene sets involving CFEM
domain, cytochrome P450/oxidoreductase activity, hypha-regulated cell wall GPI-anchored proteins and
siderophore transport, particularly at 24 HPI and comparing clade V to clade II (**Figure 7D**). Significantly
up-regulated iron-related transport genes in clade V included *FRE3*, *CFL4*, *FRP1*, and *SIT1* genes 1487,
1499, 1519, 1542, and 4097. This is consistent with the preservation of a major filamentation programme
in *C. auris*, demonstrated in the basal clade V.

Given the prominence of *SIT1* orthologues across the transcriptomic analyses of *C. auris* during
in vivo gene expression, we set out to characterise the structural and evolutionary conservation of the
expanded gene family across orthogroups, including outgroup species *C. haemulonii* and *C. albicans*. We
performed phylogenetic analysis on proteins with sequence similarity across seven orthologue clusters, of
which two main groupings of transporters with 14 transmembrane domains (TMRs) emerged (**Figure S5**).
The first group included genes with functional annotation relating to siderophore import (including 12
genes in clade I). The second group included haem transmembrane import genes (including 5 genes in
clade I). Most of the orthologues retained 14 TMRs, and few had evidence of additional annotations
except for the trichothecene mycotoxin efflux pump PFAM domain across multiple groupings. The
orthologue *SIT1* 1547 contained 12 TMRs and was up-regulated with *SIT1* 1548 containing 2 TMRs,
which was not identified as a *SIT1* orthologue in our pipeline but has previously been described as
such[19,82].

To explore the evolutionary history of *SIT1*-related genes in fungi, we performed Blastp searches
for orthologues of *C. albicans SIT1* using an e-value cut-off of $1e^{-20}$ across 226 fungal species' reference
genomes (n = 304). We identified 1,689 hits, which clustered into two branches: a smaller branch
(n = 397) annotated for siderophore transport including the 12 *C. auris* groupings which we have
designated "xenosiderophore transporter candidates" (*XTC*), and a larger branch (n = 1,292) featuring
genes predicted (by GO term) to encode haem transporters, including the 5 *C. auris* groupings, which we
have additionally designated "haem transport-related" (*HTR*, **Figure 8A**). There were no putative haem
transporters or members of the larger branch for *C. albicans*, which only contained the single *SIT1* gene.
We calculated the mean number of orthogroups per genome per species (**Figure 8B**). This showed the
most prominent expansion of the *XTC* branch in the *Metschnikowiaceae* clade, including *C. auris* (12
orthologues) and even more so in *C. haemulonii*, with 15 orthologues. No other fungal species or group of

species contained so many. *XTC* orthologues were not present in any members of the basal

Blasocladiomycota, *Chytridiomycota* and *Mucoromycota*, or in several *Ascomycota* (e.g. *Pichia*

kudriavzevii, *Schizosaccharomyces pombe* and *Pyricularia oryzae*) and Basidiomycota (e.g. *Ustilago*

maydis). Common human fungal pathogens such as *Aspergillus fumigatus*, *C. albicans* and

Nakaseomyces glabratus contained only one *XTC* orthologue (**Figure 8B**). The *HTR* clade, by contrast,

was more prominent across fungal phyla, with as many as nine in *Mucor* species, 16 in *A. flavus* and 22

in *F. solani*. We therefore propose a naming of these two expanded gene families as *XTC1-12* and *HTR1-*

5 in *C. auris.*

We note the absence of *XTC5* in clade V, *XTC10* in clade IV, and in addition to these, *XTC4*,

XTC9 and *XTC11* in clade II (**Figure 8C**). Clade II, which exhibits the greatest degree of *XTC* gene loss,

is also considered by some to be the least virulent strain. It is also noteworthy that the AK embryo

expressed *HMOX* genes during clade IV infection, while clade IV *C. auris* up-regulated *HTR1*, *HTR2* and

HTR5. This suggests a specific host-pathogen interaction axis in nutritional immunity related to haem

scavenging. No other clades of infection were associated with such iron-related functional enrichment in

the host, and no other clades expressed more than one *HTR* gene. In summary, *XTC1*, *XTC2*, *XTC3*,

XTC4, *XTC6*, *XTC7*, *XTC11* and *XTC12* were up-regulated in-host while *XTC8*, *XTC9* and *XTC12* were

down-regulated in at least one clade, while up-regulation of *XTC3* and *XTC7* was significant across all

clades. Additionally, the more virulent clades were associated with up-regulation of *XTC4*, *XTC5*, and

XTC8. The filamentous clade V up-regulated *XTC7* across comparisons with all other clades, and up-

regulated *XTC1*, *XTC4*, *XTC8* and *XTC9* compared to at least one other clade.

Discussion

We describe the use of an AK yolk-sac microinjection model for comparative functional genomics

of *C. auris* during *in vivo* in-host infection. All five clades demonstrated lethality, with an increase in

virulence during clades I and IV infections, consistent with clinical and laboratory findings[17]. In-host

growth was similar across all clades and high enough to indicate a large enough *in vivo* fungal burden for

robust bulk transcriptome interrogation across the species. The contained nature of the yolk-sac provides

a host tissue growth medium that enables the assessment of pathogen survival and virulence strategies

in vivo and a straightforward experimental unit of host tissue without the challenges of pathogen RNA

isolation such as locating foci of infection in larger animals. Thus, an AK yolk-sac microinjection model

has the potential to fill a substantial need for *in vivo* gene expression studies in general.

We investigated the host transcriptional response to *C. auris* infection, which featured an innate

immune response, as expected. The most highly up-regulated genes included two predicted haem

oxygenase (*HMOX*) genes. *HMOX* genes have been shown to mask iron from invading pathogens as a
feature of anti-*Candida* nutritional immunity[83], so presumably have a similar function against *C. auris*. This
was particularly notable given *HMOX* genes were up-regulated in response to clade IV infection, and the
clade IV isolate had a corresponding significant up-regulation of three haem transport-related *HTR* genes.
In addition to nutritional immunity genes, we found up-regulated heat shock and complement cascade
genes during infection, confirming the expected immune response to fungal infection[63,64]. Genes with
complex roles in immunity were up-regulated, such as *NLRP3*, for which *C. auris* may be less activating
than e.g. *C. albicans*[33,84]. It is notable that the yolk-sac model was not associated with histological or
transcriptional evidence of host neutrophil recruitment and activation, which may relate to the hypothesis
that *C. auris* is immunoevasive[22]. Reference genome bias resulting from the available adult gill
transcriptome assembly[85] may limit detection of host gene expression features, so future long and short
read assembly from gene expression data across a range of larval stages could further understanding of
AK immunity. Additional time-points could be an informative future area of study to develop fine-resolution
strain-specific transcription networks. Furthermore, the potential of a translucent and genetically tractable
fish embryo model could enable future visualisation of cellular dynamics and genetic pathway
manipulation[77].
337
We provided a thorough investigation of differential expression by *C. auris* during infection of host
tissue compared with *in vitro* conditions for the first time. The lack of a major distinction across time-points
(24 *vs* 48 HPI) indicates that the transcriptional activity of *C. auris* in-host was highly similar at these
stages. Six functional annotation terms (GO, KEGG pathway and PFAM domain) were significantly
enriched among genes that were up-regulated across all five clades, including sugar and siderophore
transporters. The enrichment of sugar transporters outside of nutrient-rich YPD-broth is unsurprising,
such as high-affinity glucose transporter *HGT12*, with evidence of redundancy in *C. albicans*[86–89], and
HGT2, part of an eight-gene "core filamentation response network" in *C. albicans*[90]. More surprising was
the down-regulation of putative virulence factors such as Secreted Aspartyl Protease 3 (*SAP3*), thought to
be the most important SAP in *C. auris*, and required for virulence in a mouse model[31]. Copper Superoxide
Dismutase *SOD6* was also down-regulated, previously shown to be down-regulated in iron starvation[91].
Multi-drug efflux pumps such as *MDR2* (down-regulated *in vivo*) and *MDR1* (down-regulated *in vivo* and
in clades I and IV *vs* clades II and III) are understood to drive drug resistance through multiple but unclear
mechanisms in *C. auris*[92,93]; in-host down-regulation may indicate a fitness trade-off for virulence *vs* drug
resistance. A total of twelve siderophore transporters were up-regulated during infection, and four were
down-regulated, with three family members up-regulated in more virulent clades and five in the
filamentous clade V. The prominent finding of these xenosiderophore transport candidate (*XTC*) genes
necessitates further experimental work to address their potential for anti-virulence therapeutic targeting.
356

We also identified up-regulation of several genes at the mating type locus (*MTL*) in more virulent

strains; *MTL* genes may contribute to virulence in fungal species including *Cryptococcus neoformans*[94]

through uncertain mechanisms, such as *via* the expression of non-mating genes contained within the

locus. *C. auris* does not clearly undergo meiotic recombination[95] and it is not clear what role *MTL*-

associated genes play in pathogenesis and morphology. Two of the three non-mating genes (*PAP1* and

PIKA) in the *MTL* were up-regulated by more virulent strains (I and IV) compared to the less virulent

strains (II and III). Importantly, the two more virulent clades (I and IV) contain *MTL**a*** rather than *MTLα*

(clades II & III)[29], and clades V and VI contain *MTL**a*** and *MTLα* respectively[13]. Thus, it is unclear if the up-

regulation of non-mating genes is a *MTL* specific transcriptional nuance, or a feature of more highly

virulent *C. auris* clades. The *MTL* also did not feature prominently in the filamentation response observed

in clade V, providing no evidence for an obvious role for the *MTL* in such morphological changes.

We also observed *in vivo* filamentation by a representative strain of *C. auris* clade V, the

basal/ancestral clade[96], accompanied by a genera-typical filamentous gene expression signature.

Specifically, we measured the up-regulation of *HGC1* and *UME6*, which are two of the most important

regulators of filamentation in *C. albicans*, and have recently been described as drivers of *C. auris*

filamentation and biofilm formation[97]. The expression of adhesin *SCF1* in filamenting clade V is

noteworthy because its cationic adhesion mechanism has been postulated to be consistent with adhesion

mechanisms common to marine organisms including bivalves[27]. The generalist marine-isolated pathogen

C. haemulonii has also displayed filamentation competence at lower temperatures[98], consistent with an

ancestral aquatic state. Filamentation is an important pathogenicity trait in fungi, especially *Candida*

species, although its roles have been more poorly described in aquatic infections[99].

Human histopathology reports of *C. auris* infection are rarely published and have not

demonstrated classical in-host filamentation, to our knowledge. An *ex vivo* human epithelial model using

a clade I isolate did demonstrate filamentous *C. auris* at 37 °C after injection into deeper dermis[100]. Clade

III *C. auris* has also been observed to form filamentous/hyphal forms in *Galleria mellonella* injection[58]. Fan

and colleagues have observed filamentous forms of *C. auris* across all four clades, even demonstrating a

higher virulence in the *G. mellonella* model[101], though the phenotype required passage through a murine

host with additional prolonged growth at 25 °C for at least 5 d before and after injection[25]. Other stimuli

have induced filamentation in clade I isolates, such as hydroxyurea genotoxic stress[102] and Hsp90

inhibition[103], where several up-regulated genes in *C. auris* filamentation (*SCF1*, *IFF4* and *PGA31*)

matched those in our study. Constitutive filamentous growth has also been demonstrated with a

transposon-mediated disruption of a long non-coding RNA[104], which sit adjacent to chitin deacetylase

CDA2, which was down-regulated at 24 HPI in our experiment during filamentation. These findings

indicate that hyphal formation is possible across *C. auris* clades.

We underscored how the family of predicted xenosiderophore transporter genes is significantly

expanded in several members of the *Metschnikowiaceae* clade compared to other *Saccharomycetae*[29],

raising questions as to the roles of siderophore transporters in *C. auris* biology and evolution[105].

Siderophore synthesis may be unlikely in *C. auris*, given the lack of evidence in other *Candida* species[106].

As a result, *C. albicans* Sit1, which is required for virulence[107], is thought to be a xenosiderophore

transporter for iron-chelators produced by other microbes and specific to the ferrioxamine type common in

iron-poor ocean environments[108,109]. Siderophore transporters can also act intracellularly for

compartments such as the mitochondrion[110,111], or act as efflux pumps for various molecules[112] including

trichothecene mycotoxins as identified by our functional annotations. Given that *XTC* gene up-regulation

by *C. auris* has also been detected during *ex vivo* whole blood infection (*XTC2A-B*, *B9J08_1547-8*), this

suggests that our findings may be clinically relevant[19].

C. auris encodes a large expanded gene family of 17 siderophore transporters that we named

xenosiderophore transporter candidates (*XTC*) 1-12 and haem transport-related (*HTR*) genes 1-5, the

latter sharing sequence similarity with the genomic sequence for *S. pombe* low-affinity haem transporter

Str3[113]. In other *Candida* species such as *C. albicans*, iron assimilation is understood to be central for in-

host survival[114,115], where deprivation triggers various pathogenic programmes such as alterations in cell

surface beta-glucan[116]. Haem is acquired by *C. albicans* through vesicular endocytosis and CFEM-

domain containing proteins[117,118] and does not contain *HTR* genes, suggesting a possible loss of

transmembrane haem transporters. The ubiquity of *HTR* across the fungal kingdom and lack of *XTC*

genes in the three basal fungal phyla suggest that *SIT1*/*HTC* could have arisen from a duplication of an

HTR-related gene in the *Dikarya* (the *Ascomycota* and *Basidiomycota*) with subsequent independent loss

of both *XTC* and *HTC* genes in multiple fungal lineages. These findings together underline the importance

of iron metabolism as a clinical target with existing therapies, such as iron chelators

deferiprone/deferoxamine with evidence of echinocandin antifungal synergy[119] or low toxicity

antihelminthic pyrvinium pamoate as a disruptor of iron homeostasis[120,121].

We present the first *in vivo* host tissue infection transcriptional profile for *C. auris*, a World Health

Organization critical priority fungal pathogen. By examining representative strains from all five major

clades implicated in its recent, near-simultaneous global emergence, we uncover clade-specific

differences in morphology, virulence and gene expression. Our findings demonstrate the utility of a

thermotolerant yolk-sac injection model, and support the hypothesis of a marine origin for *C. auris*.

Notably, we observed natural filamentation in clade V and elevated virulence in clades I and IV, each

linked to distinct transcriptional programs. We identify the expanded *XTC* and *HTR* families as species-

wide, infection-induced features, underscoring their potential as targets for studying fungal biology,

pathogenesis, and antifungal development. Together, our results shed light on the mechanisms by which

C. auris adapts to and persists in host tissue, offering new insights into its evolution from a likely marine

yeast ancestor and advancing our understanding of this emerging fungal pathogen and global public

health threat.

Materials and methods

Full details are given in SI Appendix. Briefly, wild-type adult AK were kept at the Aquatic

Resources Facility, Exeter, under Project Licence PP4402521. AK were kept in two 28 °C incubation

tanks, with 12-hour day and night light cycles and artificial enrichment foliage. Embryos were collected

using three plastic netted collection chambers per tank. Injection needles were prepared using

borosilicate glass capillaries (Harvard apparatus, USA) and a heated pulling system (heater level >62,

PC-10, Narishige, UK). We used tweezers to snap the tip to an estimated 1 mm graticule-measured

aperture of 10-20 μm, and sterilised needles before use with 253.7 nm UV light over 15 min

(Environmental Validation Solutions, UK). Injection bays were made with microwaved 1% agarose

(Sigma, UK) in 1 g per 100 mL ASW for up to 24 embryos in 100 mm petri dishes using custom-made

molds in an inverted 60 mm petri dish. Embryos were randomly chosen from each dish and the sequence

of injection for each biological repeat was randomly allocated by computer. *C. auris* yeasts were counted

by haemocytometer and added to 25 μL sterile filtered phenol red solution 0.5% (Sigma, UK) to a

concentration of 0.5×10^8 per mL in 100 μL. Injection needles were loaded with 5 μL injection solution

using 20 μL Microloader EpT.I.P.S. (Eppendorf, UK) immediately before injection of up to 30 embryos

before re-loading. Injections were performed with 400-500 hPa injection pressure and 7-40 hPa

compensation pressure over 100-200 ms with the FemtoJet 4i (Eppendorf UK) using a micromanipulator

(World Precision Instruments, UK) and a Microtec HM-3 stereo dissection microscope (Optech, UK). To

deliver a dose of approx. 500 yeasts in 10 nL solution, an injection bolus was administered to an

estimated diameter of 0.2 – 0.3 mm (10-15% of total embryo diameter) *via* a single chorion puncture. Any

embryos that failed to meet these criteria for injection were excluded. Control *in vitro* inoculations were

microinjected as above into 1 mL YPD in 1.5 mL Eppendorf tubes before incubation without shaking at

37 °C for 24 h contemporaneously with embryos as below.

We used an unblinded prospective virulence study comparing six groups of AK embryo yolk-sac

microinjection, including clades I-V of *C. auris*, and sham injections as the negative control. We used ≥23

embryos per condition based on a power calculation of $\sigma = 0.25$, significance = 0.05, power = 0.9, and δ

effect size of 25% mortality (two-sided t-test). After excluding embryos that were unhealthy or did not

complete successful microinjection, we included $n = 10\text{-}13$ embryos per group per experiment (total

n = 34-37 embryos per group). RNA was extracted from three embryos per condition per time-point (0, 24

and 48 HPI, *n* = 54) plus three embryos per time-point without any injection (*n* = 9, total *n* = 63) as a

control for the impact of microinjection.

We selected five *C. auris* strains to represent each major clade (**Table S1**). Culture stocks were

kept in -80 °C in Yeast extract Peptone Dextrose (YPD) broth (Sigma-Aldrich, UK) containing 25%

glycerol (ThermoFisher, UK) and plated onto YPD-agar (Sigma-Aldrich) for 48 h before inoculation and

overnight growth in 10 mL YPD broth at 30 °C in a shaking incubator at 200 RPM (Multitron, Infors HT,

UK). Five mL from each overnight culture was then centrifuged at 5000 g for 2 min before triple washing

in 1 mL autoclaved deionised water (DIW, Milli-Q, Merck, USA) to prevent aggregation[28], then re-

suspended again in 200 μL DIW.

Individual embryos were kept in 1 mL sterile filtered ASW per embryo in a 48 well flat bottom

plate (Corning, USA) sealed with Parafilm "M" (Bemis Company, USA) at 37 °C with passive

humidification using a 500 mL open glass beaker of water. Embryos were monitored for presence of heart

rate at 24-hourly intervals up to 7 d by stereo dissection microscope. Initial time of injection was

subtracted from each time-point to minimise bias between groups. Kaplan-Meier survival curves were

plotted with Log-Rank testing and Benjamini-Hochberg (BH) correction[122] using survival v.3.5.8 and

survminer v.0.4.9 packages in R v.4.4.0. Colony forming units (CFUs) were calculated by homogenisation

of embryos at 0, 24, and 48 HPI by autoclaved plastic micropestle in a 1.5 mL conical tube (Eppendorf,

UK) in 100 μL 10,000 U/mL penicillin/streptomycin (P/S) solution (Gibco, UK).

Dilutions were made with P/S solution and DIW in 1/10-10,000 and counts were adjusted

according to dilution after plating onto YPD agar plates and read after 72 h incubation at 30 °C. Statistical

testing between CFU numbers was performed with Wilcoxon testing and BH correction.

Total RNA was extracted using the Monarch® Total RNA Miniprep Kit (NEB, USA). Embryos

were flash frozen in liquid nitrogen, re-suspended in DNA/RNA Protection Reagent and disrupted with an

autoclaved plastic micropestle. Bead beating was performed in a FastPrep 24 homogeniser (MP

Biomedicals) with 0.5 mm zirconia/silica beads for 8 cycles of 35 s at 6 m/s plus 45 s on ice. Lysate was

recovered and incubated for 5 min. at 55 °C with proteinase K before addition of lysis buffer and RNA

purification following manufacturer's instructions. Library preparation and sequencing were performed by

the Exeter Sequencing Service. Samples were normalised and prepared using the NEBNext Ultra II

Directional RNA Library Prep - Poly(A) mRNA Magnetic Isolation Module following the standard protocol.

Libraries were sequenced on a Novaseq 6000 sequencer with 150bp paired-end reads. Reads were

trimmed with fastp[123] v.0.23.1 to remove reads <75 bases and trim based from the 3' end with q-score

<22. Quality control was performed with FastQC (https://github.com/s-andrews/FastQC) v.0.11.9 and

MultiQC[124] v.1.6.

We aligned sequences (x = 52.9 m reads per replicate, range 41.1-112.2 m) using the Trinity

pipeline[125] v.2.15.1 to the *C. auris* clade I B8441 reference genome (GCA_002759435.2)[29] downloaded

from GenBank (mean aligned reads in infected embryos 3.83 m, 0.81-13.1%). We further aligned clade-

specific reference genomes for each strain (**Table S1**), including clade II (GCA_003013715.2)[4], III

(GCA_002775015.1)[14], IV (GCA_008275145.1)[14] and V (GCA_016809505.1)[96,126,127]. We aligned

conditions containing *A. dispar* to the *de novo* GJEY01 adult gill transcriptome assembly[85] downloaded

from GenBank, for which we detected expression over 27,481 potential genes. Alignment was performed

with Bowtie2[128] v.2.5.1, transcript estimation with RNA Seq by Expectation Maximisation (RSEM)[129]

v.1.3.3, differential expression with EdgeR[130] v.3.40.0, Samtools[131] v.1.17 and limma[132] v.3.54.0 using a

false discovery rate (FDR) p-value cut-off of 0.001 and a log fold change cut-off of +/-2.0. Principal

components were calculated and plotted using the prcomp function in Rv4.4.0 and ggfortify v.0.4.17 R

package for fragments per kilobase per million, and correlations plots from counts per million as per the

default output of the Trinity pipeline. Log fold change was plotted with ComplexHeatmap[133].

Gene ontology terms[134,135] were assigned using DIAMOND[136] v.2.1.8 *via* the DIAMOND2GO

pipeline[137] for each of the *C. auris* reference genomes. KEGG terms[138] were assigned using the

BlastKOALA online platform using the eukaryote database[139]. Protein FAMily (PFAM) domains[140] were

identified using the PFAM-A.hmm database

(https://ftp.ebi.ac.uk/pub/databases/Pfam/current_release/Pfam-A.full.gz) and HMMER and HMM Scan[141]

with a cut-off of 1e^{-5}. Further domain-specific identification was performed with SignalP 5.0[142],

DeepTMHMM 1.0[143] and NetGPI 1.1[144]. To identify accessory genes across *C. auris* clades, we used the

Synima pipeline[145] with Orthofinder v2.5.5[146] with additional reference genomes for *C. albicans*

(SC5314_GCA_000182965.3) and *C. haemulonii* (B11899_ASM1933202v1). Multiple alignment of single

copy orthologues as defined by Orthofinder was performed with Muscle v5.1[147] and a phylogeny was

made with FastTree v2.1.11[148]. We visualised the structure using Alphafold v3[149] and ChimeraX v1.8[150].

Clade I-V *C. auris* and *C. haemulonii SIT1* genes were aligned using Muscle as above before using

RAXML-NG v1.2.2 with 1000 bootstraps and the LG4X model[151], which was visualised using ggtree

v3.12.0.

Data availability: Raw reads have been deposited *via* NCBI/GEO *via* accession GSE277854. All

datasets and code required to make figures are available *via* Github

(https://github.com/hughgifford/Arabian_Killifish_C_auris_2024).

Acknowledgements

We acknowledge funding from the MRC Centre for Medical Mycology at the University of Exeter (MR/N006364/2, MR/V033417/1), MRC Doctoral Training Grants MR/P501955/2 and MR/W502649/1, Wellcome Trust (206412/A/17/Z), Wellcome Trust Career Development Award (215239/Z/19/Z) and Wellcome Trust Fellowship (219551/Z/19/Z), National Centre for the Replacement, Refinement and Reduction of Animals in Research (NC3Rs) (NC/X001121/1), and the NIHR Exeter Biomedical Research Centre. TB also was supported *via* Elizabeth Ballou and a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society (211241/Z/18/Z) and the Lister Institute. The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care. We thank staff in the Aquatic Resource Center at the University of Exeter for husbandry of the fish and the support for developing protocols for feeding, spawning, and raising new generations of this relatively new model species, including Sam Worthington, Chloe Flint and Richard Silcox. We also thank the Exeter Sequencing Service facility and support from Wellcome Trust Institutional Strategic Support Fund (WT097835MF), Wellcome Trust Multi User Equipment Awards (WT101650MA and 218247/Z/19/Z), Medical Research Council Clinical Infrastructure Funding (MR/M008924/1) and BBSRC LOLA award (BB/K003240/1). We also thank the University of Exeter High-Performance Computing (HPC) facility funded by the UK MRC Clinical Research Infrastructure Initiative (award number MR/M008924/1). We thank Tanmoy Chakraborty, Dhara Malavia and Neil Gow for providing *C. auris* strains, Teigan Veale, Sumita Roy, Jonathan Ball and Darren Thomson for assistance in microscopy and Chloe Pelletier for advice on aggregation. We also thank Microtechnical Services, Exeter, UK, for assistance with histological sectioning.

References

1.   Meis, J. F. & Chowdhary, A. *Candida auris*: A global fungal public health threat. *The Lancet Infectious Diseases* **18**, 1298–1299 (2018).

2.   WHO. World Health Organisation (WHO) fungal priority pathogens list to guide research, development and public health action. (2022) doi:https://www.who.int/publications/i/item/9789240060241.

3.   Lee, W. G. *et al.* First three reported cases of nosocomial fungemia caused by *Candida auris*. *Journal of Clinical Microbiology* **49**, 4 (2011).

4.   Satoh, K. *et al. Candida auris* sp. Nov., a novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese hospital. *Microbiology and Immunology* **53**, 41–44 (2009).

5.   Eyre, D. W. *et al.* A *Candida auris* outbreak and its control in an intensive care setting. *New England Journal of Medicine* **379**, 1322–1331 (2018).

6.   Taori, S. K. *et al. Candida auris* outbreak: Mortality, interventions and cost of sustaining control. *Journal of Infection* **79**, 601–611 (2019).

7.   Shuping, L. *et al.* High prevalence of *Candida auris* colonization during protracted neonatal unit
outbreak, South Africa. *Emerging Infectious Diseases* (2023) doi:10.3201/eid2909.230393.

8.   Adam, R. D. *et al.* Analysis of *Candida auris* fungemia at a single facility in Kenya. *International*
Journal of Infectious Diseases **85**, 182–187 (2019).

9.   Prayag, P. S., Patwardhan, S., Panchakshari, S., Rajhans, P. A. & Prayag, A. The dominance of
Candida auris: A single-center experience of 79 episodes of candidemia from western India. *Indian*
Journal of Critical Care Medicine **26**, 560–563 (2022).

10.   Chen, J. *et al.* Is the superbug fungus really so scary? A systematic review and meta-analysis of
global epidemiology and mortality of *Candida auris*. *BMC Infectious Diseases* **20**, 827 (2020).

11.   Lyman, M. Notes from the field: Transmission of pan-resistant and echinocandin-resistant
Candida auris in health care facilities - Texas and the District of Columbia, January-April 2021. *Morbidity*
and Mortality Weekly Report **70**, (2021).

12.   Khan, T. *et al.* Emergence of the novel sixth *Candida auris* clade VI in Bangladesh. *Microbiology*
Spectrum **12**, e03540–23 (2024).

13.   Suphavilai, C. *et al.* Detection and characterisation of a sixth *Candida auris* clade in Singapore: A
genomic and phenotypic study. *Lancet Microbe* **0**, (2024).

14.   Lockhart, S. R. *et al.* Simultaneous emergence of multidrug-resistant *Candida auris* on 3
continents confirmed by whole-genome sequencing and epidemiological analyses. *Clinical Infectious*
Diseases **64**, 134–140 (2017).

15.   Gifford, H., Rhodes, J. & Farrer, R. A. The diverse genomes of *Candida auris. The Lancet*
Microbe **0**, (2024).

16.   Byun, S. A. *et al.* Virulence traits and azole resistance in Korean *Candida auris* Isolates. *Journal*
of Fungi **9**, 979 (2023).

17.   Santana, D. J., Zhao, G. & O'Meara, T. R. The many faces of *Candida auris*: Phenotypic and
strain variation in an emerging pathogen. *PLOS Pathogens* **20**, e1012011 (2024).

18.   Wang, T. W. *et al.* Functional redundancy in *Candida auris* cell surface adhesins crucial for cell-
cell interaction and aggregation. *Nature Communications* **15**, 9212 (2024).

19.   Allert, S. *et al.* From environmental adaptation to host survival: Attributes that mediate
pathogenicity of *Candida auris*. *Virulence* **13**, 191–214 (2022).

20.   Fayed, B. *et al.* Transcriptome analysis of human dermal cells infected with *Candida auris*
identified unique pathogenesis/defensive mechanisms particularly ferroptosis. *Mycopathologia* **189**, 65
(2024).

21.   Yang, B. *et al.* A correlative study of the genomic underpinning of virulence traits and drug
tolerance of *Candida auris*. *Infection and Immunity* **0**, e00103–24 (2024).

22.   Balakumar, A. *et al.* Single-cell transcriptomics unveils skin cell specific antifungal immune
responses and IL-1Ra- IL-1R immune evasion strategies of emerging fungal pathogen *Candida auris*.
PLOS Pathogens **20**, e1012699 (2024).

23.   Bruno, M. *et al.* Transcriptional and functional insights into the host immune response against the
emerging fungal pathogen *Candida auris*. *Nature Microbiology* **5**, 1516–1531 (2020).

24.   Wang, Y. *et al.* Innate immune responses against the fungal pathogen *Candida auris*. *Nature*
Communications **13**, 3553 (2022).

25. Yue, H. *et al.* Filamentation in *Candida auris*, an emerging fungal pathogen of humans: Passage
through the mammalian body induces a heritable phenotypic switch. *Emerging Microbes & Infections* **7**,
1–13 (2018).

26. Chadwick, C., Jesus, M. D., Ginty, F. & Martinez, J. S. Pathobiology of *Candida auris* infection
analyzed by multiplexed imaging and single cell analysis. *PLOS ONE* **19**, e0293011 (2024).

27. Santana, D. J. *et al.* A *Candida auris*-specific adhesin, Scf1, governs surface association,
colonization, and virulence. *Science* **381**, 1461–1467 (2023).

28. Pelletier, C., Shaw, S., Alsayegh, S., Brown, A. J. P. & Lorenz, A. *Candida auris* undergoes
adhesin-dependent and -independent cellular aggregation. *PLOS Pathogens* **20**, e1012076 (2024).

29. Muñoz, J. F. *et al.* Genomic insights into multidrug-resistance, mating and virulence in *Candida
auris* and related emerging species. *Nature Communications* **9**, 5346 (2018).

30. Richardson, J. P. *et al.* Candidalysins are a new family of cytolytic fungal peptide toxins. *mBio* **13**,
e03510–21 (2022).

31. Kim, J.-S., Lee, K.-T. & Bahn, Y.-S. Secreted aspartyl protease 3 regulated by the
Ras/cAMP/PKA pathway promotes the virulence of *Candida auris*. *Frontiers in Cellular and Infection
Microbiology* **13**, 1257897 (2023).

32. Horton, M. V. *et al. Candida auris* cell wall mannosylation contributes to neutrophil evasion
through pathways divergent from *Candida albicans* and *Candida glabrata*. *mSphere* **6**, e00406–21
(2021).

33. Weerasinghe, H. *et al. Candida auris* uses metabolic strategies to escape and kill macrophages
while avoiding robust activation of the NLRP3 inflammasome response. *Cell Reports* **42**, 112522 (2023).

34. Wang, X. *et al.* The first isolate of *Candida auris* in China: Clinical and biological aspects.
Emerging Microbes & Infections **7**, 1–9 (2018).

35. Fan, S., Li, C., Bing, J., Huang, G. & Du, H. Discovery of the diploid form of the emerging fungal
pathogen *Candida auris*. *ACS Infectious Diseases* **6**, 2641–2646 (2020).

36. Fan, S. *et al.* A biological and genomic comparison of a drug-resistant and a drug-susceptible
strain of *Candida auris* isolated from Beijing, China. *Virulence* **12**, 1388–1399 (2021).

37. Wurster, S., Albert, N. D. & Kontoyiannis, D. P. *Candida auris* bloodstream infection induces
upregulation of the PD-1/PD-L1 immune checkpoint pathway in an immunocompetent mouse model.
mSphere **7**, e00817–21 (2022).

38. Bing, J. *et al.* Rapid evolution of an adaptive multicellular morphology of *Candida auris* during
systemic infection. *Nature Communications* **15**, 2381 (2024).

39. Ben-Ami, R. *et al.* Multidrug-resistant *Candida haemulonii* and *C. auris*, Tel Aviv, Israel. *Emerging
Infectious Diseases* **23**, (2017).

40. Hager, C. L., Larkin, E. L., Long, L. A. & Ghannoum, M. A. Evaluation of the efficacy of
rezafungin, a novel echinocandin, in the treatment of disseminated *Candida auris* infection using an
immunocompromised mouse model. *Journal of Antimicrobial Chemotherapy* **73**, 2085–2088 (2018).

41. Zhao, M. *et al. In vivo* pharmacokinetics and pharmacodynamics of APX001 against *Candida
spp.* In a neutropenic disseminated candidiasis mouse model. *Antimicrobial Agents and Chemotherapy
62, e02542–17 (2018).

42. Singh, S. *et al.* The NDV-3A vaccine protects mice from multidrug resistant *Candida auris*
infection. *PLOS Pathogens* **15**, e1007460 (2019).

43.   Zhang, F. *et al.* A marine microbiome antifungal targets urgent-threat drug-resistant fungi.
Science **370**, 974–978 (2020).

44.   Muñoz, J. E. *et al.* Pathogenicity levels of Colombian strains of *Candida auris* and Brazilian
strains of *Candida haemulonii* species complex in both murine and *Galleria mellonella* experimental
models. *Journal of Fungi* **6**, 104 (2020).

45.   Abe, M. *et al.* Potency of gastrointestinal colonization and virulence of *Candida auris* in a murine
endogenous candidiasis. *PLoS One* **15**, e0243223 (2020).

46.   Forgács, L. *et al.* Comparison of in vivo pathogenicity of four *Candida auris* clades in a
neutropenic bloodstream infection murine model. *Emerging Microbes & Infections* **9**, 1160–1169 (2020).

47.   Nagy, F. *et al.* In vitro and in vivo interaction of caspofungin with isavuconazole against *Candida
auris* planktonic cells and biofilms. *Medical Mycology* **59**, 1015–1023 (2021).

48.   Forgács, L. *et al.* In vivo efficacy of amphotericin B against four *Candida auris* clades. *Journal of
Fungi (Basel, Switzerland)* **8**, 499 (2022).

49.   Singh, S. *et al.* Protective efficacy of anti-Hyr1p monoclonal antibody against systemic
candidiasis due to multi-drug-resistant *Candida auris*. *Journal of Fungi* **9**, 103 (2023).

50.   Salama, E. A. *et al.* Lansoprazole interferes with fungal respiration and acts synergistically with
amphotericin B against multidrug-resistant *Candida auris*. *Emerging Microbes & Infections* **13**, 2322649
(2024).

51.   Torres, S. R. *et al.* Impact of *Candida auris* infection in a neutropenic murine model. *Antimicrobial
Agents and Chemotherapy* **64**, e01625–19 (2020).

52.   Xin, H., Mohiuddin, F., Tran, J., Adams, A. & Eberle, K. Experimental mouse models of
disseminated *Candida auris* infection. *mSphere* **4**, e00339–19 (2019).

53.   Rosario-Colon, J., Eberle, K., Adams, A., Courville, E. & Xin, H. *Candida* cell-surface-specific
monoclonal antibodies protect mice against *Candida auris* invasive infection. *International Journal of
Molecular Sciences* **22**, 6162 (2021).

54.   Torres, S. R. *et al.* Assessment of environmental and occupational exposure while working with
multidrug resistant (MDR) fungus *Candida auris* in an animal facility. *Journal of Occupational and
Environmental Hygiene* **16**, 507–518 (2019).

55.   Borman, A. M. Of mice and men and larvae: *Galleria mellonella* to model the early host-pathogen
interactions after fungal infection. *Virulence* **9**, 9–12 (2017).

56.   Romera, D. *et al.* The *Galleria mellonella* infection model as a system to investigate the virulence
of *Candida auris* strains. *Pathogens and Disease* **78**, ftaa067 (2020).

57.   García-Carnero, L. C. *et al.* Early virulence predictors during the *Candida* species–*Galleria
mellonella* interaction. *Journal of Fungi* **6**, 152 (2020).

58.   Garcia-Bustos, V. *et al.* Characterization of the differential pathogenicity of *Candida auris* in a
Galleria mellonella infection model. *Microbiology Spectrum* **9**, e00013–21 (2021).

59.   Carvajal, S. K. *et al.* Pathogenicity assessment of Colombian strains of *Candida auris* in the
Galleria mellonella invertebrate model. *Journal of Fungi* **7**, 401 (2021).

60.   Hernando-Ortiz, A. *et al.* Virulence of *Candida auris* from different clinical origins in
Caenorhabditis elegans and *Galleria mellonella* host models. *Virulence* **12**, 1063–1075 (2021).

61.   Spettel, K. *et al. Candida auris* in Austria—what is new and what is different. *Journal of Fungi* **9**,
129 (2023).

62.   Wurster, S. *et al. Drosophila melanogaster* as a model to study virulence and azole treatment of
the emerging pathogen *Candida auris. Journal of Antimicrobial Chemotherapy* **74**, 1904–1910 (2019).

63.   Harpf, V., Rambach, G., Würzner, R., Lass-Flörl, C. & Speth, C. *Candida* and complement: New
aspects in an old battle. *Frontiers in Immunology* **11**, 1471 (2020).

64.   Pechacek, J. & Lionakis, M. S. Host defense mechanisms against *Candida auris. Expert Review*
of Anti-infective Therapy **21**, 1087–1096 (2023).

65.   Johnson, C. J., Davis, J. M., Huttenlocher, A., Kernien, J. F. & Nett, J. E. Emerging fungal
pathogen *Candida auris* evades neutrophil attack. *mBio* **9**, e01403–18 (2018).

66.   Pharkjaksu, S., Boonmee, N., Mitrpant, C. & Ngamskulrungroj, P. Immunopathogenesis of
emerging *Candida auris* and *Candida haemulonii* strains. *Journal of Fungi* **7**, 725 (2021).

67.   Shapiro, R. S. & Cowen, L. E. Uncovering cellular circuitry controlling temperature-dependent
fungal morphogenesis. *Virulence* **3**, 400–404 (2012).

68.   Leach, M. D. *et al.* Hsf1 and Hsp90 orchestrate temperature-dependent global transcriptional
remodelling and chromatin architecture in *Candida albicans. Nature Communications* **7**, 11704 (2016).

69.   Xiao, W. *et al.* Response and regulatory mechanisms of heat resistance in pathogenic fungi.
Applied Microbiology and Biotechnology **106**, 5415–5431 (2022).

70.   Arora, P. *et al.* Environmental isolation of *Candida auris* from the coastal wetlands of Andaman
Islands, India. *mBio* **12**, e03181–20 (2021).

71.   Escandón, P. Novel environmental niches for *Candida auris*: Isolation from a coastal habitat in
Colombia. *Journal of Fungi* **8**, 748 (2022).

72.   Uden, V. & Kolipinski, M. C. *Torulopsis haemulonii nov. spec.*, A yeast from the Atlantic Ocean.
Antonie Van Leeuwenhoek **28**, 78–80 (1962).

73.   Bavia, L., Santiesteban-Lores, L. E., Carneiro, M. C. & Prodocimo, M. M. Advances in the
complement system of a teleost fish, *Oreochromis niloticus. Fish & Shellfish Immunology* **123**, 61–74
(2022).

74.   Buchmann, K., Karami, A. M. & Duan, Y. The early ontogenetic development of immune cells and
organs in teleosts. *Fish & Shellfish Immunology* **146**, 109371 (2024).

75.   Tian, H. *et al.* Cytokine networks provide sufficient evidence for the differentiation of CD4+ T cells
in teleost fish. *Developmental & Comparative Immunology* **141**, 104627 (2023).

76.   Akbarzadeh, A. & Leder, E. H. Acclimation of killifish to thermal extremes of hot spring:
Transcription of gonadal and liver heat shock genes. *Comparative Biochemistry and Physiology Part A:*
Molecular & Integrative Physiology **191**, 89–97 (2016).

77.   Hamied, A. *et al.* Identification and characterization of highly fluorescent pigment cells in embryos
of the Arabian killifish (*Aphanius Dispar*). *iScience* **23**, 101674 (2020).

78.   Alsakran, A. *et al.* Stage-by-stage exploration of normal embryonic development in the Arabian
killifish, *Aphanius dispar. Developmental Dynamics* **n/a**, (2024).

79.   Minhas, R. *et al.* The thermotolerant Arabian killifish, *Aphanius dispar,* as a novel infection model
for human fungal pathogens. 2024.10.08.617174 (2024) doi:10.1101/2024.10.08.617174.

80. Gifford, H., Wilson, D., Rhodes, J. & Farrer, R. A. Seaside to bedside: Assembly in research for
emerging human fungal pathogen *Candida auris*. in *Genome Assembly* (ed. Farrer, R. A.) vol. 2955 263–
291 (Springer US, New York, NY, 2025).

81. Burden, N., Chapman, K., Sewell, F. & Robinson, V. Pioneering better science through the 3Rs:
An introduction to the national centre for the replacement, refinement, and reduction of animals in
research (NC3Rs). *Journal of the American Association for Laboratory Animal Science* **54**, 198–208
(2015).

82. Skrzypek, M. S. *et al.* The *Candida* Genome Database (CGD): Incorporation of Assembly 22,
systematic identifiers and visualization of high throughput sequencing data. *Nucleic Acids Research* **45**,
D592–D596 (2017).

83. Potrykus, J. *et al.* Fungal iron availability during deep seated candidiasis is defined by a complex
interplay involving systemic and local events. *PLOS pathogens* **9**, e1003676 (2013).

84. Kasper, L. *et al.* The fungal peptide toxin candidalysin activates the NLRP3 inflammasome and
causes cytolysis in mononuclear phagocytes. *Nature Communications* **9**, 4260 (2018).

85. Bonzi, L. C. *et al.* The time course of molecular acclimation to seawater in a euryhaline fish.
Scientific Reports **11**, 18127 (2021).

86. Varma, A. *et al.* Molecular cloning and functional characterisation of a glucose transporter,
CaHGT1, of *Candida albicans*. *FEMS microbiology letters* **182**, 15–21 (2000).

87. Luo, L., Tong, X. & Farley, P. C. The *Candida albicans* gene *HGT12* (Orf19.7094) encodes a
hexose transporter. *FEMS Immunology & Medical Microbiology* **51**, 14–17 (2007).

88. Brown, V., Sexton, J. A. & Johnston, M. A glucose sensor in Candida albicans. *Eukaryotic Cell* **5**,
1726–1737 (2006).

89. Larcombe, D. E. *et al.* Glucose-enhanced oxidative stress resistance-A protective anticipatory
response that enhances the fitness of *Candida albicans* during systemic infection. *PLOS Pathogens* **19**,
e1011505 (2023).

90. Martin, R. *et al.* A core filamentation response network in *Candida albicans* is restricted to eight
genes. *PLOS ONE* **8**, e58613 (2013).

91. Schatzman, S. S. *et al.* Copper-only superoxide dismutase enzymes and iron starvation stress in
Candida fungal pathogens. *The Journal of Biological Chemistry* **295**, 570–583 (2020).

92. Rybak, J. M. *et al.* Abrogation of triazole resistance upon deletion of *CDR1* in a clinical isolate of
Candida auris. *Antimicrobial Agents and Chemotherapy* **63**, (2019).

93. Li, J., Coste, A. T., Bachmann, D., Sanglard, D. & Lamoth, F. Deciphering the Mrr1/Mdr1 pathway
in azole resistance of *Candida auris*. *Antimicrobial Agents and Chemotherapy* **66**, e00067–22 (2022).

94. Sun, S., Coelho, M. A., David-Palma, M., Priest, S. J. & Heitman, J. The evolution of sexual
reproduction and the mating-type locus: Links to pathogenesis of *Cryptococcus* human pathogenic fungi.
Annual Review of Genetics **53**, 417–444 (2019).

95. Wang, Y. & Xu, J. Population genomic analyses reveal evidence for limited recombination in the
superbug *Candida auris* in nature. *Computational and Structural Biotechnology Journal* **20**, 3030–3040
(2022).

96. Chow, N. A. *et al.* Potential fifth clade of *Candida auris,* Iran, 2018. *Emerging Infectious Diseases*
25, 1780–1781 (2019).

97.   Louvet, M. *et al.* Ume6-dependent pathways of morphogenesis and biofilm formation in *Candida*
auris. *Microbiology Spectrum* **0**, e01531–24 (2024).

98.   Deng, Y., Li, S., Bing, J., Liao, W. & Tao, L. Phenotypic switching and filamentation in *Candida*
haemulonii, an emerging opportunistic pathogen of humans. *Microbiology Spectrum* **9**, e00779–21
(2021).

99.   Overy, D. P., Rämä, T., Oosterhuis, R., Walker, A. K. & Pang, K.-L. The neglected marine fungi,
Sensu stricto, and their isolation for natural products' discovery. *Marine Drugs* **17**, 42 (2019).

100.   Seiser, S. *et al.* Native human and mouse skin infection models to study *Candida auris*-host
interactions. *Microbes and Infection* **26**, 105234 (2024).

101.   Fan, S. *et al.* Filamentous growth is a general feature of *Candida auris* clinical isolates. *Medical*
Mycology **59**, 734–740 (2021).

102.   Bravo Ruiz, G., Ross, Z. K., Gow, N. A. R. & Lorenz, A. Pseudohyphal growth of the emerging
pathogen *Candida auris* is triggered by genotoxic stress through the S phase checkpoint. *mSphere* **5**,
e00151–20 (2020).

103.   Kim, S. H. *et al.* Genetic analysis of *Candida auris* implicates Hsp90 in morphogenesis and azole
tolerance and Cdr1 in Azole Resistance. *mBio* **10**, e02529–18 (2019).

104.   Gao, J. *et al.* LncRNA DINOR is a virulence factor and global regulator of stress responses in
Candida auris. *Nature Microbiology* **6**, 842–851 (2021).

105.   Chybowska, A. D., Childers, D. S. & Farrer, R. A. Nine things genomics can tell us about *Candida*
auris. *Frontiers in Genetics* **11**, 351 (2020).

106.   Ismail, A., Bedell, G. W. & Lupan, D. M. Siderophore production by the pathogenic yeast,
Candida albicans. *Biochemical and Biophysical Research Communications* **130**, 885–891 (1985).

107.   Heymann, P. *et al.* The Siderophore iron transporter of *Candida albicans* (Sit1p/Arn1p) mediates
uptake of ferrichrome-type siderophores and Is required for epithelial invasion. *Infection and Immunity* **70**,
5246–5255 (2002).

108.   Boiteau, R. M. *et al.* Siderophore-based microbial adaptations to iron scarcity across the eastern
Pacific Ocean. *Proceedings of the National Academy of Sciences* **113**, 14237–14242 (2016).

109.   Park, J. *et al.* Siderophore production and utilization by marine bacteria in the North Pacific
Ocean. *Limnology and Oceanography* **68**, 1636–1653 (2023).

110.   Renshaw, J. C. *et al.* Fungal siderophores: Structures, functions and applications. *Mycological*
Research **106**, 1123–1142 (2002).

111.   Johnstone, T. C. & Nolan, E. M. Beyond iron: Non-classical biological functions of bacterial
siderophores. *Dalton Transactions* **44**, 6320–6339 (2015).

112.   Nakajima, Y. *et al.* Effect of disrupting the trichothecene efflux pump encoded by *FgTri12* in the
nivalenol chemotype of *Fusarium graminearum*. *The Journal of General and Applied Microbiology* **61**, 93–
96 (2015).

113.   Xue, P. *et al.* Heme sensing and trafficking in fungi. *Fungal Biology Reviews* **43**, 100286 (2023).

114.   Fourie, R., Kuloyo, O. O., Mochochoko, B. M., Albertyn, J. & Pohl, C. H. Iron at the Centre of
Candida albicans Interactions. *Frontiers in Cellular and Infection Microbiology* **8**, 185 (2018).

115.   Alselami, A. & Drummond, R. A. How metals fuel fungal virulence, yet promote anti-fungal
immunity. *Disease Models & Mechanisms* **16**, dmm050393 (2023).

116.   Pradhan, A. *et al.* Non-canonical signalling mediates changes in fungal cell wall PAMPs that drive
immune evasion. *Nature Communications* **10**, 5315 (2019).

117.   Weissman, Z. & Kornitzer, D. A family of *Candida* cell surface haem-binding proteins involved in
haemin and haemoglobin-iron utilization. *Molecular Microbiology* **53**, 1209–1220 (2004).

118.   Ding, C. *et al.* Conserved and divergent roles of Bcr1 and CFEM proteins in *Candida parapsilosis*
and Candida albicans. *PLOS ONE* **6**, e28151 (2011).

119.   Sharma, P. & Pasrija, R. Iron chelators enhance the potency of Echinocandins against *Candida*
auris. (2025) doi:10.21203/rs.3.rs-7042978/v1.

120.   Simm, C. *et al.* Disruption of iron homeostasis and mitochondrial metabolism are promising
targets to inhibit *Candida auris*. *Microbiology Spectrum* **10**, e00100–22 (2022).

121.   Zhang, J. *et al.* Pyrvinium pamoate synergizes with azoles *in vitro* and *in vivo* to exert antifungal
efficacy against *Candida auris* and other *Candida* species. *Infection and Drug Resistance* **18**, 783–789
(2025).

122.   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical and powerful
approach to multiple testing. *Journal of the Royal Statistical Society Series B: Statistical Methodology* **57**,
289–300 (1995).

123.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. Fastp: An ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

124.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: Summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

125.   Haas, B. J. *et al.* De novo transcript sequence reconstruction from RNA-seq using the Trinity
platform for reference generation and analysis. *Nature Protocols* **8**, 1494–1512 (2013).

126.   Abastabar, M. *et al.* *Candida auris* otomycosis in Iran and review of recent literature. *Mycoses* **62**,
101–105 (2019).

127.   Muñoz, J. F. *et al.* Clade-specific chromosomal rearrangements and loss of subtelomeric
adhesins in *Candida auris*. *Genetics* **218**, iyab029 (2021).

128.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature Methods* **9**,
357–359 (2012).

129.   Li, B. & Dewey, C. N. RSEM: Accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

130.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

131.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079
(2009).

132.   Ritchie, M. E. *et al.* Limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Research* **43**, e47 (2015).

133.   Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

134.   Ashburner, M. *et al.* Gene Ontology: Tool for the unification of biology. *Nature Genetics* **25**, 25–29
(2000).

135. The Gene Ontology Consortium *et al.* The Gene Ontology knowledgebase in 2023. *Genetics* **224**,
iyad031 (2023).

136. Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life scale using
DIAMOND. *Nature Methods* **18**, 366–368 (2021).

137. Golden, C., Studholme, D. J. & Farrer, R. A. DIAMOND2GO: A rapid Gene Ontology assignment
and enrichment tool for functional genomics. 2024.08.19.608700 (2024) doi:10.1101/2024.08.19.608700.

138. Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M. & Ishiguro-Watanabe, M. KEGG for
taxonomy-based analysis of pathways and genomes. *Nucleic Acids Research* **51**, D587–D592 (2022).

139. Kanehisa, M., Sato, Y. & Morishima, K. BlastKOALA and GhostKOALA: KEGG tools for functional
characterization of genome and metagenome sequences. *Journal of Molecular Biology* **428**, 726–731
(2016).

140. Bateman, A. *et al.* The Pfam protein families database. *Nucleic Acids Research* **32**, D138–D141
(2004).

141. Eddy, S. R. Accelerated profile HMM searches. *PLOS Computational Biology* **7**, e1002195
(2011).

142. Almagro Armenteros, J. J. *et al.* SignalP 5.0 improves signal peptide predictions using deep
neural networks. *Nature Biotechnology* **37**, 420–423 (2019).

143. Hallgren, J. *et al.* DeepTMHMM predicts alpha and beta transmembrane proteins using deep
neural networks. (2022) doi:10.1101/2022.04.08.487609.

144. Gíslason, M. H., Nielsen, H., Almagro Armenteros, J. J. & Johansen, A. R. Prediction of GPI-
anchored proteins with pointer neural networks. *Current Research in Biotechnology* **3**, 6–13 (2021).

145. Farrer, R. A. Synima: A synteny imaging tool for annotated genome assemblies. *BMC
Bioinformatics* **18**, 507 (2017).

146. Emms, D. M. & Kelly, S. OrthoFinder: Phylogenetic orthology inference for comparative
genomics. *Genome Biology* **20**, 238 (2019).

147. Edgar, R. C. MUSCLE: Multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Research **32**, 1792–1797 (2004).

148. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – approximately maximum-likelihood trees for
large alignments. *PLOS ONE* **5**, e9490 (2010).

149. Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with AlphaFold 3.
Nature **630**, 493–500 (2024).

150. Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. *Protein Science* **30**, 70–82 (2021).

151. Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

Figures and tables

Figure 1: Modelling *C. auris* infection in the Arabian killifish **(A)** Illustrative time-lapse microscopy of embryo death including yolk-sac collapse during infection (clade I infection) or survival (sham injection) using Acquifer live imaging. **(B)** Survival curves for embryos injected with each *C. auris* clade or sham injection. Significance testing between groups is given in **Table S1**; hours post injection (HPI). **(C)** H&E Staining of AK yolk-sac at 48 HPI for each clade or sham injection; morphology is indicated by blue arrows, with insets indicating typical yeast (clade III) and filamentous morphology (clade V). The yolk-sac edge is oriented at the top. **(D)** Colony forming units (CFUs) per clade from embryos homogenised after injection as an indicator of injection dose and *in vivo* growth.

Figure 2: Indicative pipeline diagram for methods used. At three days post fertilization (DPF), *A. dispar* were microinjected with *C. auris* from each of five clades I-V. Transcriptional comparisons were made between host and pathogen conditions using either a shared or clade-specific reference. Functional annotations were generated to ascertain enriched features. Illustrations © the authors.

Figure 3: AK host responses to *C. auris* infection: **(A)** Number of differentially expressed genes (DEGs) for each of two comparisons, sham injection *vs* no injection (left panel), and *C. auris* injection *vs* sham injection (right panel), at either 24 or 48 h post injection (HPI). **(B)** Volcano plot showing log-fold change

Figure 4: *C. auris* gene expression *in vivo vs in vitro* **(A)** Total DEGs per clade for each comparison between 24 h post injection (HPI), 48 HPI, and growth in YPD alone. **(B)** Combined set of significant DEGs common to all five clades between *in vivo* infection and *in vivo* growth in YPD. Genes are coloured by manually annotated category (**Table S3**). **(C)** Volcano plots demonstrate mean log-fold change for DEGs identified for any clade. Genes with infinite false discovery rate (FDR) are indicated with an arrow. Genes included in the combined set are coloured by category as per panel **B**.

Figure 5: *C. auris* gene expression *in vivo vs in vitro*. **(A-C)** Upset plots for each *in vivo vs in vitro* comparison between 24 h post injection (HPI), 48 HPI, and growth in YPD alone, indicating DEGs that were significant across all five clades (far left) and for each individual clade (far right) and each set in between.

Figure 6: *C. auris* gene set, pathway and domain enrichment: Transcripts from *C. auris* reference

Figure 7: Interclade comparisons for virulence and filamentation: **(A):** The set of differentially expressed genes (DEGs) shared across all comparisons between clade I and IV (more virulent) and clade II and III (less virulent), coloured by manually annotated category, at either 24 or 48 h post infection (HPI). **(B)** Volcano plots demonstrate mean log-fold change for each DEG found in any comparison; genes included in the combined set are coloured as per **A**. **(C)** The set of DEGs shared across all comparisons between filamenting clade V all other clades (I-IV) *in vivo*, coloured by manually annotated category. **(D):** Volcano plots demonstrating mean log-fold change for all comparisons between clade V and clades I-IV at 24 HPI

and 48 HPI for each DEG found in any comparison; genes included in the combined set are coloured as per **C**.

Proposed name	Shortened name	B9441 v2 code	B8441 v3 code	B8441 v3 contig	Chromosome
XTC_1	*SIT1_1542*	B9J08_001542	B9J08_00013	CM076438.1	1
XTC_2A	*SIT1_1547*	B9J08_001547	B9J08_00018	CM076438.1	1
XTC_2B	*SIT1_1548*	B9J08_001548	B9J08_00019	CM076438.1	1
XTC_3	*SIT1_3921*	B9J08_003921	B9J08_01779	CM076438.1	1
XTC_4	*SIT1_4097*	B9J08_004097	B9J08_01954	CM076438.1	1
XTC_5	*SIT1_1948*	B9J08_001948	B9J08_03021	CM076440.1	3
XTC_6	*SIT1_2110*	B9J08_002110	B9J08_03183	CM076440.1	3
XTC_7	*SIT1_1487*	B9J08_001487	B9J08_03737	CM076440.1	3
XTC_8	*SIT1_1499*	B9J08_001499	B9J08_03749	CM076440.1	3
XTC_9	*SIT1_1519*	B9J08_001519	B9J08_03769	CM076440.1	3
XTC_10	*SIT1_1521*	B9J08_001521	B9J08_03771	CM076440.1	3
XTC_11	*SIT1_2241*	B9J08_002241	B9J08_05251	CM076444.1	7
XTC_12	*SIT1_2465*	B9J08_002465	B9J08_05475	CM076444.1	7
HTR_1	*SIT1_3908*	B9J08_003908	B9J08_01766	CM076438.1	1
HTR_2	*SIT1_0002*	B9J08_000002	B9J08_01961	CM076439.1	2
HTR_3	*SIT1_4474*	B9J08_004474	B9J08_04433	CM076442.1	5
HTR_4	*SIT1_4475*	B9J08_004475	B9J08_04434	CM076442.1	5
HTR_5	*SIT1_2581*	B9J08_002581	B9J08_05591	CM076444.1	7

Table 1: Summary of *XTC* and *HTR* genes and their chromosomal locations.