1

Asymmetric neural entrainment at resonance frequencies underlies unilateral spatial neglect

Yuka O. Okazaki[1, 2, *], Noriaki Hattori[3], Teiji Kawano[4], Megumi Hatakenaka[4], Ichiro Miyai[4], and Keiichi Kitajo[1, 2, 5*]

1. Division of Neural Dynamics, Department of System Neuroscience, National Institute for Physiological Sciences, National Institutes of Natural Sciences, Okazaki, Japan
2. Physiological Sciences Program, Department of Advanced Studies, Graduate Institute for Advanced Studies, Graduate University for Advanced Studies (SOKENDAI), Okazaki, Japan
3. Department of Rehabilitation, University of Toyama, Toyama, Japan
4. Neurorehabilitation Research Institute, Morinomiya Hospital, Osaka, Japan
5. RIKEN CBS-TOYOTA Collaboration Center, RIKEN Center for Brain Science, Wako, Japan

*Corresponding authors:
Yuka O. Okazaki, yokazaki@nips.ac.jp;
Keiichi Kitajo, kkitajo@nips.ac.jp

Keywords: Alpha oscillations, Hemispheric imbalance, Steady-state visual evoked potentials

Abstract

Unilateral spatial neglect (USN) is a common consequence of right-hemisphere stroke,
traditionally attributed to structural lesions and dysfunctional attention networks. However, the
brain is fundamentally a rhythmic and dynamical system, and how disrupted neural
synchronization underlies USN remains unknown. We recorded steady-state visual evoked
potentials (SSVEPs; 3 - 30 Hz flicker) in stroke patients with USN, without USN, and healthy
controls. Only the USN group exhibited significant hemispheric asymmetry at 9 Hz, driven by
exaggerated responses in the intact hemisphere rather than suppression in the lesioned
hemisphere. This effect appeared only during stimulation, not at rest, indicating its specificity
to sensory processing. The enhanced 9 Hz entrainment in the intact hemisphere was
accompanied by increased phase-amplitude coupling (PAC) between alpha phase and
gamma amplitude, reflecting systematic coordination of high-frequency activity. Transfer
entropy analysis further revealed increased feedforward information flow from the right visual
to the left frontal cortex, highlighting large-scale asymmetry. To explore the mechanism
underlying this frequency-specific bias, we implemented a coupled-oscillator model. The
model showed that the hemispheric asymmetry arises from resonance between intrinsic alpha
rhythms and external input, amplified by asymmetric right-to-left interhemispheric coupling.
These findings suggest that USN arises from a selective impairment of alpha-band synchrony
capacity. This study offers a novel framework conceptualizing USN as a disorder of disrupted
oscillatory dynamics underlying spatial attention, and points toward frequency-specific
neuromodulatory intervention as a potential therapeutic approach.

Significant statement

The brain behaves as a metronome-like oscillator ensemble. When a rhythmic flicker—a cortical "tuning fork" matching the visual system's intrinsic alpha frequency—is applied, it serves as a probe that visualizes the brain's internal state. Under normal conditions, neuronal oscillations in both hemispheres are moderately entrained at this frequency. In unilateral spatial neglect (USN), this resonance displays a distinctive asymmetry: the intact hemisphere becomes over-amplified, whereas the lesioned hemisphere shows no enhancement. A coupled-oscillator model shows that asymmetric interhemispheric coupling selectively boosts this intrinsic alpha resonance unilaterally. This biased resonance reflects a destabilization of the dynamic coordination of attention networks, offering a novel mechanistic view of spatial neglect in USN and pointing toward frequency-specific neuromodulation as a promising therapeutic strategy.

Introduction

Left unilateral spatial neglect (USN) is a neurological disorder in which patients fail to perceive stimuli on the left side of space despite an intact visual field. This condition is associated with damage to the right dorsal attention network, particularly the right inferior parietal lobule (1, 2), and the superior temporal gyrus (3, 4), as well as disconnection in occipital and frontal white matter pathways (5). The dorsal attention network (6), which includes the frontal eye fields and bilateral parietal sulci, encodes spatial maps of the contralateral visual field in each hemisphere. These maps are typically balanced through reciprocal inhibitory interactions(7). Disruption of this network leads to hyperactivity of the structurally intact left hemispheric circuits, which influences bottom-up sensory responses and causes a rightward bias in perception (1, 8).

Traditionally, USN has been understood in terms of structural or static disconnection. However, the brain is fundamentally a dynamical and frequency-organized system, and neural oscillations, generated by the synchronized activity of neuronal populations, are a key mechanism for temporally coordinating information flow across brain regions. Indeed, neural oscillations flexibly regulate sensory and attentional processing by dynamically adjusting intrinsic and stimulus-driven phase relationships (9-14). From this perspective, USN may not only involve anatomical disconnection but also an imbalance in the hemispheric capacity for oscillatory synchronization. Yet, despite the critical role of rhythmic synchronization in brain function, such frequency-specific asymmetries have not been systematically investigated in USN.

To address this, we employed steady-state evoked responses (SSERs) to visual stimuli as a probe to visualize the brain's frequency-dependent synchronization capacity. SSERs are elicited by rhythmic sensory stimulation and reflect neural entrainment—the periodic alignment of endogenous oscillations to external input (15). This response is strongest when the stimulus frequency matches the system's intrinsic resonance frequency (16, 17). Moreover, prior studies have shown frequency-specific reductions in SSERs, both visual and auditory, in individuals with schizophrenia (18, 19) and autism (20, 21). Consequently, SSER abnormalities may reflect impairments in underlying oscillatory mechanisms.

In this study, we presented 3–30 Hz flickering stimuli and recorded steady-state visual evoked potentials (SSVEPs) in stroke patients with and without USN, as well as in healthy controls. None of the patients exhibited clinically evident hemianopsia, and early-stage visual responses in the lesioned hemisphere were not expected to be significantly impaired. These factors minimized the possibility that observed differences could stem from low-level visual deficits. We hypothesized that frequency-specific entrainment of neural oscillations would be selectively disrupted in one hemisphere, resulting in an interhemispheric imbalance in USN.

97   Furthermore, motivated by the empirical asymmetry, we developed a computational model of
98   coupled neural oscillators (22, 23) to test which factors could account for the emergence of
99   frequency-specific differences in oscillatory responses.
100

Materials and Methods

Participants

The study consisted of three groups: USN patients (n = 21), stroke patients without USN (n =
38), and healthy control participants (n = 17). We recruited patients who had been admitted to
the Kaifukuki (convalescent) Rehabilitation Ward of Morinomiya Hospital. The inclusion criteria
for stroke patients were: (1) male, (2) first-episode stroke, and (3) right hemisphere
cortical/subcortical infarction (representative lesions shown in Supplementary Figure 1),
without clinically evident hemianopsia evaluated by the confrontation method. While lesions
partially extended to the posterior region in two patients, they exhibited preserved
neurophysiological responses to light stimulation, and were therefore included in the analysis.
For healthy participants, the inclusion criteria were: (1) male to match the patient group, (2) no
history of stroke, and (3) full scores on all neuropsychological assessments. All participants
provided written informed consent. The study was approved by the Institutional Review Boards
of RIKEN, the National Institutes of Natural Sciences, and Morinomiya Hospital.
115

Clinical assessment of stroke

Electroencephalography (EEG) measurements and clinical assessments were performed by
trained investigators approximately 1 month after stroke onset. The diagnosis of USN was
comprehensively determined by the attending physician based on regular clinical
examinations, including the Behavioral Inattention Test (BIT) (24). Motor impairment was
assessed using the Fugl-Meyer Assessment upper extremity (FMA-UE) and lower extremity
(FMA-LE) (25), and activities of daily living were assessed using the Functional Independence
Measure motor (FIM-M) and cognitive (FIM-C) scores (26). All clinical assessments were
performed by trained healthcare professionals before EEG analysis, so the evaluators were
blind to the EEG analysis results. For all scales, a lower score indicates greater disability. To
match the severity of the USN and non-USN groups, we selected 21 non-USN patients in
ascending order of the Mahalanobis square distance, d_i^2, to the USN patient within a four-
dimensional space of the clinical score:
129

$$d_i = \sqrt{(\boldsymbol{y}_i - \boldsymbol{\mu})\,\Sigma^{-1}(\boldsymbol{y}_i - \boldsymbol{\mu})^T},$$

where y_i is the i-th non-USN patient's clinical scores and μ and Σ are the mean and covariance matrix of the USN samples, respectively. The demographic information and clinical scores of each group are listed in Table 1. An unpaired t-test confirmed that there were no significant differences between the groups in any of the scores.

Table 1. Demographic information of each group and clinical scores of the USN and non-USN patients matched by Mahalanobis distance

Variables	Median (Q1-Q3)			*p*-value (*U*-test)
	Non-USN	**USN**	**Healthy control**	
Number of participants	21	21 (16)	17	—
Sex	Male	Male	Male	—
Age (years)	69.0 (62.8–78.2)	67.0 (54.8–77.5)	67.0 (58.072.2)	0.21, 0.91 (USN vs. non-USN, USN vs. Healthy
Cortical/subcortical lesion side	Right	Right	—	—
EEG recording (days poststroke)	32.0 (21.5–41.0)	38.0 (32.5–43.0)	—	0.19
FMA-UE (0–66)	49.0 (34.2–61.2)	53.0 (12.8–62.2)	—	0.93
FMA-LE (0–34)	26.0 (22.5–30.0)	28.0 (16.2–31.0)	—	0.92
FIM-M (13–91)	64.0 (45.0–71.5)	47.0 (32.8–72.8)	—	0.23
FIM-C (5–35)	30.0 (26.8–31.2)	29.0 (22.0–30.0)	—	0.13
BIT (0–222)	—	204.5 (195.5–216.5)	—	—

Experimental Design

EEG was recorded from 19 Ag/AgCl electrodes in the 10-20 system configuration using a NeuroFax EEG 1224 system (Nihon Kohden, Tokyo, Japan; sampling rate: 500 Hz; bandpass filter: 0.53–120 Hz; ground electrode: forehead; reference: left earlobe). Visual stimulation was delivered under an eyes-closed resting condition to minimize blink artifacts and reduce direct exposure to the high-intensity strobe light. A strobe light flickered at frequencies ranging from 3 to 30 Hz in 3-Hz steps. Each frequency flicker stimulus was presented for 10 s with a 10-s interval (Figure 1).

154 **Figure 1. Experimental paradigm.** The strobe light flickered for 10 seconds in 3-Hz steps, from 3 Hz to 30 Hz,
155 with 10-second inter-train intervals. Participants were lying down with their eyes closed during the EEG recording.

Preprocessing

The EEG signals were first re-referenced offline to the average of the right and left earlobe
signals. We applied wavelet-enhanced independent component analysis (wICA) to eliminate
intermittent high-amplitude artifacts (27). This method decomposes the EEG signals into
independent components (ICs), and a discrete wavelet transform is applied to each IC.
Wavelet coefficients defined as artifacts based on the level-dependent threshold of the wavelet
decomposition are zeroed, and artifact-free ICs are extracted from the inverse wavelet
transform on this result. Thus, the entire IC, which contains a significant amount of neural
signals, need not be excluded, as in the case of conventional ICA denoising.

Steady-state visual evoked potential (SSVEP)

To evaluate the neural response during the flickering stimulation, the frequency power
spectrum for each stimulus condition was obtained using the Fourier transform (Hanning
window, frequency range: 1–35 Hz). The Fourier transform was applied to the data with a 1-s
window during the 10-s flicker interval and averaged for each stimulus frequency condition.

Phase-amplitude coupling (PAC)

We evaluated PAC using the modulation index (MI) (28) to determine how high-frequency
oscillations are co-modulated by a low-frequency phase when neural oscillations are entrained
to rhythmic external stimuli. Instantaneous phase and amplitude were extracted via wavelet
transform at the center frequency, f, and time, t, with the standard deviation (SD), $\sigma_f = 4f/m$, σ_t
$= m/2\pi f$, and constant, m, set to 4 (29). We focused on the PAC between the alpha (6–12 Hz)
phase and the gamma (30–55 Hz) amplitude for 10 s during the 9-Hz flickering stimulus.

To derive the MI for each frequency pair of α phase and γ amplitude, the given α frequency
phase was divided into 20 bins, and the mean γ amplitude in each bin was calculated. This
provided a phase-amplitude distribution for each frequency pair, and the MI was obtained
according to the Kullback-Leibler divergence between the observed distribution and the

circular uniform distribution. An MI of 0 indicates that the amplitudes are the same for all phase

bins (i.e., a circular uniform distribution) and that there is zero PAC, whereas an MI of 1

indicates maximum coupling.

Transfer entropy (TE)

To quantitatively investigate the direction and strength of information transfer between brain

regions in response to flickering stimuli, we performed TE analysis (30). After applying an 8.5–

9.5 Hz FIR bandpass filter to the EEG responses to 9-Hz flickering stimuli, we extracted the

instantaneous amplitude from the analytic signal using the Hilbert transform. TE was

calculated for all pairs of electrodes for these amplitude data during the flickering stimulus

interval. To estimate the probability distributions required for TE calculation, we generated

histograms for the amplitude data, with bin sizes according to Sturges' rule (31). $T_{Y \to X}$

quantifies the extent to which knowledge of the state of system *Y* at time *t* contributes to

predicting the one-step future state of system *X* at time *t+τ*, beyond the predictability provided

by the state of *X* at time *t* alone. The formula for TE is given by:

$$T_{Y \to X(\tau)} = \sum p(x_{t+\tau}, x_t, y_t) \log_2 \frac{p(x_{t+\tau}|x_t, y_t)}{p(x_{t+\tau}|x_t)},$$

where $p(x_{t+\tau}, x_t, y_t)$ denotes the joint probability of observing the future state of *X* at time *t+τ*

along with the states of *X* and *Y* at time *t*. Here, *x* and *y* represent the observed data derived

from the systems *X* and *Y*, respectively. The terms $p(x_{t+\tau}|x_t, y_t)$ and $p(x_{t+\tau}|x_t)$ denote the

conditional probabilities of the one-step future state of *X,* given its state at time *t* with and

without considering the state of *Y* at time *t*.

In this analysis, amplitude data from the first 0.5 s of the flickering stimulus interval were

used as past observations, x_t, y_t, and amplitude data from one sample ahead in time were

used as future observations, $x_{t+\tau}$. TE was calculated by varying the delay, *τ*, from 0 to 5 s in

0.1-s steps. The optimal delay *τ* was determined as the value that maximized $T(\tau)$.

Statistical analysis

To statistically test for differences in SSVEP between the left and right hemispheres, we used

a linear mixed model with hemisphere (left and right), flicker frequency (3, 6, 9, 12, 15, 18, 21,

24, 27, and 30 Hz), and group (USN patients, non-USN patients, and healthy controls) as fixed

factors and subjects as random factors, with the exclusion of outliers 1.5 times over the

interquartile range.

To statistically test for differences in PAC between the left and right hemispheres, we

performed a cluster-based permutation test (32). The MI was calculated for each permutation
dataset generated by random reassignment of O1 and O2 electrode labels. For each
permutation, clusters were identified from adjacent data points showing statistically significant
hemispheric differences in MI, and the sum of the *t*-values within these clusters was
determined as a cluster-level statistic. Only the largest value of the cluster-level statistic
obtained for each permutation was selected, and this procedure was repeated 1000 times to
construct the distribution of the cluster-level statistic under the null hypothesis. A cluster of
hemispheric differences in the original data was considered statistically significant if it
exceeded the 95th percentile of this distribution.

The coupled phase oscillator model

To investigate potential mechanisms underlying our experimental findings, we employed the
Kuramoto model (22) as a theoretical framework for neural synchronization. The model
consists of bilateral coupled phase oscillators and an external periodic force (23). Two coupled
populations of oscillators (N = 50 each) are represented by phase vectors $\boldsymbol{\Phi_L}$= ($\phi_{L,1}$, ..., $\phi_{L,n}$)
and $\boldsymbol{\Phi_R}$ = ($\phi_{R,1}$, ..., $\phi_{R,n}$) for the left and right hemispheres, respectively. The dynamics of each
oscillator are given by:

238
$$\frac{\mathrm{d}\phi_{h,i}}{\mathrm{d}t} = \omega_{h,i} + H_{\mathrm{intra}} + H_{\mathrm{inter}} + F(t)$$

240　The natural frequencies $\omega_{h,i}$ follow a Gaussian distribution with a mean of 9 Hz and a standard
241　deviation of 2 Hz. In one configuration, we set the mean natural frequency of the right
242　hemisphere to 7 Hz to examine the effect of altered intrinsic frequency in the affected
243　hemisphere. The intrahemispheric coupling H_{intra} and interhemispheric coupling H_{inter} terms
244　are defined as:

246
$$H_{\mathrm{intra}} = \frac{K_{intra}}{N} \sum_j \sin\left(\phi_j - \phi_i\right),$$

247
$$H_{\mathrm{inter}} = \begin{cases} \dfrac{K_{R\to L}}{N} \displaystyle\sum_j \sin\left(\phi_{R,j} - \phi_{L,i}\right), & \text{for the left hemisphere,} \\ \dfrac{K_{L\to R}}{N} \displaystyle\sum_j \sin\left(\phi_{L,j} - \phi_{R,i}\right), & \text{for the right hemisphere,} \end{cases}$$

249　where $\phi_{,i}$ represents the phase of the *i*-th oscillator in left and right hemispheres. K_{intra}
250　represents intrahemispheric coupling strength, and $K_{R\to L}$ and $K_{L\to R}$ represent interhemispheric
251　coupling strengths from right to left and left to right hemispheres, respectively. The external
252　forcing term is given by:

$$F(t) = b \sin\left(2\pi\omega_f t - \phi_{h,i}\right),$$

256 where amplitude and frequency of the external periodic force are set to $b = 2$ and $\omega_f \in \{3, 6,$

257 $..., 30\}$ Hz. To quantify the degree of synchronization, we computed the instantaneous order

258 parameter within each hemisphere:

$$\rho_h(t) = \left| \frac{1}{N} \sum_{j}^{N} e^{i\phi_{h,j}(t)} \right|$$

We performed 21 independent simulations, each corresponding to a virtual subject with

different initial phases randomly drawn from a uniform distribution [0, 2pi] and natural

frequencies sampled from Gaussian distributions. The order parameter was averaged over 10

seconds with continuous stimulation. The numerical simulation used the 4th-5th-order Runge-

Kutta method with adaptive time steps, as implemented by MATLAB's ode45 function.

Results

SSVEP power in the affected hemisphere

To confirm the visual cortex of the affected (right) hemisphere could respond to visual stimuli,

we plotted SSVEP power at the O2 electrode as a function of the flicker and response

frequencies (Figure 2). When compared with the power spectrum of the baseline period, a

clear fundamental can be seen on the diagonal (black dashed line), which indicates that the

visual cortex of the affected hemisphere was precisely enhanced at the flicker frequency.

Figure 2. SSVEP power for different responses and flicker frequencies. Normalized power spectrum of the

response frequency (y-axis) at the O2 electrode (affected hemisphere for patients) as a function of flicker frequency

(x-axis). The fundamental frequency and the first harmonics (gray dashed lines) can be seen in all groups.

Hemispheric imbalance of the SSVEP

The SSVEP power of the affected (right) hemisphere (O2 electrode) was compared with that

of the intact (left) hemisphere (O1 electrode) to investigate hemispheric balance in frequency

responses. We found a significant hemisphere × flicker frequency × group interaction (F =

2.09, p < 0.005). To further investigate this interaction, a simple main effects test was

conducted for hemisphere differences at each frequency within each group, with a Bonferroni

correction applied for multiple comparisons. Within the USN group, SSVEP power of the intact

hemisphere was significantly greater than that of the affected hemisphere at 9 Hz (F = 62.36,

p < 0.001). No significant hemisphere differences were observed at other frequencies within

the USN group. In the non-USN and healthy control groups, there were no significant

hemisphere differences at any frequency after Bonferroni correction. Figure 3 shows the

hemispheric difference in SSVEP power (O1 - O2) corresponding to stimulation frequency.

This suggests that frequency-specific hemispheric bias is the factor that separates USN from

non-USN patients.

300
301

Figure 3. Hemispheric differences in SSVEP power at different flicker frequencies. Power in the intact

hemisphere was stronger than that in the affected hemisphere with 9-Hz stimulation (***p < 0.001) in the USN

group only (middle panel). Error bars represent 2 standard errors.

We applied the same analysis for the central (C3, C4) and frontal (F3, F4) electrodes to

verify the frequency response of hemispheric bias outside the visual area. Figure 4 shows the

difference in SSVEPs between C3 and C4 electrodes (Figure 4A) and between F3 and F4

electrodes (Figure 4B) at each frequency. We found no significant main effects for

hemispheres or interactions in either the central or frontal regions.

314
315
316 **Figure 4. Hemispheric differences in SSVEP power at different flicker frequencies. (A)** No hemispheric
317 difference between C3 and C4 electrodes. **(B)** No hemispheric difference between F3 and F4 electrodes. Error
318 bars represent 2 standard errors.

319
320

Hyperactivity of the intact hemisphere

To determine whether the hemispheric imbalance of SSVEP power confined to 9 Hz in the
USN group was attributed to the intact (left) or affected (right) hemisphere, we compared
SSVEP power at 9 Hz in the corresponding hemisphere in the healthy controls (Figure 5A). A
simple main effects test showed a significant effect of group on SSVEP power in the left
hemisphere ($F = 6.66$, $p < 0.005$) and in the right hemisphere ($F = 10.58$, $p < 0.001$). Post-hoc
comparisons with Bonferroni correction indicated that in the left hemisphere, both the USN
and the non-USN group had significantly higher SSVEP power than the healthy control group
($p < 0.005$ and $p < 0.05$, respectively). In the right hemisphere, the non-USN group showed
significantly higher SSVEP power than the healthy control group ($p < 0.005$), while the USN
group did not differ significantly from the healthy controls. We also examined the resting state
interval (i.e., before the photo flicker session) and found no significant differences in either the
USN or non-USN group (Figure 5B). Taken together, there was no major abnormality in the
spontaneous generation of alpha oscillations in the USN group; rather, it was the capacity to
synchronize these oscillations with other external oscillations that differed between the
hemispheres in the USN group.

337
338

Figure 5. SSVEP power during stimulation at 9 Hz and rest. (A) Box plots indicate the power of SSVEP
responses in the intact (O1 electrode) and affected (O2 electrode) hemispheres. A significantly higher response
power was observed in the intact hemisphere of the USN group than in the left hemisphere of the healthy control
group. The non-USN patients also displayed higher alpha power in both hemispheres than healthy controls without
hemispheric bias (***p < 0.001). **(B)** During the resting state interval, no significant differences in alpha power were
observed between groups. Box plots depict medians and 25th/75th percentiles, with whiskers showing the
10th/90th percentiles overlaid onto individual data points (circles). The + represents outliers. Outliers exceeding a
power value of 50 were truncated and plotted at the 50 mark.

Hemispheric imbalance of PAC

We investigated PAC given that higher frequency oscillations may be cooperatively modulated
when neural oscillations are entrained by rhythmic external stimuli. PAC was assessed using
the MI for 9-Hz flickering stimuli (Figure 6). We hypothesized that the intact left hemisphere,
which showed enhanced SSVEP, would exhibit stronger PAC. Thus, we performed a cluster-
based permutation test to examine hemispheric differences. Results showed a significant
hemispheric bias in PAC between the alpha (7–12 Hz) phase and gamma (35–45 Hz)
amplitude in the USN group. Specifically, the MI was higher in the intact hemisphere than in
the affected hemisphere ($p < 0.05$, one-tailed) peaking at the 9-Hz phase. These effects were
not observed in the non-USN or healthy control group. This suggests that gamma activity is
modulated by the alpha phase aligned to the 9-Hz flicker stimulation and that this modulation
is particularly enhanced in the intact hemisphere of the USN group.

Figure 6. Hemispheric differences in PAC in response to 9-Hz flickering stimuli. The MI measuring the
coupling between the alpha phase (7–12 Hz) and gamma power (35–45 Hz) represents the hemispheric asymmetry
in the USN group (indicated by the black outline). No such pattern was present in the non-USN or healthy control
group.

Hemispheric imbalance of TE

In USN and non-USN patients with right hemisphere damage, we aimed to clarify how
information is transferred from the responsive visual cortex to other brain regions. Here, we
used TE to examine the pattern of information transfer from the O1 and O2 electrodes to the
left and right frontal electrodes (Figure 7). Results showed that in the USN group, there was
significantly greater information flow from the right visual area (O2 electrode) than from the
left visual area (O1 electrode) to the intact left frontal areas ($t = -2.7$, $p < 0.05$, paired t-test).
This enhancement was significantly greater than that in the healthy control group ($t = -2.89$,
$p < 0.01$, unpaired t-test; Figure 7A, left panel). This indicates a hemispheric bias in information
transfer in the feedforward direction in USN patients. In contrast, no such bias was observed
in non-USN patients ($t = -0.47$, $p = 0.64$, paired t-test) or healthy controls ($t = 1.42$, $p = 0.174$,
paired t-test). There were no significant differences in information flow from the O1/O2
electrodes to the affected right frontal areas (Figure 7A, right panel) or in the feedback direction
(i.e., from the left/right frontal areas to the O1/O2 electrodes) in any group (Figure 7B). These
findings suggest that the distinct clinical symptoms of USN are not solely due to biased local
brain activity but are closely related to asymmetric information transfer, especially in the
feedforward direction.

Figure 7. Hemispheric differences in information flow between the visual and frontal areas in response to

9-Hz flickering stimuli. (A) Box plots show left-right hemispheric asymmetry in transfer entropy (TE) from visual

to frontal areas (feedforward direction). In the USN group, information flow from the right visual cortex (O2) to the

left frontal area (left panel) was significantly stronger than from the left visual cortex (O1) to the left frontal area (*p

< 0.05), and this asymmetry was significantly greater than in healthy controls (**$p < 0.01$). No significant

hemispheric differences were observed in information flow to the right frontal area (right panel). **(B)** Box plots show

left-right hemispheric asymmetry in TE from frontal to visual areas (feedback direction), with no significant

asymmetry detected in any group for either left or right frontal areas. Arrows on topographical plots indicate the

direction of information flow. Box plots show medians and 25th/75th percentiles, with whiskers at 10th/90th

percentiles. The + represents outliers.

Correlation between SSVEP power and USN severity

To investigate the relationship between neural synchronization capacity and symptom severity in USN patients, we performed Pearson's correlation analysis between SSVEP power and BIT scores. Results showed a significant positive correlation between SSVEP power at 9 Hz in the right hemisphere (O2 electrode) and BIT score (Pearson's r = 0.70, p < 0.01). That is, the higher the SSVEP power at 9 Hz in the right hemisphere, the milder the USN symptoms. In contrast, there was no significant correlation in the left hemisphere (O1 electrode; Pearson's r = 0.38, p = 0.18; Figure 8A). Additionally, to examine the frequency-specific nature of the relationship between SSVEP power and USN severity, we analyzed correlations across all stimulation frequencies. Significant correlations between SSVEP power and BIT score were observed only at 9 and 12 Hz in the right hemisphere (Pearson's r = 0.70, p < 0.01, for both frequencies, FDR-corrected; Figure 8B). No significant correlations were found at any frequency in the left hemisphere (O1 electrode). Our results showed that USN patients maintained 9 Hz SSVEP power in the right hemisphere comparable to healthy controls. However, within the USN group, this right hemisphere activity significantly correlated with symptom severity, with higher SSVEP power associated with milder USN symptoms.

Figure 8. Correlation between SSVEP power and BIT score in USN patients. (A) Scatter plots represent the relationship between SSVEP power at 9 Hz and BIT score for the left hemisphere (O1 electrode, left panel) and right hemisphere (O2 electrode, right panel). A significant positive correlation was found in the right hemisphere (r = 0.70, p < 0.007) but not in the left hemisphere (r = 0.38, p = 0.18). **(B)** Correlation coefficients between SSVEP

430 power and BIT score across frequencies (3–30 Hz) for each hemisphere. Significant correlations were observed in
431 the right hemisphere only at 9 and 12 Hz after FDR correction (**$p < 0.01$).

Hemispheric Imbalance in Model Simulations

To explain the observed hemispheric imbalance in USN patients (Figure 3, middle), we
explored three potential mechanisms using a computational model. From a base configuration
($\omega_L = \omega_R = 9\pm2$ Hz, $K_L = K_R = K_{L\to R} = K_{R\to L} = 0.2$), we modified specific parameters. In the first
configuration, we introduced asymmetric interhemispheric coupling ($K_{R\to L} = 0.8$). This
configuration most closely reproduced our experimental findings, showing a significant
hemispheric bias specifically at 9 Hz (p < 0.05, FDR corrected), while maintaining balanced
responses at other frequencies (Figure 9A). In the second configuration, we increased
intrahemispheric coupling strength in the intact hemisphere ($K_L = 0.8$) while maintaining the
baseline coupling in the affected hemisphere ($K_R = 0.2$), based on our experimental
observation of enhanced activity specifically in the intact hemisphere. This resulted in a
persistent left hemispheric bias across frequencies, with a particularly pronounced effect at 9
Hz (Figure 9B). In the third configuration, we set different natural frequencies by lowering the
right hemisphere ($\omega_R = 7$ Hz). This configuration showed a significant right hemispheric bias
at 6 Hz, close to the natural frequency of the right hemisphere, and a left hemispheric bias at
9 Hz (Figure 9C). Notably, when we removed the external periodic force in the first
configuration, no hemispheric bias was observed (see red bar in Figure 9A), indicating that
the asymmetric coupling manifests its effects only through interaction with external stimulation.

Figure 9. Hemispheric imbalance in model simulations. Upper panels **(A-C)** show schematic illustrations of three different model configurations. Lower panels show corresponding order parameter differences between hemispheres across stimulus frequencies. **(A)** Model with asymmetric interhemispheric coupling ($K_{R \to L}$ = 0.8, $K_{L \to R}$ = 0.2) reproduced the experimental findings, showing significant hemispheric bias at 9 Hz. The red bar indicates the values without external stimulation ($F(t)$ = 0). **(B)** Model with enhanced left intrahemispheric coupling (K_L = 0.8, K_R = 0.2) produced a persistent bias across frequencies. **(C)** Model with different natural frequencies between hemispheres (ω_L = 7, ω_R = 9) led to frequency-specific biases at both hemispheres' natural frequencies. Asterisks indicate significant differences (*p < 0.05, FDR corrected). Error bars represent 2 standard errors

Discussion

Our experiments revealed that while SSVEPs can be reliably elicited by flickering stimuli at various frequencies, only the USN group exhibited a significant hemispheric difference in SSVEP at 9 Hz. The computational model demonstrated that this hemispheric bias could arise from both the resonance between intrinsic and stimulus frequencies and stronger right-to-left interhemispheric coupling. In contrast to the healthy control group, this difference was attributed to an increase in activity within the intact hemisphere rather than a decrease in the affected hemisphere. The enhanced response in the intact hemisphere was accompanied by increased phase-amplitude coupling between alpha and gamma oscillations, suggesting that the entrained alpha oscillations systematically modulated local high-frequency neural activity. Furthermore, the increased information flow from the right visual cortex to the left frontal cortex revealed by TE analysis is consistent with the model predictions, and these results indicate an imbalance in information flow between hemispheres in addition to changes in local activity.

Alpha frequency-specific hemispheric bias and entrainment in USN patients

We observed hemispheric bias in entrainment to flickering stimuli in USN patients in the alpha band, which corresponds to the natural frequency of the visual system (17, 33). The entrainment of the oscillatory system to an external stimulus is caused by the periodic phase shift of neural oscillations to the delivered pulse (34). At the EEG level, this is expressed as an increase in amplitude at the stimulus frequency as multiple oscillators begin to cycle in synchrony (35).

In general, entrainment is particularly effective when the stimulation period closely matches the oscillator's natural frequency (16). This principle also applies to neural oscillations: rhythmic stimulation at frequencies that resonate with physiological rhythms influences cognitive and neural function by synchronizing task-related oscillations (36-38). Our previous research demonstrated that entrainment is most effective when the stimulation frequency closely matches the individual alpha frequency (17). We also observed that different brain

regions have distinct natural frequencies, with entrainment being most pronounced in the
alpha band in the visual cortex. Therefore, the observed bias in the 9-Hz frequency-specific
SSVEP in USN patients does not merely reflect a hemispheric difference in evoked visual
responses to rhythmic external stimuli but, rather, a bias in the synchronous function of
intrinsic neural oscillators in the visual area at a specific frequency band (i.e., the alpha band).

Our computational model provides a theoretical framework for understanding these
hemispheric differences. Among the three potential mechanisms explored, the model with
asymmetric interhemispheric coupling ($K_{R \to L} > K_{L \to R}$) best reproduced the frequency-specific
hemispheric bias observed in our experiments. While the coupling parameter K in our phase
oscillator model simply determines the strength of phase synchronization between
hemispheres, this asymmetry might reflect the underlying imbalance in interhemispheric
inhibition. Indeed, paired-pulse TMS studies have directly demonstrated enhanced inhibitory
influence from the intact to the affected hemisphere in stroke patients (39). This
interhemispheric imbalance has been further supported by evidence of altered functional
connectivity and hyperexcitability in the intact hemisphere (7, 40).

In this model, both hemispheres with a natural frequency of 9 Hz resonate with stimulation
at the same frequency, but due to the stronger right-to-left coupling, the resonant response in
the right hemisphere more strongly drives oscillations in the left hemisphere. This suggests
that the hemispheric bias observed in USN patients is driven by asymmetric interhemispheric
coupling of neural networks underlying alpha oscillations. This model prediction also aligns
with our experimental observation of enhanced transfer entropy from the right visual cortex to
the left frontal areas in USN patients.

Furthermore, it demonstrates that while the system maintains balanced activity between
hemispheres in the absence of external stimulation, the underlying asymmetry becomes
apparent with external input. This explains why spontaneous alpha oscillations remain normal
despite the emergence of a specific bias in response to 9 Hz stimulation, providing a
theoretical basis for why USN symptoms become more pronounced during sensory
processing and attentional demands.

Local hemispheric bias of entrainment and attentional dysfunction in USN patients

Hyperactivity of the SSVEP and PAC in the left visual cortex in stroke patients may result from
a lack of hemispheric reciprocal inhibition (39). This scenario is consistent with the concept
that in USN, decreased inhibition from the affected right hemisphere causes overactivity of the
visual cortex within the intact left hemisphere, which results in a left hemisphere bias in visual
processing (40, 41). However, the hemispheric bias observed in the present study was
frequency-specific and was not observed in non-USN patients (who also had right hemisphere

damage), suggesting that it may reflect not only the overactivity of the visual cortex but also
alterations in the neural networks' capacity to synchronize with external stimuli at specific
frequencies. The entrainment of neural oscillations to external stimuli is achieved via
consistent phase alignment, which, in turn, systematically influences the timing of the higher
frequency components of neural activity (42). This is evidenced by our observation of
enhanced PAC between alpha phase and gamma amplitude in the intact hemisphere of USN
patients, suggesting a systematic modulation of high-frequency activity by the entrained alpha
oscillations.

This phase coordination is influenced by both external factors and internal processes, such
as attention in anticipatory contexts. Attention adjusts the temporal sequence of excitatory and
inhibitory phases in neural oscillations, thereby influencing perception (12, 13, 43, 44). Given
the importance of alpha-band oscillations in attentional processes (45), the hemispheric bias
in oscillatory synchrony observed in USN patients may reflect critical abnormalities in cortical
excitability coordination. Therefore, the hemispheric bias in synchronization capacity at
frequencies relevant to attention likely disrupts the coordinated oscillatory activity necessary
for balanced spatial attention processing. This may contribute to the attentional deficits
observed in patients with USN.

Biased information transfer in USN patients

Locally, the USN group showed a higher SSVEP and greater PAC in the left visual cortex,
whereas TE from the right visual cortex to the left frontal cortex was higher than TE from the
left visual cortex to the left frontal cortex. Given that right frontal damage is one of the common
causes of USN, this difference may be an adaptive neural response to maintain hemispheric
communication and compensate for attentional dysfunction. Specifically, input from the right
visual cortex may bypass the affected right frontal cortex to enable enhanced information flow
to the intact left frontal cortex, thereby supporting attentional functions. This compensatory
strategy of increased information flow is consistent with functional imaging study findings on
the reorganization of functional networks following stroke; moreover, it supports the idea of
neuroplasticity, whereby patients can recover functions and maintain cognitive abilities
following neurological injury (46-48).

Distinct neural responses in non-USN and USN patients

Although both the non-USN and USN groups had right hemisphere damage, they exhibited
distinct neural responses to alpha-band rhythmic stimulation. In non-USN patients, additional
neural resources were recruited from both hemispheres in response to the damage in the right
hemisphere, resulting in enhanced neural synchronization to external stimuli across both

hemispheres. In contrast, the neural responses in USN patients are more asymmetric, with a
local increase in alpha-band SSVEP primarily observed in the intact left visual cortex. This
asymmetry may result from the damaged right hemisphere being less effectively engaged,
leading to an over-reliance on the left hemisphere (1). This suggests that the right hemisphere
damage in USN patients might affect more critical parts of the attentional network.

In non-USN patients, the observed bilateral enhancement of alpha-band synchrony likely
reflects a compensatory recruitment of residual right-hemisphere circuits that are still able to
cooperate with the intact left hemisphere. Neuroimaging studies have reported that patients
who show this kind of bilateral reorganization pattern early after stroke tend to achieve better
recovery outcomes (7, 46, 49). Thus, the bilateral pattern seen in the non-USN group may
correspond to such a favorable reorganization profile, and we propose that engaging both
hemispheres in parallel helps maintain frequency-specific rhythm coordination and protects
against the development of spatial neglect.

Importantly, this distinct pattern of hemispheric imbalance cannot be explained merely by
differences in lesion extent between USN and non-USN patients. While patients with USN
may have larger lesions than those without USN, such differences alone would predict
generalized alterations in neural responses across all frequencies in the affected hemisphere.
However, our results demonstrate a highly specific effect: the hemispheric imbalance is
observed exclusively at 9 Hz, while responses to other frequencies remain balanced between
the hemispheres. This frequency specificity, coinciding with the intrinsic alpha frequency of
the visual system, suggests a more complex mechanism involving the selective disruption of
oscillatory networks, rather than a simple consequence of lesion size. This interpretation is
supported by our computational model, which demonstrates how the interaction between
intrinsic frequency and asymmetric coupling can produce such frequency-specific effects.

Interestingly, although right-hemisphere alpha-band SSVEP power did not differ
significantly between the USN group and healthy controls, it showed a significant positive
correlation with BIT scores within the USN group. This result indicates that, even when the
quantitative power of right-hemisphere alpha oscillations appears normal, their qualitative
contribution to the spatial-attention network is altered in USN. Such a qualitative dissociation,
together with compensatory hyperactivity in the intact left hemisphere and a rerouting of
information flow from the right visual cortex to the left frontal cortex, likely skews the overall
balance of spatial attention. The interplay among these quantitative factors, and between
these dynamics and their behavioral consequences, is probably nonlinear, and the present
data cannot fully unravel their complex interactions; nevertheless, the principle that "normal
power ≠ normal function" strongly suggests that USN arises from a qualitative breakdown in
rhythm coordination, offering a new perspective on the neural basis of unilateral spatial neglect.

Limitations

This study has several limitations that must be acknowledged. First, although we revealed a 9-Hz-specific hemispheric bias in USN, the study does not pinpoint where within the cortical network the abnormal synchrony originates. Higher spatial-resolution techniques, such as magnetoencephalography or high-density intracranial recordings, could map the generators and propagation pathways of alpha-band entrainment more precisely and thus identify the circuit nodes in which synchrony breaks down. Second, the precise extent and topography of brain lesions could not be fully homogenized across patients. Differences in cortical versus subcortical involvement or the degree of white-matter disruption may influence oscillatory responses. Future work with a larger cohort—encompassing a broader range of lesion characteristics—will be necessary to combine detailed lesion mapping with frequency-resolved measures of neural synchrony. Third, the cross-sectional design precludes definitive causal inferences. Longitudinal studies that track changes in alpha-band asymmetry and behavioral recovery over time, or before and after rehabilitation interventions, will be essential to determine how dynamic reorganization of neural synchrony relates to the evolution and remediation of spatial neglect.

Conclusions

Our study demonstrates fundamental differences in neural responses between stroke patients with USN and without USN. Empirical data and computational modeling together show that asymmetric interhemispheric coupling, resonating with the brain's intrinsic alpha rhythm, selectively amplifies 9-Hz entrainment in the intact hemisphere of USN patients. This amplification is accompanied by stronger α–γ phase–amplitude coupling and asymmetric information flow, indicating that the effect propagates beyond local oscillations to reconfigure cross-frequency interactions and large-scale network dynamics. Thus, USN can be regarded as a disorder in which a qualitative breakdown of alpha-band rhythm coordination destabilizes the dynamic equilibrium of the attention network. Longitudinal studies that track the plastic evolution of these synchrony patterns during recovery and refine frequency-specific neuromodulation strategies will be crucial for translating these insights into therapy.

Acknowledgments

This work was supported in part by the NINS OPEN MIX LAB Program (NINS Program No. OML032401); a research grant from TOYOTA Motor Corporation to KK; and a grant-in-aid for General Research on Adult Diseases Including Cardiovascular Diseases and Diabetes (23FA1015) from the Japanese Ministry of Health, Labor and Welfare to IM. We also thank Dr.

637 Hroshi Yokoyama for providing the code for the transfer entropy analysis.

Data, Materials, and Software Availability

640 The datasets generated and analyzed during the current study are not publicly available due
641 to ethical considerations. Analysis code and simulation scripts used in this study are available
642 from the corresponding authors upon reasonable request (yokazaki@nips.ac.jp)

References

644
645

1.   M. Corbetta, M. J. Kincade, C. Lewis, A. Z. Snyder, A. Sapir, Neural basis and recovery
of spatial attention deficits in spatial neglect. *Nat Neurosci* **8**, 1603-1610 (2005).

2.   D. J. Mort *et al.*, The anatomy of visual neglect. *Brain* **126**, 1986-1997 (2003).

3.   H. O. Karnath, New insights into the functions of the superior temporal cortex. *Nat Rev*
Neurosci **2**, 568-576 (2001).

4.   H. O. Karnath, S. Ferber, M. Himmelbach, Spatial awareness is a function of the temporal
not the posterior parietal lobe. *Nature* **411**, 950-953 (2001).

5.   M. Thiebaut de Schotten *et al.*, Direct evidence for a parietal-frontal pathway subserving
spatial awareness in humans. *Science* **309**, 2226-2228 (2005).

6.   M. Corbetta, G. L. Shulman, Control of goal-directed and stimulus-driven attention in the
brain. *Nat Rev Neurosci* **3**, 201-215 (2002).

7.   C. Grefkes, G. R. Fink, Reorganization of cerebral networks after stroke: new insights
from neuroimaging with connectivity approaches. *Brain* **134**, 1264-1276 (2011).

8.   B. J. He *et al.*, Breakdown of functional connectivity in frontoparietal networks underlies
behavioral deficits in spatial neglect. *Neuron* **53**, 905-918 (2007).

9.   N. A. Busch, J. Dubois, R. VanRullen, The phase of ongoing EEG oscillations predicts
visual perception. *J Neurosci* **29**, 7869-7876 (2009).

10. K. E. Mathewson, G. Gratton, M. Fabiani, D. M. Beck, T. Ro, To see or not to see:
prestimulus alpha phase predicts visual awareness. *J Neurosci* **29**, 2725-2732 (2009).

11. P. Lakatos *et al.*, The spectrotemporal filter mechanism of auditory selective attention.
Neuron **77**, 750-761 (2013).

12. P. Lakatos, G. Karmos, A. D. Mehta, I. Ulbert, C. E. Schroeder, Entrainment of neuronal
oscillations as a mechanism of attentional selection. *Science* **320**, 110-113 (2008).

13. C. E. Schroeder, P. Lakatos, Low-frequency neuronal oscillations as instruments of
sensory selection. *Trends Neurosci* **32**, 9-18 (2009).

14. P. Fries, Rhythms for Cognition: Communication through Coherence. *Neuron* **88**, 220-
235 (2015).

15. A. Notbohm, J. Kurths, C. S. Herrmann, Modification of Brain Oscillations via Rhythmic
Light Stimulation Provides Evidence for Entrainment but Not for Superposition of Event-
Related Responses. *Front Hum Neurosci* **10**, 10 (2016).

16. A. Pikovsky, M. Rosenblum, J. Kurths, Synchronization - A Universal Concept in
Nonlinear Sciences. *Cambidge University Press* (2003).

17. Y. O. Okazaki, Y. Nakagawa, Y. Mizuno, T. Hanakawa, K. Kitajo, Frequency- and Area-
Specific Phase Entrainment of Intrinsic Cortical Oscillations by Repetitive Transcranial
Magnetic Stimulation. *Front Hum Neurosci* **15**, 608947 (2021).

18. G. P. Krishnan *et al.*, Steady state visual evoked potential abnormalities in schizophrenia.
Clin Neurophysiol **116**, 614-624 (2005).

19. J. S. Kwon *et al.*, Gamma frequency-range abnormalities to auditory stimulation in
schizophrenia. *Arch Gen Psychiatry* **56**, 1001-1005 (1999).

20. T. W. Wilson, D. C. Rojas, M. L. Reite, P. D. Teale, S. J. Rogers, Children and adolescents
with autism exhibit reduced MEG steady-state gamma responses. *Biol Psychiatry* **62**,
192-197 (2007).

21. D. C. Rojas *et al.*, Transient and steady-state auditory gamma-band responses in first-
degree relatives of people with autism spectrum disorder. *Mol Autism* **2**, 11 (2011).

22. Y. Kuramoto, Chemical Oscillations, Waves, and Turbulence. *Springer* (1984).

23. H. Sakaguchi, Cooperative Phenomena in Coupled Oscillator Systems under External
Fields. *Progress of Theoretical Physics* **79**, 39-46 (1988).

24. B. Wilson, J. Cockburn, P. Halligan, Development of a behavioral test of visuospatial
neglect. *Arch Phys Med Rehabil* **68**, 98-102 (1987).

25. A. R. Fugl-Meyer, L. Jaasko, I. Leyman, S. Olsson, S. Steglind, The post-stroke
hemiplegic patient. 1. a method for evaluation of physical performance. *Scand J Rehabil*
Med **7**, 13-31 (1975).

26. R. A. Keith, C. V. Granger, B. B. Hamilton, F. S. Sherwin, The functional independence
measure: a new tool for rehabilitation. *Adv Clin Rehabil* **1**, 6-18 (1987).

27. N. P. Castellanos, V. A. Makarov, Recovering EEG brain signals: artifact suppression with
wavelet enhanced independent component analysis. *J Neurosci Methods* **158**, 300-312
(2006).

28. A. B. Tort, R. Komorowski, H. Eichenbaum, N. Kopell, Measuring phase-amplitude
coupling between neuronal oscillations of different frequencies. *J Neurophysiol* **104**, 1195-
1210 (2010).

29. J. P. Lachaux *et al.*, Studying single-trials of phase synchronous activity in the brain. *Int J*
Bifurcat Chaos **10**, 2429-2439 (2000).

30. T. Schreiber, Measuring information transfer. *Phys Rev Lett* **85**, 461-464 (2000).

31. P. A. Legg, P. L. Rosin, D. Marshall, J. E. Morgan, Improving accuracy and efficiency of
mutual information for multi-modal retinal image registration using adaptive probability
density estimation. *Comput Med Imaging Graph* **37**, 597-606 (2013).

32. E. Maris, R. Oostenveld, Nonparametric statistical testing of EEG- and MEG-data. *J*
Neurosci Methods **164**, 177-190 (2007).

33. M. Rosanova *et al.*, Natural frequencies of human corticothalamic circuits. *J Neurosci* **29**,
7679-7685 (2009).

34. P. Lakatos, J. Gross, G. Thut, A New Unifying Account of the Roles of Neuronal
Entrainment. *Curr Biol* **29**, R890-R905 (2019).

35. G. Thut, P. G. Schyns, J. Gross, Entrainment of perceptually relevant brain oscillations by
non-invasive rhythmic stimulation of the human brain. *Front Psychol* **2**, 170 (2011).

36. W. Klimesch, P. Sauseng, C. Gerloff, Enhancing cognitive performance with repetitive
transcranial magnetic stimulation at human individual alpha frequency. *Eur J Neurosci* **17**,
1129-1133 (2003).

37. P. Sauseng *et al.*, Brain oscillatory substrates of visual short-term memory capacity. *Curr*
Biol **19**, 1846-1852 (2009).

38. V. Romei, J. Driver, P. G. Schyns, G. Thut, Rhythmic TMS over parietal cortex links
distinct brain frequencies to global versus local visual processing. *Curr Biol* **21**, 334-337
(2011).

39. N. Murase, J. Duque, R. Mazzocchio, L. G. Cohen, Influence of interhemispheric
interactions on motor function in chronic stroke. *Ann Neurol* **55**, 400-409 (2004).

40. G. Koch *et al.*, Hyperexcitability of parietal-motor functional connections in the intact left-
hemisphere of patients with neglect. *Brain* **131**, 3147-3155 (2008).

41. M. Corbetta, G. L. Shulman, Spatial neglect and attention networks. *Annu Rev Neurosci*
34, 569-599 (2011).

42. S. Glim *et al.*, Phase-Amplitude Coupling of Neural Oscillations Can Be Effectively Probed
with Concurrent TMS-EEG. *Neural plasticity* **2019**, 6263907 (2019).

43. J. Besle *et al.*, Tuning of the human neocortex to the temporal dynamics of attended
events. *J Neurosci* **31**, 3176-3185 (2011).

44. M. Bonnefond, O. Jensen, Alpha oscillations serve to protect working memory
maintenance against anticipated distracters. *Curr Biol* **22**, 1969-1974 (2012).

45. W. Klimesch, alpha-band oscillations, attention, and controlled access to stored
information. *Trends Cogn Sci* **16**, 606-617 (2012).

46. A. K. Rehme, G. R. Fink, D. Y. von Cramon, C. Grefkes, The role of the contralesional
motor cortex for motor recovery in the early days after stroke assessed with longitudinal
FMRI. *Cereb Cortex* **21**, 756-768 (2011).

47. R. J. Nudo, Mechanisms for recovery of motor function following cortical damage. *Curr*
Opin Neurobiol **16**, 638-644 (2006).

48. D. Saur *et al.*, Dynamics of language reorganization after stroke. *Brain* **129**, 1371-1384
(2006).

49. N. S. Ward, M. M. Brown, A. J. Thompson, R. S. Frackowiak, Neural correlates of motor
recovery after stroke: a longitudinal fMRI study. *Brain* **126**, 2476-2496 (2003).

Supplementary figure

Supplementary Figure 1: Lesion patterns in USN patients. Examples of T2-weighted axial MRI images showing
right hemisphere lesions in three patients with unilateral spatial neglect (USN). All images are labeled with 'R' to
indicate the right hemisphere.

758