Malat1 regulates female Th2 cell cytokine expression through controlling early

differentiation and response to IL2

Mags Gwynne[1, 2, *], Katie A. West[1, 2, 3, *], Stijn van Dongen[4, 5], Ioannis Kourtzelis[1, 2], Dawn

Coverley[1, 3], Sarah A. Teichmann[6, 7], Kylie R. James[8], James P. Hewitson[1, 3], Dimitris Lagos[1,

[2].

1. Hull York Medical School, University of York, United Kingdom.   **2.** York Biomedical

Research Institute, University of York, United Kingdom. **3.** Department of Biology, University

of York, United Kingdom. **4.** Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton,

United Kingdom. **5.** Current address: ExpressionEdits Ltd, Haverhill, United Kingdom. **6.**

Wellcome Sanger Institute, Hinxton, United Kingdom. **7.** Current address: Cambridge Stem

Cell Institute, University of Cambridge, Cambridge, United Kingdom. **8.** Translational

Genomics Program, Garvan Institute of Medical Research, The Kinghorn Cancer Centre,

Darlinghurst, New South Wales, Australia.

* These authors contributed equally

Corresponding author: dimitris.lagos@york.ac.uk

Abstract

Beyond sex hormone and sex chromosome-driven mechanisms, little is known about cell

intrinsic mediators of immune sexual dimorphism. We show that _Malat1_ is required for

appropriate cytokine expression in female but not male Th2 cells. _Malat1_ deficiency impairs _in_

vitro Th2 differentiation of naïve CD4[+] T cells from female mice, characterised by

transcriptome-wide effects and suppression of cytokine expression, particularly IL10.

Mechanistically, naïve CD4[+] T cells from _Malat1[-/-]_ female mice demonstrate altered early

activation kinetics, followed by impairment of the early differentiation gene expression

programme, including up-regulation of an interferon responsive gene module. This is followed

by suppression of IL2Rα and IL2Rγ expression and IL2-mediated differentiation. Male CD4$^+$

T cell activation and Th2 differentiation are not sensitive to *Malat1* loss due to stronger early

activation, higher interferon responsive gene expression during early differentiation,

maintenance of IL2Rα expression independently of *Malat1*, and lower sensitivity to exogenous

IL2 during late differentiation. *In vivo* female, but not male, *Malat1$^{-/-}$* mice demonstrate altered

Th2 differentiation characterised by a reduction in IL10$^+$ Th2 cells in both lung and spleen

following priming and challenge with *Schistosoma mansoni* eggs, a model of lung type 2

inflammation. These findings reveal *Malat1* as a determinant of immune sexual dimorphism.

Introduction

Biological sex is a known determinant of immune responses (1-4). Higher incidence or severity of autoimmune diseases (5-7) in females and infectious diseases in men (8) are commonly reported. The main drivers of immune sexual dimorphism are sex hormones (extrinsic) and X or Y chromosome-linked genes (intrinsic). For example, it is thought that females favour Th2 responses and males Th1 (2) and that this is due to hormonal regulation of immune genes such as IL4 (2, 9), and function of X chromosome-linked genes including the long intergenic non-coding RNA (lincRNA) *Xist* (10), which was recently shown to form ribonucleoprotein complexes that are directly involved in autoimmunity (11). Other mechanisms including those occurring at the post-transcriptional level, have been proposed, yet very little has been experimentally tested and validated. Overall, beyond sex hormone or sex chromosome-driven mechanisms, little is understood about cell intrinsic mediators of sexual dimorphism in Th cells.

Thousands of long non-coding transcripts, including lincRNAs, have been discovered in mammalian transcriptomes (12). In most cases, these are medium to lowly expressed transcripts displaying poor conservation across mammals. Our studies have focused on *Malat1*, a 7.5kb-long transcript discovered through an association with cancer progression and metastasis (13). *Malat1* is localised in nuclear speckles (14), which are nuclear foci enriched in factors involved in pre-mRNA splicing and transcription (15). It was initially thought that *Malat1* controls in mRNA splicing (14, 16). However, $Malat1^{-/-}$ mice develop normally and are viable and fertile without apparent defects in splicing (17-19). Although *Malat1* deletion does not seem to affect formation and size of nuclear speckles under homeostatic conditions, it affects intra-speckle organisation and nuclear retention of RNA binding proteins (20). We previously discovered that *Malat1* is a key regulator of Th1 cell responses (21). We showed that *Malat1* is down-regulated upon initiation of Th cell differentiation, yet its complete loss results in stronger immune responses through suppression of IL10 expression in mouse models of type 1 immunity to parasitic infection, including visceral leishmaniasis and malaria. In humans, *Malat1* downregulation is a hallmark of proliferative CD4[+] and CD8[+] T cells (22).

Others have linked *Malat1* to CD4[+] T cell function, particularly in Th17 cells (23, 24), and to

CD8[+] T cell function (25).

In our previous report, which focused on Th1 cells, we showed that *Malat1* loss also affected

IL10 expression in *in vitro* differentiated Th2 cells (21). However, its role in Th2 cell

differentiation and function remain under-explored. Here, we aimed to test how loss of *Malat1*

affects Th2 differentiation *in vitro* and in an *in vivo* model of type 2 inflammation, *Schistosoma*

mansoni (*S. mansoni*) egg injection (26, 27). We report that *Malat1* deficiency leads to

impaired Th2 differentiation with widespread effects across the transcriptome and a notable

suppression of IL10 expression. Both *in vivo* and *in vitro*, susceptibility to *Malat1* deficiency

occurs only in female Th2 cells. This is due to a female-specific impairment of the early

differentiation gene expression programme underpinning, suppression of IL2 receptor (IL2R)

expression in *Malat1*[-/-] cells, and increased sensitivity of female cells to IL2-driven cytokine

production.

Materials and Methods

Animals and ethics

Animal care and experimental procedures were regulated under the Animals (Scientific

Procedures) Act 1986 (revised under European Directive 2010/63/EU) and were performed

under U.K. Home Office License (project license number PP0841992 for breeding and

PP9423191for *S. mansoni* egg injections) with approval from the University of York Animal

Welfare and Ethical Review Body. Animal experiments conformed to Animal Research:

Reporting of In Vivo Experiments guidelines (28).

***S. mansoni* egg injection**

C57BL/6 CD45.2 *wildtype* (*WT*) mice were obtained from Charles River Laboratories.

Malat1[−/−] mice (complete knockouts) were obtained from the Riken Institute (18). All mice were

bred in-house, maintained under specific pathogen-free conditions, and used at 6–12 weeks

of age. Schistosome eggs were recovered from the livers of C57BL/6 mice at week 7 post-

infection following exposure to 100 *S. mansoni* cercariae. Cercariae were obtained from

schistosome-infected *Biomphalaria glabrata* snails provided by the Barrett Centre for Helminth

Control (Aberystwyth University, UK). Livers were digested overnight at 37°C with shaking

with 0.2U/ml collagenase D (Roche) in the presence of 5000U/ml polymyxin B (Merck). Eggs

were purified by centrifugation through 10ml percoll (GE Healthcare) / 20ml 0.25M sucrose

(450 x*g*, 5min, RT), washed in PBS and stored at -20°C before usage. 5000 dead *S. mansoni*

eggs in 200 µl PBS were delivered via intraperitoneal (ip) injection into mice. Two weeks later,

the mice were intravenously (iv) challenged with another 5000 eggs in 200 µl PBS. After

another week the mice were sacrificed and the lungs and spleen were extracted and

processed. Lungs were digested with 0.4 U/ml Liberase TL (Roche) and 80 U/ml DNase I type

IV in HBSS (both Sigma-Aldrich) for 45 min at 37°C. Enzyme activity was inhibited with 10

mM EDTA (pH 7.5) and single-cell suspensions were created with 100-µm nylon filters

(Falcon) in complete RPMI 1640 (Thermo Fisher Scientific) supplemented with 10% heat-

inactivated FCS (HyClone), 100 U/ml penicillin, 100 µg/ml streptomycin, and 2 mM l-glutamine

(all Thermo Fisher Scientific), then cleaned via Percoll gradient and 3ml Ammonium-Chloride-

Potassium (ACK) red blood cell lysis buffer. Spleen single cell suspensions were generated

in the same manner and only passed through a 70 µm filter and treated with 3 ml ACK buffer.

***In vitro* Th2 differentiation and *Malat1* knockdown**

Spleens and axillary, brachial, mesenteric, and inguinal lymph nodes of *WT* or *Malat1*[-/-] mice

were extracted, homogenised, through a 70 µm filter in RPMI 1640. Resulting cell pellets were

then treated with 3ml ACK lysis buffer. Naïve CD4[+] T cells were then purified via MACS column

isolation (Miltenyi Naive CD4[+] T Cell Isolation Kit, mouse, catalogue number 130-104-453)

normally resulting in purity of ~95%. For Th2 polarisation, purified naïve CD4[+] T cells (500,000

cells per well) were stimulated with 10 µg/ml plate-bound anti-CD3ε (clone 145-2C11) and 4

µg/ml soluble anti-CD28 (37.51) in RPMI 1640 in flat-bottom 96-well plates in the presence of

25 ng/ml mouse rIL4 and 5 µg/ml anti–IFN-γ (XMG1.2). Anti-CD3/anti-CD28–dependent

activation (4 days) was followed by rest in 10 U/ml human recombinant IL2 for 2 days.

Titrations of anti-CD3ε antibody or recombinant IL2 were performed as indicated.

Recombinant cytokines were from PeproTech. For *Malat1* knockdown experiments, control or

Malat1-targeting antisense oligonucleotide GapmeRs were from QIAGEN (Hilden, Germany;

LG00000002-DDA and LG00000008-DDA, respectively) and were added to naive CD4$^+$ T

cells on day 0, or differentiating Th2 cells on day 4, at a final concentration of 100 nM.

Flow Cytometry

For flow cytometry analysis, single cell suspensions were generated. For live/dead

discrimination, cells were washed twice in PBS, then stained with Zombie Aqua (BioLegend)

in PBS before resuspension in FACS (Fluorescence-Activated Cell Sorting) buffer (PBS

containing 0.5% BSA and 0.05% azide). Fc receptors were blocked with 100 µg/ml rat IgG

(Sigma-Aldrich) for 10 min at 4°C before surface staining for 30 min at 4°C. The following anti-

mouse Abs from Biolegend were used: CD45.2 BV786 (clone 104); TCRβ PE-Cy7 (H57-597);

CD19 APC-Cy7 (6D5); MHC class II (MHCII) Alexa Fluor 700 (M5/114.15.2); Ly-6G APC-Cy7

(1A8); Ly6C BV605 (HK1.4); CD64 PE (X54-5/7.1); CD11b Pacific Blue (M1/70); CD44 FITC

(IM7); CD62L PE (MEL-14); CD8α PB (53-6.7); CD4 PerCP/Cy5.5 (RM4-5); IFN-γ FITC

(XMG1.2); IL10 PE (JES5-16E3); IL4 PE-Dazzle or APC (11B11); CD25 (IL-2Ra) PerCP-

Cy5.5 or APC (PC6.1); CD69 APC (H1.2F3); CD132 (IL-2Rg) PE (TUGm2); GATA3 PE-

Dazzle 594 (16E10A23) and streptavidin PE-Cy7. These were used in combination with abs

SiglecF PerCP-eFluor 710 (1RNM44N), iNOS PE-eFluor 610 (CXNET) and goat anti-rabbit

A647 from ThermoFisher Scientific, biotinylated Ym1 from R&D systems, and anti-murine

RELMα from PeproTech. To measure intracellular cytokines in T cells following *ex vivo*

stimulation, cells were first stimulated in complete RPMI 1640 for 4 h at 37°C with 500 ng/ml

PMA, 1 µg/ml ionomycin, and 10 µg/ml brefeldin A (all Sigma-Aldrich). For all intracellular

cytokine staining, surface stained cells were fixed and permeabilized (20 min at 4°C) using

Fixation/Permeabilization Solution before washes in Perm/Wash buffer (both BD

Biosciences). Cells were then stained with intracellular Abs as above except in Perm/Wash

buffer. For transcription factor staining, surface stained cells were fixed and permeabilized (1h

or overnight at 4°C) using the FoxP3 Fixation/Permeabilization solution before washes in 1x

Permeabilization Buffer (eBioscience). Cells were stained with intracellular Abs as above

except in Permeabilization Buffer. Appropriate isotype or fluorescence minus one (FMO)

controls were used to draw gates for populations of interest. Events were acquired on an

LSRFortessa (BD Biosciences) with analysis by FlowJo (FlowJo) or a Cytoflex LX (Beckman

Coulter) before analysis with or CytExpert.

RNA extraction, cDNA synthesis, and quantitative RT-PCR

RNA was extracted from purified cell populations using QIAzol and miRNeasy RNA extraction

kits (QIAGEN) according to manufacturer's instructions. For mRNA transcripts, reverse

transcriptions were carried out with Superscript III (Thermo Fisher Scientific) and random

hexamer primers (Promega) and measured with Fast SYBR Green Master Mix (Thermo Fisher

Scientific). Quantitative PCR was performed using a StepOnePlus Real Time PCR System or

a QuantStudio 3 (both Thermo Fisher Scientific), and relative transcript levels were

determined using the $\Delta\Delta$Ct method. The primers used for the study are show in **Supplemental**

Table S1.

RNA sequencing (RNAseq) and analysis

We used short-read (Illumina) RNAseq to compare naïve CD4$^+$ T cells and *in vitro*

differentiated Th2 cells from female and male *WT* and *Malat1$^{-/-}$* mice. Pooled cDNA libraries

were sequenced on one lane of an Illumina Novaseq 6000 S4 flow-cell at 100PE. The raw

sequencing reads were aligned to the mouse genome version Ensembl GRCm38.92 with

STAR 2.5.4a. We sequenced four replicates per condition. Samples with assigned sequencing

reads lower than 500,000 were removed from downstream analysis. Of the remaining samples

(N=4 for all *Malat1$^{-/-}$* samples and *WT* male Th2, N=3 for *WT* female and male naïve CD4$^+$ T

cells and *WT* female Th2), we obtained an average of 65 million reads per sample (range: 52–

91 million). Analysis was performed in RStudio (version 1.4.1106) with R (v 4.1). Data was

normalised ('calcNormFactors' and 'estimateDisp' functions of edgeR (version 3.14.0)) and a

quasi-likelihood negative binomial generalized log-linear model ('glmQLFit' and 'glmQLFTest'

functions with coefficient = 2) was used to determine differentially expressed genes. Adjusted

p values were calculated using the Benjamini and Hochberg method.

We used Oxford Nanopore Technologies (ONT) long-read RNAseq to compare naïve CD4$^+$ T

cells and cells after 2 days of *in vitro* differentiation from female *WT* and *Malat1$^{-/-}$* mice (N=4

per group). Full length cDNA libraries were prepared using the ONT cDNA-PCR Sequencing

V14 - Barcoding kit (SQK-PCB114.24). Briefly, cDNA RT adapters were ligated to 3' ends

(polyA tails) of transcripts, prior to stand switching cDNA synthesis and a 13 cycle PCR

reaction with barcoded primers, labelling each cDNA sample with a unique DNA barcode.

Barcoded cDNAs were pooled at equimolar ratios before final adapter ligation and sequencing

on R10.4.1 flowcells in ONT PromethION sequencer (8 samples per flow cell). Live

superaccuracy basecalling and barcode demultiplexing were performed in MinKNOW

software (version 24.06.10) at the time of the run. We acquired 0.745-1.121 million reads per

sample, with 604-840 bases mean read length. Demultiplexed reads were analyzed through

the workflow Transcriptomes from Epi2Me^{TM} application, specifically designed by ONT. The

workflow generates differential gene expression data (with edgeR), as well as differential

transcript usage (DTU) data (with Dexseq), using a reference transcriptome. The Viking

cluster, provided by the University of York, was used for the analyses.

Statistics

Experiments were typically performed in 2 – 3 independent replicates in groups of N = 3 – 5

mice per condition and replicate. Statistical analyses were carried out as indicated with Prism

5 (GraphPad Software). Two-way comparisons used paired or unpaired t tests as indicated,

and multiple comparisons used one-way ANOVA, followed by Sidak's multiple comparisons

test for comparison of samples with biological relevance (e.g. *WT* female to *Malat1$^{-/-}$* female,

WT male to *Malat1$^{-/-}$* male, *WT* female to *WT* male). Confidence levels were set to 0.05 for

significance. In merged data where n>9, consensus between Anderson-Darling, Shapiro-Wilk

and D'Agostino & Pearson tests were used to determine normality. If all samples were

normally distributed one-way ANOVA was performed as described. If one or more samples

did not follow a normal distribution, a Kruskal-Wallis test followed by Dunn's multiple

comparison test were performed to identify significance, with confidence levels of 0.05. P

values are displayed as asterisks representing P value classification, where * ≤ 0.05, ** ≤ 0.01,

≤ 0.001 and * ≤ 0.0001, unless the results are borderline non-significant, in which case

the p value will be shown.

Data availability

RNAseq data have been deposited in the National Center for Biotechnology (NCBI) Gene

Expression Omnibus (GEO) database. Accession numbers: GSE279185 for Illumina RNAseq

and GSE278413 for Oxford Nanopore Technologies RNAseq data.

Results

Malat1 **loss impairs** *in vitro* **Th2 differentiation of CD4$^+$ T cells from female mice**

We first assessed the effect *of Malat1* deletion on Th2 differentiation *in vitro* in response to

stimulation with anti-CD3 and anti-CD28 antibodies in the presence of recombinant IL4 and

anti-IFNγ antibody for 4 days, followed by IL2 treatment for 2 days. This protocol allowed us

to dissect potential effects on naïve CD4$^+$ T cell activation (0-24h), early differentiation (2-4

days), and terminal differentiation in response to exogenous IL2 (**Fig. 1A**). Following *in vitro*

differentiation of naive CD4+ T cells from female mice to Th2 cells, *Malat1$^{-/-}$* cells displayed

significantly lower percentages of IL10$^+$ and IL4$^+$/IL10$^+$ cells at day 6 of differentiation, with no

differences in IL4$^+$/IL10$^-$ cells, and no differences at day 4 (**Fig. 1B – F** and **Supplemental**

Fig. 1A). This decrease in cytokine expression was not observed from cells derived from male

mice. Under these conditions, differentiation of *WT* CD4$^+$ T cells from male mice was less

efficient compared to female *WT* cells, and *Malat1* deletion did not affect this further (**Fig. 1B**

– F). RNAseq analysis of *WT* and *Malat1$^{-/-}$* naïve CD4$^+$ T cells and *in vitro* differentiated Th2

cells revealed only a small number of genes were statistically significantly differentially

expressed between *WT* and *Malat1*$^{-/-}$ cells (**Supplemental Table S2**). As shown before (21),

IL10 levels were suppressed in *Malat1*$^{-/-}$ Th2 cells compared to *WT* controls (Log$_2$ fold change

LFC = - 0.882, p = 0.034, for female-derived cells and LFC = 0.03, p = 0.932 for male-derived

cells), although this did not reach significance after multiple testing correction. This was further

confirmed by qRTPCR for the sequenced samples (**Supplemental Fig. S1B**), suggesting that

this bulk RNAseq analysis provided a conservative representation of the effect of *Malat1*

deficiency on Th2 gene expression. Focusing on genes that were differentially expressed

between Th2 and naïve CD4$^+$ T cells from *WT* mice (Th2-differentiation associated genes) we

observed a transcriptome-wide blunting of Th2 gene expression upon *Malat1* loss in female

but not male cells (**Fig. 1G – H** and **Supplemental Tables S3 – S5**). Analysis of all

(FDR<0.001) or the top (by FDR) 500 differentially expressed genes (DEGs) between *WT* Th2

and naïve CD4$^+$ T cells demonstrated that most genes that were up-regulated during *WT* Th2

differentiation were overall expressed at lower levels when comparing *Malat1*$^{-/-}$ Th2 cells to

WT Th2 cells, and, conversely, genes that were suppressed upon *WT* Th2 differentiation were

overall more highly expressed in *Malat1*$^{-/-}$ Th2 cells (**Fig. 1G** and **Supplemental Tables S3 –**

S5). This transcriptome-wide trend was not observed in male cells (**Fig. 1H**). Taking a stricter

threshold for DEGs between Th2 and naïve CD4$^+$ T cells from female mice (FDR<0.001 and

absolute LFC>2), we found 524 Th2-differentiation associated DEGs in both *WT* and *Malat1*$^{-}$

$^{/-}$, 469 DEGs only in *WT*, and 139 DEGs only in *Malat1*$^{-/-}$ Th2 cells. We noted that common

and *WT* only Th2 DEGs were mainly up-regulated, whereas *Malat1*$^{-/-}$ only Th2 DEGs were

mainly down-regulated (**Supplemental Fig. S1C – D**, **Supplemental Tables S3 – S5**).

Furthermore, when looking at common DEGs, we found that the vast majority of up-regulated

genes showed a smaller LFC in *Malat1*$^{-/-}$ Th2 cells compared to *WT* (**Supplemental Fig. S1E**).

Overall, these results indicated a *Malat1*-mediated effect in Th2 differentiation of female- but

not male-derived naïve CD4$^+$ T cells, characterised by suppression of IL10 expression and an

impairment of the Th2 gene expression programme.

The effect of *Malat1* depletion on IL10 expression is independent of activation strength and occurs during the early stages of differentiation

We next tested whether *Malat1* deficiency affects Th2 polarization under weaker activation conditions. We found that *Malat1*$^{-/-}$ cells displayed lower IL10 expression under suboptimal differentiation conditions, whereas IL4 levels were not statistically significantly different (**Fig. 2A – B**, **Supplemental Fig. S2A – B**). Of note, at lower activation levels there were no differences between *in vitro* Th2 differentiation of male and female cells. This indicated that *Malat1* deficiency can affect Th2 cytokine expression independently of activation strength.

We next tested whether *Malat1* knockdown rather than genetic deletion had a sex-specific effect on Th2 differentiation. As described previously (21), we used GapmeR oligonucleotides to suppress *Malat1* expression. Although, Th2 cells express lower *Malat1* levels than naïve CD4$^+$ T cells ((21) and **Supplemental Fig. S2C**), we found that *Malat1* levels increase between day 4 and 6 of differentiation, upon cessation of CD3 activation and addition of exogenous IL2 (**Fig. 2C**). Based on this and the observation that *Malat1* deficiency impaired cytokine induction between days 4 and 6 (**Fig. 1C – F**), we knocked down *Malat1* either on day 0, or by adding GapmeRs on day 4, at the same time with exogenous IL2 stimulation. *Malat1* knockdown reduced IL10 expression in female Th2 cells only if GapmeRs were added to naïve CD4$^+$ T cells, but not when added concurrently with exogenous IL2 stimulation. No effects on cytokine expression were observed upon *Malat1* knockdown in male cells (**Fig. 2D**). Partial knockdown of *Malat1* in cells from female mice did not affect IL4 expression (**Supplemental Fig. S2D)**. This demonstrated that the effect of *Malat1* on Th2 cytokine expression required its activity during the early differentiation stages.

Malat1 loss alters the kinetics of naïve CD4$^+$ T cell activation and impairs early Th2 differentiation

Based on the above, we tested the effect of *Malat1* deletion on the early stages of Th2 differentiation. First, we studied the first 24h of naïve CD4$^+$ T cell activation by measuring levels of T cell early activation marker CD69. As expected, *Malat1* levels were suppressed

within 24h in both male and female *WT* mice (**Supplemental Fig. S3A**). In female cells, we

found that compared to *WT* cells CD69 protein expression was higher in *Malat1$^{-/-}$* cells at 3h

post activation, the difference reaching statistical significance at 7h post activation. However,

CD69 expression was significantly lower in *Malat1$^{-/-}$* cells at 24h (**Fig. 3A – B**). At the transcript

level, *Cd69* levels peaked at 3h post activation, with significantly higher levels in *Malat1$^{-/-}$* cells

(**Supplemental Fig. S3B**). In male cells, we did not observe any statistically significant

differences between *WT* and *Malat1$^{-/-}$* cells, although we noted that male *WT* cells showed

statistically significant higher CD69 levels compared to female counterparts at 3h post

activation (**Fig. 3A – B**). We also measured levels of IL2Rα (CD25), which was induced upon

activation but found no differences between genotypes or biological sexes (**Supplemental**

Fig. S3C).

To explore the consequences of the altered activation kinetics in female *Malat1$^{-/-}$* cells, we

performed a daily timecourse (days 1 – 4), measuring transcript levels of key transcription

factors (*Gata3*, *Maf*, and *Tbet*) and differentiation-associated genes (*Il2, Il2ra, Il2rg*). We

focused on *Il2ra* and *Il2rg* as the former is induced during differentiation and the latter is an X-

linked gene, given the sex-specificity of the observed effects. We noticed suppression for

multiple of the measured transcripts in female *Malat1$^{-/-}$* differentiating cells, including *Il2*, *Il2ra*,

Il2rg, and *Gata3* on days 2 and 3 of differentiation (**Fig. 3C** and **Supplemental Fig. S3D**).

These initial findings were confirmed in independent experiments for *Il2ra*, *Il2rg*, and *Gata3*

on day 2 (**Fig. 3D**). We chose day 2 for the validation experiments as this was the earliest

timepoint the suppression of the Th2 programme was observed. In male *Malat1$^{-/-}$* cells, we

observed higher variability and similar but non-significant trends in the levels of the tested

transcripts to those observed in female cells (**Fig. 3D**). In agreement with the mRNA levels,

we found that GATA3 protein levels were suppressed at day 2 of the differentiation in *Malat1-*

/- cells (**Fig. 3E** and **Supplemental Fig, S3E**), whereas at this point we did not find any

significant differences in the levels of the IL2R subunits (**Supplemental Fig. S3F**). These

results suggested that *Malat1* was necessary for appropriate early Th2 differentiation.

Impairment of early differentiation in *Malat1*^{-/-} cells is characterised by up-regulation of an interferon-responsive gene signature

To further explore the role of *Malat1* in early Th2 differentiation we performed long-read RNAseq of *WT* and *Malat1*^{-/-} naïve CD4$^+$ T cells and differentiating cells at day 2 of differentiation. As *Malat1* has been suggested to be involved in regulation of mRNA splicing (14), we opted for long-read RNAseq to facilitate concurrent assessment of differential gene expression and differential transcript usage (DTU). As in the case of our short read RNAseq analysis, there were only few differences between *WT* and *Malat1*^{-/-} naïve CD4$^+$ T cells (16 DEGs at FDR<0.1, **Supplemental Fig. S4A** and **Supplemental Table S6**). However, comparison of *WT* and *Malat1*^{-/-} cells on day 2 of differentiation revealed 239 DEGs, 199 of which were up-regulated in *Malat1*^{-/-} cells (**Fig. 4A** and **Supplemental Table S7**). Gene set enrichment analysis (GSEA; (29)), revealed a profound over-representation of interferon alpha and interferon gamma response gene signatures amongst the up-regulated genes in *Malat1*^{-/-} cells on day 2 (**Fig. 4B**), whereas enriched terms amongst down-regulated genes, including hypoxia, only reached modest significance (**Supplemental Fig. S4B**). We note that in agreement with our qRTPCR analysis (**Fig. 3D**), trends towards down-regulation were observed for *Il2ra* (LFC = -0.589) and *Gata3* (LFC = -0.401) but these did not reach statistical significance. The most prominent feature of *Malat1*^{-/-} cells, the up-regulated interferon gene cluster (**Supplemental Fig. S4C** and **Supplemental Table S8**; clusters identified by k-means clustering) included interferon stimulated genes (e.g. *Ifit1bl1*, *Ifit3*, *Ifit1*, *Oas2*, and others) and transcription factors including *Irf7*, *Irf1*, and *Stat1* (**Fig. 4C**, **Supplemental Fig. S4C** and **Supplemental Tables S7 and S8**). Most of the day 2 *Malat1*-associated DEGs were down-regulated during differentiation of *WT* cells both when comparing day 2 with naïve cells (**Fig. 4D**) and when comparing *WT* Th2 and naïve CD4$^+$ T cells using our short-read sequencing data (**Fig. 4E**). This was also the case specifically for the interferon gene module (**Supplemental Fig. S4D**). As in the case of Th2 cells (**Fig. 1G**) *Malat1* deficiency had a transcriptome-wide effect by blunting changes in gene expression occurring in *WT* cells within the first two days of differentiation (**Supplemental Fig. S4E**).

Validation of the long-read sequencing data by qRTPCR for selected genes confirmed up-

regulation of *Ifit1bl1*, *Ifit3b*, *Iigp1*, *Ifit1*, and *Irf7* (**Fig. 4F**). Up-regulation trends for Irf1 and Stat1

did not reach statistical significance. Notably, no effects due to *Malat1* deficiency were

observed in male CD4$^+$ T cells, and levels of *Ifit1bl1*, *Ifit3b*, and *Irf7* were significantly higher

in male *WT* cells compared to female cells on day 2 (**Fig. 4F**). Comparing day 2 to naïve CD4$^+$

T cells, we found significant down-regulation for *Ifit1bl1*, *Ifit1*, and *Irf7* for all conditions.

Statistically significant *Stat1* down-regulation on day 2 was only observed for female *WT* cells

(**Supplemental Fig. S4F**). *Malat1* deficiency had weak effects on DTU affecting only 25 genes

(FDR<0.1, **Supplemental Table S9**) including some cytokine genes, but demonstrating

modest effects (**Fig. 4G** and **Supplemental Fig. S5**). In some instances, as for *Kdm2a* and

Trim12a (**Fig. 4G**) isoform usage changes due to *Malat1* loss showed opposing patterns to

those induced by early differentiation of *WT* cells. Overall, these analyses further supported

that *Malat1* is necessary for early Th2 differentiation with the predominant effect being

impaired suppression of an interferon responsive gene cluster in differentiating cells.

Female differentiating Th2 cells are sensitive to suppression of IL2R caused by *Malat1*

deficiency

Having shown that *Malat1* deficiency impairs early Th2 differentiation of CD4$^+$ T cells, we

reasoned that this could affect their readiness to respond to exogenous IL2. Having found

decreases in the mRNA levels of the IL2R subunits (**Fig. 3D**), we measured protein levels of

IL2Rα and IL2Rγ on day 4, prior to addition of exogenous IL2. We found that IL2Rα was

statistically significantly down-regulated only in female *Malat1$^{-/-}$* T cells on day 4 (**Fig. 5A – B**).

IL2Rγ was down-regulated in both female and male *Malat1$^{-/-}$* T cells (**Fig. 5C – D**). No

differences were observed at the mRNA level at this timepoint (**Supplemental Fig. S6A**).

Similarly, IL2Rα was significantly down-regulated on day 4 in female cells treated with *Malat1*-

targeting GapmeRs (**Fig. 5E**) and this was also the case for IL2Rγ (**Fig. 5F**). We note that in

male cells, there was a non-significant trend towards down-regulation for IL2Rγ, suggesting

that a more profound reduction in *Malat1* levels might be required for this effect in male cells

(**Fig. 5F**). The observed down-regulation in the IL2R subunits at this time-point in female

Malat1$^{-/-}$ cells can explain the impaired induction of cytokine expression between day 4 and

day 6 (**Fig. 1C – F**). This also indicates that starting from the same baseline at day 4 with

regards to IL4 and IL10 expression, *WT* male cells were less responsive to exogenous IL2

compared to female cells. We further tested the dependence of female and male *WT* cells on

IL2 stimulation by titrating recombinant IL2. We found that although male cells expand as well

or, at the lowest IL2 concentration, better than female cells (**Supplemental Fig. S6B**), they

demonstrated lower IL10 expression than female cells at all concentrations (**Fig. 5G**). The

observed differential regulation of IL2R receptor and dependence on IL2 for cytokine

expression could, at least in part, explain why only female Th2 differentiation was affected by

Malat1 deficiency.

***Malat1* loss impairs Th2 cell cytokine expression only in female mice in an experimental**

model of type 2 inflammation

Having demonstrated female-specific effects on Th2 cytokine expression and early

differentiation due to *Malat1* deficiency in a cell-intrinsic context *in vitro*, we tested the effect

of *Malat1* loss in an experimental model of type 2 inflammation *in vivo*. Intravenous *S. mansoni*

egg challenge in mice previously primed with schistosome eggs (**Fig. 6A**) leads to the

formation of type 2 inflammatory granulomas in the lungs and mimics helminth-induced lung

inflammation (26, 27) leading to cellular infiltration in the lungs, including Th2 cells, and

eosinophilia. Indeed, *S. mansoni* egg injection resulted in increased in lung cell numbers,

although we note that this only reached statistical significance in *WT* female mice and *Malat1^{-}*

$^{/-}$ male mice (**Supplemental Fig. S7** and **S8A**). Similar splenic cell numbers were observed

under all conditions (**Supplemental Fig. S8B**) and similar lung eosinophil numbers and

activation, measured by RELMα expression (**Supplemental Fig. S8C – D**). Notably, both lung

and splenic CD4$^+$ T cells in egg-injected female *Malat1$^{-/-}$* mice expressed lower levels of IL10

than *WT* mice (**Fig. 6B – D**). IL4$^+$/IL10$^+$ cells were also decreased in the lungs (percentage)

and spleens (percentage and numbers) of egg-injected $Malat1^{-/-}$ mice (**Fig. 6E – F**). This was also the case for triple positive IL4$^+$/IL13$^+$/IL10$^+$ Th2 cells in the lungs of egg-injected mice (**Supplemental Fig. S8E**). None of these differences were observed when comparing male $Malat1^{-/-}$ mice to *WT* controls. On the contrary the percentage of splenic IL4$^+$/IL10$^+$ cells increased in egg-injected $Malat1^{-/-}$ male mice. Under these conditions, we did not observe any differences in total IL4 expression between egg injected $Malat1^{-/-}$ and *WT* female mice, although we noted an increase in IL4-expressing splenic CD4$^+$ T cells in male $Malat1^{-/-}$ mice compared to *WT* (**Supplemental Fig. S8F – G**). IFNγ^+ CD4$^+$ T cells are also induced as a result of *S. mansoni* egg injection, and IL10 levels were also reduced in female-specific manner within this population in the lungs of inflamed mice (**Supplemental Fig. S8H – I**). With regards to activation status of CD4$^+$ T cells, we observed a borderline non-significant decrease in activated (CD44^{high}/CD64L^{low}) lung CD4$^+$ T cells in female $Malat1^{-/-}$ mice compared to *WT* and no differences in the spleen (**Supplemental Fig. S8J – K**). No differences were observed in percentage and activation status of myeloid cell populations in egg-injected mice (**Supplemental Fig. S8L – M**). Overall, these results indicated that, as observed in our *in vitro* Th2 differentiation experiments, *Malat1* deficiency resulted in a female-specific impairment of Th2 cytokine expression, predominantly demonstrated by reduced IL10 expression in Th2 cells.

Discussion

Beyond hormonal or sex chromosome-driven mechanisms, little is known about cell intrinsic mediators of sexual dimorphism in lymphocytes (1-4). We demonstrate that *Malat1*, one of the most highly abundant transcripts in mammalian cells, exerts a sex-specific function in Th2 differentiation, affecting early differentiation and end-point expression of cytokine IL10. It is of interest that the female-specific effect due to *Malat1* deficiency is observed in the *S. mansoni* egg injection model, as biological sex is a determinant of immune responses in the context of human schistosomiasis (30). We also note that, although *Malat1* loss is associated with suppression of IL10 both *in vitro* and *in vivo*, other effects of *Malat1* deficiency, for example a

modest suppression of IL4, are only observed *in vitro*. This can be due to compensatory

mechanisms and multiple effects contributing to cytokine expression *in vivo*. For example, IL4

levels in *Malat1^-/-* mice are likely to reflect both the loss of *Malat1* and reduced IL10 levels from

Th2 cells and potentially other cell types, as IL10 has been shown to suppress activation of

Th2 cells *in vivo* (31). In addition, IL10 has been shown to be expressed later than other

effector cytokines (32) and from heterogeneous populations of CD4[+] T cells (33). As such,

discrepancies between *in vitro* and *in vivo* cytokine expression can be due to the time-point

assessed and kinetics of Th2 cell emergence in the *S. mansoni* egg injection model.

Nevertheless, the observed impairment of cytokine expression during *in vitro* Th2

differentiation and in the *S. mansoni* egg injection model strongly support a cell-intrinsic

function of *Malat1* in controlling Th2 differentiation in a sex-specific manner.

Despite the limitation of not capturing the complexities of *in vivo* Th2 cell differentiation, *in vitro*

differentiation of naïve CD4[+] T cells to Th2 cells allowed us to demonstrate that *Malat1* loss

results in a sex-specific impairment of differentiation in a cell-intrinsic manner. Mechanistically,

down-regulation of IL2R upon *Malat1* loss, prior to exposure to exogenous IL2, results in a

transcriptome-wide suppression of the Th2 differentiation programme. Although down-

regulation of IL2Rγ is observed in both male and female cells, IL2Rα, which is necessary for

formation of the high affinity IL2R (34), is only suppressed in female cells. Given the direct

involvement of IL2 in Th2 differentiation (35) this can at least partly explain the selective effect

of *Malat1* loss in female cells. IL2R signaling has been linked specifically to IL10 expression,

via activation of STAT5 and MAF (36), which might explain the stronger impact observed on

IL10 over other cytokines upon loss of *Malat1*. This was also seen here, as resuspension in

exogenous IL-2 produced a larger increase in IL10 expression compared to IL4 (**Fig. 1C – D**).

In vitro, male Th2 differentiating cells are less responsive to IL2 with regards to cytokine

expression, providing an additional potential explanation of the sex-specificity of the observed

effect. Evidence for sexual dimorphism in response to IL2 in other immune cell types has been

already reported. Female group 2 Innate Lymphoid Cells demonstrate increased proliferation

in response to IL2 when compared to male ILC2 cells (37), and female NK cells produce higher

levels of IFNγ than males in response to increased levels of IL2 (38).

The observed downregulation of IL2Rα and IL2Rγ on day 4 is preceded by impaired early

differentiation (days 2-3) of *Malat1*[-/-] cells, an effect which is more pronounced in female cells

and includes down-regulation of the master transcriptional regulator GATA3 (39) and impaired

suppression of an interferon-responsive gene cluster. It is well established that GATA3

suppresses IFN expression and signaling in Th2 cells (40, 41). It is also notable that Th cells

demonstrating interferon signatures (including expression of *Ifit3*, *Irf7*, *Stat1*, but not *Ifng*) have

been described in type 2 inflammation both in humans (42) and in mouse models (43). In all

cases, these Th cells represent a distinct population to Th2 cells indicating that this signature

needs to be suppressed in Th2 cells, which in agreement with our findings in *WT* cells.

Interestingly a recent publication identified an interferon-experienced population of naïve CD4[+]

T cells that show impaired response to TCR stimulation, as seen here for *Malat1*[-/-] cells (44).

Another report demonstrated that these cells exist constitutively including in germ free mice

(45). The 50-gene signature that defines interferon-experienced naïve CD4[+] T cells (44)

displays substantial overlap with the *Malat1*-regulated gene cluster (at day 2) identified here

(26/50 genes, including *Irf7* and *Stat1*). Taken together this suggests that suppression of this

interferon signature in naïve CD4[+] T cells is necessary for optimal differentiation and regulated

by *Malat1*. It will be interesting to further investigate how *Malat1* promotes specifically down-

regulation of interferon-responsive genes in female cells during early CD4[+] T cell

differentiation. In this respect, it has been shown that *Malat1* can promote epigenetic silencing

of specific genes in CD8[+] T cells (25).

Our results suggest that the impairment of early Th2 differentiation in female *Malat1*[-/-] cells is

associated with a stronger but shorter-lived initial activation (0-24h) as measured by CD69

expression. Indeed, strong TCR stimulation in the presence of CD28 co-stimulation has been

previously linked to impaired Th2 differentiation (46). During the first 24h of activation, male

WT cells demonstrate CD69 expression kinetics similar to female *Malat1$^{-/-}$* cells, while deletion

of *Malat1* does not cause any further increase in CD69 expression in male cells. Overall, we

propose that *Malat1* downregulation upon activation of naïve CD4$^+$ T cells facilitates activation,

potentially through release of associated proteins that contribute co-transcriptional

processing. However, in female cells, complete loss of *Malat1* alters activation kinetics, which

in combination with failure to suppress expression of interferon-responsive genes, leads to

impaired Th2 differentiation and subsequent suppression of IL2R and IL2-mediated cytokine

expression. Male cells are not sensitive to *Malat1* loss, at least partly, due to their higher early

activation threshold, higher expression of interferon-responsive genes during early

differentiation, and, lower sensitivity to exogenous IL2 during the later stages of differentiation.

We cannot exclude additional female-specific mechanisms being affected by *Malat1*, for

example with regards to regulation of X Chromosome-linked genes and the role of X

chromosome inactivation during Th2 differentiation (10). Furthermore, it will be essential for

our results to be validated in other models of type 2 immunity and in human CD4$^+$ T cells.

Despite these limitations, our study reveals that female Th2 cells show a specific susceptibility

to *Malat1*, a highly expressed lincRNA that is neither responsive to sex hormones nor located

on a sex chromosome. This can have far reaching implications for our understanding of

immune sexual dimorphism and provide novel routes for sex-specific manipulation of adaptive

immunity.

Author contributions

MG and KAW contributed to experimental design, data generation, data analysis, figure

design, and manuscript writing. SvD and KRJ generated and analysed Illumina RNAseq

experiments. IK, DC, and SAT contributed to study supervision and data interpretation. JPH

contributed to study design, data generation, and data interpretation. DL conceived and

coordinated the study, contributed to study design, data interpretation, and study supervision,

and wrote the manuscript. All authors contributed to manuscript editing and approved the final manuscript. MG and KAW share the first position in the author list and are listed alphabetically.

Acknowledgements

This work was funded by the Hull York Medical School (PhD studentship to MG and additional funding to DL), the Biotechnology and Biological Sciences Research Council (BBSRC) White Rose doctoral training partnership (to KAW, reference BB/J014443/1), and the Medical Research Council (to JPH, reference MR/W018578/1). We thank Sally James and Fabiano Pais at the Genomics Lab and Data Science Hub, respectively, in the University of York Bioscience Technology Facility for support with long-read sequencing data. We thank Joanna Greenman for technical support and Allison Green for comments on the manuscript.

References

1. Gal-Oz ST, Maier B, Yoshida H, Seddu K, Elbaz N, Czysz C, et al. ImmGen report: sexual dimorphism in the immune system transcriptome. *Nat Commun.* 2019;10(1):4295.

2. Klein SL, and Flanagan KL. Sex differences in immune responses. *Nat Rev Immunol.* 2016;16(10):626-38.

3. Marquez EJ, Chung CH, Marches R, Rossi RJ, Nehar-Belaid D, Eroglu A, et al. Sexual-dimorphism in human immune system aging. *Nat Commun.* 2020;11(1):751.

4. Takahashi T, and Iwasaki A. Sex differences in immune responses. *Science.* 2021;371(6527):347-8.

5. Dodd KC, and Menon M. Sex bias in lymphocytes: Implications for autoimmune diseases. *Front Immunol.* 2022;13:945762.

6. Rubtsova K, Marrack P, and Rubtsov AV. Sexual dimorphism in autoimmunity. *J Clin Invest.* 2015;125(6):2187-93.

7. Whitacre CC, Reingold SC, and O'Looney PA. A gender gap in autoimmunity. *Science.* 1999;283(5406):1277-8.

8. Shepherd R, Cheung AS, Pang K, Saffery R, and Novakovic B. Sexual Dimorphism in

Innate Immunity: The Role of Sex Hormones and Epigenetics. *Front Immunol.*

2020;11:604000.

9. Lambert KC, Curran EM, Judy BM, Milligan GN, Lubahn DB, and Estes DM. Estrogen

receptor alpha (ERalpha) deficiency in macrophages results in increased stimulation of CD4+

T cells while 17beta-estradiol acts through ERalpha to increase IL-4 and GATA-3 expression

in CD4+ T cells independent of antigen presentation. *J Immunol.* 2005;175(9):5716-23.

10. Syrett CM, Paneru B, Sandoval-Heglund D, Wang J, Banerjee S, Sindhava V, et al.

Altered X-chromosome inactivation in T cells may promote sex-biased autoimmune diseases.

JCI Insight. 2019;4(7).

11. Dou DR, Zhao Y, Belk JA, Zhao Y, Casey KM, Chen DC, et al. Xist ribonucleoproteins

promote female sex-biased autoimmunity. *Cell.* 2024;187(3):733-49 e16.

12. Engreitz JM, Ollikainen N, and Guttman M. Long non-coding RNAs: spatial amplifiers that

control nuclear structure and gene expression. *Nat Rev Mol Cell Biol.* 2016;17(12):756-70.

13. Gutschner T, Hammerle M, Eissmann M, Hsu J, Kim Y, Hung G, et al. The noncoding

RNA MALAT1 is a critical regulator of the metastasis phenotype of lung cancer cells. *Cancer*

Res. 2013;73(3):1180-9.

14. Tripathi V, Ellis JD, Shen Z, Song DY, Pan Q, Watt AT, et al. The nuclear-retained

noncoding RNA MALAT1 regulates alternative splicing by modulating SR splicing factor

phosphorylation. *Mol Cell.* 2010;39(6):925-38.

15. Lamond AI, and Spector DL. Nuclear speckles: a model for nuclear organelles. *Nat Rev*

Mol Cell Biol. 2003;4(8):605-12.

16. Tripathi V, Shen Z, Chakraborty A, Giri S, Freier SM, Wu X, et al. Long noncoding RNA

MALAT1 controls cell cycle progression by regulating the expression of oncogenic

transcription factor B-MYB. *PLoS Genet.* 2013;9(3):e1003368.

17. Eissmann M, Gutschner T, Hammerle M, Gunther S, Caudron-Herger M, Gross M, et al.

Loss of the abundant nuclear non-coding RNA MALAT1 is compatible with life and

development. *RNA Biol.* 2012;9(8):1076-87.

18. Nakagawa S, Ip JY, Shioi G, Tripathi V, Zong X, Hirose T, et al. Malat1 is not an essential

component of nuclear speckles in mice. *RNA.* 2012;18(8):1487-99.

19. Zhang B, Arun G, Mao YS, Lazar Z, Hung G, Bhattacharjee G, et al. The lncRNA Malat1

is dispensable for mouse development but its transcription plays a cis-regulatory role in the

adult. *Cell Rep.* 2012;2(1):111-23.

20. Fei J, Jadaliha M, Harmon TS, Li ITS, Hua B, Hao Q, et al. Quantitative analysis of

multilayer organization of proteins and RNA in nuclear speckles at super resolution. *J Cell Sci.*

2017;130(24):4180-92.

21. Hewitson JP, West KA, James KR, Rani GF, Dey N, Romano A, et al. Malat1 Suppresses

Immunity to Infection through Promoting Expression of Maf and IL-10 in Th Cells. *J Immunol.*

2020;204(11):2949-60.

22. Dey S, Ashwin H, Milross L, Hunter B, Majo J, Filby AJ, et al. Downregulation of MALAT1

is a hallmark of tissue and peripheral proliferative T cells in COVID-19. *Clin Exp Immunol.*

2023;212(3):262-75.

23. Masoumi F, Ghorbani S, Talebi F, Branton WG, Rajaei S, Power C, et al. Malat1 long

noncoding RNA regulates inflammation and leukocyte differentiation in experimental

autoimmune encephalomyelitis. *J Neuroimmunol.* 2019;328:50-9.

24. Xue Y, Ke J, Zhou X, Chen Q, Chen M, Huang T, et al. Knockdown of LncRNA MALAT1

Alleviates Coxsackievirus B3-Induced Acute Viral Myocarditis in Mice via Inhibiting Th17 Cells

Differentiation. *Inflammation.* 2022.

25. Kanbar JN, Ma S, Kim ES, Kurd NS, Tsai MS, Tysl T, et al. The long noncoding RNA

Malat1 regulates CD8+ T cell differentiation by mediating epigenetic repression. *J Exp Med.*

2022;219(6).

26. Kaplan MH, Whitfield JR, Boros DL, and Grusby MJ. Th2 cells are required for the

Schistosoma mansoni egg-induced granulomatous response. *J Immunol.* 1998;160(4):1850-

6.

27. Sabin EA, Kopf MA, and Pearce EJ. Schistosoma mansoni egg-induced early IL-4

production is dependent upon IL-5 and eosinophils. *J Exp Med.* 1996;184(5):1871-8.

28. Percie du Sert N, Ahluwalia A, Alam S, Avey MT, Baker M, Browne WJ, et al. Reporting

animal research: Explanation and elaboration for the ARRIVE guidelines 2.0. *PLoS Biol.*

2020;18(7):e3000411.

29. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene

set enrichment analysis: a knowledge-based approach for interpreting genome-wide

expression profiles. *Proc Natl Acad Sci U S A.* 2005;102(43):15545-50.

30. Eloi-Santos S, Olsen NJ, Correa-Oliveira R, and Colley DG. Schistosoma mansoni:

mortality, pathophysiology, and susceptibility differences in male and female mice. *Exp*

Parasitol. 1992;75(2):168-75.

31. Coomes SM, Kannan Y, Pelly VS, Entwistle LJ, Guidi R, Perez-Lloret J, et al. CD4(+) Th2

cells are directly regulated by IL-10 during allergic airway inflammation. *Mucosal Immunol.*

2017;10(1):150-61.

32. Yssel H, De Waal Malefyt R, Roncarolo MG, Abrams JS, Lahesmaa R, Spits H, et al. IL-

10 is produced by subsets of human CD4+ T cell clones and peripheral blood T cells. *J*

Immunol. 1992;149(7):2378-84.

33. Brockmann L, Soukou S, Steglich B, Czarnewski P, Zhao L, Wende S, et al. Molecular

and functional heterogeneity of IL-10-producing CD4(+) T cells. *Nat Commun.*

2018;9(1):5457.

34. Shouse AN, LaPorte KM, and Malek TR. Interleukin-2 signaling in the regulation of T cell

biology in autoimmunity and cancer. *Immunity.* 2024;57(3):414-28.

35. Cote-Sierra J, Foucras G, Guo L, Chiodetti L, Young HA, Hu-Li J, et al. Interleukin 2 plays

a central role in Th2 differentiation. *Proc Natl Acad Sci U S A.* 2004;101(11):3880-5.

36. Rani A, Afzali B, Kelly A, Tewolde-Berhan L, Hackett M, Kanhere AS, et al. IL-2 regulates

expression of C-MAF in human CD4 T cells. *J Immunol.* 2011;187(7):3721-9.

37. Cephus JY, Stier MT, Fuseini H, Yung JA, Toki S, Bloodworth MH, et al. Testosterone

Attenuates Group 2 Innate Lymphoid Cell-Mediated Airway Inflammation. *Cell Rep.*

2017;21(9):2487-99.

38. Menees KB, Earls RH, Chung J, Jernigan J, Filipov NM, Carpenter JM, et al. Sex- and

age-dependent alterations of splenic immune cell profile and NK cell phenotypes and function

in C57BL/6J mice. *Immun Ageing.* 2021;18(1):3.

39. Zheng W, and Flavell RA. The transcription factor GATA-3 is necessary and sufficient for

Th2 cytokine gene expression in CD4 T cells. *Cell.* 1997;89(4):587-96.

40. Ferber IA, Lee HJ, Zonin F, Heath V, Mui A, Arai N, et al. GATA-3 significantly

downregulates IFN-gamma production from developing Th1 cells in addition to inducing IL-4

and IL-5 levels. *Clin Immunol.* 1999;91(2):134-44.

41. Ouyang W, Ranganath SH, Weindel K, Bhattacharya D, Murphy TL, Sha WC, et al.

Inhibition of Th1 development mediated by GATA-3 through an IL-4-independent mechanism.

Immunity. 1998;9(5):745-55.

42. Seumois G, Ramirez-Suastegui C, Schmiedel BJ, Liang S, Peters B, Sette A, et al. Single-

cell transcriptomic analysis of allergen-specific T cells in allergy and asthma. *Sci Immunol.*

2020;5(48).

43. Tibbitt CA, Stark JM, Martens L, Ma J, Mold JE, Deswarte K, et al. Single-Cell RNA

Sequencing of the T Helper Cell Response to House Dust Mites Defines a Distinct Gene

Expression Signature in Airway Th2 Cells. *Immunity.* 2019;51(1):169-84 e5.

44. Deep D, Gudjonson H, Brown CC, Rose SA, Sharma R, Paucar Iza YA, et al. Precursor

central memory versus effector cell fate and naive CD4+ T cell heterogeneity. *J Exp Med.*

2024;221(10).

45. Even Z, Meli AP, Tyagi A, Vidyarthi A, Briggs N, de Kouchkovsky DA, et al. The amalgam

of naive CD4(+) T cell transcriptional states is reconfigured by helminth infection to dampen

the amplitude of the immune response. *Immunity.* 2024;57(8):1893-907 e6.

46. Tao X, Constant S, Jorritsma P, and Bottomly K. Strength of TCR signal determines the

costimulatory requirements for Th1 and Th2 CD4+ T cell differentiation. *J Immunol.*

1997;159(12):5956-63.

Figure Legends

Figure 1. *Malat1* loss impairs *in vitro* Th2 differentiation only in female cells.

A. Schematic of *in vitro* Th2 differentiation protocol.

B. Representative FACS plots of IL10 and IL4 expression in *WT* and *Malat1^-/-^* female and male *in vitro* differentiated Th2 cells (Day 6).

C. Percentage of IL10$^+$ live TCRb$^+$ CD4$^+$ in *in vitro* differentiated Th2 cells derived from female and male mice at day 4 (prior to IL2 addition) and day 6 (experimental end-point). Levels determined by intracellular cytokine staining (N=14 per condition).

D. As in **B**, but for IL4$^+$ live TCRb$^+$ CD4$^+$ cells.

E. As in **B**, but for IL4$^+$/IL10$^+$ live TCRb$^+$ CD4$^+$ cells.

F. As in **B**, but for IL4$^+$/IL10$^-$ live TCRb$^+$ CD4$^+$ cells.

G. Log$_2$ fold change (LFC) in gene expression between *WT* female naïve and *in vitro* differentiated Th2 cells against LFC in gene expression between *WT* and *Malat1^-/-^* female *in vitro* differentiated Th2 cells. Data shown for all or the top 500 DEGs between Th2 and naïve CD4$^+$ T cells from *WT* female mice.

H. As in **E**, but for cells derived from male mice.

Figure 2. The effect of *Malat1* depletion on IL10 expression is independent of activation strength and occurs during the early stages of differentiation.

A. Percentage of IL10$^+$ live TCRb$^+$ CD4$^+$ *WT* or *Malat1^-/-^ in vitro* differentiated Th2 cells derived from female and male mice, with 10 μg/ml) or 2 μg/ml levels of anti-CD3 antibody (see Methods). Levels determined by intracellular cytokine staining (N=9).

B. As in **A**, but for IL4$^+$ live TCRb$^+$ CD4$^+$ cells.

C. *Malat1* levels in *in vitro* differentiated Th2 cells at day 4 (prior to resuspension in IL2) and at day 6 (experimental end point), determined by qRTPCR. Levels normalised to U6 and average levels of *WT* females at day 4 (N=11).

D. Percentage of IL10$^+$ live TCRb$^+$ CD4$^+$ *in vitro* differentiated *WT* Th2 cells derived from female and male mice, treated with either non-targeting control (NTC) GapmeRs or *Malat1* targeting GapmeRs. Levels determined by intracellular cytokine staining (N=6).

Figure 3. *Malat1* loss enhances naïve CD4+ T cell activation and impairs early Th2 differentiation.

A. Representative dot plots of CD69 expression in *WT* or *Malat1*$^{-/-}$ female or male CD4+ T cells at 3 hours post-activation with anti-CD3 and anti-CD28 antibodies.

B. Percentage of CD69+ live TCRb+ CD4+ *WT* or *Malat1*$^{-/-}$ *in vitro* activated CD4+ T cells derived from female or male mice at 0, 3, 7, and 24 hours post-activation. Levels determined by surface staining (N=6).

C. *Il2ra*, *Il2rg*, *Il2* and *Gata3* mRNA levels at days 1, 2, 3 and 4 of Th2 differentiation determined by qRTPCR (N=3). Levels normalised to U6 and average levels of *WT* females at day 1.

D. *Il2ra*, *Il2rg* and *Gata3* mRNA levels at day 2 of differentiation determined by qRTPCR (N=8 for WT females, N=9 for all other conditions). Levels normalised to *U6* and average levels of *WT* females.

E. Percentage and Median Fluorescence Intensity (MFI) of GATA3+ live TCRb+ CD4+ *WT* or *Malat1*$^{-/-}$ cells derived from female or male mice at day 2 of differentiation (N=8 per condition).

Figure 4. Impaired suppression of an interferon gene module during early differentiation of female *Malat1*$^{-/-}$ Th2 cells.

A. Volcano plot displaying LFC in gene expression between *WT* and *Malat1*$^{-/-}$ female cells (determined by Nanopore long-read RNAseq, N=4 per group) against -logFDR at day 2 of *in vitro* differentiation.

B. Gene set enrichment analysis (GSEA) hallmark gene set terms enriched within significantly up-regulated genes in *Malat1*$^{-/-}$ cells at day 2 of *in vitro* differentiation.

C. Heatmap displaying Z score of log$_2$CPM for top DEGs (FDR<0.01) between *WT* and *Malat1*$^{-/-}$ cells at day 2 of differentiation, in descending order of logFC.

D. LFC in gene expression between female *WT* naïve and differentiating cells at day 2 of *in vitro* differentiation against LFC between *WT* and *Malat1*$^{-/-}$ female cells at day 2 of *in vitro* Th2

differentiation. Data shown for all statistically significant DEGs between *WT* and *Malat1⁻ᐟ* cells

at day 2.

E. LFC in gene expression between female *WT* naïve and Th2 cells (as in Figure 1) against

LFC between *WT* and *Malat1⁻ᐟ* female cells at day 2 of *in vitro* Th2 differentiation. Data shown

for all statistically significant DEGs between *WT* and *Malat1⁻ᐟ* cells at day 2. Colours indicate

level of significance in differential expression when comparing day 2 vs naïve CD4⁺ T cells

(left) or Th2 vs naïve CD4⁺ T cells (right).

F. Levels of interferon induced genes (*Ifit1bl1*, *Ifit3b*, *Iigp1*, *Ifit1*) and transcription factors

(*Stat1, Irf1, Irf7*) in *WT* and *Malat1⁻ᐟ in vitro* differentiated female or male cells at day 2 of

differentiation. mRNA levels determined by qRTPCR (N=9) and normalised to *U6* and average

levels of *WT* cells from female mice.

G. Transcript usage as a percentage of total gene abundance for *Trim12a* and *Kdm2a*, two

genes that demonstrate statistically significant DTU. Levels shown between *WT* and *Malat1-*

/- females at day 2 of *in vitro* differentiation (left graphs) or between naïve (day 0) and day 2

in *WT* cells (right graphs). Levels determined by long-read RNAseq (N=4 per condition).

Tables indicate the product of each isoform. Stars indicate isoforms demonstrating statistically

significant differences in abundance as determined by Dexseq.

Figure 5. *Malat1* loss decreases expression of the IL2 receptor and impairs the

response to exogenous levels of IL2 in cells from female mice.

A. Representative dot plots of IL2Rα expression in TCRb⁺ CD4⁺ *WT* or *Malat1⁻ᐟ* cells from

female mice at day 4 of differentiation.

B. Percentage of IL2Rα⁺ live TCRb⁺ CD4⁺ *WT* or *Malat1⁻ᐟ* cells derived from female or male

mice at day 4 of differentiation (levels determined by surface staining, N=6).

C. Representative dot plots of IL2Rγ staining in *WT* or *Malat1⁻ᐟ* CD4⁺ T cells from female or

male at day 4 of differentiation.

D. Percentage of IL2Rγ⁺ live TCRb⁺ CD4⁺ *WT* or *Malat1⁻ᐟ* cells derived from female or male

mice at day 4 of differentiation (levels determined by surface staining, N=5).

E. Percentage of IL2Rα$^+$ live TCRb$^+$ CD4$^+$ *WT* cells derived from female and male mice at day

4 of differentiation that were treated with either non-targeting control (NTC) or *Malat1* targeting

GapmeRs. Levels determined by surface staining (N=5).

F. As in E, but for IL2Rγ$^+$ live TCRb$^+$ CD4$^+$ cells.

G. Normalised percentage of IL10$^+$, IL4$^+$ or IL10$^+$ IL4$^+$ live TCRb$^+$ CD4$^+$ *WT* or *Malat1$^{-/-}$ in vitro*

differentiated Th2 cells derived from female and male mice at day 6, with resuspension at day

4 in the indicated concentrations of IL2. Levels determined by intracellular staining, and

normalised to *WT* females treated with 10U/ml IL2 from each experiment (N=10).

Figure 6. *Malat1* loss causes a female-specific decrease in Th2 cell-derived IL10 levels

in vivo.

A. Schematic of *S. mansoni* egg injection experiment. 5,000 dead *S. mansoni* eggs were

injected interperitoneally (IP) into mice, then after a week the injection was repeated

intravenously (IV) via the tail vein. After another week, lungs and spleens were harvested and

processed for FACS.

B. Representative FACS plots of IL10 and IL4 expression in lung CD4$^+$ TCRb$^+$ cells from *S.*

mansoni egg injected *WT* or *Malat1$^{-/-}$* female or male mice.

C. Percentage and cell numbers of lung IL10$^+$ TCRb$^+$ CD4$^+$ cells derived from naïve or *S.*

mansoni egg injected (*Sm*) *WT* or *Malat1$^{-/-}$*, female and male mice. Levels determined by

intracellular staining (N=3 for *Malat1$^{-/-}$* male naïves, N=5 for all other naïves, N=9 for egg

injected mice).

D. As in **C**, but for splenic IL10$^+$ TCRb$^+$ CD4$^+$ cells (N=4 for *Malat1$^{-/-}$* male naïves, N=5 for all

other naïves, N=9 for egg injected mice).

E. Percentage and cell numbers of lung IL10$^+$ IL4$^+$ TCRb$^+$ CD4$^+$ cells derived from naïve or *S.*

mansoni egg injected (*Sm*) *WT* or *Malat1$^{-/-}$*, female and male mice. Levels determined by

intracellular staining (N=3 for *Malat1$^{-/-}$* male naïves, N=5 for all other naïves, N=9 for egg

injected mice).

F. As in **G**, but for splenic IL10$^+$ TCRb$^+$ CD4$^+$ cells (N=4 for *Malat1$^{-/-}$* male naïves, N=5 for all

other naïves, N=9 for egg injected mice).

Figure 1

A.

Naïve CD4+ T cell isolation	+ anti-CD3, anti-CD28, anti-IFNγ, rIL4				+ IL2	
Day 0 Naïve CD4+ T cells	**Day 1** Activation	**Day 2** Early differentiation	**Day 3** Differentiation	**Day 4** Differentiation Addition of IL2	**Day 5** Expansion and differentiation	**Day 6** Expansion and differentiation

B.

WT Female — 16.12 / 4.18 / 72.42 / 7.28

Malat1-/- Female — 9.52 / 1.27 / 82.89 / 6.32

WT Male — 7.20 / 1.28 / 86.51 / 5.01

Malat1-/- Male — 5.66 / 0.54 / 89.98 / 3.82

C. IL10+ Expression

D. IL4+ Expression

E. IL10+ IL4+ Expression

F. IL4+ IL10- Expression

- WT Female
- Malat1-/- Female
- WT Male
- Malat1-/- Male

G.

Female All Significant — Female top 500

Changes in Th2 cells due to lack of *Malat1*

Changes due to Th2 differentiation in WT cells

H.

Male All Significant — Male top 500

Figure 2

Figure 3

Figure 4

A. *Malat1⁻/⁻* vs *WT* - Day 2

B. GSEA Day 2 Up-regulated

C.

D. *Malat1⁻/⁻* vs *WT* day 2 DEGs in *WT* Day 2 v Naive

E. *Malat1⁻/⁻* vs *WT* day 2 DEGs in *WT* Th2 v Naive

F.

G.

Tx_ID	Transcript_size	Protein_size	Biotype
ENSMUST00000134564.2	3730	No protein	Retained intron
ENSMUST00000106839.9	1486	297aa	Protein coding
ENSMUST00000106837.8	1389	265aa	Protein coding
ENSMUST00000070943.7	1531	284aa	Protein coding
ENSMUST00000142641.2	562	No protein	Protein coding CDS not defined

Tx_ID	Transcript_size	Protein_size	Biotype
ENSMUST00000176483.3	1128	35aa	Protein coding
ENSMUST00000047898.14	8578	1161aa	Protein coding
ENSMUST00000075856.11	7242	1161aa	Protein coding
ENSMUST00000175682.2	2638	No protein	Retained intron
ENSMUST00000176653.2	1076	292aa	Protein coding

Figure 5

A. *WT* Female   *Malat1-/-* Female

IL2Rα / FSC

35.46   15.77

B. IL2Rα Expression Day 4

% TCRb(+) CD4(+)

WT Female, Malat1-/- Female, WT Male, Malat1-/- Male

***

C. *WT* Female   *Malat1-/-* Female   *WT* Male   *Malat1-/-* Male

IL-2Rγ / FSC

24.11   15.64   19.60   6.79

D. IL2Rγ Expression Day 4

% TCRb(+) CD4(+)

WT Female, Malat1-/- Female, WT Male, Malat1-/- Male

*   ***

E. IL2Rα Expression Day 4

% TCRb(+) CD4(+)

WT Female NTC, WT Female GapmeR, WT Male NTC, WT Male GapmeR

**

F. IL2Rγ Expression Day 4

% TCRb(+) CD4(+)

WT Female NTC, WT Female GapmeR, WT Male NTC, WT Male GapmeR

**

G.

IL10 Expression

Normalised %TCRb %CD4+

2 U/ml, 5 U/ml, 10 U/ml, 20 U/ml, 2 U/ml, 5 U/ml, 10 U/ml, 20 U/ml

***   ****   ****

IL4 Expression

Normalised %TCRb %CD4+

0.0643

2 U/ml, 5 U/ml, 10 U/ml, 20 U/ml, 2 U/ml, 5 U/ml, 10 U/ml, 20 U/ml

IL10 IL4 Expression

Normalised %TCRb %CD4+

2 U/ml, 5 U/ml, 10 U/ml, 20 U/ml, 2 U/ml, 5 U/ml, 10 U/ml, 20 U/ml

****   **

● *WT* Female
● *WT* Male

Figure 6

A.

IP injection to sensitize — Dead *S. mansoni* eggs

IV injection- eggs deposit in lungs

Infection site
Systemic

B.

Naive | Egg-injected

WT Female | Malat1⁻/⁻ Female | WT Female | Malat1⁻/⁻ Female

C. Lung IL10⁺ | Lung IL10⁺ Cell Numbers

D. Spleen IL10⁺ | Spleen IL10⁺ Cell Numbers

E. Lung IL10⁺ IL4⁺ | Lung IL10⁺ IL4⁺ Cell Numbers

F. Spleen IL10⁺ IL4⁺ | Spleen IL10⁺ IL4⁺ Cell Numbers