***Malat1* regulates female Th2 cell cytokine expression through controlling early**

differentiation and response to IL2

Mags Gwynne[*, †, ‡, ††], Katie A. West[*, †, ‡, ††], Stijn van Dongen[§, ¶], Ioannis Kourtzelis[*, †], Dawn

Coverley[†, ‡], Sarah A. Teichmann [§, #], Kylie R. James[**], James P. Hewitson[†, ‡], Dimitris [*, †].

[*] Hull York Medical School, University of York, United Kingdom. [†] York Biomedical Research

Institute, University of York, United Kingdom. [‡] Department of Biology, University of York,

United Kingdom. [§] Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, United

Kingdom. [¶] Current address: ExpressionEdits Ltd, Haverhill, United Kingdom. [#] Current

address: Cambridge Stem Cell Institute, University of Cambridge, Cambridge, United

Kingdom. [**] Translational Genomics Program, Garvan Institute of Medical Research, The

Kinghorn Cancer Centre, Darlinghurst, New South Wales, Australia.

[††] These authors contributed equally

Corresponding author: dimitris.lagos@york.ac.uk, tel.: +4401904328930

Running title: A sex-specific effect of *Malat1* on Th2 cell differentiation

Abstract

Identifying cell intrinsic regulators of immune sexual dimorphism is critical for treatment of

several immunopathologies. We show that *Malat1* is required for appropriate cytokine

expression in female but not male Th2 cells. *Malat1* deficiency impairs *in vitro* Th2

differentiation of naïve CD4[+] T cells from female mice, characterised by transcriptome-wide

effects and suppression of cytokine expression, particularly IL10. Upon IL10R blockade a

pronounced effect is also seen on IL4 and IL13. Mechanistically, naïve CD4[+] T cells from

Malat1[-/-] female mice demonstrate altered early activation kinetics and impaired early

differentiation gene expression, including up-regulation of an interferon stimulated gene (ISG)

module. This is followed by suppression of IL2Rα and IL2Rγ expression and IL2-mediated

differentiation. Mimicking the effect of *Malat1* loss by maintaining early ISG expression in *WT*

cells with IFNβ treatment partially phenocopies the effects of *Malat1* deficiency. A subset of

the effects of *Malat1* loss in female cells is also observed in male cells. However, this does

not affect endpoint Th2 differentiation. Male CD4$^+$ T cells demonstrate stronger early

activation, higher ISG expression during early differentiation, maintenance of IL2Rα

expression independently of *Malat1*, and lower sensitivity to exogenous IL2 during late

differentiation compared to female cells. *In vivo*, female, but not male, *Malat1*[-/-] mice

demonstrate altered Th2 cytokine expression characterised by a reduction in IL10$^+$ Th2 cells

in both lung and spleen following priming and challenge with *Schistosoma mansoni* eggs, a

model of lung type 2 inflammation. Overall, these findings reveal *Malat1* as a novel

determinant of immune sexual dimorphism.

Keywords: Th2 cells; Immune sexual dimorphism; lincRNAs; IL10; *Malat1*

Introduction

Biological sex is a known determinant of immune responses (1-4). Higher incidence or severity of autoimmune diseases (5-7) in females and infectious diseases in men (8) are commonly reported. The main drivers of immune sexual dimorphism are sex hormones (extrinsic) and X or Y chromosome-linked genes (intrinsic). For example, it is thought that females favour Th2 responses and males Th1 (2) and that this is due to hormonal regulation of immune genes such as IL4 (2, 9), and function of X chromosome-linked genes including the long intergenic non-coding RNA (lincRNA) *Xist* (10), which was recently shown to form ribonucleoprotein complexes that are directly involved in autoimmunity (11). Other mechanisms including those occurring at the post-transcriptional level, have been proposed, yet very little has been experimentally tested and validated. Overall, little is understood about cell intrinsic mediators of sexual dimorphism in Th cells that are not directly linked to sex hormone-mediated or sex chromosome-driven regulation.

Thousands of long non-coding transcripts, including lincRNAs, have been discovered in mammalian transcriptomes (12). In most cases, these are medium to lowly expressed transcripts displaying poor conservation across mammals. Our studies have focused on *Malat1*, a 7.5kb-long transcript discovered through an association with cancer progression and metastasis (13). *Malat1* is localised in nuclear speckles (14), which are nuclear foci enriched in factors involved in pre-mRNA splicing and transcription (15). It was initially thought that *Malat1* controls in mRNA splicing (14, 16). However, $Malat1^{-/-}$ mice develop normally and are viable and fertile without apparent defects in splicing (17-19). We previously discovered that *Malat1* is a key regulator of Th1 cell responses (20). We showed that *Malat1* is down-regulated upon initiation of Th cell differentiation, yet its complete loss results in stronger immune responses through suppression of IL10 expression in mouse models of type 1 immunity to parasitic infection, including visceral leishmaniasis and malaria. In humans, *Malat1* downregulation is a hallmark of proliferative CD4[+] and CD8[+] T cells (21). Others have linked *Malat1* to CD4[+] T cell function, particularly in Th17 cells (22, 23), and to CD8[+] T cell function (24).

In our previous report, which focused on Th1 cells, we showed that *Malat1* loss also affected

IL10 expression in *in vitro* differentiated Th2 cells (20). Although expressed by all effector Th

cells, IL10 expression is highest in Th2 cells (25), which we have also shown previously (20).

However, the role of *Malat1* in Th2 cell differentiation and function remain under-explored.

Here, we aimed to test how loss of *Malat1* affects Th2 differentiation *in vitro* and in an *in vivo*

model of type 2 inflammation, *Schistosoma mansoni* (*S. mansoni*) egg injection (26, 27). We

report that *Malat1* deficiency leads to impaired Th2 differentiation with widespread effects

across the transcriptome and a notable suppression of IL10 expression only observed in

female-derived Th2 cells. Although some effects caused by *Malat1* deficiency are shared

between male- and female-derived CD4$^+$ T cells, *Malat1* loss has more profound effects on

female Th2 cells. This is due to a female-specific impaired regulation of an early differentiation

interferon stimulated gene (ISG) expression programme, suppression of IL2 receptor (IL2R)

expression in *Malat1*$^{-/-}$ cells, and increased sensitivity of female cells to IL2-driven cytokine

production.

Materials and Methods

Animals and ethics

Animal care and experimental procedures were regulated under the Animals (Scientific

Procedures) Act 1986 (revised under European Directive 2010/63/EU) and were performed

under U.K. Home Office License (project license number PP0841992 for breeding and

PP9423191for *S. mansoni* egg injections) with approval from the University of York Animal

Welfare and Ethical Review Body. Animal experiments conformed to Animal Research:

Reporting of In Vivo Experiments guidelines (28).

***S. mansoni* egg injection**

C57BL/6 CD45.2 *wildtype* (*WT*) mice were obtained from Charles River Laboratories.

Malat1$^{-/-}$ mice (complete knockouts) were obtained from the Riken Institute (18). All mice were

bred in-house, maintained under specific pathogen-free conditions, and used at 6–12 weeks

of age. Schistosome eggs were recovered from the livers of C57BL/6 mice at week 7 post-

infection following exposure to 100 *S. mansoni* cercariae. Cercariae were obtained from

schistosome-infected *Biomphalaria glabrata* snails provided by the Barrett Centre for Helminth

Control (Aberystwyth University, UK). Livers were digested overnight at 37°C with shaking

with 0.2U/ml collagenase D (Roche) in the presence of 5000U/ml polymyxin B (Merck). Eggs

were purified by centrifugation through 10ml percoll (GE Healthcare) / 20ml 0.25M sucrose

(450 x*g*, 5min, RT), washed in PBS and stored at -20°C before usage. 5000 dead *S. mansoni*

eggs in 200 μl PBS were delivered via intraperitoneal (ip) injection into mice. Two weeks later,

the mice were intravenously (iv) challenged with another 5000 eggs in 200 μl PBS. After

another week the mice were sacrificed and the lungs and spleen were extracted and

processed. Lungs were digested with 0.4 U/ml Liberase TL (Roche) and 80 U/ml DNase I type

IV in HBSS (both Sigma-Aldrich) for 45 min at 37°C. Enzyme activity was inhibited with 10

mM EDTA (pH 7.5) and single-cell suspensions were created with 100-μm nylon filters

(Falcon) in complete RPMI 1640 (Thermo Fisher Scientific) supplemented with 10% heat-

inactivated FCS (HyClone), 100 U/ml penicillin, 100 μg/ml streptomycin, and 2 mM l-glutamine

(all Thermo Fisher Scientific), then cleaned via Percoll gradient and 3ml Ammonium-Chloride-

Potassium (ACK) red blood cell lysis buffer. Spleen single cell suspensions were generated

in the same manner and only passed through a 70 μm filter and treated with 3 ml ACK buffer.

In vitro Th2 differentiation and _Malat1_ knockdown

Spleens and axillary, brachial, mesenteric, and inguinal lymph nodes of *WT* or *Malat1−/−* mice

were extracted, homogenised, through a 70 μm filter in RPMI 1640. Resulting cell pellets were

then treated with 3ml ACK lysis buffer. Naïve CD4+ T cells were then purified via MACS column

isolation (Miltenyi Naive CD4+ T Cell Isolation Kit, mouse, catalogue number 130-104-453)

normally resulting in purity of ~95%. For Th2 polarisation, purified naïve CD4+ T cells (500,000

cells per well) were stimulated with 10 μg/ml plate-bound anti-CD3ε (clone 145-2C11) and 4

μg/ml soluble anti-CD28 (37.51) in RPMI 1640 in flat-bottom 96-well plates in the presence of

25 ng/ml mouse rIL4 and 5 µg/ml anti–IFN-γ (XMG1.2). Anti-CD3/anti-CD28–dependent

activation (4 days) was followed by rest in 10 U/ml human recombinant IL2 for 2 days.

Titrations of anti-CD3ε antibody or recombinant IL2 were performed as indicated.

Recombinant cytokines were from PeproTech. For *Malat1* knockdown experiments, control or

Malat1-targeting antisense oligonucleotide GapmeRs were from QIAGEN (Hilden, Germany;

LG00000002-DDA and LG00000008-DDA, respectively) and were added to naive CD4$^+$ T

cells on day 0, or differentiating Th2 cells on day 4, at a final concentration of 100 nM. For

IL10 receptor blockade experiments, CD4$^+$ T cells were treated either with anti-IL10R (Clone:

1B1.3A from Bio X Cell) or rat IgG at 10 µg/ml as a control, at both day 0 and day 4. For IFNβ

treatment experiments, cells were treated with 5000 U/ml mouse IFNβ (CYT-651, from

Prospec), at day 0.

Flow Cytometry

For flow cytometry analysis, single cell suspensions were generated. For live/dead

discrimination, cells were washed twice in PBS, then stained with Zombie Aqua (BioLegend)

in PBS before resuspension in FACS (Fluorescence-Activated Cell Sorting) buffer (PBS

containing 0.5% BSA and 0.05% azide). Fc receptors were blocked with 100 µg/ml rat IgG

(Sigma-Aldrich) for 10 min at 4°C before surface staining for 30 min at 4°C. The following anti-

mouse Abs from Biolegend were used: CD45.2 BV786 (clone 104); TCRβ PE-Cy7 (H57-597);

CD19 APC-Cy7 (6D5); MHC class II (MHCII) Alexa Fluor 700 (M5/114.15.2); Ly-6G APC-Cy7

(1A8); Ly6C BV605 (HK1.4); CD64 PE (X54-5/7.1); CD11b Pacific Blue (M1/70); CD44 FITC

(IM7); CD62L PE (MEL-14); CD8α PB (53-6.7); CD4 PerCP/Cy5.5 (RM4-5); IFN-γ FITC

(XMG1.2); IL10 PE (JES5-16E3); IL4 PE-Dazzle or APC (11B11); CD25 (IL-2Ra) PerCP-

Cy5.5 or APC (PC6.1); CD69 APC (H1.2F3); CD132 (IL-2Rg) PE (TUGm2); GATA3 PE-

Dazzle 594 (16E10A23); IFNAR1-PE (MAR1-5A3) and streptavidin PE-Cy7. These were used

in combination with abs SiglecF PerCP-eFluor 710 (1RNM44N), iNOS PE-eFluor 610

(CXNET) and goat anti-rabbit A647 from ThermoFisher Scientific, biotinylated Ym1 from R&D

systems, anti-murine RELMα from PeproTech, and ; IL-13 PB (eBio13A) from eBioscience.

To measure intracellular cytokines in T cells following *ex vivo* stimulation, cells were first

stimulated in complete RPMI 1640 for 4 h at 37°C with 500 ng/ml PMA, 1 µg/ml ionomycin,

and 10 µg/ml brefeldin A (all Sigma-Aldrich). For all intracellular cytokine staining, surface

stained cells were fixed and permeabilized (20 min at 4°C) using Fixation/Permeabilization

Solution before washes in Perm/Wash buffer (both BD Biosciences). Cells were then stained

with intracellular Abs as above except in Perm/Wash buffer. For transcription factor staining,

surface stained cells were fixed and permeabilized (1h or overnight at 4°C) using the FoxP3

Fixation/Permeabilization solution before washes in 1x Permeabilization Buffer (eBioscience).

Cells were stained with intracellular Abs as above except in Permeabilization Buffer.

Appropriate isotype or fluorescence minus one (FMO) controls were used to draw gates for

populations of interest. Events were acquired on an LSRFortessa (BD Biosciences) with

analysis by FlowJo (FlowJo) or a Cytoflex LX (Beckman Coulter) before analysis with or

CytExpert.

RNA extraction, cDNA synthesis, and quantitative RT-PCR

RNA was extracted from purified cell populations using QIAzol and miRNeasy RNA extraction

kits (QIAGEN) according to manufacturer's instructions. For mRNA transcripts, reverse

transcriptions were carried out with Superscript III (Thermo Fisher Scientific) and random

hexamer primers (Promega) and measured with Fast SYBR Green Master Mix (Thermo Fisher

Scientific). Quantitative PCR was performed using a StepOnePlus Real Time PCR System or

a QuantStudio 3 (both Thermo Fisher Scientific), and relative transcript levels were

determined using the ΔΔCt method. The primers used for the study are show in **Supplemental**

Table S1.

RNA sequencing (RNAseq) and analysis

We used short-read (Illumina) RNAseq to compare naïve CD4$^+$ T cells and *in vitro*

differentiated Th2 cells from female and male *WT* and *Malat1$^{-/-}$* mice. Pooled cDNA libraries

were sequenced on one lane of an Illumina Novaseq 6000 S4 flow-cell at 100PE. The raw

sequencing reads were aligned to the mouse genome version Ensembl GRCm38.92 with

STAR 2.5.4a. We sequenced four replicates per condition. Samples with assigned sequencing

reads lower than 500,000 were removed from downstream analysis. Of the remaining samples

(N=4 for all *Malat1*$^{-/-}$ samples and *WT* male Th2, N=3 for *WT* female and male naïve CD4$^+$ T

cells and *WT* female Th2), we obtained an average of 65 million reads per sample (range: 52–

91 million). Analysis was performed in RStudio (version 1.4.1106) with R (v 4.1). Data was

normalised ('calcNormFactors' and 'estimateDisp' functions of edgeR (version 3.14.0)) and a

quasi-likelihood negative binomial generalized log-linear model ('glmQLFit' and 'glmQLFTest'

functions with coefficient = 2) was used to determine differentially expressed genes. Adjusted

p values were calculated using the Benjamini and Hochberg method.

We used Oxford Nanopore Technologies (ONT) long-read RNAseq to compare naïve CD4$^+$ T

cells and cells after 2 days of *in vitro* differentiation from female *WT* and *Malat1*$^{-/-}$ mice (N=4

per group). Full length cDNA libraries were prepared using the ONT cDNA-PCR Sequencing

V14 - Barcoding kit (SQK-PCB114.24). Briefly, cDNA RT adapters were ligated to 3' ends

(polyA tails) of transcripts, prior to stand switching cDNA synthesis and a 13 cycle PCR

reaction with barcoded primers, labelling each cDNA sample with a unique DNA barcode.

Barcoded cDNAs were pooled at equimolar ratios before final adapter ligation and sequencing

on R10.4.1 flowcells in ONT PromethION sequencer (8 samples per flow cell). Live

superaccuracy basecalling and barcode demultiplexing were performed in MinKNOW

software (version 24.06.10) at the time of the run. We acquired 0.745-1.121 million reads per

sample, with 604-840 bases mean read length. Demultiplexed reads were analyzed through

the workflow Transcriptomes from Epi2Me^{TM} application, specifically designed by ONT. The

workflow generates differential gene expression data (with edgeR), as well as differential

transcript usage (DTU) data (with Dexseq), using a reference transcriptome. The Viking

cluster, provided by the University of York, was used for the analyses.

Statistics

Experiments were typically performed in 2 – 3 independent replicates in groups of N = 3 – 5

mice per condition and replicate. Statistical analyses were carried out as indicated with Prism

5 (GraphPad Software). Two-way comparisons used paired or unpaired t tests as indicated,

and multiple comparisons used one-way ANOVA, followed by Sidak's multiple comparisons

test for comparison of samples with biological relevance (e.g. *WT* female to *Malat1$^{-/-}$* female,

WT male to *Malat1$^{-/-}$* male, *WT* female to *WT* male). Confidence levels were set to 0.05 for

significance. In merged data where n>9, consensus between Anderson-Darling, Shapiro-Wilk

and D'Agostino & Pearson tests were used to determine normality. If all samples were

normally distributed one-way ANOVA was performed as described. If one or more samples

did not follow a normal distribution, a Kruskal-Wallis test followed by Dunn's multiple

comparison test were performed to identify significance, with confidence levels of 0.05. P

values are displayed as asterisks representing P value classification, where * ≤ 0.05, ** ≤ 0.01,

≤ 0.001 and * ≤ 0.0001, unless the results are borderline non-significant, in which case

the p value is shown.

Data availability

RNAseq data have been deposited in the National Center for Biotechnology (NCBI) Gene

Expression Omnibus (GEO) database. Accession numbers: GSE279185 for Illumina RNAseq

and GSE278413 for Oxford Nanopore Technologies RNAseq data.

Results

***Malat1* loss impairs *in vitro* Th2 differentiation of CD4$^+$ T cells from female mice**

We first assessed the effect *of Malat1* deletion on Th2 differentiation *in vitro* in response to

stimulation with anti-CD3 and anti-CD28 antibodies in the presence of recombinant IL4 and

anti-IFNγ antibody for 4 days, followed by IL2 treatment for 2 days. This protocol allowed us

to dissect potential effects on naïve CD4$^+$ T cell activation (0-24h), early differentiation (2-4

days), and terminal differentiation in response to exogenous IL2 (**Fig. 1A**). Following *in vitro*

differentiation of naive CD4+ T cells from female mice to Th2 cells, *Malat1$^{-/-}$* cells displayed

significantly lower percentages of IL10$^+$ and IL4$^+$/IL10$^+$ cells at day 6 of differentiation, with no

differences in IL4$^+$/IL10$^-$ cells, and no differences at day 4 (**Fig. 1B – F** and **Supplemental**

Fig. 1A). This decrease in cytokine expression was not observed from cells derived from male

mice. Under these conditions, differentiation of *WT* CD4$^+$ T cells from male mice was less

efficient compared to female *WT* cells, and *Malat1* deletion did not affect this further (**Fig. 1B**

– F). RNAseq analysis of *WT* and *Malat1*$^{-/-}$ naïve CD4$^+$ T cells and *in vitro* differentiated Th2

cells revealed only a small number of genes were statistically significantly differentially

expressed between *WT* and *Malat1*$^{-/-}$ cells (**Supplemental Table S2**). As shown before (20),

IL10 levels were suppressed in *Malat1*$^{-/-}$ Th2 cells compared to *WT* controls (Log$_2$ fold change

LFC = - 0.882, p = 0.034, for female-derived cells and LFC = 0.03, p = 0.932 for male-derived

cells), although this did not reach significance after multiple testing correction. This was further

confirmed by qRTPCR for the sequenced samples (**Supplemental Fig. S1B**), suggesting that

this bulk RNAseq analysis provided a conservative representation of the effect of *Malat1*

deficiency on Th2 gene expression. Focusing on genes that were differentially expressed

between Th2 and naïve CD4$^+$ T cells from *WT* mice (Th2-differentiation associated genes) we

observed a transcriptome-wide blunting of Th2 gene expression upon *Malat1* loss in female

but not male cells (**Fig. 1G – H** and **Supplemental Tables S3 – S5**). Analysis of all

(FDR<0.001) or the top (by FDR) 500 differentially expressed genes (DEGs) between *WT* Th2

and naïve CD4$^+$ T cells demonstrated that most genes that were up-regulated during *WT* Th2

differentiation were overall expressed at lower levels when comparing *Malat1*$^{-/-}$ Th2 cells to

WT Th2 cells, and, conversely, genes that were suppressed upon *WT* Th2 differentiation were

overall more highly expressed in *Malat1*$^{-/-}$ Th2 cells (**Fig. 1G** and **Supplemental Tables S3 –**

S5). This transcriptome-wide trend was not observed in male cells (**Fig. 1H**). Taking a stricter

threshold for DEGs between Th2 and naïve CD4$^+$ T cells from female mice (FDR<0.001 and

absolute LFC>2), we found 524 Th2-differentiation associated DEGs in both *WT* and *Malat1*$^-$

$^{/-}$, 469 DEGs only in *WT*, and 139 DEGs only in *Malat1*$^{-/-}$ Th2 cells. We noted that common

and *WT* only Th2 DEGs were mainly up-regulated, whereas *Malat1*$^{-/-}$ only Th2 DEGs were

mainly down-regulated (**Supplemental Fig. S1C – D**, **Supplemental Tables S3 – S5**).

Furthermore, when looking at common DEGs, we found that the vast majority of up-regulated

genes showed a smaller LFC in *Malat1*⁻ᐟ⁻ Th2 cells compared to *WT* (**Supplemental Fig. S1E**).

To distinguish the effect of *Malat1* from that of IL10, we blocked the IL10 receptor (IL10R)

between days 4 and 6 of differentiation. As above, in control conditions, the effect of *Malat1*

on IL4 and IL13 effector Th2 cytokines was modest. However, upon blockade of IL10R, *Malat1*

loss resulted in dramatic and statistically significant decrease in IL4 and IL13 expression in

female-derived Th2 cells, with no effect on male-derived cells (**Fig. 1I**). Overall, these results

indicated that *Malat1* loss impairs Th2 differentiation of female- but not male-derived naïve

CD4⁺ T cells, characterised by suppression of IL10 expression and an impairment of the Th2

gene expression programme.

The effect of *Malat1* depletion on IL10 expression is independent of activation strength

and occurs during the early stages of differentiation

We next tested whether *Malat1* deficiency affected Th2 polarization under weaker activation

conditions. We found that *Malat1*⁻ᐟ⁻ cells displayed lower IL10 expression under suboptimal

differentiation conditions, whereas IL4 levels were not statistically significantly different (**Fig.**

2A – B, **Supplemental Fig. S2A – B**). Of note, at lower activation levels there were no

differences in IL10 expression between *in vitro* Th2 differentiation of male and female cells.

This indicated that *Malat1* deficiency can affect Th2 cytokine expression independently of

activation strength. Under suboptimal differentiation conditions, IL10R blockade did not

enhance the effect of *Malat1* deficiency on IL4 or IL13 expression, potentially due to the low

cytokine expression levels under these conditions (**Supplemental Fig. S2C**).

We next tested whether *Malat1* knockdown rather than genetic deletion had a sex-specific

effect on Th2 differentiation. As described previously (20), we used GapmeR oligonucleotides

to suppress *Malat1* expression (**Supplemental Fig. S2D**). Although, Th2 cells express lower

Malat1 levels than naïve CD4⁺ T cells ((20) and **Supplemental Fig. S2E**), we found that

Malat1 levels increase between day 4 and 6 of differentiation, upon cessation of CD3

activation and addition of exogenous IL2 (**Fig. 2C**). As for cytokine expression (**Fig. 2A-B**),

the increase in *Malat1* levels in response to IL2 (day 4 to day 6) was more profound in female

cells. We note that there were no differences in *Malat1* expression between *WT* male and

female naïve **(Supplemental Fig. S2E)** or differentiating (day 4; **Fig. 2C**) CD4$^+$ T cells prior to

IL2 stimulation.

Based the observation that *Malat1* deficiency impaired cytokine induction between days 4 and

6 (**Fig. 1C – F**), we knocked down *Malat1* either on day 0, or by adding GapmeRs on day 4,

at the same time with exogenous IL2 stimulation. *Malat1* knockdown reduced IL10 expression

in female Th2 cells only if GapmeRs were added to naïve CD4$^+$ T cells, but not when added

concurrently with exogenous IL2 stimulation. No effects on cytokine expression were observed

upon *Malat1* knockdown in male cells (**Fig. 2D**). Partial knockdown of *Malat1* in cells from

female mice did not affect IL4 expression (**Supplemental Fig. S2F**). As in the case of *Malat1*$^-$

$^{/-}$ cells, IL10R blockade enhanced the effect of *Malat1* GapmeR-mediated knockdown on IL4

(**Fig. 2E**). This was not observed for IL13, potentially due to low IL13 expression under these

conditions (**Fig. 2E**). We noticed that treatment with non-targeting control GapmeRs

consistently resulted in lower IL13 cytokine expression in Th2 cells (compare **Fig. 2E** with **Fig.**

1I). Overall, this demonstrated that the effect of *Malat1* on Th2 cytokine expression required

its activity during the early differentiation stages.

***Malat1* loss alters the kinetics of naïve CD4$^+$ T cell activation and impairs early Th2**

differentiation

Based on the above, we tested the effect of *Malat1* deletion on the early stages of Th2

differentiation. First, we studied the first 24h of naïve CD4$^+$ T cell activation by measuring

levels of T cell early activation marker CD69 (**Supplemental Fig. S3A**). As expected, *Malat1*

levels were suppressed within 24h in both male and female *WT* mice (**Supplemental Fig.**

S3B). In female cells, we found that compared to *WT* cells CD69 protein expression was

higher in *Malat1*$^{-/-}$ cells at 3h post activation, the difference reaching statistical significance at

7h post activation. However, CD69 expression was significantly lower in *Malat1*$^{-/-}$ cells at 24h

(**Fig. 3A**). At the transcript level, *Cd69* levels peaked at 3h post activation, with significantly

higher levels in *Malat1*[-/-] cells (**Supplemental Fig. S3C**). In male cells, we did not observe any statistically significant differences between *WT* and *Malat1*[-/-] cells, although we noted that male *WT* cells showed statistically significant higher CD69 levels compared to female counterparts at 3h post activation (**Fig. 3A**). We also measured levels of IL2Rα (CD25), which was induced upon activation but found no differences between genotypes or biological sexes (**Supplemental Fig. S3D**).

To explore the consequences of the altered activation kinetics in female *Malat1*[-/-] cells, we performed a daily timecourse (days 1 – 4), measuring transcript levels of key transcription factors (*Gata3*, *Maf*, and *Tbet*) and differentiation-associated genes (*Il2, Il2ra, Il2rg*). We focused on *Il2ra* and *Il2rg* as the former is induced during differentiation and the latter is an X-linked gene, given the sex-specificity of the observed effects. We noticed suppression for multiple of the measured transcripts in female *Malat1*[-/-] differentiating cells, including *Il2*, *Il2ra*, *Il2rg*, and *Gata3* on days 2 and 3 of differentiation (**Fig. 3B – C,** and **Supplemental Fig. S3E**). These initial findings were confirmed in independent experiments for *Il2ra*, *Il2rg*, and *Gata3* on day 2 (**Fig. 3D**). We chose day 2 for the validation experiments as this was the earliest timepoint the suppression of the Th2 programme was observed. In male *Malat1*[-/-] cells, we observed higher variability and similar but non-statistically significant trends (**Fig. 3D**). GATA3 protein levels were suppressed in *Malat1*[-/-] cells at day 2 of the differentiation in both male and female *Malat1*[-/-] cells (**Fig. 3E** and **Supplemental Fig, S3F**). At this point we did not find any significant differences in the levels of the IL2R subunits (**Supplemental Fig. S3G**). These results suggested that *Malat1* was necessary for appropriate early Th2 differentiation.

Impairment of early differentiation in *Malat1*[-/-] cells is characterised by up-regulation of an ISG signature that suppresses differentiation

To further explore the role of *Malat1* in early Th2 differentiation we performed long-read RNAseq of *WT* and *Malat1*[-/-] naïve CD4[+] T cells and differentiating cells at day 2 of differentiation. As *Malat1* has been suggested to be involved in regulation of mRNA splicing (14), we opted for long-read RNAseq to facilitate concurrent assessment of differential gene

expression and differential transcript usage (DTU). As in the case of our short read RNAseq analysis, there were only few differences between *WT* and *Malat1$^{-/-}$* naïve CD4$^+$ T cells (16 DEGs at FDR<0.1, **Supplemental Fig. S4A** and **Supplemental Table S6**). However, comparison of *WT* and *Malat1$^{-/-}$* cells on day 2 of differentiation revealed 239 DEGs, 199 of which were up-regulated in *Malat1$^{-/-}$* cells (**Fig. 4A** and **Supplemental Table S7**). Gene set enrichment analysis (GSEA; (29)), revealed a profound over-representation of interferon alpha and interferon gamma response gene signatures amongst the up-regulated genes in *Malat1$^{-/-}$* cells on day 2 (**Fig. 4B**), whereas enriched terms amongst down-regulated genes, including hypoxia, only reached modest significance (**Supplemental Fig. S4B**). We note that in agreement with our qRTPCR analysis (**Fig. 3D**), trends towards down-regulation were observed for *Il2ra* (LFC = -0.589) and *Gata3* (LFC = -0.401) but these did not reach statistical significance. The most prominent feature of *Malat1$^{-/-}$* cells, the up-regulated interferon gene cluster (**Supplemental Fig. S4C** and **Supplemental Table S8**; clusters identified by k-means clustering) included ISGs (e.g. *Ifit1bl1*, *Ifit3*, *Ifit1*, *Oas2*, and others) and transcription factors including *Irf7*, *Irf1*, and *Stat1* (**Fig. 4C**, **Supplemental Fig. S4C** and **Supplemental Tables S7 and S8**). Most of the day 2 *Malat1*-associated DEGs were down-regulated during differentiation of *WT* cells both when comparing day 2 with naïve cells (**Fig. 4D**) and when comparing *WT* Th2 and naïve CD4$^+$ T cells using our short-read sequencing data (**Fig. 4E**). This was also the case specifically for the ISG module (**Supplemental Fig. S4D**). As in the case of Th2 cells (**Fig. 1G**) *Malat1* deficiency had a transcriptome-wide effect by blunting changes in gene expression occurring in *WT* cells within the first two days of differentiation (**Supplemental Fig. S4E**).

Validation of the long-read sequencing data by qRTPCR for selected genes confirmed up-regulation of *Ifit1bl1*, *Ifit3b*, *Iigp1*, *Ifit1*, and *Irf7* (**Fig. 4F**). Up-regulation trends for Irf1 and Stat1 did not reach statistical significance. Notably, no effects due to *Malat1* deficiency were observed in male CD4$^+$ T cells, and levels of *Ifit1bl1*, *Ifit3b*, and *Irf7* were significantly higher in male *WT* cells compared to female cells on day 2 (**Fig. 4F**). Comparing day 2 to naïve CD4$^+$ T cells, we found significant down-regulation for *Ifit1bl1*, *Ifit1*, and *Irf7* for all conditions.

Statistically significant *Stat1* down-regulation on day 2 was only observed for female *WT* cells

(**Supplemental Fig. S4F**). *Malat1* deficiency had weak effects on DTU affecting only 25 genes

(FDR<0.1, **Supplemental Table S9**) including some cytokine genes, but demonstrating

modest effects. In some instances, as for *Kdm2a* and *Trim12a* isoform usage changes due to

Malat1 loss showed opposing patterns to those induced by early differentiation of *WT* cells

(**Supplemental Fig. S5**).

To explore the link between sustained early ISG expression and endpoint Th2 differentiation,

we treated *WT* or *Malat1$^{-/-}$*, female or male, naïve CD4$^+$ T cells with IFNβ. This resulted in

enhanced mRNA levels in *WT* female-derived cells of several genes identified as up-regulated

in *Malat1$^{-/-}$* CD4$^+$ T cells at Day 2 of differentiation including *Ifit1bl1*, *Ifit3b*, and *Irf7*

(**Supplemental Fig. S6A**). We note that although overall ISG expression trends were similar,

there were differences between IFNβ-treated and *Malat1$^{-/-}$* cells at day 2, for example

upregulation of *Stat1* reached statistical significance in IFNβ-treated cells (**Supplemental Fig.**

S6A) but not when comparing *Malat1$^{-/-}$* to *WT* cells (**Fig. 4F**). Critically, IFNβ treatment resulted

in a statistically significant suppression of IL10 expression and a modest decrease in IL4

expression in Th2 cells (**Fig. 4G**) phenocopying the effect of *Malat1* loss. No effects were

observed in male-derived cells due to IFNβ treatment (**Fig. 4G** and **Supplemental Fig. S6B**).

Interestingly, IL10R blockade enhanced the effect of IFNβ treatment on IL13 expression

(higher statistical significance) in female cells but had no effect on IL4 (**Supplemental Fig.**

S6C). The latter can be due to the fact that the IFNβ treatment-induced ISG signature only

partially overlaps with that seen in *Malat1$^{-/-}$* cells (**Supplemental Fig. S6A**), likely resulting in

common but also distinct effects on endpoint cytokine expression. Of note the higher ISG

expression in female *Malat1$^{-/-}$* cells was not due to differences in levels of type 1 interferons

(**Supplemental Fig. S7A;** *Ifna*, *Ifnb*, and *Ifna4* were lowly detectable at mRNA level only on

day 2). We did not observe any differences in *Ifnar1* mRNA levels (**Supplemental Fig. S7B**).

IFNAR1 protein expression was higher in Day 2 differentiating CD4$^+$ T cells compared to naïve

CD4$^+$ T cells but we did not observe any differences between female and male and *WT* and

Malat1$^{-/-}$ cells (**Supplemental Fig. S7C**) that could account for the observed differences in

ISG expression. We only found a modest increase in IFNAR1 expression in male *Malat1*$^{-/-}$

naïve cells compared to *WT* controls but this was not followed by any changes in ISG

expression (**Fig. 4F and Supplemental Fig. S4**).

Overall, these analyses further supported that *Malat1* is necessary for early Th2 differentiation

with the predominant effect being impaired suppression of an ISG signature that is associated

with impaired Th2 differentiation.

Female differentiating Th2 cells are sensitive to suppression of IL2R caused by *Malat1*

deficiency

Having shown that *Malat1* deficiency impairs early Th2 differentiation of CD4$^+$ T cells, we

reasoned that this could affect their readiness to respond to exogenous IL2. Having found

decreases in the mRNA levels of the IL2R subunits (**Fig. 3D**), we measured protein levels of

IL2Rα and IL2Rγ on day 4, prior to addition of exogenous IL2. We found that IL2Rα was

statistically significantly down-regulated only in female *Malat1*$^{-/-}$ T cells on day 4 (**Fig. 5A**,

Supplemental Fig. S8A). IL2Rγ was down-regulated in both female and male *Malat1*$^{-/-}$ T cells

(**Fig. 5B**, **Supplemental Fig. S8B**). No differences were observed at the mRNA level at this

timepoint (**Supplemental Fig. S8C**). Similarly, IL2Rα was significantly down-regulated on day

4 in female cells treated with *Malat1*-targeting GapmeRs (**Fig. 5C**) and this was also the case

for IL2Rγ (**Fig. 5D**). In male cells, there was a non-significant trend towards down-regulation

for IL2Rγ (**Fig. 5D**). To explore the link between the early *Malat1*-associated ISG signature

and IL2R subunit expression on day 4, we treated *WT* cells with IFNβ. This resulted in a

statically significant suppression of IL2Rγ expression and a non-significant decrease in IL2Rα,

phenocopying in part the effect of *Malat1* loss or inhibition. IFNβ did not affect IL2R expression

in male-derived cells (**Fig. E – F**).

The observed down-regulation in the IL2R subunits at this time-point in female *Malat1*$^{-/-}$ cells

can explain the impaired induction of cytokine expression between day 4 and day 6 (**Fig. 1C**

– F). This also indicates that starting from the same baseline at day 4 with regards to IL4 and

IL10 expression, *WT* male cells were less responsive to exogenous IL2 compared to female

cells. We further tested the dependence of female and male *WT* cells on IL2 stimulation by

titrating recombinant IL2. We found that although male cells expand as well or, at the lowest

IL2 concentration, better than female cells (**Supplemental Fig. S8D**), they demonstrated

lower IL10 expression than female cells at all concentrations (**Fig. 5G**). The observed

differential regulation of IL2R receptor and dependence on IL2 for cytokine expression could,

at least in part, explain the more profound effect on female Th2 differentiation due to *Malat1*

deficiency.

***Malat1* loss impairs Th2 cell cytokine expression only in female mice in an experimental**

model of type 2 inflammation

Having demonstrated female-specific effects on Th2 cytokine expression and early

differentiation due to *Malat1* deficiency in a cell-intrinsic context *in vitro*, we tested the effect

of *Malat1* loss in an experimental model of type 2 inflammation *in vivo*. Intravenous *S. mansoni*

egg challenge in mice previously primed with schistosome eggs (**Fig. 6A**) leads to the

formation of type 2 inflammatory granulomas in the lungs and mimics helminth-induced lung

inflammation (26, 27) leading to cellular infiltration in the lungs, including Th2 cells, and

eosinophilia. Indeed, *S. mansoni* egg injection resulted in increased in lung cell numbers,

although we note that this only reached statistical significance in *WT* female mice and *Malat1*$^{-}$

$^{/-}$ male mice (**Supplemental Fig. S9** and **S10A**). Similar splenic cell numbers were observed

under all conditions (**Supplemental Fig. S10B**) and similar lung eosinophil numbers and

activation, measured by RELMα expression (**Supplemental Fig. S10C – D**). Notably, both

lung and splenic CD4$^+$ T cells in egg-injected female *Malat1*$^{-/-}$ mice expressed lower levels of

IL10 than *WT* mice (**Fig. 6B – D**). IL4$^+$/IL10$^+$ cells were also decreased in the lungs

(percentage) and spleens (percentage and numbers) of egg-injected *Malat1*$^{-/-}$ mice (**Fig. 6E –**

F). This was also the case for triple positive IL4$^+$/IL13$^+$/IL10$^+$ Th2 cells in the lungs of egg-

injected mice (**Supplemental Fig. S10E**). None of these differences were observed when

comparing male *Malat1*^{-/-} mice to *WT* controls. On the contrary the percentage of splenic

IL4$^+$/IL10$^+$ cells increased in egg-injected *Malat1*^{-/-} male mice. Under these conditions, we did

not observe any differences in total IL4 expression between egg injected *Malat1*^{-/-} and *WT*

female mice, although we noted an increase in IL4-expressing splenic CD4$^+$ T cells in male

Malat1^{-/-} mice compared to *WT* (**Supplemental Fig. S10F – G**). IFNγ$^+$ CD4$^+$ T cells are also

induced as a result of *S. mansoni* egg injection, and IL10 levels were also reduced in female-

specific manner within this population in the lungs of inflamed mice (**Supplemental Fig. S10H**

– I). With regards to activation status of CD4$^+$ T cells, we observed a borderline non-significant

decrease in activated (CD44^{high}/CD64L^{low}) lung CD4$^+$ T cells in female *Malat1*^{-/-} mice compared

to *WT* and no differences in the spleen (**Supplemental Fig. S10J – K**). No differences were

observed in percentage and activation status of myeloid cell populations in egg-injected mice

(**Supplemental Fig. S10L – M**). Overall, these results indicated that, as observed in our *in*

vitro Th2 differentiation experiments, *Malat1* deficiency resulted in a female-specific

impairment of Th2 cytokine expression, predominantly demonstrated by reduced IL10

expression in Th2 cells.

Discussion

Understanding cell intrinsic mediators of sexual dimorphism in lymphocytes is critical to

addressing differences in incidence and severity of immunopathologies between females and

males (1-4). We demonstrate that *Malat1*, one of the most highly abundant transcripts in

mammalian cells, exerts a sex-specific function in Th2 differentiation, affecting early

differentiation and end-point cytokine expression in female cells, predominantly IL10 (**Fig. 7**).

Malat1 deficiency has also some effects in male-derived cells, for example a modest

suppression of early GATA3 expression and a suppression in IL2Rγ expression, yet these do

not translate in changes in endpoint cytokine expression or gene expression programme.

Furthermore, the female-specific effect due to *Malat1* deficiency is observed in the *S. mansoni*

egg injection model, as biological sex is a determinant of immune responses in the context of

human schistosomiasis (30). We also note that, although *Malat1* loss is associated with

suppression of IL10 both *in vitro* and *in vivo*, other effects of *Malat1* deficiency, for example a

modest suppression of IL4, are only observed *in vitro*. This can be due to compensatory

mechanisms and multiple effects contributing to cytokine expression *in vivo*. For example,

through IL10R blockade we found *in vitro* that IL4 and IL13 levels in *Malat1$^{-/-}$* mice are likely

to reflect both the loss of *Malat1* and reduced IL10 levels from Th2 cells. This is in agreement

with reports demonstrating that IL10 suppresses activation of Th2 cells *in vivo* (31). We

speculate that *Malat1* might play a critical role in Th2 differentiation in type 2

immunopathologies, especially those characterised by impaired IL10 signalling (32). In

addition, IL10 has been shown to be expressed later than other effector cytokines (33) and

from heterogeneous populations of CD4$^+$ T cells (34). As such, discrepancies between *in vitro*

and *in vivo* cytokine expression can be due to the time-point assessed and kinetics of Th2 cell

emergence in the *S. mansoni* egg injection model. Nevertheless, the observed impairment of

cytokine expression during *in vitro* Th2 differentiation and in the *S. mansoni* egg injection

model strongly support a cell-intrinsic function of *Malat1* in controlling Th2 differentiation in a

sex-specific manner.

Despite the limitation of not capturing the complexities of *in vivo* Th2 cell differentiation, *in vitro*

differentiation of naïve CD4$^+$ T cells to Th2 cells allowed us to demonstrate that *Malat1* loss

results in a sex-specific impairment of differentiation in a cell-intrinsic manner. Mechanistically,

down-regulation of IL2R upon *Malat1* loss, prior to exposure to exogenous IL2, results in a

transcriptome-wide suppression of the Th2 differentiation programme. Although down-

regulation of IL2Rγ is observed in both male and female cells, IL2Rα, which is necessary for

formation of the high affinity IL2R (35), is only suppressed in female cells. *In vitro*, male Th2

differentiating cells are less responsive to IL2 with regards to cytokine expression, providing

an additional potential explanation of the sex-specificity of the observed effect. Evidence for

sexual dimorphism in response to IL2 in other immune cell types has been already reported.

Female group 2 Innate Lymphoid Cells demonstrate increased proliferation in response to IL2

when compared to male ILC2 cells (36), and female NK cells produce higher levels of IFNγ

than males in response to increased levels of IL2 (37).

We postulated that the observed downregulation of IL2Rα and IL2Rγ on day 4 is preceded by

impaired early differentiation (days 2-3) of *Malat1*$^{-/-}$ cells, an effect which is more pronounced

in female cells. To probe how early *Malat1* deficiency impairs Th2 cytokine expression we

used long-read RNA sequencing as a superior method in identifying effects on gene

expression and, particularly, splicing. *Malat1* deficiency had minimal effects on DTU at day 2

post-differentiation induction. We did not analyse our endpoint (day 6) short read RNA

sequencing experiments for DTU because (1) our data showed that it was the effect of *Malat1*

on early Th2 differentiation that was driving the phenotype of *Malat1*$^{-/-}$ Th2 cells (**Figs. 2 and**

3) and (2) any effects would most likely be a result of impaired response to IL2 (**Fig. 5**) rather

than a direct effect of *Malat1*. Indeed, rather than changes in splicing, we found that early

differentiation of female *Malat1*$^{-/-}$ cells is characterised by impaired suppression of an ISG

cluster. It is notable that Th cells demonstrating ISG (including expression of *Ifit3*, *Irf7*, *Stat1*,

but not *Ifng*) have been described in type 2 inflammation both in humans (38) and in mouse

models (39). In all cases, these Th cells represent a distinct population to Th2 cells indicating

that this signature needs to be suppressed in Th2 cells, which in agreement with our findings

in *WT* cells. Interestingly a recent publication identified an interferon-experienced population

of naïve CD4$^+$ T cells that show impaired response to TCR stimulation, as seen here for

Malat1$^{-/-}$ cells (40). Another report demonstrated that these cells exist constitutively including

in germ free mice (41). The 50-gene signature that defines interferon-experienced naïve CD4$^+$

T cells (40) displays substantial overlap with the *Malat1*-regulated gene cluster (at day 2)

identified here (26/50 genes, including *Irf7* and *Stat1*). Taken together with the observed effect

of IFNβ on Th2 differentiation here (partially phenocopying of *Malat1* deficiency), this suggests

that suppression of this ISG signature in naïve CD4$^+$ T cells is necessary for optimal Th2

differentiation and regulated by *Malat1* in female-derived cells. Of note, sexual dimorphism in

IFN responses and ISG expression have been reported (42, 43). It will be interesting to further

investigate how *Malat1* promotes specifically down-regulation of ISGs in female cells during

early CD4[+] T cell differentiation. It has been shown that *Malat1* can promote gene set-specific

Polycomb Repressor Complex-mediated epigenetic silencing in CD8[+] T cells (24). It could be

explored whether this is also the case for ISGs in naive CD4[+] T cells.

Our results suggest that the impairment of early Th2 differentiation in female *Malat1*[-/-] cells is

associated with a stronger but shorter-lived initial activation (0-24h) as measured by CD69

expression. Indeed, strong TCR stimulation in the presence of CD28 co-stimulation has been

previously linked to impaired Th2 differentiation (44). During the first 24h of activation, male

WT cells demonstrate CD69 expression kinetics similar to female *Malat1*[-/-] cells, while deletion

of *Malat1* does not cause any further increase in CD69 expression in male cells. Overall, we

propose that *Malat1* downregulation upon activation of naïve CD4[+] T cells facilitates activation,

potentially through release of associated proteins that contribute co-transcriptional

processing. However, in female cells, complete loss of *Malat1* alters activation kinetics, which

in combination with failure to suppress expression of ISGs, leads to impaired Th2

differentiation and subsequent suppression of IL2R and IL2-mediated cytokine expression.

Male cells are not sensitive to *Malat1* loss, at least partly, due to their higher early activation

threshold, higher ISG expression during early differentiation, and, lower sensitivity to

exogenous IL2 during the later stages of differentiation.

We cannot exclude additional female-specific mechanisms being affected by *Malat1*, for

example with regards to regulation of X Chromosome-linked genes and the role of X

chromosome inactivation during Th2 differentiation (10). Furthermore, it will be essential for

our results to be validated in other models of type 2 immunity and in human CD4[+] T cells.

Despite these limitations, our study reveals that female Th2 cells show a specific dependence

on *Malat1*, a highly expressed lincRNA not located on a sex chromosome and displaying

similar expression in male and female CD4[+] T cells. This can have far reaching implications

for our understanding of immune sexual dimorphism and provide novel routes for sex-specific

manipulation of adaptive immunity.

Author contributions

MG and KAW contributed to experimental design, data generation, data analysis, figure

design, and manuscript writing. SvD and KRJ generated and analysed Illumina RNAseq

experiments. IK, DC, and SAT contributed to study supervision and data interpretation. JPH

contributed to study design, data generation, and data interpretation. DL conceived and

coordinated the study, contributed to study design, data interpretation, and study supervision,

and wrote the manuscript. All authors contributed to manuscript editing and approved the final

manuscript.   MG and KAW share the first position in the author list and are listed

alphabetically.

Acknowledgements

This work was funded by the Hull York Medical School (PhD studentship to MG and additional

funding to DL), the Biotechnology and Biological Sciences Research Council (BBSRC) White

Rose doctoral training partnership (to KAW, reference BB/J014443/1), and the Medical

Research Council (to JPH, reference MR/W018578/1). We thank Sally James and Fabiano

Pais at the Genomics Lab and Data Science Hub, respectively, in the University of York

Bioscience Technology Facility for support with long-read sequencing data. We thank Joanna

Greenman for technical support and Allison Green for comments on the manuscript.

Disclosures

The authors have no financial conflicts of interest.

Data availability

Raw data are available upon request. RNAseq data have been deposited in the National

Center for Biotechnology (NCBI) Gene Expression Omnibus (GEO) database. Accession

numbers: GSE279185 for Illumina RNAseq and GSE278413 for Oxford Nanopore Technologies RNAseq data.

References

1. Gal-Oz, S.T., B. Maier, H. Yoshida, K. Seddu, N. Elbaz, C. Czysz, O. Zuk, B.E. Stranger, H. Ner-Gaon, and T. Shay, *ImmGen report: sexual dimorphism in the immune system transcriptome.* Nat Commun, 2019. **10**(1): p. 4295.

2. Klein, S.L. and K.L. Flanagan, *Sex differences in immune responses.* Nat Rev Immunol, 2016. **16**(10): p. 626-38.

3. Marquez, E.J., C.H. Chung, R. Marches, R.J. Rossi, D. Nehar-Belaid, A. Eroglu, D.J. Mellert, G.A. Kuchel, J. Banchereau, and D. Ucar, *Sexual-dimorphism in human immune system aging.* Nat Commun, 2020. **11**(1): p. 751.

4. Takahashi, T. and A. Iwasaki, *Sex differences in immune responses.* Science, 2021. **371**(6527): p. 347-348.

5. Dodd, K.C. and M. Menon, *Sex bias in lymphocytes: Implications for autoimmune diseases.* Front Immunol, 2022. **13**: p. 945762.

6. Rubtsova, K., P. Marrack, and A.V. Rubtsov, *Sexual dimorphism in autoimmunity.* J Clin Invest, 2015. **125**(6): p. 2187-93.

7. Whitacre, C.C., S.C. Reingold, and P.A. O'Looney, *A gender gap in autoimmunity.* Science, 1999. **283**(5406): p. 1277-8.

8. Shepherd, R., A.S. Cheung, K. Pang, R. Saffery, and B. Novakovic, *Sexual Dimorphism in Innate Immunity: The Role of Sex Hormones and Epigenetics.* Front Immunol, 2020. **11**: p. 604000.

9. Lambert, K.C., E.M. Curran, B.M. Judy, G.N. Milligan, D.B. Lubahn, and D.M. Estes, *Estrogen receptor alpha (ERalpha) deficiency in macrophages results in increased stimulation of CD4+ T cells while 17beta-estradiol acts through ERalpha to increase IL-4 and GATA-3 expression in CD4+ T cells independent of antigen presentation.* J Immunol, 2005. **175**(9): p. 5716-23.

10.   Syrett, C.M., B. Paneru, D. Sandoval-Heglund, J. Wang, S. Banerjee, V. Sindhava,

E.M. Behrens, M. Atchison, and M.C. Anguera, *Altered X-chromosome inactivation in T cells*

may promote sex-biased autoimmune diseases. JCI Insight, 2019. **4**(7).

11.   Dou, D.R., Y. Zhao, J.A. Belk, Y. Zhao, K.M. Casey, D.C. Chen, R. Li, B. Yu, S.

Srinivasan, B.T. Abe, K. Kraft, C. Hellstrom, R. Sjoberg, S. Chang, A. Feng, D.W. Goldman,

A.A. Shah, M. Petri, L.S. Chung, D.F. Fiorentino, E.K. Lundberg, A. Wutz, P.J. Utz, and H.Y.

Chang, *Xist ribonucleoproteins promote female sex-biased autoimmunity.* Cell, 2024. **187**(3):

p. 733-749 e16.

12.   Engreitz, J.M., N. Ollikainen, and M. Guttman, *Long non-coding RNAs: spatial*

amplifiers that control nuclear structure and gene expression. Nat Rev Mol Cell Biol, 2016.

17(12): p. 756-770.

13.   Gutschner, T., M. Hammerle, M. Eissmann, J. Hsu, Y. Kim, G. Hung, A. Revenko, G.

Arun, M. Stentrup, M. Gross, M. Zornig, A.R. MacLeod, D.L. Spector, and S. Diederichs, *The*

noncoding RNA MALAT1 is a critical regulator of the metastasis phenotype of lung cancer

cells. Cancer Res, 2013. **73**(3): p. 1180-9.

14.   Tripathi, V., J.D. Ellis, Z. Shen, D.Y. Song, Q. Pan, A.T. Watt, S.M. Freier, C.F.

Bennett, A. Sharma, P.A. Bubulya, B.J. Blencowe, S.G. Prasanth, and K.V. Prasanth, *The*

nuclear-retained noncoding RNA MALAT1 regulates alternative splicing by modulating SR

splicing factor phosphorylation. Mol Cell, 2010. **39**(6): p. 925-38.

15.   Lamond, A.I. and D.L. Spector, *Nuclear speckles: a model for nuclear organelles.* Nat

Rev Mol Cell Biol, 2003. **4**(8): p. 605-12.

16.   Tripathi, V., Z. Shen, A. Chakraborty, S. Giri, S.M. Freier, X. Wu, Y. Zhang, M.

Gorospe, S.G. Prasanth, A. Lal, and K.V. Prasanth, *Long noncoding RNA MALAT1 controls*

cell cycle progression by regulating the expression of oncogenic transcription factor B-MYB.

PLoS Genet, 2013. **9**(3): p. e1003368.

17.   Eissmann, M., T. Gutschner, M. Hammerle, S. Gunther, M. Caudron-Herger, M. Gross,

P. Schirmacher, K. Rippe, T. Braun, M. Zornig, and S. Diederichs, *Loss of the abundant*

nuclear non-coding RNA MALAT1 is compatible with life and development. RNA Biol, 2012.

9(8): p. 1076-87.

18.   Nakagawa, S., J.Y. Ip, G. Shioi, V. Tripathi, X. Zong, T. Hirose, and K.V. Prasanth,

Malat1 is not an essential component of nuclear speckles in mice. RNA, 2012. **18**(8): p. 1487-

99.

19.   Zhang, B., G. Arun, Y.S. Mao, Z. Lazar, G. Hung, G. Bhattacharjee, X. Xiao, C.J.

Booth, J. Wu, C. Zhang, and D.L. Spector, *The lncRNA Malat1 is dispensable for mouse*

development but its transcription plays a cis-regulatory role in the adult. Cell Rep, 2012. **2**(1):

p. 111-23.

20.   Hewitson, J.P., K.A. West, K.R. James, G.F. Rani, N. Dey, A. Romano, N. Brown, S.A.

Teichmann, P.M. Kaye, and D. Lagos, *Malat1 Suppresses Immunity to Infection through*

Promoting Expression of Maf and IL-10 in Th Cells. J Immunol, 2020. **204**(11): p. 2949-2960.

21.   Dey, S., H. Ashwin, L. Milross, B. Hunter, J. Majo, A.J. Filby, A.J. Fisher, P.M. Kaye,

and D. Lagos, *Downregulation of MALAT1 is a hallmark of tissue and peripheral proliferative*

T cells in COVID-19. Clin Exp Immunol, 2023. **212**(3): p. 262-275.

22.   Masoumi, F., S. Ghorbani, F. Talebi, W.G. Branton, S. Rajaei, C. Power, and F.

Noorbakhsh, *Malat1 long noncoding RNA regulates inflammation and leukocyte differentiation*

in experimental autoimmune encephalomyelitis. J Neuroimmunol, 2019. **328**: p. 50-59.

23.   Xue, Y., J. Ke, X. Zhou, Q. Chen, M. Chen, T. Huang, F. Lin, and F. Chen, *Knockdown*

of LncRNA MALAT1 Alleviates Coxsackievirus B3-Induced Acute Viral Myocarditis in Mice via

Inhibiting Th17 Cells Differentiation. Inflammation, 2022.

24.   Kanbar, J.N., S. Ma, E.S. Kim, N.S. Kurd, M.S. Tsai, T. Tysl, C.E. Widjaja, A.E. Limary,

B. Yee, Z. He, Y. Hao, X.D. Fu, G.W. Yeo, W.J. Huang, and J.T. Chang, *The long noncoding*

RNA Malat1 regulates CD8+ T cell differentiation by mediating epigenetic repression. J Exp

Med, 2022. **219**(6).

25.   Saraiva, M. and A. O'Garra, *The regulation of IL-10 production by immune cells.* Nat

Rev Immunol, 2010. **10**(3): p. 170-81.

26. Kaplan, M.H., J.R. Whitfield, D.L. Boros, and M.J. Grusby, *Th2 cells are required for*

the Schistosoma mansoni egg-induced granulomatous response. J Immunol, 1998. **160**(4): p.

1850-6.

27. Sabin, E.A., M.A. Kopf, and E.J. Pearce, *Schistosoma mansoni egg-induced early IL-*

4 production is dependent upon IL-5 and eosinophils. J Exp Med, 1996. **184**(5): p. 1871-8.

28. Percie du Sert, N., A. Ahluwalia, S. Alam, M.T. Avey, M. Baker, W.J. Browne, A. Clark,

I.C. Cuthill, U. Dirnagl, M. Emerson, P. Garner, S.T. Holgate, D.W. Howells, V. Hurst, N.A.

Karp, S.E. Lazic, K. Lidster, C.J. MacCallum, M. Macleod, E.J. Pearl, O.H. Petersen, F. Rawle,

P. Reynolds, K. Rooney, E.S. Sena, S.D. Silberberg, T. Steckler, and H. Wurbel, *Reporting*

animal research: Explanation and elaboration for the ARRIVE guidelines 2.0. PLoS Biol, 2020.

18(7): p. e3000411.

29. Subramanian, A., P. Tamayo, V.K. Mootha, S. Mukherjee, B.L. Ebert, M.A. Gillette, A.

Paulovich, S.L. Pomeroy, T.R. Golub, E.S. Lander, and J.P. Mesirov, *Gene set enrichment*

analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc

Natl Acad Sci U S A, 2005. **102**(43): p. 15545-50.

30. Eloi-Santos, S., N.J. Olsen, R. Correa-Oliveira, and D.G. Colley, *Schistosoma*

mansoni: mortality, pathophysiology, and susceptibility differences in male and female mice.

Exp Parasitol, 1992. **75**(2): p. 168-75.

31. Coomes, S.M., Y. Kannan, V.S. Pelly, L.J. Entwistle, R. Guidi, J. Perez-Lloret, N.

Nikolov, W. Muller, and M.S. Wilson, *CD4(+) Th2 cells are directly regulated by IL-10 during*

allergic airway inflammation. Mucosal Immunol, 2017. **10**(1): p. 150-161.

32. Hawrylowicz, C.M. and A. O'Garra, *Potential role of interleukin-10-secreting regulatory*

T cells in allergy and asthma. Nat Rev Immunol, 2005. **5**(4): p. 271-83.

33. Yssel, H., R. De Waal Malefyt, M.G. Roncarolo, J.S. Abrams, R. Lahesmaa, H. Spits,

and J.E. de Vries, *IL-10 is produced by subsets of human CD4+ T cell clones and peripheral*

blood T cells. J Immunol, 1992. **149**(7): p. 2378-84.

34. Brockmann, L., S. Soukou, B. Steglich, P. Czarnewski, L. Zhao, S. Wende, T. Bedke,

C. Ergen, C. Manthey, T. Agalioti, M. Geffken, O. Seiz, S.M. Parigi, C. Sorini, J. Geginat, K.

Fujio, T. Jacobs, T. Roesch, J.R. Izbicki, A.W. Lohse, R.A. Flavell, C. Krebs, J.A. Gustafsson, P. Antonson, M.G. Roncarolo, E.J. Villablanca, N. Gagliani, and S. Huber, *Molecular and functional heterogeneity of IL-10-producing CD4(+) T cells.* Nat Commun, 2018. **9**(1): p. 5457.

35.   Shouse, A.N., K.M. LaPorte, and T.R. Malek, *Interleukin-2 signaling in the regulation of T cell biology in autoimmunity and cancer.* Immunity, 2024. **57**(3): p. 414-428.

36.   Cephus, J.Y., M.T. Stier, H. Fuseini, J.A. Yung, S. Toki, M.H. Bloodworth, W. Zhou, K. Goleniewska, J. Zhang, S.L. Garon, R.G. Hamilton, V.V. Poloshukin, K.L. Boyd, R.S. Peebles, Jr., and D.C. Newcomb, *Testosterone Attenuates Group 2 Innate Lymphoid Cell-Mediated Airway Inflammation.* Cell Rep, 2017. **21**(9): p. 2487-2499.

37.   Menees, K.B., R.H. Earls, J. Chung, J. Jernigan, N.M. Filipov, J.M. Carpenter, and J.K. Lee, *Sex- and age-dependent alterations of splenic immune cell profile and NK cell phenotypes and function in C57BL/6J mice.* Immun Ageing, 2021. **18**(1): p. 3.

38.   Seumois, G., C. Ramirez-Suastegui, B.J. Schmiedel, S. Liang, B. Peters, A. Sette, and P. Vijayanand, *Single-cell transcriptomic analysis of allergen-specific T cells in allergy and asthma.* Sci Immunol, 2020. **5**(48).

39.   Tibbitt, C.A., J.M. Stark, L. Martens, J. Ma, J.E. Mold, K. Deswarte, G. Oliynyk, X. Feng, B.N. Lambrecht, P. De Bleser, S. Nylen, H. Hammad, M. Arsenian Henriksson, Y. Saeys, and J.M. Coquet, *Single-Cell RNA Sequencing of the T Helper Cell Response to House Dust Mites Defines a Distinct Gene Expression Signature in Airway Th2 Cells.* Immunity, 2019. **51**(1): p. 169-184 e5.

40.   Deep, D., H. Gudjonson, C.C. Brown, S.A. Rose, R. Sharma, Y.A. Paucar Iza, S. Hong, S. Hemmers, M. Schizas, Z.M. Wang, Y. Chen, D.R. Wesemann, V. Pascual, D. Pe'er, and A.Y. Rudensky, *Precursor central memory versus effector cell fate and naive CD4+ T cell heterogeneity.* J Exp Med, 2024. **221**(10).

41.   Even, Z., A.P. Meli, A. Tyagi, A. Vidyarthi, N. Briggs, D.A. de Kouchkovsky, Y. Kong, Y. Wang, D.A. Waizman, T.A. Rice, B. De Kumar, X. Wang, N.W. Palm, J. Craft, M.K. Basu, S. Ghosh, and C.V. Rothlin, *The amalgam of naive CD4(+) T cell transcriptional states is*

reconfigured by helminth infection to dampen the amplitude of the immune response.

Immunity, 2024. **57**(8): p. 1893-1907 e6.

42.   Gal-Oz, S.T., A. Baysoy, B. Vijaykumar, S. Mostafavi, C. Benoist, T. Shay, and P.

Immunological Genome, *Microheterogeneity in the Kinetics and Sex-Specific Response to*

Type I IFN. J Immunol, 2024. **213**(1): p. 96-104.

43.   Pujantell, M. and M. Altfeld, *Consequences of sex differences in Type I IFN responses*

for the regulation of antiviral immunity. Front Immunol, 2022. **13**: p. 986840.

44.   Tao, X., S. Constant, P. Jorritsma, and K. Bottomly, *Strength of TCR signal determines*

the costimulatory requirements for Th1 and Th2 CD4+ T cell differentiation. J Immunol, 1997.

159(12): p. 5956-63.

Figure Legends

Figure 1. *Malat1* loss impairs *in vitro* Th2 differentiation only in female cells.

A. Schematic of *in vitro* Th2 differentiation protocol.

B. Representative FACS plots of IL10 and IL4 expression in *WT* and *Malat1^{-/-}* female and male

in vitro differentiated Th2 cells (Day 6).

C. Percentage of IL10$^+$ live TCRb$^+$ CD4$^+$ in *in vitro* differentiated Th2 cells derived from female

and male mice at day 4 (prior to IL2 addition) and day 6 (experimental end-point). Levels

determined by intracellular cytokine staining (N=14 per condition).

D. As in **B**, but for IL4$^+$ live TCRb$^+$ CD4$^+$ cells.

E. As in **B**, but for IL4$^+$/IL10$^+$ live TCRb$^+$ CD4$^+$ cells.

F. As in **B**, but for IL4$^+$/IL10$^-$ live TCRb$^+$ CD4$^+$ cells.

G. Log$_2$ fold change (LFC) in gene expression between *WT* female naïve and *in vitro*

differentiated Th2 cells against LFC in gene expression between *WT* and *Malat1^{-/-}* female *in*

vitro differentiated Th2 cells. Data shown for all or the top 500 DEGs between Th2 and naïve

CD4$^+$ T cells from *WT* female mice.

H. As in **E**, but for cells derived from male mice.

I. Percentage of IL10$^+$ live TCRb$^+$ CD4$^+$ *in vitro* differentiated Th2 cells derived from *WT* or *Malat1$^{-/-}$* female and male mice at day 6, with or without treatment with 10 μg/ml of anti-IL10 receptor antibody. Levels determined by intracellular cytokine staining (N=6 per condition).

Figure 2. The effect of *Malat1* depletion on IL10 expression is independent of activation strength and occurs during the early stages of differentiation.

A. Percentage of IL10$^+$ live TCRb$^+$ CD4$^+$ *WT* or *Malat1$^{-/-}$* *in vitro* differentiated Th2 cells derived from female and male mice, with 10 μg/ml or 2 μg/ml of anti-CD3 antibody (see Methods). Levels determined by intracellular cytokine staining (N=9).

B. As in **A**, but for IL4$^+$ live TCRb$^+$ CD4$^+$ cells.

C. *Malat1* levels in *in vitro* differentiated Th2 cells at day 4 (prior to resuspension in IL2) and at day 6 (experimental end point), determined by qRTPCR. Levels normalised to U6 and average levels of *WT* females at day 4 (N=11).

D. Percentage of IL10$^+$ live TCRb$^+$ CD4$^+$ *in vitro* differentiated *WT* Th2 cells derived from female and male mice, treated with either non-targeting control (NTC) GapmeRs or *Malat1* targeting GapmeRs. Levels determined by intracellular cytokine staining (N=6).

E. Percentage of IL10$^+$, IL4$^+$, and IL13$^+$ live TCRb$^+$ CD4$^+$ *in vitro* differentiated *WT* Th2 cells derived from female and male mice treated with NTC or *Malat1* targeting GapmeRs, with and without treatment with an anti-IL10 receptor antibody. Levels determined by intracellular cytokine staining (n=6).

Figure 3. *Malat1* loss enhances naïve CD4$^+$ T cell activation and impairs early Th2 differentiation.

A. Percentage of CD69$^+$ live TCRb$^+$ CD4$^+$ *WT* or *Malat1$^{-/-}$* *in vitro* activated CD4$^+$ T cells derived from female or male mice at 0, 3, 7, and 24 hours post-activation. Levels determined by surface staining (n=6).

B. *Gata3* mRNA levels at days 1, 2, 3 and 4 of Th2 differentiation determined by qRTPCR

(n=3). Levels normalised to U6 and average levels of *WT* females at day 1.

C. *Il2ra*, *Il2rg*, and *Il2* mRNA levels at days 1, 2, 3 and 4 of Th2 differentiation determined by

qRTPCR (n=3). Levels normalised to U6 and average levels of *WT* females at day 1.

D. *Il2ra*, *Il2rg* and *Gata3* mRNA levels at day 2 of differentiation determined by qRTPCR (n=9).

Levels normalised to *U6* and average levels of *WT* females.

E. Percentage and Median Fluorescence Intensity (MFI) of GATA3$^+$ live TCRb$^+$ CD4$^+$ *WT* or

Malat1$^{-/-}$ cells derived from female or male mice at day 2 of differentiation (n=5 per condition).

Figure 4. Impaired suppression of an interferon gene module during early

differentiation of female *Malat1$^{-/-}$* Th2 cells.

A. Volcano plot displaying LFC in gene expression between *WT* and *Malat1$^{-/-}$* female cells

(determined by Nanopore long-read RNAseq, N=4 per group) against -logFDR at day 2 of *in*

vitro differentiation.

B. Gene set enrichment analysis (GSEA) hallmark gene set terms enriched within significantly

up-regulated genes in *Malat1$^{-/-}$* cells at day 2 of *in vitro* differentiation.

C. Heatmap displaying Z score of log$_2$CPM for top DEGs (FDR<0.01) between *WT* and *Malat1^{-}*

$^{/-}$ cells at day 2 of differentiation, in descending order of LFC.

D. LFC in gene expression between female *WT* naïve and differentiating cells at day 2 of *in*

vitro differentiation against LFC between *WT* and *Malat1$^{-/-}$* female cells at day 2 of *in vitro* Th2

differentiation. Data shown for all statistically significant DEGs between *WT* and *Malat1$^{-/-}$* cells

at day 2.

E. LFC in gene expression between female *WT* naïve and Th2 cells (as in Figure 1) against

LFC between *WT* and *Malat1$^{-/-}$* female cells at day 2 of *in vitro* Th2 differentiation. Data shown

for all statistically significant DEGs between *WT* and *Malat1$^{-/-}$* cells at day 2. Colours indicate

level of significance in differential expression when comparing day 2 vs naïve CD4$^+$ T cells

(left) or Th2 vs naïve CD4$^+$ T cells (right).

F. Levels of interferon induced genes (*Ifit1bl1*, *Ifit3b*, *Iigp1*, *Ifit1*) and transcription factors

(*Stat1, Irf1, Irf7*) in *WT* and *Malat1*[-/-] *in vitro* differentiated female or male cells at day 2 of

differentiation. mRNA levels determined by qRTPCR (N=9) and normalised to *U6* and average

levels of *WT* cells from female mice.

G. Percentage of IL10+ and IL4+ live TCRb+ CD4+ *in vitro* differentiated Th2 cells derived from

WT or *Malat1*[-/-] female and male mice at day 6, treated with 0 (control) or 5000 U/ml IFNβ.

Levels determined by intracellular cytokine staining (N=8 for all conditions, except control

treated *WT* male, which is n=7).

Figure 5. *Malat1* loss decreases expression of the IL2 receptor and impairs the

response to exogenous levels of IL2 in cells from female mice.

A. Percentage of IL2Rα+ live TCRb+ CD4+ *WT* or *Malat1*[-/-] cells derived from female or male

mice at day 4 of differentiation (levels determined by surface staining, N=6).

B. Percentage of IL2Rγ+ live TCRb+ CD4+ *WT* or *Malat1*[-/-] cells derived from female or male

mice at day 4 of differentiation (levels determined by surface staining, N=5).

C. Percentage of IL2Rα+ live TCRb+ CD4+ *WT* cells derived from female and male mice at day

4 of differentiation that were treated with either non-targeting control (NTC) or *Malat1* targeting

GapmeRs. Levels determined by surface staining (N=5).

D. As in **C**, but for IL2Rγ+ live TCRb+ CD4+ cells.

E. Percentage of IL2Rα+ live TCRb+ CD4+ *WT* cells derived from female and male mice at day

4 of differentiation, treated with 0 (control) or 5000 U/ml IFNβ. Levels determined by surface

staining (N=8).

F. As in **E**, but for IL2Rγ+.

G. Normalised percentage of IL10+, IL4+ or IL10+ IL4+ live TCRb+ CD4+ *WT* or *Malat1*[-/-] *in vitro*

differentiated Th2 cells derived from female and male mice at day 6, with resuspension at day

4 in the indicated concentrations of IL2. Levels determined by intracellular staining, and

normalised to average levels in *WT* female-derived cells treated with 10U/ml IL2 from each

experiment (N=10).

Figure 6. *Malat1* loss causes a female-specific decrease in Th2 cell-derived IL10 levels

in vivo.

A. Schematic of *S. mansoni* egg injection experiment. 5,000 dead *S. mansoni* eggs were

injected interperitoneally (IP) into mice, then after a week the injection was repeated

intravenously (IV) via the tail vein. After another week, lungs and spleens were harvested and

processed for FACS.

B. Representative FACS plots of IL10 and IL4 expression in lung CD4$^+$ TCRb$^+$ cells from *S.*

mansoni egg injected *WT* or *Malat1$^{-/-}$* female or male mice.

C. Percentage and cell numbers of lung IL10$^+$ TCRb$^+$ CD4$^+$ cells derived from naïve or *S.*

mansoni egg injected (*Sm*) *WT* or *Malat1$^{-/-}$,* female and male mice. Levels determined by

intracellular staining (N=3 for *Malat1$^{-/-}$* male naïves, N=5 for all other naïves, N=9 for egg

injected mice).

D. As in **C**, but for splenic IL10$^+$ TCRb$^+$ CD4$^+$ cells (N=4 for *Malat1$^{-/-}$* male naïves, N=5 for all

other naïves, N=9 for egg injected mice).

E. Percentage and cell numbers of lung IL10$^+$ IL4$^+$ TCRb$^+$ CD4$^+$ cells derived from naïve or *S.*

mansoni egg injected (*Sm*) *WT* or *Malat1$^{-/-}$,* female and male mice. Levels determined by

intracellular staining (N=3 for *Malat1$^{-/-}$* male naïves, N=5 for all other naïves, N=9 for egg

injected mice).

F. As in **E**, but for splenic IL10$^+$ TCRb$^+$ CD4$^+$ cells (N=4 for *Malat1$^{-/-}$* male naïves, N=5 for all

other naïves, N=9 for egg injected mice).

Figure 7. Schematic summary of the effect of *Malat1* loss on Th2 cell differentiation*.*

Top row shows the effects observed when comparing *Malat1$^{-/-}$* female-derived cells to *WT*.

Second row when comparing male-derived *Malat1$^{-/-}$* and *WT* cells. Third row when comparing

male to female-derived *WT* cells. Findings based on the *in vitro* process shown in Fig. 1A.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

	Activation	Early differentiation	Differentiation	Terminal differentiation
Naïve CD4+ T cell				
Malat⁻/⁻ female vs *WT* female	Accelerated CD69 induction	Impaired ISG suppression GATA3 suppression	IL2Rα and IL2Rγ suppression	IL10 suppression Transcriptome-wide suppression
Malat⁻/⁻ male vs *WT* male	No differences	GATA3 suppression	IL2Rγ suppression	No differences
WT male vs *WT* female	Accelerated CD69 induction	Enhanced ISG expression	Reduced response to IL2	Suppressed cytokine expression (*in vitro*)

Naïve CD4+ T cell	CD69 expression	GATA3 expression Suppression of ISG expression	IL2R expression Minimal cytokine expression	IL10 expression ⊣ IL4, IL13 Expression