Climate, weather, and geography shape seed mass variation and decline across western North America

Leah J. Lenzo[1], Matthew L. Forister[1], Peggy Olwell[2], Elizabeth A. Leger[1]

[1]University of Nevada, Reno; Department of Biology, Graduate Program in Ecology, Evolution, and Conservation Biology, 1664 N. Virginia St, Reno, NV 89557 U.S.A.

[2] Plant Conservation and Restoration Program, Bureau of Land Management, 1387 S Vinnell Way, Boise, Idaho, 83709 U.S.A.

Abstract

Seed mass influences the probability of germination, the timing of emergence,

and the likelihood of survival and establishment, and can affect restoration and

conservation efforts. Here, we investigated the temporal, geographic, and

environmental patterns that influence seed mass across 2,800 species of native plants

from over 13,000 populations across the western US using two decades of collections

from a national program, Seeds of Success. The majority of species exhibited low

variation in seed mass, though there were exceptions, with some species varying in

seed mass by up to 220% over their occupied area. Species with the most variation in

seed mass were collected across sites with a wide range of minimum temperatures,

from more northern latitudes, and collected across a wide range of dates, and seeds

were generally larger in areas where it was wetter and warmer. We observed an overall

decrease in seed mass by an average of 0.012 mg each year, though there were many

species-specific differences. Fifty-five percent of species decreased in seed mass, 20%

strongly so, and 17% of species increased appreciably in mass. Relationships between

seed mass and environmental variables varied widely in strength, identity, and direction

among species, including those that often co-occur. Measures of climate change (30

year changes in seasonal temperatures and precipitation) were associated with

changes in seed mass in nearly 20% of species. In sum, these results support the need

to continue native seed collections for conservation and restoration, as plant

reproductive strategies may be shifting with changing climates.

Significance

Seeds are an essential reproductive strategy for most plants, and seed mass is

predicted to respond to environmental factors, but we do not know if climate-related size

changes observed in other organisms will also be reflected in seed mass. We assessed

how weather, climate, and geography affect seed mass in thousands of native plants of

the western US. Minimum temperatures and precipitation had strong impacts on seed

mass for many species, though there were many species-specific responses. Overall,

seed mass declined for more than half of observed species. Measures of climate

change influenced seed mass in nearly one in five species, which could be evidence of

adaptation to new optimal values, or be a signal of challenges ahead.

Keywords: Seed mass; Intraspecific variation; Climate Change; Interannual variation

Introduction

Seed production is an essential component of fitness for the majority of plants.

Because adult plants are immobile, barring long distance dispersal, their reproductive

strategies must adapt to local environments in order to persist (Clausen et al. 1940;

Leimu and Fischer 2008; Baughman et al. 2019). Seed mass can strongly influence

plant fitness during the seed-to-seedling transition, a crucial life history stage (Moles

and Westoby 2004; Norden et al. 2009; Larson et al. 2015; Leger et al. 2019). Natural

selection should favor an optimal seed mass, balancing seed mass with seed number,

with local optimums determined by interacting selection pressures including

competition, dispersal, and herbivores (Venable 1992; Muller-Landau 2010; Seale and

Nakayama 2020; Tuthill et al. 2023). Yet, seed mass varies greatly within and among

individuals (Völler et al. 2012; Liu et al. 2013) possibly due to fitness benefits of

variation in seed mass such as reducing competition among siblings, or bet-hedging in

unpredictable landscapes (McGinley et al. 1987; Westoby et al. 1996). Variation in

environmental sensitivity of seed mass could lead to the loss of some species in

response to climate change if there is a mis-match between local optima and expressed

traits, or increasing dominance of highly variable or adaptable species. In the era of

climate change, understanding how environmental factors influence seed mass as a

reproductive strategy is critical for predicting species persistence (Cochrane et al. 2015;

Tangney et al. 2020).

Seed traits vary across geographic, temporal, and phylogenetic scales (Westoby

et al. 1992; Moles et al. 2007; Gorden et al. 2016). Larger seeds are typically more

common in resource-poor environments where maternal provisioning is necessary for

survival, and the opposite in resource-rich environments, where many small seeds may

increase fitness (Smith and Fretwell 1974; Adler et al. 2013). At the largest spatial

scale, environment affects seed mass across latitudinal gradients (Moles et al. 2007),

with larger seeded species found closer to the equator. Hypotheses for this pattern

include correlations with plant growth form (greater dominance of trees in the tropics),

differences in seed dispersal strategies (e.g. wind vs. animal dispersal), and net primary

productivity (greater towards the equator) (Murray et al. 2004). Latitudinal gradients in

seed mass are present but weaker in North America (Liu et al. 2013; Gorden et al.

2016), possibly due to topographic variability (Malanson et al. 2007; Albrecht and

McCarthy 2009). Further, these broad gradients may result from taxonomic turnover

across latitudes but may not explain intraspecific, among-population variation in seed

mass (Hulshof et al. 2013). It is also possible that within a community, species may

produce different seed sizes in response to the same environmental conditions, which

would be predicted if seed size plays a role in niche partitioning and community

assembly (Turnbull et al. 1999; Leishman 2001; Muller-Landau 2010). Understanding

the geographic, temporal, and taxonomic patterns of seed mass will provide valuable

insight into evolutionary pressures experienced by native plants, including ongoing

climate change (Frenne et al. 2013; Cochrane et al. 2015).

Biologists have characterized variation in seed characteristics for centuries, yet

only recently do we have the fine-scaled resolution and longitudinal data to closely link

seed traits with maternal environmental conditions (Kattge et al. 2011; Liu et al. 2019).

In North America, we have an opportunity to examine the effects of geography and

environment on seed mass through Seeds of Success (hereafter "SOS" see:

blm.gov/sos), which is the national native seed collection effort in the United States led

by the Bureau of Land Management. Through collaborative efforts and labor of

thousands of field interns and seed processing technicians, SOS has made over 27,000

native seed collections across the United States since 2001 (Barga et al. 2020) (Fig.

1*A*). The mission of SOS is to collect wildland native seed for development,

conservation, and restoration, and target species include restoration priorities as well as

taxa not previously collected. Therefore, the dataset includes multiple accessions of the

same species over multiple years from many locations, from a broad range of taxa,

collected using a consistent protocol. The scale and scope of this effort has resulted in

an invaluable source of seed for conservation and restoration across the US, and has

generated an unparalleled wealth of information about the properties of seed in wild

plant populations that can be used to inform ecological theory, global change biology,

and restoration practices (Siefert et al. 2015; Moran et al. 2016).

To assess patterns of seed mass variation within and among species, we first

took a broad approach, maximizing the inclusion of data to identify patterns of seed

mass variation at the largest possible scale, asking 1) where do plants produce bigger

seeds, and is there evidence of widespread changes in seed mass over time. Then, we

investigated environmental influence on seed mass for more well-sampled species, both

as a group and individually, asking 2) if there are environmental, climate variables, or

life history traits that have a consistent influence on seed mass across and within

species. Finally, we asked 3) if environment and geography predict not only seed mass

but also seed mass variation across the range of collections, including a phylogenetic

perspective.

Results

Seed mass ranged from 0.0046 mg (the average across-population mass of

Epipactis gigantea) to 827.7 mg (for *Pinus sabiniana*) with a mean across species of 4.4

mg and median of 1.2 mg (Fig. 1D). Certain plant families, including Fabaceae and

Ericaceae, consistently had larger seeds whereas others (e.g. Juncaceae and

Onagraceae) exhibited smaller seeds (Fig. 2). Seed mass ranged from highly

conserved across populations (e.g. CV of 0.0005 for *Wyethia magna*) to highly variable

(CV of 2.2 for *Juncus occidentalis*), with a mean CV of 0.32 and median of 0.27 (Fig. 2).

Most areas across the western US were dominated by smaller seeds with no

obvious spatial segregation (Fig. 1*A*; *SI Appendix* 1, Fig. S1). Latitude had a small,

positive effect on seed mass in models that included latitude, longitude, and elevation

as predictors (GLM: β = 0.027, p < 0.001) as well as models that included plant family

as a random factor, which had a better model fit than geography alone (GLMM: β =

0.031, p < 0.001; Fig. 1*B, SI Appendix* Q1, Table S2). Seeds were, on average, 1.03 mg

larger for every 1 degree north, in contrast to global patterns, though in models that

considered seed mass by growth form, shrubs had a contrasting response to other plant

growth forms, with smaller seeds in more northern latitudes (*SI Appendix Table* S2).

Longitude had a small negative effect on seed mass (GLM: β =−0.005, p = 0.089,

GLMM: β =−0.01, p < 0.001), with smaller seeds (0.989 mg smaller for every 1 degree)

in more eastern areas.

We examined relationships between seed mass, environment, and time for well

collected species (10,958 collections from 859 species collected from at least 3

populations) and found that, overall, seed mass decreased by 0.012 mg each year (Fig.

1C, β = -0.010, SE = 0.004, p = 0.007). When considering species independently, 38%

changed appreciably in seed mass, representing a distribution of change with greater

density in the tails of the distribution (species getting larger or smaller) than would be

expected from a normal distribution (kurtosis = 11.153, z = 10.596, p < 0.001). Of the

631 species that were collected from 3 unique populations and years, 55% were

declining in seed mass over time, with 132 declining significantly and 107 increasing

significantly (at p < 0.05) in seed mass. Multiple environmental variables had consistent

positive effects on seed mass across species (Fig. 3A). The strongest positive

relationships were between seed mass and cumulative precipitation over the previous 6

months leading up to collection (β = 0.051, SE = 0.012, p < 0.0001), as well as annual

minimum temperature (β = 0.044, SE = 0.011, p < 0.001), with generally greater seed

mass in the relatively wetter and warmer collection sites (Fig. 3A, SI Appendix Table

S4). Clay content and slope also had positive relationships with seed mass (β = 0.045,

SE = 0.011, p < 0.001, and β = 0.045, SE = 0.011, p < 0.001 respectively), with

generally larger seeds in areas with higher clay content and steeper slopes. Heat load

also had a positive relationship with seed mass (β = 0.024, SE = 0.012, p = 0.038), with

larger seeds in relatively warm collection sites. Neither growth form (shrub, grass, forb,

etc.) nor duration (annual, perennial, etc.) were retained as important effects in the

model selection process.

In the results described thus far, we maximized statistical power for the detection

of environmental effects by including all species with at least 3 collections. Now we turn

to results using fewer species, with 20 collections each (111 species, 5196 total

collections); these models allow us to examine variation among species in environment-

seed mass effects. We found several common predictors of seed mass among species,

confirming previous results (Fig. 4, *SI Appendix* Table S6). Specifically, annual

minimum temperature, clay content, heat load, cumulative precipitation six months prior

to collection, soil pH, and day of year were statistically significant in at least 10 species

models. However, the strength and direction of relationships varied widely among

species (Fig. 3*C-N*; Fig. 4). For example, soil pH had a strong positive influence on

seed mass of *Balsamorhiza hookeri*, increasing seed mass 1.095 mg as soil pH

increased by 1 (Fig. 3*C*, Fig. 4), while *Elymus elymoides* had the opposite response,

decreasing seed mass by 0.889 mg as soil pH increased by 1 (Fig 3*D*, Fig 4). We used

a resampling approach to investigate the influence of variation in sample size (the

number of populations) on species-specific results; results were robust to systematic

reductions in the number of populations included in models (*SI Appendix*).

Multiple measures of climate change across a species collection range (including

change in spring and fall maximum temperatures, change in precipitation, and change

in fall minimum temperatures) were included as important predictors in 47 species

models (out of a possible 111 species) and were significant 23 times in 22 models (Fig.

4). Of those, 11 had a statistically significant negative correlation between measures of

climate change and seed mass and 12 climate change coefficients were positive, again

illustrating heterogeneity of responses (Fig. 4, *SI Appendix* Table S5). *Atriplex*

polycarpa and *Artemisia tridentata* were particularly responsive to climate change,

including three measures of climate change in each species model, with change in

precipitation significantly affecting seed mass (*SI Appendix* Table S6).

Our models thus far have assessed mean seed mass, but because natural

selection acts on variation, we also quantified seed mass variation among populations

of a single species to understand the potential for evolutionary change. Here, we

aggregated our data at the species level to ask how environment and geography predict

seed mass coefficient of variation while considering phylogenetic relationships among

species. Species differed in the amount of variation (CV) in seed mass across collection

locations and collection years (Fig. 2). Most species exhibited low variation in seed

mass, but some, including *Artemisia frigida, Phacelia hastata, Chaenactis stevioides,*

and *Sphaeralcea munroana*, exhibited CV values above 1 indicating higher variation.

Considering the 235 species that had been collected at least 10 times and for which

phylogenetic information was available, the strongest predictors of seed mass variation

were the maximum latitude from which a species was collected ($\beta = 0.091$, SE = 0.037,

p = <0.001), the range of annual minimum temperatures experienced across the

collection area ($\beta = 0.068$, SE = 0.041, p = 0.101), and the range of collecting days

(difference between first and last day of collection) ($\beta = 0.055$, SE = 0.041, p = 0.180;

Fig. 3*B, SI Appendix* Table S7). Neither growth form nor duration were selected as

important, and, when included in models with environmental covariates or alone in a

simple model, neither predicted seed mass variation (results not shown).

Discussion

Climate change continues to affect the morphology of individual species of plants

and animals, but due to challenges of amassing quality datasets, effects across entire

assemblages of species are rarely investigated (Sheridan and Bickford 2011; Violle et

al. 2012). Further, there is increasing consensus about the importance of intraspecific

variation for understanding ecological and evolutionary processes that affect species

and community responses to climate change (Siefert et al. 2015). Yet, we often lack

consistent population-level replication for specific fitness related traits, leading to

uncertainties in identifying these intraspecific patterns (Moran et al. 2016). By assessing

seed mass variation within and among species simultaneously, our results offer a

unique perspective on plant reproductive biology and species coexistence, illuminating

how environment affects propagule size in land plants.

We found spatial patterns and environmental predictors of seed mass across all

species and common environmental predictors of seed mass within multiple species,

including precipitation, heat load, and soil characteristics like pH. Seeds were generally

larger in warmer and wetter areas, consistent with tree seed sizes observed in Boreal

forests (Liu et al. 2013). However, we observed even stronger signals within species,

though often in heterogeneous directions and magnitudes, confirming that intraspecific

seed trait-environment relationships can be idiosyncratic, and differ from broad

interspecific patterns (Gorden et al. 2016). Heterogeneity in response among species

may be an important form of resilience, which can only be seen when analyzing traits at

the scale of communities and entire floras. Conservation and restoration practices could

benefit from considering intraspecific variation in addition to interspecific variation (Sgrò

et al. 2011). While conservationists may prioritize protecting species that are more

sensitive to climate and climate change, practitioners may benefit from planting species

with seed traits that are less sensitive to climate change or selecting a mix of species

that are sensitive to different factors, thereby incorporating natural resilience into

restoration.

While seed mass was highly conserved for most species, others varied more

across space and time, including species collected across broader minimum

temperature gradients, higher latitudes, and those with longer collecting seasons. Given

that warming minimum temperatures are an early consequence of anthropogenic

climate change (Parry et al. 2007), with northern latitudes showing particular acute

change (Bunn et al. 2007), these sensitivities suggest that climate change will be

exerting selection pressure on seed mass in plants. Indeed, we observed overall

declines in seed mass across the western US over the past 20 years, though some

species increased in seed mass, consistent with heterogeneous changes observed in

seed mass in alpine plants (Zhang et al. 2019). Given the importance of seed mass for

plant regeneration (Moles and Westoby 2004) and the role of seeds as a food source

for granivores especially in arid lands (Solbreck and Knape 2017; Dylewski et al. 2020),

these small changes in seed mass could have cascading effects on ecosystems

(Sheridan and Bickford 2011).

Further highlighting the impact of climate change on terrestrial plant seeds, we

detected sensitivity to direct measures of temperature and precipitation changes for

approximately one fifth of well-collected species, though directionality was not

consistent. For example, showy milkweed, *Asclepias speciosa*, an important host plant

for the Monarch butterfly, makes larger seeds in warmer areas, while *Plantago ovata*

shows the opposite pattern. Another very important species in the region, big sagebrush

(*Artemisia tridentata),* was particularly responsive to climate change, with three of our

four measures of climate change affecting seed mass. It is possible that seed mass

sensitivity to climate change represents an adaptive response, or if changes are a result

of resource limitation, it could indicate that these species are particularly vulnerable to

climate change. Field studies of the relationship between seed mass and fitness in

highly sensitive species identified here would be important to understand if these plants

should be prioritized in conservation efforts.

We are particularly fascinated by the many cases where species had strong

responses to the same variables, but in contrasting ways (Fig. 4). For example, soil pH

had a strong positive relationship with seed mass in the widespread forb *Balsamorhiza*

hookeri, while *Elymus elymoides*, a widely distributed perennial grass commonly used in

ecological restoration had the opposite response. Other species did not change seed

mass in response to any of the variables we considered, maintaining nearly constant

seed mass across collection locations and years (e.g. *Chimaphila umbellata*). Some

species varied in seed mass across space and time but not in response to any

environmental variables we included here (e.g. *Chaenactis douglasii*). It may be that

other ecological influences such as herbivory, competition, success of insect pollination,

or polyploidy may have an effect on seed size in these species (Paige and Whitham

1987; Vaughton and Ramsey 1997; Chan et al. 2022). In the future, close monitoring of

native plant communities may uncover these ecological effects on seed mass, and our

results could inform comparative studies across species with contrasting levels of seed

mass variation.

Overall, few species exhibited high (>1) coefficient of variation, supporting our

understanding that seed mass is a highly conserved trait with some exceptions (Zhang

et al. 2020), and it is possible that species that are more variable in seed mass may be

poised to adapt to changing climates (Frei et al. 2014; Cochrane et al. 2015). However,

species demonstrating changes in seed mass in response to climate change may be

adjusting seed mass in maladaptive ways. For example, if smaller seeds confer greater

dispersal capability, but a longer, warmer growing season results in larger seeds, this

may reduce dispersal distances and lead to greater competition with siblings (Harper et

al. 1970; Greene and Johnson 1993). In contrast, species that are demonstrating

decreasing seed mass in response to warming temperatures may be experiencing

resource limitation, and changes in seed mass may reduce progeny resources and

stress tolerance (Westoby et al. 1996).

Understanding where and how species vary in seed mass can help elucidate the

reproductive strategies of land plants, and understanding the consequences of these

changes will be essential for informing applied science. This work provides a foundation

for multifaceted future investigations into these issues, and highlights the need to

increase our population level monitoring and consider intraspecific variation in key traits.

Our findings also support the need for continued collection of locally adapted native

seeds from many sites for restoration through Seeds of Success (National Academies

of Sciences, Engineering, and Medicine 2023), and the species-specific nature of seed

mass responses to climate support targeting a wide range of collection sites to collect

and preserve high quality seed for multiple species. Additionally, because the

reproductive output of many native plant species in the western US is sensitive to

climate change, the urgency to continue native seed collections for conservation is

greater than ever.

Methods

Seed mass from Seeds of Success

SOS collecting teams follow a consistent seed collection protocol (*SI Appendix*

1), collecting from at least 50 plants of all sizes per location, aiming for a representative

population sample. The majority of the SOS seed collections are made in the western

continental US, and the vast majority of those are processed at the USDA Forest

Service Seed Extractory in Bend, Oregon (BSE, *SI Appendix* 1). Seed mass is

assessed at BSE by weighing 5 subsamples of 100 seeds from each collection. After

merging BSE and national SOS datasets, we retained 13,004 unique accessions with

seed mass, collection date, and location information (Fig. 1*A*).

Environmental Data

We explored a variety of potential environmental factors that may affect seed

characteristics, and selected 14 environmental variables to move forward into analysis:

four measures of local change in temperature and precipitation, three annual weather

variables (annual minimum temperature, cumulative precipitation over the 6 months

prior to collection, and a composite of temperature and precipitation of the month prior

to collection), slope, aspect, an estimate of heat load (Dilts et al. 2015), and four soil

variables (clay content, an estimate of water content, soil pH, and organic content

(Beaudette et al. 2022) (*SI Appendix* 2, Table S1).

Statistical Analyses

We first investigated spatial patterns of seed mass by using the full SOS dataset

in generalized linear (GLM) and mixed (GLMM) models with log-transformed seed

mass. Then, we subset the data to species collected from at least 3 populations to ask

whether there were environmental influences on seed mass that could be estimated

across a large species pool, with seed mass scaled within each species. Next, we

reduced the data to species collected from at least 20 populations, and ran individual

species models, with log-transformed seed mass, asking how the local environment

influences seed mass for that subset of well-sampled species. Finally, we aggregated

our data at the species level to ask if environment and geography predict seed mass

variation for species that had been collected at least 10 times, including a phylogenetic

perspective. Models were assessed with visual inspection of QQ plots and VIF scores.

We mapped seed mass to visually assess spatial patterns separately for

grasses, forbs, and shrubs, using the full dataset of 13,004 accessions. To predict seed

mass across the western United States, we used inverse Distance Weighting (IDW)

interpolation of the full dataset using the idw function in the gstat package in R

(Pebesma and Graeler 2023). To test if there is a spatial gradient in seed mass among

all species and populations in the western United States, we ran a GLM with log-

transformed seed mass as the response variable using the full dataset with latitude,

longitude and elevation as predictors; we also ran a second model that was identical but

added growth form and duration as fixed effects. We next ran a GLMM with the same

three spatial variables and response variable but also included plant family as a random

factor to account for taxonomic variation. Not all species were collected from multiple

populations, therefore we used family, rather than species, as a random intercept effect

using the lmer function from the lme4 package in R (Bates et al. 2024). Similar to the

GLM, we also ran a version of this model that included growth form and duration as

fixed effects. After observing an effect of growth form in this model, we ran individual

GLM and GLMM models for three highly collected growth forms to understand how

seed-environment relationships varied among shrubs, grasses, and forbs.

To investigate whether there were consistent influences on seed mass across

species, we analyzed relationships between seed mass and a series of predictors for all

species that had a minimum of 3 collections in a spatial simultaneous autoregressive

error model (SAR) using the errorsarlm function from the R package spatialreg (Bivand

et al. 2024). This dataset contained 10,958 collections from 859 species from 73 plant

families, and we scaled seed mass within each species. Prior to running the SAR,

covariate selection was done using a random forest variable selection pipeline across

all species, which implemented the Boruta function from the Boruta package in R (max

runs = 100, $p < 0.0001$, *SI Appendix* Q2) (Kursa and Rudnicki 2010), considering the

suite of 14 environmental variables discussed above (*SI Appendix* 2, Table S1), along

with day of year, year, plant growth form and duration. We also assessed how seed

mass has changed each year for species collected in at least 3 years (631) with

individual models containing year as a predictor, and tabulated the standardized

coefficients (*SI Appendix* Q2).

To understand how local environmental variables affect seed mass for each of

the most well-sampled species separately, we ran individual SAR models for each

species that was collected a minimum of 20 times (N = 111 species, N = 5196 total

collections, *SI Appendix*). We initiated covariate selection using the random forest

feature selection pipeline (described above and *SI Appendix* Q3) separately for each

species. For seven species, no variables were confirmed, and thus they were not

analyzed but are included in our presentation of results. To determine whether sampling

effort affected our ability to estimate environment-seed mass relationships, we ran

additional linear models on subsets of data (*SI Appendix* Q3).

Finally, to understand the factors that affect the degree of variation in seed mass

among species, we calculated the coefficient of variation (CV) of seed mass for each

species that had been collected at least 10 times by dividing the standard deviation by

the mean seed mass per species, corrected for sampling effort (*SI Appendix* Q4). We

incorporated phylogenetic relatedness in these models using relationships described in

Mishler et al. (2020). Seventy-nine species were not present in the phylogenetic tree

even after correction and cross checking taxonomy, thus, our final model for assessing

variation in seed mass included 235 species from 6865 populations.

We aggregated environmental and climate variables across all collections for

each species for a total of 33 environmental, spatial, temporal, and life history variables

in addition to growth form and duration describing each of 235 species (*SI Appendix* Q4,

Table S1). Next, we ran all 33 variables through the random forest feature selection

pipeline (described above and *SI Appendix*). Confirmed important attributes were then

moved into a GLM with log-transformed unbiased CV as response to check VIF (all

were below 5) and model diagnostics. Then, attributes were moved into a phylogenetic

generalized least squares regression (PGLS) using Pagel's correlation of trait evolution

and the phylogeny from Mishler et al. (2020).

Acknowledgements

This work was partially supported by U.S. Department of Interior Bureau of Land Management award L20AC00317 and L22AC00506 to E. A. L. We thank J. Allen, K. Badik, E. Hanan, A. Agneray, J. McClinton, T. Bartz, R. Frederick, L. Shriver, C. Silliman, S. Swim, A. Kim, B. Nagelson, C. Phelan, H. Noble, S. Bisbing, M. Lohman, and J. Brockman for input on questions, analysis, and data visualization. The Seeds of Success program and data would not be possible without the hard work of seed collection crews, seed processing and cleaning technicians, availability of steed storage, and the expertise and knowledge of botanists and ecologists. The authors thank all their federal, state, and nongovernmental agency partners whose efforts towards native plant community restoration and conservation in the U.S. created this rich dataset.

Data Availability

The dataset will be available on Dryad.

Literature Cited

Adler PB, Fajardo A, Kleinhesselink AR, Kraft NJB. 2013. Trait-based tests of

coexistence mechanisms. Ecol Lett. 16(10):1294–1306. doi:10.1111/ele.12157.

Albrecht MA, McCarthy BC. 2009. Seedling establishment shapes the distribution of

shade-adapted forest herbs across a topographical moisture gradient. J Ecol.

97(5):1037–1049. doi:10.1111/j.1365-2745.2009.01527.x.

Barga SC, Olwell P, Edwards F, Prescott L, Leger EA. 2020. Seeds of Success: A

conservation and restoration investment in the future of U.S. lands. Conserv Sci

Pract. 2(7):e209. doi:10.1111/csp2.209.

Baughman OW, Agneray AC, Forister ML, Kilkenny FF, Espeland EK, Fiegener R,

Horning ME, Johnson RC, Kaye TN, Ott J, et al. 2019. Strong patterns of

intraspecific variation and local adaptation in Great Basin plants revealed through

a review of 75 years of experiments. Ecol Evol. 9(11):6259–6275.

doi:https://doi.org/10.1002/ece3.5200.

Beaudette D, Skovlin J, Roecker S, Brown A. 2022. soilDB: Soil Database Interface.

[accessed 2022 Oct 5]. https://CRAN.R-project.org/package=soilDB.

Bivand R, Piras G, Anselin L, Bernat A, Blankmeyer E, Chun Y, Gómez-Rubio V, Griffith

D, Gubri M, Halbersma R, et al. 2024. spatialreg: Spatial Regression Analysis.

[accessed 2024 Mar 11]. https://cran.r-

project.org/web/packages/spatialreg/index.html.

Bunn AG, Goetz SJ, Kimball JS, Zhang K. 2007. Northern high-latitude ecosystems

respond to climate change. Eos Trans Am Geophys Union. 88(34):333–335.

doi:10.1029/2007EO340001.

Chan JCS, Ooi MKJ, Guja LK. 2022. Polyploidy but not range size is associated with

seed and seedling traits that affect performance of pomaderris species. Front

Plant Sci. 12. doi:10.3389/fpls.2021.779651. [accessed 2024 May 13].

https://www.frontiersin.org/journals/plant-

science/articles/10.3389/fpls.2021.779651/full.

Clausen J, Keck DD, Hiesey WM. 1940. Experimental studies on the nature of species.

I. Effect of varied environments on western North American plants. Exp Stud Nat

Species Eff Varied Environ West North Am Plants.(520). [accessed 2023 Jan

25]. https://www.cabdirect.org/cabdirect/abstract/19421601306.

Cochrane A, Yates CJ, Hoyle GL, Nicotra AB. 2015. Will among-population variation in

seed traits improve the chance of species persistence under climate change?

Glob Ecol Biogeogr. 24(1):12–24. doi:10.1111/geb.12234.

Dilts TE, Weisberg PJ, Dencker CM, Chambers JC. 2015. Functionally relevant climate

variables for arid lands: a climatic water deficit approach for modelling desert

shrub distributions. J Biogeogr. 42(10):1986–1997. doi:10.1111/jbi.12561.

Dylewski Ł, Ortega YK, Bogdziewicz M, Pearson DE. 2020. Seed size predicts global

effects of small mammal seed predation on plant recruitment. Ecol Lett.

23(6):1024–1033. doi:https://doi.org/10.1111/ele.13499.

Frei ER, Ghazoul J, Matter P, Heggli M, Pluess AR. 2014. Plant population

differentiation and climate change: responses of grassland species along an

elevational gradient. Glob Change Biol. 20(2):441–455. doi:10.1111/gcb.12403.

Frenne PD, Graae BJ, Rodríguez-Sánchez F, Kolb A, Chabrerie O, Decocq G, Kort HD,

Schrijver AD, Diekmann M, Eriksson O, et al. 2013. Latitudinal gradients as

443 natural laboratories to infer species' responses to temperature. J Ecol.

444 101(3):784–795. doi:https://doi.org/10.1111/1365-2745.12074.

445 Gorden NLS, Winkler KJ, Jahnke MR, Marshall E, Horky J, Hudelson C, Etterson JR.

446 2016. Geographic patterns of seed mass are associated with climate factors, but

447 relationships vary between species. Am J Bot. 103(1):60–72.

448 doi:https://doi.org/10.3732/ajb.1500295.

449 Greene DF, Johnson EA. 1993. Seed Mass and Dispersal Capacity in Wind-Dispersed

450 Diaspores. Oikos. 67(1):69–74. doi:10.2307/3545096.

451 Harper JL, Lovell PH, Moore KG. 1970. The Shapes and Sizes of Seeds. Annu Rev

452 Ecol Syst. 1(1):327–356. doi:10.1146/annurev.es.01.110170.001551.

453 Hulshof CM, Violle C, Spasojevic MJ, McGill B, Damschen E, Harrison S, Enquist BJ.

454 2013. Intra-specific and inter-specific variation in specific leaf area reveal the

455 importance of abiotic and biotic drivers of species diversity across elevation and

456 latitude. J Veg Sci. 24(5):921–931. doi:10.1111/jvs.12041.

457 Kattge J, Díaz S, Lavorel S, Prentice IC, Leadley P, Bönisch G, Garnier E, Westoby M,

458 Reich PB, Wright IJ, et al. 2011. TRY – a global database of plant traits. Glob

459 Change Biol. 17(9):2905–2935. doi:https://doi.org/10.1111/j.1365-

460 2486.2011.02451.x.

461 Kursa MB, Rudnicki WR. 2010. Feature Selection with the Boruta Package. J Stat

462 Softw. 36(11). doi:10.18637/jss.v036.i11. [accessed 2023 May 8].

463 http://www.jstatsoft.org/v36/i11/.

Larson JE, Sheley RL, Hardegree SP, Doescher PS, James JJ. 2015. Seed and

seedling traits affecting critical life stage transitions and recruitment outcomes in

dryland grasses. J Appl Ecol. 52(1):199–209. doi:10.1111/1365-2664.12350.

Leger EA, Atwater DZ, James JJ. 2019. Seed and seedling traits have strong impacts

on establishment of a perennial bunchgrass in invaded semi-arid systems. J Appl

Ecol. 56(6):1343–1354. doi:10.1111/1365-2664.13367.

Leimu R, Fischer M. 2008. A Meta-Analysis of Local Adaptation in Plants. PLOS ONE.

3(12):e4010. doi:10.1371/journal.pone.0004010.

Leishman MR. 2001. Does the seed size/number trade-off model determine plant

community structure? An assessment of the model mechanisms and their

generality. Oikos. 93(2):294–302. doi:https://doi.org/10.1034/j.1600-

0706.2001.930212.x.

Liu J, Bai Y, Lamb EG, Simpson D, Liu G, Wei Y, Wang D, McKenney DW, Papadopol

P. 2013. Patterns of Cross-Continental Variation in Tree Seed Mass in the

Canadian Boreal Forest. PLOS ONE. 8(4):e61060.

doi:10.1371/journal.pone.0061060.

Liu U, Cossu TA, Dickie J. 2019 Jun 13. Royal Botanic Gardens, Kew's Seed

Information Database (SID): A compilation of taxon-based biological seed

characteristics or traits. Biodivers Inf Sci Stand. doi:10.3897/biss.3.37030.

[accessed 2023 May 1].

https://www.proquest.com/docview/2282401665/abstract/5E5AFC61B9AA47F5P

Q/1.

Malanson GP, Butler DR, Fagre DB, Walsh SJ, Tomback DF, Daniels LD, Resler LM,

Smith WK, Weiss DJ, Peterson DL, et al. 2007. Alpine treeline of Western North

America: linking organism-to-landscape dynamics. Phys Geogr. 28(5):378–396.

doi:10.2747/0272-3646.28.5.378.

McGinley MA, Temme DH, Geber MA. 1987. Parental investment in offspring in variable

environments: theoretical and empirical considerations. Am Nat. 130(3):370–398.

doi:10.1086/284716.

Mishler BD, Guralnick R, Soltis PS, Smith SA, Soltis DE, Barve N, Allen JM, Laffan SW.

2020. Spatial phylogenetics of the North American flora. J Syst Evol. 58(4):393–

405. doi:10.1111/jse.12590.

Moles AT, Ackerly DD, Tweddle JC, Dickie JB, Smith R, Leishman MR, Mayfield MM,

Pitman A, Wood JT, Westoby M. 2007. Global patterns in seed size. Glob Ecol

Biogeogr. 16(1):109–116. doi:https://doi.org/10.1111/j.1466-8238.2006.00259.x.

Moles AT, Westoby M. 2004. Seedling survival and seed size: a synthesis of the

literature. J Ecol. 92(3):372–383. doi:10.1111/j.0022-0477.2004.00884.x.

Moran EV, Hartig F, Bell DM. 2016. Intraspecific trait variation across scales:

implications for understanding global change responses. Glob Change Biol.

22(1):137–150. doi:10.1111/gcb.13000.

Muller-Landau HC. 2010. The tolerance–fecundity trade-off and the maintenance of

diversity in seed size. Proc Natl Acad Sci. 107(9):4242–4247.

doi:10.1073/pnas.0911637107.

Murray BR, Brown AHD, Dickman CR, Crowther MS. 2004. Geographical gradients in

seed mass in relation to climate. J Biogeogr. 31(3):379–388. doi:10.1046/j.0305-

0270.2003.00993.x.

National Academies of Sciences, Engineering, and Medicine. 2023. An assessment of

native seed needs and the capacity for their supply: final report. Washington,

D.C.: National Academies Press. [accessed 2024 Sep 26].

https://www.nap.edu/catalog/26618.

Norden N, Daws MI, Antoine C, Gonzalez MA, Garwood NC, Chave J. 2009. The

relationship between seed mass and mean time to germination for 1037 tree

species across five tropical forests. Funct Ecol. 23(1):203–210.

doi:https://doi.org/10.1111/j.1365-2435.2008.01477.x.

Paige KN, Whitham TG. 1987. Overcompensation in Response to Mammalian

Herbivory: The Advantage of Being Eaten. Am Nat. 129(3):407–416.

Parry M, Canziani O, Palutikof J, van der Linden P, Hanson C. 2007. Climate change

2007: impacts, adaptation and vulnerability. In: contribution of Working Group II

to the fourth assessment report of the Intergovernmental Panel on Climate

Change.

Pebesma E, Graeler B. 2023. gstat: Spatial and Spatio-Temporal Geostatistical

Modelling, Prediction and Simulation. [accessed 2024 Mar 11]. https://cran.r-

project.org/web/packages/gstat/index.html.

Seale M, Nakayama N. 2020. From passive to informed: mechanical mechanisms of

seed dispersal. New Phytol. 225(2):653–658. doi:10.1111/nph.16110.

Sgrò CM, Lowe AJ, Hoffmann AA. 2011. Building evolutionary resilience for conserving

biodiversity under climate change. Evol Appl. 4(2):326–337. doi:10.1111/j.1752-

4571.2010.00157.x.

Sheridan JA, Bickford D. 2011. Shrinking body size as an ecological response to

climate change. Nat Clim Change. 1(8):401–406. doi:10.1038/nclimate1259.

Siefert A, Violle C, Chalmandrier L, Albert CH, Taudiere A, Fajardo A, Aarssen LW,

Baraloto C, Carlucci MB, Cianciaruso MV, et al. 2015. A global meta-analysis of

the relative extent of intraspecific trait variation in plant communities. Ecol Lett.

18(12):1406–1419. doi:10.1111/ele.12508.

Smith CC, Fretwell SD. 1974. The optimal balance between size and number of

offspring. Am Nat. 108(962):499–506.

Solbreck C, Knape J. 2017. Seed production and predation in a changing climate: new

roles for resource and seed predator feedback? Ecology. 98(9):2301–2311.

doi:10.1002/ecy.1941.

Tangney R, Merritt DJ, Callow JN, Fontaine JB, Miller BP. 2020. Seed traits determine

species' responses to fire under varying soil heating scenarios. Funct Ecol.

34(9):1967–1978. doi:10.1111/1365-2435.13623.

Turnbull LA, Rees M, Crawley MJ. 1999. Seed mass and the competition/colonization

trade-off: a sowing experiment. J Ecol. 87(5):899–912.

Tuthill JE, Ortega YK, Pearson DE. 2023. Seed size, seed dispersal traits, and plant

dispersion patterns for native and introduced grassland plants. Plants.

12(5):1032. doi:10.3390/plants12051032.

Vaughton G, Ramsey M. 1997. Seed mass variation in the shrub banksia spinulosa

(proteaceae): resource constraints and pollen source effects. Int J Plant Sci.

158(4):424–431. doi:10.1086/297452.

Venable DL. 1992. Size-number trade-offs and the variation of seed size with plant

resource status. Am Nat. 140(2):287–304.

Violle C, Enquist BJ, McGill BJ, Jiang L, Albert CH, Hulshof C, Jung V, Messier J. 2012.

The return of the variance: intraspecific variability in community ecology. Trends

Ecol Evol. 27(4):244–252. doi:10.1016/j.tree.2011.11.014.

Völler E, Auge H, Prati D, Fischer M, Hemp A, Bossdorf O. 2012. Geographical and

land-use effects on seed-mass variation in common grassland plants. Basic Appl

Ecol. 13(5):395–404. doi:10.1016/j.baae.2012.06.006.

Westoby M, Jurado E, Leishman M. 1992. Comparative evolutionary ecology of seed

size. Trends Ecol Evol. 7(11):368–372. doi:10.1016/0169-5347(92)90006-W.

Westoby M, Leishman M, Lord J, Poorter H, Schoen DJ. 1996. Comparative ecology of

seed size and dispersal [and discussion]. Philos Trans Biol Sci. 351(1345):1309–

1318.

Zhang C, Li J, Willis CG, Ma Z. 2020. Among-population variation in seed mass for 190

Tibetan plant species: Phylogenetic pattern and ecological correlates. Glob Ecol

Conserv. 23:e01163. doi:10.1016/j.gecco.2020.e01163.

Zhang C, Ma Z, Zhou H, Zhao X. 2019. Long-term warming results in species-specific

shifts in seed mass in alpine communities. PeerJ. 7:e7416.

doi:10.7717/peerj.7416.

 Figure Legends

Figure 1. Seed mass of over 2,000 native plant species colored by growth form.

A) Collection locations of 13,004 filled seed lots from Seeds of Success. Each point is a

site sampled by a field crew after scouting and careful phenology tracking, representing

a collection of at least 10,000 seeds from at least 50 plants. Seed mass was log-

transformed for analysis, and sized by 20 quantile groups (5 examples shown at top).

Log transformed seed mass of 13,004 populations plotted along B) latitude and C) year

with 5 log breaks indicated on the y axis, lines represent simple linear regression of

seed mass and latitude or year on the 3 collection dataset (see Q2, Q3 and *SI Appendix*

Table S2, S4 for full generalized linear model and full spatial simultaneous

autoregressive error model results, partial effects est = 0.0266, p < 0.0001; est = -0.010,

p = 0.007 respectively). D) Log-transformed seed mass for 2093 species from 108 plant

families from a dataset of seed collections across 11 western US states. For each

species, the interquartile range of seed mass is shown by the lines and outliers are

indicated as points and are colored by growth form. The y axis is on a log 10 scale in

seed mass and species are ordered along the x axis by mean seed mass; the subset of

families represented in the 10 sample dataset are listed near the first time the largest

species appears in the distribution. *Salvia columbariae* is labeled as the median

species, for reference. Seed mass was generated from the US Forest Service's Bend

Seed Extractory (BSE) processing of the Bureau of Land Management's collections

(BSE weighs a subset of 500 seeds per collection) and as part of the Seeds of Success

program, which has been collecting native seeds since 2000.

Figure 2. Phylogenetic tree of 235 plant species from 44 families collected from at least 10 populations using phylogenetic tree from Mishler et al. (2020). Inner bar chart represents log-transformed mean seed mass of each species, outer bar chart represents log-transformed coefficient of variation of seed mass for each species. Seed mass is from the US Forest Service's Bend Seed Extractory processing the Bureau of Land Management's Seeds of Success collections. Seed mass coefficient of variation is calculated as $CV_{unbiased} = (1 + 14 \times N) \times CV$. Images of native seeds are included; species names are indicated with a nearby bullet (photos: LL).

Figure 3. Relationships between seed mass and environment for native plant

species from across the western US. Environmental influence on A) seed mass using a

spatial simultaneous autoregressive error model (N = 859 species, 10,958 populations,

73 plant families in the 3 collection dataset) and B) the variability of seed mass

estimated from a phylogenetic generalized least squares regression (N = of 235

species, 6865 total populations, in the 10 collection dataset). Covariates are centered

and scaled, and error bars represent standard error produced by the PGLS.

Relationships between soil pH and seed mass for C) *Balsamorhiza hookeri* and D)

Elymus elymoides; relationships between change in fall maximum temperatures 30

years prior to each collection and seed mass for E) *Erigeron pumilus* and F)

Hymenoclea salsola; relationships between annual minimum temperature and seed

mass for G) *Asclepias speciosa* and H) *Plantago ovata*; relationships between change

in spring maximum temperatures 30 years prior to each collection and seed mass for I)

Eriogonum heracleoides and J) *Pseudoroegneria spicata*; relationships between

cumulative precipitation six months leading up to collection and seed mass for K)

Heterotheca villosa and L) *Achnatherum thurberianum*; relationships between change in

annual precipitation 30 years prior to each collection and seed mass for M)

Hesperostipa comata and N) *Artemisia tridentata*. For C-N, relationships were analyzed

with a spatial simultaneous autoregressive error model (SAR) seed mass is log-

transformed and predictor variables are centered and scaled; species specific results

and full model structures are in *SI Appendix* Table S6. Lines represent the semi-partial

regression plots between a single predictor variable and seed mass, points are colored

by growth form.

Figure 4. Environmental and climatic influence on seed mass estimated from

105 individual spatial simultaneous autoregressive error models for species collected

from at least 20 different populations (5,010 total populations). Axes indicate individual

species models as rows and possible covariates as columns; gray boxes indicate

variables that did not enter final models. Red boxes indicate negative relationships,

grading to blue for positive relationships. Significance is indicated with *** = P <0.0001,

** = P<0.001, and * = P<0.05. Species' names are colored by growth form. Species and

variables are organized by a similarity matrix based on model results. Figure is halved

at *Penstemon acuminatus* continuing to *Stenotus acaulis* following the similarity matrix.