Title

Auxin-salicylic acid seesaw regulates the age-dependent balance between plant
growth and herbivore defense

Authors

Wen-Hao Han[1]†, Feng-Bin Zhang[1]†, Shun-Xia Ji[1], Kai-Lu Liang[1], Jun-Xia Wang[1],
Xiao-Ping Fan[2], Shu-Sheng Liu[1], Xiao-Wei Wang[1]*

Affiliations

[1]State Key Laboratory of Rice Biology and Breeding, Ministry of Agriculture Key
Lab of Molecular Biology of Crop Pathogens and Insects, Zhejiang Key Laboratory
of Biology and Ecological Regulation of Crop Pathogens and Insects, Institute of
Insect Sciences, Zhejiang University, Hangzhou 310058, China.
[2]LC-Bio Technology Co., Ltd, Hangzhou, China.
*Corresponding author. Email: xwwang@zju.edu.cn
†These authors contributed equally to this work.

Abstract

Plants must fine-tune their needs for growth and defense. According to the Plant Vigor
Hypothesis, younger, more vigorous plants tend to be more susceptible to herbivores
compared to older, mature plants, yet the molecular mechanisms underlying this
dynamic remain elusive. Here, we uncover a hormonal crosstalk framework that
orchestrates the age-related balance between plant growth and herbivore defense. We
demonstrate that the accumulation of salicylic acid (SA), synthesized by *Nicotiana*
benthamiana phenylalanine ammonia-lyase 6 (NbPAL6), dictates insect resistance in
adult plants. The expression of *NbPAL6* is driven by the key transcription factor,
NbMYB42, which is regulated by two interacting auxin response factors,
NbARF18La/b. In juvenile plants, higher auxin levels activate *Nb*miR160c, a
microRNA that silences *NbARF18La/b*, subsequently reducing *NbMYB42* expression,
lowering SA accumulation, and thus weakening herbivore defense. Furthermore,
excessive SA accumulation in juvenile plants enhances defense but antagonizes auxin
signaling, impairing early growth. Our findings suggest a seesaw-like model that
balances growth and herbivore defense depending on the plant's developmental stage.

Keywords

Plant age, Balance, Phytohormone cross-talk, Insect defense, Growth, MicroRNA,
Auxin, Salicylic acid

Teaser

The crosstalk between two phytohormones enables plants to achieve a balance
between growth and defense against insects across growth stages.

MAIN TEXT

Introduction

Plants, being sessile organisms, must constantly navigate hostile environments to survive and reproduce, facing a wide range of abiotic and biotic stresses. Over time, they have evolved sophisticated systems to detect, respond to, and defend against environmental changes. However, these defenses come at a significant cost, often resulting in trade-offs with growth and development (*1, 2*). Among the most damaging threats to plants are herbivorous insects, which can attack at any stage of plant development. While tissue-chewing insects cause direct damage by consuming plant tissues, phloem-feeding insects pierce the epidermis to delicately extract nutrient-rich phloem sap using needle-like mouthparts. In response to such threats, plants have evolved both inducible and constitutive defense mechanisms (*3, 4*). Induced defenses are triggered in response to specific stimuli, while constitutive defenses involve continuous accumulation of insecticidal components throughout the plant's growth and development (*5, 6*). Given that insect attacks are often sporadic, plants must finely balance their investment in growth with the need for defense. However, how this balance is maintained at the molecular level remains poorly understood.

In the context of plant-herbivore interactions, the Plant Vigor Hypothesis (PVH) posits that herbivores preferentially target young, vigorous plants over older, mature ones (*7, 8*). At the same time, plants tend to develop increased resistance to biological stresses, such as pathogens and herbivores, as they age—a phenomenon known as age-related resistance (ARR) (*9–13*). These observations suggest that plants may prioritize rapid growth early in life, tolerating some degree of herbivore damage, before shifting their focus to defense mechanisms later in development (*14*). However, the precise mechanisms underlying PVH and ARR in the context of herbivores remain largely unexplored. Given the dual necessity for plants to grow and defend themselves, a better understanding of how this balance is optimized holds significance to both ecological theory and agricultural production.

Phytohormones play a pivotal role in regulating various aspects of plant growth, development, and defense, often through intricate crosstalk mechanisms (*15, 16*). Auxin, the first phytohormone discovered, along with other phytohormones, orchestrates plant growth and development while also modulating stress responses (*15*). Among stress-related hormones, jasmonic acid (JA), salicylic acid (SA), abscisic acid (ABA), and ethylene (ET) are key players, particularly in response to herbivory, with JA being central to resistance against tissue-chewing insects and SA being crucial for defense against phloem-feeding insects (*14, 16*). The interaction between these phytohormones has complex effects on plants against insects (*16*). For instance, the JA-SA antagonistic relationship is a well-studied area, with each hormone modulating plant responses to different herbivores. Moreover, other phytohormones,

including auxin, ET, ABA, gibberellins (GA), brassinosteroids (BR), and cytokinins (CK) are also involved in fine-tuning plant immunity by influencing   JA and SA signaling pathways, especially in response to pathogens and abiotic stresses (*16*). The intricate mechanisms of phytohormonal interplay involve multiple layers of regulation, including transcriptional control by transcription factors (TFs), post-transcriptional regulation by small RNAs (sRNAs), and control of protein activity (*17–20*). Despite these advances, it remains unclear how plants integrate hormonal signals to balance growth and defense at different developmental stages.

The whitefly, *Bemisia tabaci* (Gennadius), is a species complex, several species of which are of agricultural importance worldwide (*21–24*). A few species of this whitefly complex are highly polyphagous, exhibiting remarkable adaptability to different hosts (*25, 26*). Being phloem-feeders, whiteflies use their piercing-sucking mouthparts delicately to extract nutrients, making juvenile, vigorous plants particularly vulnerable to attacks and yield losses (*26–28*). Understanding the underlying mechanisms that govern plant growth-defense balance could be key to developing sustainable whitefly control strategies.

Nicotiana benthamiana has become an important model organism in plant molecular biology and biotechnology, serving as a bio-factory for vaccines and pharmacological compounds (*29, 30*). Our previous studies have shown that *N. benthamiana* exhibits strong attraction to and lethal effects on several Hemiptera and Thysanoptera insects, including whiteflies, winged aphids, and thrips, highlighting its potential as a dead-end trap plant for field pest control (*31–33*). However, we observed that the insecticidal efficacy of *N. benthamiana* varied with the plant's development stage. Upon further investigation, we found that beyond environmental influences, this age-related defense capacity was closely tied to the developmental stage, suggesting a complex regulatory mechanism involving ARR. These findings underscore the utility of *N. benthamiana* as a system to study ARR and provide a foundation for its application in pest management.

In this study, we demonstrate that auxin-SA crosstalk mediates the balance between plant growth and herbivore defense at different developmental stages. We reveal that the establishment of ARR in *N. benthamiana* and several other Solanaceae species is closely tied to SA levels. Using molecular biology and biochemical approaches, we elucidate that the NbARF18La/b-NbMYB42-NbPAL6 signaling module plays a critical role in the age-related regulation of SA accumulation. Furthermore, we find that the age-dependent auxin-miRNA160c module also contributes to this process. Our findings provide new insights into PVH and lay the theoretical groundwork for developing innovative pest control strategies against whiteflies.

Results

Age-regulated resistance against phloem-feeding insects in Solanaceae plants

To compare the defense levels of plants at different ages against phloem-feeding insects, we conducted bioassays of *B. tabaci* whiteflies, *Myzus persicae* aphids, and *Frankliniella occidentalis* western flower thrips on several Solanaceae species, including *N. benthamiana*, *Solanum lycopersicum* (tomato), and *Nicotiana tabacum* (tobacco). In *N. benthamiana*, both the survival rate and fecundity of whiteflies were significantly lower on adult plants (35 days old, with 8–9 true leaves) compared to juvenile plants (21 days old, with 3–4 true leaves) **(Fig. 1, A and B)**. Similarly, whiteflies performed better on juvenile tobacco (21 days old, 3–4 true leaves) and tomato (21 days old, 3–4 true leaves) plants compared to adult plants (35 days old, 6–7 true leaves for tobacco; 8–9 true leaves for tomato) **(Fig. 1, A and B)**. Additionally, western flower thrips showed higher survival rates, and aphids produced more offspring on juvenile plants compared to adult plants **(Fig. 1, C and D)**. These results demonstrate ARR against phloem-feeding insects in Solanaceae plants.

SA-dependent ARR against phloem-feeding insects

To explore the mechanism underlying ARR, we focused on *N. benthamiana,* which exhibited the most pronounced age-dependent insect resistance. We examined whether the defense phytohormones SA and JA were involved in ARR by comparing the basal levels of SA and JA in 21-day-old and 35-day-old *N. benthamiana* plants. SA levels were higher in adult plants, while JA levels showed no significant differences between juvenile and adult plants **(Fig. 1, E and F)**. This trend was also observed in tobacco **(Fig. 1, E and F)**. Additionally, the expression of *pathogenesis-related 1* (*NbPR1*), a downstream marker of SA signaling, was significantly upregulated in adult plants, indicating enhanced SA signaling **(Fig. S1A)**.

To further investigate the role of SA in ARR, we used *N. benthamiana* transgenic plants expressing the *NahG* gene, which encodes salicylate hydroxylase from *Pseudomonas putida*, an enzyme that depletes SA accumulation (*34*). Unlike wild-type plants, *NahG* plants were unable to accumulate sufficient SA as they aged **(Fig. 1G),** and the mRNA levels of *NbPR1* were also significantly reduced in 35-day-old plants **(Fig. S1B)**. Moreover, whitefly infestation failed to induce SA accumulation in *NahG* plants **(Fig. 1H)**. Bioassays revealed that whiteflies performed better on both juvenile and adult NahG plants compared to wild-type plants **(Fig. 1I)**, indicating that both whitefly-inducible and growth-accumulated SA are critical for plant defense against whiteflies. Exogenous SA application further confirmed SA's role in ARR. Treatment with SA increased *NbPR1* expression and enhanced whitefly defense in 21-day-old wild-type plants **(Fig. 1J and Fig. S1C)**, as well as in 35-day-old *NahG* plants **(Fig. 1K)**. These results collectively confirm that ARR in *N. benthamiana* against whiteflies is SA-dependent.

SA-mediated ARR depends on the expression of the SA synthesis gene *NbPAL6*

In plants, SA is synthesized via two distinct pathways: the phenylalanine ammonia lyase (PAL) pathway and the isochorismate synthase (ICS) pathway (*35*) **(Fig. 2A)**. To assess whether gene expression in these pathways changes with plant age, we performed reverse transcription quantitative PCR (RT-qPCR). The results showed no significant differences in the transcript levels of *NbICS1* and *NbICS2*, or the the two TFs, *CaM-Binding Protein 60g* (*CBP60g*) and *Systemic Acquired Resistance Deficient 1* (*SARD1*), which activate the *ICS* pathway **(Fig. 2B)**. Similarly, components of the ICS pathway such as *Enhanced Disease Susceptibility 5* (*EDS5*) and *AvrPphB Susceptible 3* (*PBS3*) were downregulated with plant age, while *Enhanced Pseudomonas Susceptibility 1* (*EPS1*) remained unchanged **(Fig. 2B)**. Given these findings, we shifted our focus to the PAL pathway.

A search of the *N. benthamiana* genome database (https://www.nbenth.com/) identified nine *PAL* genes (*NbPAL1-NbPAL9*), which we classified into four phylogenetic branches, likely representing alleles from the allotetraploid ancestry of *N. benthamiana* **(Fig. S2 and Table S1)**. From these, we selected four *PAL* genes (*NbPAL4*, *NbPAL6*, *NbPAL7*, and *NbPAL9*) from the four branches respectively for further analysis. Expression profiling across developmental stages revealed that only *NbPAL6* showed increased transcription with plant age, while the other genes remained unchanged **(Fig. 2C)**.

To determine whether *NbPAL6* is involved in SA-mediated ARR, we employed virus-induced gene silencing (VIGS) and *Agrobacterium*-mediated overexpression to manipulate *NbPAL6* expression in *N. benthamiana* (*29*, *36*). The timing of these treatments was carefully designed to align with plant age requirements **(Fig. 2D)**. In the VIGS system, we injected the pTRV2 vector into 21-day-old plants., which led to effective gene silencing within 14 days **(Fig. 2D)**. This period of transcriptional inhibition coincides with the establishment of ARR, making VIGS an ideal system for studying gene-related ARR **(Fig. 2D)**. Compared to control plants (injected with an empty vector), 35-day-old VIGS-treated plants showed a significant reduction in target gene transcript accumulation after reaching 21 days of age. Subsequent validation experiments on the 35-day VIGS-treated plants provide valuable insights into the role of target genes in ARR establishment. Meanwhile, *Agrobacterium*-mediated overexpression can be used to enhance the mRNA level of target genes in both 21-day-old and 35-day-old plants, enabling examination of gene function in plants of different ages **(Fig. 2D)**. We injected 19-day-old plants with the pCAMBIA1305-GFP vector containing the target gene. Two days later, during the vector's active period, elevated target gene mRNA levels were observed in the 21-day-old plant. By comparing overexpressed plants with control plants (injected with an empty vector), we could evaluate the effect of target genes on resistance in 21-day-old

plants **(Fig. 2D)**. Similarly, this method can also verify the effect of target gene overexpression in 35-day-old plants **(Fig. 2D)**.

Our results showed that 35-day-old VIGS-*NbPAL6* plants could not accumulate SA to the same levels as control (VIGS-EV) plants, nor could they upregulate *NbPR1* expression **(Fig. 2, E and F; Fig. S3A)**. Moreover, these plants exhibited reduced resistance to whiteflies **(Fig. 2G)**. Conversely, overexpression of *NbPAL6* at 21 days of age significantly increased the *NbPR1* levels and improved plant defense against whiteflies **(Fig. 2H; Fig. S3, B and C)**. These findings indicate that the differential expression of *NbPAL6* between juvenile and adult stages regulates SA level and SA-mediated ARR against whiteflies in *N. benthamiana*.

Age-regulated NbMYB42 activates *NbPAL6* transcription

To identify the upstream TFs regulating *NbPAL6* expression, we screened the *NbPAL6* promoter (*NbPAL6pro*) using a yeast one-hybrid (Y1H) system with a cDNA library from *N. benthamiana*. NbMYB42, a TF belonging to the R2R3-MYB family, was identified. R2R3-MYB is known to regulate the phenylpropanoid metabolic pathway (*37–39*). Yeast transformants containing the plasmids pAbAi-*NbPAL6pro* and pGADT7-NbMYB42 were able to grow on SD/-Leu medium with 150 ng/ml AbA within 3 days, while control transformants with pAbAi-*NbPAL6pro* and pGADT7 did not **(Fig. 3A)**. This indicates that NbMYB42 can bind to the *NbPAL6* promoter. NbMYB42 was also found to localize in the nucleus **(Fig. 3B)**, suggesting its role as a nuclear TF. To assess its role in activating *NbPAL6* expression *in vivo*, we performed the GUS staining assays. The results showed that *NbPAL6pro*::*GUS* co-infiltrated with *35S*::*NbMYB42* exhibited significantly higher GUS staining levels compared to the empty vector control in *N. benthamiana* leaves **(Fig. 3C)**. Similarly, dual-luciferase reporter assays demonstrated increased luciferase activity when *NbPAL6pro::LUC* was co-expressed with *35S::NbMYB42* **(Fig. 3D)**. These results confirmed that NbMYB42 activates *NbPAL6* promoter expression.

Next, we explored whether NbMYB42 directly binds to specific sites within the *NbPAL6pro*. Using the Jasper database (https://jaspar.elixir.no/), we identified five potential *MYB* binding sites (P1–P5) **(Fig. 3E)**. To validate binding *in vivo*, we conducted chromatin immunoprecipitation (ChIP)-qPCR using plants overexpressing *NbMYB42* with glutathione S-transferase (GST) tag. We designed primers specific to the five binding sites to assess their interaction with NbMYB42 **(Table S2)**. The experiments were carried out using an anti-GFP antibody. Binding activity was observed at P1, P3, P4, and P5, but not P2, in the immunoprecipitated chromatin, indicating significantly accumulation of NbMYB42-GFP at these sites compared to the GFP control **(Fig. 3F)**. Furthermore, electrophoretic mobility shift assay (EMSA) with recombinant NbMYB42-GST, purified from *Escherichia coli*, confirmed the binding activity *in vitro*. The MYB element site P3, which showed the strongest

binding ability in the ChIP-qPCR test, was selected to prepare the biotin-labeled
probe according to its sequence (Table S2). We observed that the NbMYB42-GST
protein was able to bind directly to the P3 probe, while the mutated probe eliminated
NbMYB42-GST binding (Fig. 3G). Additionally, an unlabeled P3 probe was prepared
as the cold competitors. With the increase of competitor probes (10, 20, and 50 ×), the
shifted band corresponding to the P3-probe and the NbMYB42 complex gradually
disappeared (Fig. 3G), suggesting the binding capacity was greatly reduced by the
addition of unlabeled cold competition probe. These results collectively show that
NbMYB42 specifically binds to the MYB *cis*-element on the *NbPAL6* promoter.
252
We then studied whether NbMYB42 influences ARR against whiteflies by regulating
NbPAL6-mediated SA accumulation. Overexpressing *NbMYB42* in 21-day-old *N.*
benthamiana plants increased *NbPAL6* expression more than 150-fold compared to
control plants (Fig. 3H and Fig. S4A). The SA-downstream *NbPR1* was also
significantly upregulated (Fig. S4B). This overexpression enhanced whitefly
resistance in 21-day-old plants (Fig. 3I). Conversely, silencing *NbMYB42* in 35-day-
old plants (Fig. S4C) suppressed *NbPAL6* expression (Fig. 3J), resulting in reduced
SA and *NbPR1* levels (Fig 3. K and L). Notably, the *NbMYB42*-silenced 35-day-old
N. benthamiana plant showed significantly reduced whitefly defense compared to
control (Fig. 3M). These results indicate that NbMYB42 plays a crucial role in ARR
in *N. benthamiana* by regulating *NbPAL6* expression and SA accumulation.
264
Two interacting age-regulated NbARFs activate the expression of *NbMYB42*
We observed that *NbMYB42* expression was significantly higher in 35-day-old plants
compared to 21-day-old plants (Fig. 4A), prompting further investigation into how
NbMYB42 expression is regulated during *N. benthamiana* development. Using a Y1H
assay, we screened for TFs that regulate *NbMYB42* expression and identified two
auxin response factors (ARFs) with distinct evolutionary relationships, which we
named NbARF18La and NbARF18Lb (Fig. S5A). Yeast strains harboring the pAbAi-
NbMYB42pro and pGADT7-NbARF18La or pGADT7-NbARF18Lb constructs were
able to grow on SD/-Leu medium supplemented 150 ng/ml AbA, whereas yeast
strains containing pAbAi-*NbMYB42pro* and a control pGADT7 plasmid could not
(Fig. 4B), suggesting that both NbARF18La and NbARF18Lb bind to the *NbMYB42*
promoter.
277
Phylogenetic analysis showed that NbARF18La and NbARF18Lb are closely related
to ARF16 from *Arabidopsis thaliana* (Fig. S5A). Amino acid sequence analysis
revealed conserved B3-type DNA-binding domains in the N-termini of both
NbARF18La and NbARF18Lb and dimerization domains in their C- termini (*40–43*)
(Fig. S5B). Subcellular localization analysis confirmed that both NbARFs are
localized in the nucleus (Fig. 4C), indicating their function as nuclear TFs. To verify
the activation of *NbMYB42* by these NbARFs, we conducted GUS staining and dual-

luciferase reporter assays. Co-expression of *35S::NbARF18La* and *35S::NbARF18Lb*
with the reporter *NbMYB42pro::GUS* in *N. benthamiana* leaves significantly
increased *GUS* expression **(Fig. 4D)**, and dual-luciferase reporter assay showed a
significant up-regulation in LUC/REN ratio when *35S::NbARF18La* and
35S::NbARF18Lb were co-expressed with the *NbMYB42pro::LUC* reporter **(Fig. 4E)**.
These results collectively demonstrate that NbARF18La and NbARF18Lb activate
NbMYB42 expression.
292
DNA-binding specificity of NbARF18La and NbARF18Lb
To explore how the two NbARFs specifically bind to the *NbMYB42* promoter, we
screened for potential auxin response elements (AuxRE) using the JASPAR database
and identified three regions (P1, P2, and P3) in the promoter **(Fig. 4F)**. We designed
primers specific to these regions **(Table S2)** and performed ChIP-qPCR using GFP-
tagged NbARF18La/b and anti-GFP antibodies. The results revealed a significant
enrichment of the P2 region, but not P1 and P3, in the immunoprecipitated chromatin,
indicating that *NbARF18La/b* bind to the P2 AuxRE region (TGTCTC) in the
NbMYB42 promoter **(Fig. 4F)**.
302
To confirm the binding specificity, we conducted an EMSA assay. The recombinant
NbARF18La-His-SUMO fusion protein bound to the P2 probe of the NbMYB42
promoter, while no binding occurred with a mutant probe (TGTCTC to AAAAAA)
(Fig. 4G). Additionally, competition assays using unlabeled probes verified the
specificity of binding **(Fig. 4G)**. Similarly, the NbARF18Lb-MBP fusion protein
bound to the same promoter region (P2), but not to the mutant probe, and competition
with unlabeled probes also reduced the binding strength **(Fig. 4H)**. Together, these
results indicate that NbARF18La and NbARF18Lb specifically bind to the same
AuxRE site in the *NbMYB42* promoter.
312
NbARF18La and NbARF18Lb interaction and self-association
Previous studies have shown that ARF TFs can form complexes through self-
interactions or by interacting with other TFs (*41–44*). To explore potential interactions
between *NbARF18La* and *NbARF18Lb* or self-interactions of either of the TFs, we
performed yeast two-hybrid (Y2H) assays. Yeast cells expressing NbARF18La-
AD/NbARF18Lb-BD grew on selective selective media (SD/-Leu-Trp-His-Ade),
indicating that NbARF18La and NbARF18Lb can interact with each other **(Fig. S6A)**.
Additionally, NbARF18La was able to interact with itself in Y2H assays **(Fig. S6A)**.
321
322 We then confirmed these interactions *in vivo* using a bimolecular fluorescence
323 complementation (BiFC) assay. GFP fluorescence was detected in *N. benthamiana*
324 leaves co-expressing NbARF18La and NbARF18Lb or NbARF18La alone **(Fig.
325 S6B)**. *In vitro* pull-down assays further validated these interactions, as His-
326 NbARF18La exhibited an affinity for MBP-NbARF18Lb, but no interaction was

observed in the negative control **(Fig. S6C)**. These results demonstrate that NbARF18La and NbARF18Lb physically interact and NbARF18La can self-interact, suggesting that these interactions are important for cooperatively regulating *NbMYB42* expression.

NbARFs regulate plant resistance to whiteflies in an age-related manner

Next, we examined the role of NbARF18La and NbARF18Lb in ARR to whiteflies in *N. benthamiana*. Overexpression of both NbARFs significantly upregulated expression of *NbMYB42*, *NbPAL6*, and the downstream *NbPR1* gene in 21-days-old plants **(Fig. 5A; Fig. S7A to C)**, which led to enhanced resistance to whiteflies **(Fig. 5B)**. In contrast, silencing *NbARF* genes via VIGS in 35-day-old *N. benthamiana* plants resulted in substantially reduced expression of *NbMYB42* and *NbPAL6* **(Fig. 5C; Fig. S7D and E)**, as well as significantly decreased endogenous SA levels and reduced *NbPR1* expression **(Fig. 5D; Fig. S7F)**. Correspondingly, silencing *NbARFs* impaired the defense response against whiteflies in adult plants **(Fig. 5E)**.

To further investigate how NbARFs regulate *NbPAL6* expression, we co-expressed NbARF18La/b with a LUC reporter fused to the *NbPAL6* promoter, which significantly enhanced LUC activity **(Fig. 5F)**, indicating that NbARF18La/b regulate *NbPAL6* by modulating the expression of *NbMYB42*. Furthermore, expression of *NbARF18La/b* was significantly higher in 35-day-old plants than in 21-day-old plants **(Fig. 5G)**, indicating that the NbARF18La/b-*NbMYB42*-*NbPAL6* module is regulated in an age-dependent manner.

Auxin suppresses plant defense by repressing SA accumulation in juvenile plants

Given the findings above, we questioned why SA-mediated defense is not fully established during the juvenile stage of plants. Since defense-related ARF TFs are key components downstream of the auxin response, we postulated that auxin is involved in this process. To test this hypothesis, we first measured the levels of indole-3-acetic acid (IAA) in 21-day-old and 35-day-old plants, and found that IAA levels were significantly higher in juvenile plants **(Fig. 6A)**. In parallel, we observed elevated expression levels of the auxin synthesis genes, *TRYPTOPHAN AMINOTRANSFERASE RELATED 2* (*NbTAR2*) and *YUCCA 8* (*NbYUC8*), as well as the auxin-responsive gene *NbIAA29*, in juvenile plants **(Fig. 6B)**.

To assess the effect of auxin on plant defense, we treated 35-day-old plants, which are known for their high resistance to whiteflies, with natural IAA and the auxin-like plant growth regulator, 1-naphthalene acetic acid (NAA). Bioassays revealed that the whiteflies performed better on IAA- and NAA-treated plants, indicating that auxin negatively impacts plant defense **(Fig. 6, C and D)**. Furthermore, IAA treatment significantly suppressed both the SA level **(Fig. 6E)** and the expression of the SA-responsive gene *NbPR1* **(Fig. 6F)** in the 35-day-old plants.

To further clarify the role of auxin in repressing SA-based resistance, we treated 21-
day-old plants with two auxin biosynthesis inhibitors: ʟ-kynurenine (ʟ-Kyn), a
competitive substrate inhibitor, and Yucasin (5-(4-chlorophenyl)-4H-1,2,4-triazole-3-
thiol), an effective IAA biosynthesis inhibitor (45). Plants treated with these auxin
inhibitors showed increased resistance to whiteflies **(Fig. S8, A and B)**, as well as
higher expression levels of *NbPR1* **(Fig. S8C)**. These results confirm that auxin
suppresses SA biosynthesis and downstream signaling pathways in juvenile *N.*
benthamiana plants, thereby repressing SA-mediated defense against whiteflies.

Auxin suppresses *NbPAL6* and *NbMYB42* expression by repressing NbARF levels
To explore how auxin suppresses SA accumulation, we focused on the age-regulated
SA synthesis gene *NbPAL6*. Auxin treatment significantly reduced *NbPAL6* expression
in 35-day-old plants **(Fig. 6G)**, whereas treatment with an auxin inhibitor enhanced
NbPAL6 expression in 21-day-old plants **(Fig. S8D)**. To investigate the underlying
mechanism, we used *NbPAL6* promoter-driven GUS and LUC reporter, which showed
decreased transcript levels upon auxin treatment, indicating that auxin suppresses
NbPAL6 expression through transcriptional repression **(Fig. 6H)**.

We hypothesized that auxin may inhibit *NbPAL6* transcription by suppressing the
activity of the TF *NbMYB42*. However, *NbPAL6* promoter-linked GUS and LUC
reporter assays in IAA-treated plants revealed no significant impact on reporter
expression **(Fig. 6I)**, suggesting that auxin does not affect the transactivation ability of
NbMYB42.

Next, we assessed whether auxin affects *NbMYB42* mRNA levels. qRT-PCR revealed
a significant reduction in *NbMYB42* mRNA levels in auxin-treated 35-day-old plants
(Fig. 6J), whereas auxin inhibitor treatment enhanced *NbMYB42* expression in 21-
day-old plants **(Fig. S8E)**. A reporter-GUS assay further confirmed that auxin
represses *NbMYB42* transcription **(Fig. 6K)**. These results indicate that auxin
represses *NbMYB42* transcription, thereby regulating downstream *NbPAL6*
expression.

To understand how auxin represses *NbMYB42* transcription, we examined upstream
TFs, *NbARF18La* and *NbARF18Lb*. Surprisingly, auxin treatment did not affect the
activation of *NbMYB42* by NbARF18La/b **(Fig. 6, L and M)**. However, auxin
significantly decreased the transcript levels of *NbARF18La/b* in 35-day-old plants
(Fig. 6N), with auxin inhibitors increasing their expression in 21-day-old plants **(Fig.**
S8F). These results indicate that auxin downregulates *NbARF18La/b*, which in turn
suppresses *NbMYB42*. Further, we examined whether auxin repressed *NbARFs*
through transcriptional inhibition. Importantly, using promoter-GUS reporters, we
found that auxin did not reduce *NbARF18La/b* mRNA levels through transcriptional

repression **(Fig. 6O)**. Therefore, it appears that auxin suppresses *NbARF* mRNA
levels through a post-transcriptional regulatory mechanism.
413

Auxin-inducible miRNA *Nb*miR160c targets *NbARF* mRNA
We explored the mechanisms by which auxin suppresses the transcript levels of
NbARF18La and *NbARF18Lb* via post-transcriptional regulation. Previous studies
have shown that plant miRNAs can negatively regulate the transcripts of *ARF* genes
(*46*, *47*), and that auxin treatment can enhance miRNA accumulation in plants (*48*).
Therefore, we hypothesized that auxin-inducible, age-related miRNAs may be
responsible for the post-transcriptional regulation of these *ARF* genes. To test this
hypothesis, we analyzed miRNA sequencing data from 21-day-old *N. benthamiana*
plants and, through target prediction analysis, identified a miRNA, *Nb*miR160c,
which potentially targets the transcripts of *NbARF18La* and *NbARF18Lb*. Target site
analysis revealed that *Nb*miR160c specifically binds to the coding regions of
NbARF18La (1572 bp to 1590 bp) and *NbARF18Lb* (2099 bp to 2107 bp) **(Fig. 7A,**
B). To validate this interaction, we artificially overexpressed *Nb*miR160c in 35-day-
old plants. This overexpression significantly reduced the mRNA levels of
NbARF18La and *NbARF18Lb* **(Fig. 7C; Fig. S9, A and B)**, suggesting direct
regulation of these transcripts by *Nb*miR160c.
430

Next, to pinpoint the exact cleavage sites of *NbARF18La* and *NbARF18Lb* by
*Nb*miR160c in juvenile plants, we conducted degradome sequencing on samples from
21-day-old plants. The analysis confirmed that *Nb*miR160c cleaves *NbARF18La* and
NbARF18Lb transcripts at the predicted target sites, as shown by the accumulation of
cleavage products at these regions **(Fig. 7, D and E)**. The red dots in **Fig. 7, D and E**
highlight the cleavage sites, aligning with the *Nb*miR160c target predictions.
437

To further confirm the regulatory role of *Nb*miR160c, we cloned the target sites of
NbARF18La and *NbARF18Lb* and fused them with a GFP tag. When these constructs
were transiently expressed in *N. benthamiana* leaves, co-expression with *Nb*miR160c
resulted in the suppression of GFP fluorescence **(Fig. 7, F and G; Fig. S9, C and D)**.
However, when the *Nb*miR160c target sites were mutated, GFP expression was
unaffected, even in the presence of *Nb*miR160c **(Fig. 7, F and G; Fig. S9, C and D)**.
Additionally, in co-expression assays with GUS-tagged target sites, *Nb*miR160c
suppressed the expression of wild-type GUS-tagged *NbARF18La* and *NbARF18Lb*
target sites, while the mutant target sites were unaffected **(Fig. 7H)**. We observed
similar results using LUC-tagged reporter constructs, where *Nb*miR160c suppressed
the activity of LUC reporter fused to the wild-type target sites of *NbARF18La* and
NbARF18Lb, but not the mutant constructs **(Fig. 7I)**. Collectively, these results
confirm that *Nb*miR160c directly targets the coding regions of *NbARF18La* and
NbARF18Lb, leading to their post-transcription repression. This auxin-inducable

miRNA thereby plays a crucial role in modulating *ARF* expression in *N. benthamiana*,
particularly during the juvenile stage.
454
NbmiR160c regulates the expression of *NbMYB42* and *NbPAL6*
To explore the role of *Nb*miR160c in regulating the NbARF-NbMYB42 module-
mediated SA defense against whiteflies, we first overexpressed *Nb*miR160c in 35-
day-old plants. The overexpression resulted in a significant reduction in the mRNA
levels of *NbMYB42* and *NbPAL6* **(Fig. 8A)**, as well as in SA content and downstream
NbPR1 expression **(Fig. 8B and Fig. S9E)**. Notably, the *Nb*miR160c-expressing 35-
day-old plants exhibited reduced defense against whiteflies **(Fig. 8C)**, indicating that
*Nb*miR160c negatively regulates plant anti-insect defense.
463
To validate the role of *Nb*miR160c in repressing *NbARF18La/b* and downstream SA
regulatory pathways in juvenile plants, we overexpressed an artificial target mimic of
*Nb*miR160c (mimic160c) in 21-day-old plants to inhibit its function **(Fig. S9F)**. As
expected, *NbARF18La* and *NbARF18Lb* transcript levels were significantly
upregulated in the treated plants **(Fig. 8D)**. Moreover, the *NbMYB42* and *NbPAL6*
mRNA levels **(Fig. 8E)**, SA levels **(Fig. S9G),** and *NbPR1* expression were also
significantly elevated in the mimic160c-overexpressing plants **(Fig. S9H)**.
Correspondingly, bioassays showed that whiteflies performed poorly on the
mimic160c-overexpressing plants, indicating enhanced defense **(Fig. 8F)**.
473
We then examined the effect of *Nb*miR160c on *NbARF*-promoted *NbMYB42*
transcription in plants at juvenile and adult stages. In 35-day-old plants,
overexpression of *Nb*miR160c repressed the activity of the *NbMYB42*pro-*LUC*
reporter **(Fig. S10A)**, whereas overexpression of mimic160c enhanced *NbMYB42*pro-
LUC reporter expression in 21-day-old plants **(Fig. S10B)**. When *Nb*miR160c was co-
expressed with *NbARF18La*, *NbARF18Lb*, or both, *NbMYB42pro* reporter expression
was significantly inhibited in 35-day-old plants **(Fig. 8G)**. These results demonstrate
that *Nb*miR160c represses *NbARF*-activated *NbMYB42* transcription. Similarly, we
confirmed that *Nb*miR160c is involved in repressing *NbPAL6* transcription in a
similar manner **(Fig. S10 C to E)**. Collectively, these findings show that *Nb*miR160c
negatively regulates *NbARF18La* and *NbARF18Lb*, thereby affecting the transcription
of *NbMYB42* and its downstream target *NbPAL6*.
486
Auxin promotes the *NbmiR160c* level by enhancing its expression
Next, we investigated the relationship between the *Nb*miR160c expression, plant age,
and phytohormones. We observed that *Nb*miR160c levels were significantly higher in
21-day-old plants compared to 35-day-old plants **(Fig. 8H)**. Given that auxin content
is typically higher in juvenile plants, we speculated that auxin might promote
*Nb*miR160c expression. To test this hypothesis, we treated 35-day-old plants, which
had naturally lower *Nb*miR160c levels, with IAA. The treatment significantly

increased *Nb*miR160c levels **(Fig. 8I)**. Additionally, we cloned the promoter of the

*Nb*miR160c precursor from the *N. benthamiana* genome and fused it to GUS and

LUC reporters. Upon IAA treatment, we observed enhanced reporter expression **(Fig.**

8, J and K), confirming that auxin activates *Nb*miR160c transcription. These results

indicate that auxin promotes *Nb*miR160c expression, thereby repressing the NbARF-

NbMYB42-NbPAL6 module, which mediates whitefly defense in juvenile *N.*

benthamiana plants.

Excessive SA impacts early plant growth by antagonizing auxin

We further examined the physiological implications of auxin-mediated suppression of

SA in juvenile plants. To assess the impact of excessive SA on early plant growth, we

treated 21-day-old plants with exogenous SA. These treated plants showed stunted

growth **(Fig. 9A)** and, by the time they reached 35 days of age, had significantly

reduced weight and height compared to untreated plants **(Fig. 9, B and C)**. These

results, along with our earlier findings, suggest that while excessive SA enhances

whitefly defense in juvenile plants, it also negatively impacts plant early growth.

Since auxin and SA pathways are known to antagonize each other, we examined

whether SA inhibited early growth by interfering with the auxin pathway. Our results

show that SA treatment significantly reduced the expression of the auxin synthesis

genes *NbTAR2* and *NbYUC8* **(Fig. 9D)**, lowered auxin levels **(Fig. 9E)**, and decreased

downstream *NbIAA29* expression **(Fig. 9F)** in juvenile plants. Conversely, the NahG

plants, which have reduced SA levels, showed enhanced expression of auxin synthesis

genes compared to wild-type plants at 35 days **(Fig. 9G)**. These results demonstrate

that excessive SA in the early stages of plant development inhibits auxin

accumulation, thereby hindering normal growth.

Discussion

In nature, plants exhibit enhanced defense against herbivorous insects as they age, with juvenile plants experiencing more frequent attacks than their adult counterparts. This observation has led to the development of the concepts of ARR and PVH; however, the precise mechanisms underlying these phenomena remain largely unknown. In this study, we elucidate the fine regulation of SA-mediated ARR against phloem-feeding whiteflies in plants (Fig. 1). We highlight the importance of the PAL synthesis pathway in age-related intrinsic SA accumulation, identifying the key synthetic gene *NbPAL6* in *N. benthamiana* (Fig. 2). Additionally, we identify the age-related upstream TF NbMYB42 and NbARF18La/b that are involved in SA regulation (Fig. 3, and 4).

Furthermore, we provide insights into why SA-mediated resistance is not fully established in juvenile plants. Specifically, high levels of auxin in juvenile plants activate *Nb*miR160c, which silences the *NbARF18La/b* genes, ultimately impairing SA accumulation. This mechanism also prevents excessive SA from disrupting early plant growth. Our findings deepen the understanding of the PVH, demonstrating that juvenile plants exhibit lower herbivore tolerance while prioritizing urgent growth needs (Fig. 10).

Our previous research has established the significant role of inducible SA in tobacco defense against whiteflies (*49*). In the current study, we found that the increased intrinsic SA levels associated with age also correlate with enhanced defense against whiteflies in adult plants (Fig. 1). Although JA is crucial for plant defense against whiteflies (*49*), our findings indicate that the intrinsic JA levels did not vary with age across several Solanaceae species (Fig. 1F), indicating that JA may not be a key phytohormonal signal in plant ARR against whiteflies. Recently, the role of SA in plant defense against phloem-feeding insects has gained attention (*50–52*), and our findings broaden the understanding of SA-mediated defenses from a novel perspective.

Moreover, we found the PAL pathway, rather than the ICS pathway, in SA biosynthesis changes at the transcriptional level with age in *N. benthamiana* (Fig. 2B and C). Previous studies have shown the key role of *PAL* genes in SA accumulation in tobacco and other plant species (*38*, *53*, *54*). Here, we systematically analyzed *PAL* genes in *N. benthamiana* (Fig. S2) and determined the age-related upregulation of *NbPAL6* expression (Fig. 2C) and its role in age-related SA accumulation (Fig. 2E). It is worth noting that our conclusion does not imply that *NbPAL6* is the sole *NbPAL* gene responsible for SA synthesis in *N. benthamiana*, as other *PALs* may contribute redundantly to SA synthesis. We suspect that additional *NbPALs* may participate in SA

accumulation under various conditions, including biotic or abiotic stresses, warranting further investigation.

While extensive attention has been given to the upstream regulation of SA synthesis pathways, most studies have focused on the ICS pathway (*55*). The PAL pathway is known to contribute to the synthesis of numerous metabolic compounds in plants, such as flavonoids and lignin (*56*), but the regulation of *PAL* gene expression related to SA synthesis has been underexplored. For instance, in rice, the brown planthopper induces OsMYB30, which activates *OsPAL* gene expression, leading to increased SA and lignin accumulation and enhanced resistance (*38*). In our study, we demonstrated that the TF NbMYB42 activates *NbPAL6* expression by binding to its promoter, thereby enhancing SA levels and whitefly defense in an age-related constitutive, but not inducible, manner **(Fig. 3)**. Interestingly, in our previously published work, NbMYB42 was also shown to regulate lignin synthesis pathways in *N. benthamiana*, contributing to enhanced whitefly resistance (*39*). These observations combined highlight the necessity of MYB-PAL modules in plant SA and lignin synthesis, as well as their defense against phloem-feeding insects. In summary, our findings elucidate the mechanisms of age-related SA synthesis and regulation in plants, providing a valuable reference for future studies on the significance of age-enhanced SA in plant physiology and its induction in plant-pathogen and plant-insect interactions.

It is well established that SA and auxin signaling pathways often interact antagonistically (*57*). Elevated auxin levels correlate with reduced *PR* gene expression (*58*, *59*) and increased susceptibility to pathogens (*59*–*61*). In this study, we found that increased auxin levels can also impact plant defense against phloem-feeding insects by antagonizing SA levels **(Fig. 6, A to F)**. Previous studies have indicated that auxin signaling may affect SA biosynthesis, although the underlying mechanisms remain largely unknown (*57*, *59*). Our study revealed that auxin inhibits the expression of the key synthesis gene *NbPAL* and its transcriptional regulators, NbMYB42 and NbARF18La/b, thereby reducing SA biosynthesis **(Fig. 6, G to O)**. In addition, repression of auxin signaling by SA has been demonstrated in studies on plant growth. For example, SA treatment decreased biomass in Arabidopsis, an effect that was abolished in the *auxin resistant 3* (*axr3*) mutant background, which harbors an auxin-negative regulator (*62*). Similarly, we found that increased SA levels influenced auxin accumulation and related gene expression **(Fig. 9, D to F)**, leading to reduced growth in *N. benthamiana* plants **(Fig. 9, A to C)**. Collectively, these findings deepen our understanding of the crosstalk between SA and auxin, particularly regarding upstream biosynthesis.

Studies have shown that ARFs activate MYB TFs involved in various essential physiological processes (*63*–*66*). Our study complementarily reports that the two ARFs, NbARF18La and NbARF18Lb, positively regulate SA biosynthesis by

promoting SA-related MYB TFs, thereby enhancing plant defense against herbivores **(Fig. 4 and Fig. 5)**. Notably, the two ARFs we identified can interact with one another **(Fig. S6)**. Our findings indicate that auxin inhibits SA accumulation, positioning NbARFs as central hubs in this interaction. While ARFs are known participants in the antagonism between auxin and other phytohormones, such as ABA and JA (*40*, *67*), little is reported regarding their relationship with SA. Previous studies have shown that the auxin-induced ARF7 represses the expression of SA-downstream *PR* genes that regulate bacterial infections in tomato plants (*68*). In contrast, we found that in *N. benthamiana* auxin represses the transcriptional activators NbARF18La/b **(Fig. 6)**, thereby negatively affecting upstream SA signaling.

Additionally, this work reveals the mechanism by which auxin represses NbARFs through miR160c-mediated post-transcriptional regulation **(Fig. 7)**. Prior studies have demonstrated that several miRNAs target *ARF* transcripts to regulate growth and development. For example, in Arabidopsis, miR160 targets *ARF10/16/17* (*69*), while miR167 antagonizes *ARF6/8* (*47*). Our findings further illuminate the relatively unexplored role of miR160c in regulating ARFs, suggesting a mechanism by which it fine-tunes plant defense against phloem-feeding insects **(Fig. 7 and Fig. 8)**.

Age-related changes in sRNA levels have been shown to play an important role in plant growth and stress responses. In Arabidopsis, the decline in the levels of miR156/7 with age drives plant maturation (*70*, *71*), while an increase in miR172 with plant age promotes the transition toward adulthood and flowering (*72*). Notably, age-related miRNA gradients are also implicated in plant immunity against pathogens. For example, miR172b levels are very low during the early stage of seedling development but increase over time, enhancing pathogen immunity by reducing the expression of *TARGET OF EAT1/2* (*TOE1/2*), repressors of the immune receptor FLAGELLIN-SENSING2 (FLS2) (*73*). In this study, we demonstrate the role of age-related sRNA changes in regulating insect resistance. Specifically, the decrease in miRNA160c levels with age alleviates inhibition of its target defense-related TFs, thereby promoting insect resistance **(Fig. 8)**.

The regulation of miRNA levels in plants is complex, and influenced by various factors including phytohormones, which can either increase or decrease miRNA levels (*18*, *74*). We found that auxin increased miR160c levels by promoting its transcription **(Fig. 8, J and K)**. The connections between miRNAs and phytohormones are an emerging area of research, with multiple recent studies highlighting the role of miRNAs in regulating plant phytohormonal pathways (*74*, *75*, *76*). As a supplementary finding, we revealed that auxin-induced miRNA can target key regulators involved in the antagonism of SA synthesis **(Fig. 10)**. These findings broaden our understanding of the molecular and physiological mechanisms underlying phytohormone interactions. Our work demonstrates, for the first time, the

644 role of miRNA in mediating the crosstalk between developmental auxin and defensive
645 SA, balancing plant growth and defense **(Fig. 10)**.
646
Developing new crop varieties with improved product traits and enhanced stress
resilience is a primary goal of global crop breeding, however, this effort is often
limited by the growth-defense (G-D) trade-off (*76*). Consequently, the G-D trade-off
has garnered increasing attention. However, the G-D balance in plants is not
immutable; it can be influenced by various factors, such as developmental stage and
environmental conditions. In this study, we specifically focused on the mechanisms
underlying the trade-off between growth and insect resistance across different plant
ages **(Fig. 10)**. We conclude that the auxin-miR160c module represses the ARFs-
MYB-PAL module, mediating insect defense while ensuring early growth during the
juvenile stage. As plants mature, this repression is gradually alleviated, allowing for
the establishment of SA-mediated resistance alongside continued plant growth. These
findings are likely to inspire further strategies for regulating crop growth and stress
resistance by fine-tuning phytohormone interactions, key TFs, metabolic pathways,
and miRNA levels. Importantly, genetic optimization in breeding programs must fully
consider the age dynamics of plant-insect interactions to overcome the G-D trade-offs
and develop ideal crop varieties.

Materials and Methods

Insect and plant materials

The *Bemisia tabaci* MEAM1 whitefly (*mtCOI* GenBank accession: GQ332577) was used in this study. Whiteflies were reared on cotton plants (*Gossypium hirsutum* cv. Zhemian 1793). Whitefly adults within 7 days post-emergence were used for experiments. Green peach aphids (*Myzus persicae*) were reared on tobacco (*N. tabacum* cv. NC89), with adult winged aphids also taken at 7 days post-emergence for experiments. The western flower thrips (*Frankliniella occidentalis*) were similarly reared on cotton plants (*G. hirsutum* cv. Zhemian 1793). Adult thrips, within 7 days post-emergence, were taken in the experiments.

N. benthamiana seeds (accession LAB and line H2B-RFP), tobacco (*N. tabacum* cv. NC89), and tomato (*Solanum lycopersicum* cv. Hezuo903) were sown in a soil mixture composed of peat moss, perlite, and vermiculite in a ratio of 5:3:1 (v/v/v) and placed in climate chambers. After approximately 14 days, when the *N. benthamiana*, tobacco, and tomato seedlings had germinated in a seed starter tray, the resulting plants were transplanted into plastic pots for individual cultivation. *N. benthamiana* plants at 7 days post-transplanting (21 days old, 3–4 true leaves) were classified as juvenile plants (low resistance), while *N. benthamiana* plants at 21 days post-transplanting (35 days old, 8–9 true leaves) were referred to as adult plants (high resistance) **(Fig. 2D)**. For tobacco, 7 days post-transplanting (21 days old, 3–4 true leaves) were classified as juvenile plants and 21 days post-transplanting (35 days old, 6–7 true leaves) were classified as the adult plants. For tomatoes, 7 days post-transplanting (21 days old, 3–4 true leaves) were classified as juvenile plants and 21 days post-transplanting (35 days old, 8–9 true leaves) were classified as adult plants.

Insect-rearing plants and experimental plants were grown in climate chambers at Zhejiang University (Hangzhou, China) at 26 ± 1°C, 65 ± 10% relative humidity, and a photoperiod of 14 h light: 10 h dark. All the insect-plant interaction experiments were conducted under the same conditions.

Insect bioassay

Bioassays were conducted using the following methods and standards. The test plants were placed in specially designed nylon cages (30 cm × 30 cm × 30 cm, 120 mesh), and the test insects (5 female and 5 male adults of MEAM1 whitefly, 3 female adults of aphid, and 5 female and 5 male adults of thrip respectively) were released into each of the cages. After two days, the performance of insects (whitefly, survival rate, and fecundity; aphid, offspring number; thrips, survival rate) was observed and recorded.

For the age-related bioassays, 21-day-old and 35-day-old *N. benthamiana* plants were simultaneously prepared based on their growth timeline. Specifically, we first sowed

the initial batch of *N. benthamiana* and when these plants reached 14 days old, we
sowed the second batch. Both batches were grown in a strictly controlled environment
as described above. By the time the first batch reached 35 days, the second batch was
21 days old. These two batches of *N. benthamiana* were then used for subsequent
bioassays as mentioned above. Bioassays related to gene and miRNA functions were
performed through *Agrobacterium*-mediated transient gene expression or VIGS. Two
days after gene overexpressing or 14 days after gene silencing **(Fig. 2D)**, whiteflies
were released onto the plants, and whitefly survival and fecundity were determined
after two days. For bioassays involving the external application of chemicals to the
plants, whitefly assessments commenced two days after chemical treatment.
714
Transient expression assays
The full-length cDNAs of the selected genes (*NbPAL6*, *NbMYB42*, and *NbARF18L*s)
were amplified via PCR using gene-specific primer pairs **(Table S2)**. The full-length
CDS sequences were inserted into pCAMBIA1305 vectors using the NovoRec plus
one-step PCR cloning kit (Novoprotein, Suzhou, China) to generate gene expression
constructs (pCAMBIA1305-*NbPAL6*-GFP, pCAMBIA1305-*NbMYB42*-GFP,
pCAMBIA1305-*NbARF18La*-GFP, and pCAMBIA1305-*NbARF18Lb*-GFP). For
transient expression in *N. benthamiana*, *Agrobacterium tumefaciens* strain EHA105
containing the respective vector was cultured overnight in LB medium at 28 °C.
Bacterial cultures were harvested by centrifugation, resuspended in buffer (10 mM
MES, pH 5.7, 10 mM $MgCl_2$, and 200 μM acetosyringone) to an OD600 of 0.8, and
infiltrated into leaves of 21-day-old or 35-day-old *N. benthamiana* using a needleless
syringe **(Fig. 2D)**. The infiltrated plants were then maintained in the previously
described climate chambers. Two days later, the treated plants were used for follow-
up experiments including bioassays and sample collection **(Fig. 2D)**.
730
VIGS assays
A 300 bp fragment of the target gene from *N. benthamiana* was identified using the
VIGS tool on the Sol Genomics Network (https://vigs.solgenomics.net/). The gene
fragment was amplified from *N. benthamiana* cDNA via PCR and cloned into the
pTRV2 vector. The resulting plasmid was introduced into *A. tumefaciens* strain
EHA105 by electroporation. Following cultivation, bacterial cells were harvested,
resuspended, and adjusted to an OD600 of 0.2. The cultures were incubated at room
temperature for at least 3 h. For leaf infiltration, cultures containing pTRV1 were
mixed in equal volumes with those containing pTRV2:00 (empty vector, control),
pTRV2-*NbPAL6*, pTRV2-*NbMYB42* or pTRV2-*NbARF18L*s. These suspensions were
infiltrated into 21-day-old *N. benthamiana* leaves using a 1 mL needleless syringe
(Fig. 2D). Following infiltration, the *N. benthamiana* plants were maintained in the
same climate chamber conditions. Fourteen days later, the treated plants were used for
follow-up experiments, including bioassays and sample collection **(Fig. 2D)**.
745

Subcellular localization

The pCAMBIA1305-*NbMYB42*-GFP, pCAMBIA1305-*NbARF18La*-GFP,

pCAMBIA1305-*NbARF18Lb*-GFP, and pCAMBIA1305-GFP (empty vector)

constructs were used for subcellular localization. These constructs were infiltrated

into the leaves of *N. benthamiana* line H2B-RFP via *A. tumefaciens* strain EHA105.

Two days post-infiltration, GFP fluorescence signals were visualized using confocal

microscopy (Zeiss LSM710, Oberkochen, Germany).

RNA extraction and RT-qPCR analysis

Total RNA was extracted using *AG RNAex Pro* reagent (Accurate Biology, Changsha,

China). For mRNA, single-stranded cDNA was synthesized using the Evo *M-MLV* RT

Kit with gDNA Clean for qPCR Ver.2 (Accurate Biology Changsha, China). For plant

miRNA cDNA synthesis, the miRNA 1st strand cDNA synthesis kit (Accurate

Biology, Changsha, China) was used. RT-qPCR was conducted on a Bio-Rad CFX96

real-time PCR system (Bio-Rad, California, USA) with specific primers **(Table S2)**.

Data were analyzed with Bio-Rad CFX Manager Software, employing the $2^{-\Delta\Delta C_T}$

method. The *N. benthamiana* housekeeping gene *NbACTIN* served as the internal

control for gene expression analysis, and the expression of the *NbU6* snRNA genes

was employed as the internal control for miRNA level analysis.

Measurement of phytohormone content

For the quantification of phytohormones (SA, JA, and IAA), *N. benthamiana* leaves

were promptly excised and ground into powder in liquid nitrogen. Leaf powder (0.15

g) was mixed with 1 mL of HPLC-grade ethyl acetate (Sinopharm, Shanghai, China)

containing 10 ng D4-SA (Quality Control Chemicals, Spokane, WA, USA), 10 ng D6-

JA (Quality Control Chemicals, Spokane, WA, USA), and 5 ng D5-IAA (OlhemIm,

Shanghai, China). The mixture was fully vortexed and centrifuged. The supernatant

was collected and dried using a vacuum concentrator (Eppendorf AG, Hamburg,

Germany) at 30°C. The residue was resuspended in 110 µL of methanol: H_2O (50:50,

v/v), and the resulting supernatant was analyzed via high-performance liquid

chromatography-tandem mass spectrometry (TripleTOF 5600+; AB Sciex, Redwood

City, CA, USA).

Degradome sequencing

The degradome sequencing analysis, performed according to established protocols

from a previous study (*77*), was outsourced to Lianchuan Biotechnology Co., Ltd.

Poly(A) RNA was purified from total plant RNA using poly-T oligo magnetic beads

(Invitrogen, Carlsbad, CA, USA) through two rounds of purification. First-strand

cDNA synthesis was performed with a 3′-adapter random primer, followed by size

selection using AMPureXP beads (Beckman Coulter, Indianapolis, IN, USA) and

PCR amplification. The final cDNA library had an average insert size of 200–400 bp.

Sequencing was carried out with 50 bp single-end reads on an Illumina HiSeq 2500

(Illumina, San Diego, CA, USA). CleaveLand 3.0 and TargetFinder 50 were
employed to identify potential sliced targets of known miRNAs.
790
Y1H and Y2H assay
For the Y1H assay, approximately 2 kb of the promoter regions of the target genes
(*NbPAL6* and *NbMYB42*) were inserted into the pAbAi vector to construct the bait
plasmids. These plasmids were integrated into the Y1HGold yeast strain via
polyethylene glycol-mediated transformation, following in the Yeast User Manual
PT4087-1 (Clontech, Shiga, Japan). The CDSs of the corresponding TFs NbMYB42
and NbARF18La/b were cloned into the pGADT7 vector. The relevant vectors were
co-expressed in Y1H yeast cells and screened with varying concentrations of
Aureobasidin A (AbA, TaKaRa, Beijing, China).
800
For the Y2H assays, the full-length CDSs of the target proteins NbARF18La and
NbARF18Lb were ligated into pGBKT7 and pGADT7 (Clontech, Shiga, Japan),
respectively. The paired vectors were transformed into Y2HGold yeast strain cells via
PEG/LiAc transformation and grown on SD −Leu/−Trp medium. After 3–4 days of
incubation, the clones were transferred onto SD −Leu/−Trp/−His/−Ade medium and
incubated at 29°C for 3–4 days. The primers used for vector construction related to
the Y1H and Y2H assays are listed in **Table S2**.
808
Histochemical GUS activity assays
For the reporter constructs, the promoters of the target genes *NbPAL6* and *NbMYB42*
were introduced into the pBI121 vector containing a *GUS* gene-coding region.
Simultaneously, the full-length cDNAs of the corresponding target TFs (NbMYB42
and NbARF18La/b) were inserted into the pCAMBIA1305 vector to generate effector
constructs (pCAMBIA1305-*NbMYB42*-GFP, pCAMBIA1305-*NbARF18La*-GFP,
pCAMBIA1305-*NbARF18Lb*-GFP, as described in the Transient expression assays
section). The pCAMBIA1305 vector without the target CDS was used as the negative
control (empty effector). These vectors were transformed into *A. tumefaciens* strain
EHA105, and reporter vectors were co-transformed with the corresponding effector or
control vectors for transient expression. GUS activity was assessed using a GUS
histochemical assay kit (Coolaber, Beijing, China) according to the manufacturer's
instructions. GUS expression was visualized through staining and documented using
an iPhone 13 Pro Max (Apple, California, USA).
823
Dual-luciferase assays
For the reporter construct, the promoters of the target genes (*NbPAL6* and *NbMYB42*)
were inserted into the reporter vector pGreenII 0800-LUC. Effector constructs,
including pCAMBIA1305-*NbMYB42*-GFP, pCAMBIA1305-*NbARF18La*-GFP,
pCAMBIA1305-*NbARF18Lb*-GFP, and the empty vector pCAMBIA1305, were the
same as those used in the Histochemical GUS activity assays. These vectors were

transformed into *A. tumefaciens* EHA105 (pSoup). In this system, LUC expression
was driven by the target gene promoters, while REN was driven by the *CaMV 35S*
promoter as an internal control. The reporter and corresponding effector constructs
were transiently transformed into *N. benthamiana* leaves. After a three-day
incubation, LUC signals were measured using the Dual-Luciferase Reporter Assay Kit
(Vazyme, Nanjing, China) and analyzed on a FlexStation 3 Multi-Mode Microplate
Reader (Molecular Devices, San Jose, CA, USA). Relative luciferase activity was
calculated based on LUC: REN ratios.
838
ChIP assay
Leaves transformed with pCAMBIA1305-*NbMYB42*-GFP, pCAMBIA1305-
NbARF18La-GFP, or pCAMBIA1305-*NbARF18Lb*-GFP (as described in the
Transient expression assay), were collected for ChIP assays. The experiment followed
established protocols using the same transient gene expression system (*78*). Briefly,
transformed *N. benthamiana* leaves were harvested and fixed with 1% formaldehyde.
The fixation was quenched by adding 1 × PBS buffer containing 0.125 M glycine
under vacuum for 5 min. Chromatin DNA was sonicated to fragments of
approximately 200–300 bp. The chromatin complexes were immunoprecipitated (IP)
by Protein G Agarose (Beyotime, Shanghai, China) with an anti-GFP monoclonal
antibody (Beyotime, Shanghai, China). The co-IP DNA was recovered at 65°C,
purified, and analyzed by qPCR with SYBR qPCR Mix (Accurate Biology, Changsha,
China) with gene-specific primers (Table S2). The relative fold enrichment was
calculated using leaves transformed with the pCAMBIA1305-GFP (empty vector) as
a control.
854
EMSA assay
To prepare the test TF proteins for EMSA, the CDS of NbMYB42 was fused into the
GST-tagged pGEX-6P-1 vector to generate NbMYB42-GST. The NbARF18La CDS
was fused into the His-tagged pET28a-SUMO vector, generating NbARF18La-His-
SUMO, and the NbARF18Lb CDS was fused into the MBP-tagged pMAL-c5x vector,
generating NbARF18Lb-MBP. These fusion proteins were expressed in *E. coli* BL21
cells using 0.2 mM isopropyl-β-D-thiogalactopyranoside (IPTG) and purified. For
protein purification, NbMYB42-GST was purified through GST-affinity
chromatography (Beyotime, Shanghai, China), NbARF18Lb-MBP using MBPSep
Dextrin Agarose Resin 6FF (Yeasen, Shanghai, China) and NbARF18La-His with Ni
NTA Beads 6FF (Smart-Lifesciences, Changzhou, China).
866
Putative binding sites in the target gene promoters were predicted by Jasper
(https://jaspar.elixir.no/). For the NbMYB42 and *NbPAL6* promoter interaction, biotin-
labeled probes were designed based on the P3 sequence (GGTGGTTGTTGAGAGG)
of the *NbPAL6* promoter (Fig. 3G; Table S2), with a mutated version
(AAAAAAAAAAAAAAAA) serving as the mutant probe. For the interaction

between NbARF18La/b and the *NbMYB42* promoter, biotin-labeled probes targeting
the AuxRE element P2 sequence (TGTCTC) were designed **(Fig. 4, G and H; Table**
S2), with a mutated version (AAAAAA) as the control. Cold competitors (10-, 20-,
and 50-fold excess) of unlabeled probes were included in the reactions. Probes were
labeled using the EMSA Probe Biotin Labeling Kit (Beyotime, Shanghai, China).
Approximately 300 ng of the fusion protein was incubated with 30 ng of labeled
probes, and specific binding was assessed via the Chemiluminescent EMSA Kit
(Beyotime, Shanghai, China) following the manufacturer's instructions.
Chemiluminescent imaging was performed with the ChemiDoc Touch system (Bio-
Rad, California, USA). Labeled probes incubated with GST, His-SUMO, and MBP
proteins served as negative controls.
883
Pull-down assay
Pull-down assays were conducted to determine the interaction of NbARF18La and
NbARF18Lb. The recombinant proteins NbARF18La-His and NbARF18Lb-MBP
obtained in the above-mentioned EMSA assay were further used in this test. For MBP
pull-down, MBPSep Dextrin Agarose Resin (Yeasen, Shanghai, China) is prepared by
washing them with PBS and then incubating them with MBP-bait protein
NbARF18Lb or control MBP protein. After washing, the beads are incubated with
His-tagged interacting protein NbARF18La at 4°C for 3 h. Then the MBP beads
underwent four washes with washing buffer (20 mM phosphate buffer, 500 mM NaCl,
100 mM imidazole, pH 7.4) and were then collected using collecting buffer (20 mM
phosphate buffer, 500 mM NaCl, 500 mM imidazole, pH 7.4). The eluted samples
were separated by SDS-PAGE and detected by an anti-MBP (Affinity Bioscience,
Jiangsu, China) antibody and an anti-His antibody (Proteintech, Wuhan, China).
897
Bimolecular fluorescence complementation (BiFC) assay
The CDSs of *NbARF18La* and *NbARF18Lb* were inserted into p2YN and p2YC
vectors, respectively. *A. tumefaciens* EHA105 cells carrying p2YN constructs were
combined with cells containing p2YC constructs at a 1:1 ratio. This mixture was then
infiltrated into the leaves of the *N. benthamiana* H2B-RFP line. After three days of
incubation, fluorescence was visualized under a laser confocal microscope (Zeiss
LSM710, Oberkochen, Germany).
905
Exogenous application of phytohormones
The target compound was dissolved in methanol (for SA, IAA, and NAA) or dimethyl
sulfoxide (DMSO) (for Yucasin and $_L$-Kyn) to produce the stock solution. Lanolin
was selected as a slow-release medium. The lanolin was heated to a liquid state and
the stock solution was added to the lanolin to create the final mixture: SA at 1 mM;
IAA and NAA at 1, 25, and 50 μM; Yucasin and $_L$-Kyn at 50 μM. All the compounds
were purchased from Aladdin, Shanghai, China. The lanolin solution containing the
target compounds was transferred to a 1 mL sterile syringe and stored at -20°C.

Exogenous SA was applied to 21-day-old and 35-day-old *N. benthamiana* plants. IAA
and NAA were applied to 35-day-old plants, while Yucasin and $_L$-Kyn were applied to
21-day-old plants. For application, 0.25 mL of lanolin was squeezed from the syringe
and gently smeared along the midvein at the base of the leaves using latex-gloved
fingers. After two days **(Fig. 2D)**, the treated plants were used in bioassays or for
sample collection for further experiments.

Transient expression and functional inhibition of miRNA in *N. benthamiana*
To overexpress target miRNA *Nb*mi160c in *N. benthamiana*, the precursor sequences
of *Nb*miR160c were synthesized by GenScript (Nanjing, China) and inserted into the
pCAMBIA1300 vector through specific primers **(Table S2; Fig. S9A)**. The constructs
were infiltrated into 35-day-old *N. benthamiana* plants, and after two days **(Fig. 2D)**,
the treated plants were used in bioassays or for sample collection for further
experiments.

To inhibit the *Nb*miR160c function, a target mimicry (mimic160c) expression system
was introduced. The precursor sequences of mimic160c were synthesized by
GenScript (Nanjing, China) and inserted into the pCAMBIA1300 vector using
specific primers **(Table S2; Fig. S9A and F)**. The constructs were infiltrated into 21-
day-old *N. benthamiana* plants, and after two days **(Fig. 2D)**, the treated plants were
used for bioassays or sample collection for further experiments.

MiRNA targeting validation
GFP, GUS, and LUC reporter systems were used to investigate the cleavage of
NbARF18La and *NbARF18Lb* targets by *Nb*miR160c. For the GFP reporter, the
*Nb*miR160c target site of *NbARF18La* and *NbARF18Lb* mRNA, as well as its mutant
form **(Fig. S9C)** were cloned into the pCAMBIA1305-GFP vector to construct GFP
sensors through specific primers **(Table S2; Fig. S9D)**. The constructs of the target
site and mutant target site were co-infiltrated with pCAMBIA1300-*Nb*miR160c
through *A. tumefaciens* strain EHA105. After two days of infiltration, GFP
fluorescence was observed using confocal microscopy, as described above. For the
GUS and LUC reporters, the *Nb*miR160c target site of *Nb*ARF18La and *Nb*ARF18Lb
mRNA, along with its mutant form, were cloned into the pCAMBIA121-GUS and
pGreenII 0800-LUC vectors, respectively, to construct GUS and LUC sensor through
specific primers **(Table S2; Fig. S9D)**. These constructs were co-infiltrated with
pCAMBIA 1300-*Nb*miR160c. After two days of infiltration, GUS and LUC activity
was examined using the previously described method.

MiRNA dual-luciferase assay
Co-expression of the LUC reporter and *Nb*miR160c was carried out to evaluate the
influence of *Nb*miR160c on downstream genes. The effectors pCAMBIA1300-pre-

*Nb*miR160c, pCAMBIA1300-premimic160c, and pCAMBIA1300 (empty vector
control) were co-expressed with LUC reporters linked to the *NbMYB42* or *NbPAL6*
promoters. After two days of infiltration, LUC activity was examined using the
method mentioned method. The combination of effectors and reporters is shown in
Fig. 10 A to D.
The system was also used to evaluate the possible influence of *Nb*miR160c on
NbPAL6 transcription when NbARFs were overexpressed. The *LUC* gene linked to
the *NbPAL6* promoter or *NbMYB42* promoter was used as the reporter respectively.
pCAMBIA1305-NbARFs plus pCAMBIA1300-pre-*Nb*miR160c, and
pCAMBIA1305-NbARFs plus pCAMBIA1300 (empty vector control) were co-
infiltrated with the reporter into the leaves of *N. benthamiana*. LUC activity for the
different combinations was detected and compared. The combinations of effectors and
reporters are shown in **Fig. 8G** and **Fig. S10 E**.
969
970 Statistical analysis
971 All data were analyzed using IBM SPSS Statistics 22.0 and visualized using
972 GraphPad Prism 9. Statistical significance was determined using Student's *t*-test (two-
973 tailed) or one-way ANOVA, followed by Fisher's least significant difference (LSD)
974 test.

Reference

1. D. Walters, M. Heil, Costs and trade-offs associated with induced resistance. *Physiol. Mol. Plant Pathol.* **71**, 3–17 (2007).

2. D. Cipollini, C. B. Purrington, J. Bergelson, Costs of induced responses in plants. *Basic Appl. Ecol.* **4**, 79–89 (2003).

3. M. Erb, P. Reymond, Molecular interactions between plants and insect herbivores. *Annu. Rev. Plant Biol.* **70**, 527–557 (2019).

4. G. A. Howe, G. Jander, Plant immunity to insect herbivores. *Annu. Rev. Plant Biol.* **59**, 41–66 (2008).

5. J. A. Gatehouse, Plant resistance towards insect herbivores: a dynamic interaction. *New Phytol.* **156**, 145–169 (2002).

6. A. R. Zangerl, C. E. Rutledge, The probability of attack and patterns of constitutive and induced defense: a test of optimal defense theory. *Am. Nat.* **147**, 599–608 (1996).

7. P. W. Price, The plant vigor hypothesis and herbivore attack. *Oikos* **62**, 244–251 (1991).

8. S. W. Makhabu, C. Skarpe, H. Hytteborn, Z. D. Mpofu, The plant vigour hypothesis revisited – how is browsing by ungulates and elephant related to woody species growth rate? *Plant Ecol.* **184**, 163–172 (2006).

9. T. Rankenberg, B. Geldhof, H. Van Veen, K. Holsteens, B. Van De Poel, R. Sasidharan, Age-dependent abiotic stress resilience in plants. *Trends Plant Sci.* **26**, 692–705 (2021).

10. M. Develey-Rivière, E. Galiana, Resistance to pathogens and host developmental stage: a multifaceted relationship within the plant kingdom. *New Phytol.* **175**, 405–416 (2007).

11. L. Hu, L. Yang, Time to fight: molecular mechanisms of age-related resistance. *Phytopathology* **109**, 1500–1508 (2019).

12. J. H. Kim, H. R. Woo, J. Kim, P. O. Lim, I. C. Lee, S. H. Choi, D. Hwang, H. G. Nam, Trifurcate feed-forward regulation of age-dependent cell death involving *miR164* in *Arabidopsis*. *Science* **323**, 1053–1057 (2009).

13. Y.-B. Mao, Y.-Q. Liu, D.-Y. Chen, F.-Y. Chen, X. Fang, G.-J. Hong, L.-J. Wang, J.-W. Wang, X.-Y. Chen, Jasmonate response decay and defense metabolite accumulation contributes to age-regulated dynamics of plant insect resistance. *Nat. Commun.* **8**, 13925 (2017).

14. Z. He, S. Webster, S. Y. He, Growth–defense trade-offs in plants. *Curr. Biol.* **32**, R634–R639 (2022).

15. M. Vanstraelen, E. Benková, Hormonal interactions in the regulation of plant development. *Annu. Rev. Cell Dev. Biol.* **28**, 463–487 (2012).

16. A. Robert-Seilaniantz, M. Grant, J. D. G. Jones, Hormone crosstalk in plant disease and defense: more than just jasmonate-salicylate antagonism. *Annu. Rev. Phytopathol.* **49**, 317–343 (2011).

17.   M. L. Berens, H. M. Berry, A. Mine, C. T. Argueso, K. Tsuda, Evolution of
hormone signaling networks in plant defense. *Annu. Rev. Phytopathol.* **55**, 401–425
(2017).
18.   X. Song, Y. Li, X. Cao, Y. Qi, MicroRNAs and their regulatory roles in
plant–environment interactions. *Annu. Rev. Plant Biol.* **70**, 489–525 (2019).
19.   J. Curaba, M. B. Singh, P. L. Bhalla, miRNAs in the crosstalk between
phytohormone signalling pathways. *J. Exp. Bot.* **65**, 1425–1438 (2014).
20.   M. Altmann, S. Altmann, P. A. Rodriguez, B. Weller, L. Elorduy Vergara, J.
Palme, N. Marín-de La Rosa, M. Sauer, M. Wenig, J. A. Villaécija-Aguilar, J. Sales,
C.-W. Lin, R. Pandiarajan, V. Young, A. Strobel, L. Gross, S. Carbonnel, K. G.
Kugler, A. Garcia-Molina, G. W. Bassel, C. Falter, K. F. X. Mayer, C. Gutjahr, A. C.
Vlot, E. Grill, P. Falter-Braun, Extensive signal integration by the phytohormone
protein network. *Nature* **583**, 271–276 (2020).
21.   S.-S. Liu, P. J. De Barro, J. Xu, J.-B. Luan, L.-S. Zang, Y.-M. Ruan, F.-H.
Wan, Asymmetric mating interactions drive widespread invasion and displacement in
a whitefly. *Science* **318**, 1769–1772 (2007).
22.   S. Liu, J. Colvin, P. J. De Barro, Species concepts as applied to the whitefly
Bemisia tabaci systematics: how many species are there? *J. Integr. Agric.* **11**, 176–186
(2012).
23.   P. J. De Barro, S.-S. Liu, L. M. Boykin, A. B. Dinsdale, *Bemisia tabaci*: a
statement of species status. *Annu. Rev. Entomol.* **56**, 1–19 (2011).
24.   S. Kanakala, M. Ghanim, Global genetic diversity and geographical
distribution of *Bemisia tabaci* and its bacterial endosymbionts. *PLoS One* **14**,
e0213946 (2019).
25.   X.-W. Wang, P. Li, S.-S. Liu, Whitefly interactions with plants. *Curr. Opin.*
Insect Sci. (2017).
26.   M. Inbar, D. Gerling, Plant-mediated interactions between whiteflies,
herbivores, and natural enemies. *Annu. Rev. Entomol.* **53**, 431–448 (2008).
27.   X.-W. Wang, S. Blanc, Insect transmission of plant single-stranded DNA
viruses. *Annu. Rev. Entomol.* **66**, 389–405 (2021).
28.   S. Morin, P. W. Atkinson, L. L. Walling, Whitefly–plant interactions: an
integrated molecular perspective. *Annu. Rev. Entomol.* **69**, 503–525 (2024).
29.   J. Bally, H. Jung, C. Mortimer, F. Naim, J. G. Philips, R. Hellens, A.
Bombarely, M. M. Goodin, P. M. Waterhouse, The rise and rise of *Nicotiana*
benthamiana : a plant for all reasons. *Annu. Rev. Phytopathol.* **56**, 405–426 (2018).
30.   M. M. Goodin, D. Zaitlin, R. A. Naidu, S. A. Lommel, *Nicotiana*
benthamiana: its history and future as a model for plant–pathogen interactions.
Mol. Plant-Microbe Interact. **2015 (1)**, 28–39 (2015).
31.   W. Han, J. Wang, F. Zhang, S. Ji, Y. Zhong, Y. Liu, S. Liu, X. Wang, A new
feature of the laboratory model plant *Nicotiana benthamiana* : dead-end trap for
sustainable field pest control. *Plants People Planet*, ppp3.10484 (2024).

32.   X. Wang, H. Han, R. Xie, B. Zhang, W. Ge, X. Ji, S. Liu, W. Wang,
Metabolic and molecular insights into *Nicotiana benthamiana* trichome exudates: an
ammunition depot for plant resistance against insect pests. *Plant, Cell Environ.* (2024)
doi: 10.1111/pce.15135.
33.   X.-Y. Zhang, W.-H. Han, F.-B. Zhang, J.-X. Wang, S.-S. Liu, X.-W. Wang,
Attraction of *Nicotiana benthamiana* to *Bemisia tabaci* is related to a chemical signal
in plant volatile, undecane. *Pest Manage. Sci.* (2024) doi: 10.1002/ps.8411.
34.   T. Gaffney, L. Friedrich, B. Vernooij, D. Negrotto, G. Nye, S. Uknes, E.
Ward, H. Kessmann, J. Ryals, Requirement of salicylic acid for the induction of
systemic acquired resistance. *Science* **261**, 754–756 (1993).
35.   Y. Peng, J. Yang, X. Li, Y. Zhang, Salicylic acid: biosynthesis and signaling.
Annu. Rev. Plant Biol. **72**, 761–791 (2021).
36.   C. J. Arntzen, Using tobacco to treat cancer. *Science* **321**, 1052–1053 (2008).
37.   B. Craven-Bartle, B. Pascual, F. M. Ca, A Myb transcription factor regulates
genes of the phenylalanine pathway in maritime pine. *Plant J.* (2013).
38.   J. He, Y. Liu, D. Yuan, M. Duan, Y. Liu, Z. Shen, C. Yang, Z. Qiu, D. Liu, P.
Wen, J. Huang, D. Fan, S. Xiao, Y. Xin, X. Chen, L. Jiang, H. Wang, L. Yuan, J. Wan,
An R2R3 MYB transcription factor confers brown planthopper resistance by
regulating the phenylalanine ammonia-lyase pathway in rice. *Proc. Natl. Acad. Sci.*
U.S.A. **117**, 271–277 (2020).
39.   F.-B. Zhang, S.-X. Ji, J.-G. Yang, X.-W. Wang, W.-H. Han, Genome-wide
analysis of MYB family in *Nicotiana benthamiana* and the functional role of the key
members in resistance to *Bemisia tabaci*. *Int. J. Biol. Macromol.* **235**, 123759 (2023).
40.   C. Cancé, R. Martin-Arevalillo, K. Boubekeur, R. Dumas, Auxin response
factors are keys to the many auxin doors. *New Phytol.* **235**, 402–419 (2022).
41.   D. R. Boer, A. Freire-Rios, W. A. M. van den Berg, T. Saaki, I. W. Manfield,
S. Kepinski, I. López-Vidrieo, J. M. Franco-Zorrilla, S. C. de Vries, R. Solano, D.
Weijers, M. Coll, Structural basis for DNA binding specificity by the auxin-dependent
ARF transcription factors. *Cell* **156**, 577–589 (2014).
42.   T. J. Guilfoyle, The PB1 domain in auxin response factor and Aux/IAA
proteins: a versatile protein interaction module in the auxin response. *Plant Cell* **27**,
33–43 (2015).
43.   D. A. Korasick, Defining a two-pronged structural model for PB1
(Phox/Bem1p) domain interaction in plant auxin responses. *J. Biol. Chem.* **290**,
12868-78 (2015).
44.   S.-B. Li, Z.-Z. Xie, C.-G. Hu, J.-Z. Zhang, A review of auxin response
factors (ARFs) in plants. *Front. Plant Sci.* **7** (2016).
45.   K. Jiang, T. Asami, Chemical regulators of plant hormones and their
applications in basic research and agriculture. *Biosci. Biotechnol. Biochem.* **82**, 1265–
1300 (2018).
46.   E. Marin, V. Jouannet, A. Herz, A. S. Lokerse, D. Weijers, H. Vaucheret, L.
Nussaume, M. D. Crespi, A. Maizel, miR390, *Arabidopsis TAS3* tasiRNAs, and their

AUXIN RESPONSE FACTOR targets define an autoregulatory network quantitatively
regulating lateral root growth. *Plant Cell* **22**, 1104–1117 (2010).
47. N. Liu, S. Wu, J. Van Houten, Y. Wang, B. Ding, Z. Fei, T. H. Clarke, J. W.
Reed, E. Van Der Knaap, Down-regulation of *AUXIN RESPONSE FACTORS 6* and *8*
by microRNA 167 leads to floral development defects and female sterility in tomato.
J. Exp. Bot. **65**, 2507–2520 (2014).
48. J.-J. Wang, H.-S. Guo, Cleavage of *INDOLE-3-ACETIC ACID*
INDUCIBLE28 mRNA by microRNA847 upregulates auxin signaling to modulate
cell proliferation and lateral organ growth in Arabidopsis. *Plant Cell* **27**, 574–590
(2015).
49. Y.-X. Liu, W.-H. Han, J.-X. Wang, F.-B. Zhang, S.-X. Ji, Y.-W. Zhong, S.-S.
Liu, X.-W. Wang, Differential induction of JA/SA determines plant defense against
successive leaf-chewing and phloem-feeding insects. *J. Pest Sci.* (2024) doi:
10.1007/s10340-024-01821-x.
50. T. Züst, A. A. Agrawal, Mechanisms and evolution of plant resistance to
aphids. *Nat. Plants* **2**, 15206 (2016).
51. R. Blundell, J. E. Schmidt, A. Igwe, A. L. Cheung, R. L. Vannette, A. C. M.
Gaudin, C. L. Casteel, Organic management promotes natural pest control through
altered plant resistance to insects. *Nat. Plants* **6**, 483–491 (2020).
52. D. Mertens, M. Fernández de Bobadilla, Q. Rusman, J. Bloem, J. C. Douma,
E. H. Poelman, Plant defence to sequential attack is adapted to prevalent herbivores.
Nat. Plants **7**, 1347–1353 (2021).
53. M. B. Shine, J. Yang, M. El-Habbak, P. Nagyabhyru, D. Fu, D. Navarre, S.
Ghabrial, P. Kachroo, A. Kachroo, Cooperative functioning between phenylalanine
ammonia lyase and isochorismate synthase activities contributes to salicylic acid
biosynthesis in soybean. *New Phytol.* **212**, 627–636 (2016).
54. D. Ogawa, N. Nakajima, S. Seo, I. Mitsuhara, H. Kamada, Y. Ohashi, The
phenylalanine pathway is the main route of salicylic acid biosynthesis in *Tobacco*
mosaic virus-infected tobacco leaves. *Plant Biotechnol.* **23**, 395–398 (2006).
55. P. Ding, Y. Ding, Stories of salicylic acid: a plant defense hormone. *Trends*
Plant Sci. **25**, 549–565 (2020).
56. N. Dong, H. Lin, Contribution of phenylpropanoid metabolism to plant
development and plant–environment interactions. *J. Integr. Plant Biol.* **63**, 180–209
(2021)
57. A. Robert-Seilaniantz, M. Grant, J. D. G. Jones, Hormone crosstalk in plant
disease and defense: more than just jasmonate-salicylate antagonism. *Annu. Rev.*
Phytopathol. **49**, 317–343 (2011).
58. J.-E. Park, J.-Y. Park, Y.-S. Kim, P. E. Staswick, J. Jeon, J. Yun, S.-Y. Kim, J.
Kim, Y.-H. Lee, C.-M. Park, GH3-mediated auxin homeostasis links growth
regulation with stress adaptation response in *Arabidopsis*. *J. Biol. Chem.* **282**, 10036–
10046 (2007).

59.   A. Robert-Seilaniantz, D. MacLean, Y. Jikumaru, L. Hill, S. Yamaguchi, Y.
Kamiya, J. D. G. Jones, The microRNA miR393 re-directs secondary metabolite
biosynthesis away from camalexin and towards glucosinolates. *Plant J.* **67**, 218–231
(2011).
60.   L. Navarro, R. Bari, P. Achard, P. Lisón, A. Nemri, N. P. Harberd, J. D. G.
Jones, DELLAs control plant immune responses by modulating the balance of
jasmonic acid and salicylic acid signaling. *Curr. Biol.* **18**, 650–655 (2008).
61.   S. A. McClerklin, S. G. Lee, C. P. Harper, R. Nwumeh, J. M. Jez, B. N.
Kunkel, Indole-3-acetaldehyde dehydrogenase-dependent auxin synthesis contributes
to virulence of *Pseudomonas syringae* strain DC3000. *PLoS Pathog.* **14**, e1006811
(2018).
62.   J. V. Canet, A. Dobón, F. Ibáñez, L. Perales, P. Tornero, Resistance and
biomass in Arabidopsis: a new model for salicylic acid perception. *Plant Biotechnol.
J.* **8**, 126–141 (2010).
63.   R. Ghelli, P. Brunetti, N. Napoli, A. De Paolis, V. Cecchetti, T. Tsuge, G.
Serino, M. Matsui, G. Mele, G. Rinaldi, G. A. Palumbo, F. Barozzi, P. Costantino, M.
Cardarelli, A newly identified flower-specific splice variant of *AUXIN RESPONSE
FACTOR8* regulates stamen elongation and endothecium lignification in Arabidopsis.
Plant Cell **30**, 620–637 (2018).
64.   X.-F. Xu, B. Wang, Y.-F. Feng, J.-S. Xue, X.-X. Qian, S.-Q. Liu, J. Zhou, Y.-
H. Yu, N.-Y. Yang, P. Xu, Z.-N. Yang, AUXIN RESPONSE FACTOR17 directly
regulates *MYB108* for anther dehiscence. *Plant Physiol.* **181**, 645–655 (2019).
65.   G. Qu, D. Peng, Z. Yu, X. Chen, X. Cheng, Y. Yang, T. Ye, Q. Lv, W. Ji, X.
Deng, B. Zhou, Advances in the role of auxin for transcriptional regulation of lignin
biosynthesis. *Funct. Plant Biol.* **48**, 743–754 (2021).
66.   W. Jiang, Y. Xia, X. Su, Y. Pang, ARF2 positively regulates flavonols and
proanthocyanidins biosynthesis in *Arabidopsis thaliana*. *Planta* **256**, 44 (2022).
67.   Y. Li, S. Han, Y. Qi, Advances in structure and function of auxin response
factor in plants. *J. Integr. Plant Biol.* **65**, 617–632 (2023).
68.   X. Kong, C. Zhang, H. Zheng, M. Sun, F. Zhang, M. Zhang, F. Cui, D. Lv, L.
Liu, S. Guo, Y. Zhang, X. Yuan, S. Zhao, H. Tian, Z. Ding, Antagonistic interaction
between auxin and SA signaling pathways regulates bacterial infection through lateral
root in *Arabidopsis*. *Cell Rep.* **32**, 108060 (2020).
69.   J.-W. Wang, L.-J. Wang, Y.-B. Mao, W.-J. Cai, H.-W. Xue, X.-Y. Chen,
Control of root cap formation by microRNA-targeted auxin response factors in
Arabidopsis. *Plant Cell* **17**, 2204–2216 (2005).
70.   J. Gao, K. Zhang, Y.-J. Cheng, S. Yu, G.-D. Shang, F.-X. Wang, L.-Y. Wu, Z.-
G. Xu, Y.-X. Mai, X.-Y. Zhao, D. Zhai, H. Lian, J.-W. Wang, A robust mechanism for
resetting juvenility during each generation in *Arabidopsis*. *Nat. Plants* **8**, 257–268
(2022).
71.   S. Bergonzi, M. C. Albani, E. V. Loren Van Themaat, K. J. V. Nordström, R.
Wang, K. Schneeberger, P. D. Moerland, G. Coupland, Mechanisms of age-dependent

response to winter temperature in perennial flowering of *Arabis alpina*. *Science* **340**,
1094–1097 (2013).
72.   C.-M. Zhou, T.-Q. Zhang, X. Wang, S. Yu, H. Lian, H. Tang, Z.-Y. Feng, J.
Zozomova-Lihová, J.-W. Wang, Molecular basis of age-dependent vernalization in
Cardamine flexuosa. *Science* **340**, 1097–1100 (2013).
73.   Y. Zou, S. Wang, Y. Zhou, J. Bai, G. Huang, X. Liu, Y. Zhang, D. Tang, D.
Lu, Transcriptional regulation of the immune receptor FLS2 controls the ontogeny of
plant innate immunity. *Plant Cell* **30**, 2779–2794 (2018).
74.   T. Li, N. Gonzalez, D. Inzé, M. Dubois, Emerging connections between small
RNAs and phytohormones. *Trends Plant Sci.* **25**, 912–929 (2020).
75.   P. Singh, P. Dutta, D. Chakrabarty, miRNAs play critical roles in response to
abiotic stress by modulating cross-talk of phytohormone signaling. *Plant Cell Rep.* **40**,
1617–1630 (2021).
76.   E. Shen, T. Zhao, Q. Zhu, Are miRNAs applicable for balancing crop growth
and defense trade-off? *New Phytol.* **243**, 1670–1680 (2024).
77.   W.-H. Han, J.-X. Wang, F.-B. Zhang, Y.-X. Liu, H. Wu, X.-W. Wang, Small
RNA and degradome sequencing reveal important microRNA function in *Nicotiana*
tabacum response to *Bemisia tabaci*. *Genes* **13**, 361 (2022).
78.   T. Zhu, X. Zhou, L. Zhang, H. Zhang, P. Zhang, X. You, P. E. Jameson, T.
Ma, L. Guo, Ethylene-induced *NbMYB4L* is involved in resistance against tobacco
mosaic virus in *Nicotiana benthamiana*. *Mol. Plant Pathol.* **23**, 16-31 (2021).
1203

Acknowledgments：

We thank Rong Jin from the Agricultural Experiment Station, Zhejiang University, for greenhouse management. We also thank Xiao-Dan Wu from the Analysis Center of Agrobiology and Environmental Sciences, Zhejiang University, for technical support in the detection and analysis of phytohormones.

Funding:

Financial support for this study was provided by:

National Natural Science Foundation of China (31925033, 32161143008)

National Key Research and Development Program (2022YFC2601002)

Author contributions:

Conceptualization: W.H., F.Z., J.W., and X.W.

Methodology: F.Z., W.H., and X.F.

Investigation: F.Z., W.H., S.J., J.W., X.F., and K.L.

Visualization: W.H. and F.Z.

Supervision: S.L. and X.W.

Writing—original draft: W.H. and F.Z.

Writing—review & editing: W.H., F.Z., S.J., and X.W.

Competing interests:

The authors declare that they have no competing interests.

Data and materials availability:

All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

 Figures

1228
Fig. 1. Plant age-related resistance against insects is regulated by SA. (**A** and **B**)
Survival rate and fecundity of whiteflies on 21-day-old and 35-day-old *N.*
benthamiana, tobacco, and tomato plants. (**C**) Survival rate of thrips on 21-day-old
and 35-day-old *N. benthamiana*, tobacco, and tomato plants. (**D**) Fecundity of aphids
on *N. benthamiana*, tobacco, and tomato plants at 21-day and 35-day ages. (**E** and **F**)
SA and JA levels in *N. benthamiana* and tobacco at different ages. (**G**) Temporal
dynamics of SA content in wild-type and NahG *N. benthamiana* plants. (**H**) SA levels
in 21-day-old wild-type and NahG *N. benthamiana* plants upon whitefly attack. (**I**)
Survival rate and fecundity of whiteflies on 21-day-old and 35-day-old wild-type and
NahG *N. benthamiana* plants. (**J**) Survival rate and fecundity of whiteflies on SA-
treated (1 mM) 21-day-old wild-type *N. benthamiana* plants. (**K**) Survival rate and

fecundity of whiteflies on SA-treated (1 mM) 35-day-old NahG *N. benthamiana*

plants. Values are means $\pm$ SE, $n = 40$ for **A**, **B**, **C**, **D**, **I**, **J**, and **K**, $n = 7$ for **E**, **F**, and

G, $n = 8$ for **H**. Student's *t*-test (two-tailed) was used for significant difference

analysis. n.s., not significant. *$P < 0.05$, **$P < 0.01$.

1244

Fig. 2. ARR depends on the expression of the SA synthetic gene *NbPAL6*. (**A**) SA
synthesis pathways in plants. (**B**) Expression levels of genes in the ICS pathways in
21-day-old and 35-day-old *N. benthamiana* plants. (**C**) Expression levels of *NbPAL*
genes in 21-day-old and 35-day-old *N. benthamiana* plants. (**D**) Experimental design
of VIGS and gene overexpression in *N. benthamiana* plants. In the VIGS system, the

pTRV2 vector is injected into 21-day-old plants, activating the silencing vector to
inhibit target gene expression. Gene silencing typically became evident 14 days post-
injection, aligning with the timeline for ARR establishment in *N. benthamiana*,
making VIGS an effective tool for studying ARR-related gene functions. In
comparison to control plants (receiving only the pTRV2 vector), 35-day-old plants
subjected to gene silencing show reduced accumulation of target gene transcripts after
reaching 21 days. Additionally, *Agrobacterium*-mediated gene overexpression
enhances target gene mRNA levels in both 21-day-old and 35-day-old plants,
allowing for gene function assessment across different developmental stages. For
instance, when the pCAMBIA1305-GFP vector containing the target gene is
introduced at 19 days, mRNA levels increase in the treated 21-day-old plants within
two days. By comparing overexpressed plants to controls (those receiving the
pCAMBIA1305-GFP vector without the target gene), the impact of target genes on
resistance in 21-day-old plants can be assessed. This method was also used to evaluate
the role of gene overexpression in enhancing resistance in 35-day-old plants. (**E**) SA
levels in *NbPAL6*-silenced 35-day-old *N. benthamiana* plants. (**F**) Expression level of
NbPR1 in *NbPAL6*-silenced 35-day-old *N. benthamiana* plants. (**G**) Survival rate and
fecundity of whiteflies on *NbPAL6*-silenced 35-day-old *N. benthamiana* plants. (**H**)
Survival rate and fecundity of whiteflies on *NbPAL6*-overexpressed 21-day-old *N.*
benthamiana plants. Values are mean ± SE, $n = 12$ for **B**, **C**, and **F**; $n = 6$ for **E**; $n = 40$
for **G** and **H**. Student's *t*-test (two-tailed) was used for significant difference analysis.
n.s., not significant. *$P < 0.05$, **$P < 0.01$.

1272

**Fig. 3. NbMYB42 directly binds to the *NbPAL6* promoter to activate its
transcription.** (**A**) Yeast one-hybrid assay of the interaction between the *NbPAL6*
promoter and NbMYB42. Gold yeast cells co-transformed with pAbAi-*NbPAL6pro*
and pGADT7-*NbMYB42* plasmids were cultured on SD/-Leu medium with or without
150 ng ml⁻¹AbA for 3 days. The empty vector pGADT7 served as a negative control.

(**B**) Subcellular localization of NbMYB42 in *N. benthamiana*. NbMYB42-GFP (green

fluorescent protein) was transiently expressed in *N. benthamiana* line H2B-RFP, with

red fluorescent protein (RFP) as a nucleus marker and the empty *35S::GFP* vector

used as a control. Scale bar: 20 μm. (**C**) Analysis of the interaction between the

NbPAL6 promoter and NbMYB42 in *N. benthamiana* leaves using a GUS staining

assay. Diagrams on the left illustrate the reporter and effector vectors: *35S, CaMV 35S*

promoter; *EGFP*, enhanced green fluorescent protein; *GUS*, β-glucuronidase.

Representative photographs from several replicates are shown. (**D**) Analysis of the

interaction between the *NbPAL6* promoter and NbMYB42 in *N. benthamiana* leaves

using a dual-luciferase reporter assay. Diagrams on the left illustrate the reporter and

effector vectors. *REN, Renilla luciferase*; *LUC, firefly luciferase*. The activity of REN

was used as an internal control. (**E**) Schematic of the *NbPAL6* promoter, indicating

possible binding sites P1 to P5 for the TF NbMYB42. (**F**) ChIP-qPCR assay of the

enrichment of NbMYB42-GFP at the *NbPAL6* promoter with an anti-GFP antibody.

PCR primer pairs were designed based on the position shown in (**E**). Element P3

(GGTGGTTGTTGAGAGG) is significantly enriched in NbMYB42-GFP plants. (**G**)

EMSA results show NbMYB42-GST binding to the *NbPAL6* promoter *in vitro*. The

NbMYB42-GST protein was incubated with a biotin-labeled probe to assess binding

capacity. Labeled probes were incubated with and without GST protein as negative

controls, and 10-, 20-, and 50-fold excess unlabeled probes were used as cold

competitors. Biotin-mProbe was the labeled probe containing mutations in the P3

element of the *NbPAL6* promoter. (**H**) Expression level of *NbPAL6* in *NbMYB42*-

overexpressed 21-day-old *N. benthamiana* plants. (**I**) Survival rate and fecundity of

whiteflies on *NbMYB42*-overexpressed 21-day-old *N. benthamiana* plants. (**J**)

Expression level of *NbPAL6* in *NbMYB42*-silenced 35-day-old *N. benthamiana* plant.

(**K**) SA levels in *NbMYB42*-silenced 35-day-old *N. benthamiana* plants. (**L**)

Expression level of SA downstream gene *NbPR1* in *NbMYB42*-silenced 35-day-old *N.*

benthamiana plants. (**M**) Survival rate and fecundity of whiteflies on *NbMYB42*-

silenced 35-day-old *N. benthamiana* plants. Values are mean ± SE, $n = 4$ for **D** and **F**;

$n = 5$ for **H**; $n = 40$ for **I** and **M**; $n = 12$ for **J**; $n = 3$ for **K**; $n = 6$ for **L**. Student's *t*-test

(two-tailed) was used for significant difference analysis. n.s., not significant. $*P <$

0.05, $**P < 0.01$.

1310

Fig. 4. NbARF18La and NbARF18Lb activate *NbMYB42* expression. (A)
Expression levels of *NbMYB42* in 21-day-old and 35-day-old *N. benthamiana* plants.
(**B**) Yeast one-hybrid assay of the interaction between the *NbMYB42* promoter and
NbARF18La/b. Gold yeast cells co-transformed with pAbAi-*NbMYB42pro* and
pGADT7-*NbARF18La/b* plasmids were cultured on SD/-Leu medium with or without
150 ng ml⁻¹ AbA for 3 days. The empty vector pGADT7 was used as a negative
control. (**C**) Subcellular localization of NbARF18La and NbARF18Lb in *N.
benthamiana*. NbARF18La/b-GFP fusion proteins were transiently expressed in *N.
benthamiana* line H2B-RFP. Red fluorescent protein (RFP) served as a nucleus
marker, and the empty *35S*::*GFP* vector was used as a control. Scale bar: 20 μm. (**D**)
GUS staining assay analyzing the interaction between the *NbMYB42* promoter and
NbARF18La/b in *N. benthamiana* leaves. Diagrams on the top show the reporter and
effector vectors. (**E**) Dual-luciferase reporter assay analyzing the interaction between
the *NbMYB42* promoter and NbARF18La/b in *N. benthamiana* leaves. REN activity
was used as an internal control. (**F**) Top: Schematic of the *NbMYB42* promoter

showing the distribution of possible NbARF binding sites. Bottom: ChIP-qPCR assay

of the enrichment for NbARF18La/b-GFP protein at the *NbMYB42* promoter with an

anti-GFP antibody. The PCR primer pair is based on the positions shown above.

Element P2 (TGTCTC) is significantly enriched in NbARF18La/b-GFP plants. (**G**

and **H**) EMSA showing binding of NbARF18La-His-SUMO and NbARF18Lb-MBP

to the *NbMYB42* promoter *in vitro*. Proteins were incubated with a biotin-labeled

fragment of the *NbMYB42* promoter (P2, TGTCTC). The labeled probe was incubated

with and without His-SUMO or MBP protein as negative controls and 10-, 20- and

50-fold excess unlabeled probes were used as cold competitor probes. Biotin-mProbe

refers to the labeled probe with mutations. Values are mean $\pm$ SE, $n = 9$ for **A**; $n = 4$

for **E**; $n = 3$ for **F**. Student's *t*-test (two-tailed) was used for significant difference

analysis. n.s., not significant. **$P < 0.01$.

1338

Fig. 5. NbARF18La and NbARF18Lb participate in whitefly defense. (**A**)
Expression levels of *NbMYB42* and *NbPAL6* in *NbARF18La/18b*-overexpressed 21-
day-old *N. benthamiana* plants. (**B**) Survival rate and fecundity of whiteflies on
NbMYB18a/b-overexpressed 21-day-old *N. benthamiana* plants. (**C**) Expression levels
of *NbMYB42* and *NbPAL6* in *NbARF18La/b*-silenced 35-day-old *N. benthamiana*
plants. (**D**) SA levels in *NbMYB18La/b*-silenced 35-day-old *N. benthamiana* plants.
(**E**)Survival rate and fecundity of whiteflies on the *NbMYB18La/b*-silenced 35-day-
old *N. benthamiana* plants. (**F**) LUC activity was driven by the *NbPAL6* promoter
when co-expressed with different effectors. (**G**) Expression levels of *NbARF18La/b* in
21-day-old and 35-day-old *N. benthamiana* plants. Values are mean ± SE, n = 10 for
A and **C**; n = 40 for **B** and **E**; n = 3 for **D** and **F**; n = 12 for **G**. Student's *t*-test (two-
tailed) was used for significant difference analysis. n.s., not significant. *P < 0.05,
**P < 0.01.

1352
Fig. 6. Auxin suppresses NbARF18La/b-MYB42 mediated SA accumulation. (**A**)
IAA levels in 21-day-old and 35-day-old *N. benthamiana* plants. (**B**) Expression level
of *NbTAR2*, *NbYUC8*, and *NbIAA29* in 21-day-old and 35-day-old *N. benthamiana*
plants. (**C**, **D**) Survival rate and fecundity of whiteflies on IAA/NAA-treated (1, 25,
and 50 μM) 35-day-old *N. benthamiana* plants. (**E**) SA levels in IAA-treated (25 μM)
35-day-old *N. benthamiana* plants. (**F**) Expression level of *NbPR1* in IAA-treated (1,
25, and 50 μM) 35-day-old *N. benthamiana* plants. (**G**) Expression level of *NbPAL6*
in IAA-treated (1, 25, and 50 μM) 35-day-old *N. benthamiana* plants. (**H**) Top:
Schematic of the *GUS* reporter gene. Bottom: *GUS* reporter gene expression in mock-

and IAA-treated (25 μM) *pNbPAL6::GUS N. benthamiana* plants. (**I**) Top: Schematic

of the *GUS* and *LUC* reporter genes and effector NbMYB42. Bottom: GUS / LUC

activity co-expressed with *NbMYB42* in mock- or IAA-treated (25 μM)

pNbPAL6::GUS N. benthamiana plants. (**J**) Expression level of *NbMYB42* in IAA-

treated (1, 25, and 50 μM) 35-day-old *N. benthamiana* plants. (**K**) Top: Schematic of

the *GUS* reporter genes. Bottom: *GUS* reporter gene expression in mock- and IAA-

treated (25 μM) *pNbMYB42::GUS N. benthamiana* plants. (**L**) Top: Schematic of the

GUS and *LUC* reporter gene and effector NbARF18La. Bottom, GUS / LUC activity

co-expressed with *NbARF18La* in mock- and IAA-treated (25 μM) *pNbMYB42::GUS*

N. benthamiana plants. (**M**) Top: Schematic of the *GUS* and *LUC* reporter gene and

effector NbARF18Lb. Bottom: GUS / LUC activity co-expressed with *NbARF18Lb* in

mock- and IAA-treated (25 μM) *pNbMYB42::GUS N. benthamiana* plants. (**N**)

Expression level of *NbARF18La/b* in IAA-treated (1, 25, and 50 μM) 35-day-old *N.*

benthamiana plants. (**O**) Top: Schematic of the *GUS* reporter gene. Bottom: *GUS*

reporter gene expression in mock- and IAA-treated (25 μM) *pNbARF18La/b::GUS N.*

benthamiana plants. Values are mean ± SE, $n = 3$ for **A** and **E**; $n = 12$ for **B**; $n = 15$ for

C and **D**; $n = 5$ for **F**; $n = 4$ for **G**, **J** and **N**; $n = 6$ for **H**, **K**, and **O** (*GUS* normalized

gene expression), $n = 3$ for **H** and **K** (Relative LUC activity); $n = 6$ for **I**, **L**, and **M**

(Relative LUC activity). Student's *t*-test (two-tailed) was used for significant

difference analysis. n.s., not significant. *$P < 0.05$, **$P < 0.01$.

Fig. 7. Auxin-induced microRNA160c targets *NbARF18La/b* transcripts. (A)
*Nb*miR160c targets *N. benthamiana NbARF18La* mRNA. CDS: Coding sequence;
UTR: Untranslated regions; ATG: Initiation codon; TAA: Termination codon. (**B**)
*Nb*miR160c targets *N. benthamiana NbARF18Lb* mRNA. (**C**) Expression of
NbARF18La/b in *Nb*miR160c-expressing 35-day-old *N. benthamiana* plants. (**D**)
Target plots (t-plots) of *Nb*miR160c and its target *NbARF18La* mRNA, validated via
degradome sequencing. The red dot indicates the cleavage site of the *NbARF18La*
transcript at 1572 bp, as analyzed by GSTAr (1.0) cleavage site prediction software.
(**E**) Target plots (t-plots) of *Nb*miR160c and its target *NbARF18Lb* mRNA, showing
cleavage at 2099 bp. (**F**, **G**) Camera (Left) and confocal (Right) images of GFP-
NbARF18La/b target site and GFP-mutant site expression in *N. benthamiana* when

*Nb*miR160c was overexpressed. Top: GFP; Middle, Bright-field; Bottom: GFP/bright-

field overlay. Scale bars, 40 μm. (**H**) GUS phenotype observed by histochemical

staining showing GFP-*NbARF18La*/*b* target site and GFP-mutant site expression in *N.*

benthamiana with overexpressed *Nb*miR160c. (**I**) LUC activity displaying LUC-

NbARF18Lb target site and LUC-mutant site expression. Values are mean ± SE, $n =$

10 for **C**; $n = 3$ for **I**. Student's *t*-test (two-tailed) was used for significant difference

analysis in **C**. **$P < 0.01$. One-way ANOVA followed by Fisher's least significant

difference (LSD) test was used for significant difference analysis in **I**. Bars with

different lowercase letters indicate significant differences between treatments at $P <$

0.05.

1404

Fig. 8. Auxin-induced microRNA160c negatively regulates plant defense. (A)
Expression of *NbMYB42* and *NbPAL6* in *Nb*miR160c-expressing 35-day-old *N. benthamiana* plants. (**B**) SA levels in *Nb*miR160c-expressing 35-day-old *N. benthamiana* plants. (**C**) Survival rate and fecundity of whiteflies on *Nb*miR160c-expressing 35-day-old *N. benthamiana* plants. (**D**) Expression of *NbARF18La/b* in mimic160c-expressing 21-day-old *N. benthamiana* plants. (**E**) Expression of *NbMYB42* and *NbPAL6* in mimic160c-expressing 21-day-old *N. benthamiana* plants.
(**F**) Survival rate and fecundity of whiteflies on the mimic160c-expressed 21-day-old *N. benthamiana* plant. (**G**) Left: Schematic diagram showing effector and reporter vectors used in transient transcriptional activity assays in *N. benthamiana* leaves. Right: The LUC activity driven by the *NbMYB42* promoter when coexpressed with different effectors. (**H**) Expression level of *Nb*miR160c in 21-day-old and 35-day-old *N. benthamiana* plants. (**I**) *Nb*miR160c levels in IAA-treated (1, 25, and 50 μM) 35-day-old plants. (**J**) Top: Schematic of the *GUS* reporter gene; Bottom: *GUS* reporter gene expression and activity in mock- and IAA-treated (25 μM) *pNbmiR160c::GUS* *N. benthamiana* plants. (**K**) Top: Schematic of the *LUC* reporter gene; Bottom: *LUC*

reporter gene expression and activity in mock- and IAA-treated (25 μM)

pNbmiR160c::LUC N. benthamiana plants. Values are mean ± SE; $n = 10$ for **A**; $n = 3$

for **B** and **I**; $n = 15$ for **C** and **F**; $n = 8$ for **D**; $n = 5$ for **E**; $n = 6$ for **H** and **K**; $n = 4$ for

G and **J**. Student's t-test (two-tailed) was used for significant difference analysis in **A**,

B, **C**, **D**, **E**, **F**, **H**, **I**, **J**, and **K**. $*P < 0.05$, $**P < 0.01$. One-way ANOVA followed by

Fisher's least significant difference (LSD) test was used for significant difference

analysis in **G**. Bars with different lowercase letters indicate significant differences

between treatments at $P < 0.05$.

1429

Fig. 9. Excessive SA impacts early plant growth by antagonizing auxin. (**A**)

Photographs of 35-day-old *N. benthamiana* plants treated with SA starting at 21 days

old. (**B**, **C**) Aboveground fresh weight (**B**) and plant height (**C**) of SA-treated (1 mM)

N. benthamiana plants. (**D**) Expression levels of auxin synthesis genes *NbTAR2* and

NbYUC8 in 21-day-old *N. benthamiana* plants treated with 1 mM SA. (**E**) Auxin

levels in 21-day-old *N. benthamiana* plants after successive SA treatment (1 mM). (**F**)

Expression levels of auxin downstream gene *NbIAA29* in SA-treated (1 mM) 21-day-

old *N. benthamiana* plants. (**G**) Expression of auxin pathway genes in 35-day-old wild

type and NahG *N. benthamiana* plants. Values are mean ± SE, $n = 12$ for **B** and **C**; $n =$

8 for **D** and **F**; $n = 3$ for **E**; $n = 6$ for **G**. Student's *t*-test (two-tailed) was used for

significant difference analysis. *P < 0.05, **P < 0.01.

1441
Fig. 10. Auxin-SA seesaw mediates the age-related balance between plant growth
and herbivore defense. At juvenile stages, plants exhibit high auxin levels,
promoting the expression of *Nb*miRNA160c. This miRNA targets *NbARF18La/b*
transcripts, leading to their degradation via the plant ARGONAUTE1 (AGO1)
protein. As a result, NbARF18La/b is degraded at an early age, inhibiting downstream
expression of *NbMYB42* and *NbPAL6*. This repression of the NbARF18La/b-
NbMYB42-NbPAL6 module limits anti-whitefly SA accumulation in juvenile plants.
As plants mature, the auxin levels decrease, lifting the repression of NbARF18La/b.
These two interacting ARF TFs directly bind to the promoter of the TF *NbMYB42*,
activating its expression. NbMYB42 subsequently binds to the promoter of *NbPAL6*,
enhancing its expression. NbPAL6 promotes SA biosynthesis, boosting SA-mediated
defense against phloem-feeding insect whiteflies. The age-related rise in SA levels
determines the ARR of plants. This model illustrates a seesaw-like mechanism for
balancing growth and herbivore defense, with auxin dominance in early stages and
SA-driven defense taking precedence as the plant ages.

Supplementary Materials

This PDF file includes:

Supplementary Text

Figs. S1 to S10

Other Supplementary Material for this manuscript includes the following:

Tables S1 to S2

1463

1464 **Supplementary Figures**

1465

Fig. S1. Expression of the *NbPR1* gene downstream of SA at different ages and
SA levels. (**A**) Expression level of the SA downstream gene *NbPR1* in the *N.*
benthamiana plants at 21 and 35 days old. (**B**) Expression level of *NbPR1* in 35-day-
old NahG *N. benthamiana* plant. (**C**) Expression level of *NbPR1* in 21-day-old *N.*
benthamiana plants treated with 1 mM SA. Values are mean ± SE, $n = 10$ for **A** $n = 6$
for **B**; $n = 8$ for **C**. Student's *t*-test (two-tailed) was used for significant difference
analysis. **$P < 0.01$.

1473

1474 **Fig. S2. Phylogenetic tree of PAL protein.** Amino acid sequences of PAL proteins
1475 from *N. benthamiana* (Nb), *N. tabacum* (Nt), *N. attenuata* (Na), *Arabidopsis thaliana*
1476 (At), *Oryza sativa* (Os), *Solanum tuberosum* (St), *Vitis vinifera* (Vv), and *Camellia*
1477 *sinensis* (Cs) were aligned by ClustalW. The Maximum Likelihood tree was
1478 constructed with MEGA X with 1000 bootstrap replicates.

1479

Fig. S3. Gene expression level after silencing and overexpressing *NbPAL6*. (**A**)

Expression level of *NbPAL6* in *NbPAL6*-silenced 35-day-old *N. benthamiana* plants.

(**B**) Expression level of *NbPAL6* in *NbPAL6*-overexpressed 21-day-old *N.*

benthamiana plants. (**C**) Expression level of SA downstream *NbPR1* gene in *NbPAL6*-

overexpressed 21-day-old *N. benthamiana* plants. Values are mean ± SE, $n = 12$ for **A**

$n = 7$ for **B**; $n = 5$ for **C**. Student's *t*-test (two-tailed) was used for significant

difference analysis. **P < 0.01.

1487

Fig. S4. Gene expression level after silencing and overexpressing *NbMYB42*. (A)
Expression level of *NbMYB42* in the *NbMYB42*-overexpressed 21-day-old *N.*
benthamiana plants. (**B**) Expression level of SA downstream *NbPR1* gene in the
NbMYB42-overexpressed 21-day-old *N. benthamiana* plants. (**C**) Expression level of
NbMYB42 in *NbMYB42*-silenced 35-day-old *N. benthamiana* plants. Values are mean
± SE, $n = 8$ for **A**; $n = 6$ for **B**; $n = 12$ for **C**. Student's *t*-test (two-tailed) was used for
significant difference analysis. **$P < 0.01$.

1495

Fig. S5. Phylogenetic relationship and conserved DNA binding domain of

NbARF18La/b. (**A**) Phylogenetic tree of ARF proteins. Homologous genes of

NbARF18Ls were identified using NCBI Blast

(https://blast.ncbi.nlm.nih.gov/Blast.cgi). Protein sequences were aligned using

ClustalW, and a phylogenetic tree with 1000 bootstrap replicates was constructed by

the Maximum Likelihood method in MEGA X. (**B**) Multi-sequence alignment

analysis of ARF proteins. The B3 DNA binding domain (DBD) and C-terminal

dimerization domain (CTD) regions are underlined. Protein sequence alignment

results were visualized by GenDoc.

1505

Fig. S6. NbARF18La interacts with NbARF18Lb. (**A**) Yeast two-hybrid (Y2H)
assay between NbARF18La and NbARF18Lb. Yeast cells co-transformed with
NbARF18La and NbARF18Lb plasmids grow on a selective medium (SD/-W-L-H-A)
for 3 days, indicating an interaction between NbARF18La and NbARF18Lb.
Additionally, yeast cells co-transformed with NbARF18La plasmids (pGADT7-
NbARF18La and pGBKT7-NbARF18La) also grew on selective medium (SD/-W-L-
H-A), showing that NbARF18La exhibits self-interaction. (**B**) Bimolecular
fluorescence complementation (BiFC) assays confirmed the interaction between
NbARF18La and NbARF18Lb in *N. benthamiana* epidermal cells. BiFC assays show
that NbARF18La interacts with NbARF18Lb in nuclei, and NbARF18La interacts
with itself in nuclei. The BiFC signal (green fluorescence) observed in the nuclei
indicates an interaction between NbARF18La and NbARF18Lb, as well as a self-
interaction of NbARF18La. Confocal microscopy was used to visualize the
fluorescence. Scale bars, 40 μm. (**C**) The interaction between NbARF18La and
NbARF18Lb was further validated by an *in vitro* pull-down assay. Recombinant
NbARF18Lb-MBP and NbARF18La-His-SUMO proteins were incubated with MBP
beads in a pull-down binding buffer. The pulled-down proteins were detected by
immunoblotting using anti-MBP and anti-His antibodies. A mixture of MBP and
NbARF18La-His-SUMO was used as a negative control.

1525

Fig. S7. Gene expression levels after overexpressing and silencing *NbARF18La/b*.
(**A**, **B**) Expression levels of *NbARF18La/b* in the *NbARF18La/b*-overexpressed 21-
day-old *N. benthamiana* plants. (**C**) Expression level of SA downstream gene *NbPR1*
in the *NbARF18La/b*-overexpressed 21-day-old *N. benthamiana* plants. (**D**, **E**)
Expression levels of *NbARF18La* and *NbARF18Lb* in the *NbARF18La/b*-silenced 35-
day-old *N. benthamiana* plants. (**F**) Expression level of SA downstream gene *NbPR1*
in the *NbARF18La/b*-silenced 35-day-old *N. benthamiana* plants. Values are mean ±
SE, $n = 6$ for **A**, **B** and **D**; $n = 9$ for **C**; $n = 10$ for **E** and **F**. Student's *t*-test (two-tailed)
was used for significant difference analysis. **$P < 0.01$.

1535

Fig. S8. Auxin synthesis inhibitors enhance the NbARF18La/b-MYB42 module in juvenile plants. (**A**) Survival rate and fecundity of whiteflies on *N. benthamiana* plants treated with L-Kyn and Yucasin (50 μM) at 21 days old. (**B**) SA levels in *N. benthamiana* plants treated with the IAA inhibitors L-Kyn and Yucasin (50 μM) at 21 days old. (**C**) Expression level of SA downstream gene *NbPR1* in L-Kyn- and Yucasin-treated (50 μM) 21-day-old *N. benthamiana* plants. (**D**) Expression level of *NbPAL6* in L-Kyn- and Yucasin-treated (50 μM) 21-day-old *N. benthamiana* plants. (**E**) Expression level of *NbMYB42* in L-Kyn- and Yucasin-treated (50 μM) 21-day-old *N. benthamiana* plants. (**F**) Expression levels of *NbARF18La/b* in the L-Kyn- and Yucasin-treated (50 μM) 21-day-old *N. benthamiana* plant. Values are mean ± SE, *n* = 15 for **A** and **B**; *n* = 5 for **C**, **D**, **E**, and **F**. Student's *t*-test (two-tailed) was used for significant difference analysis. n.s., not significant; *$P < 0.05$, **$P < 0.01$.

1548

Fig. S9. Overexpression of mimicRNA160c increased the salicylic acid level in
juvenile plants. (**A**) Schematic diagram of the *Nb*miR160c expression cassette *p35S-*
*Nb*miR160c. This vector was used to overexpress *Nb*miR160c in *N. benthamiana*
plants through *Agrobacterium*-mediated transformation. The blue box represents the
miRNA precursor, and the red box represents *Nb*miR160c. HygR: Hygromycin B
phosphotransferase; NOS: Nos Terminator; RB: Right border; LB: Left border. (**B**)
Expression level of *Nb*miR160c in the *Nb*miR160c-expressed 35-day-old plants. (**C**)
Sequence alignment of *Nb*miR160c with the predicted target sites in *NbARF18La* and
NbARF18Lb, along with their mutated versions (*NbARF18La*-m and *NbARF18Lb*-m)
used in subsequent experiments. Colons represent G: U pairs. (**D**) Schematic diagram
of the GFP/GUS/LUC sensor carrying the *Nb*miR160c target site and the mutant
target site. TGA: Termination codon; UTR: Untranslated regions. (**E**) Expression level
of SA downstream gene *NbPR1* in the *Nb*miR160c-expressed 35-day-old plants. (**F**)

1562   Design of mimicry sequences (mimic160c) for *Nb*miR160c. (**G**) SA levels in

1563   mimic160c-expressed 21-day-old plants. (**H**) Expression level of SA downstream

1564   gene *NbPR1* in the mimic160c-expressed 21-day-old plants. Values are mean ± SE, *n*

1565   = 4 for **B**; *n* = 8 for **E**; *n* = 3 for **G**; *n* = 6 for **H**. Student's *t*-test (two-tailed) was used

1566   for significant difference analysis. **P < 0.01.

1567

Fig. S10. *Nb*miR160c affects the transcription of *NbMYB42* and *NbPAL6*. (A to

E) Schematic diagram showing the effector and reporter vectors used in transient

transcriptional activity assays in *N. benthamiana* leaves. The data show the relative

LUC activity normalized to REN activity. Values are mean ± SE, $n = 3$. Student's *t*-

test (two-tailed) was used for significant difference analysis in **A**, **B**, **C**, and **D**. *$P <$

0.05, **$P < 0.01$. One-way ANOVA followed by Fisher's least significant difference

(LSD) test was used for significant difference analysis in **E**. Bars with different

lowercase letters indicate significant differences between treatments at $P < 0.05$.

Supplementary Tables

Table S1. Basic information on identifying *NbPAL* genes.

Gene Name	Gene ID	CDS (bp)	Protein Length (aa)	Molecular Weight (Da)
NbPAL1	NbL01g24700.1	2040	679	73888.46
NbPAL2	NbL05g03880.1	2154	717	78005.01
NbPAL3	NbL10g04670.1	2154	717	78248.28
NbPAL4	NbL10g08440.1	2166	721	78734.94
NbPAL5	NbL12g19650.1	2154	717	78393.48
NbPAL6	NbL13g11330.1	2154	717	78233.31
NbPAL7	NbL14g10160.1	2139	712	77447.38
NbPAL8	NbL15g02170.1	2139	712	77438.39
NbPAL9	NbL15g24630.1	2154	717	78151.22

1578

1579 **Table S2.** List of primers used in this study.

Primer name	Sequence (5'-3')
q-mir160c-F	TGCCTGGCTCCCTGTATTTT
q-NbU6-F	GGAACGATACAGAGAAGATTAGCA
q-NbACTIN-F	CAATCCAGACACTGTACTTTCTCTC
q-NbACTIN-R	AAGCTGCAGGTATCCATGAGACTA
q-NbPR1-F	TGCAAATCAGAGAGCTGGGG
q-NbPR1-R	CACTTTTCCCGGTGCACAAG
q-NbICS1-F	CGTGCTTATGGGGCAATTCG
q-NbICS1-R	ATTGCCATCTGGTACGTGCA
q-NbICS2-F	CTCGCGAGTACTCATGTCCC
q-NbICS2-R	ATCAATGTCCGCAGCTGTCA
q-NbEPS1-F	GCCGGCGTAGAATTCACCTA
q-NbEPS1-R	GCCGACGTTATTGAGTGGGA
q-NbCBP60g-F	CGACCTTGGCAAGCAGTTTC
q-NbCBP60g-R	GGAGAAACTGGCTGGATGCT
q-NbSARD1-F	AGTTGGCCACTGTTGACACA
q-NbSARD1-R	AGTGTAGTTTGCTCCGGTGG
q-NbEDS5-F	CAAACACTGCTGCTGCACAT
q-NbEDS5-R	AGGGACTTCAGCAACATCCG
q-NbPBS3-F	AACCCCGAAATTACCGACCC
q-NbPBS3-R	AGCACCCGTGACAATTACGT
q-NbEPS1-F	GCCGGCGTAGAATTCACCTA
q-NbEPS1-R	GCCGACGTTATTGAGTGGGA
q-NbMYB42-F	GGTGAGAGCTCCTTGTTGTGA
q-NbMYB42-R	TCTCCAATTGCCATGGCCAT
q-NbPAL4-F	ATGCACTCAAGGGTAGCCAC
q-NbPAL4-R	CAGCTCCACCATCACCTCTG
q-NbPAL6-F	AGGAACAAGGCATTGCATGG
q-NbPAL6-R	TTTCCAATGGAGGCAAGTGC
q-NbPAL7-F	ACAATCAGGGACAAGGGCAG
q-NbPAL7-R	GACCTGTGAGTAAACCGGCA
q-NbPAL9-F	TGTTTACCTCTCCGTGGCAC
q-NbPAL9-R	CAGCATTAAGGGCCTCACCA
q-NbARF18La-R	TGATCCCATCTACTGGCCCA
q-NbARF18La-F	GAGACCAGTTCGACAAGCCA
q-NbARF18Lb-F	CAACTCGTGCTTTTCGGCAA
q-NbARF18Lb-R	AGTGCAGAACCTGAGCCATC
q-NbTAR2-F	GCTGAGACTGGAAACCGACA
q-NbTAR2-R	CGAATAAAATGGGGCAGCGG
q-NbYUC8-F	GCAAATGGCGCTGAAGTTGA
q-NbYUC8-R	TGCCAGAATTTCCACAGCCA

Primer name	Sequence (5'-3')
q-NbIAA29-F	CGTTTAAGGCTTGCGACGAG
q-NbIAA29-R	CCAATGGCCACTCCTTCCAT
pBI121-pNbPAL6-F	GACCATGATTACGCCAAGCTTGAATTCGTTCGGAAATTATGA
pBI121-pNbPAL6-R	<u>ACCACCCGGGGATCCTCTAGA</u>AAATGGAAAATATGAAAGAAAAAGA
pBI121-pNbARF18La-F	<u>GACCATGATTACGCCAAGCTT</u>CGCATGCAATCAAATATTATTACTCT
pBI121-pNbARF18La-R	<u>ACCACCCGGGGATCCTCTAGA</u>TTGCCTTATCCAGTACACTTCT
pBI121-pNbARF18Lb-F	GACCATGATTACGCCAAGCTTGAGGTGAGGATTGCCCAAGC
pBI121-pNbARF18Lb-R	<u>GGACTGACCACCCGGGGATCC</u>CTTGATTTCAACAGAATTTTACAGA
pBI121-pNbMYB42-F	GCTATGACCATGATTACGCCAAGCTTTAGTGATTTGACTGTGCGTGA
pBI121-pNbMYB42-R	GACTGACCACCCGGGGATCCTCTAGATTTTGCTTTAATCACTTGCT
pBI121-pmir160-F	AGCTATGACCATGATTACGCCAAGCTTAATTTGACTATTACATAATTCCT
pBI121-pmir160-R	GGACTGACCACCCGGGGATCCCGAGGAAAATACGAAGTGAA
TRV-NbARF18La-F	GC<u>TCTAGA</u>ACCCAGTTTCCAGCGAGTTG
TRV-NbARF18La-R	CG<u>GGATCC</u>TCTTGAAGACCTTTCTGCTCA
TRV-NbARF18Lb-F	GC<u>TCTAGA</u>AATCCAGCTGGTGGGAACTG
TRV-NbARF18Lb-R	CG<u>GGATCC</u>AAAAGGCATCTTGAACCTCATCC
TRV-NbPAL6-F	GC<u>TCTAGA</u>TCTATGGTTCTATTTGAGGCT
TRV-NbPAL6-R	CG<u>GGATCC</u>GGCCCTAATGACCTCAATTTGA
TRV-NbMYB42-F	GAGTAAGGTTACCGAATTCTCTAGAAAAGACACTCCAAAAACAATCTTGA
TRV-NbMYB42-R	<u>GTGAGCTCGGTACCGGATCC</u>TTGGTCATCGATTGTAATAGTCGT
1305-NbARF18La-F	<u>TAAGTCCGGAGCTAGCTCTAGA</u>ATGGAGGAGGTTATGGAGAAATGT
1305-NbARF18La-R	CGGTCCTCGAGACGTCTCTAGATGCAAACATGCTAAGTGGTCC
1305-NbARF18Lb-F	<u>TAAGTCCGGAGCTAGCTCTAGA</u>ATGATTACTTTTATGGATCCAAAGGAC
1305-NbARF18Lb-R	CGGTCCTCGAGACGTCTCTAGACTCCCTGAGTCCAACATTATCG
1305-NbPAL6-F	GC<u>TCTAGA</u>ATGGAGTATGCAAATGGAGATTGT

Primer name	Sequence (5'-3')
1305-NbPAL6-R	CG<u>GGATCC</u>GCAGAGTGGAAGAGGAGC
1305-NbMYB42-F	GC<u>TCTAGA</u>ATGGTGAGAGCTCCTTGTTGT
1305-NbMYB42-R	CG<u>GGATCC</u>AAATTCCGGTAAATCTAGT
pABAi-pNbMYB42-F	AAAAAAAATGATGAATTGAAAAGCTTTAGTGATTTGACTGTGCGTGA
pABAi-pNbMYB42-R	TCGAGGTCGACAGATCCCCGGGTACCTTTTGCTTTAATCACTTGCT
pABAi-pNbPAL6-F	AAAAAAAATGATGAATTGAAAAGCTTGAATTCGTTCGGAAATTATGA
pABAi-pNbPAL6-R	TCGAGGTCGACAGATCCCCGGGTACCAAATGGAAAATATGAAAGAAAAAGA
pGAD-NbARF18La-F	CCGGGTGGGCATCGATACGGGATCCATGGAGGAGGTTATGGAGAAATGT
pGAD-NbARF18La-R	TATCTACGATTCATCTGCAGCTCGAGCTATGCAAACATGCTAAGTGGTCC
pGAD-NbARF18Lb-F	<u>CCGGGTGGGCATCGATACGGGATCC</u>ATGATTACTTTTATGGATCCAAAGGAC
pGAD-NbARF18Lb-R	TATCTACGATTCATCTGCAGCTCGAGTTACTCCCTGAGTCCAACAT
pGAD-NbMYB42-F	CG<u>GGATCC</u>ATGGTGAGAGCTCCTTGTTGT
pGAD-NbMYB42-R	CC<u>CTCGAG</u>TCAAAATTCCGGTAAATCTAGT
pGBK-NbARF18La-F	<u>TGGCCATGGAGGCCGAATTC</u>ATGGAGGAGGTTATGGAGAAATGT
pGBK-NbARF18La-R	<u>CGCTGCAGGTCGACGGATCC</u>CTATGCAAACATGCTAAGTGGTCC
pGBK-NbARF18Lb-F	TGGCCATGGAGGCCGAATTCATGATTACTTTTATGGATCC
pGBK-NbARF18Lb-R	CGCTGCAGGTCGACGGATCCCTACTCCCTGAGTCCAACAT
pGreen0800-pNbPAL6-F	TCGAGGTCGACGGTATCGATAAGCTTGAATTCGTTCGGAAATTATGA
pGreen0800-pNbPAL6-R	<u>GGCCGCTCTAGAACTAGTGGATCC</u>AAATGGAAAATATGAAAGAAAAAGA
pGreen0800-pNbMYB42-F	TCGAGGTCGACGGTATCGATAAGCTTTAGTGATTTGACTGTGCGTGA
pGreen0800-pNbMYB42-R	GCGGCCGCTCTAGAACTAGTGGATCCTTTTGCTTTAATCACTTGCT
pGreen0800-pmir160-F	ACTCACTATAGGGCGAATTGGGTACCAATTTGACTATTACATAATTCCT
pGreen0800-pmir160-R	CGGCCGCTCTAGAACTAGTGGATCCCGAGGAAAATACGAAGTGAA
pET28a-SUMO-NbARF18La-F	<u>GCTCACAGAGAACAGATTGGTGGA</u>ATGGAGGAGGTTATGGAGAAATGTG

Primer name	Sequence (5'-3')
pET28a-SUMO-NbARF18La-R	TGCCGCCACTACCACCACCACTACCACCGCCTTTTTCAAACTGCGGATGGCTCCATGCAAACATGCTAAGTGGTCCTGC
pMAL-NbARF18Lb-F	<u>TCCAGGGGCCCCTGGGATCC</u>ATGATTACTTTTATGGATCC
pMAL-NbARF18Lb-R	CTGCCGCCACTACCACCACCACTACCACCGCCTTTTTCAAACTGCGGATGGCTCCACTCCCTGAGTCCAACATTATCGC
pGEX-NbMYB42-F	CTGGAAGTTCTGTTCCAGGGGCCCGGCTCCATGGTGAGAGCTCCTTGTTGTGA
pGEX-NbMYB42-R	CACTACCACCACCACTACCACCGCCTTTTTCAAACTGCGGATGGCTCCAAAATTCCGGTAAATCTAGTAAGTCCCCG
1300-mir160-F	GG<u>GGTACC</u>CAGCAGCAGCCACAGCAAAA
1300-mir160-R	CG<u>GGATCC</u>GCTGCTGATGCTGATGCCAT
1300-mimic160-F	GGTACCTGGCCATCCCCTA
1300-mimic160-R	GGATCCCGGAAGCAAATTTACA
Bifc-NbARF18La-F	CCATTTACGAACGATAGTTAATTAACATGGAGGAGGTTATGGAGAAATGT
Bifc-NbARF18La-R	CCACCACTGCCACCTCCTCCACTAGTTGCAAACATGCTAAGTGGTCC
Bifc-NbARF18Lb-F	<u>CCATTTACGAACGATAGTTAATTAAC</u>ATGATTACTTTTATGGATCCAAAGGAC
Bifc-NbARF18Lb-R	CCACCACTGCCACCTCCTCCACTAGTCTCCCTGAGTCCAACATTATCG
35S-GFP-mir160-F	AGGACCGGTCCCGGGGGATCCATGGTGAGCAAGGGCGAG
GFP-NbARF18La-site-R	GGGCGGCCGCTTTAAGATCTATGCCTGGCTCTCTGTATGC
GFP-NbARF18La-mutant site-R	GGGCGGCCGCTTTAAGATCTATGTCTCGCGCTCTCTAGGC
GFP-NbARF18Lb-site-R	GGGCGGCCGCTTTAAGATCTATGCCTGGCTCCCTGCAT
GFP-NbARF18Lb-mutant site-R	GGGCGGCCGCTTTAAGATCTATGTCTTGCGCCTTGAATGC
35S-GUS-mir160-F	GGAGAGAACACGGGGGACTCTAGAATGTTACGTCCTGTAGAAACCCC
GUS-NbARF18La-site-R	AACGATCGGGGAAATTCGAGCTCTCAATGCCTGGCTCTCTGTATGCC
GUS-NbARF18La-mutant site-R	AACGATCGGGGAAATTCGAGCTCTCAATGTCTCGCGCTCTCTAGGCC
GUS-NbARF18Lb-site-R	AACGATCGGGGAAATTCGAGCTCTCAATGCCTGGCTCCCTGCATGCC
GUS-NbARF18Lb-mutant site-R	AACGATCGGGGAAATTCGAGCTCTCAATGTCTTGCGCCTTGAATGCC
35S-LUC-mir160-F	CATGCCATGGAAGACGCCAA
LUC-NbARF18La-site-R	GCTCTAGAATGCCTGGCTCTCTGTATGCCATTACACGGCGATCTTTCCG

Primer name	Sequence (5'-3')
LUC-NbARF18La-mutant site-R	GCTCTAGAATGTCTCGCGCTCTCTAGGCCATTACACGGCGAT CTTTCCG
LUC-NbARF18Lb-site-R	GCTCTAGAATGCCTGGCTCCCTGCATGCCATTACACGGCGAT CTTTCCG
LUC-NbARF18Lb-mutant site-R	GCTCTAGAATGTCTTGCGCCTTGAATGCCATTACACGGCGAT CTTTCCG
ChIP-pNbMYB42-P1-F	AGTGATTTGACTGTGCGTGA
ChIP-pNbMYB42-P1-R	AAAACGGCACAAATGGTCCG
ChIP-pNbMYB42-P2-F	AACTTGAGAACCCACGACCC
ChIP-pNbMYB42-P2-R	CCTGCATGGAAATTCATGAAATCT
ChIP-pNbMYB42-P3-F	ACTCCTTCAGATTTTGACGTGA
ChIP-pNbMYB42-P3-R	ACCTTATTTACGAGGATTACACTGAGT
ChIP-pNbPAL6-P1-F	TCGGTCTCAACGCGTTAATGA
ChIP-pNbPAL6-P1-R	ACTTTCCCTTCACCAACATGA
ChIP-pNbPAL6-P2-F	ATGACGTGGTGCTGTCGTTG
ChIP-pNbPAL6-P2-R	TTTCCCTCGTAAAATACCGCA
ChIP-pNbPAL6-P3-F	CGCCAGTAAGTTATTATCGCGC
ChIP-pNbPAL6-P3-R	CGTTGGATTGCACATGGAGA
ChIP-pNbPAL6-P4-F	TCTCCATGTGCAATCCAACG
ChIP-pNbPAL6-P4-R	TGTTGAGCCAAAATGTGAGTGT
ChIP-pNbPAL6-P5-F	TGGCTCAACAAACTCAGGAAATT
ChIP-pNbPAL6-P5-R	AATGTGTGTTGCGAACTTCTC
EMSA-pPAL6-F	GTGAGGGTTGAAGGGTGGTTGTTGAGAGGAAGCCCGCTT ACTTTTTTTGG
EMSA-pPAL6-R	GGTTTTTTTCATTCGCCCGAAGGAGAGTTGTTGGTGGGAA GTTGGGAGTG
EMSA-pPAL6-mF	GTGAGGGTTGAAGAAAAAAAAAAAAAAAAAAAAGCCCGCTT ACTTTTTTTGG
EMSA-pPAL6-mR	GGTTTTTTTCATTCGCCCGAAAAAAAAAAAAAAAAAAAAGAA GTTGGGAGTG
EMSA-pMYB42-F	GGACAATTTTTCTTTATTTTCTTTTTTGTCTCCAGTAAAATAA CG
EMSA-pMYB42-R	GCAATAAAATGACCTCTGTTTTTTCTTTTATTTCTTTTTAACA GG
EMSA-pMYB42-mF	GGACAATTTTTCTTTATTTTCTTTTTAAAAAACAGTAAAATA ACG
EMSA-pMYB42-mR	GCAATAAAATGACAAAAAATTTTTCTTTTATTTCTTTTTAACA GG

1580