1
2  **Increased dietary levels of folic acid reduce survival and alter climbing behaviors 24 hours**
3  **after hypoxia in male and female *Drosophila melanogaster***
4

5  Siddarth Gunnala MBS[1], Alisha Harrison BS[1], Amber Juba BA[1], Paula Ashcraft BSc[2], Meher
6  Garg[2], Teodoro Bottiglieri PhD[3], Lori M Buhlman PhD[1,4], Nafisa M. Jadavji[2,5,6]
7
8
**Affiliations**
[1] Biomedical Sciences Program, College of Graduate Studies, Midwestern University, Glendale,
AZ, USA
[2] Division of Integrative and Molecular Physiology, Department of Biomedical Sciences,  School
of Medicine, Southern Illinois University, Carbondale, IL, USA
[3] Center of Metabolomics, Institute of Metabolic Disease, Baylor Scott & White Research
Institute, Dallas, TX, USA
[4] College of Veterinary Medicine, Midwestern University, Glendale, AZ, USA
[5]Department of Child Health, College of Medicine Phoenix, University of Arizona, Phoenix, AZ,
USA
[6]Department of Neuroscience, Carleton University, Ottawa, Canada
20
21
**Corresponding Author:**
Nafisa M. Jadavji, PhD, FAHA
Department of Physiology
SIU School of Medicine
Life Science II, room 2076
1135 Lincoln Drive
Mail Code 6512
Carbondale, IL 62901
nafisa.jadavji@mail.mcgill.ca
31

**Abstract**

Hypoxia is a major component of ischemic stroke. The prevalence of ischemic stroke is

expected to increase as the global population ages and risk factors, like obesity, are on the rise.

Nutrition is a modifiable risk factor for ischemic stroke. Increased dietary intake of folic acid (FA)

has become an increasing problem in the U.S and other countries as more people are consuming

at or above the recommended daily amount of FA. The impact of too much dietary FA on hypoxia

is not well understood. This study aimed to investigate how increased dietary levels of FA impact

hypoxia outcomes using *Drosophila melanogaster* as a model. Adult female and male $w^{1118}$

*Drosophila melanogaster* flies were placed on control (CD) and 100 μM folic acid (FAD) diets.

At 5 to 6 days old progeny were exposed to hypoxia for two hours prior to returning to normoxic

conditions to model reperfusion. We observed escape behavior in hypoxia larvae, confirming

exposure to hypoxia Using liquid chromatography-tandem mass spectrometry, elevated FA levels

were observed in FAD compared to controls. We report increased acute hypoxia-induced mortality

in FAD flies. Furthermore, FAD flies were not motivated to climb after hypoxia. Under normoxic

conditions FAD flies had a higher velocity when descending during a climb. There was no impact

of FA on neurodegeneration in brain tissue. The data collected from this study suggests that

hypoxia has a negative effect on survival and climbing behavior in a fly model. Together these

data suggest that increase dietary intake of FA can have negative health outcomes after hypoxia.

**Keywords**
Ischemic stroke, Folic acid, Hypoxia, *Drosophila melanogaster*, Nutrition

57
**Introduction**
58
59
Folic acid (FA) is well known for its role in pregnancy as the demand for it increases

significantly. Inadequate levels of FA during pregnancy lead to infant neural tube defects

(NTD)[1]. In 1998, the U.S and Canadian governments began fortifying foods with FA to reduce

the prevalence of NTDs. Furthermore, the movement towards optimizing health through proper

diet and supplementation started to gain popularity. Research followed suit with most published

work focusing predominantly on the adverse effects and risks associated with deficiencies in FA

[2]. However, the impact of increased levels of FA on other health outcomes is not well

characterized.

There have been reports indicating that too much dietary intake of FA can be detrimental

to health [3–5]. This includes promotion of tumorigenesis [3,4], negative interactions when levels

of B12 are reduced [5] and negative outcomes in offspring when mothers are fed too much folic

acid during pregnancy [6]. The Centers for Disease Control recommends 400 μg per day of FA

daily for adults [7].  Over-supplementation is defined as a consumption of FA over 1,000 μg per

day in adults [8]. A study in the US reported that 54.9% of the sampled population excluding

pregnant and lactating women were taking $\geq$ 400 μg/d of FA [9] .

In populations where no mandatory FA fortification is not in place, supplementation with

FA has been shown to have positive effects when individuals are at high risk of stroke [10,11].

Results from an *in vitro* study investigating FA supplementation of 4μg/mL prior to hypoxia in

human monocyte THP-1 cells showed anti-inflammatory effects, through downregulation of

interleukin-1beta (IL-1β) and tumor necrosis factor-alpha (TNF-α) protein levels [12]. In this same

study, vascular endothelial growth factor (VEGF) mRNA levels are decreased in a dose-dependent

manner in response to FA and hypoxia treatments. Additionally, FA pretreatment attenuates

nuclear hypoxia inducible factor-1 alpha (HIF-1α) signal, which could lead to worse chronic

hypoxia management due to less angiogenesis [12]. Other *in vitro* studies have shown that FA

reduced the inflammatory response by inhibiting reactive oxygen species [13] and increased cell

survival  [14] post hypoxia. In mice a combination of increased B-vitamins and choline

ameliorated hypoxia-induced memory deficit [15]. However, the impact of FA over-

supplementation *in vivo* requires more investigation. The objective of our present study is to

understand the role of increased dietary intake of FA and hypoxia on survival, larvae behavior,

and climbing behaviors *in vivo* using the *Drosophila melanogaster* model.

**Methods**
91

92    **Experimental timeline and *Drosophila melanogaster* model**

93           Our experimental manipulations are summarized in **Figure 1**. Manipulations included

adult $w^{1118}$ flies (Bloomington Drosophila Stock Center, Indiana University, Bloomington, IN,

USA) were placed on either folic acid diet supplemented (FAD) or control diets for a period of

about ten days. This ten-day period is roughly the amount of time it takes for larval development

to finish and allow the offspring to emerge from pupa cases as adult flies [16]. After about 5-6

days post-eclosion, adult flies are either subjected to hypoxia for a period of 2 hours or left in

atmospheric oxygen levels (normoxia). Twenty-four hours after the exposure to hypoxia, the flies

were subjected one of the following experiments: mortality, climbing assay, larvae escape time, or

brains tissue. Non-hypoxia control and FAD flies whole bodies were collected for FA and related

metabolite measurements. The flies were housed in a controlled laboratory environment, 25ºC with

consistent 12-hour light-dark cycles.

**Increasing Dietary Intake of Folic Acid in Fly Food**
105

A dose of 100 µM of folic acid was chosen for this study. Prior work using 50 µM or 100

µM of folic acid combined with hypoxia demonstrated high fly mortality using the 100 µM dose

(data not shown).

Weekly fly food would be prepared for fly vial dispensation, there were two conditions

control (CD, 0 µM FA), or additionally supplemented with 100 µM FA (FAD) (MilliporeSigma,

Burlington, MA, USA) in the liquid form and then solidified. The FA was dissolved in 0.0001 N

NaOH (ThermoFisher Scientific, Waltham, MA, USA), so control diet was had 0.0001 N of NaOH

added.

FA was added to standard fly food consisting of a mixture of corn syrup, cornmeal,

molasses, agar, soy flour, and yeast using standard methods and protocols.  The stock food was

prepared using reverse osmosis water, treated with propionic acid for preservation and allowed to

anneal at room temperature prior to transfer into a fridge before future portioning out into

individual fly vials. Food vials were stored at 4ºC until used and were replaced every 7 days for

the duration of experimental studies.

**Hypoxia Treatment**
An Eppendorf CellXpert (C170i) was used to induce hypoxia in flies.  Flies 5-6 days post

eclosion were subjected to 1% $O_2$, 26°C, and 0.1% $CO_2$ for 2 hours. After hypoxia exposure, flies

were transferred to normoxic (20% $O_2$) conditions for 24 hours to model reperfusion injury (ref).

**Behavioral Escape Response of Larvae to Hypoxia**
To confirm exposure to hypoxia we performed larvae climbing experiments [17,18]. These

experiments were performed in the hypoxia chamber, using the transparent cover to count larvae

that moved out of food vial as demonstrated in Figure 3A. Vials containing larvae at all stages

were placed in the hypoxia chamber via their respective FA treatment vials, along with flashlights

to illuminate the interior of the chamber. The chamber was activated and allowed to reach the

desired point of hypoxia. At that stage, the number of larvae in each of the 12-29 vials per condition

group that exhibit a behavioral escape response was recorded at time 0, 5, and 10 minutes. We

measured larvae from control and FAD diets, as well as normoxia and hypoxia conditions.

**Folic Acid Metabolite Measurements**

Whole body tissue lysates of FA-treated 5–6-day post-eclosion female and male flies were

tested. Six samples per experimental group were used for measurements. Body tissue lysates were

measured for FA, tetrahydrofolic acid (THF), and 5,10-methenyltetrahydrofolate using a UHPLC–

MS/MS system consisting of a Waters Acquity UHPLC coupled to a Xevo-TQS triple quadrupole

mass spectrometer equipped with an electrospray ionization probe (Waters Corporation, Milford,

MA, USA), as previously described [19].

**Survival Assay**

Starting 24 hours post-hypoxia exposure (n = 204 to 465 individual flies), the number of

dead flies was recorded, and surviving flies were recorded daily until all flies expired.

Approximately 20 to 25 vials of flies from hypoxia and normoxia (control) groups were kept for a

survival assay lasting about 90 days based on when the last fly expires.

**Climbing Assay**

We measured climbing behavior in flies to negative geotaxis in flies [20]. The behavioral

assay was performed on flies 24 hours after exposure to hypoxia or normoxia. Flies from each

experimental group were placed individually into one of 16 transparent 5 mm diameter, 80 mm

long polycarbonate tubes.  The MB5 Multibeam Activity Monitor (Trikinetics Inc. Waltam, MA,

USA) holds 16 polycarbonate tubes in a vertical position with 9 infrared beams passing through

each tube at 3 mm vertical distances for a total of 51 mm, leading to a total of 17 unique positions

in the tube. The Multi-Beam Monitor (MBM) device records the position of the fly in the tube

each second for 20-minutes, counting a movement as any positional change that was captured per

second through singular or multiple beam breaks. Every time the fly ascends to a peak and to

reverse travel for a minimum of 3 mm. The total upward movement is determined by counting the

number of beam breaks as a fly ascends. Individual fly behavior was measured, there were 14 to

29 flies used per experimental group.

**Immunofluorescence**

Brain tissue from 9 to 14 flies was used for immunofluorescence analysis to assess

molecular mechanisms involved hypoxia and increased dietary intake of FA. Brain tissue was

fixed with using 3.7% formaldehyde and then blocked using normal goat serum (NGS, 1:10)

diluted in PBS-T. The active caspase-3 primary antibody (ac3; 1:100, Cell Signaling Technologies,

Danvers, MA, USA, Catalog: # 9662S, RRID: AB_331439) was diluted in PBT and incubated

with brain tissue overnight at 4°C. The next day, whole brains were incubated in an anti-rabbit

Alexa Fluor 555-conjugated secondary antibody (1:100, Cell Signaling Technologies Danvers,

MA, USA, Catalog: 4413S) and then mounted for microscope analysis. Z-stacks of active caspase-

3 per brain were captured at 400X magnification with a TCS PSE confocal microscope (Lecia

Microsystems, Wetzlar, Germany) and analyzed with Image Pro Premier 3D image processing

software (Media Cybernetics, Inc., Rockville,

170
171 MD, USA).
172

**Statistics**
All experimental procedures were conducted by individuals blinded to experimental

groups. The larvae escape time was analyzed using two-way ANOVA with repeated measures with

IBM SPSS Version 28.0.1.1. The survival curve after exposure to hypoxia or normoxia with

varying levels of FA was analyzed using a Hillslope curve. This test utilized probability curves

derived from an application of the Hill equation on the control flies using the HEPB (Hill Equation

with Probability Band) program [21]. This analysis provided the predicted day at which each

condition's group's population will have a 50% life expectancy (LE50). The LE50 value of each

condition group was then compared to the true control being flies kept in normoxic conditions with

a control diet. A 95% prediction interval for the confidence band was applied to the data based on

an algorithm by N.C Shammas that approximates critical values in the distribution{Citation}.

Active-caspase 3 immunofluorescence volume data was quantified using an un-paired t-test via

GraphPad Prism 8. Climbing attempts, total height climbed and percent expired flies on day 1

post-hypoxia were analyzed using a three or two-way ANOVA and significant main effects were

followed up with Tukey's multiple comparisons test with GraphPad Prism 8. The mean and the

standard error of the mean of each group is presented on the graphs unless otherwise indicated.

189
190
**Results**

**Increased Levels of Whole-Body Folic Acid in FAD Flies**
To confirm over increased levels of FA, we measured folic acid related metabolites. For

these set of experiments, we measured levels of metabolites in flies that were not exposed to

hypoxia (normoxic flies). We measured FA, tetrahydrofolic acid (THF), and 5,10-

methylenetetrahydrofolate (5,10-$CH_2$-THF) levels using mass spectrometry. FA levels were

elevated in female and male FAD flies (**Figure 2A**; $F_{(1,20)} = 122.1$, $p < 0.0001$). There was no

effect of sex ($F_{(1,20)} = 1.70$, $p = 0.21$) or interaction between FA levels and sex ($F_{(1,20)} = 1.83$, $p =$

0.1913).

Female flies of both control diet and FAD had higher THF levels (**Figure 2B**; $F_{(1,20)} =$

26.03, $p < 0.0001$; control, $p = 0.0039$, FAD, $p = 0.02$).  Diet had no effect on THF levels ($F_{(1,20)}$

$= 0.2140$, $p = 0.65$), and no interaction between FA levels and sex were detected ($F_{(1,20)} = 0.25$, $p$

$= 0.62$).

Control flies had higher levels of 5,10-$CH_2$-THF (**Figure 2C**; $F_{(1,20)} = 14.33$, $p = 0.001$).

This effect was driven by males ($p < 0.0001$).  Untreated female flies had reduced levels of 5,10-

$CH_2$-THF compared to untreated males ($p = 0.005$).  There was a significant interaction between

FA levels and sex ($F_{(1,20)} = 10.07$, $p = 0.005$), and there was no effect of sex on THF levels ($F_{(1,20)}$

$= 2.098$, $p = 0.16$).

**Larval Escape from Food During Hypoxia**

The escape behavior of larvae from food occurs under hypoxic conditions [17,18]. We used

the larvae escape behavior to confirm that our flies were subjected to hypoxia in the incubator we

were using.  A cartoon and representative image of larvae under normoxia and hypoxia conditions

is shown in **Figure 3A**. When exposed to hypoxia, larvae from both control and FAD groups

crawled up out of the food when compared to normoxic larvae (**Figure 3B**; $F_{(1,57)} = 187.88$, $p <$

$0.001$). There was no effect of diet ($F_{(1,57)} = 0.25$, $p = 0.62$), and there was no interaction between

hypoxia and diet ($F_{(1,57)} = 0.16$, $p = 0.69$).  We did not observe larvae escape from food in vials

under normoxic conditions.

**Reduced FAD Fly Survival 24 hours After Hypoxia**

Beginning twenty-four hours after hypoxia treatment, the percent of surviving flies was

recorded for 90 days. The 50% life expectancy ($LE_{50}$; is the day at which 50% mortality occurred).

This value for the control/normoxia flies was 39 days, while the $LE_{50}$ for FAD/normoxia flies was

45 days **(Figure 4A).** The Hillslope curve $LE_{50}$ was determined utilizing the 95% prediction

interval upper and lower probability bands from the control/normoxia data. $LE_{50}$ for upper and

lower bands are 47 and 30 days, respectively **(Figure 4A)**. $LE_{50}$ values less than 30 days are

considered statistically significant [21]. The $LE_{50}$ control/hypoxia was 8 days ($p < 0.05$),

FAD/normoxia was 45 days, FAD/hypoxia 5 days ($p < 0.05$) **(Figure 4A)**.

228   Acute mortality in response to hypoxia was observed the first 24 hours after hypoxia. FAD

flies had increased mortality (**Figure 4B;** $F_{(1,52)} = 6.46$, $p = 0.01$) Additionally there was a

significant interaction between fly groups in different FA and oxygen levels ($F_{(1,52)} = 6.27$, $p =$

0.015), and this effect was exacerbated in as well as flies exposed to hypoxia groups in different

oxygen levels ($F(1,52) = 118.6$, $p < 0.0001$).  Tukey's multiple comparisons test demonstrated a

difference between control diet normoxia and hypoxia flies ($p < 0.0001$) as well as a difference

between FAD normoxia and hypoxia flies ($p < 0.0001$). There was a difference between control

diet hypoxia flies and FASD hypoxia flies ($p = 0.004$).

236
237 **Reduced Climbing Behaviors in Flies Post-Hypoxia**
238   Twenty-four hours after hypoxia treatment, climbing behavior was measured in hypoxic

239 and normoxic male and female flies. There were two specific measurements we made, the first

240 was the total movement and the second was the motivation to climb.

241 *Total Movement*

242   Data with both male and female flies combined because no sex difference was observed

($F_{(1,139)} = 0.39$, $p = 0.53$). Flies exhibited a significant difference between hypoxia and normoxia

in the number of total movements (**Figure 5A**; $F_{(1,143)} = 4.76$, $p = 0.031$). There was no effect of

FA ($F_{(1,143)} = 0.65$, $p = 0.42$) or any interaction between fly groups of different FA and oxygen

levels ($F_{(1,143)} = 0.77$, $p = 0.38$).

*Motivation to Climb*

There were no differences between female and male flies ($F_{(1,129)} = 1.22$, $p = 0.27$), so they

have been grouped together. Control and FAD flies exposed to hypoxia had reduced motivation to

climb, as indicated by number of climbs (**Figure 5B**; $F_{(1,133)} = 12.60$, $p = 0.001$). There was no

effect of FA ($F_{(1,133)} = 0.58$, $p = 0.45$) or interaction between hypoxia and diet ($F_{(1,133)} = 2.76$, $p =$

0.09). Tukey's multiple pairwise comparisons demonstrated an interaction between normoxia and

hypoxia FAD flies ($p = 0.001$), as well as an interaction between FASD normoxia and control diet

hypoxia flies ($p = 0.02$)

*Descending Velocity*

There were no differences between female and male flies ($F_{(1,101)} = 2.53$, $p = 0.061$), so

they have been grouped together. Flies exhibited a significant difference between FAD and CD

flies in their descending velocity (**Figure 5C**; $F_{(1,113)} = 5.32$, $p = 0.023$). Normoxia FAD flies had

a higher velocity than CD flies ($p = 0.087$). Hypoxic control flies had a higher velocity compared

to normoxia control flies ($p = 0.0036$). There was no effect of oxygen levels ($F_{(1,113)} = 0.65$, $p =$

3.08 or any interaction between fly groups of different FA and oxygen levels ($F_{(1,113)} = 1.86$, $p =$

0.18).

**No Changes in Neurodegeneration in Brain Tissue of FAD flies After Hypoxia**
264
265  We measured neurodegeneration in female and male flies that were exposed to hypoxia in

266  brain tissue. These flies were maintained on FAD and control diets. Due to differences in brain

267  size between male and female brains, active-caspase 3 volume was quantified separately between

268  hypoxia between the sexes. Unpaired t-test of male (**Figure 6A**, $p = 0.70$) and female (**Figure 6B**,

269  $p = 0.84$) brains separately demonstrated no significant differences between control diet and FAD.

270
271  **Discussion**
272

Hypoxia is a major component of ischemic stroke. The impact of too much dietary FA

intake on hypoxia is not well understood. This study highlights interactions between hypoxia and

increased dietary intake of FA *in vivo*. We have demonstrated that increased levels of dietary FA

exacerbate impact of hypoxia on mortality at the 24-hour time point. Hypoxia and FA also had a

negative effect on climbing behavior, flies had a decreased motivation to climb, and reduced

movement. There is also a change in their descending velocity. We did not observe an effect of

hypoxia and FAD on active-caspase 3 levels in the fly brain tissue. In future studies we plan to

incorporate other markers of neurodegeneration.  For example, hypoxia inducible factor 1 alpha

(HIF-1$\alpha$) is a transcription factor that is turned on after exposure to hypoxia. It has been thought

to organize the cells' response to hypoxia by regulating angiogenesis, cell survival and other vital

processed [22]. Another marker is Nrf-2 (Nuclear factor erythroid 2-related factor 2), which is

transcription factor involved in the oxidative stress signaling cascade [23].

In our study, we confirmed that the flies were consuming very high amounts of FA from

their diet by measuring FA metabolites in whole body tissue.  However, we were unable to

determine FA body distribution, or how much was absorbed through the gut. Adding 1C enzymatic

data or isolated measurements of these metabolites from fly heads, for example, would help narrow

down possible explanations. There are notable sex differences in levels of 1C metabolites like

tetrahydrofolate and 5,10-CHF-THF. The lack of differences in THF levels may imply enzyme

function leading to THF at this level of 1C metabolism is less affected by FA levels and maybe

more influenced by sex. Increased levels of FA impact developing organism as demonstrated in

mouse studies [6,24,25] as well as older individuals [26]. Cancer has also been reported to increase

as a result of increased dietary FA intake [4,27]. FA dietary deficiencies cause hippocampal

changes in a mouse model [28], demonstrating sex differences in an invertebrate model,

*Drosophila melanogaster,* in conditions of increased dietary FA rather than deficiency.

In our present study, we observed that the highest mortality per day occurred within 24

hours of hypoxia. We found that FAD flies had increased mortality compared to control diet

hypoxia flies. There was no impact of FAD on mortality in normoxic or hypoxic groups throughout

the remainder of the survival study. The sheer stress of hypoxic cell damage occurred contributed

to increased acute mortality in FAD flies. Increase FA levels in hypoxic conditions has been

reported to attenuate IL-10, anti-inflammatory cytokine, levels *in* vitro [12]. An inadequate

response to hypoxic cell damage due to attenuation of hypoxia response mechanisms could be

demonstrated *in vivo* via increased mortality. The relatively indifferent mortality rates after the

first twenty-four hours could be due to robust compensation from the cellular damage that occurs

after hypoxia exposure. Prior research has demonstrated such novel brain regions in fly brain,

namely regions of the brain that interact with the ventral nerve cord and femoral chordotonal

organs that function as proprioceptors for locomotion in *Drosophila melanogaster* [29,30].

Mechanistic insights of how FA diet impacts gross development could be involved in the study of

the ventral nerve cord after hypoxia.  Furthermore, *Drosophila melanogaster* are relatively

resilient to hypoxia [31], and fine-tuning the duration and strength of hypoxia for a could cause

sub-lethal effects of hypoxia, and trigger adaptive compensation in behavioral and metabolic

metrics [31–34].

To dissect the impact of FA and hypoxia on mortality, an additional study that separates

male and female *Drosophila melanogaster* to measure the impact of sex on mortality would

provide additional insight, as we and others have described sex differences [28,35–37]. A sex

difference study in flies would not only provide data on differences of adaptability to FA over-

supplementation after hypoxia between sexes, but it would also show what relative age points in

male and female flies are most affected by chronic increased levels of FAD. The results of that

potential study would be valuable to corroborate with the 1C metabolite data towards finding

potential implications of 1C metabolism aberrancies leading to the exacerbations of mortality. This

could potentially imply that there are different risks of hypoxic cell damage exacerbation caused

by increased dietary of FA levels between males and females, potentially at different ages in their

life cycles.

It is unclear whether strong metabolic support for 1C metabolism occurs in neurons

associated with proprioception and locomotion. Indeed, suppression of 1C metabolism enzymes,

serine hydroxymethyltransferase-2 (SHMT2) and NAD-dependent methylenetetrahydrofolate

dehydrogenase-methenyltetrahydrofolate cyclohydrolase (NMDMC) resulted in motor

impairment and mitochondrial defects in flies [38]. Measurement of 1C enzymes in our model

would provide insight into how they are utilized in cellular maintenance for neuronal cells

associated with locomotion and proprioception. This data could be used to compare the response

of increased FA levels to other cells of various denominations of importance for the fly's ability

to survive.

Several studies have highlighted the values of flies as a model for disease of aging [39].

While ischemic stroke can occur at different age groups, there is significantly higher prevalence

of it occurring in older demographics [40]. The present study used day 5 to 6 old flies which do

not model the current clinical demographics of ischemic stroke. It is important to acknowledge

when trying to study such a disease through models. The mortality of flies may be different if

exposure to hypoxia occurs later in the fly life cycle.  If the present study were to have used older

flies, the hypoxia response mechanisms may not be as robust and could have lasting effects

throughout the remainder of the fly life cycle, leading to potential effects demonstrated in climbing

behavior. It would be valuable to perform neurodegeneration measurements, such as active

caspase-3 immunofluorescence experiments, at different time points through the adult *Drosophila*

life. That data could be relatable to understanding how *Drosophila melanogaster* aging manages

aberrances of FA diet and hypoxia.

As the basic science field and public health begin to progressively see more demographics

come under the conditions of nutritional over-supplementation, our research will hopefully provide

avenues for future research to better understand what the impact of chronic dietary increase of

FAD has on neurological diseases *in vivo*.

**Acknowledgements**
None

**Abbreviations**
1C, one-carbon
FA, folic acid
FAD, folic acid supplemented diet
IL-1β, interleukin-1beta
TNF-α, tumor necrosis factor-alpha protein levels
VEGF, vascular endothelial growth factor
HIF-1α, hypoxia inducible factor-1 alpha

**Ethics Declarations**
Not applicable

**CrediT authorship contribution statement**
Siddarth Gunnala: data curation, investigation, methodology, writing-original draft, writing-review and editing. Alisha Harrison: investigation, methodology, writing- review and editing. Paula Ashcraft: methodology. Meher Garg: methodology. Teodoro Bottiglieri: investigation, methodology, writing- review and editing. Lori M Buhlman: investigation, supervision, writing-review and editing. Nafisa M Jadavji: conceptualization, data curation, formal analysis, methodology, project administration, resources, supervision, validation, visualization, writing-original draft, writing- review and editing.

**Funding Sources**
None

**References**

[1] J.D. Van Gool, H. Hirche, H. Lax, L. De Schaepdrijver, Folic acid and primary prevention of neural tube defects: A review, Reproductive Toxicology 80 (2018) 73–84. https://doi.org/10.1016/j.reprotox.2018.05.004.

[2] A.L. Boyles, E.A. Yetley, K.A. Thayer, P.M. Coates, Safe use of high intakes of folic acid : research challenges and paths forward, 74 (2016) 469–474. https://doi.org/10.1093/nutrit/nuw015.

[3] K.C. Strickland, N.I. Krupenko, S.A. Krupenko, Molecular mechanisms underlying the potentially adverse effects of folate, Clin Chem Lab Med 51 (2013) 607–616. https://doi.org/10.1515/cclm-2012-0561.

[4] C.M. Ulrich, J.D. Potter, N. Ulrich, Folate supplementation: Too much of a good thing?, Cancer Epidemiology Biomarkers and Prevention 15 (2006) 189–193. https://doi.org/10.1158/1055-9965.EPI-152CO.

[5] G.E. Mullin, Folate: is too much of a good thing harmful?, Nutrition in Clinical Practice : Official Publication of the American Society for Parenteral and Enteral Nutrition 26 (2011) 84–7. https://doi.org/10.1177/0884533610392238.

[6] R.H. Bahous, N.M. Jadavji, L. Deng, M. Cosín-Tomás, J. Lu, O. Malysheva, K.-Y. Leung, M.-K. Ho, M. Pallás, P. Kaliman, N.D.E. Greene, B.J. Bedell, M.A. Caudill, R. Rozen, High dietary folate in pregnant mice leads to pseudo-MTHFR deficiency and altered methyl metabolism, with embryonic growth delay and short-term memory impairment in offspring, Human Molecular Genetics 26 (2017) 888–900. https://doi.org/10.1093/hmg/ddx004.

[7] CDC, Folic Acid: Sources and Recommended Intake, Folic Acid (2025). https://www.cdc.gov/folic-acid/about/intake-and-sources.html (accessed June 30, 2025).

[8] B.J. Merrell, J.P. McMurry, Folic Acid, in: StatPearls, StatPearls Publishing, Treasure Island (FL), 2024. http://www.ncbi.nlm.nih.gov/books/NBK554487/ (accessed November 15, 2024).

[9] C.M. Pfeiffer, M.R. Sternberg, Z. Fazili, E.A. Yetley, D.A. Lacher, R.L. Bailey, C.L. Johnson, Unmetabolized folic acid is detected in nearly all serum samples from US children, adolescents, and adults., The Journal of Nutrition 145 (2015) 520–31. https://doi.org/10.3945/jn.114.201210.

[10] G.J. Hankey, B vitamins for stroke prevention, Stroke and Vascular Neurology 3 (2018) 51–58. https://doi.org/10.1136/svn-2018-000156.

[11] Y. Huo, J. Li, X. Qin, Y. Huang, X. Wang, R.F. Gottesman, G. Tang, B. Wang, D. Chen, M. He, J. Fu, Y. Cai, X. Shi, Y. Zhang, Y. Cui, N. Sun, X. Li, X. Cheng, J. Wang, X.C. Yang, T. Yang, C. Xiao, G. Zhao, Q. Dong, D. Zhu, X. Wang, J. Ge, L. Zhao, D. Hu, L. Liu, F.F. Hou, K. Cao, L. Chen, P. Gao, R. Gao, X. Ji, N. Li, C. Ma, W. Wang, X. Yang, X. Dai, F. Fan, X. Gao, R. Hui, H. Jiang, J. Jiang, X. Jiang, W. Kong, B. Liu, G. Sun, L. Sun, B. Wang, D. Yin, W. Yang, H. Zhang, C. Zhang, L. Zhao, L. Wang, Y. Chen, A. Huang, Y. Li, J. Wang, R. Xie, C. Yao, D. Zhao, Z. Zhao, L.J. Wei, F. Chen, J. Dong, J. Gu, J. Guo, L. Shen, W. Zhang, Z. Zhang, Z. Wang, X. Li, Y.L. Wan, G. Hu, J. Tan, S. Dong, C. Tong, H. Xu, J. Liu, J. Xu, D. Li, C. Tong, H. Li, C. Liu, J. Xu, T. Zhou, J. Zhou, C. Wu, C. Liu, Y. Wang, L. Xu, Z. Zhou, Z. Zhou, S. Liu, C. Tang, Z. Zhang, J. Tang, Z. Xu, Z. Xiong, Q. Zhu, P. Wu, Efficacy of folic acid therapy in primary prevention of stroke among adults with hypertension in China: The CSPPT randomized clinical trial, JAMA - Journal of the

American Medical Association 313 (2015) 1325–1335.
https://doi.org/10.1001/jama.2015.2274.

[12]  X. Huang, Z. He, X. Jiang, M. Hou, Z. Tang, X. Zhen, Y. Liang, J. Ma, Folic Acid
Represses Hypoxia-Induced Inflammation in THP-1 Cells through Inhibition of the
PI3K/Akt/HIF-1α Pathway, Plos One 11 (2016) e0151553–e0151553.
https://doi.org/10.1371/journal.pone.0151553.

[13]  J. Ma, X. Zhen, X. Huang, X. Jiang, Folic acid supplementation repressed hypoxia-induced
inflammatory response via ROS and JAK2/STAT3 pathway in human promyelomonocytic
cells, Nutr Res 53 (2018) 40–50. https://doi.org/10.1016/j.nutres.2018.03.007.

[14]  F. Cheng, J. Lan, W. Xia, C. Tu, B. Chen, S. Li, W. Pan, Folic Acid Attenuates Vascular
Endothelial Cell Injury Caused by Hypoxia via the Inhibition of ERK1/2/NOX4/ROS
Pathway, Cell Biochem Biophys 74 (2016) 205–211. https://doi.org/10.1007/s12013-016-
0723-z.

[15]  L. Yu, Y. Chen, W. Wang, Z. Xiao, Y. Hong, Multi-Vitamin B supplementation reverses
hypoxia-induced tau hyperphosphorylation and improves memory function in adult mice,
Journal of Alzheimer's Disease 54 (2016) 297–306. https://doi.org/10.3233/JAD-160329.

[16]  M.A. Fernández-Moreno, C.L. Farr, L.S. Kaguni, R. Garesse, Drosophila melanogaster as a
Model System to Study Mitochondrial Biology, Methods Mol Biol 372 (2007) 33–49.
https://doi.org/10.1007/978-1-59745-365-3_3.

[17]  J.A. Wingrove, P.H. O'Farrell, Nitric oxide contributes to behavioral, cellular, and
developmental responses to low oxygen in Drosophila, Cell 98 (1999) 105–114.
https://doi.org/10.1016/S0092-8674(00)80610-8.

[18]  D.B. Morton, Behavioral responses to hypoxia and hyperoxia in Drosophila larvae:
Molecular and neuronal sensors, Fly 5 (2011) 119–125.
https://doi.org/10.4161/fly.5.2.14284.

[19]  J. Nandania, M. Kokkonen, L. Euro, V. Velagapudi, Simultaneous measurement of folate
cycle intermediates in different biological matrices using liquid chromatography–tandem
mass spectrometry, Journal of Chromatography B 1092 (2018) 168–178.
https://doi.org/10.1016/j.jchromb.2018.06.008.

[20]  S.T. Madabattula, J.C. Strautman, A.M. Bysice, J.A. O'Sullivan, A. Androschuk, C.
Rosenfelt, K. Doucet, G. Rouleau, F. Bolduc, Quantitative Analysis of Climbing Defects in
a Drosophila Model of Neurodegenerative Disorders, J Vis Exp (2015) 52741.
https://doi.org/10.3791/52741.

[21]  S.R. Gadagkar, G.B. Call, Computational tools for fitting the Hill equation to dose-response
curves, J Pharmacol Toxicol Methods 71 (2015) 68–76.
https://doi.org/10.1016/j.vascn.2014.08.006.

[22]  K.J. Banasiak, G.G. Haddad, Hypoxia-induced apoptosis: Effect of hypoxic severity and
role of p53 in neuronal cell death, Brain Research 797 (1998) 295–304.
https://doi.org/10.1016/S0006-8993(98)00286-8.

[23]  N.M. Jadavji, J. Emmerson, W.G. Willmore, A.J. MacFarlane, P. Smith, B-vitamin and
choline supplementation increases neuroplasticity and recovery after stroke, Neurobiology
of Disease 103 (2017) 89–100.

[24]  D. Wiens, M.C. Desoto, Is high folic acid intake a risk factor for autism?—a review, Brain
Sciences 7 (2017). https://doi.org/10.3390/brainsci7110149.

[25] L.K. Murray, M.J. Smith, N.M. Jadavji, Maternal oversupplementation with folic acid and its impact on neurodevelopment of offspring, Nutrition Reviews 76 (2018) 708–721. https://doi.org/10.1093/nutrit/nuy025.

[26] E.M. Moore, D. Ames, A.G. Mander, R.P. Carne, H. Brodaty, M.C. Woodward, K. Boundy, K.A. Ellis, A.I. Bush, N.G. Faux, R.N. Martins, C.L. Masters, C.C. Rowe, C. Szoeke, D.A. Watters, Among vitamin B12 deficient older people, high folate levels are associated with worse cognitive function: Combined data from three cohorts, Journal of Alzheimer's Disease 39 (2014). https://doi.org/10.3233/JAD-131265.

[27] S. Deghan Manshadi, L. Ishiguro, K.-J. Sohn, A. Medline, R. Renlund, R. Croxford, Y.-I. Kim, Folic Acid supplementation promotes mammary tumor progression in a rat model., PloS One 9 (2014) e84635–e84635. https://doi.org/10.1371/journal.pone.0084635.

[28] C. Bennett, J. Green, M. Ciancio, J. Goral, L. Pitstick, M. Pytynia, A. Meyer, N. Kwatra, N.M. Jadavji, Dietary folic acid deficiency impacts hippocampal morphology and cortical acetylcholine metabolism in adult male and female mice, Nutritional Neuroscience (2021) 1–9. https://doi.org/10.1080/1028415X.2021.1932242.

[29] K.M. Moon, J. Kim, Y. Seong, B.-C. Suh, K. Kang, H.K. Choe, K. Kim, Proprioception, the regulator of motor function, BMB Rep 54 (2021) 393–402. https://doi.org/10.5483/BMBRep.2021.54.8.052.

[30] L. Venkatasubramanian, R.S. Mann, The development and assembly of the Drosophila adult ventral nerve cord, Curr Opin Neurobiol 56 (2019) 135–143. https://doi.org/10.1016/j.conb.2019.01.013.

[31] D. Zhou, G.G. Haddad, Genetic Analysis of Hypoxia Tolerance and Susceptibility in *Drosophila* and Humans, Annu. Rev. Genom. Hum. Genet. 14 (2013) 25–43. https://doi.org/10.1146/annurev-genom-091212-153439.

[32] G.G. Haddad, R.J. Wyman, A. Mohsenin, Y.-A. Sun, S.N. Krishnan, Behavioral and Electrophysiologic Responses of Drosophila melanogaster to Prolonged Periods of Anoxia, J Insect Physiol 43 (1997) 203–210. https://doi.org/10.1016/s0022-1910(96)00084-4.

[33] P. Azad, D. Zhou, E. Russo, G.G. Haddad, Distinct mechanisms underlying tolerance to intermittent and constant hypoxia in Drosophila melanogaster, PLoS One 4 (2009) e5371. https://doi.org/10.1371/journal.pone.0005371.

[34] W.A. Van Voorhies, Metabolic function in *Drosophila melanogaster* in response to hypoxia and pure oxygen, Journal of Experimental Biology 212 (2009) 3132–3141. https://doi.org/10.1242/jeb.031179.

[35] F. Sadre-Marandi, T. Dahdoul, M.C. Reed, H.F. Nijhout, Sex differences in hepatic one-carbon metabolism, BMC Systems Biology 12 (2018). https://doi.org/10.1186/s12918-018-0621-7.

[36] P. García-Rodríguez, F. Ma, C.D. Río, M. Romero-Bernal, A.M. Najar, M. de la L. Cádiz-Gurrea, F.J. Leyva-Jimenez, L. Ramiro, P. Menéndez-Valladares, S. Pérez-Sánchez, A. Segura-Carretero, J. Montaner, Diet Supplementation with Polyphenol-Rich Salicornia ramosissima Extracts Protects against Tissue Damage in Experimental Models of Cerebral Ischemia, Nutrients 14 (2022) 5077. https://doi.org/10.3390/nu14235077.

[37] P. Habib, J. Jung, G.M. Wilms, A. Kokott-Vuong, S. Habib, J.B. Schulz, A. Voigt, Posthypoxic behavioral impairment and mortality of Drosophila melanogaster are associated with high temperatures, enhanced predeath activity and oxidative stress, Exp Mol Med 53 (2021) 264–280. https://doi.org/10.1038/s12276-021-00565-3.

[38] I. Celardo, S. Lehmann, A.C. Costa, S.H. Loh, L. Miguel Martins, dATF4 regulation of
mitochondrial folate-mediated one-carbon metabolism is neuroprotective, Cell Death Differ
24 (2017) 638–648. https://doi.org/10.1038/cdd.2016.158.
[39] S.L. Helfand, B. Rogina, Genetics of Aging in the Fruit Fly, Drosophila melanogaster,
Annual Review of Genetics 37 (2003) 329–348.
https://doi.org/10.1146/annurev.genet.37.040103.095211.
[40] D. Amoah, M. Schmidt, C. Mather, S. Prior, M.P. Herath, M.-L. Bird, An international
perspective on young stroke incidence and risk factors: a scoping review, BMC Public
Health 24 (2024) 1627. https://doi.org/10.1186/s12889-024-19134-0.
523
524

**Figure Captions**

**Figure 1.** Experimental timeline for experiments. Adult female and male $w^{1118}$ Drosophila were placed on one of 2 diets: control or 100μM folic acid diet. Offspring from the crosses were used for experiments. At 5 to 6 days of age female and male flies were exposed to 2 hours of hypoxia at 1% oxygen, after which they were moved to normoxic conditions to model reperfusion. Daily mortality measurements were taken for 90 days post hypoxia. Twenty-four hours post-hypoxia negative geotaxis (climbing) assays were performed, or brain tissue was dissected for immunofluorescence experiments.

**Figure 2.** Impact of increased dietary folic acid (FAD) on whole body folic acid (A), Tetrahydrofolate (THF) (B), and 5,10-CH-THF (C) content in flies maintained on control or 100μM folic acid supplemented diets. The data represent mean ± standard error of mean (SEM) of 6 flies per group. A,C ** $p < 0.01$, **** $p < 0.0001$ Tukey's pairwise comparisons. B, **** $p < 0.0001$ 2-way ANOVA sex main effect.

**Figure 3.** Impact of hypoxia and increased dietary intake of folic acid (FAD) on larvae escape behavior. (A) Visual diagram of fly vials depicting differences of exploratory and feeding behaviors for fly larvae in conditions of normoxia and hypoxia. (B) Quantification of the number of larvae escaping at time 0, 5, and 10 after onset of hypoxic conditions (1%) compared to normoxic larvae. There was a significant impact of hypoxia on larvae escape behaviors ($p < 0.001$). The data represent the mean ± standard error of the mean (SEM) of 12-29 vials per group.

**Figure 4.** Survival study of flies maintained on control and increased dietary intake of folic acid (FAD) subjected to 2 hours of hypoxia or normoxia. (A) Ninety-day percent expired flies. (B) Percentage of expired flies 24 hours after hypoxia. * $p < 0.05$, ****$p < 0.00001$, Tukey's pairwise comparison between indicated groups. The data represent the survival of flies over two studies of 204 to 456 flies. For B, the data represent the mean ± standard error of the mean (SEM) of 12-29 vials per group.

**Figure 5.** Impact of hypoxia and increased dietary intake of folic acid (FAD) on fly climbing behaviors. (A) Total climbing movement, * $p = 0.031$ oxygen treatment main effect. (B) Motivation to climb (number of climbs), *** $p = 0.005$ Tukey's pairwise comparison. (C) Descending velocity, * $p = 0.036$, ** $p = 0.0087$ Tukey's pairwise comparison. The data represents the mean ± standard error of the mean (SEM) of 29 to 41 flies per group.

**Figure 6.** Relative volume of active caspase-3 in brain tissue of control diet and increased dietary intake of folic acid (FAD) flies 24 hours after exposure to hypoxia, A, females and B, males. The data represents the mean ± standard error of the mean (SEM) of 9 to13 brains used per diet group.