Cortex-wide characterization of decision-making neural dynamics during spatial navigation

Samuel P. Haley[1,2], Daniel A. Surinach[3], Angela K. Nietz[1], Russell E. Carter[1], Lucas S. Zecker[1],
Laurentiu S. Popa[1], Suhasa B. Kodandaramaiah[2,3], Timothy J. Ebner[1,2#]

[1]Department of Neuroscience, University of Minnesota, Minneapolis, MN, 55455, USA
[2]Graduate Program in Biomedical Engineering, University of Minnesota, Minneapolis, MN,
55455, USA
[3]Department of Mechanical Engineering, University of Minnesota, Minneapolis, MN, 55455, USA

Abbreviated Title: Cortex wide neural dynamics of decision-making during spatial navigation

[#]Correspondence: Timothy J. Ebner, M.D., Ph.D.
Department of Neuroscience
University of Minnesota
Lions Research Building, Room 421
2001 Sixth Street S.E.
Minneapolis, MN 55455
Phone: 612-626-9204
Fax: 612-626-9201
Email: ebner001@umn.edu

Competing Interests Statement: The authors declare no competing financial interests.

Summary

Decision-making during freely moving behaviors involves complex interactions among many

cortical and subcortical regions. However, the spatiotemporal coordination across regions to

generate a decision is less understood. Using a head-mounted widefield microscope, cortex-

wide calcium dynamics were recorded in mice expressing GCaMP7f as they navigated an 8-

maze using two paradigms. The first was an alternating pattern that required short term memory

of the previous trial to make the correct decision and the second after a rule change to a fixed

path in which rewards were delivered only on the left side. Identification of cortex-wide activation

states revealed differences between the two paradigms. There was a higher probability for a

visual/retrosplenial cortical state during the alternating paradigm and higher probability of a

secondary motor and posterior parietal state during left-only. Three state sequences (motifs)

illustrated both anterior and posterior activity propagations across the cortex. The anterior

propagating motifs had the highest probability around the decision and propagating motifs

peaked following the decision. The latter, likely reflecting internal feedback to influence future

actions, were more common in the left-only paradigm. Therefore, the probabilities and

sequences of cortical states differ when working memory is required versus a fixed trajectory

reward paradigm.

Introduction

Decisions are generated through complex interactions among various neocortical regions. The perception-action cycle is a common model used to describe decision-making behavior as a hierarchical flow of activation from posterior sensory regions to anterior motor regions[1]. As an organism collects and processes sensory information about its environment, this information is passed to higher-order cortical regions to weigh decision outcomes and execute commands. While this perception-action cycle predominantly follows the classical dorsal stream of activation with an anterior (feedforward) flow of activation[2], it has also been hypothesized that the system operates in the reverse direction to provide feedback and influence future decision-making behavior[1,3,4]. Although coordinated feedforward/feedback signals have been proposed and cerebral cortical connectivity supports this concept (for reviews see[4-6]), there is limited evidence for bidirectional activation across the cortex as most studies have focused on information flows between two regions. Visualizing and quantifying reciprocal patterns of communication among many brain regions simultaneously is fundamental to understanding how the cortex functions as a cohesive system to generate decisions and behavior.

Cortical lesions and optogenetic inactivation in animals show that decision accuracy requires several key cortical regions. The retrosplenial cortex (RSP), posterior parietal cortex (PPC), and secondary motor cortex (M2) are all critically involved in various decision and spatial navigation tasks[4,7-11]. Inhibition of these regions effects decision accuracy only for specific assays, suggesting alternative circuits are utilized depending on the rules of the task[12,13]. While lesion and inhibition studies have provided essential demonstrations of the importance of single regions, approaches are needed that simultaneously assess the contributions of multiple regions to decision-making in real-time. Mesoscopic optical imaging of cortical dynamics is one approach to capture the roles of diverse regions during decision tasks[7,14,15].

While wide-field calcium imaging has been proven to be a powerful method to investigate neocortical dynamics at the mesoscale level[12-23], most investigations leverage this technique in head-fixed subjects. Expanding wide-field imaging in freely moving mice would allow investigation of more natural and complex behaviors as neural activity during free, unrestrained behavior can differ substantially from head-fixed recordings[24-26]. Therefore, there is a need to examine cortical engagement in freely moving behaviors to capture naturally occurring spatiotemporal patterns of neural activation.

Taking advantage of a recently developed miniaturized head-mounted microscope (mini-

mScope[27,28]), we recorded cerebral cortex-wide calcium activity of freely-moving mice during

two different paradigms in an 8-maze decision task. The two paradigms required different

behavioral strategies to isolate the cortical circuitry engaged during decision-making. One

paradigm required memory recall of the previous decision to correctly make the next decision

(alternating), while the other paradigm simply required a fixed navigation pattern keeping all

other aspects of the task consistent (left-only). While several forms of analyses have been used

for widefield calcium imaging[7,14], we utilized a form of brain state analysis that reduces widefield

calcium data into discrete "states" to characterize cortex-wide activity patterns and how they

change during the task[20,28–31]. The probabilities of the state activations and two state transitions

differ between the two paradigms. By quantifying motifs using sequences of 3 states, we found

not only anterior propagating motifs of cortical activation states consistent with the dorsal stream

but also posterior propagating motifs. Differential utilizations of these feedforward and feedback

spatiotemporal flows during the decision support the perception-action cycle and reveal a

framework for the interactions among cortical regions underlying decision-making during spatial

navigation.

Results

Wide-field imaging in freely moving mice and 8-maze paradigms

Mesoscale calcium imaging in mice expressing GCaMP7f using the head-mounted mini-

mScope (**Figure 1A**) produces clear, high-quality images of a large region of the dorsal

neocortex (**Figure 1B**). Post-mortem histological GFP labeling reveals dense, even expression

of neurons throughout the cortex for both anterior (**Figure 1C**) and posterior (**Figure 1D**) cortical

regions. Six mice were trained in a modified T-maze we will refer to as the 8-maze (**Figure 1E**).

An automatic swing door was used to isolate a single decision at the top of the maze and

sucrose rewards were delivered on either side of the maze. The animals' behavior was first

shaped to follow a figure-8 pattern, followed by two paradigms (**Figure 1F**). The first paradigm

("alternating") required the mice to execute an alternating figure-8 pattern to receive a reward

and the second paradigm ("left-only") implemented a rule change in which the animal was only

rewarded on the left side of the maze. Before initiating imaging in the alternating paradigm, the

mice were fully trained to over 80% accuracy for three weeks. All mice were tested in both
paradigms, starting with the alternating task, and followed directly by the change to left-only.

Behavior in the 8-maze

Mouse behavior during both the alternating and left-only paradigms was evaluated using
a combination of manually scored and automatic pose estimation. Head position was used to
determine instantaneous speed at each frame in the maze (**Figure 2A-B**). The average
movement speed during correct alternating paradigm trials was significantly lower than during
correct left-only paradigm trials (Alternating: 0.14 ± 0.01 m/s, Left-only: 0.18 ± 0.02 m/s, p =
0.03, t=2.98, df=5, Paired t-test; **Figure 2C**), consistent with additional time being allotted to
decide on which maze arm to select. Across the 7 recording days of the alternating paradigm,
the average preference for the left side of the maze was almost exactly half (50 ± 3.4%; **Figure
2D**), indicating no innate preference to one side of the maze over the other during this
paradigm. After the left-only task was implemented, the mice developed a significant preference
for the left side over the subsequent 8 recording days (left 73.2 ± 4.0%, p<0.0001, t=17.27, df
=5, Paired t-test). Since the animals' behavior was shaped to the alternating task prior to the
recording trials, the decision accuracy was initially greater than 75% and remained stable for the
next 6 recording days, with a slight decrease to 73% on day 7 (**Figure 2E**). Following the rule
change to the left-only paradigm, decision accuracy was lower on the first day, followed by a
steady increase in accuracy over the 8 testing days of the left-only paradigm. For the first 3 days
of the left-only paradigm, accuracy was significantly lower than alternating (Day 1 p=0.0025,
t=4.014, df=31; Day 2 p=0.0039, t=3.85, df=31; Day 3 p=0.0018, t=4.126, df=31, 2-way ANOVA
with multiple comparisons, Bonferroni correction). However, the following 5 days revealed a
steady increase in decision accuracy, comparable to the alternating task (**Figure 2E**). The
alternating paradigm showed a significantly higher average decision accuracy than the left-only
(**Figure 2F**, p=0.007, t=4.425, df=5, Paired t-test), due to the initial lower accuracy on left-only
trials as the mice learned the rule change. Nevertheless, during both behavioral paradigms, the
mice displayed an understanding of the task with total average accuracy above 70%.

Cortical dynamics at key behavioral events

To assess patterns of Ca^{2+} fluorescence modulation of the cortical activity, we identified two
salient behavioral events: making the correct decision and getting to the reward spout. Plots of
the average z-scored ΔF/F for 7 CCF regions during a 6 s time window centered on either the

moment the animal crossed the decision point or interacted with the spout during correct trials

illustrate distinct, consistent activation patterns for both positions in the maze (**Figure 3A-B**).

These CCF regions were selected because they represent diverse cortical areas as well as

modulation in calcium fluorescence. At the time the mouse crosses the decision point (t = 0s),

the activity of both motor regions (M1 and M2) decreases at -0.4 s. The somatosensory regions

(SS_ll and SS_trunk) exhibit a local minimum average fluorescence at -0.4 s, followed by a

general increase in activity. Similar decreases in activation at +0.13 s occur at the retrosplenial

cortices (RSP_lat and RSP_dorsal) as well as earlier and later increases in fluorescence.

Finally, a distinct increase in V1 activity precedes the decision point at -0.53 s.

When the fluorescence data is aligned to when the animals initially interacted with the

spout (t=0 s), different activity patterns emerge. Immediately following interaction with the spout

for correct decisions (**Figure 3B**), the motor regions show a prolonged increase in activity with

peaks at +0.4 s. The somatosensory regions exhibit a biphasic activity pattern, peaking in

activity prior to the spout at -0.4 s that reach a minimum after at +1.8 s. Retrosplenial cortical

activity peaks at -0.7 s, followed by a decrease in activation. In contrast, V1 activation generally

decreases around the spout interaction, with a minimum at +0.5 s. As described in the Methods

and Materials, bootstrapping analysis demonstrates that all the average fluorescence peaks and

troughs in **Figure 3A-B** exceeded the 95% confidence interval of the randomized data, so we

conclude these are not chance events.

These $\Delta F/F$ signals were verified to reflect real cortical modulation and are not the result

of confounding variables. First, to ensure the distinct cortical patterns following spout

interactions were specific to the rewarding stimulus and not a confound of reduced movement

speed, the average z-scored fluorescence at all stops outside of the spout region were

compared with the peak fluorescence following correct spout interactions. Stops outside the

spout regions were defined as any 1 sec segment where the animals average speed was

<0.05m/s and was not within a defined region around the spout (**Supplemental figure 1A**).

Statistical comparisons reveal that each CCF regions exhibited a $\Delta F/F$ signal that was unique to

the reward spout, not reduced locomotion (**Supplemental figure 1B**, M2 $p<0.0001$, t=23.71,

df=2358; M1 $p<0.0001$, t=28.33, df=2358; SS_ll $p<0.0001$, t=24.38, df=2358; SS_tr $p<0.0001$,

t=17.81, df=2358; RSP_lat $p<0.0001$, t=13.71, df=2358; RSP_dor $p<0.0001$, t=9.187, df=2358;

V1 $p<0.0001$, t=20.86, df=2358, Unpaired t-test, Bonferroni corrected alpha).

Next, to verify the patterns of activation in the central corridor were not an auditory

confound from the tones (see Methods and Materials), we conducted alternating trials in the

absence of tones and aligned the z-scored ΔF/F to the moment the animals crossed the center

IR beam, where a tone would normally play. The average ΔF/F in each CCF region during

alternating no-tone trials show similar patterns to those observed when the tones were present

indicating these patterns were specific to the region of the maze, not simply a result of the tone

(**Supplemental figure 1C**). Further, no-tone trials exhibit a clear activation of V1 at the IR beam

followed by activation of somatosensory regions, matching the trials with a tone thus

demonstrating the patterns of cortical activation measured are not an auditory confound. Finally,

the average accuracy during tone and no-tone trials were nearly identical suggesting that the

tones were not a meaningful factor in the animals' decisions (Alt tone 0.86 ± 0.04; Alt no-tone

0.86 ± 0.09; $p=0.96$, $t=0.05$, $df=5$, paired t-test). With this, we do not make any further

interpretations about how the tones effect decision behavior in the task, despite their presence

in the 8-maze.

Cortical activation states

While robust, repeatable ΔF/F modulation occurs at different CCF regions during

behavioral events, the spatial activation patterns and their temporal sequences across the

cortex are difficult to investigate in this manner. Therefore, a clustering method was used to

classify each ΔF/F frame as one of a discrete set of cortical activation patterns to more

accurately capture cortex-wide activations[28] (see Methods and Materials and **Figure 4**). Using

k-means clustering across all trials, an average of 12.2 (range 9-14) cortical activation patterns

were identified across the 6 subjects, and 11 global clusters across all subjects (**Supplemental**

figure 2). The spatial patterns of these clusters consist of localized regions of increased

fluorescence, with a fraction of the remaining cortex having a relative decreased fluorescence.

Across clusters, the regions of increased activation extend from the most anterior to posterior

regions of the imaging field. Each ΔF/F frame was assigned to 1 of the 11 global clusters, which

we refer to as cortical activation states, resulting in sequences of cortical activation states as a

mouse navigates the maze. Aligning the cortical activation states to the moment the animals

reached the decision point reveals patterns (**Figure 5**). For example, around the time of the

decision point, activation of the V1/RSP state 8 (yellow) precedes activation of the M2/PPC

state 11 (red). Many of these patterns of state activations are present across all mice.

Qualitative differences between the two paradigms can be observed, with state 8 more

prevalent prior to crossing the decision point during alternating than left-only trials (**Figure 5A-**

B). Comparing the aligned cortical state data to the aligned ΔF/F time courses reveals

similarities (see **Figure 3**), for example the strong activation of V1 preceding the decision point,

verifying the k-means clustering extracts meaningful patterns from the ΔF/F data. When the

state time courses are aligned to when the animals reach the spout, different sequences of

states are evident, for example a prominent state M2 state 3 (blue), that also correlate with the

ΔF/F time courses (**Supplemental figure 3**).

As the two paradigms require different numbers of left and right decisions, we tested for

differences in the state time courses for left and right-hand spout interactions during alternating

trials alone (**Supplemental figure 1D-E**). Comparison of the average probability curves for each

state reveals a main effect difference between the left and right spouts for 3 of the 11 states

(**Supplemental figure 1F-G**, 2-way ANOVA with multiple comparisons, Bonferroni correction,

State 1: $p=0.0003$, $F_{(89,445)} =1.703$; State 2: $p=0.011$, $F_{(89,445)}=1.432$; State 11: $p=0.046$,

$F_{(89, 445)} = 1.302$). However, post-hoc testing did not identify any significant time points to

describe these main effect differences. Therefore, we interpret any state activation differences

between the sides of the maze to be trivial, and any major changes in state probabilities are

related to the paradigm.

To test for differences in state activations between the two behavioral paradigms, the

peri-event activation probability curves for a 3 s window centered on the moment the mouse

reached the decision point were calculated for each state and compared across behavioral

paradigms (**Figure 5C**, see Methods and Materials for detailed description of statistical

comparisons and exclusion criteria). State 8 has a significantly higher activation probability

during the alternating paradigm ($F_{(45,225)}=2.95$, $p<0.0001$), suggesting this task requires

greater utilization of cortical regions involved in visual processing and spatial navigation.

Alternatively, state 11, a combined M2 and PPC state, has significantly increased probability

during left-only trials near the decision point ($F_{(45,225)}=2.38$, $p<0.0001$), indicating greater

involvement of motor planning in M2 in concert with the PPC as the mice traverse the maze to

obtain the reward. We also compared state probability curves during the spout interaction

(**Supplemental figure 3C**), finding the only significant difference is an increased activation

probability of the RSP/V1 state 8 during the left-only trials prior to reaching the spout ($F_{(45,225)}$

= 2.552, p<0.0001). Since the spout interaction occurs following the decision, the decreased

probability of state 8 during the alternating paradigm may indicate that more cortical regions

used in working memory are engaged even after the decision during this paradigm.

State activation frequency by behavioral region

To understand the relationship between cortical state and maze location, we determined

the probability of each state activation when the animal was within the 7 maze regions (**Figure**

6A-B). These maze regions, which are symmetrical on the two sides, can be used to segment

the task into 4 behaviors: decision-making in the central corridor, approaching the reward

spouts, receiving the reward, and returning to the central corridor. By comparing the actual

probability of state activation by maze region with shuffled data (alpha <0.0001, see Methods

and Materials), we identified 25 significant state probabilities across the 7 maze regions (**Figure**

6C-D). Consistent with the state activation plots (**Figure 5**), state 8 has a high probability of

activation before the decision point in the central corridor and state 11 has a high probability

after the decision point in regions 2 and 5 along the path to the reward. For both paradigms, the

probability of the M1/M2 state 3 increases in the two spout regions (regions 3 and 6). As this

activation occurs in the right spout region where no reward was delivered during left-only trials,

this suggests that state 3 encodes anticipatory and/or error information.

Cortical two state transitions

In addition to the cortical activation state probabilities as a function of maze position, an

analysis of the two state transitions assessed how information is passed in the cortex during

decision-making behavior. Using the state data of a 6 second time window centered on crossing

the decision point, the probability of state x transitioning to state y was calculated, resulting in a

11x11 matrix for both the alternating and left-only paradigms (**Figure 6E-F**). Self-self state

transitions (matrix diagonal) were omitted as they do not describe state transitions. The

probability of each state transition was compared with bootstrapped, shuffled data to determine

significance (alpha < 0.05, see Methods and Materials). During alternating trials, significant

transitions include the RSP/V1 state 9 to frontal state 1 and the reverse transition (**Figure 6G**),

suggesting feedforward and feedback communications between frontal and visual/RSP regions.

During the left-only task, significant transitions involve frontal states 10 and 11 to posterior

states 7 and 5, respectively (**Figure 6H**). These left-only transitions suggest a feedback

mechanism by which frontal and lateral regions influence PPC and V1 processing of sensory

information. More generally, these two state transitions demonstrate the occurrence of
directional flows of cortical information, with posterior to anterior (state 9 to 1) and anterior or
anterior/lateral to posterior (states 1 to 9, 11 to 5, and 10 to 7).

To understand where these two state transitions occur in the maze, we plotted the onset
and offset locations of these four significant transitions. The 9,1 and 1,9 transitions occur in
similar locations in the maze during the alternating paradigm with onset and offset locations
throughout the central corridor and following the decision point, predominantly on the left side of
the maze (**Supplemental figure 4**)**.** The 10,7 transition shows similar distribution both before
and after the decision point during the left-only paradigm. The 11,5 transition shows a higher
frequency of expression with state 11 onsets concentrated prior to the decision point and state 5
offsets after rounding the corner (**Figure 6I**). Therefore, during the left-only task, anterior to
posterior cortical communication between M2/PPC and visual regions are engaged while
making the decision.

Cortical activation state motifs

To provide further insights into the flow of activation across the cortex as the mouse
navigates the maze, we characterized temporal sequences of cortical activation states. Referred
to as cortical state motifs, we identified 3-state sequences that were both statistically different
from randomized sequences and exceeded a minimum probability of occurrence (see Methods
and Materials). This resulted in 23 significant state motifs during alternating trials, and 30 during
left-only trials, with 13 common motifs across both paradigms. Among these significant
motifs,14 are characterized by a primarily anterior propagating sequence of activations and 13
by a posterior propagating sequence (**Figure 7A**). In addition to commonly expressed motifs,
both paradigms display unique anterior and posterior propagating motifs.

Rather than randomly distributed probabilities across all possible motif permutations,
there are peaks in probability for specific motifs (**Figure 7B**). Both specific sequences and
propagation characteristics highlight the differences in the two paradigms. Comparing the
probabilities of the most prevalent motifs for both paradigms, 8,7,11 represents a complete
posterior to anterior propagation along the cortex during the left-only task, while the 8,7,5 motif a
partial propagation for the alternating task (**Figure 7B**). Anterior motifs are active for 16.5% of
the 6-second time window centered on the decision point across all trials, and posterior motifs
are active for 11.6% of the time. Across all 6 subjects, anterior motifs are expressed in 55.2 ±

9.3% of correct trials, and posterior motifs in 24.8 ± 14.4%. Thus, around the decision, these

directional flows of cortical activity are both commonly expressed and found in all mice.

Plotting the mouse head position at the times of onset and offset for some of the more

prevalent motifs provides qualitative evidence that anterior and posterior propagating motifs are

utilized at different positions along the maze. For example, anterior propagating motifs for the

V1/RSP state 8 (8,7,11, or 8,7, 5) have onset locations at the bottom and top of the central

corridor with offset locations either at the top of the central corridor or as the mouse rounds the

corner to either reward spout (**Figure 7C**). The respective reverse motifs (11,7,8 and 5,7,8)

have a different spatial distribution, with onsets occurring before and after the decision point,

and offsets clustered just prior to the spout.

We then plotted each activation of the anterior propagating and posterior propagating

motifs during the 6 s time window centered to the decision point (**Figure 7D**) and calculated the

average peri-event probability for all motifs for either direction with respect to the decision point.

Anterior propagating motifs have peak probability near the moment the animals cross the

decision point, and posterior propagating motifs have highest probabilities around 1.5 seconds

prior to and following the decision point (**Figure 7E**). This result suggests that sequential

feedforward and feedback mechanisms are engaged as the mice navigate and make decisions

along the maze. Comparison of these average curves reveals significantly different expression

probabilities among the two flow directions ($p<0.0001$, $F(89,2225)=2.623$, 2-way ANOVA with

multiple comparisons, Bonferroni corrected). By comparing the average probability of the 14

anterior motifs and 13 posterior motifs during the two paradigms, there was no significant

difference between paradigms for anterior motifs ($p=0.86$, $t=0.17$, $df=83$, paired t-test), but a

significant increase in posterior flowing motifs during the left-only task ($p=0.028$, $t=2.25$, $df=77$,

paired t-test, **Figure 7F**). Therefore, anterior propagation of activity is necessary for decision

generation and maze navigation during both paradigms, while posterior propagation is utilized

more often by the left-only paradigm further supporting that unique cortical feedback patterns

arise when the load of working memory is reduced.

Next, we determined activation duration for all anterior and posterior motifs during a 6 s

time window relative to the decision point for each correct lap. The average durations of the

anterior and posterior propagating motifs are nearly identical (1.20 ± 0.16 sec, 1.19 ± 0.12 sec

respectively, **Figure 7G,** $p=0.88$, $t=0.1496$, $df=25$, Unpaired t-test) indicating the differences in

utilization of forward and posterior motifs cannot be simply explained by duration of activation.

Instead, these motifs suggest a cortex wide temporal parcellation scheme that directs task-

dependent information in two directions, with the anterior propagation primarily feedforward

information and the posterior propagation primarily feedback.

Discussion

We used widefield Ca^{2+} imaging to record neural activity during an 8-maze decision task

to investigate differences in cortical involvement during two variants of the task. Since activation

of specific cortical regions during goal-directed navigation is dependent on task context[32], and

cognitively-demanding tasks utilize more spatially diverse cortical regions[13], we hypothesized

that the cortical circuits engaged will depend on the degree to which working memory is

required. We implemented a k-means clustering algorithm to characterize spatiotemporal

patterns of cortical activation, referred to as cortical activation states[27,28], and calculated

probabilities of these states as well as sequential combinations of multiple states. These

findings support the theory of a bidirectional perception-action cycle by highlighting how the

cerebral cortex dynamically modifies how information is passed among regions to most

effectively execute decision-making behavior.

Spatiotemporal patterns of cortical state activations

The cortical state probability curves demonstrated significant differences in state

activation patterns during the decision phase of either behavioral paradigm. State 8,

characterized by activations in V1 and RSP, has a significantly higher activation probability

during alternating trials than during left-only trials. In contrast, a M2/PPC state 11 has an

increased probability during the left-only paradigm. These changes in state utilizations across

behavioral paradigms were observed both with the peri-decision probability curves (**Figure 5C**)

as well as the state probabilities in maze regions (**Figure 6C-D**). The peak probability of

V1/RSP (state 8) activation prior to the decision point in the central corridor during the

alternating paradigm likely reflects a difference in how visual information is utilized during

memory recall under this paradigm. During the alternating task, the required knowledge of the

previous decision may increase engagement of widespread cortical regions that initiate an

anterior flow of activation beginning at the most posterior cortical regions. Given the roles of

RSP in spatial navigation and spatial memory[33,34], a task involving memory of recent spatial

location will also likely require greater use of this cortical region.

The reduced probability of the V1/RSP activation (state 8) and increase of M2/PPC

(state 11) during the left-only paradigm suggests an alteration in cortex-wide involvement during

the decision-making process. To explain this difference, at least partially, this result is consistent

with the role of frontal cortical regions in establishing reward contingencies[35–39]. As an important

neural hub for integrating decision and reward information, M2 contributes to the determination

of the best decision given the possible outcomes[40,41]. The increase in state 11 suggests that the

action-reward contingency is more defined during the simplified left-only paradigm. Furthermore,

the average movement speed was higher during the left-only compared to alternating trials

(**Figure 2C**), suggesting a reduced cognitive load on the animals when generating a decision

and a more efficient transformation of perception to the correct decision. Given the PPCs

involvement in the transformation of sensory inputs to motor regions to guide movement and

make decisions[42], the increased use of state 11 during the left-only paradigm may reflect the

initiation of a simplified flow of activation that starts at slightly more anterior regions, rather than

at V1. Perhaps the perception-action circuit during the left-only paradigm relies less on the

processing of visual and spatial information and shifts the anterior flow of activation forward.

Cortical two state transitions and motif flow patterns

The dorsal stream is a long-standing tenet that visual information is processed

sequentially in higher-order brain regions as it is passed in the anterior direction[2,43–45]. More

generally, this posterior-anterior flow of information is implicated in mediating the transformation

of sensorimotor information during sensory-guided tasks[2]. Wide-field calcium imaging has

shown that both the spatial and temporal pattern of posterior to anterior activation is refined

during learning of an auditory task[46]. More recently, anterior to posterior propagations of

information have been hypothesized to be involved with various cognitive functions[1,3,6,47,48].

From these perspectives, we sought to identify differences in two state transitions between the

two paradigms. During the alternating paradigm, transitions between RSP state 9 and M2 state

1 are significantly more likely than chance, consistent with task demands and involvement of

RSP in spatial memory and spatial navigation[33,34]. The combination of these mirrored transitions

reflects the engagement of the reciprocal circuit between M2 and RSP[49] that might be

necessary during the recall of previous decisions. During left-only trials, transitions from M2

states 10 and 11 to visual and somatosensory states 7 and 5, respectively (**Figure 6E-H**), were

significantly greater than chance, suggesting increased anterior to posterior interaction during

this paradigm. As described above, V1/RSP state 8 shows more pronounced activation during

the alternating paradigm. Interestingly, this increase in transitions from frontal to posterior states

including V1 and PPC occurs in left-only, not alternating trials. The seemly incongruence of

these two results can be explained by the direction in which the activation flows. The higher

frequency of these anterior-posterior transitions (states 11 to 5, and 10 to 7) suggests the left-

only paradigm involves more internal feedback to guide subsequent behavior. Because the

animals displayed a significant increase in average movement speed during the left-only

paradigm, this increased utilization of feedback may contribute to a more efficient generation of

the decisive action. These differences in two state transitions between paradigms highlight an

additional mechanism on how cortical dynamics can change after implementation of new rule to

match altered task demands and/or behavioral strategies.

386   The state motifs analysis further characterized these directional flows of activation,

finding that flows in the anterior and posterior directions are common. Therefore, while we

observed the generally accepted flow of information from sensory cortices to PPC and to frontal

cortices, the results demonstrate that the previously hypothesized posterior flows also occur as

the mice navigate the maze. There are differences in both the probability of common states as

well as the state motifs between the two paradigms. For example, comparison of the most

prevalent motifs between the two tasks suggests the complete posterior to anterior

transformation of information described by the 8,7,11 motif is more common during the left-only

task, while the partial transformation shown in the 8,7,5 motif is more common for the

alternating task (**Figure 7B**). Previous studies found that once a task is learned, the sequence

of cortical activity to execute movement becomes temporally compressed, indicating a more

efficient utilization of task information[19]. Perhaps the decrease in cognitive resources required

during the left-only task allows for fewer steps in the progression of information anteriorly

reflecting a more efficient sensorimotor transformation in a single motif. With more steps

required for the alternating paradigm, as it relies on working memory.

401   These spatiotemporal sequences of activations not only provide new evidence that

information flows through the cortex in both directions, but also that these anterior and posterior

flows occur at distinct times during the perception-action cycle. Anterior propagating motifs are

most likely to occur immediately before crossing the decision point, suggesting engagement of
the dorsal stream pathway as the animal processes sensory information in the central corridor.
Conversely, the peak probability of posterior propagating motifs occur after the decision was
made, similar to the two state transitions. Perhaps this peak in posterior propagating motifs
following the decision, reflects an internal feedback mechanism by which higher-order regions of
the frontal cortex project to posterior sensory regions to update an internal model on task
performance and prepare for or guide future actions, for example the upcoming reward. Most of
the posterior motifs involve M2, consistent with both its intracortical connectivity and numerous
roles in cognitive and executive processes. The latter include engagement in perceptual tasks,
decision-making and choice, executing and learning complex movements, spatial navigation,
and feedback to predict optical flow (for review see[4]). These roles have led to the proposal that
M2 provides top-down information about these processed to cortical and subcortical regions[4,6],
consistent with the posterior flows observed.

Interestingly, the average duration for the anterior and posterior flow motifs were almost
identical at 1.2 s, indicating a uniform and relatively slow bidirectional movement of information.
Comparing flow direction during the two paradigms reveals that an anterior flow is necessary
during both paradigms, further supporting the classic interpretation of sensory information to
action via the dorsal stream. However, posterior flow motifs are significantly more likely during
the left-only paradigm providing evidence that the left-only paradigm uses a different cortical
processing scheme that relies more on internal feedback for future action.

Conclusions

In conclusion, we performed widefield Ca^{2+} imaging in freely moving mice navigating two
versions of an 8-maze and uncovered well-defined patterns of cortical activation at key
behavioral aspects of the task. Differences in the probabilities of cortical activation states, two
state transitions, and sequences of states distinguish the alternating and left-only reward
paradigms. The transformation of visual and spatial information into decision-making involves
the integration of more spatially diverse cortical regions during a task requiring working memory
of the previous decision. Different probabilities and sequences of cortical state activation also
characterize the paradigms. The alternating task utilizes patterns of activation flow from
posterior to anterior including extensive involvement of the visual areas, RSP, PPC, and M2.
The left-only paradigm involves a higher frequency of posterior flows of activation from M2 to

PPC, suggesting a simplified sensorimotor circuit. While the posterior to anterior flows provides

widefield imaging support for the widely accepted dorsal steam hypothesis, the demonstration of

equally prominent reverse flows emphasizes the importance of directional information exchange

across the cortex during the decision-action cycle.

Methods and Materials

All experiments were approved by the Institutional Animal Care and Use Committee (IACUC) at

the University of Minnesota.

8-Maze and behavior task

Based on these previous modified T-maze designs[50–56], we constructed an automated 8-

maze that was able to detect mouse position and deliver a drop of 5% sucrose solution on either

side of the maze after a desired rule was followed. The 8-maze was an 80 x 54 cm rectangular

arena constructed using clear acrylic and custom 3D-printed connection pieces (Creality CR-

10s, HATCHBOX 1.75 mm PLA) (see **Figure 1E**). The walls of the arena were 12 cm tall, and

corridors were 8 cm wide. The internal walls and floor of the maze were covered with a white,

textured vinyl sticker to create traction for the mouse and promote higher visual contrast.

Intersections at the top and bottom of the center corridor had motorized swing doors to shape

the mouse's path in an alternating path during behavior training sessions. Using an Arduino

script, servo motors (Aideepen, MG996R) and infrared break beam sensors (Adafruit Industries,

#2168) controlled the position of 3D printed swing doors relative to the mouse's position in the

maze. To reduce potential IR penetrating light artifacts, break beam sensors were covered with

a vinyl sticker that had a pinhole to allow just enough light through for the sensor to function.

Two variants of the 8-maze task were implemented to target different aspects of

cognitive processing and examine for different patterns of cortical activation. The first paradigm

required the mice to execute an alternating figure-8 pattern to receive a reward (**Figure 1F,**

middle). A solenoid valve was used to deliver a 27 µL reward of 5% sucrose through a rubber

tipped needle spout on either side of the maze when the correct decision was made (The Lee

Company, #LHDA0533315H). The second paradigm implemented a rule change where the

animal was only rewarded on the left side of the maze (**Figure 1F, bottom**). During this left-only

paradigm, the right side of the maze remained open to exploration but did not deliver a reward.

The distance the animals were required to travel was equal for both paradigms, and each side

of the maze was identical. All mice were tested in both paradigms, starting with alternating, and

followed directly by the change to left-only. This way, any mouse-specific changes in behavior

could be isolated and the differences in cortical activity associated with the rule change could be

determined.

Speakers were mounted on the maze walls on either side of the decision point (**see**

Figure 1E) and played one of 3 innocuous tones at specific times during testing. During the

alternating paradigm, a brief 3520 Hz tone played on the right speaker when the mouse was in

the central corridor to signify a right-hand turn is necessary for a reward and a 1047 Hz tone

was played on the left speaker when a left-hand turn was necessary for reward. 15 cm after the

decision point on either side of the maze a 2637 Hz reinforcement tone was played only when

the correct decision was made. Tones remained consistent for the left-only task, with the only

difference being that only the 1047Hz was played in the central corridor to signal only for left-

hand turns. Two behavior cameras were mounted 1.5 m above the maze (30fps, FLIR blackfly).

Prior to each trial, the maze was thoroughly cleaned with 70% ethanol to eliminate inter-trial

olfactory cues. The entire maze was enclosed in an arena with black curtains hanging on each

of the walls to reduce the amount of light in the maze. Only ambient room light from underneath

the maze was present during testing trials.

Mouse training in the 8-maze lasted for 3 weeks, broken into separate 1-week phases.

The first phase simply involved handling the mouse for a 5-minute session per day. The second

phase consisted of placing the mouse in the 8-maze with a weighted dummy head-mounted

microscope and providing a sucrose reward whenever the mouse reached the spout. During the

third phase, a functional mini-mScope was attached, including the data and power wires. Mice

were then taught to follow a figure-8 alternating pattern to receive a sucrose reward using both

automatic swinging doors (5-minute trials per day). To verify the behavior was learned, the door

at the top of the maze was removed to give the mouse free choice, and decision accuracy was

measured. After shaping was completed and the mice achieved a decision accuracy $\geq$75%,

cortical Ca^{2+} imaging was initiated during the alternating task for 7 days, followed by the left-only

task for 7-8 days. Only a single recording session per day per mouse was run to ensure the

animals were consistently water deprived and not satiated by the water delivered from the

reward spouts.

Behavior analysis

Mouse behavior in the 8-maze task was analyzed using a combination of manual scoring

and automatic pose estimation software (DeepLabCut)[57]. Manual scoring consisted of finding

the exact frame where the animal's nose crossed the decision threshold defined in **Figure 1E**,

and the frame when the nose reached the reward spout during correct trials. If the mouse did

not stop at the spout, this frame was defined as the point where the nose is closest to the spout

for the particular lap. Using DeepLabCut, the position of the mouse's nose, head, body, and

base of the tail were tracked and quantified. For speed and position analysis, the head position

was used as it had the most accurate measure of tracking.

Animals, cranial windows, and viral vector injections

In this study, 6 C57BL/6 mice (2 male, 4 female, average age of 225 days at first day of

testing) were used. Animals were housed in a 12-hour reverse light/dark cycle and all behavioral

testing was done during the dark phase. Mice were water restricted to 1 mL per day and were

maintained at a weight at least 80% of their pre-restriction weight. The mice were implanted with

a version of our previously described cranial polymer window[16,17,27,28]. Following a 1 week

recovery from implantation, the mice were then retro-orbitally injected[58] with 4.4×10^{11}

genocopies of a GCaMP7f viral vector in 100µl of sterile saline (AAV-PHP.eb-Syn-jGCaMP7f-

WPRE; Addgene #104488). The viral vector incubated for at least 3 weeks prior to Ca^{2+} imaging

recordings.

Immunohistochemistry

To harvest the brains for tissue processing, the animals were deeply anesthetized using

5% isoflurane and transcardially perfused first with ice-cold phosphate buffered saline (PBS)

followed by 4% paraformaldehyde (PFA). Extracted brains were fixed in PFA for 2 days before

sectioning at 50 µm thickness using a vibratome. Brain slices were then gently shaken in a

blocking solution (10% Donkey serum in 0.1M TBS) for 1 hour. The slices were next incubated

in a 1:1000 dilution of the primary antibody for GFP staining (rabbit anti-GFP, Invitrogen, REF#

A6455) overnight at 4°C followed by three 10-minute washes in tris buffered saline (TBS). The

slices were then incubated in 1:1000 dilution of the secondary antibody overnight at 4°C

(Donkey anti-rabbit Alexa Fluor Plus 488, Invitrogen, REF# A32790). Using a confocal

microscope (Leica Stellaris 8), images of the sections were taken at 20X magnification with a

488 nm laser. The images were then tiled together to get a single high-resolution image for each
section (see **Figure 1C-D**).

Widefield imaging with the mini-mScope

A miniaturized head-mounted microscope (mini-mScope) was used to record Ca^{2+}
activity during both the alternating and left-only trials in the maze (n = 34 alternating, n = 29 left-
only, from 6 subjects). The mini-mScope body was adapted from a previous design[27,28] and 3D
printed from polymethyl methacrylate (PMMA) resin using a stereolithography printer
(FormLabs). To ensure that no light could pass through the PMMA, an additional coating of
black paint was applied to the exterior (DYKEM BRITE-MARK). Underneath the mini-mScope
body, 3 magnets were inserted for attachment to the magnets in the cortical window implant.
The body housed 4 blue LEDs (470 nm, Digikey) radially distributed around a central shaft and
pointed down to the dorsal surface of the cortex. To eliminate wavelengths of light outside of the
range that excites GCaMP7f, bandpass excitation filters were added for each LED (450-490 nm,
Chroma). A miniaturized imaging sensor, the miniFAST was adopted from previous work[59] to be
used for Ca^{2+} imaging. The housing for the CMOS sensor was machined from
polyoxymethylene (Delrin). The focal plane was adjusted by sliding the CMOS housing along
the mini-mScope shaft until a clear brain image was displayed, and a screw was used to anchor
the CMOS in place. The emitted light from GCaMP7f was directed to a collecting lens in the
central shaft of the mini-mScope body and was passed through a bandpass emission filter (500-
550 nm, Chroma) before reaching the CMOS sensor.

Widefield Ca^{2+} recordings

The CMOS sensor was set to record at 30 frames per second (fps) with the gain set to
25 for a total trial duration of 420 seconds. To achieve stable LED power and mitigate heating
issues, the LEDs were flashed at 15 Hz for 17 ms using TTL pulses generated from a
microcontroller (Teensy 3.5). The data acquisition unit (DAQ) for the miniFAST CMOS sensor
synchronized the behavior cameras with the CMOS sensor also using TTL pulses. Prior to each
recording trial, the PET window on the animal's implant was gently cleaned with saline to
remove any dust or debris from the imaging field. During the first 120 s of each trial, no Ca^{2+}
data was collected to allow the LED light power to stabilize. Following this stabilization period,
the mini-mScope was attached to the mouse and the animal was placed in the center lane of

the 8-maze facing the decision intersection. During the recordings, a manually controlled

motorized commutator was used to relieve torsion in the coaxial and LED power cables[60].

Ca²⁺ imaging data pre-processing

All data processing was completed using custom MATLAB scripts (MATLAB 2022b).

The 30 fps recordings were first down sampled to 15 fps to remove all frames in which the LED

was off. The video was then rotated 180 degrees to have the rostral cortex at the top of the

image. Each frame was converted to grayscale and resized using an 80% bilinear compression.

A mask was drawn to exclude pixels in the background and central sinus. The ΔF/F calculation

utilized a previously described method that normalizes the data to changes in global illumination

fluctuations to account for differences in LED power during each trial[61]. Each pixel of the image

stack was divided by its mean over the entire recording. From this, the mean fluorescence of

each frame divided by the mean of the entire stack was subtracted yielding a normalized image

stack. The data were spatially filtered by averaging a 14 pixel-radius and temporally filtered with

a 0.1-5 Hz bandpass filter (MATLAB 2022b). The filtered ΔF/F data were then z-scored for each

recording trial.

ΔF/F bootstrapping

The fluorescence data was bootstrapped by randomly selecting 470 frames from the

entire dataset, for each CCF region, and calculating the average fluorescence value. This

sample size matches the number of correct laps across all imaging trials. Using 1000 replicates,

a 95% confidence interval was calculated by finding the mean randomized fluorescence ± 2

SDs. Any mean fluorescence value of the behavior-aligned data outside of this interval indicates

the temporal activation produces a significantly different fluorescence pattern than random

chance. All average fluorescence peaks and troughs in **Figure 4** exceeded the 95% confidence

interval of the randomized data, so we can conclude these are not chance events.

Identifying cortical activation patterns using K-means clustering

Using a clustering method adapted from a previous study[28], the Ca²⁺ data was reduced

into distinct cortical activation patterns. A k-means clustering step was implemented at three

levels: 1) trial level, 2) mouse level, and 3) global level. For trial-level clustering (**Figure 4A**), the

correlation coefficients between each ΔF/F frame were calculated (MATLAB *corrcoef* function)

and the resulting correlation values were clustered using a k-means algorithm (5000 iterations;
5 replicates).

To determine the optimal number of clusters for a particular trial, a data reconstruction
algorithm was utilized to guarantee that the resulting data reduction described the majority of
the variability in the real ΔF/F data. By sequentially testing increasing numbers of clusters
(starting with 2 clusters), the average of all frames within each cluster was calculated, and the
resulting images represented the spatial weighting matrix. The temporal weighting matrix was
calculated by multiplying the 2-dimensional ΔF/F stack with the pseudoinverse of the spatial
weighting matrix (**Figure 4A**). This resulted in a time course for each cluster representing how
much it contributes to each ΔF/F frame. Data reconstruction was completed by multiplying the
temporal component for each frame with the spatial component. A correlation between each
real ΔF/F frame and the reconstructed frame was calculated, and the average across all frames
as used to determine if the clusters sufficiently described the real ΔF/F data. When the average
correlation between real and reconstructed frames reached a threshold of 0.85, successive
testing of cluster numbers was aborted, and that number of clusters was used for the within-trial
k-means analysis. Occasionally, the reconstruction correlation with the real ΔF/F data plateaued
before reaching the 0.85 threshold. In this case, the optimal cluster number was determined as
the cluster number when the correlation changed less than 3% over the last 5 iterations. Using
the output from the trial-level clustering, a second k-means clustering step was completed to
classify a set of clusters for all trials within individual animals. This second step followed the
analysis outlined for the trial-based clustering, except the optimal number of clusters was
determined by the average number of clusters for the particular animal. In this clustering step,
the k-means parameters were set to 10^6 iterations and 100 replicates.

The final k-means clustering step combined all the output from the within-mouse
clustering to define a universal catalog of clusters that we refer to as cortical activation states.
Due to varying fields of view for each animal, the Allen Brain Atlas Common Coordinate
Framework (CCF)[62], was applied to a representative brain image for each mouse. The average
pixel value within each CCF region was calculated from each output image of the mouse-level
clustering. Reducing image size to the 66 average atlas regions made it possible to compare
average cluster images across different mice by ensuring all regions were anatomically
consistent. The correlation of each reduced image was calculated and clustered using k-means

with 10^6 iterations and 500 replicates as parameters. The optimal number of states was

determined by a t-distance calculation that maximizes the within cluster correlation and

minimizes the across cluster correlation[28,63].

Cortical state bootstrapping

To quantify significant two state transitions, random 90-frame segments of the cortical

state time course were selected from the datasets of either behavioral paradigm. The number of

random 90-frame segments matched the number of correct laps across all imaging trials. These

segments were shuffled within each mouse to disrupt any naturally occurring transitions, and

the transition probabilities of randomized data were calculated. This was repeated over 1000

iterations to create a distribution of random transition probabilities for each of the 110 possible

transition pairs. Any transition probability of the real, behaviorally aligned data > 2 SDs above

the mean probability of randomized data was considered significant.

To assess statistical significance for the state frequency of maze regions, the chance

level probabilities were determined from the frequencies of randomly shuffled data to uncouple

any real patterns of state activation and location in the maze. Again, by performing 1000

iterations of the shuffling, a distribution for each state-region probability was created and

significance was determined if the real probability exceeded 5 SDs of this distribution.

Characterizing cortical state motifs

To find cortical state motifs, the state data was first smoothed using a custom script

that found the most common state within a 5-frame sliding window (MATLAB 2022b). This

resulted in the elimination of many single-frame transitions, while maintaining the general

patterns of state activation. During the time window ± 3 s after crossing the decision point, the

frequency of each 3-state permutation was calculated, and probability was determined by

normalizing to the total number of sequences. Prolonged activations of individual states were

reduced to a single frame to isolate the transitions between different states. For this analysis,

we chose to only utilize the time window around the decision point, as this should yield the most

information about the spatiotemporal patterning of corresponding cortical motifs. Since the

animal's decision has already been made when it reaches the reward spout, we chose not to

analyze this time window as it would not provide information about how the decision is

generated. Only sequences of 3 different states were considered motifs as it describes a
complete transformation of neural activation.

The motif probabilities generated from the real data were then compared to randomized
bootstrapped data to determine which were statistically significant. A 90-frame segment was
randomly selected from either the alternating or left-only data to generate a matrix that is the
identical size of the real behaviorally aligned data. This data was then smoothed in the same
way as the real aligned data. All states within each mouse were then shuffled individually to
preserve the state distribution per mouse. Motif probability of the randomized data was then
calculated, and the results were obtained over 1000 iterations. Any motifs from the real, aligned
data >5 SDs above the mean randomized probability were considered significant. To isolate
only the most probable motifs, a minimum probability of 0.005 was applied.

Statistical analysis

All statistical analyses were performed using Graphpad PRISM 10.2.3. State probability curves
were compared using a 2-way ANOVA with Bonferroni post-hoc correction. Each paradigm was
represented by a probability curve for each of the 6 subjects. If a state was not present for a
particular animal, a probability of zero was used. Samples were matched between the two
paradigms since the probabilities were calculated using the same animals, during the same time
period with respect to behavior. Any comparisons that showed a significant main effect but
lacked significant post-hoc time points or had less than 2 consecutive post-hoc points were
excluded from interpretation as we determined these differences to be too minor to make any
impactful claims.

Data and Code Availability: All calcium imaging recordings, behavioral recordings, and
MATLAB scripts used in data analysis are available upon request.

Author Contributions:

Conceptualization and Funding: SH, DS, RC, SK, TE

Technology development and experiments: SH, DS, SK, LZ

Data analysis: SH, DS

Interpretation and figures: SH, TE

Project supervision: SH, AN, RC, TE

Manuscript writing and editing: SH, TE, RC, AN, LP, LZ, DS, SK

Acknowledgements: The authors and members of the Ebner lab would like to thank Lijuan

Zhuo for assisting with rodent surgeries and general laboratory support during this project. We

thank members of the Kodandaramaiah lab for their work in developing the mini-mScope and

constructing much of the data analysis pipeline for this project. This work was funded in part by

NIH grants P30 DA048742 (TJE and SK), R01 NS111028 (SK and TJE), RF1 NS126044 (SK

and TJE).

References:

1. Fuster JM. Chapter 8 - Prefrontal Cortex in Decision-Making: The Perception–Action Cycle.
In: Dreher JC, Tremblay L, eds. *Decision Neuroscience*. Academic Press; 2017:95-105.
doi:10.1016/B978-0-12-805308-9.00008-7

2. Goodale MA, Milner AD. Separate visual pathways for perception and action. *Trends*
Neurosci. 1992;15(1):20-25. doi:10.1016/0166-2236(92)90344-8

3. Fuster JM. Upper processing stages of the perception–action cycle. *Trends Cogn Sci*.
2004;8(4):143-145. doi:10.1016/j.tics.2004.02.004

4. Yang JH, Kwan AC. Secondary motor cortex: broadcasting and biasing animal's decisions
through long-range circuits. *Int Rev Neurobiol*. 2021;158:443-470.
doi:10.1016/bs.irn.2020.11.008

5. Alexander AS, Tung JC, Chapman GW, et al. Adaptive integration of self-motion and goals in
posterior parietal cortex. *Cell Rep*. 2022;38(10):110504. doi:10.1016/j.celrep.2022.110504

6. Hwang EJ, Sato TR, Sato TK. A Canonical Scheme of Bottom-Up and Top-Down Information
Flows in the Frontoparietal Network. *Front Neural Circuits*. 2021;15.
doi:10.3389/fncir.2021.691314

7. Ren C, Komiyama T. Characterizing Cortex-Wide Dynamics with Wide-Field Calcium Imaging.
J Neurosci. 2021;41(19):4160-4168. doi:10.1523/JNEUROSCI.3003-20.2021

8. Barthas F, Kwan AC. Secondary motor cortex: where 'sensory' meets 'motor' in the rodent
frontal cortex. *Trends Neurosci*. 2017;40(3):181-193. doi:10.1016/j.tins.2016.11.006

9. Coen P, Sit TPH, Wells MJ, Carandini M, Harris KD. Mouse frontal cortex mediates additive
multisensory decisions. *Neuron*. 2023;111(15):2432-2447.e13.
doi:10.1016/j.neuron.2023.05.008

10. Liu J, Liu D, Pu X, et al. The Secondary Motor Cortex-striatum Circuit Contributes to
Suppressing Inappropriate Responses in Perceptual Decision Behavior. *Neurosci Bull*.
2023;39(10):1544-1560. doi:10.1007/s12264-023-01073-2

11. Sul JH, Jo S, Lee D, Jung MW. Role of rodent secondary motor cortex in value-based action
selection. *Nat Neurosci*. 2011;14(9):1202-1208. doi:10.1038/nn.2881

12. Allen WE, Kauvar IV, Chen MZ, et al. Global Representations of Goal-Directed Behavior in
Distinct Cell Types of Mouse Neocortex. *Neuron*. 2017;94(4):891-907.e6.
doi:10.1016/j.neuron.2017.04.017

13. Pinto L, Rajan K, DePasquale B, Thiberge SY, Tank DW, Brody CD. Task-Dependent Changes in
the Large-Scale Dynamics and Necessity of Cortical Regions. *Neuron*. 2019;104(4):810-
824.e9. doi:10.1016/j.neuron.2019.08.025

14. Nietz AK, Popa LS, Streng ML, Carter RE, Kodandaramaiah SB, Ebner TJ. Wide-Field Calcium
Imaging of Neuronal Network Dynamics In Vivo. *Biology*. 2022;11(11):1601.
doi:10.3390/biology11111601

15. Cardin JA, Crair MC, Higley MJ. Mesoscopic imaging: shining a wide light on large-scale
neural dynamics. *Neuron*. 2020;108(1):33-43. doi:10.1016/j.neuron.2020.09.031

16. Ghanbari L, Carter RE, Rynes ML, et al. Cortex-wide neural interfacing via transparent
polymer skulls. *Nat Commun*. 2019;10:1500. doi:10.1038/s41467-019-09488-0

17. West SL, Aronson JD, Popa LS, et al. Wide-Field Calcium Imaging of Dynamic Cortical
Networks during Locomotion. *Cereb Cortex N Y NY*. 2021;32(12):2668-2687.
doi:10.1093/cercor/bhab373

18. Musall S, Sun XR, Mohan H, et al. Pyramidal cell types drive functionally distinct cortical
activity patterns during decision-making. *Nat Neurosci*. 2023;26(3):495-505.
doi:10.1038/s41593-022-01245-9

19. Makino H, Ren C, Liu H, et al. Transformation of Cortex-wide Emergent Properties during
Motor Learning. *Neuron*. 2017;94(4):880-890.e8. doi:10.1016/j.neuron.2017.04.015

20. MacDowell CJ, Briones BA, Lenzi MJ, Gustison ML, Buschman TJ. Differences in the
expression of cortex-wide neural dynamics are related to behavioral phenotype. *Curr Biol*.
Published online February 2024:S0960982224001465. doi:10.1016/j.cub.2024.02.004

21. Musall S, Kaufman MT, Juavinett AL, Gluf S, Churchland AK. Single-trial neural dynamics are
dominated by richly varied movements. *Nat Neurosci*. 2019;22(10):1677-1686.
doi:10.1038/s41593-019-0502-4

22. Makino H. Arithmetic value representation for hierarchical behavior composition. *Nat
Neurosci*. 2023;26(1):140-149. doi:10.1038/s41593-022-01211-5

23. Gilad A. Wide-field imaging in behaving mice as a tool to study cognitive function.
Neurophotonics. 2024;11(3):033404. doi:10.1117/1.NPh.11.3.033404

24. Parker PRL, Abe ETT, Leonard ESP, Martins DM, Niell CM. Joint coding of visual input and
eye/head position in V1 of freely moving mice. *Neuron*. 2022;110(23):3897-3906.e5.
doi:10.1016/j.neuron.2022.08.029

25. Parker PRL, Martins DM, Leonard ESP, et al. A dynamic sequence of visual processing
initiated by gaze shifts. *Nat Neurosci*. 2023;26(12):2192-2202. doi:10.1038/s41593-023-
01481-7

26. Aghajan ZM, Acharya L, Moore JJ, Cushman JD, Vuong C, Mehta MR. Impaired spatial
selectivity and intact phase precession in two-dimensional virtual reality. *Nat Neurosci*.
2015;18(1):121-128. doi:10.1038/nn.3884

27. Rynes ML, Surinach D, Linn S, et al. Miniaturized head-mounted microscope for whole
cortex mesoscale imaging in freely behaving mice. *Nat Methods*. 2021;18(4):417-425.
doi:10.1038/s41592-021-01104-8

28. Surinach D, Rynes ML, Saxena K, Ko E, Redish AD, Kodandaramaiah SB. Distinct mesoscale
cortical dynamics encode search strategies during spatial navigation. *bioRxiv*. Published
online March 28, 2023:2023.03.27.534480. doi:10.1101/2023.03.27.534480

29. Bagi B, Brecht M, Sanguinetti-Scheck JI. Unsupervised discovery of behaviorally relevant
brain states in rats playing hide-and-seek. *Curr Biol*. 2022;32(12):2640-2653.e4.
doi:10.1016/j.cub.2022.04.068

30. Mohajerani MH, Chan AW, Mohsenvand M, et al. Spontaneous cortical activity alternates
between motifs defined by regional axonal projections. *Nat Neurosci*. 2013;16(10):1426-
1435. doi:10.1038/nn.3499

31. Chia XW, Tan JK, Ang LF, Kamigaki T, Makino H. Emergence of cortical network motifs for
short-term memory during learning. *Nat Commun*. 2023;14:6869. doi:10.1038/s41467-023-
42609-4

32. Arlt C, Barroso-Luque R, Kira S, et al. Cognitive experience alters cortical involvement in
goal-directed navigation. *eLife*. 2022;11:e76051. doi:10.7554/eLife.76051

33. Powell AL, Nelson AJD, Hindley E, Davies M, Aggleton JP, Vann SD. The rat retrosplenial
cortex as a link for frontal functions: A lesion analysis. *Behav Brain Res*. 2017;335:88-102.
doi:10.1016/j.bbr.2017.08.010

34. Cooper BG, Manka TF, Mizumori SJY. Finding your way in the dark: The retrosplenial cortex
contributes to spatial memory and navigation without visual cues. *Behav Neurosci*.
2001;115(5):1012-1028. doi:10.1037/0735-7044.115.5.1012

35. Ghods-Sharifi S, Haluk DM, Floresco SB. Differential effects of inactivation of the
orbitofrontal cortex on strategy set-shifting and reversal learning. *Neurobiol Learn Mem*.
2008;89(4):567-573. doi:10.1016/j.nlm.2007.10.007

36. Joel D, Weiner I, Feldon J. Electrolytic lesions of the medial prefrontal cortex in rats disrupt
performance on an analog of the Wisconsin Card Sorting Test, but do not disrupt latent
inhibition: implications for animal models of schizophrenia. *Behav Brain Res*.
1997;85(2):187-201. doi:10.1016/s0166-4328(97)87583-3

37. Oberto V, Gao H, Biondi A, Sara SJ, Wiener SI. Activation of prefrontal cortex and striatal
regions in rats after shifting between rules in a T-maze. *Learn Mem Cold Spring Harb N*.
2023;30(7):133-138. doi:10.1101/lm.053795.123

38. Cardinal RN, Pennicott DR, Lakmali C, Sugathapala, Robbins TW, Everitt BJ. Impulsive Choice
Induced in Rats by Lesions of the Nucleus Accumbens Core. *Science*. 2001;292(5526):2499-
2501. doi:10.1126/science.1060818

39. Womack S, Geduldig PS, Foley EA, et al. Medial Frontal Cortex Lesions: Deficits and
Treatment with Nimodipine. *Exp Neurol*. 1993;124(2):387-389. doi:10.1006/exnr.1993.1211

40. Siniscalchi MJ, Wang H, Kwan AC. Enhanced Population Coding for Rewarded Choices in the
Medial Frontal Cortex of the Mouse. *Cereb Cortex N Y NY*. 2019;29(10):4090-4106.
doi:10.1093/cercor/bhy292

41. Chantranupong L, Beron CC, Zimmer JA, Wen MJ, Wang W, Sabatini BL. Dopamine and
glutamate regulate striatal acetylcholine in decision-making. *Nature*. 2023;621(7979):577-
585. doi:10.1038/s41586-023-06492-9

42. Andersen RA, Cui H. Intention, action planning, and decision making in parietal-frontal
circuits. *Neuron*. 2009;63(5):568-583. doi:10.1016/j.neuron.2009.08.028

43. Krumin M, Lee JJ, Harris KD, Carandini M. Decision and navigation in mouse parietal cortex.
eLife. 2018;7:e42583. doi:10.7554/eLife.42583

44. Guo ZV, Li N, Huber D, et al. Flow of Cortical Activity Underlying a Tactile Decision in Mice.
Neuron. 2014;81(1):179-194. doi:10.1016/j.neuron.2013.10.020

45. Goard MJ, Pho GN, Woodson J, Sur M. Distinct roles of visual, parietal, and frontal motor
cortices in memory-guided sensorimotor decisions. *eLife*. 2016;5:e13764.
doi:10.7554/eLife.13764

46. Gilad A, Helmchen F. Spatiotemporal refinement of signal flow through association cortex
during learning. *Nat Commun*. 2020;11:1744. doi:10.1038/s41467-020-15534-z

47. Buschman TJ, Miller EK. Top-down versus bottom-up control of attention in the prefrontal
and posterior parietal cortices. *Science*. 2007;315(5820):1860-1862.
doi:10.1126/science.1138071

48. Semedo JD, Jasper AI, Zandvakili A, et al. Feedforward and feedback interactions between
visual cortical areas use different population activity patterns. *Nat Commun*. 2022;13:1099.
doi:10.1038/s41467-022-28552-w

49. Yamawaki N, Radulovic J, Shepherd GMG. A Corticocortical Circuit Directly Links
Retrosplenial Cortex to M2 in the Mouse. *J Neurosci*. 2016;36(36):9365-9374.
doi:10.1523/JNEUROSCI.1099-16.2016

50. Van Heusden FC, Palacín I Bonsón S, Stiedl O, Smit AB, Van Kesteren RE. Longitudinal
Assessment of Working Memory Performance in the APPswe/PSEN1dE9 Mouse Model of
Alzheimer's Disease Using an Automated Figure-8-Maze. *Front Behav Neurosci*.
2021;15:655449. doi:10.3389/fnbeh.2021.655449

51. Hoxha M, Sabariego M. Delayed Alternation Task for the Study of Spatial Working and Long
Term Memory in Rats. *BIO-Protoc*. 2020;10(5). doi:10.21769/BioProtoc.3549

52. van der Meer MAA, Redish AD. Theta Phase Precession in Rat Ventral Striatum Links Place
and Reward Information. *J Neurosci*. 2011;31(8):2843-2854. doi:10.1523/JNEUROSCI.4869-
10.2011

53. Pedigo SF, Song EY, Jung MW, Kim JJ. A computer vision-based automated Figure-8 maze for
working memory test in rodents. *J Neurosci Methods*. 2006;156(1-2):10-16.
doi:10.1016/j.jneumeth.2006.01.029

54. Torres-Flores M, Peña-Ortega F. Amyloid Beta Alters Prefrontal-dependent Functions Along
with its Excitability and Synaptic Plasticity in Male Rats. *Neuroscience*. 2022;498:260-279.
doi:10.1016/j.neuroscience.2022.07.006

55. Okada S, Igata H, Sasaki T, Ikegaya Y. Spatial Representation of Hippocampal Place Cells in a
T-Maze with an Aversive Stimulation. *Front Neural Circuits*. 2017;11. Accessed January 19,
2024. https://www.frontiersin.org/articles/10.3389/fncir.2017.00101

56. Shoji H, Hagihara H, Takao K, Hattori S, Miyakawa T. T-maze Forced Alternation and Left-right
Discrimination Tasks for Assessing Working and Reference Memory in Mice. *J Vis Exp JoVE*.
2012;(60):3300. doi:10.3791/3300

57. Mathis A, Mamidanna P, Cury KM, et al. DeepLabCut: markerless pose estimation of user-
defined body parts with deep learning. *Nat Neurosci*. 2018;21(9):1281-1289.
doi:10.1038/s41593-018-0209-y

58. Yardeni T, Eckhaus M, Morris HD, Huizing M, Hoogstraten-Miller S. Retro-orbital injections in
mice. *Lab Anim*. 2011;40(5):155-160. doi:10.1038/laban0511-155

59. Juneau J, Duret G, Chu JP, et al. MiniFAST: A sensitive and fast miniaturized microscope for in
vivo neural recording. Published online November 5, 2020:2020.11.03.367466.
doi:10.1101/2020.11.03.367466

60. Oladepo I, Saxena K, Surinach D, Lehman M, Kodandaramaiah SB. Computer vision guided
open-source active commutator for neural imaging in freely behaving animals. Published
online June 1, 2024:2024.05.28.596351. doi:10.1101/2024.05.28.596351

61. Vanni MP, Murphy TH. Mesoscale Transcranial Spontaneous Activity Mapping in GCaMP3
Transgenic Mice Reveals Extensive Reciprocal Connections between Areas of Somatomotor
Cortex. *J Neurosci*. 2014;34(48):15931-15946. doi:10.1523/JNEUROSCI.1818-14.2014

62. Wang Q, Ding SL, Li Y, et al. The Allen Mouse Brain Common Coordinate Framework: A 3D
Reference Atlas. *Cell*. 2020;181(4):936-953.e20. doi:10.1016/j.cell.2020.04.007

63. Geerligs L, Van Gerven M, Güçlü U. Detecting neural state transitions underlying event
segmentation. *NeuroImage*. 2021;236:118085. doi:10.1016/j.neuroimage.2021.118085

Figure 1: Widefield calcium imaging and the 8-maze apparatus. A. Mini-mScope attached to a mouse while in the 8-maze. **B.** Example brain imaging quality recorded from the mini-mScope. **C-D.** Coronal slices at 20x magnification stained for GFP from an anterior region (**C**) and posterior region (**D**). **E.** Overhead view of the 8-maze consisting of infrared break beam sensors that control the position of the motorized swing door and the delivery of a sucrose solution reward. **F.** Schematic of 8-maze behavioral paradigms. (Top) During the shaping phase, an automatic swing door was used at both the top and bottom of the maze to shape the animals' movement in an alternating pattern. Each lap, doors swung in the opposite position to allow access to the opposite side of the maze. (Middle) The alternating phase utilized only 1 swing door so a free decision could be made at the top of the maze. Rewards were delivered only when an alternating pattern was met. (Bottom) The left-only paradigm rewarded only on the left side of the maze. however, the mouse had the ability to choose the right side and receive no reward.

Figure 2: Behavior in the 8-maze. A-B. Head position during a representative 5-min alternating (**A**) and left-only (**B**) trial. Data points are color coded by instantaneous speed. **C.** Average movement speed ± SD per mouse for the two behavioral paradigms (Alt=alternating and LO=left-only). **D**. Average proportion of left decisions across all trials per mouse ± SD. **E.** Decision accuracy relative to testing day for both behavioral paradigms. **F.** Average accuracy across all trials for alternating and left-only (Mean ± SD).

Figure 3: Cortical dynamics at key behavioral events. A-B. Plot of ΔF/F traces for all correct trials aligned to the time of crossing the decision point (**A**) and interacting with the spout (**B**). Individual mice are separated by horizontal white lines in the plots. Mean ± SD traces shown below each plot. Vertical red dashed lines indicate significant peaks compared to randomized data, dotted blue lines are significant troughs. Vertical solid black line indicates t=0 for the respective behavior. **C.** Atlas legend of the 7 CCF regions plotted in A-B.

Figure 4: Identifying cortical activation patterns using K-means clustering. A. Trial-level clustering methods. ΔF/F stack represents all images collected in a 5-minute recording session (~4500 frames). Correlation of each frame with all other frames is calculated, then clustered using k-means to determine within-trial cortical activation states. Optimal number of states determined with an 85% percent reconstruction threshold. Similar steps followed to cluster all trials for a single animal. **B.** Across-mouse clustering methods. Average cluster images are reduced to atlas regions, then correlation coefficients are calculated. Values are clustered using k-means and a t-distance algorithm was used to determine the optimal number of states. Final state images are determined by averaging all frames belonging to each of the 11 global states.

Figure 5: Cortical activation states aligned to the decision point. A-B. Cortical activation state time course ± 3 s with respect to the moment the mouse crossed the decision point during alternating trials (**A**) and left-only trials (**B**). Each correct lap is plotted by row across all 6 animals, with 470 total correct laps. For state time course aligned to spout interaction, see **Supplemental Figure 3**. **C.** Peri-event probability curves for each state with respect to the decision point. The probability of each state activating was calculated and compared between the alternating (black) and left-only (red) paradigms. Post-hoc time points where the two paradigms are significantly different are highlighted in gray. Main effect significance is indicated with (*).

Figure 6: State probability by region and state transitions. **A.** Legend for the 7 maze regions. **B**. Legend for the 11 cortical activation states. **C-D.** State activation probability for each maze region during the alternating paradigm (**C**) and left-only paradigm (**D**). Black boxes indicate probabilities significantly greater than chance (>mean+5 SDs from randomized) that are present during both paradigms. White boxes indicate significant probabilities unique for the respective paradigm. **E-F**. Probabilities of each cortical activation state transitioning to every other state during the alternating paradigm (**E**) and left-only paradigm (**F**). Significant transition pairs shown in yellow dotted boxes. **G-H.** Average state images corresponding to the significant transitions during the alternating paradigm (**G**) and left-only paradigm (**H**). **I.** Mouse locations for the 11,5 transition during the left-only paradigm. Green dots indicate location at state 11 onset and blue dots indicate position at state 5 offset. Onset and offset pairs are linked with red line.

Figure 7: Cortical state motifs. A. All significant 3-state motifs separated into anterior and posterior propagation patterns. Unique alternating (Alt) and left-only (LO) motifs were found to be significantly more likely than chance for one paradigm but not the other. Common motifs were significant from chance during both paradigms. **B.** Probabilities of all state motif permutations. Real probabilities shown as black line. Bootstrapped probabilities +5SD shown in orange. Specific motifs of interest are highlighted with arrows. **C.** Representative plots of mouse locations during 2 common forward propagating motifs and their respective reverse motifs. Onset locations shown as green points and offset locations shown as blue points. Behavioral paradigm is signified by the line color connecting the onset and offset points (black=alternating; red=left-only). **D.** Plot of anterior propagating and posterior propagating motif activations during a 6-second time window centered on the decision point for each correct trial (Green=anterior flowing motifs, Purple=posterior flowing motifs). **E.** Average motif activation probability curve over the same time window from (**D**). Average anterior propagating motif curve shown as green line, average posterior propagating motif curve shown as a purple line. Significant post-hoc points highlighted in gray area. **F.** Average probability of all anterior and posterior propagating motifs between the two paradigms. **G.** Average motif duration for motifs of either direction. Data are shown as mean ±SD.

Supplemental figure 1: Confounding variable verification. A. Mouse head position at the start of every reduced locomotion period (<0.05 m/s) while outside of the two spout regions. **B.** Average ΔF/F z-scores for each of the 7 CCF regions of interest during reduced locomotion periods in the spout regions (gray) and outside of the spout regions (black). Data are presented as mean ± SD. Significant comparisons indicated with (*). **C.** ΔF/F z-score time course of each CCF region of interest over a 6 second time window centered on the moment the animal crossed the center IR beam, just before the decision point, during no-tone alternating trials. **D**. State time course over a 6 second time window centered on the moment the animal interacts with the spout during correct trials, separated by left and right-hand decisions. **F-G**. Average probability of each state at the left spout (**F**), and right spout (**G**), during the same 6 second time window as D-E.

Supplemental figure 2: K-means clustering results. A. Average images for each mouse-level cluster for each of the 6 subjects. **B.** Global cortical state images from across-mouse clustering. Color scale applies to both A and B.

Supplemental figure 3: Cortical activation states aligned to the spout. A-B. Cortical activation state time course for all 470 correct trials during a 6 second time window centered on the spout interactions for the alternating (**A**) and left-only (**B**) paradigms. **C.** Average state probabilities for each of the 11 cortical activation states during alternating (black lines) and left-only (red lines). Significant post-hoc time points indicated with gray shaded areas. 2-way ANOVA main effect indicated with (*). **D.** Cortical activation state key.

Supplemental figure 4: Mouse locations during significant 2-state transitions. Mouse head position during the onset of a state transition (green dots) and offset (blue dots). Onset/offset pairs are connected with a black line during the alternating paradigm, and a red line during the left-only paradigm.