1
2 **A common stay-on-goal mechanism in anterior cingulate cortex for information and effort**
3 **choices**
4
5 Abbreviated title: Anterior cingulate cortex in information and effort
6

7 Valeria V. González[1], Melissa Malvaez[1], Alex Yeghikian[1], Sydney Wissing[1], Melissa Sharpe[2], Kate
8 M. Wassum[1,3-5] & Alicia Izquierdo[1,3-5]
9
[1]Department of Psychology, University of California-Los Angeles, Los Angeles, California, USA.
[2]Department of Psychology, University of Sydney, Sydney, Australia.
[3]The Brain Research Institute, University of California-Los Angeles, Los Angeles, California, USA.
[4]Integrative Center for Learning and Memory, University of California-Los Angeles, Los Angeles,
California, USA.
[5]Integrative Center for Addictions, University of California-Los Angeles, Los Angeles, California,
USA.

17
Author contributions:
AI: Conceptualization, methodology, formal analysis, data curation, resources, writing-original
draft, writing- review & editing, software, supervision, and funding. **AY:** Investigation and data
curation. **KW:** Resources, writing- review & editing, and supervision. **MM:** Methodology,
supervision. **MS:** Resources, writing- review & editing, supervision, and funding. **SW:**
Investigation and data curation. **VVG**: Conceptualization, methodology, formal analysis,
investigation, data curation, writing- original draft preparation, writing- review & editing,
software, visualization, and funding.
26

27 Corresponding authors:
28 Dr. Valeria V. González (gvaleria@reed.edu) or
29 Dr. Alicia Izquierdo (aizquie@psych.ucla.edu)

30
31 Figures: 3   Abstract: 201
32 Tables: 0   Significance Statement: 119
33 Multimedia: 0   Introduction: 469
34   Discussion: 798
35

36 Acknowledgements:
37 We thank members of the Izquierdo, Sharpe, and Wassum labs for early feedback and
38 suggestions on these experiments.
39
40 Authors report no conflict of interest
41
Funding sources:
This work was supported by R01 DA047870 (Izquierdo), the UC Chancellor's Postdoctoral
Fellowship program (González), and the BRI Carol Moss Spivak Fellowship (González).

Abstract

Humans and non-humans alike often make choices to gain information, even when the

information cannot be used to change the outcome. Prior research has shown the anterior

cingulate cortex (ACC) is important for evaluating options involving reward-predictive

information. Here we studied the role of ACC in information choices using optical inhibition to

evaluate the contribution of this region during specific epochs of decision making. Rats could

choose between an uninformative option followed by a cue that predicted reward 50% of the

time vs. a fully informative option that signaled outcomes with certainty, but was rewarded only

20% of the time. Reward seeking during the informative S+ cue decreased following ACC

inhibition, indicating a causal contribution of this region in supporting reward expectation to a

cue signaling reward with certainty. Separately in a positive control experiment and in support

of a known role for this region in sustaining high-effort behavior for preferred rewards, we

observed reduced lever presses and lower breakpoints in effort choices following ACC inhibition.

The lack of changes in reward latencies in both types of decisions indicate the motivational value

of rewards remained intact, revealing instead a common role for ACC in maintaining persistence

toward certain and valuable rewards.

Keywords: frontal cortex, effort-based decision-making, information choices

Significance Statement

We often make choices to gain information, even when the information cannot be used to

change the outcome. Here we investigated the precise timing of the role of the anterior

cingulate cortex (ACC) in decisions that involve seeking certain versus uncertain rewards. By

optically inhibiting ACC neurons, we demonstrate that this region is crucial for maintaining

persistence toward rewards signaled with certainty, without altering the motivational value of

the reward itself. In a positive control experiment, we also confirm that ACC is important in

effort-based choice. The findings reveal a common role for ACC in maintaining persistence

toward certain and valuable rewards, necessary for making optimal decisions. These results

have implications for understanding psychiatric disorders involving maladaptive reward-seeking

behavior.

Individuals often are willing to pay a cost for information even when the information

cannot be used to change the outcome. Prior research has shown that information-seeking

behavior is modulated by a network involving frontocortical regions such as anterior cingulate

cortex (ACC) and orbitofrontal cortex, along with subcortical regions like the striatum and

pallidum (Bromberg-Martin & Monosov, 2020). Indeed, macaque ACC activity is one the earliest

predictors of future choices suggesting it is important in establishing the drive to actively seek

information from the environment (White et al., 2019). By comparison, research on the specific

brain regions tracking information value in rodents is scarce (Bussell et al., 2023), as are

inquiries about the causal mechanisms involved. Recently, we found that ACC was critical in

information choice preference and that its mechanism was sex-dependent: previously

rewarded information choices strongly predicted future choices in control animals, but not in

female rats following ACC inhibition. These results revealed a causal role for ACC in decisions

involving information (González et al., 2024).

To model information choices in nonhumans, several groups (Macias et al., 2021; Molet

et al., 2012; Orduña, 2015; Pisklak et al., 2015; Stagner & Zentall, 2010; Vasconcelos et al.,

2015) administer a task where rats choose between a more profitable option followed by a

discrete cue, rewarded 50% of the time (*No information*) vs. a less profitable option followed by

a cue that signals a certain outcome (*Information*). One of these cues (S+) always results in

reward (100%) and the other (S-) is never rewarded (0%). In this procedure, the information

follows the choice and does not affect the outcome and thus, any preference for that option

suggests information is inherently rewarding (Cunningham & Shahan, 2019; Macias et al., 2021;

Orduña, 2015; Stagner & Zentall, 2010). A preference for information could be due to an

overweighting of the cue that signals certain reinforcement (S+) and/or an underweighting of

the cue paired with the absence of reward (S-) (González & Blaisdell, 2021; González et al.,

2023; McDevitt et al., 2016). However, behavioral manipulations involving either the S+ or S-

have led to largely equivocal results, with more studies garnering support for overweighting the

S+ (Fortes et al., 2016; Stagner et al., 2011).

To more precisely uncover the timing of the role of ACC in information choices, we

studied decision making following optogenetic inhibition of this region during discrete trial

epochs: On separate sessions during the presentation of the informative cue that fully predicts

reward (S+), or during the intertrial interval (ITI). As a positive control for the effectiveness of

optogenetic inhibition of ACC, we evaluated animals on effort-based choices involving high vs.

low valued rewards, where its role has been documented extensively (Bailey et al., 2016;

Floresco & Ghods-Sharifi, 2007; Hart et al., 2020; Hauber & Sommer, 2009; Mashhoori et al.,

2018; Porter et al., 2019).

111   **Materials and Methods**

Animals

Twenty-four Long Evans rats (Rattus Norvegicus) *(N* = 14 females) were acquired from

Envigo (Bioanalytical Systems, Inc., Indianapolis, IN). Two animals were not included in the

analyses due to unilateral expression of the virus: one due to wrong placement of the optic

fiber and one on losing the headcap during the experiment (Total *N* = 20, 10 females). A subset

of animals (n=11) was also tested on a progressive-ratio (PR) task. Subjects were aged between

post-natal day (PND) 90 and 120 at the start of the experiment. Subjects were paired-housed

Figure 1. Information choice task, Effort-based choice task, timelines, and histology. (A) Info Choice task. Rats choose between two levers (left or right). If the rat chooses the 'Info' option, on 20% of the trials, tone S+ plays for 30s terminating in the delivery of one sugar pellet; the other 80% of trials tone S- plays, terminating in no food. If the rat chooses the 'No-Info' option, a third tone S3 plays for 30s, terminating in one sugar pellet on 50% of the trials. A session consisted on 54 trials evenly split between choice, forced-info and forced no-info trials. Only one lever was available on forced trials. **(B) Timeline Info task.** Rats underwent surgery for virus and fiber implant prior to any training. Conditions with optogenetic inhibition were during the S+ and ITI, counterbalanced and alternated with a no-laser session (baseline) between conditions. During the S+ condition, ACC inhibition occurred in choice and forced-info trials, during the duration of the cue S+ (30 s). During the ITI condition, ACC inhibition occurred on 10% of the 10-s ITI epochs. **(C) Timeline PR task.** Rats underwent surgery for virus and fiber implant prior to any training. During PR Test and PR Choice, ACC inhibition occurred during sucrose pellet retrieval. PR Test was just as the training session. In PR Choice and PR Choice Control, animals were presented with a ramekin of 20 g of chow. The order of testing was counterbalanced across animals. A subset of animals experienced both tasks, in counterbalanced order. **(D) Photomicrograph.** Representative placement of inhibitory opsin ArchT (green) and fiber placement at Anterior-Posterior (AP) level +3.7 for ACC relative to Bregma. **(E) Reconstructions.** Reconstructed placements of inhibitory opsin ArchT virus (green) and optic fiber placement (black) across all animals, for AP level +4.2, +3.7 and +3.2 relative to Bregma, targeting ACC.

before and separated after surgeries, housed in transparent plastic tubs with wood shaving

bedding in a vivarium maintained on a reverse 12-hr light cycle. Experiments were conducted at

a minimum of six days per week during the dark portion of the light cycle. A progressive food

restriction schedule was imposed prior to the beginning of the experiment to maintain rats at

85% of their initial free-feeding weights. Water was provided *ad libitum* in the homecage. The

procedures used in this experiment were conducted under approval and following the

guidelines established by the Chancellor's Animal Research Committee at UCLA.

Viral constructs

An adeno-associated virus AAV9 driving an inhibitory opsin archaerhodopsin T under the

CaMKIIa promoter was expressed in neurons in ACC (n=15, AAV9-CaMKIIa-ArchT)-Green,

packaged by Addgene, Addgene, viral prep #99039). A virus lacking the Arch-T gene and instead

containing the fluorescent tag eYFP (n=5, AAV9-CaMKIIa-EYFP, packaged by Addgene) was

infused into ACC as a null virus control. All animals underwent identical surgical procedures,

and all behavior was conducted in a counterbalanced within-subject design with each animal

serving as its own control between inactivation sessions. This allowed us to account for non-

specific effects of surgery, exposure to AAV9, and non-specific effects of laser stimulation.

Behavioral apparatus

Behavioral testing was conducted in Med Associates (East Fairfield, VT) operant

conditioning chambers. Each chamber contained two retractable levers inserted to the left and

right of a lower food delivery port in the front wall. A photobeam entry detector was positioned

at the entry to the food port to detect head entries. A pellet dispenser connected to the food

port delivered a single 45-mg sucrose pellet (Bio-Serv, Frenchtown, NJ). Tone, clicker and white

noise generator were attached to individual speakers on the wall opposite the lever and

magazine. A 3-watt, 24-volt house-light on the top of the back wall provided illumination. For

optogenetic manipulations, the chambers were outfitted with an Intensity Division Fiberoptic

Rotary Joint (Doric Lenses, Quebec, QC, Canada) connecting the output fiber optic patch cords

to a 532nm Green laser (Shanghai Laser & Optics Century Co., Ltd. (SLOC), China) positioned

outside of the chamber.

Surgical procedures

After completing the training phase, rats were anesthetized with isoflurane for bilateral

infusion of ACC with either an inhibitory opsin archaerhodopsin T (AAV9-CaMKII-ArchT)-Green

(Addgene, Cambridge, MA, viral prep #99039) or eYFP (AAV9-CamKII-EYFP; Addene, Cambridge,

MA). Craniotomies were created for the injection sites but also four extra holes were used to

screw anchors. After the four screws were fixed in the skull, animals were infused with two

bilateral sites of injections in ACC (0.3 µL at 0.1 µL/min AP: +3.7, ML: ±0.8, DV: -2.4 for a total

volume of 0.3 µL per side). After infusion, the blunt needle was left in place for 10 min to allow

diffusion. Using the same craniotomies, a stainless-steel fiber optic cannula ferrule (size:

1.25mm, 200 um Core, NA: 0.37, length: 2.5 mm; Thor Labs, Newton, NJ) was inserted in each

side (AP: +3.7, ML: ± 1.0, DV: -2.1, inserted in a 15° angle). All measurements were taken from

Bregma. Optic fibers were secured in place with cyanoacrylate, bone cement (C&B Metabond,

Parkell Inc., Edgewood, NY) and dental acrylic (Patterson Dental, St. Paul, MN). Animals were

provided with post-operative Carprofen (5 mg/kg, s.c.; Zoetis, Parsippany, NJ) daily for 5 d.

Behavioral procedure

Pretraining. Before training, 10 sucrose pellets (0.5 g) were provided in the homecage to

accommodate rats to the food rewards. On the first day, rats were trained to eat pellets from

the pellet tray by delivering 1 pellet every 20 ± 15 s in the chamber for a total of 40 pellets. On

days 2 and 3, rats were trained to lever press the left and right lever (lever presented in

alternate order) in an autoshaping procedure. Reinforcements were delivered following each

lever press. After pretraining, animals were either trained on the information choice task or the

effort-based choice task, in counterbalanced order.

Information choice Training. The training stage of the task was comprised of two types of trials:

choice and forced trials. In choice trials, rats were required to choose between two levers

available simultaneously. A choice trial started with the simultaneous insertion of both levers

and the houselight turning on. A choice was made by pressing a lever one time. After the choice

was made, if the animal chose the Info option, both levers retracted and the houselight turned

off. If the lever associated with the Info option was pressed, 20% of the time an auditory cue

(S+) was presented for 30 s always ending with food; the other 80% of the time another

auditory cue (S-) was on for 30 s never ending with food. If the rat chose the No-Info option, a

third auditory cue (S3) was presented for 30 s and ended with food on half of the trials. On

forced trials, only one lever was presented at a time, following the same contingencies

described above. All trials were separated by a 10-s intertrial-interval (ITI) during which all

stimuli were off. A single session constituted 54 trials, 18 choice trials, and 36 forced trials. The

assignment of sounds to S+, S- or S3 and the lever side for each option was counterbalanced

across animals. After 10 sessions, rats were connected to the patch cords without any

stimulation to habituate them to move around and perform the task with the optical tether

(200 μm, 0.22 NA, Doric Lenses) for another 4 sessions, with no light delivery. Training

continued for 14 sessions.

Information choice test. During two consecutive sessions, rats were tested on the task as

described above but this time receiving optogenetic inhibition during either the 30s S+

presentation or on 10% of ITI periods (to equate to the duration of other trial epochs and

minimize potential damage due to prolong photoinhibition). Green light was delivered into ACC,

which inhibited pyramidal neurons in this region during these trial epochs in rats expressing

ArchT and not eYFP controls. During each presentation, a green light (532 nm; 10 mW) was

delivered via a laser (532nm; Shanghai Laser and Optics Century Co., Shanghai, China)

connected through a ceramic mating sleeve (2.5 mm; Precision Fiber Products, Chula Vista,

California) to the ferrule implanted in the rat. Light leakage from laser output was prevented

using 5-cm long black shrinktube shielding over the connected patch cord and cannula ferrules.

Between each condition, animals received a baseline session, in which no inhibition was

administered but animals were connected to the optical tether as before. The order of

inhibition during the three different epochs was counterbalanced (**Figure 1B**).

Progressive-ratio (PR) Training (for Effort-based choice). Animals were trained on a progressive

ratio (PR) schedule where the required number of presses for each pellet increased according to

the formula:

$$n_i = 5e^{(i/5)} - 5$$

As described in Hart et al. (2020), the effort requirement increased for the rat on earning 5

pellets where n_i is equal to the number of presses required on the i^{th} ratio, rounded to the

nearest whole number after 5 successive schedule completions. Each session started with the

insertion of the left lever. The completion of a ratio requirement was marked by a light, which

remained on until the animal collected the pellet from the food tray. A session lasted 90 min.

The training lasted a minimum of 5 sessions and continued until stability was reached, defined

as the average number of lever presses in a session not varying by more than 15% across the

last two sessions.

PR Test. Animals experienced two test sessions in which ACC was inhibited, using the same

parameters described in the Info choice test. In one test, animals underwent a PR session

similar to the training phase, during which inhibition occurred during pellet retrieval, and where

sucrose pellets were the only option. Our previous imaging of single cells in rat ACC revealed a

phasic increase in activity during reward collection of the preferred reward (Hart et al., 2020),

thus, we predicted that inhibiting ACC during this trial epoch would be particularly effective in

changing effort-based decision making. In another test session, animals received a PR Choice

session where the task occurred as described, but rats also had free access to 20 g of chow in a

ceramic ramekin positioned on the right side of the food port (opposite the lever location,

along the same wall as the pellet tray). For both tests, the onset of the inhibition occurred at

pellet delivery and lasted for 2 seconds or until the animal collected the pellet or pressed the

lever (whichever occurred first). Animals also underwent a PR Choice control session identical

to PR Choice but without laser inhibition. The order of test sessions was counterbalanced

(**Figure 1C**).

Histology

At the conclusion of the experiment, rats were euthanized with an overdose of sodium

pentobarbital (Euthasol, 0.8 mL, i.p.; VetOne, Paris, France) and transcardially perfused with

phosphate buffered saline (PBS) followed by 10% buffered formalin acetate. Brains were

extracted and post-fixed in this solution for 24 hours followed by 30% sucrose immersion.

Tissue was sectioned in 40-µM thick slices and cover slipped with DAPI mounting medium

(Prolong gold, Invitrogen, Carlsbad, CA) visualized using a BZ-X710 microscope (Keyence, Itasca,

IL), and analyzed with BZ-X Viewer software. A representative image of the viral expression and

implant tracks in the Arch-T group is shown in **Figure 1D**. Reconstructions of the viral expression

across all Arch-T expressing animals are shown in green with black dots indicating the location

of the bottom of the optic fiber implants across animals (**Figure 1E**).

Data Analysis

All data processing was conducted using R Studio. All analyses were performed via

custom-written code in MATLAB (MathWorks, Inc., Natick, MA). For the Info choice task, we

used three main variables for analyses: 1) preference for the Info option (Info choice divided by

the total number of choices), 2) latency to choose (the time from the onset of the trial until

lever press), and 3) port entry rate (the number of entries into the food port) during the 30-s

cue duration. For all analyses, we used mixed-effects General Linear Models (GLM) (*fitglme*

function; Statistics and Machine Learning Toolbox) to more rigorously handle data

interdependencies (de Melo et al., 2022; Yu et al., 2022) such as cue exposure depending on

individual rat's choice. Preference data were analyzed first in omnibus GLM that included all

factors (sex, trial epoch, laser on/off) and all groups (virus: Arch-T vs. control). Sex was included

as a moderator in training and a covariate at test. Follow-up analyses were further pursued

pending significant interactions in the full model with planned comparisons: S+, S-, S3 and PR,

PR Choice. Coding of variables for GLMs was as follows: 0 = females and 1 = males; 0 = control

and 1 = ArchT; 0 = ITI, 1 = S+; 0= laser off (no laser), 1= laser on (laser on in the current trial).

Individual rat was entered as a random factor. Latency analysis included the described variables

in addition to the within-subject variable: trial-type (forced-info, forced-no info and choice).

Port entry rate analysis included cue as a crossed, within-subject variable (S+, S- and S3).

For the PR task we used three main measures: 1) Mean lever presses in a session, 2)

Breakpoint (BR), that is, highest incomplete PR ratio, and 3) Latency to collect reward. As in the

information task, we used GLM planned comparisons to probe significant interactions, when

observed.

259   **Results**

Information choice training: Rats acquire a stable preference phenotype, and respond faster

in choice trials.

Preference. We found that rats exhibit a stable preference phenotype that does not change by

the end of training. Training data were analyzed using preference across sessions. A GLM was

performed for the outcome variable, information (*Info*) preference, using session as within-

subject factor, sex as a between-subject factor, and individual rat as a random factor using the

formula: *y ~ [1 + sex*session + (1 + session| rat)].* We found no significant predictors or

interactions with session [GLM: $\beta_{session}$ = 0.005, $t(290)$ = 0.65, p = 0.51], sex [GLM: β_{sex} = 0.025,

$t(290)$ = 0.40, p = 0.68], or session*sex [GLM: $\beta_{session*sex}$ = 0.0002, $t(290)$ = 0.02, p = 0.98].

The absence of a significant effect of session may be due to the high variability of choices in the

development of a preference. To address this, we classified animals based on the phenotype

exhibited during the last three sessions of training: Info preferring (preference above 0.6), No

Information preferring (*No Info*) (preference below 0.4), and Indifferent (between 0.4 and 0.6).

Using phenotype as a fixed factor moderator in the formula *γ ~ [1 + sex*phenotype*session + (1*

+ session| rat)], we found a significant effect of session [GLM: $\beta_{session}$ = 0.033, $t(286)$ = 2.71, p =

0.007] and a significant session*phenotype interaction [GLM: $\beta_{session*phenotype}$ = -0.023, $t(286)$ = -

2.68, p = 0.007]. This interaction indicates that preference changed differentially across sessions

of training. Animals began their preference around the indifference point 0.5 but then diverged,

with a group of animals remaining at the indifference point and the other two groups

developing a strong preference towards/against the Info option (**Figure 2A**). When selecting the

last three sessions of training to check for stability, use of the same formula led to no significant

predictors or interactions, confirming animals' preference was stable by the end of training.

Latency to choose. We found that rats are faster to make choices than to complete forced trials,

corroborating prior research (González et al., 2023; González et al., 2024). We conducted a GLM

on median latencies using trial type (forced Info, forced NoInfo, and Choice) as within-subject

factors, sex as between-subject factor, and individual data as a random factor, formula: *γ ~ [1 +*

trial type*sex + (1 + trial type| rat)]. The results yielded a significant main predictor of trial type

[GLM: $\beta_{trial\ type}$ = 5.19, t(68) = 3.79, p < .001], but no effect of sex [GLM: β_{sex} = 1.43, t(68) = 0.52, p

= 0.60], or interaction [GLM: $\beta_{trial\ type*sex}$ = -.167, t(68) = -0.60, p = 0.55]. Indeed, rats are faster

during choice trials (Median$_{Choice}$= 4.4 s, SEM ± 0.81) than forced trials (Median$_{Forced\ Info}$= 11.4 s,

SEM ± 2.46; Median$_{Forced\ NoInfo}$= 7.6 s, SEM ± 2.64) **(Figure 2B).**

Port entry Rate. Port entry rate is an indication of differential reward expectation for S+ vs. S-.

We analyzed the median port entry rate (entrance to the feeder port) for each 30-s cue

duration for the last three sessions of training. A GLM was conducted for median port entries

using sex as a between-subject factor, and individual rat crossed with cue (S+, S- and S3) as a

random factor, according to the following formula: $y \sim [1 + sex*cue + (1|rat:cue)]$. Cue was

included in this way to account for the fact that rats experienced each cue a different number

of times given the inherent contingencies of the task: S+ was presented on only 20% of all Info

trials but it also depended on the rat's choices (for instance, a rat that chooses the No Info

option exclusively would have an overrepresentation of the S3 cue and very few S+ trials). We

found no effect of cue or sex on port entry rate ($p > 0.64$). We have previously reported the

greatest responses for the S+ and lowest responses to S- (González et al., 2024). However, in

that study, the cue duration was 60s, affording a longer window for differences to emerge.

Information choice test: Optogenetic inhibition of ACC neurons decreases port entry rate

during S+ cue, but does not affect overall information preference or latencies

Preference. Info preference during test sessions was analyzed for both epochs. Mean values

were used for S+ and ITI inhibition sessions. The GLM formula was $y \sim [1 +$

virus*laser*epoch+sex+ (1 + epoch+laser| rat)], with epoch (S+ or ITI inhibition) and laser

entered as within-subject factors, and sex as a covariate. None of these factors or their

interactions significantly predicted Info preference. Given that we observed high variability in

the initial preference, any detectable changes in preference depended on each rat's baseline.

Therefore, we calculated the absolute change in preference during each optogenetic inhibition

session minus the preference during baseline, identical to the analysis in González et al. (2024).

The same GLM formula was used but instead we used the *absolute preference change* as the

outcome of interest. No significant predictors or interactions between factors was found.

Similarly, these same factors did not predict *phenotype*. In summary, optogenetic inhibition of

ACC did not affect overall preference or change in preference phenotype (**Figure 2C**).

*Latencie*s. Optogenetic inhibition of ACC could affect performance measures such as latencies,

not just preference *per se*. To test this, we analyzed median latencies to choose during test

sessions. Here we used the formula: *γ ~ [1 + virus*epoch*trial type*laser +sex+ (1 + epoch +*

trial type+ laser| rat)]; where virus and sex were between-subject factors, epoch (S+, ITI, and

Baseline), trial type (Choice, forced Info and forced No info), and laser (laser on, off) were

within-subject factors, and individual rat was a random factor. The results revealed no

significant predictors or interactions.

Port entry rate. Median reward port entries were analyzed for S+ and ITI epochs using the

following GLM formula: *γ ~ [1 + virus *cue*laser + (1 + laser| rat:cue)]*, and separately for the

baseline condition with laser not included in the model because laser was not introduced: *γ ~ [1*

+ virus *cue + (1 | rat:cue)]. The analysis of baseline sessions resulted in no significant

predictors ($p > .28$) or interactions ($p = .82$) (**Figure 2D**). Similarly, the analysis of S+ inhibition

(with cue left out of the model) resulted in no significant effects ($p > .17$) or interactions ($p >$

0.88) (**Figure 2E**). However, inhibition during the ITI epoch resulted in a significant effect of

laser [GLM: $\beta_{laser} = 9.87$, $t(97) = 2.66$, $p = .009$], a virus*laser interaction [GLM: $\beta_{virus\ *laser} = -$

14.97, $t(97) = -3.45$, $p = .0008$], and virus*cue*laser interaction [GLM: $\beta_{virus*cue*laser} = 5.21$, $t(97)$

$= 2.72$, $p = .007$]. Due to these interactions, we separately investigated the effect of inhibition

in the ITI epoch by virus, where we used this GLM formula for predicting median port entry

rate: $y \sim [1 + cue*laser + (1 + laser| rat:cue)]$.

The analysis for the ArchT, but not controls, resulted in a laser*cue interaction [GLM:

$\beta_{laser*cue}$ = 2.13, $t(73)$ = 2.15, p = .035], indicating the inhibition differentially affected

responsivity to cues. Planned comparisons for each cue resulted in a significantly lower port

entry rate during the subsequent S+ [GLM: β_{laser} = -4.65, $t(20)$ = -2.33, p = .03], but not S- (p

= .20) or S3 (p= .89) (**Figure 2F**). The analysis of the control virus group resulted in a significant

effect of laser [GLM: β_{laser} = 9.119, $t(24)$ = 2.48, p = .020], with animals checking the food port

more (not less) following laser on trials, irrespective of the cue presented. This effect was not

explained by animals being distracted by residual light, given that the responses during cue

presentation occurred after the laser was experienced. In sum, ACC inhibition led to a

prospective decrease in reward checking specifically during the S+, indicating that animals did

not exhibit the appropriate expectation of a certain reward.

Figure 2. Optogenetic inhibition of ACC prospectively reduces reward expectation during the Info (S+) cue. **(A)** Establishment of phenotype in animals during training. **(B)** Density of the total number of observations for latency to choose by trial type collapsed across conditions during training. Dashed lines refer to median latency. Latency to choose in choice trials was faster than in forced trials. **(C)** Mean absolute preference change from baseline (no laser) during inhibition of S+ cue or ITI presentation, with individual rats represented as scatter. **(D)** Mean port entries during the 30 s cue presentations (S-, S+ and S3) for ArchT (left-green) and Control (right-black) animals during baseline sessions, individual diamonds and circles correspond to median responses for each animal. **(E)** S+ inhibition: Mean port entries during each cue presentation on laser on (green vertical stripes) and off (no stripes) for ArchT (left, green) and control (right, black) animals, with individual rats represented as scatter. **(F)** ITI inhibition: Mean responses during each cue presentation following laser on (green vertical stripes) and off (no stripes) for ArchT (left, green) and control (right, black) animals, with individual rats represented as scatter. A decrement in port entries during S+ was observed following inhibition in the ArchT group, but not the Control group. *p<0.05. ArchT n=11, Control n=5.

Progressive Ratio: ACC inhibition reduced effort and decreased breakpoint, without affecting latency to retrieve reward

Total lever presses. We analyzed total lever presses during sessions with laser on during the reward collection period. This epoch was selected based on previous work from our lab

implicating phasic increases in population activity in ACC for higher-valued rewards as essential

for effort-based decision making (Hart et al., 2020). In these analyses, virus was a between-

subject factor, test (PR Test and PR Choice Test) was a within-subject factor, sex was a

covariate, and individual rat was entered a random factor. The GLM formula was as follows: *y ~*

[1 + virus* test + sex + (1 + test| rat)]. We found a main effect of test [GLM: β_{test} = -1229.6, *t*(17)

= -4.02, *p* = .0008], an effect of virus [GLM: β_{virus} = -2191, *t*(17) = -2.42, *p* = .027] but no effect of

sex (*p* = .30) nor interactions (*p*= .19), indicating that responses during the PR Test were higher

than during PR Choice Test, but also that overall responses were lower for the ArchT virus

group compared with the control animals. We also compared performance during sessions with

no laser (last session of training and PR Control Choice). For this, the GLM formula was identical

to the previous analysis, and here we found a significant effect of test [GLM: β_{test} = -678.4, *t*(17)

= -4.86, *p* = .0001], also resulting in a higher level of responding during PR training than PR

Control Choice, but no effect of virus (*p* = .09), sex (*p* = .22), or interactions (*p* = .28). These

results collectively show the expected effect that animals tend to work more vigorously when

sucrose pellets (i.e., in PR training or PR Test, but not PR Choice or PR Control Choice) are the

only option. More importantly, these data indicate that ACC inhibition reduced effort, and that

this decrement in effort was not due to a general reduction in motivation *per se*, given that rats

still lever press more when sucrose was the only reward (i.e., the decrement was proportional

to their initial performance in the given test). If ACC inhibition was affecting motivation, we

would expect a similar level decrement in PR Test and PR Choice.

Breakpoint. The last incomplete PR step was noted as the breakpoint. We analyzed breakpoint

for the two tests with inhibition (PR Test and PR Choice), with an identical GLM formula as for

lever presses: $\gamma \sim [1 + virus* test + sex + (1 + test| rat)]$. This analysis also yielded a main effect

of virus [GLM: β_{virus} = -33.30, $t(17)$ = -2.32, p = .033], and test [GLM: β_{test} = -25, $t(17)$ = -4.92, p

= .0001], but no effect of sex (p = .46) nor interactions (p = .55). For the test sessions without

laser (PR Training and PR Control Choice) we used the same formula as above. Here we found

an effect of test [GLM: β_{test} = -13.5, $t(17)$ = -7.04, p = 2.00^{e-06}], but no other effects (p > .05) or

interactions (p = .18). Taken together, this confirms that the decrement in effort following ACC

inhibition depended on the availability of another source of reward.

Latencies. We analyzed median latencies to collect the reward. As before, we analyzed the two

tests where inhibition occurred using the following GLM formula: $\gamma \sim [1 + virus* test + sex + (1 +$

test| rat)], where γ was median latency to collect reward, virus was a between-subject factor,

test was a within-subject factor, sex was entered as covariate, and individual rat was entered as

a random factor. Comparing PR Test and PR Choice Test, we found a significant effect of sex

[GLM: β_{sex} = -0.40, $t(17)$ = -2.45, p = .026], where females were faster at collecting reward than

males, yet no other significant predictors (p > .089) or interactions (p = .22) were observed.

Similarly, for PR training and PR Control Choice no significant effects (p > .13) or interactions

were found (p = .74). Collectively, these results suggest that the decrement in effort cannot be

explained by a general reduction in motivation to the rewards, where we would expect an

Figure 3. Optogenetic inhibition of ACC during PR and PR Choice decreases effort without affecting motivation.
(A) Mean total lever presses during PR test and PR choice for ArchT (green) and control (black) animals. Individual rats represented as scatter. There was an overall decrement in responses in PR Choice compared with PR test with this decrease significantly more pronounced in ArchT animals. **(B)** Mean total lever presses during control Choice (PR choice with laser off) and last session of training for ArchT (green) and control (black) animals. Individual rats are represented as scatter. Lever presses during Control choice test were lower than during training, with no difference between virus group. **(C)** Mean BP during PR test and PR choice for ArchT (green) and control (black) animals. Responses were lower during PR choice than PR test, but also lower for ArchT animals. **(D)** Mean BP during Control choice and Training (no inhibition) for ArchT (green) and control (black) animals. Individual rats are represented as scatter. No differences between test nor virus were found. **(E-F)** Mean latency to collect reward during inhibited trials (PR Test, PR Choice) and during noninhibited trials (control choice and training) for ArchT (green) and control (black) animals, with individual dots corresponding to median latency for each rat. No difference in reward latencies were observed. *p<0.05. ArchT n=6, Control n=5.

431   increase in latencies to collect pellets.

432   **Discussion**

We investigated the role of ACC in information and effort choices. Previous research has

highlighted the involvement of ACC in decision-making processes, particularly in evaluating

effort costs (Bailey et al., 2016; Cai & Padoa-Schioppa, 2021; Fatahi et al., 2020; Hart et al.,

2020; Hart et al., 2017; Hosokawa et al., 2013), so we used this behavior as a positive control

for the efficacy of our optogenetic manipulation in Info choices. In previous studies, we and

others have found ACC to be important in maintaining a representation of the value of *both*

options in a 2-alternative choice paradigm (Hart et al., 2020; Hart et al., 2017; Mashhoori et al.,

2018). The present results from both tasks suggest that ACC is required to maintain the

appropriate reward expectation, either for the higher-valued reward of sucrose (in the effort

task), or for the more certain reward (in the Info task). Interestingly, inhibiting ACC neurons

decreases this stay-on-goal behavior without changing motivational value *per se* because

reward latencies were unaffected by inhibition. Based on the results of this experiment, we

suggest that this may be the mechanism by which stable preference is maintained in Info

choices (González et al., 2024).

Despite the effect of inhibition on reward expectation at the trial level, we were

surprised to find no change in global Info preference. However, in order to equate overall

inhibition time between S+ and ITI epochs, we only inhibited ACC on 10% of the 10-s ITIs. Had

we inhibited ACC neurons on more of these ITIs, we may have observed an effect on overall

preference. In keeping with the literature, we did observe distinct behavioral phenotypes for

information choices: most rats either exhibited a strong preference for the Info option or

preferred the No-info option. Almost no animals remained indifferent to the choice options.

This divergence suggests that information may be valuable, but when there is a cost, such as via

a potential loss of reward opportunities, there are individual differences in that cost sensitivity.

Future experiments could modulate or titrate this cost sensitivity as a continuous variable.

We and others (Cox et al., 2023) have previously proposed that ACC is important in

accessing the relative value of each option, which requires a representation of both outcomes.

In support of this, we find that ACC inhibition decreased reward expectation for a certain

reward by affecting goal maintenance. Recent work by Vasquez et al. (2024) proposed that ACC

is important for attention-related processes, contributing to the ability to "stay on task"

(Vázquez et al., 2024). Recent work by Regalado et al. (2024) is also consistent with our

findings. This group investigated the role of neural activity in ACC when mice self-initiate trials

to learn cue-reward contingencies. Authors found that neural activity in OFC-to-ACC projections

increased progressively with consecutive unrewarded cues and peaked at a reward-associated

cue, keeping the animal "on track" in the task. Most relevant to the present study is that ACC

maintains the relevant goal information to allow for such goal-directed behavior.

On the effort choice task, ACC inhibition had no impact on the speed of reward

collection, but did reduce effort. This aligns with the extensive literature implicating ACC in

effort-related decision-making, supporting the idea that ACC activity is integral to sustaining

high-effort behavior for high-value rewards (Azab & Hayden, 2017; Bailey et al., 2016; Cai &

Padoa-Schioppa, 2021; Fatahi et al., 2020; Hosking et al., 2014). Consistent with what was

observed in the Info choice task, the lack of changes in latencies to collect reward suggest that

the motivational value of the reinforcer remained intact. Furthermore, the decrement in effort

during ACC inhibition was proportional to the type of test, that is, the effort was lower when an

alternative source of reward was introduced (PR Choice) compared to when only sugar pellets

were available (PR Test). If ACC inhibition was involved in altering motivation, a similar level of

effort would have been observed if either one or two sources of reward were present. Instead,

the pattern of responses was preserved and did not affect speed of retrieval, suggesting a

disruption in maintaining the goal of the better option (i.e., sucrose). We have previously

shown sucrose is indeed the better option when the choice is between chow and sucrose

pellets (Hart et al., 2020; Hart et al., 2017).

In summary, while ACC inhibition did not directly alter Info preference or reward

latencies, it significantly affected reward expectation during an informative (S+) cue and

attenuated effort-based choice. These results underscore a nuanced role for ACC in

information-seeking behavior and effort-based choice via a common stay-on-goal mechanism.

Future studies should explore more precise temporal and spatial dynamics of ACC activity to

further elucidate its role in these complex behaviors.

References

Azab, H., & Hayden, B. Y. (2017). Correlates of decisional dynamics in the dorsal anterior
cingulate cortex. *PLoS Biol*, *15*(11), e2003091.
https://doi.org/10.1371/journal.pbio.2003091
Bailey, M. R., Simpson, E. H., & Balsam, P. D. (2016). Neural substrates underlying effort, time,
and risk-based decision making in motivated behavior. *Neurobiol Learn Mem*, *133*, 233-
256. https://doi.org/10.1016/j.nlm.2016.07.015
Bromberg-Martin, E. S., & Monosov, I. E. (2020). Neural circuitry of information seeking. *Curr*
Opin Behav Sci, *35*, 62-70. https://doi.org/10.1016/j.cobeha.2020.07.006
Bussell, J. J., Badman, R. P., Márton, C. D., Bromberg-Martin, E. S., Abbott, L. F., Rajan, K., & Axel,
R. (2023). Representations of information value in mouse orbitofrontal cortex during
information seeking. *bioRxiv*, 2023.2010.2013.562291.
https://doi.org/10.1101/2023.10.13.562291
Cai, X., & Padoa-Schioppa, C. (2021). Neuronal activity in dorsal anterior cingulate cortex during
economic choices under variable action costs. *eLife*, *10*, 2021.2006.2014.448291-
442021.448206.448214.448291. https://doi.org/10.7554/eLife.71695
Cox, J., Minerva, A. R., Fleming, W. T., Zimmerman, C. A., Hayes, C., Zorowitz, S., Bandi, A.,
Ornelas, S., McMannon, B., Parker, N. F., & Witten, I. B. (2023). A neural substrate of sex-
dependent modulation of motivation. *Nat Neurosci*, *26*(2), 274-284.
https://doi.org/10.1038/s41593-022-01229-9
Cunningham, P. J., & Shahan, T. A. (2019). Rats engage in suboptimal choice when the delay to
food is sufficiently long. *J Exp Psychol Anim Learn Cogn*, *45*(3), 301-310.
https://doi.org/10.1037/xan0000211
de Melo, M. B., Daldegan-Bueno, D., Menezes Oliveira, M. G., & de Souza, A. L. (2022). Beyond
ANOVA and MANOVA for repeated measures: Advantages of generalized estimated
equations and generalized linear mixed models and its use in neuroscience research. *Eur*
J Neurosci, *56*(12), 6089-6098. https://doi.org/10.1111/ejn.15858
Fatahi, Z., Ghorbani, A., Ismail Zibaii, M., & Haghparast, A. (2020). Neural synchronization
between the anterior cingulate and orbitofrontal cortices during effort-based decision
making. *Neurobiology of Learning and Memory*, *175*, 107320-107320.
https://doi.org/10.1016/J.NLM.2020.107320
Floresco, S. B., & Ghods-Sharifi, S. (2007). Amygdala-prefrontal cortical circuitry regulates effort-
based decision making. *Cereb Cortex*, *17*(2), 251-260.
https://doi.org/10.1093/cercor/bhj143
Fortes, I., Vasconcelos, M., & Machado, A. (2016). Testing the boundaries of "paradoxical"
predictions: Pigeons do disregard bad news. *J Exp Psychol Anim Learn Cogn*, *42*(4), 336-
346. https://doi.org/10.1037/xan0000114
González, V. V., & Blaisdell, A. P. (2021). The Role of Inhibition in the Suboptimal Choice Task.
Journal of Experimental Psychology-Animal Learning and Cognition, *47*(4), 429-444.
https://doi.org/10.1037/xan0000300

González, V. V., Izquierdo, A., & Blaisdell, A. P. (2023). Theoretical Mechanisms of Paradoxical
Choices Involving Information. *Comparative Cognition & Behavior Reviews*, *18*, 11-31.
https://doi.org/10.3819/ccbr.2023.180002
González, V. V., Zhang, Y., Ashikyan, S. A., Rickard, A., Yassine, I., Romero-Sosa, J. L., Blaisdell, A.
P., & Izquierdo, A. (2024). A special role for anterior cingulate cortex, but not
orbitofrontal cortex or basolateral amygdala, in choices involving information. *Cerebral*
Cortex, *34*(4). https://doi.org/10.1093/cercor/bhae135
Hart, E. E., Blair, G. J., O'Dell, T. J., Blair, H. T., & Izquierdo, A. (2020). Chemogenetic Modulation
and Single-Photon Calcium Imaging in Anterior Cingulate Cortex Reveal a Mechanism for
Effort-Based Decisions. *J Neurosci*, *40*(29), 5628-5643.
https://doi.org/10.1523/JNEUROSCI.2548-19.2020
Hart, E. E., Gerson, J. O., Zoken, Y., Garcia, M., & Izquierdo, A. (2017). Anterior cingulate cortex
supports effort allocation towards a qualitatively preferred option. *Eur J Neurosci*, *46*(1),
1682-1688. https://doi.org/10.1111/ejn.13608
Hauber, W., & Sommer, S. (2009). Prefrontostriatal circuitry regulates effort-related decision
making. *Cereb Cortex*, *19*(10), 2240-2247. https://doi.org/10.1093/cercor/bhn241
Hosking, J. G., Cocker, P. J., & Winstanley, C. A. (2014). Dissociable contributions of anterior
cingulate cortex and basolateral amygdala on a rodent cost/benefit decision-making task
of cognitive effort. *Neuropsychopharmacology*, *39*(7), 1558-1567.
https://doi.org/10.1038/npp.2014.27
Hosokawa, T., Kennerley, S. W., Sloan, J., & Wallis, J. D. (2013). Single-neuron mechanisms
underlying cost-benefit analysis in frontal cortex. *J Neurosci*, *33*(44), 17385-17397.
https://doi.org/10.1523/JNEUROSCI.2221-13.2013
Macias, A., Gonzalez, V. V., Machado, A., & Vasconcelos, M. (2021). The functional equivalence
of two variants of the suboptimal choice task: choice proportion and response latency as
measures of value. *Anim Cogn*, *24*(1), 85-98. https://doi.org/10.1007/s10071-020-
01418-8
Mashhoori, A., Hashemnia, S., McNaughton, B. L., Euston, D. R., & Gruber, A. J. (2018). Rat
anterior cingulate cortex recalls features of remote reward locations after disfavoured
reinforcements. *eLife*, *7*. https://doi.org/10.7554/eLife.29793
McDevitt, M. A., Dunn, R. M., Spetch, M. L., & Ludvig, E. A. (2016). When good news leads to
bad choices. In *Journal of the Experimental Analysis of Behavior* (Vol. 105, pp. 23-40).
Molet, M., Miller, H. C., Laude, J. R., Kirk, C., Manning, B., & Zentall, T. R. (2012). Decision
making by humans in a behavioral task: do humans, like pigeons, show suboptimal
choice? *Learn Behav*, *40*(4), 439-447. https://doi.org/10.3758/s13420-012-0065-7
Orduña, R. E. T. a. V. (2015). Rats are optimal in a choice task in which pigeons are not.
Behavioural Processes, *119*, 22-27. https://doi.org/10.1016/j.beproc.2015.07.010
Pisklak, J. M., McDevitt, M. A., Dunn, R. M., & Spetch, M. L. (2015). When good pigeons make
bad decisions: Choice with probabilistic delays and outcomes. *J Exp Anal Behav*, *104*(3),
241-251. https://doi.org/10.1002/jeab.177
Porter, B. S., Hillman, K. L., & Bilkey, D. K. (2019). Anterior cingulate cortex encoding of effortful
behavior. *J Neurophysiol*, *121*(2), 701-714. https://doi.org/10.1152/jn.00654.2018

Regalado, J. M., Asensio, A. C., Haunold, T., Toader, A. C., Li, Y. R., Neal, L. A., & Rajasethupathy,
P. (2024). Neural activity ramps in frontal cortex signal extended motivation during
learning. *eLife*, *13*. https://doi.org/https://doi.org/10.7554/eLife.93983.2
Stagner, J. P., Laude, J. R., & Zentall, T. R. (2011). Sub-optimal choice in pigeons does not depend
on avoidance of the stimulus associated with the absence of reinforcement. *Learning*
and Motivation, *42*(4), 282-287. https://doi.org/10.1016/j.lmot.2011.09.001
Stagner, J. P., & Zentall, T. R. (2010). Suboptimal choice behavior by pigeons. *Psychon Bull Rev*,
17(3), 412-416. https://doi.org/10.3758/PBR.17.3.412
Vasconcelos, M., Monteiro, T., & Kacelnik, A. (2015). Irrational choice and the value of
information. *Sci Rep*, *5*, 13874. https://doi.org/10.1038/srep13874
Vázquez, D., Maulhardt, S. R., Stalnaker, T. A., Solway, A., Charpentier, C. J., & Roesch, M. R.
(2024). Optogenetic Inhibition of Rat Anterior Cingulate Cortex Impairs the Ability to
Initiate and Stay on Task. *The Journal of Neuroscience*, *44*(20), e1850232024.
https://doi.org/10.1523/JNEUROSCI.1850-23.2024
White, J. K., Bromberg-Martin, E. S., Heilbronner, S. R., Zhang, K., Pai, J., Haber, S. N., &
Monosov, I. E. (2019). A neural network for information seeking. *Nat Commun*, *10*(1),
5168. https://doi.org/10.1038/s41467-019-13135-z
Yu, Z., Guindani, M., Grieco, S. F., Chen, L., Holmes, T. C., & Xu, X. (2022). Beyond t test and
ANOVA: applications of mixed-effects models for more rigorous statistical analysis in
neuroscience research. *Neuron*, *110*(1), 21-35.
https://doi.org/10.1016/j.neuron.2021.10.030
593