Beneficial reversal of dominance maintains a resistance polymorphism under fluctuating insecticide selection

Marianthi Karageorgi[1,7], Anastasia S. Lyulina[1,2], Mark C. Bitter[1], Egor Lappo[1], Sharon I. Greenblum[3], Zach K. Mouza[1], Caitlynn T. Tran[1], Andy V. Huynh[1], Hayes Oken[4], Paul Schmidt[4,6,7], and Dmitri A. Petrov[1,5,6,7]

1. Department of Biology, Stanford University, Stanford, CA 94305, USA
2. Department of Applied Physics, Stanford University, Stanford, CA 94305, USA
3. Department of Energy Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA
4. Department of Biology, University of Pennsylvania, Philadelphia, PA 19104, USA
5. CZ Biohub, San Francisco, CA 94158
6. These authors contributed equally
7. Corresponding authors: M.K. (mkarag@stanford.edu), P.S. (schmidtp@upenn.edu), and D.A.P (dpetrov@stanford.edu)

Abstract

Large-effect functional genetic variation is commonly found in natural populations, even though

natural selection should erode such variants. Theory suggests that under fluctuating selective

pressures, beneficial reversal of dominance - where alleles are dominant when beneficial and

recessive when deleterious - can protect these loci from selection, allowing them to persist.

However, empirical evidence for this mechanism remains elusive because testing requires direct

measurements of selection and dominance in natural conditions. Here, we show that insecticide-

resistant alleles at the *Ace* locus in *Drosophila melanogaster* persist worldwide at intermediate

frequencies and exhibit beneficial reversal of dominance. By combining laboratory and large-

scale field mesocosm experiments with insecticide manipulation, and mathematical modeling,

we show that the benefits of the resistant *Ace* alleles are dominant while their fitness costs

recessive. We further show that fluctuating insecticide selection generates chromosome-scale

genomic perturbations at sites linked to the resistant *Ace* alleles, revealing broader genomic

consequences of this mechanism. Overall, our results suggest that beneficial reversal of

dominance contributes to the maintenance of functional genetic variation and impacts patterns of

genomic diversity via linked fluctuating selection.

Main text

Large-effect functional variants responding to temporally fluctuating selective pressures persist

in natural populations across taxa, from ancient to recently evolved polymorphisms to

anthropogenic pressures[1,2]. A striking example comes from studies of seasonal dynamics in

Drosophila, where alleles can change frequencies by tens of percent per season[3-6] yet remain

polymorphic over millions of generations[3]. However, their maintenance is puzzling because

population genetics theory suggests that the conditions for preserving functional genetic

variation under temporally fluctuating selection are restrictive[7]. One mechanism that can relax

these conditions is beneficial reversal of dominance[8,9], where alleles are partially or completely

dominant when beneficial in favorable environments but become partially or completely

recessive when deleterious in unfavorable environments[10,11]. This mechanism acts as a

stabilizing force, protecting recessive deleterious alleles from selection while enabling rapid

adaptation when selection shifts. An extension of Wright's physiological theory of dominance[12]

further suggests that this reversal can occur when alternative adaptive alleles at a locus function

as loss-of-function variants in their unfavored environments[10]. However, empirical evidence for

the mechanism remains scarce because testing requires direct measurements of selection and

dominance of adaptive alleles on fitness in natural conditions.

The *Ace* locus in insects provides an ideal genetic system to test whether beneficial

reversal of dominance can maintain functional variation under fluctuating selection[13]. *Ace*

encodes acetylcholinesterase (AChE), a key enzyme in the insect nervous system responsible for

breaking down acetylcholine[13,14]. Organophosphate insecticides, introduced in the 1950s and

1960s, act as irreversible inhibitors of AChE, causing insect death by binding to its active site

and preventing removal of acetylcholine from the synapses upon firing[15]. These insecticides have

imposed seasonally fluctuating selection pressures on resistant insect populations, as seen in

seasonal resistance fluctuations in field studies[16–18]. Following Wright's physiological theory of

dominance, resistant alleles should be loss-of-function and thus recessive in pesticide-free

environments, while sensitive alleles should become loss-of-function and thus recessive in

pesticide-rich environments[19,20]. Consistent with this prediction, previous studies across resistant

insect populations suggest that resistant *Ace* alleles follow patterns of beneficial reversal of

dominance often showing dominant benefits and/or recessive costs for fitness-associated

traits[19,21], though direct evidence is lacking.

Here, we test whether beneficial reversal of dominance under temporally fluctuating

selection contributes to the maintenance of the resistant *Ace* alleles in natural populations of *D.*

melanogaster, and investigate the broader genomic consequences of the mechanism. In

Drosophila melanogaster, resistant *Ace* alleles have repeatedly evolved in natural populations in

response to organophosphate selection[22,23], and enzymatic assays suggest the resistance

mutations are beneficial under insecticide exposure and costly in its absence[22,24]. However, the

phenotypic and fitness consequences of these mutations have not been studied. We combine

population genomic analysis, laboratory assays, and a novel field mesocosm system that enables

real-time tracking of genome-wide allele frequencies and resistance under controlled insecticide

manipulation[5] to test for beneficial reversal of dominance of the resistant *Ace* alleles. The field

mesocosm system establishes a direct experimental approach to observe beneficial reversal of

dominance and its genomic consequences in response to controlled selective pressures in semi-

natural conditions.

Resistant *Ace* alleles persist in natural populations and respond to fluctuating insecticide

selection

We first examined whether resistant *Ace* alleles exhibit signatures consistent with beneficial

reversal of dominance under temporally fluctuating selection in natural *D. melanogaster*

populations. This mechanism predicts that resistant *Ace* alleles should fluctuate in response to

temporally fluctuating selection while persisting at intermediate frequencies. Previous studies

identified resistant *Ace* alleles in *D. melanogaster* with mutations at four conserved sites near the

AChE active site (I161V, G265A, F330Y, G368A), with <u>V</u>G<u>F</u>G and <u>VA</u>Y<u>G</u> as the most common

resistance haplotypes[22,23] (inset, **Fig. 1a**). Our analysis of inbred lines from Pennsylvania

(LNPA[5]) and North Carolina (DGRP[25]) confirmed these as the most common resistant

haplotypes in resistant *Ace* alleles (**Extended Data Fig. 1a, b**), with the LNPA also carrying

<u>V</u>GF<u>A</u> at ~13% frequency. Because I161V is present across the most common haplotypes, we

used it as a shared marker to track resistant *Ace* alleles in natural populations.

While resistant *Ace* alleles experience temporally fluctuating selection, they are also

subject to spatial variation in insecticide exposure. To disentangle these effects, we analyzed

genomic data from worldwide *D. melanogaster* populations from the Drosophila Evolution Over

Space and Time (DEST) database[26] and country-level pesticide use data from the Food and

Agricultural Organization of the United Nations (FAO)[27] (see **Methods f**or details). Spatially,

I161V frequency increases with pesticide use intensity across countries (linear mixed-effect

model, $p = 0.006$) (**Fig. 1a, b**) and shows stronger spatial variation compared to control SNPs

matched for recombination and average frequency ($p = 0.005$) (**Fig. 1c**). Temporally, seasonal

sampling at Linvilla Orchards, a non-organic farm where samples were collected in spring and

fall from 2009 to 2015, revealed a consistent fall-to-spring decline in I161V frequency

(generalized linear model, $p = 0.005$), suggesting either fitness costs of the resistant *Ace* alleles in

winter (insecticide-free time period) or seasonal migration (**Fig. 1d**). In addition, the I161V

polymorphism showed stronger temporal fluctuations than matched control SNPs ($p = 0.053$)

(**Fig. 1e**). Therefore, these findings are consistent with beneficial reversal of dominance, showing

that resistant *Ace* alleles persist at intermediate frequencies under temporally fluctuating

selection.

Figure 1. Organophosphate-resistant *Ace* alleles persist at intermediate frequencies in natural *Drosophila melanogaster* populations and respond to spatially and temporally fluctuating insecticide selection. a, Allele frequency map of I161V (pie charts), a marker of organophosphate-resistant *Ace* alleles, overlaid on insecticide use intensity (shading). Inset shows common resistant *Ace* alleles with three different resistance haplotypes (R1: <u>V</u>GFG, R2: <u>V</u>GF<u>A</u>, R3: <u>VA</u>Y<u>G</u>) in *D. melanogaster* (See also **Extended Data Fig. 1** for frequencies of resistant haplotypes in DGRP and LNPA panels of fully sequenced inbred lines). **b,** I161V frequency increases with increased pesticide use (linear mixed-effect model). **c,** I161V shows higher variance in frequency compared to matched control SNPs across populations (orange dashed line indicates I161V). **d,** Seasonal changes in I161V frequency at Linvilla Orchards showing decline from fall (dark blue) to spring (pink) samples during insecticide-free periods (turquoise, mid-fall to mid-spring) (generalized linear model, mean ± s.e.m.). **e,** Cumulative frequency changes at Linvilla Orchards samples show elevated allele frequency fluctuations of I161V compared to matched control SNPs (orange dashed line indicates I161V).

Lab experiments reveal dominant benefits and recessive (or co-dominant) costs of resistant

Ace **alleles for fitness-associated phenotypes**

We then examined whether the resistant *Ace* alleles exhibit beneficial reversal of dominance for

fitness associated phenotypes relevant to organophosphate-rich and -free environments. In

organophosphate-rich environments, resistance is primarily under selection and the resistant *Ace*

alleles should be beneficial because the resistance mutations alter the shape of AChE's active site

to prevent pesticide binding[22]. In organophosphate-free environments, life history traits are

primarily under selection and the resistant *Ace* alleles should be deleterious because the

resistance mutations impair AChE activity and stability[24]. Following Wright's theory of

dominance, the resistant alleles should be dominant in organophosphate-rich environments

where sensitive alleles become loss-of-function, and recessive in organophosphate-free

environments where they themselves become loss-of-function[20] (**Fig. 2a,b**). The underlying basis

of this mechanism is that even one copy of the functional allele in each favored environment

should provide sufficient gene activity to carry out the required function.

To test these predictions, we used the LNPA panel of inbred lines (**Extended Data Fig.**

1a,b) to generate recombinant populations with homozygous sensitive, homozygous resistant,

and heterozygous genotypes for each resistant allele (S: IGFG, R1: <u>V</u>GFG, R2: <u>V</u>GF<u>A</u>, R3:

<u>VAY</u>G), an approach[28] that controls for the genomic background, and then measured resistance

to the organophosphate insecticide malathion and two key life history traits, viability (egg to

adult survival) and fecundity (number of eggs laid by gravid females) (see **Methods** for details).

For malathion resistance, the R2 and R3 alleles significantly increased LD_{50} (dose causing 50%

egg-to-adult mortality) by ~5.8x and ~7.8x respectively (p < 0.001), and were strongly dominant

over the sensitive S allele, with dominance *h* of 0.78 and 0.77 for R2 and R3, respectively (p <

0.001 for h > 0.5 in both cases). The R1 allele showed no significant resistance (p = 0.646) with

inconclusive dominance effects (p = 0.751) (**Fig. 2, Extended Data Fig. 2, Extended Data**

Table 1). For viability and fecundity, all homozygous resistant genotypes exhibited lower

phenotypic fitness values compared to the homozygous sensitive genotype, for either or both of

them, with viability reductions ranging from 12.4% to 16.9% (all p < 0.001) and fecundity

reductions from 16.2% (R2R2, p = 0.37) to 46.8% (R3R3, p < 0.001). The costs driven by the

three resistant alleles were either significantly recessive for at least one of the phenotypes for the

strongest resistance alleles R2 and R3 (viability R2: h = 0.17, p = 0.027, fecundity R3: h = -0.02,

p = 0.001), or indistinguishable from codominance for R1 (**Fig. 2g-l, Extended Data Tables 2-**

3). Therefore, these findings support the beneficial reversal of dominance of the resistant *Ace*

alleles for key fitness associated phenotypes in organophosphate-rich and -free environments.

Figure 2

Figure 2. Beneficial reversal of dominance of organophosphate-resistant *Ace* alleles for fitness-associated phenotypes relevant to malathion-rich and malathion-free environments. **a-b**, Under the beneficial dominance reversal, R alleles should be beneficial and partially or completely dominant in malathion-rich environments ($W_{SR} > (W_{SS}+W_{RR})/2$, orange), and deleterious and partially or completely recessive in malathion-free environments ($W_{SR} < (W_{SS}+W_{RR})/2$, blue). **c,** Selection (*s*) and dominance (*h*) coefficients for each fitness-associated phenotype are estimated based on the viability selection model. If s>1, the resistant R *Ace* allele is beneficial; if s<1, it is deleterious. If h=1, the resistant R *Ace* allele is dominant to the sensitive S allele; if h=0, the reverse is true. **d-l**, Comparison of malathion LD_{50} (**d,e,f**), egg-adult viability (n=10 technical replicates, 50 eggs/replicate) (**g,h,i**), and fecundity (n=8–10 technical replicates, 5 females/bottle/3 days) (**j,k,l**) among *Ace* genotypes. Malathion LD_{50} was calculated from a four-parameter logistic model fitted to egg-adult viability measurements on media with a range of malathion concentrations (n=2 technical replicates, 50 eggs/replicate, see **Extended Data Fig. 2**). Egg-adult viability and fecundity data were analyzed with logistic and linear regression models.

Field mesocosm experiments reveal dominant benefits and recessive costs of resistant *Ace*

alleles for fitness

We next tested whether beneficial reversal of dominance of the resistant *Ace* alleles drives

evolutionary responses to fluctuating insecticide selection in semi-natural conditions and

examined its genomic consequences. Given the dominant benefits and recessive costs observed

in the laboratory experiments, we predicted a rapid increase in the frequency of the resistant *Ace*

alleles and in the level of overall resistance at insecticide concentrations similar to the LD_{50} of

the sensitive homozygous genotype. We also predicted a frequency-dependent decline of the

resistant *Ace* alleles in the absence of insecticide because recessive costs become exposed in

resistant homozygotes when the resistant *Ace* alleles are at high frequencies. For the genomic

consequences, we expected that large-amplitude fluctuations of the resistant *Ace* alleles due to

fluctuating insecticide selection would drive significant effects on linked genomic variation, as

theoretical models suggest[29]. These effects on linked sites should extend across large genomic

distances, with their magnitude scaling with the rate of adaptive allele frequency shifts[30].

To test these predictions, we conducted a field mesocosm experiment to track the

evolutionary dynamics of the resistant *Ace* alleles under controlled seasonal insecticide selection

(**Fig. 3a,** see **Methods**). Using an experimental orchard in Pennsylvania, we established 20

replicate *D. melanogaster* populations derived from a baseline population constructed from the

LNPA panel of inbred lines on June 21, 2021 (**Extended Data Fig. 1**). These lines were outbred

for 4 generations in large indoor cages before ~1000 males and females were used to seed each

outdoor cage (see **Methods**). After ~4 weeks of untreated propagation, we divided the

populations into ten malathion-treated and ten untreated outdoor cages. To mimic natural

insecticide exposure, we applied stepwise increasing concentrations of commercial malathion in

the growth media of the treated populations (1.5–2.5 ppm from July 21 to August 24, followed

by 7.5 ppm until September 6, 2021), exerting strong selection for resistant *Ace* alleles based on

our lab LD_{50} measurements (**Extended Data Fig. 2**). Following insecticide removal, all

populations were maintained without malathion through December 22, 2021. Throughout the

experiment, we tracked population dynamics, malathion resistance by estimating LD_{50}, as well as

trajectories of resistant *Ace* alleles and linked sites in response to fluctuating malathion treatment

at eight timepoints (~every one to two generations) (see **Methods**).

175   Surprisingly, the malathion treatment did not alter the seasonal population dynamics in

176   the malathion-treated cages (**Fig. 3a**). In all cages, population sizes increased through the

177   summer months until reaching maximum size and then slowly declined in late Fall as abiotic

178   conditions worsened. This suggests that population size in all cages is primarily limited by high

179   levels of pre-adult mortality, potentially masking the additional mortality imposed by malathion.

180   Consistent with our predictions, the frequency of the resistant *Ace* alleles increased

rapidly in the malathion-treated cages, followed by a frequency-dependent decline after

insecticide removal. By tracking the resistant *Ace* alleles, we found that the R2 and R3 allele

frequencies increased during malathion selection (R2: t = 13.93, p < 0.001; R3: t = 27.59, p <

0.001) (**Fig. 3b**), with the speed of change consistent with the selection coefficients (*hs*) on the

order of 150-200% per generation. In contrast, the R1 allele showed no significant increase

during treatment, consistent with its lack of detectable malathion resistance in lab assays (**Fig.**

3b, Fig. 2d). After insecticide removal, only the most common resistant allele (R3) declined (t =

-5.18, p < 0.001), which reflects fitness costs that became evident at high frequencies (**Fig. 3b,**

Extended Data Table 5).

To understand how these allele frequency changes influenced population-level resistance,

we tracked malathion resistance throughout the experiment. The frequency increase of the

resistant *Ace* alleles coincided with a rapid rise in malathion resistance, with LD_{50} increasing

from ~2.2 ppm at the start of treatment to ~12.6 ppm (t = 24.26, p < 0.001) (**Fig. 3d, Extended**

Data Fig. 3). Resistance subsequently declined to ~10.6 ppm after malathion removal (t = -3.96,

p = 0.001), consistent with fitness costs in the absence of selection (**Fig. 3d, Extended Data Fig.**

3, Extended Data Table 4.1). To assess the contribution of the *Ace* locus in the evolution of

resistance, we applied a viability selection model parametrized by lab-measured genotype

viabilities (see **Methods**). This model closely recapitulated both allele frequency trajectories and

resistance evolution, strongly suggesting that *Ace* is the major-effect locus driving rapid

evolution of resistance to malathion selection (**Extended Data Fig. 4**).

In the absence of insecticide selection in the untreated cages, the resistant *Ace* allele

frequencies remained stable supporting the prediction that their costs are hidden in

heterozygosity (**Fig. 3c**). None of the resistant *Ace* alleles exhibited significant frequency

changes (**Fig. 3c, Extended Data Table 6**), suggesting low selection coefficients (*s*) or

dominance parameters (*h*) in the absence of malathion. Consistently, LD_{50} values remained low

and unchanged throughout the experiment (~2.2 ppm, t = 1.07, p = 0.288) (**Fig. 3d, Extended**

Data Fig. 3, Extended Data Table 4.2), indicating that overall resistance levels were unaffected

in the untreated populations.

To further obtain bounds on evolutionary parameters for the resistant *Ace* alleles in

organophosphate-free conditions, we fit the allele frequency data of the treated and untreated

cages to a viability selection model where all resistant alleles share the same fitness cost (see

Methods). The model that fit both datasets best had low cost (*s* ~ 2.5 to 5%) and low dominance

values ($h < 0.38$) (**Fig. 3f**). At higher values of h, there is no single selection coefficient that can

explain the patterns of allele frequency change for both treated and untreated cages, and the

likelihood of the data is higher the closer h is to zero. This analysis confirmed our prediction that

the fitness cost of the resistant *Ace* alleles is recessive in semi-natural conditions.

Figure 3

Figure 3. Beneficial reversal of dominance of resistant *Ace* alleles for fitness in an ecologically realistic environment. a, Twenty field mesocosm cages were seeded with an outbred population of *D. melanogaster* generated via outcrossing for 4 generations 76 inbred lines carrying the sensitive S and resistant R1, R2, and R3 alleles. Ten cages received a malathion pulse after the initial population establishment. Eight timepoints were sampled to track the evolution of malathion resistance and genome-wide allele in the treated and untreated populations. Malathion did not affect estimated adult fly population sizes (mean ± s.e.m.) in treated populations. **b-c,** Trajectories of *Ace* alleles in treated (**b**) and untreated (**c**) populations. In treated populations, the malathion pulse led to a significant increase in the frequency of the resistant R2 and R3 alleles, followed by a decline in the frequency of R3. *Ace* allele frequencies remained stable in the untreated populations. Light lines/points show individual allele trajectories, dark lines/points represent the mean ± s.e.m. (n = 10 populations per treatment group). **d-e,** The malathion pulse led to a significant increase and subsequent decline of resistance (LD$_{50}$, ppm) in the treated populations while resistance remained stable in the untreated populations. For LD$_{50}$ calculations see **Methods** and **Extended Data Fig. 3**. (**continued on next page**)

Figure 3 (continued). Light lines/points show individual populations (orange, treated; blue, untreated), dark lines/points represent mean ± s.e.m. (n=10 populations per treatment group).**f,** Inferred selection (s) and dominance (h) coefficients from changes in the resistant *Ace* allele frequency. The colored ranges show the selection coefficients corresponding to the 0.95 confidence intervals of the slope distribution, obtained by bootstrapping cages within each treatment group. The overlap between the orange (treated cages) and dark blue (untreated cages) restricts the possible values of *s* and *h*. Within that overlap, lower values of *h* better explain both sets of data, as shown by higher values of the joint likelihood calculated from the bootstrap distributions (see **Methods**). All results of statistical tests can be found in **Extended Data Tables 4-6**.

Fluctuating insecticide selection generates chromosome-scale genomic perturbations at

sites linked to the resistant *Ace* alleles

We then examined the genomic consequences of rapid frequency fluctuations at the *Ace* locus on

linked genetic regions. Our experimental design presented us with a rare opportunity to study

these effects directly because the linkage patterns of genome-wide SNPs with the *Ace* alleles at

the beginning of the experiment are known from the inbred line genotypes. To characterize

forward and reverse sweeps driven by the R2 and R3 alleles during and after malathion

treatment, respectively, we used a two-step approach (see **Methods**). First, we employed

quasibinomial logistic regression to estimate genome-wide allele frequency changes during

malathion treatment (July 26 to September 21, 2021) and post-treatment (September 21 to

December 22, 2021). Next, we quantified changes in allele frequencies in SNPs tightly linked to

the sweeping R2 and R3 alleles across the genome (r2 > 0.03 and frequency > 60% in resistant

lines, *Ace*-linked) and those that were largely unlinked (r2 < 0.01, control).

During malathion treatment, we observed a chromosome-wide forward sweep in the

malathion-treated cages driven by the R2 and R3 resistant alleles (**Fig. 4a**). The sweep extended

across the chromosome arms 3R (~4-17Mbp), including the *Ace* locus (9.6Mb), and 3L (~5-

19Mbp), with *Ace*-linked sites showing consistently elevated effect sizes above control sites. The

sweep's width is consistent with the extremely rapid rise of the *Ace* alleles over 2-4 generations

and the maximum recombination rate of ~5cM/Mbp in *D. melanogaster*, implying the effects

conservatively extending over a region of ~5-10 Mbp. Our analysis also indicated the presence

of two additional loci potentially responding to malathion selection. On chromosome arm 3L

(~19.5Mb), control sites showed a strong directional perturbation while *Ace*-linked sites

uniformly decreased in frequency, suggesting interference from another adaptive locus with the

Ace sweep (**Fig. 4a,** asterisk**)**. A similar pattern was observed on chromosome arm 2R, where

both *Ace*-linked and control sites showed comparable frequency perturbations, indicating a

second putative locus under selection (**Fig. 4a** asterisk). The study of these additional adaptive

loci to malathion treatment is beyond the scope of this work. As expected, in the untreated cages

where the frequencies of the R2 and R3 alleles remained stable (**Fig. 3d**), *Ace*-linked sites

showed no elevated frequency changes compared to the control sites (**Fig. 4b**).

After malathion treatment ceased and selection reversed, we detected an ~5Mbp reverse

sweep driven by the R2 and R3 alleles in the malathion-treated cages (**Fig. 4c**). Individual SNPs

within 1Mbp of the *Ace* locus demonstrated a strong reversal effect, in which sites that increased

in frequency during treatment tended to decrease afterwards, with this reversal being

significantly stronger for *Ace*-linked compared to control sites (**Fig. 4e**). As expected, in the

untreated populations, *Ace*-linked sites showed neither significant median frequency changes

(**Fig. 4d**) nor the reversal behavior (**Fig. 4f**). Notably, recombination through the course of the

experiment degraded the initial haplotype structure, suggesting the true extent of the reverse

sweep may exceed 5 Mbp. Together, these findings confirmed our predictions that large-

amplitude frequency fluctuations of adaptive alleles under temporally varying selection can drive

significant effects on linked genomic variation.

Figure 4

Figure 4. Strong fluctuating selection of resistant *Ace* alleles caused chromosome-scale forward sweep during malathion treatment and reverse sweep after treatment. **a-d**, Effect sizes (median allele frequency changes) for *Ace*-linked (orange/blue) and unlinked control (grey) SNPs in 1Mbp windows across the genome were estimated using quasibinomial logistic regression (see **Methods**). Changes during malathion treatment in treated (**a**) and untreated (**b**) populations. Changes after treatment cessation in treated (**c**) and untreated (**d**) populations. The brackets represent the inferred width of the sweep. Stars indicate additional adaptive loci to malathion treatment. **e-f**, Comparison of effect size for *Ace*-linked (orange/blue) and unlinked control (grey) SNPs during (y-axis) and after malathion treatment (x-axis) in malathion-treated (**e**) and untreated (**f**) populations within 1Mb from *Ace*. Each panel shows SNPs grouped by their distance from *Ace* (in Mb): [-1, -0.5], [-0.5, 0], [0, 0.5], and [0.5, 1]. In panel **e**, red stars indicate regions where the regression slopes significantly differ between *Ace*-linked and unlinked control SNPs.

Discussion

Our study provides compelling empirical evidence for beneficial dominance reversal maintaining

a large-effect resistance polymorphism at the *Ace* locus under temporally fluctuating insecticide

selection in *D. melanogaster*. Previous studies have suggested the presence of beneficial reversal

of dominance of resistant *Ace* alleles in *D. melanogaster* and other resistant insect populations

based on the extension of Wright's physiological theory[20], enzymatic studies[22,24], and analysis of

fitness-associated phenotypes[19]. While these approaches can provide insights into the functional

effects of resistance mutations, they do not necessarily predict organismal and fitness

consequences[31]. For example, our results with the R1 allele illustrate this limitation - while

enzymatic studies predicted it would confer similar levels of resistance as the R2 allele to

malaoxon (the active breakdown product of malathion)[22], we found a very small resistance

benefit to malathion at the organismal and fitness levels. The novelty of our work with the *Ace*

locus in *D. melanogaster* lies in combining laboratory assays, field mesocosm experiments with

insecticide manipulation, and mathematical modeling to directly measure selection and

dominance effects on the organismal level and on fitness in semi-natural conditions.

The beneficial dominance reversal we observed at the *Ace* locus suggests this mechanism

may be widespread in maintaining functional genetic variation at large-effect loci under

temporally fluctuating selection. Indeed, a growing body of literature indicates that beneficial

reversal of dominance acts as a common mechanism for maintaining large-effect genetic

variation across different forms of antagonistic selection between environments (e.g., fitness

components, sexes, or niches/environments)[10,11]. For example, one allele of the large-effect

RXFP2 locus in Soay sheep has a dominant beneficial effect on male mating success but a

recessive deleterious effect on male survival[32]. Similarly, the late maturation allele of the large-

effect *VGLL3* locus in Atlantic salmon is partially dominant and likely beneficial in females, yet

recessive and likely deleterious in males[33]. In the context of temporally fluctuating selection,

large-effect loci coding for protein polymorphisms (e.g., enzymes, transcription factors) that are

specialized for alternative contexts/substrates may commonly display beneficial reversal of

dominance, as environmental shifts render these variants functionally impaired in their unfavored

conditions. This pattern follows naturally from Wright's physiological theory of dominance[12],

which predicts that the loss of protein function in alternative environments leads to the shifting

identity of the recessive, deleterious allele[10,12].

Finally, our study examined the genomic consequences of beneficial reversal of

dominance at the *Ace* locus under temporally fluctuating selection. Our field mesocosm

experiment revealed substantial genomic effects, with the rapid frequency changes at the *Ace*

locus driving both a chromosome-wide forward sweep during malathion treatment and a reverse

sweep after treatment removal. These findings provide empirical validation of theoretical

predictions that beneficial dominance reversal of large-effect alleles can generate large-

amplitude frequency fluctuations with significant effects on linked genetic variation[2,29]. Given

that the *D. melanogaster* genome contains a fair number (tens to hundreds) of large-effect alleles

displaying large frequency shifts due to fluctuating selection[5,6] and the likelihood that this is the

case for many other species[34,35], fluctuating selection at large-effect loci could act as a major

force shaping levels of genetic variation in natural populations[29].

Acknowledgements This project was supported by grants from the National Institute of General

Medical Sciences of the National Institutes of Health (Award number 1K99GM143455-01 to

M.K, Grant 5R01GM100366-08 and 5R01GM137430-02 to P.S., Award Number

5R35GM118165-07 to D.A.P), the Sarah Hotchkis Ketterer Graduate Fellowship from Stanford

University (to A.S.L.), the National Science Foundation (Grant PRFB 2109407 to M.C.B.), and

the Stanford Biology Summer Undergraduate Research Program (to Z.K.M. and A.H.). The

authors thank Marcus Feldman for helpful discussions and assistance with modeling *Ace* allele

frequency evolution, Danae Papadopetraki for assistance with statistical analyses of the

phenotypic and genomic data, Axel Wiberg for feedback on statistical analyses of the genomic

data, Emily Behrman for feedback on the day degree model for generation estimation, Jessica

Rhodes for helpful discussions and assistance with statistical analyses of the DEST data, and

Anastasia Andreeva for assistance with phenotyping. The authors are very grateful to Jean Villa,

Jessica Rhodes, Meike Wittmann, Noah Whiteman, and Marcus Feldman for feedback that

improved the earlier versions of the manuscript.

Author Contributions M.K was responsible for conceptualization, investigation, formal analysis, writing original draft, writing-review and editing, visualization, and funding acquisition; A.L. was responsible for formal analysis, visualization, and writing-review and editing; M.B. was responsible for formal analysis and investigation; E.L. was responsible for formal analysis, visualization, and writing-review and editing; S.G. was responsible for formal analysis; A.H., CT, and Z.K.M. were responsible for investigation; P.S. and D.A.P. jointly supervised the research. P.S. was responsible for conceptualization, investigation, formal analysis, writing-review and editing, and funding acquisition. D.A.P. was responsible for conceptualization, investigation, formal analysis, writing-review and editing, and funding acquisition.

Methods

Population genomic analysis for variation of organophosphate-resistant *Ace* alleles across space and time

We first conducted a literature review to compile all known naturally segregating target site resistance mutations in the acetylcholinesterase (*Ace* gene) protein of *D. melanogaster* that confer resistance to organophosphates. Mutero and colleagues in 1994[36] first identified the resistance mutations F115S, I199V, I199T, G303A and F368Y, and numbered them based on the precursor acetylcholinesterase (AChE) protein. Menozzi and colleagues in 2004[22] identified combinations of the resistance mutations I161(199)V, G265(303)A, F330(368)Y and G368(406)A in worldwide organophosphate-resistant *D. melanogaster* strains and numbered them based on the mature AChE protein (numbers in the brackets refer to the precursor AChE protein). They did not identify the resistance mutation F77(115)S. At position 77, the mutation TTC to T**C**C changes phenylalanine to serine (Chr3R: 9,070,101, v5.39). At position 161, the mutation ATC to **G**TC changes the isoleucine to valine (Chr3R:9,069,721, v5.39). At position 265, the mutation GGC to G**C**C changes the glycine to alanine (Chr3R:9,069,408 , v5.39). At position 330, the mutation TTT to T**A**T changes the phenylalanine to tyrosine (Chr3R:9,069,054 , v5.39). At position 368, the mutation GGC to G**C**C/G**CT** changes the glycine to alanine (Chr3R: 9,063,921 , v5.39). The AChE protein's numbering changed between the studies after its crystallization in 2000, which defined the N-terminal signal peptide and the C-terminal sequence[37]. The mutation G368A was missanotated in the study of adaptation of *D. melanogaster* to organophosphate insecticides by Karasov and colleagues[23].

Organophosphate-resistant *Ace* alleles in DGRP and LNPA inbred lines

We characterized the presence of organophosphate-resistant *Ace* alleles in two fully

sequenced inbred line panels, the Drosophila Genetic Reference Panel (DGRP) and our LNPA

panel of inbred lines. The DGRP panel consists of 205 inbred lines derived from wild-caught

females collected at the Raleigh, North Carolina Farmers Market in 2003[25]. The LNPA panel

consists of 76 inbred lines derived from wild-caught individuals collected in Spring and Fall

from Linvilla Orchards, Media PA USA[5]. Both panels contain the two common resistant *Ace*

alleles (Ace-R1: VGFG and Ace-R3: VAYG) at high frequency, the sensitive *Ace* allele (Ace-S:

IGFG), and other rare resistant alleles. The LNPA panel also contains the resistant *Ace* allele

Ace-R2: VGFA at high frequency. We used the LNPA panel for our lab and field mesocosm

experiments.

<u>Spatial and seasonal variation of organophosphate-resistant *Ace* alleles in natural *Drosophila*</u>

<u>*melanogaster* populations</u>

To examine the spatial variation of the I161V polymorphism, a marker of common

organophosphate-resistant *Ace* alleles (Ace-R1: VGFG, Ace-R2: VGFA, Ace-R3: VAYG), we

first analyzed the relationship between country-level pesticide use and I161V frequency in

Drosophila melanogaster populations. We obtained pesticide use data from the Food and

Agriculture Organization (FAO) FAOSTAT database[27], which provides country-level estimates

of pesticide use per area of cropland. For population genomic data, we used the Drosophila

Evolution over Space and Time (DEST) database, which contains pooled genome-wide allele

frequency estimates from 753 population samples of *D. melanogaster* collected worldwide. For

our analyses, we included pesticide use data and genomic data for the period of 2000-2022.

To visualize the spatial distribution of insecticide use and frequency of the I161V

polymorphism across locations, we created a choropleth map where countries are shaded based

on their pesticide use per cropland area, with pie charts indicating the frequency of the resistant

(I161V) and sensitive alleles at sampled locations. For each location, we calculated median allele

frequencies across available samples for the period 2000-2022. To test whether insecticide use

predicts the frequency of the resistance allele I161V, we fitted a linear mixed effects model using

the lmer() function in the R package lme4 (version 1.1.33). The model included I161V frequency

as the response variable and insecticide use per cropland area as a fixed effect. We included a

random effect for cities.

In addition, to further test whether I161V shows evidence of spatial selection rather than

drift, we compared its variance across samples collected in spring to matched control SNPs. We

selected control SNPs based on three criteria: (1) located within 2Mb but at least 100kb away

from *Ace* to avoid linked SNPs, (2) within the ln(3R)K cosmopolitan inversion to match

inversion status, and (3) average frequency across samples within 4% of the median frequency of

the I161V resistance mutation. We used the cumulative distribution function (CDF) to quantify

how extreme the I161V variance was compared to the control SNPs.

To examine the temporal variation of the I161V polymorphism, we first tested two

hypotheses: (1) whether the frequency of the resistant allele decreases during winter months

when insecticide pressure is relaxed, potentially due to fitness costs, and (2) whether the

frequency of the resistant allele increases during the growing seasons when insecticides are

applied. We analyzed time series data from Linvilla Orchards, a non-organic orchard in

Pennsylvania, USA. Samples were collected near apple trees from 2009 to 2015, with paired

spring and fall collections each year, allowing us to test both hypotheses. To account for

sampling variance in pooled sequencing data, we calculated the effective number of

chromosomes sampled (Nc) following Machado et al. (2016), where $Nc = (1/N + 1/R)^{-1}$, with

N being the number of chromosomes in the pool ($2 \times$ number of flies) and R the read depth at the

site. We then tested for systematic seasonal changes using generalized linear models (GLM) with

binomial error structure. The models used the number of resistant and susceptible alleles

(weighted by Nc) as the response variable and season as a categorical predictor. For the winter

decline hypothesis, we paired each fall sample with the following spring sample. For the summer

increase hypothesis, we paired each spring sample with the following fall sample from the same

year. This approach allowed us to directly test for consistent directional changes in allele

frequencies across seasons while accounting for both biological and technical sources of variance

in the data.

407   Finally, to test whether the temporal dynamics of I161V reflect selection rather than drift,

we compared its frequency fluctuations to those of matched control SNPs at Linvilla. To

quantify temporal variation, we calculated the cumulative absolute change in I161V allele

frequency over time as the sum of frequency differences between consecutive timepoints. This

metric captures the total magnitude of frequency changes, regardless of direction. We selected

control SNPs that had similar starting conditions as I161V, with initial frequencies matching the

I161V frequency (0.42) at the first Linvilla timepoint within ±4%. To ensure a valid comparison,

control SNPs were selected from within the same genomic region and inversion status as I161V,

while excluding potentially linked sites (same criteria as for the spatial analysis). We then used

the cumulative distribution function to assess whether I161V's temporal variation was

significantly greater than that observed in the matched control SNPs, which would suggest that

its frequency changes are driven by seasonal selection rather than drift.

Laboratory experiments to estimate the dominance of resistant *Ace* alleles for fitness-

associated phenotypes in the presence and absence of malathion

To assess whether the dominance of each resistant *Ace* allele (Ace-R1: VGFG, Ace-R2: VGFA,

Ace-R3: VAYG) over their sensitive counterpart (Ace-S: IGFG) reverses for fitness-associated

phenotypes that are primarily under selection in organophosphate-rich and -free environments,

we first used our LNPA panel of inbred lines to generate recombinant populations with

homozygous sensitive, homozygous resistant, and heterozygous *Ace* genotypes for each *Ace*

allele. This step was meant to bring potential deleterious, recessive mutations in the inbred lines

into heterozygosity. We then measured malathion resistance, viability and fecundity in

homozygous and heterozygous populations for each *Ace* genotype.

Generation of recombinant populations with homozygous sensitive, homozygous resistant, and

heterozygous *Ace* genotypes

To generate recombinant populations for each *Ace* allele, we first intercrossed 2-3 inbred

lines with the same *Ace* allele using a round robin cross scheme (e.g., Line_1 SS x Line_2 SS,

Line_2 SS x Line_3 SS, Line_3 SS x Line_1 SS) to bring all founders with the same *Ace* allele

together (10 non-mated females x 10 males per round robin cross). We then combined 16 F1

flies (8 non-mated females and 8 males) from each of the round robin crosses per *Ace* allele to

establish the next generation. Subsequently, we maintained the homozygous populations in vials

and allowed them to recombine for four generations until we performed the phenotype

measurements. We used lines 12LN6-70_(B33), 12LN6-92_(B3), LNPA-3 for the sensitive *Ace*

allele (Ace-S: IGFG), lines 12LN6-86_(B29) and LNPA-53 for the R1 resistant *Ace* allele (Ace-

R1: VGFG), lines LNPA-14, LNPA-17 and LNPA-87 for the R2 resistant *Ace* allele (Ace-R2:

VGFA), and lines 12LN6-45_(B26), 12LN6-82_(B4), and 12LN6-94_(B19) for the R3 resistant

Ace allele (Ace-R3: VAYG).

<u>Measurements of fitness-associated phenotypes</u>

For all experiments, we first set up breeding cages with ~ 500 4-to-7-day-old flies of

homozygous or heterozygous populations for the sensitive and the resistant *Ace* alleles. To

generate the heterozygous recombinant populations, we crossed SS non-mated females with

homozygous males of each resistant allele (R1R1, R2R2, R3R3).

To compare the level of malathion resistance among the different alleles, we measured

egg-to-adult viability on diets that ranged in malathion concentration from 0 ppm to 30 ppm. For

this, we collected eggs of each population that were laid within a period of 24 hours and then

introduced 50 eggs of each population into narrow vials containing approximately 7 ml of

Drosophila growth media supplemented with the respective malathion concentration. We tested

each of the malathion concentrations in duplicate vials. We used commercial malathion of 50%

strength (Southern Ag Malathion 50% E.C.) and dissolved it in the *Drosophila* growth media.

This medium was kept overnight at 4°C and used the next day for the assay. We monitored adult

eclosion every evening over a period of 10 days until all adult flies had enclosed.

To compare viability among the different genotypes, we introduced 50 eggs of each

population into narrow vials containing ~7 ml of *Drosophila* growth media. Each of the

genotypes were tested in 10 replicate vials. We monitored adult eclosion every evening over a

period of 7 days until all adult flies had enclosed.

To compare fecundity among the different genotypes, we used density-controlled (~50

eggs/vial) adult female progeny of each population. We introduced five 3-day-old females of

each genotype in egg-laying bottles fitted with egg-laying caps filled with Drosophila growth

media. Each day over a period of 3 days we counted the number of eggs laid. Each of the

genotypes were tested in 8-10 replicate egg-laying bottles. The egg-laying caps were examined

under a stereoscope where we counted the eggs laid each day over the 3-day period.

<u>Statistical analysis of fitness-associated phenotypes</u>

We assessed fecundity, viability, and malathion resistance for the three genotypes (SS,

SR, RR) of each *Ace* allele (R1, R2, R3) using R (version 4.2.2). For fecundity and viability, we

fitted linear and logistic regression models, respectively, calculating estimated marginal means

and performing pairwise comparisons adjusted for multiple testing using the Benjamini-

Hochberg procedure. We then implemented a bootstrap procedure (1,000 resamples) to estimate

deviation from additivity, selection coefficient, and dominance coefficient. The deviation from

additivity was calculated as W[SR] - (W[SS] + W[RR]) / 2, where W represents fitness

(fecundity or viability) for each genotype. The selection coefficient s was computed as (W[RR] –

W[SS]) / W[SS], indicating the relative fitness of the resistant genotype compared to the

sensitive genotype. The dominance coefficient h was calculated as (W[SR] – W[SS]) / (W[RR] –

W[SS]), measuring the degree to which the heterozygote phenotype is closer to either

homozygote. Significance was assessed by testing whether results differed from zero (additivity

and selection) or 0.5 (dominance). For malathion resistance, we estimated LD_{50} values for each

genotype using four-parameter logistic dose-response models $f(x) = c + (d-c) / (1 + e^{\wedge}(b(x-e)))$,

where e is $LD_{50,}$ d (upper limit) = 100%, c (lower limit) = 0%, and b is the relative slope around

e. For this, we used the L.4() function in the drc package (version 3.0.1)[38] in R (version 4.2.2)

with egg to adult viability at the range of malathion concentrations as the response variable. We

then used a bootstrap procedure (1,000 resamples) to compare LD_{50} values between genotypes

and calculate the same coefficients, substituting LD_{50} values for W in the formulas. For all

analyses, bootstrap resampling was used to calculate confidence intervals and p-values, with

significance assessed against null hypotheses of 0 (additivity and selection) and 0.5 (dominance).

Field mesocosm experiments to track the evolution of malathion resistance and genome-

wide allele frequencies in the presence and absence of fluctuating malathion selection

Experimental orchard and field mesocosm cages

To mimic the natural surroundings and weather conditions that flies are exposed to in nature, we

conducted the study in an experimental orchard at the University of Pennsylvania in

Philadelphia. The orchard consisted of cleared land with walk-in mesocosm cages (2m x 2m x

2m) constructed of fine mesh and built around metal frames (BioQuip PO 1406C). We planted a

dwarf peach tree in each mesocosm cage to provide flies with shade. All peaches were removed

before ripening to prevent flies from feeding or laying eggs on them. To provide flies with a

feeding and breeding substrate, we added 1.5lb aluminum loaf pans with 400ml Drosophila

growth media every two days during the experiment. To avoid an overcrowded developmental

environment for the eggs, each loaf pan was covered with a screen mesh lid every two days

during the experiment, and eggs were allowed to develop until eclosion ceased, at which point

adult flies were released into the general population. The loaf pans were placed on a shelving

unit oriented away from direct sunlight to protect them from rain and direct sun.

Establishment of experimental *D. melanogaster* populations in field mesocosm cages

To construct the baseline *D. melanogaster* population for the experiment, we crossed the

76 fully sequenced LNPA inbred lines from Pennsylvania. We used inbred lines to facilitate the

use of haplotype inference to attain high effective sequencing coverage[39]. To initiate the baseline

population, we combined 10 females and 10 males from each line into a single breeding cage.

After 4 generations of mating and density-controlled rearing in laboratory conditions, we

introduced 500 females and 500 males of a single age cohort into each mesocosm cage on June

21, 2021.

Malathion treatment over the course of the experiment

A diagram of the experimental design is shown in **Fig. 3a**. In the first step (*Establishment*

phase: June 21 to July 20, 2021), we established 20 replicate populations in mesocosm cages and

allowed them to expand in the absence of interventions. After establishment, the populations

were split into ten malathion-treated and ten malathion-untreated populations. To mimic the

insecticide exposure of flies in nature, the malathion-treated populations were exposed to

malathion in the summer. In nature, insecticides are mainly applied during summer for pest

control and their application ceases in the fall[40]. During the malathion treatment phase of the

experiment, we sequentially exposed the populations to stepwise increasing concentrations of

malathion, from 1.5 (July 21 to July 27, 2021) to 2.5ppm (July 28 to August 24, 2021) and

7.5ppm (August 25 to September 6, 2021). We used commercially available malathion grade of

50% strength (Southern Ag Malathion 50% E.C.) and dissolved it in the *Drosophila* growth

media. We selected malathion doses based on the Malathion 50% E.C. label instructions to

control small insect pests in the field and a previous long-term selection experiment for

malathion resistance in *D. melanogaster*[41]. The medium supplemented with malathion was

cooked once per week and stored at 4˚C until deployed. After the malathion treatment, the

experiment lasted until early winter (September 7 to December 22, 2021). Over the course of the

experiment, we performed high-resolution temporal sampling (~ bi-weekly sampling during the

summer and monthly during the fall) to track evolution of malathion resistance and genome-wide

allele frequencies.

Tracking population size in the field mesocosms

To make relative comparisons of population sizes between the malathion-treated and untreated

cages, we photographed 4 equal-size quadrats of the ceiling in each cage at dusk (approximately

2.5% of the ceiling for each quadrat). To count the number of adult flies in each photograph, we

designed and validated an algorithm for semi-automated image processing. The number of adult

flies in each of the 4 photographs was multiplied by 40 to correct for total surface area and obtain

4 population size estimates per cage.

<u>Statistical analysis of population data</u>

To assess the effects of time and malathion treatment on population size, we fitted a linear

mixed-effect model with the estimated population size as the response variable. We defined our

model to investigate the quadratic relationship of time with population size and to account for the

random effects attributable to differences in cages. The regression analysis was performed in R

(version 4.2.2) using lme4 (version 1.1.33).

Tracking malathion resistance in the field mesocosms

To characterize the rate and direction of the evolution of malathion resistance in the malathion-

treated and untreated populations, we repeatedly assayed the evolving populations. Prior to the

malathion resistance assay, we collected F1 eggs from each mesocosm cage and transferred them

in 2 groups of 200 eggs to bottles with media (i.e., density controlled) in a constant common

garden environment (25°C, 12L:12D, 50% humidity) in the laboratory to distinguish evolution

from phenotypic plasticity. We then performed the malathion resistance assay with density and

age-controlled individuals from each mesocosm cage in the F2 generation (25°C, 12L:12D, 50%

humidity). The common garden experiment for assaying malathion resistance took place at

Stanford University with F1 eggs in bottles with media shipped overnight from the University of

Pennsylvania.

Malathion resistance measurements

To characterize the level of malathion resistance in the malathion-treated and untreated

populations, we tracked egg to adult survival in the F2 generation on diets that ranged in

malathion concentration from 0 ppm to 15 ppm over the course of the experiment. For the

collection of F2 eggs, we placed ~ 400 4-to-7-day-old F1 individuals in breeding cages with egg

lay caps for 5-7 hours. We then introduced 50 F2 eggs in a narrow vial containing ~7ml of

Drosophila growth media supplemented with malathion. We tested each of the malathion

concentrations in duplicate vials. We used commercial malathion of 50% strength (Southern Ag

Malathion 50% E.C.) and dissolved it in the *Drosophila* growth media. This medium was kept

overnight at 4°C and used the next day for the assay. We monitored adult eclosion every evening

over a period of 10 days until all adult flies had eclosed.

Statistical analysis of malathion resistance data

To assess the temporal dynamics of malathion resistance in both malathion-treated and untreated

cages, we fitted separate linear mixed-effects models using the lmer() function in the R package

lme4 (version 1.1.33). The response variable in these models was malathion resistance, measured

as the LD_{50} value, the dose that causes 50% egg-to-adult mortality. For the malathion-treated

cages, we modeled resistance changes over two distinct time periods: (1) from July 26, 2021, to

September 21, 2021, when populations were evolving under malathion treatment, and (2) from

September 21, 2021, to December 22, 2021, after malathion treatment had ceased. In contrast,

for the malathion-untreated cages, we assessed resistance changes over the full experimental

period, from July 26, 2021, to December 22, 2021. To avoid biases, we excluded data from July

13, 2021, because the LD_{50} estimate from this time point was underestimated due to a technical

error. In each model, we included time point as fixed effect, while a random intercept was

included for each cage to account for between-cage variability. The main outcome measure of

malathion resistance for each cage at each time point was the mean of 1–2 technical replicate

LD_{50} measurements. To estimate LD_{50} for each technical replicate per cage per time point, we

fitted four-parameter logistic dose-response models using the L.4() function in the drc package

(version 3.0.1) in R (version 4.2.2) with egg to adult viability at the range of malathion

concentrations as the response variable.

Track genome-wide allele frequencies in the field mesocosms over time

To track the trajectories of the resistant and sensitive *Ace* alleles and characterize the extent of

the sweep caused by the *Ace* alleles, we tracked genome-wide allele frequencies of the evolving

populations across the experiment using the software tool HAF-pipe. HAF-pipe estimates

genome-wide allele frequencies in the evolving populations via inference of known founder

haplotypes from low-coverage pooled sequencing data [39]. As part of our experimental design, we

performed whole-genome, low-coverage pooled sequencing of 100 randomly selected F1

females from each of the 20 evolving populations at eight time points through the experiment. To

measure biological and technical variation, we performed whole-genome, low-coverage pooled

sequencing of different sets of 100 randomly selected F1 females from the 20 evolving

populations at two time points collected at Stanford (biological replicates) and of same samples

of 100 randomly selected F1 females but processed independently (technical replicates). Last, to

evaluate the accuracy of HAF-pipe, we performed whole-genome, high-coverage pooled

sequencing of four biological replicate samples from the baseline population and for the evolving

populations at an early and a late time point. The data for the untreated cages were also used for

a recent independent study from Bitter and colleagues [6].

<u>Sample collection, DNA extraction, library preparation, and sequencing</u>

We collected samples from each population approximately every two generations during

the summer phase and every generation during the fall phase. To collect the samples, we placed a

new loaf pan with Drosophila growth media in each mesocosm cage one day before collection.

The next day, we transferred five groups of 200 F1 eggs to bottles with media, where we

controlled the density and maintained them in a constant laboratory environment of 25°C,

12L:12D, and 50% humidity. Three bottles with F1 eggs were kept at UPenn and two bottles

with F1 eggs were shipped overnight to Stanford. At Stanford, F1 adults were allowed to lay

eggs, and 200 F2 eggs were placed in vials for density control. For both the F1 and F2

generation, we harvested 4-to-7-day-old individuals after eclosion and preserved them in 100%

ethanol in 1.5ml vials at -20°C until further processing.

For each sample, we prepared genomic DNA from pools of 100 ethanol-preserved

females. We first submerged the samples twice in a rehydration buffer (400 mM NaCl, 20 mM

Tris-HCl pH 8.0, 30 mM EDTA) for 15 min at room temperature and then isolated genomic

DNA using the Monarch Genomic DNA purification kit (Cat No. 3010L, New England Biolabs).

The manufacturer's protocol was modified to include a step for tissue homogenization with two

3mm glass beads (Cat No. 11-312C, Fisher Scientific) in a bead beater for five minutes to ensure

complete homogenization of the flies so that the genomic DNA from each fly was equally

represented in the genomic DNA pool. We quantified the genomic DNA using the Qubit dsDNA

BR (Broad-Range) Assay kit (Cat No. Q32853, Life Technologies) on a Qubit

spectrophotometer and estimated its purity on a NanoDrop spectrophotometer.

We then prepared sequencing libraries using the Illumina DNA prep, (M) Tagmentation

kit (Cat No. 20018705, Illumina Inc.) and the Illumina UD Indexes sets (Cat No 20027123,

20027214, 20042666, 20042667, Illumina Inc.) for unique dual indexing. We modified the

manufacturer's protocol by preparing libraries in 1/5 of the recommended reaction volume to

maximize kit efficiency and pooling equimolarly amplified libraries in sets of eight after the

library amplification step. After the cleanup, we quantified the pooled libraries using a Qubit

spectrophotometer and an Agilent Bioanalyzer. Finally, the pooled libraries were 150bp paired-

end sequenced on the NovaSeq 6000 S2 platform at the Sequencing Service Center at the

Stanford Genomics Center and on a NovaSeq 6000 SP platform at Admera. All sequences will

be deposited in SRA.

Processing of DNA sequencing data for genome-wide allele frequency estimation

We trimmed the raw sequencing reads of adapter sequences and bases with quality score

< 20. We then mapped the trimmed reads to the *D. melanogaster* reference genome (v5.39) using

bwa and default parameters [42], and aligned reads were deduplicated using Picard tools. We

randomly sampled reads from the aligned bams to obtain an equivalent, genome-wide coverage

of 8x across all samples. We computed haplotype-derived allele frequency (HAF) estimates

using HAF pipe. Briefly, after using an expectation-maximization algorithm to estimate

haplotype frequencies in overlapping genomic windows from mapped sequence reads (originally

developed by Kessner and colleagues [43]) at each site previously identified as polymorphic in the

inbred reference panel, we summed the estimated frequencies of haplotypes carrying the

alternate allele to obtain HAFs . We conducted haplotype inference in window sizes that varied

proportionally to the expected length of un-recombined haplotype blocks, which varied as a

function of the estimated number of generations since the original outbreeding. HAFs were

compiled to only include those sites with an average minor allele frequency (MAF) > 0.02 in the

baseline population, and present in at least one evolved sample at a MAF > 0.01.

664
665   **Track trajectories of sensitive and resistant *Ace* alleles in the evolving populations over**

666   **time**

667
We first estimated the frequency of each of the 76 founder haplotypes in the *Ace* region

(Chr3R:9,063,921-9,069,721) in each of the malathion-treated and malathion-untreated

populations at each time point, focusing on the founders containing the four known resistance

mutations . To obtain an accurate measure of the frequency of each haplotype, we calculated the

mean frequency across the 10 overlapping genomic windows that included each site within the

region of interest.

We then estimated the relative frequency of each *Ace* allele, defined by the combination

of the four known resistance mutations, in each population at each time point. Given our

knowledge of the resistance mutation combinations for each founder, we computed the relative

frequency of each *Ace* allele (Ace-S: IGFG, Ace-R1: <u>V</u>GFG, Ace-R2: <u>V</u>GF<u>A</u>, Ace-R3: <u>VA</u>Y<u>G</u>,

Ace-R4:IGF<u>A</u>, Ace-R5:IG<u>Y</u>G, Ace-R6:<u>VA</u>FG, Ace-R7: <u>VA</u>Y<u>A</u>) by summing the frequencies of

the haplotype blocks from the 76 founders that shared the same *Ace* allele. For founders

heterozygous for the *Ace* alleles, we divided the contributions of their haplotype block evenly

between the two corresponding *Ace* allele frequencies. *Ace* alleles with a frequency < 0.01%

(Ace-R4:IGF<u>A</u>, Ace-R5:IG<u>Y</u>G, Ace-R6:<u>VA</u>FG, Ace-R7: <u>VA</u>Y<u>A</u>) in the evolving populations

were excluded from further analysis.

<u>Statistical analysis of *Ace* allele frequency data</u>

To assess the temporal dynamics of *Ace* allele frequencies (Ace-S: IGFG, Ace-R1:

<u>V</u>GFG, Ace-R2: <u>V</u>GF<u>A</u>, Ace-R3: <u>VA</u>Y<u>G</u>), we fitted separate linear mixed-effects models using

the lmer() function in the R package lme4 (version 1.1.33). The response variables in these

models were the frequencies of each *Ace* allele. For the malathion-treated cages, we modeled *Ace*

allele frequency changes over two distinct time periods: (1) from July 26, 2021, to September 21,

2021, when populations were evolving under malathion treatment, and (2) from September 21,

2021, to December 22, 2021, after malathion treatment had ceased. In contrast, for the

malathion-untreated cages, we assessed *Ace* allele frequency changes from the beginning of the

treatment to the end of the experiment, from July 26, 2021 to December 22, 2021.

Model *Ace* allele frequency evolution

We simulated the evolution of *Ace* allele frequencies and malathion resistance in our mesocosm

experiments using a viability selection model[44]. Additionally, based on the changes in *Ace* allele

frequencies in malathion-free environments across both sets of cages, we estimated the possible

ranges for the fitness costs of and dominance of malathion resistance.

The impact of viability on *Ace* allele frequency dynamics and malathion resistance in the field

mesocosm experiment

We investigated whether selection acting on viability recapitulates the allele frequency

and malathion resistance dynamics observed in the field. We focused on viability as it is a key

fitness-related trait in both malathion-rich and malathion-free environments. Furthermore, we

had laboratory measurements of viabilities for *Ace* genotypes across ranges of malathion

concentrations, allowing us to parametrize our model and test its predictions.

Within our model, the population frequencies of the *Ace* alleles change only due to selection that

acts on viability. We assumed that the effects of other forces (mutation, recombination, and

genetic drift) were negligible in comparison over the timescales that we considered (~10

generations). Under these assumptions, the frequency dynamics of the sensitive (f_0) and resistant

(f_1, f_2, f_3) *Ace* alleles follow from

715
$$f_i(t+1) = \frac{f_i(t) \sum_j w_{ij} f_j(t)}{\overline{w}(t)},$$

716 where t is time measured in generations, w_{ij} is viability of individuals carrying both i and j

717 alleles, and $\underline{w}$ is the mean population viability defined as

718
$$\overline{w}(t) = \sum_{i,j} f_i(t) f_j(t) w_{ij}.$$

We used laboratory measurements to determine individual viabilities w_{ij} in the absence

and presence of malathion (at 2.5 and 7.5 ppm). We used the marginal mean values from the

linear model for viability for each genotype and the predicted viability values from the dose-

response analysis (see previous section on 'Statistical analysis of fitness-associated phenotypes').

Due to the lack of measurements for heterozygous resistant genotypes, we assumed that

combinations of resistant alleles are codominant, i.e. that $w_{12} = 0.5 \cdot w_{11} + 0.5 \cdot w_{22}$; $w_{13} = 0.5 \cdot$

$w_{11} + 0.5 \cdot w_{33}$; $w_{23} = 0.5 \cdot w_{22} + 0.5 \cdot w_{33}$. The initial *Ace* allele frequencies, $f_i(0)$, were

estimated as the median values from the first sampling time point (July 13, 2021) for each

treatment group.

To estimate *D. melanogaster* generation times at our eight sampling time points, we used

a degree-day model based on Behrman and Schmidt[53]. This model assumes *D. melanogaster*

requires 112 degree-days to complete one generation, with development occurring between 12°C

and 29°C. We calculated daily degree-days as [(daily max. temp. + daily min. temp.) / 2] - 12°C,

adjusting temperatures outside the 12-29°C range to these limits. Cumulative degree-days were

tracked, with each 112 degree-day accumulation marking a new generation. This allowed us to

estimate generations at each sampling time point throughout the experiment.

We then calculated the population resistance R as a weighted sum,

736
$$R(t) = \sum_{i,j} g_{ij}(t) w_{ij},$$

737   where $g_{ij}(t)$ are genotype frequencies and w_{ij} are the corresponding viabilities. The initial

738   genotype frequencies were estimated from the initial allele frequencies, assuming Hardy-

739   Weinberg equilibrium:

740
$$g_{ij}(0) = f_i(0) f_j(0).$$

741   The consequent genotype frequencies followed from the viability selection model,

742
$$g_{ij}(t) = \begin{cases} \dfrac{2f_i(t)f_j(t)w_{ij}}{\overline{w}}, i \neq j, \\ \dfrac{f_i(t)f_j(t)w_{ij}}{\overline{w}}, i = j. \end{cases}$$

<u>Inferring the ranges of possible fitness costs and dominance coefficients from empirical allele</u>

<u>frequency trajectories</u>

To estimate the fitness cost and dominance of malathion resistance in the malathion-free

environment, we analyzed the empirical frequency trajectories of these alleles in both the

untreated and treated populations after the removal of malathion. This approach allowed us to

constrain the range of possible parameters that could explain the observed frequencies of the

resistant alleles in both the treated and untreated populations. Assuming the resistant alleles share

the same selection (s) and dominance (h) coefficients, and further that the cost of resistance is

small (s<<1), the frequency dynamics follow

753                        $\partial_t f = s(f \cdot (1 - f) \cdot (f + h(1 - 2f)))$,

allowing us to find s that would be consistent with the observed slope $\partial_t f$ of the frequency

trajectory for a given h[54].

To do that, we first bootstrapped the combined frequency trajectories of resistant alleles

in individual cages within each treatment group 10,000 times. For each set of bootstrapped

trajectories, we used linear regression (LinearRegression().fit function from the scikit-learn

Python library[55] to infer their slope. For a range of $h \in [0,1]$, we then calculated the bounds on s

by finding the values that correspond to the lower and upper bound of the 95% confidence

interval of the bootstrap distribution. To further constrain the range of possible parameters, we

similarly inferred the full distribution of s for each set of cages, and evaluated their overlap by

computing the joint likelihood function,

$$\mathcal{L} = \prod p(s),$$

where $p(s)$ is the probability density corresponding to a fixed value of s. Larger values of L therefore correspond to more likely combinations of h and s given the observed frequency dynamics.

Characterize extent of forward and reverse sweep caused by *Ace* resistant alleles

To assess how fluctuating organophosphate selection on the R2 and R3 *Ace* resistant alleles affected genome-wide frequency trajectories of linked SNPs during and post-malathion treatment, we employed a two-step approach. First, we used quasibinomial logistic regression to estimate allele frequency shifts for genome-wide SNPs during malathion treatment (July 26 to September 21, 2022) and post-treatment (September 21 to December 22, 2021). Second, we developed a linked SNP selection procedure using linkage information from the inbred lines to identify SNPs linked (Ace-linked) and not linked (control) to the sweeping R2 and R3 *Ace* alleles across the genome. We used these sets of SNPs to quantify the extent of both the forward *Ace* sweep during malathion treatment and the reverse *Ace* sweep post-treatment.

<u>Quasi-binomial logistic regression to estimate allele frequency differences in malathion-treated and untreated populations during and after the malathion treatment</u>

To quantify allele frequency changes during malathion treatment (July 26 to September 21, 2022) and post-treatment (September 21 to December 22, 2021) in both treated and untreated populations, we used a quasibinomial logistic regression approach[5,48]. This method models the number of reads mapped to the alternative allele 'α' (relative to the reference allele 'A') at each biallelic SNP for each pooled sample. To account for both the expected accuracy of HAFs as

787 well as pseudo-replication due to a fixed number of individuals in each sample [49], we estimated

788 an effective read depth reads (at each biallelic SNP for each pooled sample per population using

789
$$n_{eff} = \frac{((2 \times N \times EC) - 1)}{2 \times N + EC}$$

790 where N is the number of individuals per sample (100) and EC is the expected effective read

791 coverage based on the accuracy of HAF-pipe.

792 We then used a logit link implemented in a quasi-binomial logistic regression to model the

793 counts of the alternative allele 'α' ($C_a = HAF * neff$) and reference allele 'A' ($C_A = (1 - $

794 $HAF) * neff)$ as a function of time in the malathion-treated and -untreated populations

795
$$logit(p_a) = log \left(\frac{p_a}{1 - p_a}\right) = b_0 + b_1 \times time + e$$

796 where p_a is probability of observing the allele 'α', while the coefficients b_0 and b_1 correspond to

797 the intercept and the effect of time, respectively. We fitted the model using the glm() function in

798 R (version 4.2.2) with family = 'quasibinomial'. To compute the allele frequency change (effect

799 size) between two time points (x_1 and x_2), we calculated

800
$$\Delta p = p_2 - p_1 = \frac{e^{(b_0 + b_1 \times x_2)}}{1 + e^{(b_0 + b_1 \times x_2)}} - \frac{e^{(b_0 + b_1 \times x_1)}}{1 + e^{(b_0 + b_1 \times x_1)}}$$

<u>Analysis of allele frequency changes of Ace-linked SNPs</u>

To quantify the extent of the forward and reverse *Ace* sweeps, we identified SNPs linked

to the sweeping R2 and R3 *Ace* alleles (*Ace*-linked), as well as unlinked (control) SNPs, using

whole genome data from the 76 sequenced Pennsylvania inbred lines (LNPA). First, we labeled

R2 and R3 *Ace* alleles as "R" and the remaining alleles (R1 and S) as "S", creating a binary

"R"/"S" *Ace* locus across the inbred lines. Among the inbred lines, 54 were homozygous "S", 5

heterozygous "R/S", and 17 homozygous "R". To select linked SNPs, we used the inbred lines

data to compute r^2 between the binary "R"/"S" *Ace* locus and every reported biallelic SNP in

the inbred LNPA lines as the squared correlation coefficient of allelic indicator variables[50]. We

initially retained SNPs with r2 > 0.03, a threshold frequency previously used to deem two loci

highly linked[6]. However, to avoid issues with low-frequency SNPs in finite populations, we

added an allele count condition. We selected SNPs with r2 > 0.03 that also have an allele present

in ≥60% of "R/R" lines and in ≤40% of "S/S" and "S/R" lines. This ensures selection of high-

frequency, well-assorted SNPs between "R" and "S" lines, which are more informative for sweep

analysis. This approach yielded 5672 Ace-linked SNPs across chromosomes 2, 3, and X. The

60% threshold was chosen as the highest value providing enough Ace-linked SNPs to uniformly

cover the genome, with consistent results when varied. Additionally, for each Ace-linked SNP,

we selected an unlinked control SNP with r2 < 0.01 with the binary "R"/"S" *Ace* locus. To ensure

comparable initial conditions, unlinked control SNPs were chosen within 500kb of their linked

counterpart and had an allele frequency within 0.05 of the Ace-linked SNP at the start of

malathion treatment in each cage. Finally, to ensure that we track the SNP alleles that are linked

specifically with the resistant lines, we transformed the alleles frequencies based on the sign of r

— the non-squared correlation coefficient of allelic indicator variables. If r between *Ace* and a

SNP is positive, then the alternative allele is assorted together with the resistant *Ace* alleles in the

inbred lines, if the sign of r is negative, then the reference allele is linked to resistant *Ace* alleles,

and in this case the alternative/reference labels were flipped, and the allele frequencies f_t were

transformed to 1-f_t. This process yielded 4793 matched unlinked control SNPs, fewer than the

linked SNPs due to the stringent matching criteria. To visualize the results, we aggregated SNPs

into 1Mb windows along the genome and used bootstrap resampling to compute median GLM-

estimated effect sizes and confidence intervals for Ace-linked and unlinked control SNPs

separately for both forward and reverse sweeps. We excluded the areas around the centromeres

and chromosome ends where less than 20 *Ace*-linked or control SNPs were observed. To

characterize the width of each sweep, we performed a two-sample Mann-Whitney U-test within

every 1Mb window across the chromosome and recorded the width of the contiguous range of

windows around *Ace* in which the median for *Ace*-linked SNPs is significantly greater than the

median for control SNPs at the 0.05 confidence level.

To quantify the reversal phenomenon during the *Ace* sweep post-malathion treatment, we

analyzed allele frequency changes for the same Ace-linked and unlinked control SNPs during the

post-treatment period (September 21 to December 22, 2021). We binned SNPs into

500Kb windows based on the genomic distance from the *Ace* locus on chromosome 3R and then

compared post-treatment to during-treatment frequency changes by plotting them against each

other within each bin. To quantify the reversal, we performed a joint linear regression for Ace-

linked and control SNPs and recorded which bins have shown a slope for Ace-linked SNPs

significantly more negative compared to control at the 0.05 confidence level.

Data and Code availability

The data and associated code for this study will become available after manuscript acceptance in

public repositories: Dryad (https://doi.org/10.5061/dryad.w0vt4b937) for the data and GitHub

(https://github.com/mkarageorgi/Beneficial_Dominance_Reversal) for the code. The sequences

of the LNPA inbred lines used to construct the baseline *D. melanogaster* population for the field

mesocosm experiment are available on NCBI under accession ID PRJNA722305. The raw

sequencing reads for pooled samples from the malathion-treated cages, collected during the field

mesocosm experiment, will be deposited in the SRA BioProject under accession number

PRJNA1031645. The raw sequencing reads for pooled samples from the untreated cages are

already available in the same SRA BioProject (PRJNA1031645).

Extended Data Figure 1

b,

Ace haplotype	LNPA relative frequency (%)	DGRP relative frequency (%)
IGFG	63.16	54.15
VGFA	13.16	0.49
VAYG	12.50	30.73
VGFG	7.89	12.20
IGYG	1.32	0.98
IGFA	0.66	NA
VAFG	0.66	NA
VAYA	0.66	NA
IAFG	NA	1.46

Extended Data Figure 1. Sensitive and resistant *Ace* alleles in the panel of LNPA and DGRP inbred lines. a, Heatmap showing clustering of seventy-six *D. melanogaster* isogenic lines in the LNPA panel based on the sensitive or resistant *Ace* allele they carry (Ace-S: IGFG, Ace-R1: <u>V</u>GFG, Ace-R2: <u>V</u>GF<u>A</u>, Ace-R3: <u>VAY</u>G). The x-axis in the heatmap shows the genomic position on chromosome 3R around the *Ace* locus. Each row represents the *Ace* haplotype information of one isogenic line. Each column represents one biallelic exonic SNP at the *Ace* locus. Reference alleles are colored in dark grey, alternative alleles are colored in light grey, heterozygotes in medium grey. For the resistance mutations, alleles coding for the resistance mutations are colored in orange, alternative in grey, and heterozygotes in yellow. **b,** Table showing frequency of the sensitive and each of the resistant *Ace* alleles carrying different resistance haplotypes in the panels of LNPA and DGRP inbred lines.

Extended Data Figure 2

a

b

Dose response curve analysis for each Ace genotype

Genotype	LD50	std.error	statistic	p.value
SS	1.39	0.18	7.79	3.7768e-12
R1R1	1.56	0.20	7.68	6.3407e-12
R2R2	8.16	0.40	20.41	7.7563e-40
R3R3	10.81	0.45	23.96	3.3625e-46
SR1	2.34	0.19	12.49	3.0000e-23
SR2	6.80	0.26	25.89	2.1374e-49
SR3	8.84	0.30	29.95	1.3764e-55

Extended Data Figure 2. Comparison of malathion resistance among the *Ace* genotypes. a, Lab measurements of egg to adult viability in recombinant populations carrying different *Ace* genotypes on standard Drosophila media with a range of malathion concentrations (n= 2 technical replicates per genotype for each malathion doses, 50 eggs per replicate). Four-parameter logistic models fitted to data of each *Ace* genotype. The shaded areas highlight viability of each genotype at malathion doses (2.5ppm and 7.5 ppm) applied in the field mesocosm experiment. **b,** Table with estimated LD_{50} values for each *Ace* genotype from four-parameter logistic dose-response model (see also **Methods**).

Extended Data Figure 3

Extended Data Figure 3. Tracking evolution of malathion resistance in the field mesocosm experiment. Egg-adult survival on diets with a range of malathion concentrations as measured after two generations of common garden rearing throughout the field mesocosm experiment. Light lines and points show the mean egg-adult survival for individual malathion-treated (orange) and untreated (blue) populations (n = 2 technical replicates, 50 eggs per replicate). Dark lines and points represent the mean ± s.e.m. from all populations for each treatment group (n = 10 populations per treatment group). Further information on experimental design can be found in **Methods**.

Extended Data Figure 4

Extended Data Figure 4. Modeling the impact of viability selection on *Ace* allele frequency dynamics and malathion resistance in the field mesocosm experiment. **a-b,** Comparison between the observed *Ace* allele frequency trajectories over 10 generations in the malathion-treated (**a**) and untreated (**b**) populations from the field mesocosm experiment (mean ± s.e.m.) and the expected trajectories from simulations based on a viability selection model using laboratory-derived parameters. See **Methods** for details about the viability selection model. **c-d,** Comparison between the observed egg-to-adult survival across a range of malathion concentrations (**c**) and the evolution of malathion resistance (LD_{50}, ppm) (**d**) over 10 generations in the field mesocosm experiment (mean ± s.e.m.), and the expected results from the same model simulations. Dashed lines represent the expected values from the viability selection model, while solid lines represent the observed data from the field experiment. Generations were estimated based on a degree-day model starting from the first sampling point (July 13, 2021). In all plots, points correspond to each generation and estimated time point, and are connected by lines for visualization.

Extendend Data Table 1

Effect estimates from linear model analysis for LD50

Parameter	LD50	Std.Error	P.value
SS mean	1.39	0.19	-
SR1 mean	2.34	0.13	-
R1R1 mean	1.56	0.14	-
SS vs. SR1	-0.93	0.34	0.0063
SS vs. R1R1	-0.17	0.37	0.646
SR1 vs. R1R1	0.76	0.39	0.0516
SR1 - (SS + R1R1)/2 (additivity)	0.76	0.32	1e-16
(R1R1 - SS) / R1R1 (selection, s)	-0.25	0.29	0.751
(SR1 - SS) / (R1R1 - SS) (dominance, h)	-1.29	5.10	0.751
SR2 mean	6.80	0.24	-
R2R2 mean	8.16	0.36	-
SS vs. SR2	-5.38	0.36	0
SS vs. R2R2	-6.72	0.42	0
SR2 vs. R2R2	-1.34	0.45	0.003
SR2 - (SS + R2R2)/2 (additivity)	1.87	0.35	1e-16
(R2R2 - SS) / R2R2 (selection, s)	3.83	1.05	1e-16
(SR2 - SS) / (R2R2 - SS) (dominance, h)	0.78	0.06	1e-16
SR3 mean	8.84	0.32	-
R3R3 mean	10.81	0.49	-
SS vs. SR3	-7.41	0.36	0
SS vs. R3R3	-9.37	0.79	0
SR3 vs. R3R3	-1.96	0.80	0.0145
SR3 - (SS + R3R3)/2 (additivity)	2.59	0.48	1e-16
(R3R3 - SS) / R3R3 (selection, s)	5.64	1.45	1e-16
(SR3 - SS) / (R3R3 - SS) (dominance, h)	0.77	0.07	1e-16

Extendend Data Table 2

Effect estimates from logistic model analysis for viability

Parameter	Viability	Std.Error	Lower.CI	Upper.CI	P.value
SS mean	0.89	0.01	-	-	-
SR1 mean	0.86	0.02	-	-	-
R1R1 mean	0.76	0.02	-	-	-
SS vs. SR1	1.42	0.27	-	-	0.0702
SS vs. R1R1	2.61	0.47	-	-	<.0001
SR1 vs. R1R1	1.84	0.30	-	-	4e-04
SR1 - (SS + R1R1)/2 (additivity)	0.03	0.02	-0.02	0.07	0.214
(R1R1 - SS) / R1R1 (selection, s)	-0.15	0.03	-0.2	-0.09	0.001
(SR1 - SS) / (R1R1 - SS) (dominance, h)	0.29	0.20	-0.15	0.64	0.214
SR2 mean	0.87	0.02	-	-	-
R2R2 mean	0.74	0.02	-	-	-
SS vs. SR2	1.28	0.25	-	-	0.205
SS vs. R2R2	2.96	0.53	-	-	<.0001
SR2 vs. R2R2	2.31	0.39	-	-	<.0001
SR2 - (SS + R2R2)/2 (additivity)	0.05	0.02	0.01	0.1	0.027
(R2R2 - SS) / R2R2 (selection, s)	-0.17	0.05	-0.28	-0.09	0.001
(SR2 - SS) / (R2R2 - SS) (dominance, h)	0.17	0.15	-0.15	0.45	0.027
SR3 mean	0.86	0.02	-	-	-
R3R3 mean	0.78	0.02	-	-	-
SS vs. SR3	1.42	0.27	-	-	0.0702
SS vs. R3R3	2.43	0.44	-	-	<.0001
SR3 vs. R3R3	1.72	0.29	-	-	0.0018
SR3 - (SS + R3R3)/2 (additivity)	0.02	0.02	-0.02	0.06	0.286
(R3R3 - SS) / R3R3 (selection, s)	-0.13	0.02	-0.18	-0.08	0.001
(SR3 - SS) / (R3R3 - SS) (dominance, h)	0.32	0.19	-0.15	0.67	0.286

Extendend Data Table 3

Effect estimates from linear model analysis for fecundity

Parameter	Fecundity	Std.Error	Lower.CI	Upper.CI	P.value
SS mean	200.22	11.99	-	-	-
SR1 mean	165.33	14.01	-	-	-
R1R1 mean	136.50	14.86	-	-	-
SS vs. SR1	34.89	19.81	-	-	0.14
SS vs. R1R1	63.72	20.42	-	-	0.01
SR1 vs. R1R1	28.83	20.42	-	-	0.17
SR1 - (SS + R1R1)/2 (additivity)	-3.03	17.26	-36.62	31.59	0.879
(R1R1 - SS) / R1R1 (selection, s)	-0.32	0.09	-0.48	-0.12	0.003
(SR1 - SS) / (R1R1 - SS) (dominance, h)	0.55	0.40	-0.07	1.32	0.881
SR2 mean	187.60	13.64	-	-	-
R2R2 mean	167.78	14.37	-	-	-
SS vs. SR2	12.62	19.81	-	-	0.53
SS vs. R2R2	32.44	20.33	-	-	0.37
SR2 vs. R2R2	19.82	19.81	-	-	0.49
SR2 - (SS + R2R2)/2 (additivity)	3.60	18.42	-33.15	43.16	0.834
(R2R2 - SS) / R2R2 (selection, s)	-0.16	0.09	-0.33	0.03	0.093
(SR2 - SS) / (R2R2 - SS) (dominance, h)	0.39	4.09	-2.3	3.23	0.842
SR3 mean	202.00	12.71	-	-	-
R3R3 mean	106.44	11.99	-	-	-
SS vs. SR3	-1.78	17.47	-	-	0.92
SS vs. R3R3	93.78	16.95	-	-	<.0001
SR3 vs. R3R3	95.56	17.47	-	-	<.0001
SR3 - (SS + R3R3)/2 (additivity)	48.67	12.84	22.41	73.56	1e-16
(R3R3 - SS) / R3R3 (selection, s)	-0.47	0.07	-0.59	-0.31	0.001
(SR3 - SS) / (R3R3 - SS) (dominance, h)	-0.02	0.19	-0.48	0.29	0.001

Extendend Data Table 4

Extended Data Table 4.1 Effect of time on malathion resistance (LD50) in treated cages

	During treatment				After treatment			
Predictors	**Estimates**	**std. Error**	**Statistic**	**p**	**Estimates**	**std. Error**	**Statistic**	**p**
Intercept	-3.02 ***	0.46	-6.58	**<0.001**	15.56 ***	1.09	14.29	**<0.001**
Time	2.48 ***	0.10	24.26	**<0.001**	-0.57 ***	0.14	-3.96	**0.001**
Random Effects								
σ^2		0.91				0.98		
τ_{00}		0.00 Treatment_Cage				0.24 Treatment_Cage		
ICC						0.20		
N		10 Treatment_Cage				10 Treatment_Cage		
Observations		40				30		
Marginal R^2 / Conditional R^2		0.938 / NA				0.302 / 0.441		
AIC		122.513				101.066		

*p<0.05 ** p<0.01 *** p<0.001*

Extended Data Table 4.2 Effect of time on malathion resistance (LD50) in untreated cages

	Beginning of treatment to end of experiment			
Predictors	**Estimates**	**std. Error**	**Statistic**	**p**
Intercept	2.27 ***	0.15	14.91	**<0.001**
Time	0.03	0.02	1.07	0.288
Random Effects				
σ^2		0.16		
τ_{00} Treatment_Cage		0.03		
ICC		0.15		
N Treatment_Cage		10		
Observations		59		
Marginal R^2 / Conditional R^2		0.017 / 0.166		
AIC		83.651		

*p<0.05 ** p<0.01 *** p<0.001*

Extendend Data Table 5

Extended Data Table 5.1 Effect of time on S (IGFG) frequency in treated cages

Predictors	During treatment				After treatment			
	Estimates	std. Error	Statistic	p	Estimates	std. Error	Statistic	p
Intercept	0.88 ***	0.02	49.15	**<0.001**	0.04	0.05	0.74	0.468
Time	-0.13 ***	0.00	-30.81	**<0.001**	0.02 *	0.01	2.64	**0.014**
Random Effects								
σ^2		0.00				0.00		
τ_{00}		0.00 $_{cage}$				0.00 $_{cage}$		
N		10 $_{cage}$				10 $_{cage}$		
Observations		50				30		
Marginal R^2 / Conditional R^2		0.951 / NA				0.194 / NA		
AIC		-148.876				-97.774		

* p<0.05   ** p<0.01   *** p<0.001

Extended Data Table 5.2 Effect of time on R1 (VGFG) frequency in treated cages

Predictors	During treatment				After treatment			
	Estimates	std. Error	Statistic	p	Estimates	std. Error	Statistic	p
Intercept	0.08 ***	0.00	16.13	**<0.001**	0.04 **	0.01	3.09	**0.005**
Time	-0.00 ***	0.00	-4.05	**<0.001**	0.00	0.00	0.46	0.651
Random Effects								
σ^2		0.00				0.00		
τ_{00}		0.00 $_{cage}$				0.00 $_{cage}$		
ICC						0.25		
N		10 $_{cage}$				10 $_{cage}$		
Observations		50				30		
Marginal R^2 / Conditional R^2		0.251 / NA				0.005 / 0.250		
AIC		-276.043				-162.492		

* p<0.05   ** p<0.01   *** p<0.001

Extended Data Table 5.3 Effect of time on R2 (VGFA) frequency in treated cages

Predictors	During treatment				After treatment			
	Estimates	std. Error	Statistic	p	Estimates	std. Error	Statistic	p
Intercept	0.01	0.01	0.93	0.356	0.14 **	0.04	3.39	**0.002**
Time	0.04 ***	0.00	13.93	**<0.001**	0.02 **	0.01	3.03	**0.006**
Random Effects								
σ^2		0.00				0.00		
τ_{00}		0.00 $_{cage}$				0.00 $_{cage}$		
N		10 $_{cage}$				10 $_{cage}$		
Observations		50				30		
Marginal R^2 / Conditional R^2		0.798 / NA				0.240 / NA		
AIC		-178.169				-107.043		

* p<0.05   ** p<0.01   *** p<0.001

Extended Data Table 5.4 Effect of time on R3 (VAYG) frequency in treated cages

Predictors	During treatment				After treatment			
	Estimates	std. Error	Statistic	p	Estimates	std. Error	Statistic	p
Intercept	0.03	0.01	1.88	0.067	0.78 ***	0.05	15.18	**<0.001**
Time	0.09 ***	0.00	27.59	**<0.001**	-0.04 ***	0.01	-5.18	**<0.001**
Random Effects								
σ^2		0.00				0.00		
τ_{00}		0.00 $_{cage}$				0.00 $_{cage}$		
ICC		0.04						
N		10 $_{cage}$				10 $_{cage}$		
Observations		50				30		
Marginal R^2 / Conditional R^2		0.937 / 0.940				0.481 / NA		
AIC		-170.172				-96.128		

* p<0.05   ** p<0.01   *** p<0.001

Extendend Data Table 6

Extended Data Table 6.1 Effect of time on S (IGFG) haplotype frequency in untreated cages

Predictors	Beginning of treatment to end of experiment			
	Estimates	std. Error	Statistic	p
Intercept	0.63 ***	0.02	39.49	**<0.001**
Time	0.00	0.00	0.21	0.838
Random Effects				
σ^2		0.00		
$\tau_{00 \ cage}$		0.00		
ICC		0.03		
N_{cage}		10		
Observations		69		
Marginal R^2 / Conditional R^2		0.001 / 0.032		
AIC		-196.762		

*p<0.05 ** p<0.01 *** p<0.001*

Extended Data Table 6.2 Effect of time on R1 (VGFG) haplotype frequency in untreated cages

Predictors	Beginning of treatment to end of experiment			
	Estimates	std. Error	Statistic	p
Intercept	0.06 ***	0.00	14.68	**<0.001**
Time	0.00	0.00	1.91	0.061
Random Effects				
σ^2		0.00		
$\tau_{00 \ cage}$		0.00		
N_{cage}		10		
Observations		69		
Marginal R^2 / Conditional R^2		0.051 / NA		
AIC		-370.061		

*p<0.05 ** p<0.01 *** p<0.001*

Extended Data Table 6.3 Effect of time on R2 (VGFA) haplotype frequency in untreated cages

Predictors	Beginning of treatment to end of experiment			
	Estimates	std. Error	Statistic	p
Intercept	0.07 ***	0.01	9.08	**<0.001**
Time	-0.00	0.00	-0.02	0.985
Random Effects				
σ^2		0.00		
$\tau_{00 \ cage}$		0.00		
ICC		0.02		
N_{cage}		10		
Observations		69		
Marginal R^2 / Conditional R^2		0.000 / 0.024		
AIC		-285.686		

*p<0.05 ** p<0.01 *** p<0.001*

Extended Data Table 6.4 Effect of time on R3 (VAYG) haplotype frequency in untreated cages

Predictors	Beginning of treatment to end of experiment			
	Estimates	std. Error	Statistic	p
Intercept	0.24 ***	0.01	17.99	**<0.001**
Time	-0.00	0.00	-0.86	0.392
Random Effects				
σ^2		0.00		
$\tau_{00 \ cage}$		0.00		
ICC		0.05		
N_{cage}		10		
Observations		69		
Marginal R^2 / Conditional R^2		0.010 / 0.064		
AIC		-222.940		

*p<0.05 ** p<0.01 *** p<0.001*

References

1.   Hedrick, P. W. Genetic polymorphism in heterogeneous environments: The age of

genomics. *Annu. Rev. Ecol. Evol. Syst.* **37**, 67–93 (2006).

2.   Johnson, O. L., Tobler, R., Schmidt, J. M. & Huber, C. D. Fluctuating selection and the

determinants of genetic variation. *Trends Genet.* **39**, 491–504 (2023).

3.   Bergland, A. O., Behrman, E. L., O'Brien, K. R., Schmidt, P. S. & Petrov, D. A. Genomic

evidence of rapid and stable adaptive oscillations over seasonal time scales in Drosophila.

PLoS Genet. **10**, e1004775 (2014).

4.   Machado, H. E. *et al.* Broad geographic sampling reveals the shared basis and

environmental correlates of seasonal adaptation in Drosophila. *Elife* **10**, (2021).

5.   Rudman, S. M. *et al.* Direct observation of adaptive tracking on ecological time scales in

Drosophila. *Science* **375**, eabj7484 (2022).

6.   Bitter, M. C. *et al.* Continuously fluctuating selection reveals fine granularity of adaptation.

Nature **634**, 389–396 (2024).

7.   Haldane, J. B. S. & Jayakar, S. D. Polymorphism due to selection of varying direction. *J.*

Genet. **58**, 237–242 (1963).

8.   Hedrick, P. W. Genetic variation in a heterogeneous environment. II. Temporal

heterogeneity and directional selection. *Genetics* **84**, 145–157 (1976).

9.   Wittmann, M. J., Bergland, A. O., Feldman, M. W., Schmidt, P. S. & Petrov, D. A.

Seasonally fluctuating selection can maintain polymorphism at many loci via segregation

lift. *Proc Natl Acad Sci U S A* **114**, E9932–E9941 (2017).

10.   Connallon, T. & Chenoweth, S. F. Dominance reversals and the maintenance of genetic

variation for fitness. *PLOS Biology* **17**, e3000118 (2019).

11. Grieshop, K., Ho, E. K. H. & Kasimatis, K. R. Dominance reversals: the resolution of

genetic conflict and maintenance of genetic variation. *Proceedings of the Royal Society B*

(2024) doi:10.1098/rspb.2023.2816.

12. Wright, S. Physiological and evolutionary theories of dominance. *Am. Nat.* **68**, 24–53

(1934).

13. Fournier, D. & Mutero, A. Modification of acetylcholinesterase as a mechanism of

resistance to insecticides. *C, No. I, pp.* **19**, 31 (1994).

14. Hall, J. C. & Kankel, D. R. Genetics of acetylcholinesterase in Drosophila melanogaster.

Genetics **83**, 517–535 (1976).

15. Aldridge, W. N. Some properties of specific cholinesterase with particular reference to the

mechanism of inhibition by diethyl p-nitrophenyl thiophosphate (E 605) and analogues.

Biochem. J. **46**, 451–460 (1950).

16. Lenormand, T., Bourguet, D., Guillemaud, T. & Raymond, M. Tracking the evolution of

insecticide resistance in the mosquito Culex pipiens. *Nature* **400**, 861–864 (1999).

17. Miyo, T., Akai, S. & Ogumma, Y. Seasonal fluctuation in susceptibility to insecticides

within natural populations of Drosophila melanogaster: empirical observations of fitness

costs of insecticide resistance.

18. Miyo, T., Oguma, Y. & Charlesworth, B. Seasonal fluctuation in susceptibility to

insecticides within natural populations of Drosophila melanogaster. II. Features of genetic

variation in susceptibility to organophosphate insecticides within natural populations of D.

melanogaster. *Genes Genet. Syst.* **81**, 273–285 (2006).

19. Bourguet, D. & Raymond, M. The molecular basis of dominance relationships: the case of

some recent adaptive genes. *J. Evol. Biol.* **11**, 103–122 (1998).

20. Muralidhar, P. & Veller, C. Dominance shifts increase the likelihood of soft selective

sweeps. *Evolution* **76**, 966–984 (2022).

21. Labbé, P. *et al.* GENE-dosage effects on fitness in recent adaptive duplications: ace-1 in the

mosquito Culex pipiens. *Evolution* **68**, 2092–2101 (2014).

22. Menozzi, P., Shi, M. A., Lougarre, A., Tang, Z. H. & Fournier, D. Mutations of

acetylcholinesterase which confer insecticide resistance in Drosophila melanogaster

populations. *BMC Evol. Biol.* **4**, 4 (2004).

23. Karasov, T., Messer, P. W. & Petrov, D. A. Evidence that adaptation in Drosophila is not

limited by mutation at single sites. *PLoS Genet.* **6**, e1000924 (2010).

24. Shi, M. A. *et al.* Acetylcholinesterase alterations reveal the fitness cost of mutations

conferring insecticide resistance. *BMC Evol. Biol.* **4**, 5 (2004).

25. Mackay, T. F. C. *et al.* The Drosophila melanogaster Genetic Reference Panel. *Nature* **482**,

173–178 (2012).

26. Kapun, M. *et al.* Drosophila Evolution over Space and Time (DEST): A new population

genomics resource. *Mol. Biol. Evol.* **38**, 5782–5805 (2021).

27. FAO. Pesticides use and trade, 1990–2022. (2024).

28. Paaby, A. B., Bergland, A. O., Behrman, E. L. & Schmidt, P. S. A highly pleiotropic amino

acid polymorphism in the Drosophila insulin receptor contributes to life-history adaptation.

Evolution **68**, 3395–3409 (2014).

29. Wittmann, M. J., Mousset, S. & Hermisson, J. Modeling the genetic footprint of fluctuating

balancing selection: from the local to the genomic scale. *Genetics* **223**, iyad022 (2023).

30. Kaplan, N., Hudson, R. & Langley, C. The 'hitchhiking effect' revisited. *Genetics* **123**,

887–899 (1989).

31. Langmüller, A. M. *et al.* Fitness effects for Ace insecticide resistance mutations are

determined by ambient temperature. *BMC Biol.* **18**, 157 (2020).

32. Johnston, S. E. *et al.* Life history trade-offs at a single locus maintain sexually selected

genetic variation. *Nature* **502**, 93–95 (2013).

33. Barson, N. J. *et al.* Sex-dependent dominance at a single locus maintains variation in age at

maturity in salmon. *Nature* **528**, 405–408 (2015).

34. Pfenninger, M. & Foucault, Q. Population genomic time series data of a natural population

suggests adaptive tracking of fluctuating environmental changes. *Integr. Comp. Biol.* **62**,

1812–1826 (2022).

35. Pfenninger, M., Foucault, Q., Waldvogel, A.-M. & Feldmeyer, B. Selective effects of a

short transient environmental fluctuation on a natural population. *Mol. Ecol.* **32**, 335–349

(2023).

36. Mutero, A., Pralavorio, M., Bride, J. M. & Fournier, D. Resistance-associated point

mutations in insecticide-insensitive acetylcholinesterase. *Proc. Natl. Acad. Sci. U. S. A.* **91**,

5922–5926 (1994).

37. Harel, M. *et al.* Three-dimensional structures of Drosophila melanogaster

acetylcholinesterase and of its complexes with two potent inhibitors. *Protein Sci.* **9**, 1063–

1072 (2000).

38. Ritz, C., Baty, F., Streibig, J. C. & Gerhard, D. Dose-Response Analysis Using R. *PLoS*

One **10**, e0146021 (2015).

39. Tilk, S. *et al.* Accurate Allele Frequencies from Ultra-low Coverage Pool-Seq Samples in

Evolve-and-Resequence Experiments. *G3* **9**, 4159–4168 (2019).

40. Miyo, T. & Oguma, Y. Negative correlations between resistance to three organophosphate

962  insecticides and productivity within a natural population of Drosophila melanogaster

963  (Diptera: Drosophilidae). *J. Econ. Entomol.* **95**, 1229–1238 (2002).

964 41. Singh, R. S. & Morton, R. A. Selection for malathion-resistance in Drosophila

965  melanogaster. *Can. J. Genet. Cytol.* **23**, 355–369 (1981).

966 42. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

967  transform. *Bioinformatics* **25**, 1754–1760 (2009).

968 43. Kessner, D., Turner, T. L. & Novembre, J. Maximum likelihood estimation of frequencies

969  of known haplotypes from pooled sequence data. *Mol. Biol. Evol.* **30**, 1145–1158 (2013).

970 44. Gillespie, J. H. *Population Genetics*. (Johns Hopkins University Press, Baltimore, MD,

971  2004).

972 45. Behrman, E. L. & Schmidt, P. How predictable is rapid evolution? (2022)

973  doi:10.1101/2022.10.27.514123.

974 46. Ewens, W. J. *Mathematical Population Genetics 1: Theoretical Introduction*. (Springer,

975  New York, NY, 2010).

976 47. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *arXiv [cs.LG]* (2012).

977 48. Wiberg, R. A. W., Gaggiotti, O. E., Morrissey, M. B. & Ritchie, M. G. Identifying

978  consistent allele frequency differences in studies of stratified populations. *Methods Ecol.*

979  *Evol.* **8**, 1899–1909 (2017).

980 49. Feder, A. F., Petrov, D. A. & Bergland, A. O. LDx: estimation of linkage disequilibrium

981  from high-throughput pooled resequencing data. *PLoS One* **7**, e48588 (2012).

982 50. VanLiere, J. M. & Rosenberg, N. A. Mathematical properties of the r2 measure of linkage

983  disequilibrium. *Theor. Popul. Biol.* **74**, 130–137 (2008).