Human gut commensal *Alistipes timonensis* modulates the host lipidome and delivers anti-inflammatory

outer membrane vesicles to suppress colitis in an *Il10*-deficient mouse model

Ethan A. Older[a†], Mary K. Mitchell[b†], Andrew Campbell[a], Xiaoying Lian[a], Michael Madden[a], Yuzhen Wang[c],

Lauren E. van de Wal[b], Thelma Zaw[b], Brandon N. VanderVeen[d], Rodney Tatum[a], E. Angela Murphy[d], Yan-Hua

Chen[a], Daping Fan[c], Melissa Ellermann[b*], and Jie Li[a*]

[a]Department of Chemistry and Biochemistry, University of South Carolina, Columbia, South Carolina 29208,

United States.

[b]Department of Biological Sciences, University of South Carolina, Columbia, South Carolina 29208, United

States.

[c]Department of Cell Biology and Anatomy, School of Medicine, University of South Carolina, Columbia, South

Carolina 29209, United States.

[d]Department of Pathology, Microbiology and Immunology, School of Medicine, University of South Carolina,

Columbia, South Carolina 29209, United States.

[†]These authors contributed equally.

*To whom correspondence may be addressed.

Email: mellermann@sc.edu; li439@mailbox.sc.edu

Abstract

Correlative studies have linked human gut microbes to specific health conditions. *Alistipes* is one such microbial genus negatively linked to inflammatory bowel disease (IBD). However, the protective role of *Alistipes* in IBD is understudied, and the underlying molecular mechanisms remain unknown. In this study, colonization of *Il10*-deficient mice with *Alistipes timonensis* DSM 27924 delays colitis development. Colonization does not significantly alter the gut microbiome composition, but instead shifts the host plasma lipidome, increasing phosphatidic acids while decreasing triglycerides. Outer membrane vesicles (OMVs) derived from *Alistipes* are detected in the plasma of colonized mice, carrying potentially immunomodulatory metabolites into the host circulatory system. Fractions of *A. timonensis* OMVs suppress LPS-induced *Il6*, *Il1b*, and *Tnfa* expression *in vitro* in murine macrophages. We detect putative bioactive lipids in the OMVs, including immunomodulatory sulfonolipids (SoLs) in the active fraction, which are also increased in the blood of colonized mice. Treating *Il10*-deficient mice with purified SoL B, a representative SoL, suppresses colitis development, suggesting its contribution to the anti-inflammatory phenotype observed with *A. timonensis* colonization. Thus, *A. timonensis* OMVs represent a potential mechanism for *Alistipes*-mediated delay of colitis in *Il10*-deficient mice via delivery of immunomodulatory lipids and modulation of the host plasma lipidome.

Keywords:

Introduction

The trillions of microbes inhabiting the human gut are inherently linked to human health through interactions with their host environment. These complex interactions affect human health in various negative and positive ways; for example, *Helicobacter pylori* can lead to the formation of gastric ulcers and later gastric cancer whereas, *Akkermansia muciniphila* is inversely associated with obesity, diabetes, cardiometabolic diseases and inflammation.[1,2] The composition of the gut microbiome is delicately balanced and it dynamically responds to external stimuli. When this balance is significantly disturbed, it can impair crucial functions such as maintaining immune homeostasis and can contribute to the development and exacerbation of various diseases including cancer, cardiovascular disease, and chronic inflammation.[3–5] This disturbance in microbiome composition and resulting functions is known as microbial dysbiosis. Inflammatory bowel disease (IBD) is particularly affected by microbial dysbiosis.[6] While numerous studies have reported major shifts in the gut microbiome composition with IBD pathogenesis, there is a lack of mechanistic understanding regarding how specific bacterial populations might protect against IBD progression.

Another hallmark of IBD is dysregulation of the circulating plasma lipid profile, known as dyslipidemia.[7–9] Normally, circulating lipids in the plasma play a role in maintaining homeostasis in processes throughout the body including energy metabolism, cardiovascular health, and managing inflammation.[10,11] Imbalances in these lipids lead to perturbations in their respective processes and often have far reaching effects.[11] For example, dysregulation of specific lipids such as sphingosine-1-phosphate and triglycerides have been correlated with IBD pathogenesis.[12,13] Gut microbes have also been found to modulate the host lipidomic profile and impact host physiology, further complicating the interplay between the gut microbiome and human health.[8,14]

Alistipes is an emerging microbial genus that is consistently found in the human gut microbiome.[15] As a non-pathogenic commensal, *Alistipes* has shown increasing human health relevance and has been linked to protective effects against numerous inflammatory diseases by correlative gut microbial sequencing studies.[16–19] Importantly, *Alistipes* abundance has been negatively associated with disease severity in IBD patients and its metabolites were shown to be decreased in an *Il10*-deficient mouse model of experimental colitis, supporting that *Alistipes* may play a protective role against IBD through its metabolite production.[20] *Alistipes* colonization in the DSS and oxazolone-induced mouse models of experimental colitis has been found to attenuate inflammation

through the induction of endogenous anti-inflammatory cytokines including IL-10.[21,22] While this interaction has been documented, it remains unclear how *Alistipes* induces this phenotype and to what extent *Alistipes*-derived microbial functional metabolites mediate interactions between *Alistipes* and the host.

It has been proposed that gut bacteria secrete outer membrane vesicles (OMVs) to interact with their hosts. OMVs are spherical nanoparticles derived from the outer membranes of Gram-negative bacteria.[23] Their size ranges between 20 – 300 nm in diameter and they are typically composed of a lipid bilayer containing phospholipids, lipopolysaccharide (LPS), and outer membrane proteins. They are deliberately produced in response to stress signals including temperature, oxidation, and nutrient availability, and are particularly important in mediating interactions among microbes, hosts, and other organisms.[24,25] Notably, OMVs can be selectively packaged by their producers and serve as delivery vehicles for their cargo which can include metabolites, proteins, DNA, and RNA.[26–29] This provides a means to carry microbial components through the protective mucus layer and enable otherwise gut-restricted elements to cross the colonic epithelium and reach host cells via the cardiovascular system, initiating various host responses.[28,30–32]

Here, we show that colonization of *Il10*-deficient mice with *Alistipes timonensis* DSM 27924, a producer of sulfonolipids (SoLs), delays the development of colitis. We show that *Alistipes* colonization does not significantly affect the gut microbiome composition during colitis development, but results in a shift of the host plasma lipidome consistent with clinical observations. We find OMVs derived from *A. timonensis* in the blood plasma of colonized mice which carry potentially immunomodulatory *A. timonensis* metabolites to the host circulatory system. We then show that fractionated OMVs stimulate an anti-inflammatory response *in vitro* in murine macrophages and we detect immunomodulatory SoLs in the active fraction. We finally demonstrate that a purified SoL B fraction isolated from *A. timonensis* OMVs also delays colitis in *Il10*-deficient mice. Taken together, this study reports for the first time that *A. timonensis* delays colitis in the *Il10*-deficient mouse model and proposes *A. timonensis*-derived SoL B, which is delivered to host circulation by OMVs, as a potentially contributing factor to the anti-inflammatory action of *A. timonensis* colonization.

Results

Colonization of *Alistipes timonensis* DSM 27924 in *Il10*-deficient mice delayed colitis progression

To determine whether *Alistipes* impacts intestinal inflammation development, we utilized the well-

established *Il10*-deficient (IL10 KO) mouse model of piroxicam-accelerated colitis.[33–35] This mouse model

captures the genetic and environmental interactions and contributions that drive IBD development.[34] Further,

Alistipes colonization in alternative models of experimental colitis has shown that colonization induces innate IL-

10 production which is a major contributor to anti-inflammatory phenotypes.[21,22] Thus, the use of the IL10 KO

model will allow us to examine the effects of *Alistipes* and its metabolites on colitis progression without influence

from endogenous IL-10 production. *Alistipes timonensis* DSM 27924 was engrafted into the gut microbiomes of

the colitis-susceptible IL10 KO mice at the start of colitis induction via a diet supplemented with piroxicam. *A.*

timonensis DSM 27924 was selected because it was previously shown to colonize the gut of mice.[36] Additionally,

while the genus *Alistipes* has been negatively correlated with inflammatory diseases, such as IBD, this species

of *Alistipes* has not been previously positively or negatively linked to any disease.[16,20]

Two groups of female IL10 KO mice were used: group 1, piroxicam (uncolonized) and group 2, piroxicam

+ *A. timonensis* (*A. timonensis*-colonized). A confirmatory second independent cohort of female IL10 KO mice

was also used, and all results were found to be consistent between the two cohorts. Thus, the data from these

two independent cohorts were pooled for all further analysis. PCR amplification of the *A. timonensis* 16S rRNA

gene confirmed the presence and persistence of *A. timonensis* after oral gavage, supporting successful

colonization and confirming that *A. timonensis* was not present prior to engraftment (**Supplemental Figure S1a**).

Uncolonized mice developed colitis as indicated by decreased body weight over time (**Supplemental Figure**

S1b,c) and increased gross pathology and histology scores at necropsy (**Figure 1a**). In contrast, *A. timonensis*-

colonized mice showed reduced weight loss compared to uncolonized mice as well as significantly decreased

gross pathology and histology scores (**Supplemental Figure S1b,c**) (**Figure 1a**), suggesting that *A. timonensis*

may act to suppress colitis development in the IL10 KO mice. Consistent with model expectations, gross

pathology and histology scoring indicated that suppression of intestinal inflammation was localized to the colon

and was more pronounced in the distal colon (**Supplemental Files 1 and 2**).[33,34] Inflammatory cytokine profiling

of RNA extracted from the distal colon via reverse transcription quantitative PCR (RT-qPCR) revealed that *A.*

timonensis-colonized mice had decreased expression of pro-inflammatory cytokines *Il1b*, *Tnfa*, and *Il6*, as well

as pro-inflammatory marker *Nos2* (**Figure 1b-e**), further supporting that *A. timonensis* colonization suppressed

intestinal inflammation.

Colitis in this model is driven by enhanced colonic epithelial apoptosis leading to a loss of gut barrier

integrity.[34] Thus, we examined cellular tight junctions using immunofluorescence microscopy and found that tight

junction morphology was restored in *A. timonensis*-colonized colon samples (**Figure 2a**). We also measured the

expression of tight junction proteins, revealing that *A. timonensis*-colonized mice had significantly increased

expression of *claudin-3* concurrent with decreased expression of *claudin-7*. (**Figure 2b**). Claudin-3 has been

previously reported to decrease in colitis, reflecting a loss of gut barrier integrity and leakage of gut contents into

the host circulation.[37] Claudin-7 has been linked to intestinal epithelial stem cell renewal and regulation and its

decreased expression in *A. timonensis*-colonized mice here may suggest a reduced need for cellular repair in

the epithelium due to reduced colitis after *A. timonensis* colonization.[38]

Figure 1. *A. timonensis* colonization delays piroxicam-accelerated colitis development in IL10 KO mice.

(a) Representative H&E images of uncolonized (Uncol.) and *A. timonensis*-colonized (*A. tim.*) mouse colons

showing less inflammation in colonized mice. One-sided Student's *t*-test confirmed significantly decreased gross

pathology and histology scores in Uncol. (n = 9 mice) compared to *A. tim.* (n = 9 mice) mice. (b-e) The expression

of pro-inflammatory factors *Il1b*, *Tnfa*, *Il6*, *and Nos2* were significantly decreased in the distal colons of *A.*

timonensis-colonized (*A. tim., n* = 9 mice) mice compared to uncolonized (Uncol*., n* = 9 mice) mice, supporting

that *Alistipes* colonization mediated a suppressive effect against colitis progression. Significance was determined

using two-sided Student's *t*-test. For all *p*-values, * 0.05 > *p* > 0.0 and **** 0.0001 > *p*.

Figure 2. *A. timonensis* colonization modulates tight junction protein morphology and gene expression

in IL10 KO mouse colons. (a) Immunofluorescent staining of tight junction proteins claudin-3 and claudin-7

(red) illustrating restoration of tight junction morphology and reduction of tight junction repair with *A. timonensis*

colonization, respectively. Nuclei were stained with DAPI (blue). A: Apical side of colon tissue; Arrow: tight

143   junction morphologies. (b) Gene expression of *claudin-3* and *claudin-7* in uncolonized (Uncol., n = 9 mice) and

144   *A. timonensis*-colonized (*A. tim.*, n = 9 mice) mice, which is consistent with immunofluorescence staining.

145   Significance was determined using two-sided Student's *t*-test. For all *p*-values, * $0.05 > p > 0.01$.

147   **The anti-inflammatory effects of *A. timonensis* in IL10 KO mice are independent from reversal of**

148   **microbial dysbiosis**

149   Gut dysbiosis is a hallmark of IBD, with prolonged dysbiosis leading to reduced intestinal homeostasis

and ultimately chronic inflammation.[6,39] In the IL10 KO mouse model, dysbiosis further exacerbates inflammation

due to increased leakage of gut contents through the damaged colonic epithelium.[34] Considering this, we first

hypothesized that *A. timonensis* colonization may mediate an anti-colitic effect by preventing or reversing

perturbations in gut microbiota composition, resulting in protection against IBD-related gut dysbiosis. Thus, we

performed 16S rRNA sequencing to profile the gut microbiome composition in *A. timonensis*-colonized (*A.*

timonensis, n = 10 mice) and uncolonized (Uncolonized, *n* = 11 mice) IL10 KO mice - both groups of which were

treated with piroxicam. An age-matched control group of IL10 KO mice (Control, *n* = 17 mice) that were neither

treated with piroxicam nor colonized with *A. timonensis* was also included (**Supplemental File 3**). Comparing

uncolonized mice to control mice, microbiome composition analysis revealed a clear shift in the microbial profile

(**Figure 3a**) accompanied by a significant decrease in alpha diversity (**Figure 3b**). This loss in microbial diversity

is indicative of microbial dysbiosis in colitis development, consistent with the literature.[40,41]

161   However, despite the attenuated inflammation observed with *A. timonensis* engraftment, the microbiota

profile of *A. timonensis*-colonized mice showed no clear distinction from that of uncolonized mice and had no

significant change in alpha diversity compared to uncolonized mice (**Figure 3a,b**). To verify this observation, we

analyzed the second cohort independently and observed a consistent lack of significant microbiome

compositional changes between *A. timonensis*-colonized and uncolonized mice (**Figure 3a**). At the family level,

we observed clear differences between Control samples and both uncolonized and *A. timonensis*-colonized

samples but observed only minor changes between uncolonized and *A. timonensis*-colonized samples, with one

prominent family that *A. timonensis* belongs to, Rikenellaceae, notably more abundant in *A. timonensis*-colonized

samples (**Figure 3c**). Since *Alistipes* is a member of the phylum Bacteroidetes, we also looked at changes in

genera within Bacteroidetes; we observed that an increase in the abundance of *Alistipes* in *A. timonensis*-colonized compared to uncolonized samples was the only major difference (**Figure 3d**). This confirmed that the only major change in the gut microbiome profile upon *A. timonensis* colonization was an expected increase in *Alistipes*.

In contrast with many other gut microbes such as *Bacteroides uniformis* that has been shown to reduce colitis through significant modulation of the gut microbiome composition when colonized, colonization with *A. timonensis* appears to mediate its anti-colitic effect independently of major microbiome remodeling.[42] This unexpected yet intriguing result suggests that *A. timonensis* itself plays a more direct role in the observed anti-colitic effect and focused the study towards exploring more direct host-bacterial mechanisms by which this strain modulates host inflammation.

Figure 3. Microbiome composition analysis indicates *A. timonensis* colonization does not reverse dysbiosis. (a) Beta diversity measured by principal coordinate analysis (PCoA) using Jaccard distances shows clear separation between both independent cohorts (red and green triangles and circles) and the control group (blue squares) but no separation between *A. timonensis*-colonized (red) versus uncolonized (green) microbiome profiles. (b) Alpha diversity measured by Shannon diversity index indicates a significant loss of microbial diversity from control ($n = 17$ mice) to uncolonized ($n = 11$ mice) and *A.timonensis*-colonized ($n = 10$ mice) mice that was not restored by *A. timonensis* colonization. Significance was determined using two-sided Student's *t*-test. For all *p*-values, * $0.05 > p > 0.01$, ** $0.01 > p > 0.001$, and ns: not significant. (c) Heatmap of microbiome composition at the family level showing a clear difference in control samples compared to uncolonized and *A. timonensis*-colonized mice. (d) Genera composition of the phylum Bacteroidetes highlights the major expansion of *A. timonensis* in colonized compared to uncolonized samples.

A. timonensis colonization modulates host lipidomic profile associated with colitis

Considering that the observed anti-colitic phenotype was reflected in host cytokine and host tight junction protein and gene expression, we hypothesized that *A. timonensis* colonization may directly induce a change in host signaling pathways. Notably, host immune signaling pathways are tightly regulated by circulating blood immunomodulatory lipids.[8,43] The levels of these lipids are directly affected by the gut microbiome in various ways, such as through direct microbial biotransformation of primary bile acids to secondary bile acids or through bacterial sphingolipid-dependent downregulation of endogenous sphingolipid biosynthesis as well as the production and circulation of other microbial metabolites.[44,14] To examine the effects of *A. timonensis* colonization on circulating host lipids, lipidomic analysis of the mouse blood plasma was conducted (**Supplemental File 4**). Using orthogonal partial least squares discriminant analysis, we observed that the host lipid profile was significantly different between *A. timonensis*-colonized and uncolonized mice (**Figure 4a**). Among the significantly changed lipids, phospholipids were found to be more abundant in *A. timonensis*-colonized compared to uncolonized samples, while triglycerides were more abundant in uncolonized compared to *A. timonensis*-colonized samples. (**Figure 4b**). These results were consistent with clinical observations that triglycerides are increased in both ulcerative colitis and Crohn's disease.[13]

A recent study in acutely stimulated immune cells demonstrated that arachidonic acid (AA) is mobilized

first from membrane phospholipids, which is then replenished by triacylglycerol AA after inflammation has

subsided.[45] We observed a similar trend in that AA-containing phosphatidic acids were accumulated in *A.*

timonensis-colonized compared to uncolonized samples while triglycerides were decreased in *A. timonensis*-

colonized samples (**Figure 4c,d**) (**Supplemental Table 1**). This may represent the replenishment of AA from

triglycerides to phospholipids in association with an *A. timonensis*-mediated suppression of the inflammatory

response following piroxicam challenge. These results indicate that *A. timonensis* colonization indeed modulates

host signaling pathways by affecting immunomodulatory lipid regulation and suggests that a factor derived from

A. timonensis may be involved in mediating this effect. To explore this potential factor from a molecular point of

view, we considered the production and translocation of metabolites by *A. timonensis*.

Figure 4. Plasma lipidomics reveals colonization-driven change in circulating lipid profile. (a) Orthogonal

partial least squares-discriminant analysis (OPLS-DA) shows clear difference in plasma lipid profiles between

uncolonized (n = 9 mice) and *A. timonensis*-colonized (n = 9 mice) mice. (b) The top 20 significantly differentially

abundant lipids reveal a marked depletion of triglycerides (TGs) and accumulation of phosphatidic acids (PAs)

in *A. timonensis*-colonized mice compared to uncolonized mice. (c,d) *A. timonensis* colonization affected the

balance of two pools of arachidonic acid-containing lipids: triglycerides (c), which were less abundant in *A.*

timonensis-colonized mice, and phospholipids (d), which were more abundant in *A. timonensis*-colonized mice.

Significance was determined using two-sided Student's *t*-test. For all *p*-values, * 0.05 > *p* > 0.01 and ** 0.01 > *p*

> 0.001.

***A. timonensis* secreted outer membrane vesicles (OMVs) bearing biologically active lipids into the**

plasma after colonization

OMVs are ubiquitously produced by Gram-negative bacteria and are known to carry bacterially-derived

elements away from the producing cell.[27,28,31] These bacterial delivery vehicles represent a potential mechanism

for metabolite translocation to the host circulation. Thus, we collected the plasma of uncolonized and *A.*

timonensis-colonized mice and purified OMVs using density gradient ultracentrifugation (**Supplemental Figure**

S2). To ensure detection, aliquots of plasma from each mouse were pooled together into representative samples.

Indeed, OMV-like nanoparticles were purified from the pooled plasma and visualized by transmission electron

microscopy (TEM) (**Figure 5a**) (**Supplemental Figure S3**), suggesting the presence of OMVs in the samples.

To determine if these OMVs were related to *A. timonensis* colonization, we used dynamic light scattering (DLS)

to measure the size distributions of OMVs isolated from the plasma of uncolonized and *A. timonensis*-colonized

mice in comparison to OMVs isolated from a monoculture of *A. timonensis*. DLS showed that OMVs from plasma

are distributed into two distinct populations (10-55 nm and 55-330 nm) and that OMVs from *A. timonensis*-

colonzied mouse plasma showed an increased abundance of particles corresponding to the diameter range

matching that of OMVs from *A. timonensis* culture (**Figure 5b**). This data indicates that *A. timonensis* colonization

resulted in more OMVs in the plasma which are potentially derived from *A. timonensis.* To confirm that these

OMVs came from *A. timonensis*, we conducted targeted metabolomic analysis looking for biomarkers of *A.*

timonensis. We and others have previously shown that sulfonolipids (SoLs) are a class of molecules that are

uniquely produced by the gut microbes *Alistipes* and *Odoribacter*.[35,36,46] SoLs are further known to localize to the

outer membrane of producing bacteria,[47] strongly suggesting they may be found in OMVs. Indeed, our

metabolomic analysis revealed a significantly higher abundance of sulfobacin B (SoL B), the major SoL produced

by *Alistipes*,[35] in *A. timonensis*-colonized plasma OMV samples compared to uncolonized plasma OMV samples

(**Figure 5c**). Considering our DLS results, this metabolomic evidence, and our microbiota composition analysis which showed that increased abundance of SoL-producing *A. timonensis* was the only major change after colonization, the OMVs we isolated from the plasma of *A. timonensis*-colonized mice likely contain an abundance of *A. timonensis*-derived OMVs.

Microbial metabolites tend to be restricted within the gut if they are not actively delivered to host circulation.[31,48] Since the OMVs were found to carry SoL B, we considered if *A. timonensis*-derived lipids, including SoLB, could be delivered to the blood of mice after *A. timonensis* colonization. Accordingly, targeted metabolomic analysis of the host plasma lipidome showed a significant increase of SoL B in the plasma and also indicated a significant increase in the abundance of sulfobacins A (SoL A) and F (SoL F) in *A. timonensis*-colonized compared to uncolonized plasma (**Figure 6a-d**). The identity of SoLs was confirmed in these samples by retention time and MS/MS fragmentation matching compared to in-house standards and literature references (**Figure 6e,f**).[49] This result suggested that *A. timonensis*-derived OMVs indeed delivered *A. timonensis*-derived metabolites including SoLs to the host circulatory system. The delivery of these non-gut restricted lipids thus represents a potential method by which *A. timonensis* colonization suppresses the colitis phenotype. To support this, we considered if the increased abundance of SoLs in the plasma was due to passive diffusion or if *A. timonensis* OMVs containing SoLs were actively translocated to the host circulation. Since the SoL-producer *Alistipes* colonizes the gut of the mice, we assumed that the production of SoL B is primarily in the gut and thus the concentration of SoL B should be greatest in the feces. If SoL B passively diffuses into host circulation, then we would expect a higher abundance of SoL B in the feces compared to the plasma. If SoL B is actively translocated to host circulation, then we would expect the abundance of SoL B to be greater in the plasma than the feces. We compared the fold changes in SoL B abundance between fecal material and plasma in both uncolonized and *A. timonensis*-colonized samples. We found that SoL B increased by 1.8-fold in uncolonized mice while it increased by 3.9-fold in *A. timonensis*-colonzied mice (**Figure 6g**). In both cases, SoL B was greater in the plasma than in the feces, suggesting that SoL B is actively translocated from the gut into host circulation. Taken together with the presence of SoL B in *A. timonensis* OMVs and increased abundance of these OMVs in *A. timonensis*-colonized mice, it is likely that OMVs serve as a vehicle for active delivery of *A. timonensis*-derived SoL B to host circulation.

Figure 5. OMVs purified from the plasma of *A. timonensis*-colonized mice are abundant in *A. timonensis*-derived OMVs. (a) TEM was used to visually confirm the presence of OMVs after negative staining. x30.0k magnification was used to confirm the diameter of OMVs in the expected range. (b) DLS was used to measure the size distribution of plasma OMVs from uncolonized (green) and *A. timonensis*-colonized (red) mice alongside OMVs purified from *A. timonensis* culture (black). DLS analysis showed two clear vesicle populations with diameter ranges of 10-55 nm and 55-320 nm. The rightmost larger diameter population corresponds with *A. timonensis* culture OMVs and reflects the increased abundance of OMVs derived from *A. timonensis* colonization. (c) Lipids were extracted from purified blood OMVs, measured using targeted metabolomics, and SoL B was found to be significantly increased in plasma OMVs from *A. timonensis*-colonized ($n = 4$ mice) compared to uncolonized ($n = 4$ mice) mice. Significance was determined using two-sided Student's *t*-test. For all *p*-values, ** $0.01 > p > 0.001$.

Figure 6. Sulfonolipid abundance significantly increased in the plasma of *A. timonensis*-colonized mice.

(a-c) Abundance of SoL B (a), SoL A (b), and SoL F (c) measured in uncolonized (n = 10 mice) compared to *A. timonensis*-colonized (n = 10 mice) mouse plasma by targeted metabolomics. (d) Corresponding chemical structures of SoL B, A, and F. (e) Retention time comparison of in-house SoL B standard and observed SoL B peak in *A. tim.* samples supporting identification as SoL B. (f) MS/MS fragmentation used to positively identify SoL B based on characteristic fragmentation including the ~81 m/z fragment corresponding to the loss of the HSO_3^- headgroup. (g) The plasma-to-feces ratio of SoL B abundance was significantly greater in *A. timonensis*-colonized (n = 7) compared to uncolonized (n = 7) samples, suggesting that the increased production of SoL B

by *A. timonensis* in the gut led to increased SoL B abundance in the plasma via active translocation of SoL B

from the gut to the plasma by OMVs. Significance was determined using two-sided Student's *t*-test. For all *p*-

values, * 0.05 > *p* > 0.01, ** 0.01 > *p* > 0.001, *** 0.001 > *p* > 0.0001, and **** 0.0001 > *p*.

A. timonensis OMV fraction containing bioactive lipids exerts an anti-inflammatory effect *in vitro* in

murine macrophages

Because *Alistipes*-derived OMVs were found in the host circulation, we considered if these OMVs

contributed to the observed anti-colitic phenotype. To obtain enough OMVs for an *in vitro* assay, we purified

OMVs from a culture of *A. timonensis*. Due to the complex chemical composition of OMVs, we considered if one

or more of their components possessed anti-inflammatory activity. We thus fractionated the OMV samples based

on polarity and tested each fraction's ability to suppress LPS-induced inflammation in macrophages. We

observed that one fraction, Fraction 2 (F2), significantly suppressed LPS-induced inflammation as measured by

fold change of expression of the cytokines *Il1b* and *Il6*, while *Tnfa* showed a similar trend (**Figure 7a**).

Importantly, the leakage of gut antigens, such as LPS, into the blood, leads to immune system activation, which

is one of the driving factors of colitis in the IL10 KO mouse model.[34] Thus, the suppression of LPS-induced pro-

inflammatory cytokine expression by F2 suggested that one or more components in *A. timonensis* OMVs may

modulate the immune system by partially blocking the LPS-mediated inflammatory response.

To investigate these potential biologically active components, the molecular composition of F2 was

analyzed by untargeted metabolomics using high resolution mass spectrometry (HRMS/MS) followed by

molecular networking using the widely used Global Natural Product Social (GNPS) molecular networking tool.[50]

A bioactivity-based molecular network was then constructed by correlating the abundance change of molecular

features with the expression change of inflammatory marker *Il1b* using spearman correlation.[51] The correlative

bioactivity score was mapped to the node color parameter while the abundance was mapped to the node size.

Individual nodes represent molecular features in F2 and edges between nodes indicate similarity in the MS/MS

fragmentation pattern of the nodes (**Figure 7b**). Interrogating the network, we identified several clusters with

both promising bioactivity score and high chemoinformatic confidence scores, supporting the presence of

bioactive components in F2. The cluster containing the largest node in F2 corresponded to SoLs, and the largest

node corresponded to SoL B, which was confirmed by MS/MS comparison to our in-house standard

(**Supplemental Figure S4**). Notably, in our previous study, pure SoL B was found to suppress LPS-induced

inflammation by binding to and displacing LPS from the canonical binding pocket of the TLR4/MD-2 complex,

supporting the observed anti-inflammatory activity of F2.[35]

To identify other potential contributors from F2, we further analyzed the HRMS/MS data using the

chemoinformatic platform SIRIUS, which afforded computationally predicted chemical formulas based on exact

mass measurements as well as predicted compound classes and putative structural identities based on MS/MS

fragmentation patterns.[52–55] In addition to confirming the known biologically active SoL cluster (**Supplemental**

Figure S5), we prioritized five other clusters, clusters 2 through 6, representing families of potentially bioactive

compounds and predicted their compound classes, including glycerophosphoethanolamines, triacylglycerols, *N*-

acyl amines, glycerophosphoserines, and another family of N-acyl amines, respectively. Based on these classes,

we further identified assigned structural features for clusters 2 through 4 (**Figure 7b,c**), then selected

representative nodes for putative structural identifications based on their MS/MS fingerprint matching to known

and predicted database spectra (**Supplemental Tables S2-S5**) (**Supplemental Figures S6-S8**).[53–55] This result

demonstrates the diverse chemical composition, especially lipid molecules, of *A. timonensis* OMVs. Furthermore,

our observation that SoL B was a major compound in the bioactive molecular network with good bioactivity score

suggests that other compounds with similar bioactivity scores may also contribute to the anti-colitic effect of *A.*

timonensis colonization.

Figure 7. Bioactive molecular networking of *A. timonensis* OMV fractions suggests multiple compound classes contributing to an overall anti-inflammatory effect against LPS-induced inflammation. (a) Relative fold change expression pro-inflammatory cytokines *Il1b, Il6,* and *Tnfa* after treating mouse peritoneal macrophages 1 µg/mL LPS, or 1 µg/mL LPS with fractions of *A. timonensis* OMVs at 20 µg/mL (*n* = 3 wells per

treatment). Fraction 2 (F2) showed the strongest suppression among other fractions and was further investigated

for its bioactive components. Significance was determined using two-sided Student's *t*-test. For all *p*-values, *

0.05 > *p* > 0.01, *** 0.001 > *p* > 0.0001, and ns: not significant. (b) Untargeted metabolomics and subsequent

bioactive molecular networking analysis of Fraction 2 of *A. timonensis* OMVs revealed multiple clusters of

molecules including SoLs which were abundant in this active fraction. (c) Predicted chemical structures of the

major clusters, suggesting other potential contributors to the biological activity of Fraction 2.

Sulfonolipids isolated from *A. timonensis* OMVs delayed colitis progression in *Il10*-deficient mice.

Our bioinformatic analysis suggesting SoLs, particularly SoL B, as a major contributing class to the

observed anti-inflammatory activity aligns with the existing literature reporting that SoL B is an

immunomodulatory lipid with anti-inflammatory effects on cells *in vitro* and in some *in vivo* applications.[35,56] Thus,

we were encouraged to determine if SoL B delivered by *A. timonensis*-derived OMVs plays a role in delaying

colitis in the IL10 KO model. From the bioactive *A. timonensis* OMV F2, we further purified SoL B (**Supplemental**

Fig. S9a) and determined this fraction to be free of LPS contamination using an LAL endotoxin assay

(**Supplemental Fig. S9b**). We used this purified SoL B to treat IL10 KO mice (SoL B; n = 6) at a dosage of 1

mg/kg by intraperitoneal (I.P.) injection. As a vehicle control, IL10 KO mice (Veh.; n = 5) were treated with 1X

PBS alongside the SoL B group. Concurrently with the first I.P. injection, both groups were switched to a

piroxicam-containing diet to induce colitis. Mouse weight was measured over the course of 17 days, which

revealed that mice in the SoL B group showed lower weight change at necropsy compared to mice in the Vehicle

group (**Supplemental Fig. S10a**) and showed significantly lower inflammation in gross pathological parameters

with a similarly trending but non-significant pattern in histological parameters (**Figure 8a**) (**Supplemental File**

5). In the distal colon, SoL B treatment further significantly suppressed the expression of pro-inflammatory

markers *Tnfa* and *Lcn2*, while other cytokines *Il6* and *Nos2* showed lowered but non-significant outcomes

(**Figure 8b-e**). In addition, treatment showed a reduced but non-significant expression of *Ifng*, *Il17a*, and *Il12b*,

suggesting SoL B attenuated Th1 and Th17 adaptive immune responses in the host (**Supplemental Fig. S10b**).

Taken together, these data indicate that SoL B treatment by I.P. injection at a dosage of 1 mg/kg can suppress

colitis development in IL10 KO mice, supporting that SoL B likely plays a role in the anti-inflammatory phenotype

observed with *A. timonensis* colonization.

Figure 8. IL10 KO mice treated with purified SoL B by I.P. injection exhibit delayed colitis progression.

(a) Representative histology images illustrate the difference in colon pathology associated with purified SoL B

treatment (SoL B) compared to vehicle control (Vehicle). Gross pathological analysis revealed significantly lower

levels of inflammation in SoL B-treated mice. Histological analysis also supports lower inflammation in the SoL

B-treated mice. (b-e) RT-qPCR showed expression of pro-inflammatory markers were decreased in SoL B-

treated mice (SoL B) compared to vehicle-treated mice (Veh.), supporting that SoL B treatment suppresses

colitis development.

Discussion

In this work, we demonstrate that colonization of *A. timonensis* DSM 27924 in an *Il10*-deficient (IL10 KO)

mouse model of IBD suppresses colitis development which is related to host lipidome modulation rather than

through counteracting microbial dysbiosis. We found that OMVs derived from the plasma of *A. timonensis*-

colonized mice carry immunomodulatory SoLs leading to significantly increased SoL abundance in the plasma

of *A. timonensis*-colonized mice. We then show that fractions of *A. timonensis* OMVs suppress LPS-induced

inflammation and find immunomodulatory SoLs in the active fraction. We further leverage analytical tools

integrating bioactivity and metabolomics to probe the chemical composition of the active *A. timonensis* OMV

fraction, potentially expanding the biologically active metabolites of *A. timonensis* OMVs. Overall, we

demonstrate that *A. timonensis* colonization in IL10 KO mice suppresses colitis progression via OMV-mediated

delivery of immunomodulatory *A. timonensis* metabolites including SoLs to mouse plasma, potentially resulting

in host lipidomic modulation.

Alistipes is a genus with emerging clinical relevance and there are conflicting reports about the impact of

various *Alistipes* species on different diseases, particularly those relating to inflammation such as IBD.[16] Several

studies have negatively correlated *Alistipes* abundance in the gut microbiome to IBD, all suggesting that *Alistipes*

may play a role in protecting against IBD.[20,35] As such, we sought to explore the molecular mechanisms by which

Alistipes affects the host in IBD. We used IL10 KO mice as a model of IBD since these mice are genetically

predisposed to developing intestinal inflammation and disease onset in these mice can be synchronized by

administration of the non-steroidal anti-inflammatory drug (NSAID) piroxicam.[33,34] Using this colitis model, we

observed that colonization with *A. timonensis* DSM 27924 by oral gavage resulted in suppressed colitis

development through significantly reduced gross pathology and histology scores, as well as significantly reduced

inflammatory cytokine expression and favorable modulation of tight junction protein expression. These results

indicate that *A. timonesis* colonization attenuates colitis induced by chronic NSAID feeding.

Typically, this type of interaction is accompanied by major pathological restructuring of the gut

microbiome, which can further exacerbate the host pro-inflammatory immune response.[57,58] However, our

microbial composition analysis by 16S rRNA sequencing revealed that *A. timonensis* colonization had little effect

on the overall gut microbiome composition except for an increase in the abundance of *A. timonensis* itself. This

result was unusual and intriguing, indicating that instead of inducing a compensatory microbial shift, *A.*

timonensis alone directly influences the host to exert an anti-colitic effect, possibly through the modulation of

host signaling pathways. Similarly, the human gut commensal *Bacteroides thetaiotaomicron* is known to produce

a lipid molecule, α-galactosylceramide, that induces a host anti-cancer immune response through natural killer

T cell activation.[59]

To investigate the influence of *Alistipes* on host immune signaling molecules, targeted lipidomic analysis

of host circulating lipids revealed a significantly different lipid profile upon *A. timonensis* colonization

characterized by increased phosphatidic acids and decreased triglycerides. The majority of differential lipids

belonged to the phosphatidic acid and triglyceride classes. Notably, in acutely stimulated immune cells,

arachidonic acid (AA) is mobilized first from membrane phospholipids, including phosphatidic acids, which is

then replenished by triacylglycerol AA after inflammation has subsided.[45] We observed a consistent trend in that

AA-containing phosphatidic acids were accumulated in *A. timonensis*-colonized compared to uncolonized

samples while triglycerides were decreased in *A. timonensis*-colonized samples. This may represent the

replenishment of AA from triglycerides to phospholipids in an *A. timonensis*-mediated suppression of the

inflammatory response in response to piroxicam challenge. Taken together, the clear shift in host lipidome profile

alongside the changes in balance between AA-containing phospholipids and triglycerides support that *A.*

timonensis-colonization indeed affects host immune signaling.

Since *A. timonensis* colonization did not induce a microbiome shift but did affect the profile of host

circulating lipids, we considered if *A. timonensis* was producing and translocating immunomodulatory factors to

the host circulation. Like other Gram-negative bacteria that produce OMVs as a mechanism to interact with their

environments, *A. timonensis* also produces OMVs. OMVs are known to cross the mucosal barrier and epithelial

cell layer, effectively delivering microbial factors to the host circulation.[28,31,32] We found that OMVs isolated and

purified from the blood of *A. timonensis*-colonized mice were enriched in SoL B, the major sulfonolipid of *A.*

timonensis and a component of its outer membranes, suggesting that the purified OMVs were derived from *A.*

timonensis and indicating that *A. timonensis* indeed delivers factors to the host circulation via OMVs.[47] We noted

that the fold-change increase of SoL B in the whole plasma of *A. timonensis*-colonized compared to uncolonized

samples was significantly larger than that of the feces, indicating that SoL B is a non-gut restricted microbial

metabolite and suggesting that *A. timonensis*-derived SoL B may be actively translocated from the gut to the

host circulation by OMVs. Further HRMS/MS analysis of the whole plasma also revealed the increased

abundance of SoLs A and F in *A. timonensis*-colonized blood samples, indicating that *A. timonensis* OMVs

deliver other factors in addition to SoL B to the host circulation. Importantly, SoL A and B were previously

identified to exert anti-inflammatory activity against LPS-induced inflammation, supporting that *A. timonensis* may mediate its anti-colitic effect through delivery of functional metabolites to the host circulation via OMVs.[35]

To determine if *A. timonensis* OMVs could recapitulate the anti-colitic phenotype of *A. timonensis* colonization, we fractionated *A. timonensis* OMVs into different fractions by normal phase chromatography, tested their biological activity, and pinpointed the component responsible for the activity. In an *in vitro* assay measuring LPS-induced pro-inflammatory cytokine expression, we found that Fraction 2 significantly suppressed the expression of *Il1b* and *Il6*. Because the inflammatory response in IL10 KO mice is characterized by leakage of gut contents, such as LPS, into the circulation, we hypothesized that Fraction 2's ability to suppress LPS-induced inflammation in macrophages represents a mechanism by which *A. timonensis* OMVs suppress colitis after *A. timonensis* colonization. To identify bioactive components in the active Fraction 2, we conducted untargeted metabolomics and bioactive molecular networking, which revealed an abundance of SoL B that was negatively correlated with the expression of *Il1b*, supporting its hypothesized involvement in the observed anti-colitic phenotype. While our previous work supports the role of SoL B in the observed phenotype, our understanding of the complexity and intricacy of host-microbial interactions suggested that additional *A. timonensis*-derived components may also play a role.[35] Thus, from the bioactivity-based molecular network, we prioritized five additional clusters with good bioactivity scores and numerous members for chemoinformatic analysis to predict compound classes and potential structure identities using SIRIUS, a machine learning-based chemoinformatics platform.[52–55] Our analysis revealed several other lipid-type compound classes and matched spectral features to several compounds, many of which had similar or better bioactivity score than SoL B. This result significantly expands the pool of potentially bioactive metabolites delivered to host blood by *A. timonensis* OMVs and opens the door for further exploration of *A. timonensis*' effect on inflammation. The presence of potentially bioactive metabolites in *A. timonensis* OMVs suggests that the intact OMVs may also exhibit biological activity. While this activity could be relevant to the effect of *A. timonensis* colonization in IL10 KO mice, OMVs are highly complex in nature. Thus, we focused our study on the identification of bioactive components and fractions of OMVs in the present study, which serves as a foundational step toward exploring the role of intact OMVs in *A. timonensis* colonization in subsequent investigations.

Finally, to confirm the role of SoL B in the observed anti-inflammatory phenotype with *A. timonensis*

colonization, we purified SoL B from the active *A. timonensis* OMV fraction and tested its efficacy in an

independent cohort of IL10 KO mice. We found that purified SoL B does suppress colitis in the IL10 KO model

compared to vehicle-treated control mice. Compared to *A. timonensis* colonization, mice treated with SoL B saw

an almost equivalent level of reduction in gross pathological parameters. This report is the first to demonstrate

the ability of SoL B to suppress colitis in IL10 KO mice and supports that SoL B is a major contributor in mediating

the anti-inflammatory response observed with *A. timonensis* colonization. Further, the presence of SoL B in *A.*

timonensis OMVs coupled with its effectiveness *in vivo* strongly supports our hypothesis that *A. timonensis*

modulates the inflammatory response in IL10 KO mice through delivery of immunomodulatory lipids by OMVs.

By establishing OMVs as a delivery vehicle for immunomodulatory SoLs and other potentially bioactive

lipids, we show that *A. timonensis* can translocate functional metabolites from the gut into the plasma and affect

the balance of circulating host immune signaling molecules. This provides a potential explanation for *Alistipes'*

influences on distant organs such as the liver and cardiovascular system.[17,19] For example, circulating *Alistipes*

OMVs in the blood may directly deliver active metabolites to the liver or cardiovascular tissue inducing a long-

range protective effect against liver and cardiovascular inflammation. While this study is focused on *Alistipes*,

we envision that investigating the OMVs of other biologically relevant human gut microbiota may serve as a vital

strategy to reveal additional biologically active non-gut restricted microbial metabolites in host-microbe

interactions.

Materials and Methods

Bacterial strain cultivation. *Alistipes timonensis* DSM 27924 was obtained from the German Collection of

Microoogranisms and Cell Cultures (DSMZ). For the production of OMVs, we used *Alistipes timonensis* DSM

27924. *A. timonensis* was cultivated under anaerobic conditions on solid brain heart infusion agar plates or in

liquid reinforced clostridium media at 37°C.

Animals. *Il10*-deficient (IL10 KO) mice on the C57BL/6 background were originally purchased from The Jackson

Laboratory. All mice were bred and maintained in specific pathogen free conditions at the University of South

Carolina and were maintained on a 12-hour light/dark cycle with unlimited access to water and food (Inotiv

Teklad, 8604). All animal protocols were approved by the University of South Carolina Institutional Animal Care

and Use Committee.

***Il10*-deficient mouse model and treatments.** At 7.5-9 weeks of age, female IL-10 KO mice were divided into

one of the following experimental groups: 1) piroxicam treated (n = 10 mice distributed into 3 cages); 2) piroxicam

treated and colonized with *A. timonensis* (n = 11 mice distributed into 3 cages). Female control IL10 KO mice (n

= 17 mice distributed into 6 cages) at 7.5-9 weeks old were caged separately from piroxicam-treated IL10 KO

mice. Colitis development was initiated as previously described.[35] Briefly, on day 0, all mice were switched to a

diet supplemented with 100 ppm of piroxicam (Cayman Chemical, 13368; Inotiv Teklad, TD.210442). From day

0, mice in the first group were gavaged with RCM (reinforced clostridium media) to serve as a vehicle control,

while *A. timonensis*-colonized mice were gavaged with 7.48x10^7 CFU/mL of *A. timonensis* in RCM. All mice were

gavaged three times per week for 2.5 weeks until necropsy at 17 days post piroxicam initiation.

For purified SoL B treatment, female IL10 KO mice at 8.5-10 weeks of age were divided into two groups: vehicle

control (n = 5 mice distributed into 3 cages – originally 3 cages of 6 females, but 1 mouse died due to reasons

unrelated to the model) and SoL B treated (n = 6 mice distributed into 3 cages). SoL B was prepared at a

concentration of 0.2 mg/mL in 1X PBS with <1% DMSO and 100 µL was delivered to mice at a dosage of 1

mg/kg by I.P. injection. Control mice were injected with 100 µL of 1X PBS vehicle. On day 0, all mice were

switched to a piroxicam-supplemented diet as above prior to the first I.P. injection. All mice were injected three

times per week for 2.5 weeks until necropsy at 17 days post piroxicam induction.

Mouse body weight was tracked as an indicator for piroxicam-accelerated colitis progression. At necropsy,

intestinal tissues were collected for evaluating colitis severity, intestinal contents were collected for microbiome

and metabolite analysis, and plasma was collected for lipidomics and OMV detection. Colitis severity including

gross pathology analysis, histopathology analysis, and proinflammatory cytokine gene expression were

performed as previously described.[35] The relative abundance of mammalian mRNA transcripts was calculated

using the $\Delta\Delta C_T$ method and normalized to *Eef2* levels. The oligonucleotides used for qRT-PCR were: *Eef2*

forward: TGTCAGTCATCGCCCATGTG, reverse: CATCCTTGCGAGTGTCAGTGA; *Tnfa* forward:

AGCCAGGAGGGAGAACAGAAAC, reverse: CCAGTGAGTGAAAGGGACAGAACC; *Nos2* forward

TTGGGTCTTGTTCACTCCACGG, reverse: CCTCTTTCAGGTCACTTTGGTAGG; *Il6* forward:

GAAATGATGGATGCTACCAAACTG, reverse: CTCTCTGAAGGACTCTGGCTTTG; *Il1b* forward:

CTCAATGGACAGAATATCAACCAAC, reverse: GGCTGTGCCGTCTTTCATTAC.

Immunofluorescence light microscopy. Fresh frozen colon tissues were cross-sectioned at 10 μm and fixed

with 100% Acetone for 10 mins at -20°C before rinsing with PBS. Sections were blocked with 5% bovine serum

albumin (BSA) for 1 h at room temperature. Sections were incubated with anti-claudin-3 or anti-claudin-7

antibody diluted at 1:100 in BSA overnight at 4°C. Sections were rinsed with PBS, and the secondary antibody

ALEXA FLUOR™ 488 diluted at 1:600 in BSA was added to the sections for 1 h in the dark. Then, sections were

rinsed with PBS and mounted with FluoroGel + DAPI. Sections were examined and photographed using a Zeiss

Axio Imager M2 fluorescent light microscope equipped with Zen imaging software.

Genomic DNA extraction and 16S rRNA gene sequencing. Genomic DNA (gDNA) was extracted from

Control, uncolonized, and *A. timonensis*-colonized mouse feces using a QIAamp Fast DNA Stool Mini Kit

(Qiagen). gDNA was quantified using a BioSpectrometer (Eppendorf) and sent to the University of Alabama

Birmingham Microbiome Institutional Research Core for 16S rRNA sequencing. Sequencing data was processed

and analyzed using Qiime2 (v2024.5) and R (v4.4.1) along with the R package Qiime2R to import Qiime2 objects

to R.[60]

Plasma lipidomic analysis. Flash frozen mouse plasma samples were sent to MetwareBio for lipid-focused

extraction and targeted lipidomic analysis and quantification of host lipids. Fragmentation and retention time data

corresponding to SoLs was provided to facilitate targeted analysis and quantification of detectable SoLs.

Abundance data was analyzed using R (v4.4.1) along with the R package MetaboAnalystR (v4.0).[61]

OMV purification. *A. timonensis* was cultivated as described above and OMVs were obtained following

established isolation procedures with minimal modification.[62] Briefly, bacterial cells were harvested by

centrifugation at 12,000 x *g* for 30 minutes at 4°C. The supernatants were then filtered through 0.2 μm bottle top

aPES membrane filters (Thermo Scientific) to remove residual bacterial cells. Filtered supernatants were then

aliquoted into polycarbonate tubes for ultracentrifugation at 100,000 x *g* for 2 hours at 4°C. After

ultracentrifugation, the supernatant was decanted, replaced with fresh filtered supernatant, and centrifuged again

at 100,000 x *g* for 2 hours at 4°C until all the filtered supernatant was processed. The resulting OMV pellets were

then resuspended in filter sterilized 1X phosphate buffered saline (PBS) pH 8.0 and passed again through 0.2

µm syringe filters to ensure bacterial cells had been completely removed. Diluted OMV samples were then

pooled and concentrated in Spin-X UF 20 100 kDa PES membrane centrifugal filters (Corning). Protein

concentrations of the purified OMV samples were determined using a Bradford assay and samples were

prepared for electron microscopy at 200 ug/mL protein concentration.

Density gradient purification of OMVs from mouse plasma 667 µL aliquots of plasma from *A. timonensis*-

colonized IL10 KO mice were added to 3.333 mL of 60% iodixanol (OptiPrep) stock to create a 50% iodixanol

OMV load solution. 4 mL of 40%, 30%, 20%, and 10% Iodixanol solutions were prepared in 1X PBS. Iodixanol

solutions were then layered in order of decreasing Iodixanol concentration, beginning with the 50% iodixanol

OMV load solution, into polycarbonate tubes for ultracentrifugation, and 1 mL 1X PBS was added to the top of

each sample. The samples were then ultracentrifuged at 100,000 x *g* for 18 hours at 4°C, forming a continuous

density gradient. After ultracentrifugation, each sample was fractionated into 21 1 mL fractions, taken from the

top. The presence of SoLs in each fraction was determined by HPLC-MS and HPLC-MS/MS analysis. SoL-

containing fractions were then pooled and concentrated in Spin-X UF 20 100 kDa PES membrane centrifugal

filters (Corning).

Size Exclusion Chromatography of OMVs from mouse plasma. Concentrated OMV solutions were then

briefly centrifuged to remove cell debris and aggregates, before being subjected to Size Exclusion

Chromatography. Apex 4B Size Exclusion Columns (Everest Biolabs) were washed with 10 mL of 1X PBS before

0.5 mL of concentrated OMVs were applied to each column. The first 2 mL of eluate was discarded, and the

following eight 0.5 mL fractions were collected. OMV-containing fractions were determined via Bradford Assay

and confirmed by HPLC-MS/MS targeting SoL B. OMV-containing fractions were then pooled and stored at -

20°C prior to analysis.

TEM visualization. 10 µL aliquots of OMV samples either undiluted or prepared at 200 µg/mL protein

concentration were treated with 10 µL of 4% OsO4 (ThermoFisher Scientific) for five minutes. Then 10 µL of the

treated OMV sample was loaded onto copper formvar/carbon coated 200 mesh grids and allowed to settle for 5

minutes. Excess solution was dabbed away and the grids were negative stained with 10 µL of 5% uranyl acetate

(ThermoFisher Scientific) for 5 minutes before excess solution was dabbed away. Grids were allowed to dry and

samples were viewed using a Hitachi HT7800 TEM (Hitachi High-Tech Co.).

Dynamic light scattering of OMV samples. 1mL aliquots of OMV samples from plasma or *A. timonensis* culture were prepared at 100 µg/mL protein in 1X PBS and added to plastic cuvettes. Particle size distribution was then measured by dynamic light scattering using a Zetasizer Nano ZS (Malvern Panalytical). Experiments were performed in triplicate and all measurements were repeated three times.

OMV fractionation and SoL isolation. Approximately 200 µg of OMVs were dried and lyophilized prior to extraction with methanol. The methanol extract was deposited on silica and dried overnight prior to loading for open normal phase column chromatography. The OMV extract sample was eluted using five column volumes each of 15:1, 7:1, 5:1, 3:1, and 1:1 dichloromethane:methanol which was collected and dried by rotary evaporation and by lyophilization. Fraction samples were prepared at 20 ug/mL in dimethylsulfoxide for subsequent *in vitro* assays and HRMS/MS analysis. SoL B was purified from OMV fractions and tested for LPS contamination as previously described.[35]

Preparation of primary mouse macrophages. Primary mouse macrophages were prepared by first introducing 3 mL of 3% thioglycolate to mice via intraperitoneal injection. After 3 days, 10 mL of chilled PBS was introduced intraperitoneally to flush out macrophages. The cell suspension was then separated by centrifugation at 300 x g for 5 min. Cells were seeded in culture dishes containing DMEM with 10% FBS for 1 hour before being rinsed with serum-free DMEM two times to remove unattached cells.

Cell treatment. Samples of *A. timonensis* OMV fractions or *A. timonensis* OMV fractions + LPS were added to the culture media of primary mouse macrophages for 6 hours before the treatment media was removed. The cells were then washed twice with 1X PBS and detached from the cell culture plate with 0.25% trypsin. Cells were removed from the suspension by centrifugation and split equally for mRNA extraction and metabolite extraction. *A. timonensis* OMV fractions were prepared at 20 ug/mL and LPS was prepared at 1 ug/mL.

mRNA extraction and RT-qPCR. Cells were lysed with TRI Reagent (Zymo Research) and total RNA was extracted from the cell lysate using a Direct-Zol™ RNA MiniPrep Kit (Zymo Research) according to the manufacturer's protocol. The quality and quantity of RNA was then determined using a Nanodrop One (ThermoFisher) and 1,000 ng of mRNA from each sample was reverse-transcribed to cDNA using iScript cDNA Synthesis Kit (Bio-Rad). qPCR was conducted on a CFX96 system (Bio-Rad) using iQ SYBR Green Supermix (Bio-Rad). All primers used for qPCR analysis were synthesized by Integrated DNA Technologies. The 18S RNA

was used as a housekeeping gene. The relative amount of target mRNA was determined using the comparative

threshold (Ct) method by normalizing target mRNA Ct values to those of 18S RNA. PCR thermal cycling

conditions were 3 min at 95°C, and 40 cycles of 15 s at 95°C and 58 s at 60°C. Samples were run in triplicate.

Relative gene expression levels were calculated using the $\Delta\Delta C_T$ method and expression levels of 18S were used

to normalize the results.

HPLC-MS/MS analysis of OMV fractions and mouse plasma. OMV samples were extracted by addition of an

equal volume of methanol. The organic solvent was then removed and dried under a gentle stream of N_2. The

residue was reconstituted in methanol for HPLC-MS/MS analysis using a Vanquish HPLC (ThermoFisher

Scientific) coupled to a Q-Exactive HF-X hybrid Quadrupole-Orbitrap mass spectrometer using electrospray

ionization in positive mode. Analytical scale HPLC separation was performed on a Waters Xbridge BEH column

(2.1 x 100 mm) using a basic mobile phase consisting of solvent A: 60:40 acetonitrile:water + 0.1% formic acid

+ 10 mM ammonium formate and solvent B: 10:90 acetonitrile:isopropanol + 0.1% formic acid + 10 mM

ammonium formate delivered at a flow rate of 0.2 mL/min in a gradient starting from 37% B to 98% B over 20

minutes, followed by 98% B for 5 minutes to wash the column, back to 37% B over 3.5 minutes, and finally 37%

B for 7.5 minutes to re-equilibrate the column. MS scans were obtained in the orbitrap analyzer which was

scanned from 200 to 2000 *m/z* at a resolution of 60,000. MS/MS was conducted using data-dependent

acquisition with a resolution of 30,000, isolation window of 2.0 *m/z*, and dynamic exclusion time of 15 seconds.

Bioactive Molecular Networking and chemoinformatic analysis. HPLC-HRMS/MS data was processed

using MZmine3 following the GNPS FBMN workflow with minimal changes.[63] Molecular networks were

constructed using the quickstart GNPS FBMN setting with no changes.[50] Raw data used for this analysis was

deposited in the University of California, San Diego Center for Computational Mass Spectrometry MassIVE

database (https://doi.org/doi:10.25345/C55M62K1G). Bioactivity scores were assigned using a custom R script

which calculated Pearson correlation coefficients between each molecular feature and the activity of each

fraction.[51] Finally, bioactive molecular networks were visualized in Cytoscape v3.9.1.[64] Representative nodes

were selected for further chemoinformatic analysis with SIRIUS v6.0.6 using the CANOPUS and CSI:FingerID

tools to predict compound classes as well as representative node structures and substructure annotations.[52,53,55]

Statistics and reproducibility. For all experiments, sample sizes were chosen to achieve statistical significance, and data from outliers were excluded based on Grubbs' test for outliers. For animal experiments, randomization was applied to assign animals to groups and cages and blinding was applied in the analysis of histopathology data.

Author contributions

E.A.O., M.K.M., M.E., and J.L. designed the study. E.A.O., M.K.M., A.C., X.L., M.M., Y.W., L.E.V., T.Z., B.V., and R.T. conducted the study and acquired the data. E.A.O., M.K.M., M.E., and J.L. analyzed the data. E.A.O. and J.L. drafted the manuscript. All authors interpreted the data, read, and critically revised the manuscript, and approved the final version.

Acknowledgements

We acknowledge Drs. Michael Walla and William Cotham of the University of South Carolina Mass Spectrometry Facility for their assistance with HRMS/MS data acquisition and processing. We also acknowledge Dr. Diego Altomare, the director of the University of South Carolina Microarray Core Facility for his help with high throughput RT-qPCR for measuring cytokine expression, as well as Mr. Cole Espinosa for his assistance in *A. timonensis* OMV purification. We acknowledge the Instrumentation Resource Facility at the University of South Carolina School of Medicine for their assistance with processing and staining intestinal tissues for histopathology analysis. We acknowledge the University of Alabama Birmingham Microbiome Institutional Research Core for their assistance with 16S sequencing of IL10 KO mouse fecal samples.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Data availability statement

All data generated or analyzed during this study are included in the published article and its supplementary information files. Raw MS/MS data used for the bioactivity-based molecular networking analysis was deposited in the University of California, San Diego Center for Computational Mass Spectrometry MassIVE database (https://doi.org/doi:10.25345/C55M62K1G). Raw MS/MS data was also published alongside raw 16S sequencing data in Zenodo (https://doi.org/10.5281/zenodo.13987894).

Funding

657   This work was partially supported by the National Institutes of Health (NIH) through Grant

658   1R35GM150565 awarded to J.L. and another NIH Grant 1R01AI184916 awarded to M.E.

References

1. FitzGerald, R. & Smith, S. M. An overview of *Helicobacter pylori* infection. in *Helicobacter Pylori* vol. 2283 (Springer US, New York, NY, 2021).

2. Cani, P. D. & de Vos, W. M. Next-generation beneficial microbes: The case of *Akkermansia muciniphila*. *Front. Microbiol.* **8**, (2017).

3. Lau, K. *et al.* Bridging the gap between gut microbial dysbiosis and cardiovascular diseases. *Nutrients* **9**, 859 (2017).

4. Sheflin, A. M., Whitney, A. K. & Weir, T. L. Cancer-promoting effects of microbial dysbiosis. *Curr Oncol Rep* **16**, 406 (2014).

5. Sultan, S. *et al.* Metabolic influences of gut microbiota dysbiosis on inflammatory bowel disease. *Frontiers in Physiology* **12**, (2021).

6. Franzosa, E. A. *et al.* Gut microbiome structure and metabolic activity in inflammatory bowel disease. *Nat Microbiol* **4**, 293–305 (2019).

7. Chen, H. *et al.* Association of serum lipids with inflammatory bowel disease: a systematic review and meta-analysis. *Front. Med.* **10**, (2023).

8. Kayama, H. & Takeda, K. Emerging roles of host and microbial bioactive lipids in inflammatory bowel diseases. *European Journal of Immunology* **53**, 2249866 (2023).

9. Fan, F. *et al.* Lipidomic profiling in inflammatory bowel disease: Comparison between ulcerative colitis and Crohn's disease. *Inflammatory Bowel Diseases* **21**, 1511–1518 (2015).

10. Calzada, C., Vors, C., Penhoat, A., Cheillan, D. & Michalski, M.-C. Role of circulating sphingolipids in lipid metabolism: Why dietary lipids matter. *Front. Nutr.* **9**, (2023).

11. Quehenberger, O. & Dennis, E. A. The human plasma lipidome. *New England Journal of Medicine* **365**, 1812–1823 (2011).

12. Verstockt, B. *et al.* Sphingosine 1-phosphate modulation and immune cell trafficking in inflammatory bowel disease. *Nat Rev Gastroenterol Hepatol* **19**, 351–366 (2022).

13. Koutroumpakis, E. *et al.* Association between long-term lipid profiles and disease severity in a large cohort of patients with inflammatory bowel disease. *Dig Dis Sci* **61**, 865–871 (2016).

14. Brown, E. M. *et al. Bacteroides*-derived sphingolipids are critical for maintaining intestinal homeostasis and

symbiosis. *Cell Host and Microbe* **25**, 668-680.e7 (2019).

15. Lloyd-Price, J. *et al.* Strains, functions and dynamics in the expanded Human Microbiome Project. *Nature*

550, 61–66 (2017).

16. Parker, B. J., Wearsch, P. A., Veloo, A. C. M. & Rodriguez-Palacios, A. The genus *Alistipes*: Gut bacteria

with emerging implications to inflammation, cancer, and mental health. *Frontiers in Immunology* **11**, 906

(2020).

17. Sung, C. M. *et al.* Predicting clinical outcomes of cirrhosis patients with hepatic encephalopathy from the

fecal microbiome. *Cellular and Molecular Gastroenterology and Hepatology* **8**, 301-318.e2 (2019).

18. Iida, N. *et al.* Commensal bacteria control cancer response to therapy by modulating the tumor

microenvironment. *Science* (2013) doi:10.1126/science.1240527.

19. Zuo, K. *et al.* Disordered gut microbiota and alterations in metabolic patterns are associated with atrial

fibrillation. *GigaScience* **8**, giz058 (2019).

20. Dziarski, R., Park, S. Y., Kashyap, D. R., Dowd, S. E. & Gupta, D. Pglyrp-regulated gut microflora *Prevotella*

falsenii, *Parabacteroides distasonis* and *Bacteroides eggerthii* enhance and *Alistipes finegoldii* attenuates

colitis in mice. *PLOS ONE* **11**, e0146162 (2016).

21. Ishikawa, D. *et al.* Anti-inflammatory effects of Bacteroidota strains derived from outstanding donors of fecal

microbiota transplantation for the treatment of ulcerative colitis. *Inflammatory Bowel Diseases* izae080

(2024) doi:10.1093/ibd/izae080.

22. Lin, X. *et al.* Gut commensal *Alistipes shahii* improves experimental colitis in mice with reduced intestinal

epithelial damage and cytokine secretion. *mSystems* **10**, e01607-24 (2025).

23. Schwechheimer, C. & Kuehn, M. J. Outer-membrane vesicles from Gram-negative bacteria: Biogenesis and

functions. *Nature Reviews Microbiology* **13**, 605–619 (2015).

24. Kulp, A. & Kuehn, M. J. Biological functions and biogenesis of secreted bacterial outer membrane vesicles.

Annu Rev Microbiol **64**, 163–184 (2010).

25. Lynch, J. B., Alegado, R. A. & O'Toole, G. A. Spheres of hope, packets of doom: The good and bad of outer

membrane vesicles in interspecies and ecological dynamics. *Journal of Bacteriology* **199**, (2017).

26. Bonnington, K. E. & Kuehn, M. J. Protein selection and export via outer membrane vesicles. *Biochimica et*

Biophysica Acta (BBA) - Molecular Cell Research **1843**, 1612–1619 (2014).

27. Rocha, F. G., Ottenberg, G., Eure, Z. G., Davey, M. E. & Gibson, F. C. Sphingolipid-containing outer

membrane vesicles serve as a delivery vehicle to limit macrophage immune response to *Porphyromonas*

gingivalis. Infection and Immunity **89**, e00614-20 (2021).

28. Yoon, H., Ansong, C., Adkins, J. N. & Heffron, F. Discovery of *Salmonella* virulence factors translocated via

outer membrane vesicles to murine macrophages. *Infection and Immunity* **79**, 2182–2192 (2011).

29. Renelli, M., Matias, V., Lo, R. Y. & Beveridge, T. J. DNA-containing membrane vesicles of *Pseudomonas*

aeruginosa PAO1 and their genetic transformation potential. *Microbiology* **150**, 2161–2169 (2004).

30. Hickey, C. A. *et al.* Colitogenic *Bacteroides thetaiotaomicron* antigens access host immune cells in a

sulfatase-dependent manner via outer membrane vesicles. *Cell Host and Microbe* **17**, 672–680 (2015).

31. Bomberger, J. M. *et al.* Long-distance delivery of bacterial virulence factors by *Pseudomonas aeruginosa*

outer membrane vesicles. *PLOS Pathogens* **5**, e1000382 (2009).

32. Schaack, B. *et al.* Microbiota-derived extracellular vesicles detected in human blood from healthy donors.

International Journal of Molecular Sciences **23**, 13787 (2022).

33. Berg, D. J. *et al.* Rapid development of colitis in NSAID-treated IL-10-deficient mice. *Gastroenterology* **123**,

1527–1542 (2002).

34. Hale, L. P., Gottfried, M. R. & Swidsinski, A. Piroxicam treatment of IL-10-deficient mice enhances colonic

epithelial apoptosis and mucosal exposure to intestinal bacteria. *Inflammatory Bowel Diseases* **11**, 1060–

1069 (2005).

35. Older, E. A. *et al.* Biosynthetic enzyme analysis identifies a protective role for TLR4-acting gut microbial

sulfonolipids in inflammatory bowel disease. *Nat Commun* **15**, 9371 (2024).

36. Walker, A. *et al.* Sulfonolipids as novel metabolite markers of *Alistipes* and *Odoribacter* affected by high-fat

diets. *Scientific Reports* **7**, 11047 (2017).

37. Lu, Z., Ding, L., Lu, Q. & Chen, Y.-H. Claudins in intestines: Distribution and functional significance in health

and diseases. *Tissue Barriers* **1**, e24978 (2013).

38. Xing, T. *et al.* Tight junction protein claudin-7 Is essential for intestinal epithelial stem cell self-renewal and

differentiation. *Cell Mol Gastroenterol Hepatol* **9**, 641–659 (2019).

39. Zhao, M. *et al.* Immunological mechanisms of inflammatory diseases caused by gut microbiota dysbiosis: A

review. *Biomedicine & Pharmacotherapy* **164**, 114985 (2023).

40. Maharshak, N. *et al.* Altered enteric microbiota ecology in interleukin 10-deficient mice during development

and progression of intestinal inflammation. *Gut Microbes* **4**, 316–324 (2013).

41. Yang, Y. *et al.* Dietary eggshell membrane powder improves survival rate and ameliorates gut dysbiosis in

interleukin-10 knockout mice. *Front. Nutr.* **9**, (2022).

42. Yan, Y. *et al. Bacteroides uniformis*-induced perturbations in colonic microbiota and bile acid levels inhibit

TH17 differentiation and ameliorate colitis developments. *npj Biofilms Microbiomes* **9**, 1–14 (2023).

43. Chiurchiù, V., Leuti, A. & Maccarrone, M. Bioactive lipids and chronic inflammation: Managing the fire within.

Frontiers in Immunology **9**, 38 (2018).

44. Funabashi, M. *et al.* A metabolic pathway for bile acid dehydroxylation by the gut microbiome. *Nature* **582**,

566–570 (2020).

45. Bermúdez, M. A., Rubio, J. M., Balboa, M. A. & Balsinde, J. Differential mobilization of the phospholipid and

triacylglycerol pools of arachidonic acid in murine macrophages. *Biomolecules* **12**, 1851 (2022).

46. Hou, L. *et al.* Identification and biosynthesis of pro-inflammatory sulfonolipids from an opportunistic pathogen

Chryseobacterium gleum. ACS Chemical Biology **17**, 1197–1206 (2022).

47. Godchaux, W. & Leadbetter, E. R. Sulfonolipids are localized in the outer membrane of the gliding bacterium

Cytophaga johnsonae. Archives of Microbiology **150**, 42–47 (1988).

48. Shalon, D. *et al.* Profiling the human intestinal environment under physiological conditions. *Nature* **617**, 581–

591 (2023).

49. Beemelmanns, C. *et al.* Synthesis of the rosette-inducing factor RIF-1 and analogs. *Journal of the American

Chemical Society* **136**, 10210–10213 (2014).

50. Nothias, L.-F. *et al.* Feature-based molecular networking in the GNPS analysis environment. *Nat Methods*

17, 905–908 (2020).

51. Nothias, L.-F. *et al.* Bioactivity-based molecular networking for the discovery of drug leads in natural product

bioassay-guided fractionation. *J. Nat. Prod.* **81**, 758–767 (2018).

52. Dührkop, K. *et al.* SIRIUS 4: a rapid tool for turning tandem mass spectra into metabolite structure

information. *Nature Methods* **16**, 299–302 (2019).

53. Dührkop, K. *et al.* Systematic classification of unknown metabolites using high-resolution fragmentation

mass spectra. *Nat Biotechnol* **39**, 462–471 (2021).

54. Heinonen, M., Shen, H., Zamboni, N. & Rousu, J. Metabolite identification and molecular fingerprint

prediction through machine learning. *Bioinformatics* **28**, 2333–2341 (2012).

55. Hoffmann, M. A. *et al.* High-confidence structural annotation of metabolites absent from spectral libraries.

Nat Biotechnol **40**, 411–421 (2022).

56. Maeda, J. *et al.* Inhibitory effects of sulfobacin B on DNA polymerase and inflammation. *International Journal*

of Molecular Medicine **26**, 521–527 (2010).

57. Deng, L. *et al.* Colonization with ubiquitous protist *Blastocystis* ST1 ameliorates DSS-induced colitis and

promotes beneficial microbiota and immune outcomes. *npj Biofilms Microbiomes* **9**, 1–12 (2023).

58. Dong, D. *et al. Clostridioides difficile* aggravates dextran sulfate solution (DSS)-induced colitis by shaping

the gut microbiota and promoting neutrophil recruitment. *Gut Microbes* **15**, 2192478 (2023).

59. Wieland Brown, L. C. *et al.* Production of α-galactosylceramide by a prominent member of the human gut

microbiota. *PLoS Biol* **11**, e1001610 (2013).

60. Bisanz, J. E. qiime2R: Importing QIIME2 artifacts and associated data into R sessions. (2018).

61. Pang, Z., Chong, J., Li, S. & Xia, J. MetaboAnalystR 3.0: Toward an optimized workflow for global

metabolomics. *Metabolites* **10**, 186 (2020).

62. Sheikh, A., Zechmann, B., Sayes, C. M., Taube, J. H. & Greathouse, K. L. A preparation of bacterial outer

membrane with osmium tetroxide and uranyl acetate co-stain enables improved structural determination by

transmission electron microscopy. *Microscopy* **72**, 515–519 (2023).

63. Schmid, R. *et al.* Integrative analysis of multimodal mass spectrometry data in MZmine 3. *Nat Biotechnol* **41**,

447–449 (2023).

64. Shannon, P. *et al.* Cytoscape: A software environment for integrated models of biomolecular interaction

networks. *Genome Res.* **13**, 2498–2504 (2003).