Sphingolipid imbalance aggravates tau pathology by endomembrane rigidification and rupture

Jessica Tittelmeier[1], Carl Alexander Sandhof[2,3], Nicole Martin[1], Deike El-Kabarity[1], Soki-Bradel Ngonza-Nito[4], Ronald Melki[4] and Carmen Nussbaum-Krammer[1,2,*]

[1]Chair of Neuroanatomy, Institute of Anatomy, Faculty of Medicine, Ludwig-Maximilians University of Munich (LMU), Pettenkoferstrasse 11, 80336 Munich, Germany

[2]Center for Molecular Biology of Heidelberg University (ZMBH) and German Cancer Research Center (DKFZ), DKFZ-ZMBH Alliance, Im Neuenheimer Feld 282, 69120 Heidelberg, Germany

[3]Chemical Neurobiology Laboratory, Center for Genomic Medicine, Departments of Neurology and Psychiatry, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114, USA

[4]Institute Francois Jacob (MIRCen), CEA, and Laboratory of Neurodegenerative Diseases, CNRS, Fontenay-Aux-Roses, France

*Correspondence should be addressed to carmen.nussbaum@med.uni-muenchen.de

Summary

Endolysosomal dysfunction is a hallmark of Alzheimer's disease (AD) and related tauopathies, yet

underlying mechanisms remain poorly understood. This study investigates the role of sphingolipid

metabolism in maintaining endolysosomal membrane integrity and its impact on tau aggregation

and toxicity in *Caenorhabditis elegans* and human cell culture models. Fluorescence recovery after

photobleaching and C-Laurdan dye imaging revealed that silencing sphingolipid metabolism genes

reduced endolysosomal vesicle membrane fluidity, increasing their rupture. The accumulation of

aggregated tau in endolysosomal vesicles further aggravated endomembrane rigidification and

damage and promoted seeded tau aggregation, potentially by facilitating the escape of tau seeds

from the endolysosomal system. Supplementation with unsaturated fatty acids improved

membrane fluidity, suppressing endolysosomal rupture and seeded tau aggregation in cell models,

and alleviating tau-associated neurotoxicity in *C. elegans*. This study provides mechanistic insights

into how impaired sphingolipid homeostasis drives endolysosomal membrane damage and

contributes to the progression of tau pathology, suggesting that restoring membrane fluidity may

offer a therapeutic strategy for AD.

Keywords: Alzheimer's disease / tau propagation / sphingolipid metabolism / endolysosomal

system / lipid homeostasis

Introduction

The gradual accumulation of microtubule-associated protein tau (MAPT/tau) aggregates is a hallmark of Alzheimer's disease (AD) and related tauopathies. Misfolded tau species exhibit prion-like behavior by self-replicating and spreading from cell to cell[1,2]. This contributes to the progression of pathology and neurotoxicity, eventually culminating in widespread neuronal dysfunction and degeneration[3]. At the molecular level, tau aggregates replicate by templating the conversion of native tau into an amyloid conformation, promoting its accumulation into fibrillar aggregates. For this to occur at the cellular level, seeding-competent tau species (or tau "seeds") must be released from a donor cell and taken up into the cytosol of the neighboring receiving cell in order to come into direct contact with the cytosolic native tau protein[4-6].

In this process, the autophagy-lysosomal pathway (ALP), which is an important clearance route for tau[7], appears to play a central role. Studies have demonstrated impaired ALP function in the brains of tauopathy patients as well as in animal and cell models, showing that the accumulation of abnormal autolysosomal and endolysosomal vesicles correlating with neuronal toxicity[8-11]. Inhibition of autophagic clearance of tau increases its secretion and spreading[12]. In addition, the accumulation of misfolded tau within endolysosomes leads to a destabilization and rupture of these vesicles[13-18]. While intact endolysosomes normally restrict tau seeds from reaching cytosolic monomers, membrane rupture allows their escape. Recent studies highlight this escape as a critical rate-limiting step in seeded tau propagation[13-18]. Endolysosomal damage not only promotes the propagation of pathological aggregates, but also causes the release of hydrolytic enzymes into the cytosol, leading to cellular damage and death[19]. Although recent efforts to identify pathways involved in endolysosomal damage and repair have intensified, many aspects of these processes remain unknown.

To identify cellular factors that are critical for the integrity of endolysosomal vesicles, we recently performed an unbiased genome-wide RNAi screen in *C. elegans*[20]. One of the pathways identified was sphingolipid (SL) metabolism. SLs constitute a large and diverse class of lipids that are involved in various physiological processes[21,22]. They consist of two main building blocks, a long-chain base with a serine backbone and an attached long acyl chain. This characteristic chemical structure mediates unique biophysical properties. SL biosynthesis and degradation rely on a specialized enzymatic machinery distinct from that of other lipids[23] (Figure 1A). Functionally, SLs, along with glycerolipids and sterols, serve as structural components of cell membranes,

contributing to the stability and fluidity of membranes and to the organization of microdomains
such as lipid rafts. Beyond their structural role, SLs serve as bioactive molecules participating in
cellular signaling pathways that control cell growth, differentiation, apoptosis, and intercellular
communication[24]. Abnormalities in SL metabolism have been observed during aging and in
neurodegenerative conditions, including AD, underscoring their potential significance in the
pathogenesis of these disorders[25,26]. Interestingly, recent evidence suggests that sphingolipid
accumulation disturbs the endolysosomal pathway and induces or potentiates endolysosomal
membrane rupture[27,28]; however, the mechanisms underlying membrane destabilization remain
unclear.
Here we investigated the role of SL metabolism in the maintenance of endolysosomal vesicle
integrity. Our results revealed that silencing of SL metabolism genes led to a marked reduction in
membrane fluidity, particularly in endolysosomal vesicles, thereby increasing their fragility and
rupture. The accumulation of aggregated tau had an additive effect and led to a further reduction
in endomembrane fluidity, which aggravated the damage to the endolysosomal compartment.
Increased membrane rigidity also facilitated seeded tau aggregation. Improvement of membrane
fluidity through supplementation of polyunsaturated fatty acids (PUFAs) counteracted tau
propagation in cells and tau-associated neurotoxicity in *C. elegans*.
This study highlights the interplay between lipid metabolism and proteostasis and provides insights
into how disturbances in sphingolipid homeostasis contribute to endolysosomal dysfunction and
tau pathology. In addition, it provides a possible mechanistic explanation for the beneficial effect
of PUFA supplementation that has been observed in AD patients and suggests that restoring
membrane fluidity may help mitigate tau toxicity and disease progression.
98
99

Results

Knockdown of sphingolipid metabolism genes induces lysosomal rupture

Several of the strongest hits identified in the unbiased genome-wide RNAi screen we performed to identify modulators of endolysosomal vesicles integrity were genes involved in SL metabolism (Figure 1A). Knockdown (KD) of these genes (the serine incorporator *R11H6.2*, the 3-ketodihydrosphingosine reductase *Y37E11AM.3*, the hydroxyacyl-CoA dehydratase *hpo-8*, the cytochrome b5 reductases *hpo-19* and *T05H4.4*, the saposin *spp-10*, the sphingosine kinase *sphk-1*, and the sphingosine-1-phosphate lyase *spl-1*) via RNA interference (RNAi) induced endolysosomal membrane damage as detected by a galectin puncta assay[29] (Figure 1B-D). Under steady-state conditions, human galectin-3 fused to superfolder-GFP (sfGFP::LGALS3) exhibits an even distribution within the cytosol of hypodermal cells. Upon endolysosomal vesicle damage and exposure of their lumen to the cytosol, sfGFP::LGALS3 binds exposed β-galactoside and re-localizes, yielding visible puncta[29].

The screen identified distinct steps in sphingolipid metabolism catalyzed by non-redundant genes or targeted by a single RNAi clone, as in the case of *hpo-19/T05H4.4* which are both depleted by *hpo-19* RNAi. However, other steps in this pathway, which are mediated by two or more proteins may have been missed in the screen. Indeed, simultaneous KD of selective redundant genes using co-RNAi revealed additional genes functioning in *de novo* sphingolipid biosynthesis that also induced endolysosomal rupture, including the serine palmitoyltransferases *sptl-1* and *-3* and the ceramide synthases *hyl-1* and *-2* (Figure S1A). In addition, the co-KD of all four glucocerebrosidases *gba-1-4* or only *gba-2-4*, also triggered endolysosomal rupture (Figure S1B). Hence, these findings suggest that the KD of genes involved in both biosynthesis and degradation of SLs contributes to endolysosomal rupture, highlighting the tightly regulated nature of sphingolipid metabolism. Any disruption in this balance poses challenges to endolysosomal integrity.

KD of sphingolipid metabolism genes particularly affects membrane fluidity of endolysosomal vesicles

SLs are important components of eukaryotic cell membranes and genetic modulation of SL metabolism likely affects the lipid composition and biophysical properties of membranes. Similar to their mammalian isoforms, the N-acyl chains of *C. elegans* SLs contain predominantly long and

saturated fatty acid chains[30,31], which tend to pack tightly within the lipid bilayer. As such, a higher

proportion of SLs generally reduces membrane fluidity, leading to increased order, rigidity, or

viscosity[32]. Consequently, KD of the hits involved in SL degradation should lead to SL

accumulation and reduced membrane fluidity, while KD of the hits involved in SL biosynthesis

should have the opposite effect.

To evaluate the membrane fluidity of endolysosomal membranes, fluorescence recovery after

photobleaching (FRAP) was employed on *C. elegans* that express the Lysine/Arginine Transporter

1 (LAAT-1) tagged with mCherry, which localizes to the lysosomal membrane[33]. FRAP enables

the assessment of the lateral mobility of LAAT-1 within the lysosomal membrane, thereby

indirectly measuring its fluidity. We focused our analysis on the hypodermis, where we expressed

the sfGFP::LGALS3 reporter and observed endolysosomal rupture following KD of the SL

metabolism genes (Figure 1C, D). Interestingly, KD of all of our hits significantly increased the

time required to recover half of the maximum fluorescence intensity (t_{half}), indicating less fluid

lysosomal membrane in the hypodermis (Figure 2A-C, E, F). With some gene KDs, we also

observed a decrease in the maximal recovered signal after bleaching (Figure 2D, G). The decrease

in lysosomal membrane fluidity was not specific to the hypodermis as the KD of *spl-1* also

increased the t_{half} of LAAT-1::mCherry in intestinal lysosomes (Figure S2A-D). To discern

whether the detrimental effect of KD on membrane fluidity is confined to lysosomal membranes

or extends to other cellular membranes, we utilized a *C. elegans* strain expressing a plasma

membrane-anchored GFP[34]. Intriguingly, this reporter showed no change in the t_{half} with any KD

and only *spl-1* KD led to decrease in the maximal recovered signal (Figure 2H-K). Hence, the

endolysosomal membrane seems to be particularly sensitive to a disruption of SL metabolism in

contrast to the plasma membrane. Dysregulation of both, synthesis and degradation of SLs reduces

the fluidity of the endolysosomal membrane and makes it more susceptible to rupture.

KD of sphingolipid metabolism genes affects membrane fluidity in human cells

Next, we asked whether dysregulation of SL metabolism exerts similar effects in human cells. To

this end, we employed HEK293T cells and C-Laurdan dye to evaluate membrane fluidity (Figure

3A). This approach leverages the unique properties of C-Laurdan, which emits fluorescent light of

varying wavelengths in response to changes within the phospholipid bilayer, particularly reflective

of membrane fluidity[35,36]. The shift in emission profile between liquid-disordered and liquid-

ordered phases allows a quantitative assessment of the membrane order by calculating the

ratiometric relationship of the fluorescence intensity recorded in two spectral channels, known as

a generalized polarization (GP) value[35] (Figure 3A). Decreasing GP values are indicative of

increasingly fluid membrane while increasing GP values denote increased rigidity. Of note, we

exposed the cells to C-Laurdan for an extended time of 2 hours (instead of 30 min as recommended

in the original protocol), to promote internalization of the dye and thus increase the staining of

endolysosomal membranes relative to the plasma membrane[35]. Upon KD of *SPHK2*, one of the

human homologs of the *C. elegans sphk-1* gene, the C-Laurdan fluorescence GP ratio augmented

significantly, indicating increased membrane rigidity (Figure 3B, D and Figure S3A). Of note,

Lipofectamine 2000 was used to deliver siRNA into the cells and lipid-based transfection reagents

have been shown to impact endolysosomal membranes[37]. Indeed, Lipofectamine alone slightly

increased membrane fluidity (Figure 3C). Taken together, FRAP and C-Laurdan analyses revealed

that disrupted SL homeostasis led to more ordered and rigid endolysosomal membranes, in both

C. elegans and human cells.

Fibrillar tau increases membrane rigidity and exacerbates endolysosomal damage

To investigate whether disruption of SL homeostasis also affects endolysosomal integrity in

human cells, we generated a HEK293T cell line stably expressing the same sfGFP-LGALS3

reporter used for the galectin puncta assay in *C. elegans*. To further mimic the *C. elegans* model,

in which aggregation-prone tau fragments are chronically transferred from the touch receptor

neurons into the hypodermis where the sfGFP-LGALS3 reporter is expressed[20], recombinant

1N4R tau fibrils were added to sfGFP-LGALS3 expressing HEK293T cells on top of the KD of

SPHK2. Interestingly, both the addition of tau fibrils alone and the KD of *SPHK2* alone already

caused rupture of endolysosomes in sfGFP-LGALS3 cells (Figure 3D, E and Figure S3B). The

combination of both had an additive effect (Figure 3D, E and Figure S3B).

We therefore asked whether the KD of SLs alone would also already rupture hypodermal

endolysosomes in *C. elegans* and whether accumulation of transmitted tau would exacerbate this

process. To answer this question, we determined the amount of hypodermal sfGFP-LGALS3 foci

in F3$^{\Delta K281}$::mCherry *C. elegans* compared to control animals expressing only the mCherry tag. KD

of *Y37E11AM.3* alone did not increase lysosomal rupture compared to the EV control RNAi, only

the presence of tau spreading resulted in more lysosomal rupture relative to the control (Figure

3F). KD of the remaining hits resulted in near 100% animals with hypodermal sfGFP-LGALS3
foci in both the F3$^{\Delta K281}$::mCherry and mCherry control strains. To test whether the additional
negative impact of the F3 tau fragment might be masked by the strength of the RNAi clones,
different dilutions of the RNAi bacteria were tested. Under conditions where the RNAi KD did
not saturate the readout of the assay, KD of *spl-1, hpo-8,* and *hpo-19* alone already affected
lysosomal integrity (Figure 3G). Moreover, F3$^{\Delta K281}$::mCherry transgenic animals showed a
significantly higher level of endolysosomal rupture upon KD of *spl-1, hpo-8,* and *hpo-19* compared
to the mCherry control (Figure 3G). Diluting the *R11H6.2* RNAi clone showed no significant
differences, and KD of *sphk-1* still resulted in 100% endolysosomal rupture-positive animals in
both genetic backgrounds even after dilution of the RNAi bacteria to below 5% and could therefore
not be assessed. In conclusion, chronic transmission of the F3$^{\Delta K281}$::mCherry fragment from the
touch receptor neurons to the hypodermis impaired the endolysosomal system exacerbating the
rupture due to perturbation of SL homeostasis.
Since disruption of SLs metabolism led to an increase in membrane rigidity resulting in
endolysosomal rupture, and fibrillar tau also ruptured endolysosomal membranes (Figure 3D, E)[14],
we asked whether tau itself could also affect membrane fluidity. To test this, membrane fluidity
was assessed for somas of the posterior (PLM) touch receptor neurons of *C. elegans* using C-
Laurdan dye. Indeed, expression of the tau fragment led to a decrease in membrane fluidity
compared to expression of the mCherry tag alone (Figure 3H, I). Also, HEK293T cells incubated
with tau fibrils for 5 hours exhibited an increased GP index (Figure 3J, K). Importantly, this was
specific to fibrillar tau, as monomeric tau had no effect on membrane fluidity (Figure 3L and
Figure S3C). Thus, aggregated forms of tau can cause rigidification of the endolysosomal
membrane, resulting in increased fragility and rupture.
216

Membrane rigidification exacerbates seeded tau aggregation
The packing of fatty acids within the membrane influences membrane fluidity. Saturated fatty
acids (SFAs) have straight hydrocarbon chains forming a tightly packed, highly ordered and rigid
membrane. To investigate the role endomembrane lipid homeostasis plays in tau propagation, we
introduced an SFA to rigidify membranes. Incubating HEK293T cells with the SFA palmitic acid
(PA) conjugated to BSA increased membrane rigidity (Figure 4A, B). In addition, cells incubated

with PA and tau fibrils had an even more rigid membrane than cells incubated with tau fibrils alone

(Figure 4C, D).

An important step in tau propagation is tau seeds escape from the endolysosomal compartment,

which can then propagate themself by recruiting native monomeric tau. Therefore, we asked if the

PA-induced loss of membrane fluidity impairs endolysosomal integrity. Worms grown on plates

supplemented with PA displayed an increase in endolysosomal rupture (Figure 4E). In cells, the

addition of PA alone did not lead to endolysosomal rupture; however due to compounding impact

of PA and tau fibrils on membrane fluidity, the addition of PA increased fibrillar tau-induced

endolysosomal rupture (Figure 4F, G). Hence, loss of membrane fluidity mediated by fibrillar tau

and/or SFAs negatively affects endolysosomal integrity, resulting in increased endolysosomal

rupture.

Impairment of endolysosomal integrity may enhance tau propagation as tau could escape from the

endolysosomal compartments, reach the cytosol and recruit native monomeric tau, promoting

seeded tau aggregation. Given the observed membrane rigidification after pre-loading with SFAs,

we investigated whether membrane lipid composition influences tau propagation using a tau

biosensor cell line expressing Venus-tagged full-length P301S mutant 0N4R tau (tau-Venus)[38,39].

PA-induced membrane rigidification led to an increase in tau-Venus foci upon seeding with tau

fibrils (Figure 4H, I). Moreover, KD of *SPHK2* also increased the formation of tau-Venus foci due

to SPHK2 KD-mediated membrane rigidification (Figure S4C, D). This highlights a positive

correlation between membrane rigidity and seeded tau aggregation. Changes in membrane fluidity,

which compromise endolysosomal integrity may promote the entry of seeds into the cytosol. This

would consequently foster the seeded propagation of tau aggregation.

Increasing membrane fluidity by PUFAs rescues neurotoxic phenotypes

Given that membrane stiffening exacerbates tau propagation, we wondered whether we could

prevent this by improving membrane viscosity. While SFA allow for tightly packed lipids,

unsaturated fatty acids have one or more double bonds in their hydrocarbon chain that introduce

kinks in their structure, resulting in loosely packed lipids and increased membrane fluidity.

Therefore, we investigated whether we could rescue tau pathology using polyunsaturated fatty

acids (PUFAs). Incubating HEK293T cells with the ω-3 PUFA α-linolenic acid (ALA) increased

membrane fluidity (Figure 5A and Figure S5A). Moreover, pre-loading with ALA prevented tau-

induced membrane rigidification and decreased the GP-index of membranes even after the addition
of tau fibrils (Figure 5B, C). Importantly, pre-loading cells with ALA also counteracted fibrillar
tau-mediated endolysosomal rupture (Figure 5D, E and Figure S5B).

Since ALA prevented endolysosomal rupture, we expected it to also prevent seeded tau
aggregation. Indeed, tau-Venus cells pre-loaded with ALA exhibited significantly fewer foci when
exposed to tau fibrils (Figure 5F and Figure S5C). These data show that ALA supplementation
counteracts tau-mediated decrease in membrane fluidity, thereby preventing endolysosomal
rupture and decreasing seeded tau aggregation.

Finally, we asked whether ALA would also reduce the toxicity associated with aggregated tau.
Expression of F3$^{\Delta K281}$::mCherry in touch receptor neurons resulted in an age-dependent decline in
the response to gentle touch from day 4, corresponding to the first day of adulthood to day 8,
corresponding to day 5 of adulthood (Figure 5H, and Figure S5D)[20]. Addition of ALA to the
growth medium mitigated this loss of function (Figure 5H and Figure S5D) and also reduced
neurotoxicity (Figure 5I, J). Thus, supplementation with ALA not only reduced endolysosomal
damage but also attenuated neurotoxic phenotypes associated with aggregated tau.

Discussion

In a previous genome-wide screen in *C. elegans*, we identified genes involved in SL metabolism as crucial for maintaining endolysosomal integrity[20]. In this study we explored the mechanisms by which disruption of SL homeostasis leads to endolysosomal rupture and how it affects tau pathology. We found that imbalanced SL metabolism leads to decreased membrane fluidity, and subsequent endolysosomal rupture. This promoted seeded tau aggregation in a biosensor cell line exposed to fibrillar tau, implying that dysregulation of SL metabolism might boost disease progression by enhancing propagation of tau aggregation.

The central nervous system (CNS) is particularly rich in lipids. Approximately 50% of the brain's dry weight consists of lipids with high concentrations of cholesterol, phospholipids and sphingolipids [40,41]. Alterations in lipid and SL metabolism have been observed not only during the aging process, but also in various late-onset neurodegenerative diseases including AD[26,42]. While aging is the major risk factor for AD, one of the most significant genetic risk factors for AD is the apolipoprotein E (ApoE) polymorphism, specifically the ApoE4 allele[43]. ApoE is the primary lipoprotein in the brain, crucial for lipid transport and metabolism. Besides cholesterol, the ApoE4 variant has been shown to influence SL metabolism, thereby affecting the pathogenesis of AD from its early stages[26,42,44]. While disturbances in SL homeostasis are increasingly recognized as a contributing factor to the onset and progression of the disease[26,42], the underlying mechanisms are poorly understood.

Our results show that reducing the expression of genes involved in both, biosynthesis and degradation of SLs significantly reduced the fluidity of endolysosomal membranes, making them more fragile and prone to rupture. KD of the saposin *spp-10*, the sphingosine kinase *sphk-1*, and the sphingosine-1-phosphate lyase *spl-1* blocks the degradation of complex SLs and should lead to an increase in their relative abundance, and therefore to a decrease in membrane fluidity. KD of the very-long-chain (3R)-3-hydroxyacyl-CoA dehydratase *hpo-8*, which is involved in fatty acid elongation, has been demonstrated to promote the incorporation of long-chain UFAs into phospholipids, thereby restoring plasma membrane fluidity[45]. The absence of *hpo-8* leads to membrane rigidification[45]. Conversely, KD of the serine incorporator *R11H6.*2 and the 3-ketodihydrosphingosine reductase *Y37E11AM.3*, which function in the SL de novo synthesis pathway, should theoretically decrease total SL content and consequently increase membrane fluidity. The cytochrome b5 reductases *hpo-19* and *T05H4.4*, required for lipid desaturation, are

involved in the conversion of dihydroceramide to ceramide and the biosynthesis of PUFAs[46]; KD

of these genes is expected to reduce the levels of SLs and PUFAs, which could theoretically either

increase or decrease membrane fluidity, depending on which metabolic pathway is predominantly

affected.

Our observation that KD of all these genes results in decreased membrane fluidity is seemingly

counterintuitive. However, SLs form a complex metabolic network and can interconvert, allowing

a dynamic adaptation of SL levels. In addition, the cell likely compensates for the loss of certain

SLs by increasing the biosynthesis of other lipid molecules. This dynamic interplay makes it

difficult to predict how individual genes affect the endolysosomal membrane lipid composition.

Genes that are involved in apparently opposing steps of SL metabolism could therefore still lead

to an accumulation of similar SL species.

Future lipidomic analyses are necessary to determine the SL and lipid content of endolysosomal

membranes in our model systems. Such studies will provide a clearer understanding of how genetic

manipulation of SL metabolism affects endomembrane lipid composition and could pinpoint the

exact species likely responsible for the observed membrane rigidification.

AD brains exhibit elevated levels of sphingosine, ceramide 1-phosphate, and ceramide compared

to control subjects[42]. Multivariate analysis and machine learning also identified these SL species

as key contributors to AD[42]. These and other analyses were performed in different brain regions

but without cell type or subcellular resolution. We found that disruption of *C. elegans* SL

homeostasis primarily affected endolysosomal membranes and had little effect on plasma

membrane fluidity. The lysosomal membrane is particularly rich in sphingolipids[47], which seems

to make it more susceptible to disruption of SL homeostasis. Studies in yeast have shown that

increased membrane lipid saturation affects the vacuole —the functional counterpart of

mammalian lysosomes—as well as the nuclear envelope, whereas the plasma membrane and

mitochondria remained relatively unaffected[48]. These findings indicate that cellular membranes

vary in their resilience to lipid saturation stress, potentially due to differences in lipid composition

and protective mechanisms[48,49]. To better understand the implications of SL alterations in AD, it

would be important to examine whether the observed changes primarily originate from

endolysosomal membranes or also from other organelles, such as the endoplasmic reticulum and

the nucleus, or from the plasma membrane.

Intriguingly, expression of the aggregation-prone F3 tau fragment also enhanced membrane
rigidity with SL homeostasis disruption having an additive effect, exacerbating endomembrane
damage. Thus, alterations in SL metabolism could be not only a cause but also a consequence of
AD related pathology[26,42]. The accumulation of misfolded tau in lysosomes could impair the
lysosomal clearance capacity. This dysfunction may reduce the activity of acid hydrolases, leading
to an accumulation of biomolecules, including membrane lipids. Impairment of membrane lipid
metabolism could in turn further compromise lysosomal integrity and function. Both processes -
accumulation of SLs and protein aggregates - may create a vicious self-promoting cycle of
lysosomal dysfunction that eventually leads to neurodegeneration.
Supplementation of SFAs exacerbated the negative impact of the F3 tau fragment on
endomembrane fluidity. Given the high levels of saturated and trans fats in the Western diet[50], it
is plausible that dietary factors also contribute to the dysregulation of lipid homeostasis in the
brain. The Mediterranean UFA-rich diet has been associated with healthy aging, while reduced
UFA brain levels have been correlated with increased tau and Aβ pathology and decreased
cognitive function in AD[51,52]. Restoring membrane fluidity through the supplementation of PUFAs
mitigated tau propagation in cells and reduced tau-associated neurotoxicity in *C. elegans*. These
results suggest a potential mechanism for the beneficial effects of PUFA supplementation observed
in AD patients[53,54]. Future studies are needed to evaluate whether these benefits are related to an
increase in the fluidity of endolysosomal membranes and the consequent prevention of their
damage by protein aggregates and/or impaired SL homeostasis.
In sum, this study revealed an important role of endolysosomal membrane homeostasis in disease
progression providing a possible mechanism for the detrimental effect of SL imbalance and
beneficial effect of PUFA supplementation that have been observed in AD patients.
355
356

Methods

Maintenance of *C. elegans* and age synchronization

All animals were cultured using standard methods[55]. If not otherwise indicated, worms were grown on nematode growth medium (NGM) plates seeded with *E. coli* strain OP50 at 20 °C. For RNAi, NGM medium were supplemented with Ampicillin, Tetracycline, and IPTG, seeded with the respective HT115 *E. coli* RNAi clones and grown at 20 °C. Animals were age-synchronized by bleaching. Briefly, gravid adults were dissolved in 20% sodium hypochlorite solution. The surviving *C. elegans* embryos were hatched overnight in M9 buffer with gentle rocking at 20 °C. The next day, appropriate amounts of L1 larvae were added to the plates.

***C. elegans* fatty acid supplementation**

Polyunsaturated fatty acids were dissolved in EtOH and stored in the dark at -20 °C under a N_2 atmosphere to prevent oxidation. NGM was cooled after autoclaving to 55 °C. To enhance fatty acid distribution, NP-40 substitute (0.001% v/v) was added to the media and subsequently either 0.3 mM of the desired fatty acid solution or the ethanol solvent as control and stirred for 5 min before plate pouring [56]. Palmitic acid (PA) was supplemented by enriching the OP50 bacterial food source. Overnight cultures were grown in the presence of 2 mM of PA dissolved in EtOH or EtOH. The next day, OP50 bacteria were pelleted by centrifugation (10 min at 4500x g) and washed 2x with M9 buffer[57]. 10x concentrated bacteria (in M9) were then seeded on NGM plates.

Quantification of lysosomal rupture (galectin puncta assay) in *C. elegans*

Quantification of animals showing hypodermal lysosomal rupture was done as previously described[29,58]. Briefly, age-synchronized animals expressing the sfGFP::LGALS3 construct in the hypodermis were seeded on OP50 or RNAi NGM plates. Scoring was done using a Leica M205 FA widefield binocular microscope. 20-50 worms were analyzed per replicate.

Mounting of live animals for imaging

Synchronized animals were mounted on 8-10% agarose in M9 buffer pads with a drop of mounting mix (2% (w/v) levamisole and 50% (v/v) nanosphere size standards (Thermo Fisher)) and covered with a coverslip.

Fluorescence recovery after photobleaching (FRAP)

FRAP measurements to estimate membrane fluidity were done following a published protocol[59].

Animals were synchronized by bleaching and grown until the second day of adulthood on the

indicated RNAi or empty vector (EV) control plates. Directly before the FRAP experiments,

animals were mounted at the Zeiss LSM 780 confocal microscope. FRAP measurements of either

GFP enriched in the intestinal plasma membrane, or LAAT-1::mCherry located in intestinal or

hypodermal lysosomal membranes were taken with 40X Water immersion objective. For the

lysosomal membrane, mCherry-positive membranes were photobleached over a circular area

(seven pixel radius) using 10 iterations of the 561 nm DPSS laser with 100% laser power

transmission. Images were collected at a 12-bit intensity resolution over 128×128 pixels (digital

zoom 6X) using a pixel dwell time of 3.15 μsec with 2% laser power transmission. For the plasma

membrane, GFP-positive membranes were photobleached over a circular area (10 pixel radius)

using 20 iterations of the 488 nm Argon laser with 100% laser power transmission. Images were

collected at a 12-bit intensity resolution over 128×128 pixels (digital zoom 6X) using a pixel

dwell time of 3.15 μsec with 10% laser power transmission. At least 6 pre-bleaching images were

collected before the region of interest was bleached. The recovery of fluorescence was traced for

25s. Fluorescence recovery and T_{half} were calculated as previously described in Svensk et al.[60].

Touch sensory assay and neurotoxicity in touch receptor neurons

Touch sensory assay was done as previously described[61]. Briefly, touch sensitivity was tested by

gently stroking the tip of an eyebrow hair attached to a toothpick transversely across the anterior

or posterior half of an animal. A touch-sensitive animal was one that stopped/moved away from

the stimulus. 10 worms were examined for each strain in 3 biological replicates with 5 strokes in

the anterior and posterior. For neurotoxicity scoring of the PLM neurons, live animals were imaged

using an Olympus IXplore SpinSR Confocal microscope with an UplanS Apo 60x/1.30 Silicon oil

objective. The neurotoxicity score was determined by adding a score of 1 for each of the following

phenotypes: soma outgrowth, process branching, process break, wavy process. 15 animals were

examined at each day of age in 3 independent biological replicates.

Cell culture

Cells were cultured in DMEM containing high glucose, GlutaMAX Supplement, pyruvate (Gibco), and 10% FBS (Gibco) at 37 °C and 5% CO_2. Regular mycoplasma tests were performed (GATC Biotech). HEK293T cells expressing Venus-tagged full-length P301S mutant 0N4R tau (0N4R tauP301S-Venus) were kindly provided by Dr. William A. McEwan, Cambridge University[38].

Cloning of mammalian expression constructs and generation of sfGFP-LGALS3 HEK293T cells

HEK293T cells were cultured in DMEM containing high glucose, GlutaMAX supplement, pyruvate (Gibco), and 10% FBS (Gibco) at 37 °C and 5% CO_2. Cells expressing sfGFP-LGALS3 were generated as follows. sfGFP::LGALS3 was amplified from the *C. elegans* expression plasmid pPD49.26 by PCR and cloned into the pLenti PGK Puro DEST (w529-2) vector backbone (Addgene plasmid #19068) by MultiSite Gateway cloning according to the manufacturers protocol. pLenti PGK Puro DEST (w529-2) was a gift from Eric Campeau & Paul Kaufman[62]. After each cloning step plasmids isolated from bacterial colonies grown on selective media were extracted with the GenEluteTM Five-Minute Plasmid Miniprep Kit (Qiagen) and validated by sequencing. HEK293T sfGFP-LGALS3 were generated using a 3rd generation lentiviral system. Lentivirus was produced by transiently transfecting HEK293T cells with the pLenti PGK Puro DEST, MDL-gagpol, RSV-REV and IVS-VSVG at a ratio of 5:3:1:1, respectively. Cell media was changed 24 hours after transfection. Viral supernatant was collected 72 h post-transfection. After centrifugation at 2000 x g for 5 min, the supernatant containing the virus was concentrated by adding the PEG-it Virus Precipitation Solution (Systems Biosciences) followed by overnight incubation at 4 °C. The precipitated virus was pelleted by centrifugation (1500 x g, 30 min, 4°C), resuspended in DMEM containing 25 mM HEPES buffer and stored at -80 °C. The lentivirus was added to HEK293T followed by treatment with Puromycin to select successfully transduced HEK293T cells stably expressing sfGFP-LGALS3.

Quantification of lysosomal rupture (galectin puncta assay) in cells

Cells were imaged on an Olympus IXplore SpinSR Confocal microscope with an UplanS Apo 60x/1.30 Silicon oil objective. To detect the number of foci in an image, a difference of gaussian blur was applied to max projections to isolate foci from cytsosolic sfGFP signal. Thresholding was

applied using 'moments' threshold on FIJI. Number of foci was counted using the 'analyze

particles' function. Cell number was counted manually to obtain number of foci/cell for each

image.

C-Laurdan dye measurement of membrane fluidity in worms and cells

Live worms were stained with 10 mM C-Laurdan dye (6-dodecanoyl-2-

dimethylaminonaphthalene) (Thermo Fisher Scientific) as previously described[63]. Cells were also

stained with C-Laurdan at 15 µM for 2 hours then fixed in 4% PFA in PBS. Images were acquired

with an DMI6000 confocal microscope and Leica software with a 63x oil-immersion objective,

with a zoom factor of 4. Samples were excited with a 405 nm laser and the emission recorded

between 400 and 460 nm (ordered phase) and between 470 and 530 nm (disordered phase).

Quantitative assessment of the membrane order was achieved by calculating the ratiometric

relationship of the fluorescence intensity recorded in two spectral channels (GP value) by using an

automated ImageJ macro, according to published guidelines[35].

Knockdown of target genes by RNAi in cells

SPHK2 or a scrambled control siRNA was purchased from Thermo Scientific Dharmacon

(ONTarget plus siRNA and Control Pool). Cells were seeded in DMEM with 10% FBS. siRNA

was diluted in siRNA buffer (Final concentration 20 nM) before mixing with Lipofectmine2000

for 20 min prior to being added to cells. To minimize lipofectamine impact on the endolysosomal

system, media was exchanged after 6 hours.

SDS-Page and immunoblotting.

Cells were pelleted by centrifugation followed by lysis in lysis buffer (10mM Tris, 100mM NaCl,

0.2% Triton X-100, 10mM EDTA) on ice for 20 min. The lysates were transferred into fresh

Eppendorf tubes and centrifuged (1000 g for 1 min at 4 °C) in a tabletop centrifuge to remove

cellular debris. The protein concentration was determined using protein assay dye reagent

concentrate (Bio-Rad). Proteins were separated under denaturing conditions by SDS–PAGE and

transferred onto a PVDF membrane (Carl Roth) by standard wet blotting protocols. Samples were

probed with rabbit polyclonal anti-SPHK2 (1:5000, 9C5E1, Santa Cruz) primary antibody. Anti-

GAPDH antibody (1:5000, clone GAPDH-71.1, Sigma-Aldrich) was used as loading control.

HRP-conjugated anti-mouse and anti-rabbit IgG secondary antibodies (Bio-Rad) were used for
subsequent ECL-based detection (Bio-Rad).

Fatty acid treatment in cells

For α-linolenic acid (ALA), a stock solution was made in EtOH and stored in the dark at -20°C
under a N_2 atmosphere to prevent oxidation[64]. Palmitic acid (PA) was dissolved in 0.1 M NaOH
according to Cousin et al.[65]. ALA and PA were then conjugated to BSA in 10% fatty acid free
solution by shaking for 1 hour at 37°C to a concentration of 6 mM for ALA and 5 mM for PA[66].
Prior to preloaded cells with BSA-PA, the solution was heated to 65 °C for 15 min. For assays
with C-Laurdan, cells were preloaded with FAs for 3 hours before the addition of C-Laurdan for
another 2 hours. If preloading was followed by seeding with tau, preloading with FAs was done
for 3 hours before fibrillar tau was added for an additional 5 hours. C-Laurdan was spiked in for
the last 2 hours. No media changes were done between steps. For experiments with HEK293T tau-
Venus cells and HEK293T sfGFP-LGALS3 cells, preloading with FAs was done for 3 hours before
seeding with 1N4R tau fibrils. Cells were then fixed after 48 hours.

Monomeric and fibrillar tau

Full-length human 1N4R tau was expressed and purified as previously described[67]. The purified
monomeric tau was dialyzed against PBS buffer containing 1 mM DTT, flash frozen in liquid
nitrogen and stored at -80 °C. The concentration of monomeric tau was determined
spectrophotometrically using an extinction coefficient at 280 nm of 7575 M^{-1} cm^{-1}. Monomeric
1N4R tau was assembled into fibrils at 40 μM in the presence of 10 μM heparin, in PBS buffer
containing 1 mM DTT under continuous shaking (600 rpm) for 5 days at 37 °C in an Eppendorf
Thermomixer. The quality of 1N4R tau fibrils produced after 5 days was assessed by ThioflavinT
binding, sedimentation, and transmission electron microscopy (TEM). The resulting 1N4R tau
fibrils were spun at 25 °C and 75 000 g for 30 min. The pelleted fibrils were resuspended in PBS
buffer with 1 mM DTT at 50 μM equivalent monomeric tau concentration. 1N4R tau fibrils were
labeled by incubation with the 2 molar equivalents of lysine-reactive ATTO 550 (ATTO-TEC,
GMBH) for 1 h at room temperature. The unreacted fluorophore was removed by two cycles of
centrifugation at 75 000 g for 10 min and resuspension of the pellet in PBS. 1N4R tau fibrils were
fragmented by sonication for 5 min in 2-ml Eppendorf tubes in a Vial Tweeter powered by an

ultrasonic processor UIS250v (250 W, 2.4 kHz; Hielscher Ultrasonic, Teltow, Germany) to

generate fibrillar particles with an average size 50 nm as assessed by TEM analysis. For seeding

experiments, either monomeric or fibrillar tau was added directly into the cell culture media to a

final concentration of 400 nM. Cells were then fixed after 48 hours.

Statistical analysis

Statistical analysis was performed with GraphPad Prism (GraphPad Software, Version 6h and

10.1.1). Data was tested for normal distribution by D'Agostino and Pearson omnibus normality

test. Data presentation, sample size (number of biological repeats and sample size per repeat and

condition), and the applied statistical tests are indicated in the figure legends for each experiment.

For all cell culture data, data represents 3 independent replicates with 10 images each replicate.

Significance levels: non-significant (n.s.) $p > 0.05$, $*p \leq 0.05$, $**p \leq 0.01$, and $***p \leq 0.001$.

Acknowledgements

We gratefully acknowledge the technical support of Sarah Wübbel. We extend our gratitude to Dr. Sunil Yeruva for his assistance with the use of the DMI6000 confocal microscope. We appreciate the help of Dr. Hana Nuskova and Dr. Aurelio Teleman (DKFZ, Heidelberg) with the fatty acid conjugation protocol. The tauP301S-Venus expressing HEK293T cell line was kindly provided by Dr. William A. McEwan (Cambridge University)[38]. The pLenti PGK Puro DEST (w529-2) vector was a gift from Dr. Eric Campeau and Dr. Paul Kaufman[62]. We also thank Dr. Bin Liu and Dr. Marja Jäättelä (University of Copenhagen) for sharing the strain BIJ34 and a pPD49.26 expression plasmid coding for sfGFP::LGALS3 and Dr. Xiaochen Wang (Chinese Academy of Sciences, Beijing) for nematodes expressing the qxIs352 transgene. Some strains were provided by the Caenorhabditis Genetics Center (CGC), which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440).

Author contributions

Conceptualization J.T., C.A.S., C.N.K.

Formal Analysis J.T., C.A.S., N.M.

Funding acquisition R.M., C.N.K.

Investigation J.T., C.A.S., N.M., D.E.

Methodology J.T., C.A.S., N.M.

Project administration C.N.K.

Resources S.B.N.N.

Supervision R.M., C.N.K.

Visualization J.T., C.N.K.

Writing – original draft J.T., C.A.S., C.N.K.

Writing – review & editing J.T., C.A.S., N.M., D.E., R.M., C.N.K.

Declaration of interests

The authors declare no competing interest.

Supplemental information

Document S1. Figures S1–S5.

557   **Data availability**

558   All data supporting the findings of this study are available within the paper and its supplementary

559   information. *C. elegans* strains and plasmids generated in this study are available upon request.

References

1. Clavaguera, F., Bolmont, T., Crowther, R.A., Abramowski, D., Frank, S., Probst, A., Fraser, G., Stalder, A.K., Beibel, M., Staufenbiel, M., et al. (2009). Transmission and spreading of tauopathy in transgenic mouse brain. Nat Cell Biol *11*, 909-913. 10.1038/ncb1901.

2. Walker, L.C., and Jucker, M. (2024). The prion principle and Alzheimer's disease. Science *385*, 1278-1279. 10.1126/science.adq5252.

3. Biel, D., Brendel, M., Rubinski, A., Buerger, K., Janowitz, D., Dichgans, M., Franzmeier, N., and Alzheimer's Disease Neuroimaging, I. (2021). tau-PET and in vivo Braak-staging as prognostic markers of future cognitive decline in cognitively normal to demented individuals. Alzheimer's research & therapy *13*, 137. 10.1186/s13195-021-00880-x.

4. Frost, B., Jacks, R.L., and Diamond, M.I. (2009). Propagation of tau misfolding from the outside to the inside of a cell. J Biol Chem *284*, 12845-12852. M808759200 10.1074/jbc.M808759200.

5. Sanders, D.W., Kaufman, S.K., DeVos, S.L., Sharma, A.M., Mirbaha, H., Li, A., Barker, S.J., Foley, A.C., Thorpe, J.R., Serpell, L.C., et al. (2014). Distinct tau prion strains propagate in cells and mice and define different tauopathies. Neuron *82*, 1271-1288. 10.1016/j.neuron.2014.04.047.

6. Shrivastava, A.N., Redeker, V., Pieri, L., Bousset, L., Renner, M., Madiona, K., Mailhes-Hamon, C., Coens, A., Buee, L., Hantraye, P., et al. (2019). Clustering of tau fibrils impairs the synaptic composition of alpha3-Na(+)/K(+)-ATPase and AMPA receptors. EMBO J *38*. 10.15252/embj.201899871.

7. Caballero, B., Wang, Y., Diaz, A., Tasset, I., Juste, Y.R., Stiller, B., Mandelkow, E.M., Mandelkow, E., and Cuervo, A.M. (2018). Interplay of pathogenic forms of human tau with different autophagic pathways. Aging cell *17*. 10.1111/acel.12692.

8. Menzies, F.M., Fleming, A., and Rubinsztein, D.C. (2015). Compromised autophagy and neurodegenerative diseases. Nat Rev Neurosci *16*, 345-357. 10.1038/nrn3961.

9. Nixon, R.A., Wegiel, J., Kumar, A., Yu, W.H., Peterhoff, C., Cataldo, A., and Cuervo, A.M. (2005). Extensive involvement of autophagy in Alzheimer disease: an immuno-electron microscopy study. J Neuropathol Exp Neurol *64*, 113-122.

10. Hou, W.C., Massey, L.A., Rhoades, D., Wu, Y., Ren, W., Frank, C., Overkleeft, H.S., and Kelly, J.W. (2024). A PIKfyve modulator combined with an integrated stress response inhibitor to treat lysosomal storage diseases. Proc Natl Acad Sci U S A *121*, e2320257121. 10.1073/pnas.2320257121.

11. Lim, S.H.Y., Hansen, M., and Kumsta, C. (2024). Molecular Mechanisms of Autophagy Decline during Aging. Cells *13*. 10.3390/cells13161364.

12. Caballero, B., Bourdenx, M., Luengo, E., Diaz, A., Sohn, P.D., Chen, X., Wang, C., Juste, Y.R., Wegmann, S., Patel, B., et al. (2021). Acetylated tau inhibits chaperone-mediated autophagy and promotes tau pathology propagation in mice. Nature communications *12*, 2238. 10.1038/s41467-021-22501-9.

13. Chen, J.J., Nathaniel, D.L., Raghavan, P., Nelson, M., Tian, R., Tse, E., Hong, J.Y., See, S.K., Mok, S.A., Hein, M.Y., et al. (2019). Compromised function of the ESCRT pathway promotes endolysosomal escape of tau seeds and propagation of tau aggregation. J Biol Chem. 10.1074/jbc.RA119.009432.

14. Flavin, W.P., Bousset, L., Green, Z.C., Chu, Y., Skarpathiotis, S., Chaney, M.J., Kordower, J.H., Melki, R., and Campbell, E.M. (2017). Endocytic vesicle rupture is a conserved

mechanism of cellular invasion by amyloid proteins. Acta Neuropathol. 10.1007/s00401-
017-1722-x.

15. Tuck, B.J., Miller, L.V.C., Katsinelos, T., Smith, A.E., Wilson, E.L., Keeling, S., Cheng,
S., Vaysburd, M.J., Knox, C., Tredgett, L., et al. (2022). Cholesterol determines the
cytosolic entry and seeded aggregation of tau. Cell reports *39*, 110776.
10.1016/j.celrep.2022.110776.

16. Dimou, E., Katsinelos, T., Meisl, G., Tuck, B.J., Keeling, S., Smith, A.E., Hidari, E., Lam,
J.Y.L., Burke, M., Lovestam, S., et al. (2023). Super-resolution imaging unveils the self-
replication of tau aggregates upon seeding. Cell reports *42*, 112725.
10.1016/j.celrep.2023.112725.

17. Polanco, J.C., Hand, G.R., Briner, A., Li, C., and Götz, J. (2021). Exosomes induce
endolysosomal permeabilization as a gateway by which exosomal tau seeds escape into the
cytosol. Acta Neuropathol *141*, 235-256. 10.1007/s00401-020-02254-3.

18. Rose, K., Jepson, T., Shukla, S., Maya-Romero, A., Kampmann, M., Xu, K., and Hurley,
J.H. (2024). tau fibrils induce nanoscale membrane damage and nucleate cytosolic tau at
lysosomes. Proc Natl Acad Sci U S A *121*, e2315690121. 10.1073/pnas.2315690121.

19. Kroemer, G., and Jaattela, M. (2005). Lysosomes and autophagy in cell death control.
Nature reviews. Cancer *5*, 886-897. 10.1038/nrc1738.

20. Sandhof, C.A., Martin, N., Tittelmeier, J., Schlueter, A., Pezzali, M., Schoendorf, D.C.,
Lange, T., Reinhardt, P., Ried, J.S., Liang, S., et al. (2024). A Novel *C. elegans* Model for
tau Spreading Reveals Genes Critical for Endolysosomal Integrity and Seeded tau
Aggregation. bioRxiv, 2024.2011.2013.619586. 10.1101/2024.11.13.619586.

21. Breslow, D.K., and Weissman, J.S. (2010). Membranes in balance: mechanisms of
sphingolipid homeostasis. Mol Cell *40*, 267-279. 10.1016/j.molcel.2010.10.005.

22. Jimenez-Rojo, N., Leonetti, M.D., Zoni, V., Colom, A., Feng, S., Iyengar, N.R., Matile, S.,
Roux, A., Vanni, S., Weissman, J.S., and Riezman, H. (2020). Conserved Functions of
Ether Lipids and Sphingolipids in the Early Secretory Pathway. Curr Biol *30*, 3775-3787
e3777. 10.1016/j.cub.2020.07.059.

23. Gault, C.R., Obeid, L.M., and Hannun, Y.A. (2010). An overview of sphingolipid
metabolism: from synthesis to breakdown. Adv Exp Med Biol *688*, 1-23. 10.1007/978-1-
4419-6741-1_1.

24. Futerman, A.H., and Hannun, Y.A. (2004). The complex life of simple sphingolipids.
EMBO Rep *5*, 777-782. 10.1038/sj.embor.7400208.

25. Czubowicz, K., Jesko, H., Wencel, P., Lukiw, W.J., and Strosznajder, R.P. (2019). The
Role of Ceramide and Sphingosine-1-Phosphate in Alzheimer's Disease and Other
Neurodegenerative Disorders. Mol Neurobiol *56*, 5436-5455. 10.1007/s12035-018-1448-
3.

26. van Echten-Deckert, G., and Walter, J. (2012). Sphingolipids: critical players in
Alzheimer's disease. Prog Lipid Res *51*, 378-393. 10.1016/j.plipres.2012.07.001.

27. Huotari, J., and Helenius, A. (2011). Endosome maturation. EMBO J *30*, 3481-3500.
10.1038/emboj.2011.286.

28. Yong, J., Villalta, J.E., Vu, N., Kukurugya, M.A., Olsson, N., Lopez, M.P., Lazzari-Dean,
J.R., Hake, K., McAllister, F.E., Bennett, B.D., and Jan, C.H. (2024). Impairment of lipid
homeostasis causes lysosomal accumulation of endogenous protein aggregates through
ESCRT disruption. eLife *12*. 10.7554/eLife.86194.

29. Aits, S., Kricker, J., Liu, B., Ellegaard, A.M., Hamalisto, S., Tvingsholm, S., Corcelle-Termeau, E., Hogh, S., Farkas, T., Holm Jonassen, A., et al. (2015). Sensitive detection of lysosomal membrane permeabilization by lysosomal galectin puncta assay. Autophagy *11*, 1408-1424. 10.1080/15548627.2015.1063871.

30. Hanel, V., Pendleton, C., and Witting, M. (2019). The sphingolipidome of the model organism Caenorhabditis elegans. Chem Phys Lipids *222*, 15-22. 10.1016/j.chemphyslip.2019.04.009.

31. Cheng, X., Jiang, X., Tam, K.Y., Li, G., Zheng, J., and Zhang, H. (2019). Sphingolipidomic Analysis of C. elegans reveals Development- and Environment-dependent Metabolic Features. Int J Biol Sci *15*, 2897-2910. 10.7150/ijbs.30499.

32. D'Auria, L., and Bongarzone, E.R. (2016). Fluid levity of the cell: Role of membrane lipid architecture in genetic sphingolipidoses. J Neurosci Res *94*, 1019-1024. 10.1002/jnr.23750.

33. Liu, B., Du, H., Rutkowski, R., Gartner, A., and Wang, X. (2012). LAAT-1 is the lysosomal lysine/arginine transporter that maintains amino acid homeostasis. Science *337*, 351-354. 10.1126/science.1220281.

34. Morck, C., Olsen, L., Kurth, C., Persson, A., Storm, N.J., Svensson, E., Jansson, J.O., Hellqvist, M., Enejder, A., Faergeman, N.J., and Pilon, M. (2009). Statins inhibit protein lipidation and induce the unfolded protein response in the non-sterol producing nematode Caenorhabditis elegans. Proc Natl Acad Sci U S A *106*, 18285-18290. 10.1073/pnas.0907117106.

35. Owen, D.M., Rentero, C., Magenau, A., Abu-Siniyeh, A., and Gaus, K. (2011). Quantitative imaging of membrane lipid order in cells and organisms. Nat Protoc *7*, 24-35. 10.1038/nprot.2011.419.

36. Barucha-Kraszewska, J., Kraszewski, S., and Ramseyer, C. (2013). Will C-Laurdan dethrone Laurdan in fluorescent solvent relaxation techniques for lipid membrane studies? Langmuir *29*, 1174-1182. 10.1021/la304235r.

37. Wittrup, A., Ai, A., Liu, X., Hamar, P., Trifonova, R., Charisse, K., Manoharan, M., Kirchhausen, T., and Lieberman, J. (2015). Visualizing lipid-formulated siRNA release from endosomes and target gene knockdown. Nat Biotechnol *33*, 870-876. 10.1038/nbt.3298.

38. McEwan, W.A., Falcon, B., Vaysburd, M., Clift, D., Oblak, A.L., Ghetti, B., Goedert, M., and James, L.C. (2017). Cytosolic Fc receptor TRIM21 inhibits seeded tau aggregation. Proc Natl Acad Sci U S A *114*, 574-579. 10.1073/pnas.1607215114.

39. Nachman, E., Wentink, A.S., Madiona, K., Bousset, L., Katsinelos, T., Allinson, K., Kampinga, H., McEwan, W.A., Jahn, T.R., Melki, R., et al. (2020). Disassembly of tau fibrils by the human Hsp70 disaggregation machinery generates small seeding-competent species. J Biol Chem *295*, 9676-9690. 10.1074/jbc.RA120.013478.

40. Yoon, J.H., Seo, Y., Jo, Y.S., Lee, S., Cho, E., Cazenave-Gassiot, A., Shin, Y.S., Moon, M.H., An, H.J., Wenk, M.R., and Suh, P.G. (2022). Brain lipidomics: From functional landscape to clinical significance. Sci Adv *8*, eadc9317. 10.1126/sciadv.adc9317.

41. Hornemann, T. (2021). Mini review: Lipids in Peripheral Nerve Disorders. Neurosci Lett *740*, 135455. 10.1016/j.neulet.2020.135455.

42. Uranbileg, B., Isago, H., Sakai, E., Kubota, M., Saito, Y., and Kurano, M. (2024). Alzheimer's disease manifests abnormal sphingolipid metabolism. Front Aging Neurosci *16*, 1368839. 10.3389/fnagi.2024.1368839.

43. Corder, E.H., Saunders, A.M., Strittmatter, W.J., Schmechel, D.E., Gaskell, P.C., Small,
G.W., Roses, A.D., Haines, J.L., and Pericak-Vance, M.A. (1993). Gene dose of
apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families.
Science *261*, 921-923. 10.1126/science.8346443.
44. Kosicek, M., Zetterberg, H., Andreasen, N., Peter-Katalinic, J., and Hecimovic, S. (2012).
Elevated cerebrospinal fluid sphingomyelin levels in prodromal Alzheimer's disease.
Neurosci Lett *516*, 302-305. 10.1016/j.neulet.2012.04.019.
45. Ruiz, M., Devkota, R., Kaper, D., Ruhanen, H., Busayavalasa, K., Radovic, U., Henricsson,
M., Kakela, R., Boren, J., and Pilon, M. (2023). AdipoR2 recruits protein interactors to
promote fatty acid elongation and membrane fluidity. J Biol Chem *299*, 104799.
10.1016/j.jbc.2023.104799.
46. Zhang, Y., Wang, H., Zhang, J., Hu, Y., Zhang, L., Wu, X., Su, X., Li, T., Zou, X., and
Liang, B. (2016). The cytochrome b5 reductase HPO-19 is required for biosynthesis of
polyunsaturated fatty acids in Caenorhabditis elegans. Biochim Biophys Acta *1861*, 310-
319. 10.1016/j.bbalip.2016.01.009.
47. Fabri, J., de Sa, N.P., Malavazi, I., and Del Poeta, M. (2020). The dynamics and role of
sphingolipids in eukaryotic organisms upon thermal adaptation. Prog Lipid Res *80*,
101063. 10.1016/j.plipres.2020.101063.
48. Romanauska, A., and Kohler, A. (2023). Lipid saturation controls nuclear envelope
function. Nat Cell Biol *25*, 1290-1302. 10.1038/s41556-023-01207-8.
49. Harayama, T., and Riezman, H. (2018). Understanding the diversity of membrane lipid
composition. Nat Rev Mol Cell Biol *19*, 281-296. 10.1038/nrm.2017.138.
50. Rakhra, V., Galappaththy, S.L., Bulchandani, S., and Cabandugama, P.K. (2020). Obesity
and the Western Diet: How We Got Here. Mo Med *117*, 536-538.
51. Snowden, S.G., Ebshiana, A.A., Hye, A., An, Y., Pletnikova, O., O'Brien, R., Troncoso, J.,
Legido-Quigley, C., and Thambisetty, M. (2017). Association between fatty acid
metabolism in the brain and Alzheimer disease neuropathology and cognitive performance:
A nontargeted metabolomic study. PLoS Med *14*, e1002266.
10.1371/journal.pmed.1002266.
52. Bischoff-Ferrari, H.A., Gangler, S., Wieczorek, M., Belsky, D.W., Ryan, J., Kressig, R.W.,
Stahelin, H.B., Theiler, R., Dawson-Hughes, B., Rizzoli, R., et al. (2025). Individual and
additive effects of vitamin D, omega-3 and exercise on DNA methylation clocks of
biological aging in older adults from the DO-HEALTH trial. Nat Aging. 10.1038/s43587-
024-00793-y.
53. Shinto, L., Quinn, J., Montine, T., Dodge, H.H., Woodward, W., Baldauf-Wagner, S.,
Waichunas, D., Bumgarner, L., Bourdette, D., Silbert, L., and Kaye, J. (2014). A
randomized placebo-controlled pilot trial of omega-3 fatty acids and alpha lipoic acid in
Alzheimer's disease. J Alzheimers Dis *38*, 111-120. 10.3233/JAD-130722.
54. Wood, A.H.R., Chappell, H.F., and Zulyniak, M.A. (2022). Dietary and supplemental long-
chain omega-3 fatty acids as moderators of cognitive impairment and Alzheimer's disease.
Eur J Nutr *61*, 589-604. 10.1007/s00394-021-02655-4.
55. Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71-94.
56. Svensk, E., Stahlman, M., Andersson, C.H., Johansson, M., Boren, J., and Pilon, M. (2013).
PAQR-2 regulates fatty acid desaturation during cold adaptation in C. elegans. PLoS Genet
9, e1003801. 10.1371/journal.pgen.1003801.

57.   Devkota, R., Svensk, E., Ruiz, M., Stahlman, M., Boren, J., and Pilon, M. (2017). The
adiponectin receptor AdipoR2 and its Caenorhabditis elegans homolog PAQR-2 prevent
membrane rigidification by exogenous saturated fatty acids. PLoS Genet *13*, e1007004.
10.1371/journal.pgen.1007004.
58.   Sandhof, C.A., Hoppe, S.O., Druffel-Augustin, S., Gallrein, C., Kirstein, J., Voisine, C.,
and Nussbaum-Krammer, C. (2020). Reducing INS-IGF1 signaling protects against non-
cell autonomous vesicle rupture caused by SNCA spreading. Autophagy *16*, 878-899.
10.1080/15548627.2019.1643657.
59.   Devkota, R., and Pilon, M. (2018). FRAP: A Powerful Method to Evaluate Membrane
Fluidity in Caenorhabditis elegans. Bio Protoc *8*, e2913. 10.21769/BioProtoc.2913.
60.   Svensk, E., Devkota, R., Stahlman, M., Ranji, P., Rauthan, M., Magnusson, F.,
Hammarsten, S., Johansson, M., Boren, J., and Pilon, M. (2016). Caenorhabditis elegans
PAQR-2 and IGLR-2 Protect against Glucose Toxicity by Modulating Membrane Lipid
Composition. PLoS Genet *12*, e1005982. 10.1371/journal.pgen.1005982.
61.   Chalfie, M., Sulston, J.E., White, J.G., Southgate, E., Thomson, J.N., and Brenner, S.
(1985). The neural circuit for touch sensitivity in Caenorhabditis elegans. J Neurosci *5*,
956-964.
62.   Campeau, E., Ruhl, V.E., Rodier, F., Smith, C.L., Rahmberg, B.L., Fuss, J.O., Campisi, J.,
Yaswen, P., Cooper, P.K., and Kaufman, P.D. (2009). A versatile viral system for
expression and depletion of proteins in mammalian cells. PLoS One *4*, e6529.
10.1371/journal.pone.0006529.
63.   Jeong, J.H., Han, J.S., Jung, Y., Lee, S.M., Park, S.H., Park, M., Shin, M.G., Kim, N.,
Kang, M.S., Kim, S., et al. (2023). A new AMPK isoform mediates glucose-restriction
induced longevity non-cell autonomously by promoting membrane fluidity. Nature
communications *14*, 288. 10.1038/s41467-023-35952-z.
64.   Oliveira, A.F., Cunha, D.A., Ladriere, L., Igoillo-Esteve, M., Bugliani, M., Marchetti, P.,
and Cnop, M. (2015). In vitro use of free fatty acids bound to albumin: A comparison of
protocols. Biotechniques *58*, 228-233. 10.2144/000114285.
65.   Cousin, S.P., Hugl, S.R., Wrede, C.E., Kajio, H., Myers, M.G., Jr., and Rhodes, C.J. (2001).
Free fatty acid-induced inhibition of glucose and insulin-like growth factor I-induced
deoxyribonucleic acid synthesis in the pancreatic beta-cell line INS-1. Endocrinology *142*,
229-240. 10.1210/endo.142.1.7863.
66.   Nuskova, H., Cortizo, F.G., Schwenker, L.S., Sachsenheimer, T., Diakonov, E.E., Tiebe,
M., Schneider, M., Lohbeck, J., Reid, C., Kopp-Schneider, A., et al. (2023). Competition
for cysteine acylation by C16:0 and C18:0 derived lipids is a global phenomenon in the
proteome. J Biol Chem *299*, 105088. 10.1016/j.jbc.2023.105088.
67.   Tardivel, M., Begard, S., Bousset, L., Dujardin, S., Coens, A., Melki, R., Buee, L., and
Colin, M. (2016). Tunneling nanotube (TNT)-mediated neuron-to neuron transfer of
pathological tau protein assemblies. Acta neuropathologica communications *4*, 117.
10.1186/s40478-016-0386-4.
68.   Chitwood, D.J., Lusby, W.R., Thompson, M.J., Kochansky, J.P., and Howarth, O.W.
(1995). The glycosylceramides of the nematode Caenorhabditis elegans contain an
unusual, branched-chain sphingoid base. Lipids *30*, 567-573. 10.1007/BF02537032.
69.   Zhang, H., Abraham, N., Khan, L.A., Hall, D.H., Fleming, J.T., and Gobel, V. (2011).
Apicobasal domain identities of expanding tubular membranes depend on
glycosphingolipid biosynthesis. Nat Cell Biol *13*, 1189-1201. 10.1038/ncb2328.

70.   Zhu, H., Shen, H., Sewell, A.K., Kniazeva, M., and Han, M. (2013). A novel sphingolipid-
TORC1 pathway critically promotes postembryonic development in Caenorhabditis
elegans. eLife *2*, e00429. 10.7554/eLife.00429.
71.   Scholz, J., Helmer, P.O., Nicolai, M.M., Bornhorst, J., and Hayen, H. (2021). Profiling of
sphingolipids in Caenorhabditis elegans by two-dimensional multiple heart-cut liquid
chromatography - mass spectrometry. J Chromatogr A *1655*, 462481.
10.1016/j.chroma.2021.462481.

 Figure Legends

Figure 1. Sphingolipid metabolism dysregulation by KD is linked to of endolysosomal vesicle

rupture.

(A) General overview of sphingolipid metabolism with a particular focus on the genes identified

in the screen we performed. Sphingolipids constitute a group of amphipathic lipids featuring a

polar head group and a sphingoid base backbone that is N-acetylated with a (very) long-chain fatty

acid ((V)LCFA) side chain. In contrast to mammals, where the sphingoid base is conventionally

derived from palmitic acid and serine, *C. elegans* sphingolipids usually contain a characteristic

C17iso branched chain sphingoid base [68-70]. Its synthesis involves the branched chain FA (BCFA)

elongation pathway to yield C15iso-CoA, which then condensates with L-serine to form 3-

ketosphinganine. This reaction is catalyzed by serine palmitoyltransferase (encoded by *sptl-1*, *-2*,

and *-3*). The serine incorporator (SERINC) protein family (encoded by *R11H6.2*) is believed to

assist in the incorporation of L-serine into specific membranes. 3-ketodihydrosphingosine

reductase (KDSR, in *C. elegans* predicted to be encoded by Y37E11AM.3) then catalyzes the

reduction of 3-keto sphinganine to sphinganine. The (V)LCFA side chain is primarily comprised

of a straight saturated FA chain, ranching from 20-26 carbon atoms in length, with or without

hydroxylation [30,71]. It can be also derived from BCFAs, such as C15iso and C17iso [68,69]. However,

most of the side chain FA moieties originate from palmitoyl-CoA via the canonical de novo FA

biosynthesis pathway, involving the sequential addition of C2 moieties from malonyl-CoA through

the LCFA elongation cycle [68,69]. Each elongation cycle comprises four reactions (condensation,

reduction, dehydration, reduction), with the third reaction requiring very-long-chain (3R)-3-

hydroxyacyl-CoA dehydratase (encoded by *hpo-8*) [69]. Finally, ceramide synthases (encoded by

hyl-1 and *hyl-2*) catalyze the addition of various acyl side chains to the sphingoid base to yield

dihydroceramide, which is then desaturated to ceramide. The latter reaction is catalyzed by

dihydroceramide desaturases, which require electrons from NAD(P)H provided by cytochrome b5

reductases (encoded by *hpo-19* and *T05H4.4*). All complex sphingolipids, such as sphingomyelin

and glycosphingolipids (including cerebrosides and gangliosides) originate from ceramide.

Degradation of complex sphingolipids takes place in the lysosome. Essential for this process are

saposins or sphingolipid activator proteins (PSAPs, encoded by *spp-10*), which serve as crucial

bridges between the lipid substrate and hydrophilic hydrolases. Glucocerebrosidases (encoded by

gba-1, gba-2, gba-3, and *gba-4*) hydrolyse glucosylceramide into ceramide and glucose.

Sphingosine may be either recycled and metabolized back into ceramide or phosphorylated by
sphingosine kinase (encoded by *sphk-1*) to generate sphingosine-1-phosphate (S1P). S1P lyase
(encoded by *spl-1*) irreversibly cleaves S1P into phospho-ethanolamine and hexadecenal. The *C.*
elegans genes identified in the primary screen, along with their human orthologs, are framed with
color. Genes identified in subsequent co-RNAi experiments and their human orthologs are framed
in gray.
(B) Schematic of lysosomal rupture detected by the galectin puncta assay.
(C) Widefield fluorescence images of animals expressing hypodermal sfGFP::LGALS3.
Numerous foci are visible upon RNAi-mediated KD of sphingolipid metabolisms genes. Zoomed
in image is indicated in overview image by a white box.
(D) Mean percentage of animals positive for lysosomal rupture (defined as three or more
sfGFP::LGALS3 foci in the hypodermis). Data shown as means of three technical replicate plates
with 17-30 animals per plate ± SEM. Statistical analysis comparing RNAi conditions to the empty
vector control (EV ctrl) was done using one-way ANOVA with Dunnett's post-hoc test. *** = p
< 0.001.
844
Figure 2. Knockdown of genes involved in sphingolipid metabolism decreases lysosomal
membrane fluidity.
(A) Representative confocal single plane images from a FRAP experiment in animals expressing
a mCherry tagged Lysosomal Lysine/Arginine Transporter 1 (LAAT-1::mCherry) in the
hypodermis grown either on empty vector control (EV ctrl), *spl-1* or *sphk-1* RNAi plates. Dashed
circles outline the bleach spots.
(B) Combined FRAP curves of LAAT-1::mCherry in hypodermal lysosomal membranes. Curves
are normalized to the pre-bleach intensity as 100% and the first post-bleach intensity as 0%.
(C, D) Increase (C) in the mean time until half of the maximal signal is recovered (t_{half},) and
decrease (D) in the maximal % recoverable fluorescence values upon KD of *sphk-1* and *spl-1*
indicate a reduction in lysosomal membrane fluidity.
(E) FRAP curves of LAAT-1::mCherry in hypodermal lysosomal membranes of animals grown
either on empty vector or the indicated RNAi plates. Curves are normalized to the pre-bleach
intensity set as 100% and the first post-bleach intensity as 0%.
(F) Mean t_{half} upon KD of sphingolipid metabolism genes.

(G) Maximal % recoverable fluorescence values upon KD of sphingolipid metabolism genes.

(H) Representative confocal single plane images from a FRAP experiment in animals expressing

prenylated GFP for lipid membrane anchorage in the intestine.

(I) Combined FRAP curves of prenylated GFP enriched on the intestinal plasma membrane of

animals grown either on empty vector, *sphk-1* or *spl-1* RNAi plates. Curves are normalized to the

pre-bleach intensity as 100% and the first post-bleach intensity as 0%.

(J, K) Mean t_{half} (J) and maximal % recoverable fluorescence values (K). Data represented as

means ± SEM of 5 – 12 FRAP measurements per condition collected in five biological replicates.

Statistical analysis comparing RNAi conditions to the empty vector control was done using one-

way ANOVA with Dunnett's post-hoc test. n.s.: not significant, * = $p < 0.05$, ** = $p < 0.01$, ***

= $p < 0.001$

Figure 3. KD of SPHK2 and aggregated tau reduces membrane fluidity leading to lysosomal

rupture.

(A) Scheme of the fluorescence properties of C-Laurdan. The dye is excited using illumination at

405 nm. The dye fluoresces with a peak emission wavelength around 450 nm (red) when residing

in the ordered phase and ~500 nm in the disordered phase (blue). Two-channel acquisition is

conducted in the wavelength bands indicated by shaded boxes.

(B) Pseudo-color images showing the GP index of C-Laurdan dye at each pixel position in

HEK293T cells upon KD of SPHK2.

(C) Increase in GP Index of HEK293T cells upon KD of SPHK2. Statistical analysis was done

using one-way ANOVA with Bonferroni's post-hoc test.

(D) Max. intensity projection of confocal z-stacks of HEK293T cells expressing sfGFP-LGALS3

upon KD of SPHK2 and seeding with 1N4R tau fibrils.

(E) Average number of foci per number of cells. n= 3 independent experiments, 10 images each.

Statistical analysis done using oKruskal-Wallis with a Dunn's post-hoc test.

(F) Quantification of the percentage of animals with ≥ 3 hypodermal sfGFP::LGALS3 foci,

representing endolysosomal damage, expressing either the F3$^{\Delta K281}$::mCherry (red) or only the

mCherry tag (grey) in touch receptor neurons under knockdown of the indicated genes.

(G) Dilution of the indicated RNAi cultures with the empty vector control culture was done to

reduce the strength of the knockdown. Data represented as mean ± SEM. Besides knockdown of

sphk-1 in **C** n = 3 - 7 with **F** 40 – 50 or **G** 20 – 30 animals per replicate. Since even a dilution down

to 5% of the *sphk-1* RNAi resulted in 100% of animals being scored as positive in two independent

replicates this condition was not repeated further. Statistical analysis comparing RNAi conditions

to the empty vector control done using one-way ANOVA with Dunnett's post-hoc test.

(H) Worms expressing F3$^{\Delta K281}$::mCherry or mCherry in touch receptor neurons stained with C-

Laurdan. The mCherry signal was used to select a region of interest (ROI) around the soma of the

posterior touch receptor neurons (PLM) to determine the GP Index.

(I) Increase in rigidity of membranes in animals expressing F3$^{\Delta K281}$::mCherry compared to

mCherry. Statistical analysis was done using a t-test. n=4, total of at least 28 worms.

(J) Pseudocolored HEK293T cells exposed to tau 1N4R fibrils visualizing the C-Laurdan dye GP

Index.

(K) Increase rigidity of membranes of cells seeded with tau fibrils as indicated by the GP index.

(L) GP index of cells seeded with monomeric tau. Statistical analysis for C-Laurdan in cells was

done using a t-test. n= 3 independent experiments, 10 images each. Scale bar 10 μm. n.s.: not

significant, * = p < 0.05, ** = p < 0.01, *** = p < 0.001.

Figure 4. Saturated fatty acids decrease membrane fluidity and exacerbate seeded tau

aggregation.

(A) Left: saturated fatty acid membrane scheme. Right: pseudocolored HEK293T cells pre-loaded

with 50 μM PA conjugated to BSA (BSA-PA) displaying the C-Laurdan dye GP Index.

(B) GP Index indicates increased rigidity of membranes with BSA-PA. Statistical analysis was

done using the Mann-Whitney-test. n= 3 independent experiments, 10 images each.

(C) Pseudocolored HEK293T cells pre-loaded with BSA-PA and seeded with 1N4R tau fibrils.

(D) GP Index indicates an additive impact of tau fibrils on membrane rigidity when added to

HEK293T cells pre-loaded with BSA-PA. Statistical analysis was done using t-test. n= 3

independent experiments, 10 images each.

(E) Quantification of hypodermal sfGFP-LGALS3 foci in animals grown on plates supplemented

with PA reveal an increase in lysosomal rupture. Statistical analysis was done using Mann-

Whitney test. n= 3 independent experiments, 15 worms per replicate.

(F) Max. intensity projection of confocal z-stacks of HEK293T cells expressing sfGFP-LGALS3

pre-loaded with BSA-PA and seeded with 1N4R tau fibrils.

(G) Quantification of sfGFP-LGALS3 foci in HEK293T cells following indicated treatments.

(H) Max. intensity projection of confocal z-stacks of Venus-tagged full-length P301S tau

biosensor cells pre-loaded with BSA-PA and seeded with 1N4R tau fibrils.

(I) Quantification of Venus-P301S tau foci following indicated treatments. Statistical analysis

comparing BSA + tau to other conditions was done using one-way ANOVA with Dunnett's post-

hoc test. n= 3 independent experiments, 10 images each. * = $p < 0.05$, ** = $p < 0.01$, *** = $p <$

0.001. Scale bar 10 µm.

Figure 5. PUFA supplementation restores lysosomal membrane integrity and reduces seeded

tau aggregation and toxicity.

(A) Left: unsaturated fatty acid membrane scheme. Right: C-Laurdan GP Index of HEK293T cells

pre-loaded with 150 µM ALA conjugated to BSA (BSA-ALA) indicating an increase in membrane

fluidity. Statistical analysis was done with a t-test. n= 3 independent experiments, 10 images each.

(B) Pseudocolored HEK293T cells pre-loaded with BSA-ALA and seeded with 1N4R tau fibrils.

(C) GP Index indicates that BSA-ALA reduces tau-induced rigidification of the membrane.

Statistical analysis was done using the Mann-Whitney U-test. n= 3 independent experiments, 10

images each.

(D) Max. intensity projection of confocal z-stacks of HEK293T cells expressing sfGFP-LGALS3

pre-loaded with BSA-ALA and seeded with 1N4R tau fibrils. Scale bar 10 µm.

(E) Quantification of sfGFP-LGALS3 foci following indicated treatments. n= 3 independent

experiments, 10 images each.

(F) Max. intensity projection of confocal z-stacks of tau-Venus biosensor cell line pre-loaded with

BSA-ALA and seeded with 1N4R tau fibrils. Scale bar 10 µm.

(G) Quantification of tau-Venus foci following indicated treatments. Statistical analysis

comparing BSA + tau to other conditions was done using Kruskal-Wallis with Dunn's post-hoc

test. n = 3 independent experiments, 10 images each.

(H) Posterior response to touch during aging of animals expressing F3$^{\Delta K281}$::mCherry grown on

plates supplemented with ALA or only the ethanol solvent. Statistical analysis was done using

two-way ANOVA with Bonferroni's multiple comparison test.

(I) Max. intensity projection of confocal z-stacks of day 6 old animals expressing

F3$^{\Delta K281}$::mCherry.

(J) Neurotoxicity score of PLM neurons of animals grown on control or ALA plates. Statistical

analysis was done using 2-way ANOVA with a Holm-Sidak's multiple comparison test. n= 3

independent experiments, 15 worms per replicate. Scale bar 20 μm. * = $p < 0.05$, ** = $p < 0.01$,

*** = $p < 0.001$.

a Biosynthesis

Degradation

a

C-Laurdan Ordered Disordered

Peak emission wavelength:
Ordered: ~450 nm
Disordered: ~500 nm

b

Control   Lipofectamine

Control siRNA   SPHK2 siRNA

c

d

sfGFP-LGALS3

Control siRNA   SPHK2 siRNA   Control siRNA + Tau fibrils   SPHK2 siRNA + Tau fibrils

e

f

EV ctrl, RH116H.2, Y37E11AM.3, hpo-8, hpo-19, spp-10, sphk-1, spl-1

g

25% RH116H.2, 10% hpo-8, 25% hpo-19, 5% sphk-1, 25% spl-1

F3^(ΔK281)::mCherry
mCherry

h

mec-4p::mCherry   PLM   ROI selected for soma of PLM   mCherry   F3^(ΔK281)::mCherry

PLM R/L
AVM R/L
AVM   PVM

i

j

PBS   Tau

k

l

a

Unsaturated fatty acids

GP Index
***
BSA   BSA-ALA

b
GP Index
0.4

-0.6
BSA + Tau fibrils   BSA-ALA + Tau fibrils

c
GP Index
***
BSA +Tau   BSA-ALA +Tau

d
sfGFP-LGALS3

BSA   BSA-ALA   BSA + Tau fibrils   BSA-ALA + Tau fibrils

e
Foci/cell
***   ***
BSA   BSA-ALA   BSA + Tau   BSA-ALA + Tau

f
Tau-Venus

BSA   BSA-ALA   BSA + Tau fibrils   BSA-ALA + Tau fibrils

g
Foci/cell
***   ***
BSA   BSA-ALA   BSA + Tau   BSA-ALA + Tau

h
Posterior touch response
n.s.   *   **
Control ALA   Control ALA   Control ALA
Day 4   Day 6   Day 8

i
F3^{ΔK281}::mCherry
Control

ALA

j
Neurotoxicity score
n.s.   **   *
Control ALA   Control ALA   Control ALA
Day 4   Day 6   Day 8