Title

Single-nucleus transcriptional and chromatin accessibility analyses of maturing mouse

Achilles tendon uncover the molecular landscape of tendon stem/progenitor cells

Authors

Hiroki Tsutsumi[1#], Tomoki Chiba[1#], Yuta Fujii[1,2], Takahide Matsushima[1], Tsuyoshi Kimura[3,4],

Akinori Kanai[5], Akio Kishida[3], Yutaka Suzuki[5] and Hiroshi Asahara[1,2*]

Affiliations

1 Department of Systems BioMedicine, Graduate School of Medical and Dental Sciences,

Institute of Science Tokyo, Tokyo, Japan

2 Department of Molecular and Cellular Biology, The Scripps Research Institute, La Jolla,

USA

3 Laboratory for Biomaterials and Bioengineering, Institute of Integrated Research, Institute

of Science Tokyo, Tokyo, Japan

4 Department of Biomedical Engineering, Faculty of Life Science, Toyo University, Saitama,

Japan

5 Laboratory of Systems Genomics, Department of Computational Biology and Medical

Sciences, Graduate School of Frontier Sciences, the University of Tokyo, Chiba, Japan

#: These authors contributed equally to this work

Contact info

*Correspondence: asahara@scripps.edu

Abstract

Tendons and ligaments are crucial connective tissues linking bones and muscles, yet

achieving full functional recovery after injury remains challenging. We investigated the

characteristics of tendon stem/progenitor cells (TSPCs) by focusing on the declining tendon

repair capacity with growth. Using single-cell RNA sequencing on Achilles tendon cells from

2- and 6-week-old mice, we identified *Cd55* and *Cd248* as novel surface antigen markers for

TSPCs. Combining single-nucleus ATAC and RNA sequencing analyses revealed that *Cd55*

and *Cd248* positive fractions in tendon tissue are TSPCs, with this population decreasing at

6 weeks. We also identified candidate upstream transcription factors regulating these

fractions. Functional analyses of isolated CD55/CD248 positive cells demonstrated high

clonogenic potential and tendon differentiation capacity, forming functional tendon-like tissue
in vitro. This study establishes CD55 and CD248 as novel TSPC surface antigens, potentially
advancing tendon regenerative medicine and contributing to the development of new
treatment strategies for tendon and ligament injuries.
41
Introduction
Tendons are connective tissues involved in the contractile movement of muscle to bone.
Tendons are rich in extracellular matrix components, such as type 1 collagen and
proteoglycans, which both have elastic and viscous properties to withstand overload
(Andarawis-Puri et al., 2015; Asahara et al., 2017; Connizzo et al., 2013).
Tendons have a hierarchical structure. Collagen fibrils aggregate to form fibers, which then
bundle into fascicles. Each fascicle is surrounded by a thin connective tissue layer called the
endotenon. Fascicles are further enclosed by the epitenon, and in some cases, an additional
outer layer called the paratenon. The number of fascicles and the presence of a paratenon
can vary among different tendons and species (Jozsa et al., 1991; Walia et al., 2019).
Tendon and ligament injuries potentially account for the majority of musculoskeletal
disorders and can lead to arthritis and spondylitis. (Gracey et al., 2020). The most popular
treatment approaches are surgery and conservation (Steinmann et al., 2020). However, full
functional recovery is often not achieved due to scarring and fibrosis in the injured area
(Guilak et al., 2014; Thomopoulos S et al., 2015). Furthermore, the risk of postoperative
rupture is high (Leong et al., 2020; Loiacono et al., 2019). In this context, tendon regeneration
focused on tendon stem/progenitor cells (TSPCs) is attracting attention (Leong et al., 2016).
Tendons have few cellular components, including tendon cells (tenocytes) and TSPCs (Tang
et al., 2016; Bi et al., 2007). Tenocytes are responsible for tendon homeostasis, while TSPCs
self-renew and differentiate into tenocytes.
Tendon Stem/Progenitor Cells (TSPCs) were first reported in 2007 by Bi Y et al. (Bi et al.,
2007). They demonstrated that TSPCs possess self-renewal capacity, colony-forming
ability, and multi-differentiation potential *in vitro*. Furthermore, TSPCs were reported to
simultaneously express stem cell markers such as Cd44 and Stem cells antigen-1 (*Ly6a*),
as well as tendon-related genes like Scleraxis (*Scx*) and Collagen type I alpha 1 chain
(*Col1a1*), and were identified as a subset of tendon cells within the tendon fascicle.
Subsequent research on TSPC localization has yielded diverse perspectives. Harvey et al.
reported that *Tppp3*/*Pdgfra*-positive cells in the epitenon are induced upon tendon injury
and contribute to repair (Harvey et al., 2019). Additionally, Yin et al. and Tempfer et al.
demonstrated the presence of TSPCs around blood vessels (Yin et al., 2016; Tempfer et
al., 2009).

Markers used to characterize TSPCs include CD73, CD105, and CD90, which are known

criteria for mesenchymal stem cells (MSCs) (Lui et al., 2015). Additionally, TSPCs have

been reported to express POU class 5 homeobox 1 (*Oct-4*), *Nanog*, Nucleostemin, stage-

specific embryonic anitigen-4 *(SSEA-4)*, *c-Myc*, SRY-box transcription factor *(Sox*2),

Fucosyltransferase 4 (*Fut-4*), and other genes (Zhang et al., 2016). Expression of CD146

and CD44 has been confirmed as well (Ruzzini et al., 2014). The *Tppp3*/*Pdgfra*-positive

TSPCs reported by Harvey et al. highly express *Cd34*, which is generally considered to

have low expression in MSCs (Harvey et al., 2019; Tachibana, et al., 2022). *Cd34* is known

to be highly expressed in mouse embryonic limb buds at E14.5 compared to E11.5 (Havis

et al., 2014), making it a potential marker for TSPCs. However, these markers are not

specific to TSPCs, thus, a definitive method to distinguish TSPCs from mature tendons *in*

vivo has not been established yet (Li et al., 2019; Wang et al., 2008; Chen et al., 2012; Cho

et al., 2018; Fang et al., 2022).

Regarding tendon regenerative capacity, Howell et al. reported an interesting observation.

Tendons in juvenile mice can regenerate functional tissue after injury, but this ability is lost

in mature mice, resulting in scar tissue formation (Howell et al., 2017). This finding

suggests the possibility of abundant TSPCs in juvenile mouse tendons.

Given this background, we hypothesized that evaluating tendon tissue heterogeneity at

the single-cell level using juvenile mouse tendons would enable a more detailed

characterization of TSPCs. This approach is expected to advance the identification and

characterization of TSPCs, which has been challenging with conventional methods.

Therefore, we performed scRNA-seq using cells collected from 2- and 6-week-old mouse

Achilles tendons and investigated clusters that co-express known TSPCs markers, such as

Tppp3, *Pdgfra*, and *Ly6a*. Then, we analyzed the expression of surface antigens in these

clusters and identified *Cd55* and *Cd248* as novel candidate surface antigens for TSPCs.

Furthermore, snATAC-seq and snRNA-seq were performed simultaneously using cells

collected from Achilles tendons to evaluate the validity of *Cd55* and *Cd248* as surface

antigens for TSPCs and to identify the landscape of transcription factors (TFs) involved in

tendon maturation. We sorted mouse Achilles tendon cells based on CD55 and CD248 and

confirmed their phenotypes, demonstrating high clonogenicity and highly efficient induction

into tendon cells. These results suggested that CD55 and CD248 are novel surface antigens

of TSPCs and may be useful for understanding the process of tendon maturation and for

applications in cell therapy.

Material and Methods

Single cell isolation

Achilles tendons were digested to obtain a single-cell suspension. In total, 40 or 60 (2-
week and 6-week, respectively) mouse Achilles tendons were processed together as a
single sample. After washing with 1× phosphate-buffered saline (PBS) several times and
cutting 1-mm-wide sections using scissors, tendons were digested for 1 h at 37°C with
continuous shaking at 1,200 rpm and 37°C in a dissociation solution consisting of 30
mg/mL collagenase (Wako, Osaka, Japan). After digestion, the single-cell suspension was
filtered for debris using a 40 μm cell strainer and washed twice with 1× PBS. The samples
were centrifuged for 15 min at 1,500 rpm and 4°C. Dead cells were removed using a Dead
Cell Removal Kit (Miltenyi Biotec, Bergisch Gladbach, Germany).
118

scRNA-seq library construction and sequencing
Cells were resuspended in PBS with 1% fetal bovine serum (FBS) at a concentration of
10,000 cells per μL. Then, 10,000 cells per sample were loaded on a Chromium Controller
(10x Genomics, Pleasanton, CA, USA) for single cell capture. Libraries were prepared
using Single Cell 3′ Library & Gel Bead Kit v3 (10x Genomics) following the manufacturer's
instructions. A single-cell emulsion (Gel Bead-In-EMulsions, GEMs) was created by making
barcoded cDNA unique to each individual emulsion. A recovery agent was added to break
GEM and cDNA was then amplified. A library was produced via end repair, dA-tailing,
adapter ligation, post-ligation cleanup with SPRIselect, and sample index PCR. The quality
and concentration of the amplified cDNA were evaluated using the Bioanalyzer (Agilent
2100) on a High Sensitivity DNA chip (Agilent, #5065-4401; Santa Clara, CA, USA).
Sequencing was performed using the HiSeq X system (Illumina Inc., San Diego, CA, USA)
to generate 28/90 bp paired-end reads.
132

Cell dissociation, nuclei isolation, and snRNA-seq and snATAC-seq library
construction and sequencing
After obtaining cells from the mouse Achilles tendons, nuclei were isolated following the
Chromium Next GEM Single Cell Multiome ATAC + Gene Expression Reagent Bundle
(10XGenomics) and all of the buffers were made according to the manufacturer's
instructions. Briefly, after centrifugation, the supernatant was discarded, and the cells were
re-suspended in 1 mL of 1% FBS in PBS. Approximately 500,000 cells were transferred to
a new tube for further lysis. To remove the supernatant, cells were centrifuged again for 5
min at 300 rcf and 4°C and resuspended in 100 mL of chilled lysis buffer. Cells were lysed
for 9 min on ice and 1 mL of wash buffer was added to stop the reaction. Then, the
suspension was filtered through a 40 μm cell strainer, cells were centrifuged and
resuspended in 66.2 mL of chilled Diluted Nuclei Buffer aiming to target 7000 nuclei. The

final concentration of nuclei was determined, followed by transposition, GEM generation,

barcoding, and library construction according to the Chromium Next GEM Single Cell

Multiome ATAC + Gene Expression (10XGenomics). Libraries were sequenced with the

parameters recommended by the manufacturer, using the NovaSeq 6000 (Illumina Inc.) to

generate 28/90 bp paired-end reads for gene expression and 50/49 bp paired-end reads for

ATAC sequencing.

scRNA-seq and snRNA-seq analyses

Sequencing reads were processed with the Cellranger_arc (10X Genomics, v2.0.0) using

the mouse reference mm10. From the gene expression matrix, downstream analyses were

carried out using R. Quality control, filtering, data clustering and visualization, and a

differential expression analysis were carried out using Seurat (Butler et al., 2018). For each

dataset, cells with unique feature counts over 2,500 or less than 200 and >5%

mitochondrial counts were filtered. Then, heterotypic doublets (assuming 5% of barcodes

represent doublets) were removed using DoubletFinder (McGinnis et al., 2019).

Unsupervised shared nearest neighbor (SNN) clustering was performed with varying

resolution and the results were visualized using uniform manifold approximation and

projection (UMAP) (Becht et al., 2019). Differentially expressed genes (DEGs) among each

cell cluster were determined using the FindAllMarkers function in Seurat. DEGs expressed

in at least 25% of cells within the cluster and with a log(fold change) of more than 0.25

were identified as marker genes for the cluster.

Pseudotime analysis

Monocle3 (Trapnell et al., 2014) was used to convert the snRNA-seq dataset into a cell

dataset object (CDS), preprocess data, correct for batch effects, embed with dimensional

reduction, and perform pseudotemporal ordering. Cicero (Pliner et al., 2018) was used to

generate pseudo-temporal trajectories for the snATAC-seq dataset.

Cell-cell communication analysis

Intercellular communication networks were quantitatively inferred and analyzed using

scRNA-seq data. The R package CellChat (Jin et al., 2021) was used to visualize the

interactions among different cell groups. Two hundred twenty-nine signaling pathway

families were grouped as a library to analyze cell–cell communication.

GO analysis

The R package ClusterProfiler (Yu et al., 2012) was used to perform a gene enrichment

analysis. The p-values were corrected by the Benjamini & Hochberg method.

Motif enrichment analysis (ChiP seeker)

Genomic regions containing snATAC-seq peaks were annotated using ChIPSeeker (Yu et

al., 2015) and the UCSC database on mouse (mm10).

snATAC-seq analysis

The Cell Ranger ATAC pipeline (1.2.0) (Satpathy et al., 2019) was used to preprocess the

data resulting from sequencing. First, Tn5 sites were mapped to the mouse reference

transcriptome mm10, and duplicate reads and background cells were removed. This

returned barcoded fragment files, which were loaded into Signac (Stuart et al., 2021) for

downstream analyses using the standard Signac/Seurat pipeline. Macs2 (Zhang et al.,

2008) was run on the fragment files to call peaks using the Signac 'CallPeaks' function.

Fragments were mapped to the Macs2 called peaks and assigned to cells using the Signac

'FeatureMatrix' function. Nucleosome signal strength and transcription start site (TSS)

enrichment for each cell were calculated using the Signac 'NucleosomeSignal' and

'TSSEnrichment' functions, respectively. Outliers in the QC metric categories were removed

as per Signac's standard processing guidelines. Latent semantic indexing (LSI), a form of

dimensional reduction, was performed using the Signac 'RunTFIDF' and 'RunSVD'

functions. The UMAP hyperparameters were varied to produce consistent object shapes

(using R). Once hyperparameters were chosen, the Signac/Seurat's 'RunUMAP' function

was run on the LSI dimensions chosen earlier for UMAP embedding. The Signac/Seurat

'FindNeighbors' function was run using the same LSI dimensions used for UMAP to

compute the nearest neighbor graph. Signac/Seurat 'FindClusters' was then run at varying

resolutions. A gene activity matrix was constructed by counting ATAC peaks within the gene

body and 2 kb upstream of the transcriptional start site for protein-coding genes annotated

in the Ensembl database. The gene activity matrix was log-normalized prior to label transfer

with the aggregated snRNA-seq Seurat object using a canonical correlation analysis. The

aggregated snATAC-seq object was filtered using a 97% confidence threshold for cell-type

assignment following label transfer to remove heterotypic doublets. The filtered snATAC-

seq object was reprocessed with TFIDF, SVD, and batch effect correction followed by

clustering and annotation based on lineage- specific gene activity. Differential chromatin

accessibility between cell types was assessed with the Signac FindMarkers function for

peaks detected in at least 20% of cells using a likelihood ratio test and a log-fold-change

threshold of 0.25. Bonferroni-adjusted p-values were used to determine significance at an
FDR < 0.05.
217
Integration of snRNA-seq and snATAC-seq data
snRNA-seq and snATAC-seq data were integrated using the cluster-label transfer
procedure as implemented in Signac and Seurat. Each snRNA-seq sample. as clustered
individually and its cluster labels were projected onto the matching, individually clustered
snATAC-seq sample or vice versa. Anchors were identified for condition-matched snRNA-
and snATAC-seq samples using the FindTransferAnchors function and a canonical
correlation analysis (CCA) was performed using the snRNA expression values and the
snATAC-imputed gene expression values. The anchors were used to transfer cluster-label
identifiers between the two data types using the TransferData function. Each cell in the
query was assigned the cluster label with the highest prediction score, and label transfer
was considered successful for query cells with prediction scores above 0.3.
229
SCENIC analysis
To identify TFs and characterize cell states, a cis-regulatory analysis was performed using
the R package SCENIC (Lake et al., 2018), which infers gene regulatory networks based
on co-expression and DNA motif analyses. The network activity was then analyzed for each
cell to identify recurrent cellular states. In short, TFs were identified using GENIE3 and
compiled into modules (regulons), which were subsequently subjected to cis-regulatory
motif analysis using RcisTarget with two gene-motif rankings: 10 kb around the TSS and
500 bp upstream. Regulon activity in every cell was then scored using AUCell.
238
Immunohistochemistry
Tissue samples were fixed in 4% paraformaldehyde overnight at 4°C, washed in 1× PBS,
cryo-preserved in 20% sucrose overnight, and embedded in optimal cutting temperature
(OCT) compound (4583; Sakura Finetek Japan Co., Ltd., Tokyo, Japan). Then, the bio-
tendon was cryo-sectioned at a thickness of 10 μm and desiccated by air-drying overnight.
Immunohistochemical staining was performed using a Vectastain ABC-AP Rabbit IgG Kit
(AK- 5001; VECTOR LABORATORIES, INC., Burlingame, CA, USA) and Vector Red (SK-
5100; VECTOR LABORATORIES, INC.) according to the manufacturer's instructions. Anti-
CD55 (1/500 dilution) (ab231061; Abcam, Cambridge, UK) and Anti-CD248 antibodies
(1/500 dilution) (LS-B2712; LSBio, Seattle, WA, USA) were used as the primary antibodies.
249
Fluorescence-activated cell sorting (FACS)

Cells harvested from the Achilles tendon were incubated for 60 min on ice with APC-CD55

(#131812; BioLegend, San Diego, CA, USA) and a primary antibody against CD248 (#LS-

B2712) using a FACS buffer (PBS containing 1 [v/v%] FBS). After washing, cells were

reacted with fluorescent-conjugated rabbit secondary antibody. Cell sorting was performed

using MoFlo XDP FACS (Beckman Coulter, Brea, CA, USA).

Primary cell culture

Primary TSPCs were isolated from the Achilles tendons of 2-week-old mice with

collagenase digestion described above. Single-cell suspensions were cultured in the

culture medium (MEMα + 20% FBS + 1% penicillin-streptomycin + 1 [v/v%] 100× non-

essential amino acid solution [NEAA; Gibco], 1 [v/v%] 100× GlutaMAX [Gibco]). At 80%–

90% confluence, cells were trypsinized, centrifuged, resuspended in culture medium as

passage 1 cells, and incubated in 5% CO_2 at 37°C, with fresh medium every 2–3 days.

Colony formation assay

For the colony formation assay, single-cell suspensions of TSPCs (1000 cells/well) were

seeded and incubated in six-well plates for 12–14 days in the growth medium and fixed

with 4% paraformaldehyde (PFA) (Sigma-Aldrich, St. Louis, MO, USA). Then, 0.1% crystal

violet solution (Wako) was used to stain the cells. Colonies of >30–50 cells were defined as

a single colony unit, and the number of clusters was counted using the ImageJ package

ColonyArea (Guzmán et al., 2014)

RT-PCR

Total RNA was extracted using TRIzol reagent (Invitrogen, Grand Island, NY, USA). The

PrimeScript RT Reagent Kit (Takara, Tokyo, Japan) was used for reverse transcription of

mRNA. Complementary DNA was quantitated by qRT-PCR using a Thunderbird SYBR mix

(Toyobo Co., Osaka, Japan). *B2MG* expression served as a control and changes in gene

expression were quantified using the ΔΔCT method.

Tenocytes, cartilage, and osteocyte differentiation

For the differentiation experiment, TSPCs were cultured in 6-well plates (50,000

cells/well). Osteogenic, chondrogenic, and tenogenic differentiation were induced using a

corresponding differentiation medium. The osteogenic differentiation medium contained a

culture medium supplemented with 10 nM dexamethasone (Sigma-Aldrich), 5 mM β-

glycerophosphate (APEXBIO), and 0.05 mM L-ascorbic acid 2-phosphate (Sigma-Aldrich).

The chondrogenic differentiation medium contained a culture medium supplemented with

100 nM dexamethasone and 10 ng/mL BMP2 (Sigma-Aldrich). The tenogenic differentiation

medium contained a culture medium supplemented with 10 ng/mL TGF-β1 (Peprotech,

Rocky Hill, NJ, USA), 10 ng/mL GDF-5 (R&D Systems, Minneapolis, MN, USA), and 0.05

mM L-ascorbic acid 2-phosphate. After a 2 week induction period, cells were harvested for

RT-PCR.

Bio-cultured tendon (Bio-tendon)

Cultured cells were embedded in a 3D-culture cocktail (Tsutsumi et al., 2022). The 3D-

culture cocktail was constructed by mixing collagen gel [final concentrations: 2 mg/mL

Cellmatrix (Type I-A, Nitta Gelatin Inc., Osaka, Japan) and 1× collagen neutralization buffer

(Type I-A, Nitta Gelatin Inc.)], pro-survival cocktail (final concentrations: 100 nM B-cell

lymphoma extra-large (Bcl-Xl) BH4 4-23 (197217-1MG, Calbiochem, United States), 100

μM carbobenzoxy-valyl-alanyl-aspartyl-[*O*-methyl]-fluoromethylketone (Z-VAD-FMK)

(Promega, Madison, WI, USA)), and culture medium. The 3D chamber was coated with 1%

gelatin and incubated at 37°C and 5% CO_2 for 30 min. After washing thrice with 1× PBS,

1.0×10^6 cells were transferred to a 3D-culture cocktail mixture and incubated at 37°C and

5% CO_2 for 60 min for gelation. Following gelation, tenogenic differentiation medium was

added to the chamber. Following 24 h of incubation, the 3D-cultured samples were set into

a mechanical cell stretch system device (Shellpa Pro, Menicon Co., Ltd./Life Science

Department, Aichi, Japan). Cyclic mechanical stretch was performed for 1 week by

gradually increasing the stretch loading rate as follows: 1% (day 1), 2% (day 2), 3% (day 3),

4% (day 4), and 5% (days 5–7). The cyclic mechanical stretch was programed at 0.25 Hz

for 18 h/day, followed by resting for 6 h/day at 37°C and 5% CO_2. Daily medium changes

were also required.

Decellularization by HHP and chemical treatment

The cultured bio-tendon was placed into a plastic pack filled with saline and sealed to

prevent implosion and leakage during the procedure. The pack was then pressurized at

1,000 MPa at 30°C for 10 min using an HHP machine (Dr. CHEF; Kobe Steel, Ltd., Hyogo,

Japan). After pressurization, the bio-tendon was washed thrice with 30% ethanol (EtOH) by

continuous shaking for 5 min at each step. Finally, 1-ethyl-3-(3-dimethylaminopropyl)

carbodiimide (EDC)/*N*-hydroxysuccinimide (NHS)-based cross-linking was performed by

adding 70 mM EDC (Wako) and 70 mM NHS (Wako) with 30% EtOH for 24 h at 4°C. The

processed bio-tendon was incubated in 1× PBS at 4°C until the experiment.

Electron microscopy

Tendon/ligament-like tissues were dissected and fixed in 2.5% glutaraldehyde in 0.1 M

phosphate buffer (PB) overnight. For transmission electron microscopy (TEM), the

specimens (n = 3) were washed with 0.1 M PB, post-fixed in 1% osmium buffered with 0.1 M

PB for 2 h, and dehydrated in a graded series of ethanol. Then, the specimens were

embedded in Epon 812, sliced into ultrathin sections (70 nm), collected on copper grids, and

double-stained with uranyl acetate and lead citrate. The specimens were observed using

TEM (JEM-1400Flash; JEOL, Ltd., Tokyo, Japan). For scanning electron microscopy (SEM),

the specimens (n = 2) were dried in a critical-point drying apparatus (HCP-2; Hitachi, Ltd.)

with liquid CO_2 and were sputter-coated with platinum. Then, the specimens were observed

using SEM (JSM-7900F/JED-2300; JEOL, Ltd.). Fiber alignment in the microscopic sections

was analyze using the OrientationJ image processing tool (an ImageJ plugin).

Stretch test

The mechanical properties were measured using a creep meter (RE-3305S; Yamaden,

Tokyo, Japan). After measuring the initial length (mm), diameter (mm), and thickness (mm)

using a micrometer, the samples were fixed with two grips, which were pulled at a constant

speed of 0.05 mm/s until failure, and the tensile strength (N) and failure strain (mm) were

measured. The cross-sectional area (mm^2) was calculated using the initial diameter and

thickness. The stiffness was manually determined from the slope in the linear region of the

failure-stress curve.

The tensile strength (MPa), failure strain (%), stiffness, and Young's modulus were

calculated using the following formulas:

Tensile strength (MPa) = tensile strength (N)/cross-sectional area (mm^2).

Failure strain (%) = failure strain (mm)/initial length (mm).

Stiffness = Stress (N)/Strain (mm)

Young's modulus = stiffness × initial length (mm)/cross-sectional area (mm^2)

Statistical analysis

All values are presented as means ± SEM. Statistically significant differences were

assessed using unpaired two-tailed Student's t-tests and one-way analysis of variance

(ANOVA) with Tukey's post hoc tests. Statistical significance was set at $p < 0.05$.

Results

Identification of Novel Surface Antigens for Tendon Stem/Progenitor Cells (TSPCs)

In mice, the healing capacity for tendon injuries is high up to 2 weeks of age, but decreases as the musculoskeletal system matures at 6 weeks (Howell et al., 2017). To investigate whether this phenomenon is due to changes in the cellular population of tendon tissue, including fluctuations in progenitor cells, we performed single-cell RNA sequencing on mouse Achilles tendons at these two time points.

Achilles tendons were harvested from mice at 2 and 6 weeks of age. Following collagenase digestion and dead cell removal, we employed massively parallel, droplet-enabled scRNA-seq analysis (10X Genomics Chromium). Data processing was conducted using the Cell Ranger pipeline (10X Genomics). We analyzed 10,314 cells from 2-week-old mice (median 3,167 genes/cell and 41,611 mean reads/cell) and 6,513 cells from 6-week-old mice (median 732 genes/cell and 60,386 mean reads/cell). After doublet removal using Doubletfinder, we merged the datasets and performed unbiased clustering using Seurat, identifying a total of 15 clusters (Fig. 1A). The top differentially expressed genes (DEGs) for each cluster, based on log2 fold change and statistical significance, are summarized in Fig. 1B.

We identified two clusters (0 and 11) with high expression of *Mkx* and *Scx*, transcription factors involved in tendon development and growth. Cluster 0 showed higher expression of ECM-related genes such as *Col1a1* and *Fmod* compared to cluster 11, suggesting these were tenocytes (TC). Cluster 11, with high Periostin (*Postn)* expression, was classified as myotendinous junction (MTJ). Other clusters were characterized as follows: cartilage (CA, high SRY-box transcription factor 9 (*Sox9*) expression), lymphocytes (LC1, LC2, high Lysozyme 2 (*Lyz2*) and Complement component 1, q subcomponent, alpha polypeptide (*C1qa*)), endothelial cells (EC, high Platelet and endothelial cell adhesion molecule 1 (*Pecam1*)), red blood cells (RBC, high Hemoglobin alpha, adult chain 1 (*Hba-a1*)), smooth muscle cells (SM, high Myosin light chain 9 (*Myl9*)), proliferating cells (PC, high *Mki67* and Stathmin 1 (*Stmn1*)), Schwann cells (SW, high Myelin basic protein (*Mbp*)), vascular endothelial cells (VEC, high Multimerin 1 (*Mmrn1*)), and macrophages (MC, high Protein tyrosine phosphatase receptor type C (*Ptprc*)). In cluster3, *Col1a1* and *Fmod* were expressed; however, *Gm42418* and *AY036118* were highly expressed. These long non-coding RNAs are related to RN45s and therefore cluster3 represents ribosomal contamination (Isola et al., 2024).

To identify progenitor populations within these clusters, we analyzed expression patterns of previously reported markers *Tppp3* and *Pdgfra* (Harvey et al., 2019; Tachibana, et al., 2022), along with the known stem/progenitor cell marker *Ly6a* (Holmes et al., 2007; Sung et al., 2008; Hittinger et al., 2013; Sidney et al., 2014; Fang et al., 2022). We identified subclusters within clusters 1 and 4 showing high expression of these genes, which we

defined as SP1 and SP2. SP2 exhibited the highest expression of these genes, suggesting

it had the strongest progenitor characteristics.

SP2 also showed strong expression of genes associated with early tendon development in

mouse embryos, as identified by RNA-seq of mouse limb tendon cells at E14.5 (Havis et

al., 2014) and the Eurexpress mouse embryo transcriptome atlas database (Diez-Roux et

al., 2011) (Supplemental Figure 1A). Gene Ontology analysis revealed upregulation of

tendon development-related pathways in SP2, including TGFβ production and collagen-

containing extracellular matrix (Supplemental Figure 1B). Evaluation of cell numbers in

each cluster showed a decrease in SP2, TC, and MTJ at 6 weeks (Fig. 1C).

We then examined DEGs in SP2 compared to SP1 (Fig. 1D). Among these, we identified

Cd34, *Cd55*, and *Cd248* as genes encoding surface antigens. While *Cd34* has been

previously reported in tendon stem progenitor populations (Harvey et al., 2019; Havis et al.,

2014), to our knowledge, *Cd55* and *Cd248* have not been associated with this context.

Expression of *Cd55* and *Cd248* was localized to areas corresponding to SP1 and SP2 (Fig.

1E). We evaluated the correlation coefficients between the expression of these genes and

previously suggested TSPC markers (*Cd73*, *Cd90*, *Cd105*, *Cd44*, *Cd146*) as well as

Tppp3, *Pdgfra*, and *Ly6a* at 2 weeks of age. *Cd34*, *Cd55*, and *Cd248* showed high

correlation, suggesting these genes as new candidate surface antigens for TSPCs (Fig.

1F).

Comparing gene expression between 2 and 6 weeks, we observed a decrease in *Mkx* and

Scx expression, consistent with previous reports (Grinstein et al., 2019). Furthermore, the

expression of our candidate tendon progenitor cell markers *Cd34*, *Cd55*, and *Cd248* all

decreased at 6 weeks. Inflammatory cells, including macrophages, increased at 6 weeks

(Fig. 1G). Gene expression analysis revealed a shift from M2 macrophage-related genes

(Mannose receptor C-type 1 (*Mrc1*) and Cathepsin C (*Ctsc*)) at 2 weeks to M1

macrophage-related genes (Chemokine (C-C motif) ligand 6 (*Ccl6*) and *Chemokine (C-C*

motif) ligand 9 (*Ccl9*)) at 6 weeks. Analysis of cell-cell communication changes in postnatal

tendon and surrounding tissues (Lui et al., 2019) using CellChat (Jin et al., 2021) showed a

significant decrease in overall interaction at 6 weeks compared to 2 weeks. These results

demonstrate that the cellular profile of tendon tissue changes dramatically at 6 weeks, with

decreased expression of stem progenitor markers and reduced interaction with surrounding

tissues.

Single Nucleus RNA + ATAC Analysis of 2-Week-Old Mouse Achilles Tendon Cells

Gene regulation is mediated by the binding of transcription factors to cis-regulatory

elements proximal to the gene. Consequently, epigenetic changes such as chromatin

accessibility play a crucial role in gene expression (Miller et al., 2013). Moreover,
transcription factors are typically expressed at low levels, potentially leading to false
negatives in scRNA-seq due to detection limits. Chromatin accessibility changes often
precede gene expression changes, potentially allowing us to predict transcriptional
changes (Ranzoni et al., 2020). Therefore, we performed simultaneous ATAC-seq and
RNA-seq at the single-cell level to identify gene expression changes and their associated
epigenetic alterations, aiming to elucidate the post-natal growth mechanisms of tendon
tissue and evaluate the validity of *Cd55* and *Cd248* as markers.
We extracted nuclei from cells isolated from 2-week-old mouse Achilles tendons and
conducted droplet-enabled multi-omics analysis (10X Genomics Chromium), performing
simultaneous snATAC-seq and snRNA-seq as a reference. We first analyzed the snATAC-
seq data. After mapping reads to the genome and calling peaks, we annotated the location
of peaks in terms of genomic features. The peaks were associated with promoters, introns,
exons, and intergenic regions.
We evaluated 6,571 cells (median high-quality fragments per cell: 17,408) and clustered
them using latent semantic indexing and Uniform Manifold Approximation and Projection
(UMAP) with the R package Signac (Stuart et al., 2021) (Figure 2A). Given the limited
knowledge of cell-specific chromatin accessibility, we assessed gene activity by computing
counts per cell within the gene body and promoter of protein-coding genes. Using this gene
activity data, we performed unsupervised clustering and identified 17 clusters (ground-truth
annotation). Clusters expressing tendon-related genes such as *Mkx* and *Scx* were
identified as clusters 1 and 6, defined as A1: TC1 and A6: TC2, respectively. Clusters 0, 14,
and 15 were identified as expressing stem progenitor markers including *Cd34*, *Ly6a*,
Pdgfra, *Cd55*, and *Cd248*, and were defined as A2-0: SP1, A2-14: SP2, and A2-15: SP3,
respectively. The characteristic gene activities and annotations for each cluster are
summarized in Fig. 2B.
Next, we analyzed the snRNA-seq data. We evaluated 10,314 cells (median 3,167
genes/cell and 41,611 mean reads/cell). After doublet removal, we performed unbiased
clustering using Seurat and identified 17 clusters (Supplemental Figure 2A). Clusters
expressing tendon-related genes *Mkx* and *Scx* were clusters 1 and 13, defined as R2-1:
TC_1 and R2-13: TC_2, respectively. Clusters 2, 3, and 7 were identified as expressing
stem/progenitor-related genes and defined as R2-2: SP_1, R2-3: SP_2, and R2-7: SP_3,
respectively. Using the Signac package's "FindTransferAnchors" function, we calculated
predicted IDs for snATAC-seq clusters based on snRNA-seq annotations (predicted
annotation) (Fig. 2C). Evaluation of the two annotation methods for snATAC-seq revealed
that both methods allowed annotation of major cell types, and correlation between the two

was maintained. Thus, the validity of predicted annotations based on snRNA-seq was

confirmed (Figure 2D).

R2-7: SP3, considered the most undifferentiated stem/progenitor fraction in snRNA-seq,

could be divided into A2-0: SP1 and A2-14: SP2 in snATAC-seq.

Comparing gene activity of tendon-related and stem/progenitor genes in each cluster of the

ground truth annotation, A2-14: SP2 showed high expression of *Ly6a*, *Pdgfra*, and *Tppp3*,

and inverse correlation with *Mkx*, *Scx*, and *Tnmd*. Furthermore, the newly identified surface

antigens *Cd55* and *Cd248* also showed high expression in cluster 14. In contrast, *Cd34* did

not show a high expression pattern in A2-14: SP2. These results suggest that A2-14: SP2

is the most immature cluster, with *Cd55* and *Cd248* showing characteristic expression (Fig.

2E). Subcluster analysis of R7, considered the most immature fraction in snRNA-seq,

revealed two clusters based on *Cd55*/*Cd248* and *Cd34* expression, similar to the snATAC-

seq analysis. The former (R7-1) showed high expression of *Ly6a*, *Tppp3*, and *Pdgfra* (Fig.

2F). Trajectory analysis using R2-7 and A2-14 as roots in snRNA-seq and snATAC-seq,

respectively, showed increased expression of tendon-related genes *Mkx* and *Scx* along the

trajectory, while expression of *Tppp3*, *Cd55*, and *Cd248* decreased (Figure 3A). Evaluation

of peaks in each cluster based on gene expression levels in snRNA-seq revealed multiple

peaks with heights correlated to gene expression for *Mkx* and *Scx*. Similar evaluation of

Cd55 and *Cd248* showed multiple peaks upstream of the TSS for *Cd248* (Fig. 3B). These

results indicate correlation between snRNA-seq and snATAC-seq, and consistent with

scRNA-seq results, suggest that A2-14 represents the most undifferentiated TSPCs based

on *Cd55* and *Cd248* expression.

Comparison of Single Nucleus RNA+ATAC Analysis between 2-week and 6-week

Mouse Achilles Tendon Cells

To confirm the presence of the identified *Cd55*/*Cd248* fraction at 6 weeks, we performed

snRNA-seq and snATAC-seq on 6-week-old mouse Achilles tendon cells. Using the 2-week

snATAC-seq data as a reference, we annotated the 6-week snATAC-seq clusters. Clusters

A2-14 and A2-0 corresponded to clusters A6-12 and A6-0, respectively (Supplemental

Figure 4A, 4B). Gene activity evaluation revealed that, similar to the 2-week data, the 6-

week A6-12 and A6-0 clusters showed *Cd55*/*Cd248* high and *Cd34* dim, and Cd55/*Cd248*

dim and *Cd34* high patterns, respectively. Furthermore, annotation of the 6-week snRNA-

seq data using the 2-week snRNA-seq as a reference revealed clusters with expression

patterns similar to those observed in snATAC-seq. In both cases, clusters with high *Cd55*

and *Cd248* expression showed high gene activity (snATAC-seq) and gene expression

(snRNA-seq) of *Tppp3*, *Pdgfra*, and *Ly6a*. These results confirm that the *Cd55* and *Cd248*

high clusters identified in the 2-week snRNA-seq+snATAC-seq analysis are similarly

detected at 6 weeks.

We also compared the snATAC-seq data between 2 and 6 weeks. No significant

differences were observed in genomic annotations between the two time points

(Supplemental Figure 6A). After merging the datasets (Supplemental Figure 6B), we

compared gene activity of tendon/stem-related genes to evaluate changes in gene activity.

We found that the activity of *Tppp3*, *Pdgfra*, *Ly6a*, *Cd55*, *Cd248*, and *Cd34* all decreased at

6 weeks (Supplemental Fig 6C). This was consistent with the decreased gene expression

observed when comparing 2-week and 6-week data.

Estimation of Transcription Factor Activity in 2-week Mouse Achilles Tendon Cells

To estimate transcription factor activity in each cluster, we used the Single-Cell Regulatory

Network Inference and Clustering (SCENIC) package (Lake et al., 2018) to calculate gene

regulatory network activity from scRNA-seq gene expression data. SCENIC constructs

gene expression networks centered on transcription factors and infers transcription factor

activity in each cluster. We summarized the predicted transcription factor activities in

tendon and stem/progenitor-related clusters R2-7, R2-2, R2-3, R2-1, and R2-13 identified

by snRNA-seq (Figure 4).

Next, we evaluated the gene activity of these identified transcription factors in each

snATAC-seq cluster, along with motif activity calculated by chromVAR (Schep et al., 2017).

We also assessed the expression levels of transcription factors in each snRNA-seq cluster.

In the most immature fractions, represented by cluster A2-14 in snATAC-seq and cluster

R2-7 in snRNA-seq, KLF transcription factor 3 (*Klf3*), KLF transcription factor 4 (*Klf4*), and

cAMP responsive element binding protein 5 (*Creb5*) showed consistent behavior in gene

activity, motif activity, and gene expression. Additionally, Signal transducer and activator of

transcription 2 (*Stat2*) and cAMP responsive element binding protein 3 like 1 (*Creb3l1*)

showed high values in A2-0/A2-15 and A2-1, respectively. *Stat2* gene expression was not

observed, likely due to false positives resulting from low expression levels. These results

suggest a correlation between the transcription factor activity predicted by SCENIC and the

gene activity and motif activity derived from snATAC-seq.

Furthermore, SCENIC can predict candidate transcription factors regulating each gene

(Table 1). *Cd55* and *Cd248* were predicted to be regulated by *Klf3* and *Klf4*, while *Mkx* and

Scx were predicted to be under the control of *Creb3l1*. These predictions were consistent

with the data calculated from gene activity and motif activity in snATAC-seq.

***In vitro* Evaluation of CD55 and CD248**

Immunostaining evaluation of CD55 and CD248, the identified candidate stem/progenitor markers, revealed expression in the tendon sheath (Figure 5C, 5A). This was consistent with previous reports of localized expression of *Tppp3*/*Pdgfra*-positive cells in the tendon sheath and our analysis showing co-expression of *Tppp3*/*Pdgfra* and *Cd55*/*Cd248*. Given that *Cd55* and *Cd248* expression appeared to reflect *Tppp3*, *Pdgfra*, and *Ly6a* expression more sensitively than *Cd34*, we extracted tendon cells from 2-week-old mice and sorted them by FACS to determine the biological phenotype of Cd55 and Cd248 positive cells (Figure 5B).

Gene expression analysis of sorted cells showed that CD55/CD248 positive cells, compared to negative cells, had lower expression of *Mkx*, *Scx*, *Col1a1*, and *Creb3l1*, but higher expression of *Ly6a*, *Tppp3*, *Pdgfra*, *Creb5*, and *Klf3*, consistent with snRNA-seq analysis (Supplemental Figure 6). To assess stem/progenitor capacity, we performed colony formation assays. CD55/CD248 sorted cells showed significantly increased colony formation compared to CD55/CD248 negative cells (control) (Figure 5C).

We then evaluated the differentiation tendencies of these cells towards tendon, cartilage, and bone. Tendon differentiation resulted in clusters of spindle-shaped cells from CD55/CD248 positive cells, suggesting tenogenic differentiation (Figure 5D). Gene expression analysis revealed that while CD55/CD248 positive cells initially had lower *Mkx* and *Scx* expression than negative cells, this pattern reversed after differentiation (Figure 5E). Increased expression of other tendon-related genes such as *Col1a1* and *Bgn* was also observed. In contrast, no significant increases in *Sox9* and Collagen type II alpha 1 chain (*Col2a1*)(chondrogenic) or RUNX family transcription factor 2 (*Runx2*), and Alkaline phosphatase, biomineralization associated (*Alpl*) (osteogenic) expression were observed compared to negative cells during cartilage and bone differentiation assays (Supplemental Figure 7). These results suggest that CD55/CD248 positive cells possess tenogenic differentiation capacity.

To further evaluate the tenogenic potential of CD55/CD248 positive cells, we created tendon-like tissue (bio-tendon) using our previously reported 3D stretch stimulation culture system. SEM and TEM analysis to assess collagen fiber density and thickness showed that bio-tendons derived from CD55/CD248 positive cells had an increased proportion of collagen fibers parallel to the stretch direction and increased collagen fiber diameter. TEM also revealed characteristic banding patterns and triple helix structures in these bio-tendons, indicating mature collagen organization (Wieczorek et al., 2015) (Figure 5F-H). Stretch tests to evaluate the mechanical capacity of the bio-tendons showed high tensile strength (Figure 5I).

These results demonstrate that CD55/CD248 positive cells have a tendency to

differentiate into tendon cells.

Discussion

Tendons are known for their low cellular content, making complete functional recovery

after injury challenging and increasing the risk of re-rupture. As a result, cell therapy has

garnered attention as a novel treatment strategy, distinct from current conservative and

surgical approaches. However, this requires a thorough understanding of Tendon

Stem/Progenitor Cells (TSPCs). Research on TSPCs has been limited due to the lack of

identified characteristic surface antigens.

To our knowledge, no multi-omics analysis comparing juvenile and mature mouse tendon

cells has been conducted. In this study, we performed single-cell RNA-seq along with

snRNA-seq+snATAC-seq on nuclei isolated from 2-week and 6-week mouse Achilles

tendons. This approach allowed us to identify candidate surface antigens for TSPCs and

elucidate the dynamism of transcription factors involved in post-developmental tendon

growth. As a result, we discovered a novel combination of *Cd55* and *Cd248* as surface

antigens showing characteristic expression in the TSPC fraction.

While scRNA-seq analysis also showed *Cd34* expression characteristic of the TSPC

fraction, consistent with previous reports, snRNA-seq + snATAC-seq analysis revealed that

the *Cd34* high-expression cluster could be further classified into two clusters based on

Cd55 and *Cd248* expression patterns. The cluster showing high expression of both *Cd55*

and *Cd248* also exhibited high expression of *Tppp3*, *Pdgfra*, and *Ly6a*, suggesting that

isolating *Cd55* and *Cd248* positive fractions may more sensitively capture immature

populations compared to *Cd34*.

snRNA-seq+snATAC-seq analysis provided simultaneous information on gene expression

levels and open chromatin regions for each cluster. Notably, *Cd248* showed several peaks

upstream of the TSS in high-expression clusters, likely reflecting its expression regulation

mechanism. We also identified several peaks that increased proportionally with expression

for *Mkx* and *Scx*. Given the many unknowns in *Mkx* and *Scx* expression regulation

mechanisms (Guerquin et al., 2013; Otabe et al., 2015), we used the SCENIC package to

predict upstream transcription factors based on expressed genes in each snRNA-seq

cluster. Combined with gene activity and motif activity data from snATAC-seq, Creb3l1 was

identified as a transcription factor regulating *Mkx* and *Scx* expression. *Creb3l1* has been

reported to increase in expression throughout development (Liu et al., 2015).

Klf3, *Klf4*, and *Creb5* were identified as characteristic transcription factors in the TSPC

fraction. Recent reports have highlighted the role of the *Klf* family in limb development (Kult

et al., 2021), suggesting its potential importance in tendon differentiation. *Creb5* has been

reported to show increased expression from E11 to E13 (Liu et al., 2015), suggesting a role

in early developmental stages. Furthermore, *Creb5* has been reported to regulate

Proteoglycan 4 (*Prg4*) expression (Zhang et al., 2021). Further investigation into the

functions of these transcription factors in TSPCs is necessary.

CD55 was identified by Hoffmann et al. in 1969 as a surface antigen functioning as a

complement inhibitory factor on erythrocytes. CD55 is known to be expressed from early

developmental stages and characterizes the initial differentiation stage of hematopoietic

stem cells (Guo et al., 2013). It is also expressed in MSCs and has been reported as an

early progenitor marker for mouse mammary epithelial cells (Pal et al., 2017). CD248 was

identified in 1992 as an antigen for the FB5 antibody reacting with vascular wall cells

(Rettig et al., 1992). CD248 is a transmembrane glycoprotein expressed in pericytes and

fibroblasts during developmental stages. Regarding their relationship, CD55 and CD248

have been reported to be expressed in stromal cells during the early stages of arthritis

(Choi et al., 2017), but their detailed mechanisms in tendons remain unclear.

Previously suggested TSPC surface antigen candidates such as *Cd73*, *Cd90*, and *Cd105*

showed poor correlation with the expression of genes like Tppp3. Moreover, the initial

report on TSPCs (Bi et al., 2007) described TSPCs as *Scx+Cd34-*. The high expression of

CD55 and CD248 in the tendon sheath, similar to *Tppp3* (Harvey et al., 2019; Staverosky

et al., 2019), suggests the possibility of cells with tenogenic differentiation potential

exhibiting *Scx+Cd34-* expression patterns within the tendon. Future studies using lineage

tracing experiments with mice labeled for CD55 and CD248 are necessary to analyze the

developmental functions of CD55 and CD248 positive cells and their roles in injury healing

in more detail.

Clinically, to our knowledge, few studies have sorted TSPCs based on surface antigens

and examined their *in vitro* tendon tissue-generating ability. In this study, we found that cells

sorted for CD55/CD248 showed higher clonogenicity compared to CD55-/CD248- cells and

demonstrated superior tendon tissue-generating ability in an artificial tendon model. In the

future, CD55/CD248 double-positive TSPC cells may prove useful in clinical applications

such as *in vitro* artificial tendon creation (Tsutsumi et al., 2022) and cell therapy for tendon

injuries (Huang et al., 2021).

Reference

Andarawis-Puri, N., Flatow, EL., Soslowsky, LJ. (2015). Tendon basic science: Development,

repair, regeneration, and healing. J Orthop Res. 33(6), 780-4.

Asahara, H., Inui, M., Lotz, MK. (2017). Tendons and Ligaments: Connecting Developmental

Biology to Musculoskeletal Disease Pathogenesis. J Bone Miner Res. 32(9), 1773-1782.

Becht, E., McInnes, L., Healy, J., Dutertre, CA., Kwok, IWH., Ng, LG., Ginhoux, F., Newell.

EW. (2019). Dimensionality reduction for visualizing single-cell data using UMAP. Nat

Biotechnol. 37, 38-44.

Bi Y, Ehirchiou D, Kilts TM, Inkson CA, Embree, MC., Sonoyama, W., Li, L., Leet, AI., Seo,

BM., Zhang, L., Shi, S., Young, MF. (2007). Identification of tendon stem/progenitor cells and

the role of the extracellular matrix in their niche. Nat Med. 13(10), 1219-27.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., Satija, R. (2018). Integrating single-cell

transcriptomic data across different conditions, technologies, and species. Nat Biotechnol.

36(5), 411-420.

Cao, Y., Liu, Y., Liu, W., Shan, Q., Buonocore, SD., Cui, L. (2002). Bridging tendon defects

using autologous tenocyte engineered tendon in a hen model. Plast Reconstr Surg. 110(5),

1280-9.

Chen, B., Wang, B., Zhang, WJ., Zhou, G., Cao, Y., Liu, W. (2012). In vivo tendon engineering

with skeletal muscle derived cells in a mouse model. Biomaterials. 33(26), 6086-97.

Chen, L., Liu, JP., Tang, KL., Wang, Q., Wang, GD., Cai, XH., Liu, XM. (2014). Tendon derived

stem cells promote platelet-rich plasma healing in collagenase-induced rat achilles

tendinopathy. Cell Physiol Biochem. 34(6), 2153-68.

Cho, YT., Hsu, WY., Lin, LF., Lin, YN. (2018). Kinesio taping reduces elbow pain during

resisted wrist extension in patients with chronic lateral epicondylitis: a randomized, double-

blinded, cross-over study. BMC Musculoskelet Disord. 19(1), 193.

Connizzo, BK., Yannascoli, SM., Soslowsky, LJ. (2013). Structure-function relationships of

postnatal tendon development: a parallel to healing. Matrix Biol. 32(2), 106-16.

Craft, A. M., Ahmed, N., Rockel, J. S., Baht, G. S., Alman, B. A., Kandel, R. A., Grigoriadis,

A. E. and Keller, G. M. (2013). Specification of chondrocytes and cartilage tissues from

embryonic stem cells. Development 140, 2597-2610.

Diez-Roux, G., Banfi, S., Sultan, M., Geffers, L., Anand, S., Rozado, D., Magen, A., Canidio,

E., Pagani, M., Peluso, I., Lin-Marq, N., Koch, M., Bilio, M., Cantiello, I., Verde, R., De, Masi,

C., Bianchi, SA., Cicchini, J., Perroud, E., Mehmeti, S., Dagand, E., Schrinner, S., Nürnberger,

A., Schmidt, K., Metz, K., Zwingmann, C., Brieske, N., Springer, C., Hernandez, AM., Herzog,

S., Grabbe, F., Sieverding, C., Fischer, B., Schrader, K., Brockmeyer, M., Dettmer, S., Helbig,

C., Alunni, V., Battaini, MA., Mura, C., Henrichsen, CN., Garcia-Lopez, R., Echevarria, D.,

Puelles, E., Garcia-Calero, E., Kruse, S., Uhr, M., Kauck, C., Feng, G., Milyaev, N., Ong, CK.,

Kumar, L., Lam, M., Semple, CA., Gyenesei, A., Mundlos, S., Radelof, U., Lehrach, H.,

Sarmientos, P., Reymond, A., Davidson, DR., Dollé, P., Antonarakis, SE., Yaspo, ML.,

Martinez, S., Baldock, RA., Eichele, G., Ballabio, A. (2011). A High-Resolution Anatomical

Atlas of the Transcriptome in the Mouse Embryo. PLoS Biol. 9(1), e1000582.

Gracey, E., Burssens, A., Cambré, I., Schett, G., Lories, R., McInnes, IB., Asahara, H.,

Elewaut, D. (2020). Tendon and ligament mechanical loading in the pathogenesis of

inflammatory arthritis. Nat Rev Rheumatol. 16(4):193–207.

Fang, F., Xiao, Y., Zelzer, E., Leong, KW., Thomopoulos, S. (2022). A mineralizing pool of

Gli1-expressing progenitors builds the tendon enthesis and demonstrates therapeutic

potential. Cell Stem Cell. 29(12), 1669-1684.

Franken, NA., Rodermond, HM., Stap, J., Haveman, J., van Bree C. (2006). Clonogenic

assay of cells in vitro. Nat Protoc. 2006. 1(5), 2315-9

Grinstein, M., Dingwall, HL., O'Connor, LD., Zou, K., Capellini, TD., Galloway, JL. (2019). A

distinct transition from cell growth to physiological homeostasis in the tendon. Elife. 8, e48689.

Harvey, T., Flamenco, S., Fan, CM. (2019). A Tppp3 + Pdgfra + tendon stem cell population

contributes to regeneration and reveals a shared role for PDGF signalling in regeneration

and fibrosis. Nat Cell Biol. 21(12), 1490-1503.

Guilak, F., Butler, DL., Goldstein, SA., Baaijens, FP. (2014). Biomechanics and

mechanobiology in functional tissue engineering. J Biomech. 47(9), 1933-40

Gumucio, JP., Sugg, KB., Mendias, CL. (2015). TGF-β superfamily signaling in muscle and

tendon adaptation to resistance exercise. Exerc Sport Sci Rev. 43(2), 93-9.

Guzmán, C., Bagga, M., Kaur, A., Westermarck, J., Abankwa, D. (2014). ColonyArea: an
ImageJ plugin to automatically quantify colony formation in clonogenic assays. PLoS One.
9(3), e92444.

Havis, E., Bonnin, MA., Olivera-Martinez, I., Nazaret, N., Ruggiu, M., Weibel, J., Durand, C.,
Guerquin, MJ., Bonod-Bidaud, C., Ruggiero, F., Schweitzer, R., Duprez, D. (2014).
Transcriptomic analysis of mouse limb tendon cells during development. Development.
141(19), 3683-96.

Hittinger, M., Czyz, ZT., Huesemann, Y., Maneck, M., Botteron, C., Kaeufl, S., Klein, CA.,
Polzer, B. (2013). Molecular profiling of single Sca-1+/CD34+,- cells--the putative murine lung
stem cells. PLoS One. 8(12), e83917.

Hoffmann EM. (1969). Inhibition of complement by a substance isolated from human
erythrocytes. II. Studies on the site and mechanism of action. Immunochemistry. 6(3), 405-
19.

Holmes, C., Stanford, WL. (2007). Concise review: stem cell antigen-1: expression, function,
and enigma. Stem Cells. 25(6), 1339-47.

Howell, K., Chien, C., Bell, R., Laudier, D., Tufa, SF., Keene, DR., Andarawis-Puri, N., Huang,
AH. (2017). Novel Model of Tendon Regeneration Reveals Distinct Cell Mechanisms
Underlying Regenerative and Fibrotic Tendon Healing. Sci Rep. 7, 45238.

Isola, JVV., Ocañas, SR., Hubbart, CR., Ko, S., Mondal, SA., Hense, JD., Carter, HNC.,
Schneider, A., Kovats, S., Alberola-Ila, J., Freeman, WM., Stout, MB. (2024). A single-cell
atlas of the aging mouse ovary. Nat Aging. 4(1), 145-162.

Jozsa, L., Kannus, P., Balint, JB., Reffy, A. (1991). Three-dimensional ultrastructure of human
tendons. Acta Anat (Basel). 142(4), 306-12.

Kendal, AR., Layton, T., Al-Mossawi, H., Appleton, L., Dakin, S., Brown, R., Loizou, C.,
Rogers, M., Sharp, R., Carr, A. (2020). Multi-omic single cell analysis resolves novel stromal
cell populations in healthy and diseased human tendon. Sci Rep. 10(1): 13939.

Leong, DJ., Sun, HB. (2016). Mesenchymal stem cells in tendon repair and regeneration:
basic understanding and translational challenges. Ann. N. Y. Acad. Sci. 1383, 88–96.
755

756 Leong, NL., Kator, JL., Clemens, TL., James, A., Enamoto-Iwamoto, M., Jiang, J. (2020).
757 Tendon and Ligament Healing and Current Approaches to Tendon and Ligament
758 Regeneration. J Orthop Res. 38(1), 7-12.
759

760 Li, Y., Wu, T., Liu, S. (2021). Identification and Distinction of Tenocytes and Tendon-Derived
761 Stem Cells. Front Cell Dev Biol. 9, 629515.
762

763 Liu, H., Xu, J., Liu, CF., Lan, Y., Wylie, C., Jiang, R. (2015). Whole transcriptome expression
764 profiling of mouse limb tendon development by using RNA-seq. J Orthop Res. 33(6), 840–
765 848.
766

767 Loiacono, C., Palermi, S., Massa, B., Belviso, I., Romano, V., Gregorio, AD., Sirico, F., Sacco,
768 AM. (2019). Tendinopathy: Pathophysiology, Therapeutic Options, and Role of Nutraceutics.
769 A Narrative Literature Review. Medicina (Kaunas). 55(8), 447.
770

771 Lui, PP. (2015). Markers for the identification of tendon-derived stem cells in vitro and tendon
772 stem cells in situ - update and future development. Stem Cell Res Ther. 6(1), 106.
773

774 Lui, PPY., Wong, CM. (2019). Biology of Tendon Stem Cells and Tendon in Aging. Front
775 Genet. 10, 1338.
776

777 McGinnis, CS., Murrow, LM., Gartner, ZJ. (2019). DoubletFinder: Doublet Detection in Single-
778 Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst. 8(4), 329-337.e4.
779

780 Mendias, CL., Gumucio, JP., Bakhurin, KI., Lynch, EB., Brooks, SV. (2012). Physiological
781 loading of tendons induces scleraxis expression in epitenon fibroblasts. J Orthop Res. 30(4),
782 606-12.
783

784 Morikawa, S., Mabuchi, Y., Kubota, Y., Nagai, Y., Niibe, K., Hiratsu, E., Suzuki, S., Miyauchi-
785 Hara, C., Nagoshi, N., Sunabori, T., Shimmura, S., Miyawaki, A., Nakagawa, T., Suda, T.,
786 Okano, H., and Matsuzaki, Y. (2009). Prospective identification, isolation, and systemic
787 transplantation of multipotent mesenchymal stem cells in murine bone marrow. J. Exp.
788 Med. 206, 2483-2496.

Nusspaumer, G., Jaiswal, S., Barbero, A., Reinhardt, R., Ishay Ronen, D., Haumer, A., Lufkin, T., Martin, I. and Zeller, R. (2017). Ontogenic identification and analysis of mesenchymal stromal cell populations during mouse limb and long bone development. Stem Cell Rep. 9, 1124-1138.

Pliner, HA., Packer, JS., McFaline-Figueroa, JL., Cusanovich, DA., Daza, RM., Aghamirzaie, D., Srivatsan, S., Qiu, X., Jackson, D., Minkina, A., Adey, AC., Steemers, FJ., Shendure, J., Trapnell, C. (2018). Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell Chromatin Accessibility Data. Mol. Cell. 71, 858–871. e8.

Ruzzini, L., Abbruzzese, F., Rainer, A., Longo, UG., Trombetta, M., Maffulli, N., Denaro, V. (2014). Characterization of age-related changes of tendon stem cells from adult human tendons. Knee Surg Sports Traumatol Arthrosc. 22(11), 2856-66.

Satpathy, AT., Granja, JM., Yost, KE., Qi, Y., Meschi, F., McDermott, GP., Olsen, BN., Mumbach, MR., Pierce, SE., Corces, MR., Shah, P., Bell, JC., Jhutty, D., Nemec, CM., Wang, J., Wang, L., Yin, Y., Giresi, PG., Chang, ALS., Zheng, GXY., Greenleaf, WJ., Chang, HY. (2019). Massively parallel single-cell chromatin landscapes of human immune cell development and intratumoral T cell exhaustion. Nat Biotechnol. 37(8), 925-936.

Sidney, LE., Branch, MJ., Dunphy, SE., Dua, HS., Hopkinson, A. (2014). Concise review: evidence for CD34 as a common marker for diverse progenitors. Stem Cells. 32(6), 1380-9.

Somerville, JM., Aspden, RM., Armour, KE., Armour, KJ., Reid, DM. (2004). Growth of C57BL/6 mice and the material and mechanical properties of cortical bone from the tibia. Calcif Tissue Int. 74(5), 469-75.

Staverosky, JA., Pryce, BA., Watson, SS., and Schweitzer, R. (2019). Tubulin polymerization-promoting protein family member 3, Tppp3, is a specific marker of the differentiating tendon sheath and synovial joints. Dev Dyn. 238(3), 685-92.

Steinmann, S., Pfeifer, CG., Brochhausen, C., Docheva, D. (2020). Spectrum of Tendon Pathologies: Triggers, Trails and End-State. Int J Mol Sci. 21(3), 844.

Still, C, 2nd., Chang, WT., Sherman, SL., Sochacki, KR., Dragoo, JL., Qi, LS. (2021). Single-

cell transcriptomic profiling reveals distinct mechanical responses between normal and
diseased tendon progenitor cells. Cell Rep Med. 2(7), 100343.

Stuart, T., Hanna, J., Gutruf, P. (2021). Single-cell chromatin state analysis with Signac. Nat
Methods. 18(11), 1333-1341.

Sung, JH., Yang, HM., Park, JB., Choi, GS., Joh, JW., Kwon, CH., Chun, JM., Lee, SK., Kim,
SJ. (2008). Isolation and characterization of mouse mesenchymal stem cells. Transplant
Proc. 40(8), 2649-54.

Tachibana, N., Chijimatsu, R., Okada, H., Oichi, T., Taniguchi, Y., Maenohara, Y., Miyahara,
J., Ishikura, H., Iwanaga, Y., Arino, Y., Nagata, K., Nakamoto, H., Kato, S., Doi, T.,
Matsubayashi, Y., Oshima, Y., Terashima, A., Omata, Y., Yano, F., Maeda, S., Ikegawa, S.,
Seki, M., Suzuki, Y., Tanaka, S., Saito, T. (2022). RSPO2 defines a distinct undifferentiated
progenitor in the tendon/ligament and suppresses ectopic ossification. Sci Adv. 8(33),
eabn2138.

Takakura, N., Yoshida, H., Ogura, Y., Kataoka, H., Nishikawa, S., and Nishikawa, S. (1997).
PDGFR alpha expression during mouse embryogenesis: immunolocalization analyzed by
whole-mount immunohistostaining using the monoclonal anti-mouse PDGFR alpha antibody
APA5. J. Histochem. Cytochem. 45, 883-893.

Tang, JB., Wu, YF., Cao, Y., Chen, CH., Zhou, YL., Avanessian, B., Shimada, M., Wang, XT.,
Liu, PY. (2016). Basic FGF or VEGF gene therapy corrects insufficiency in the intrinsic
healing capacity of tendons. Sci Rep. 6, 20643.

Tempfer, H., Wagner, A., Gehwolf, R., Lehner, C., Tauber, M., Resch, H., Bauer, HC. (2009).
Perivascular cells of the supraspinatus tendon express both tendon- and stem cell-related
markers. Histochem Cell Biol. 131(6), 733-41.

Thomopoulos, S., Parks, WC., Rifkin, DB., Derwin KA. (2015). Mechanisms of tendon injury
and repair. J Orthop Res. 33(6), 832-9.

Trapnell, C., Cacchiarelli. D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, NJ., Livak,
KJ., Mikkelsen, TS., Rinn, JL. (2014). The dynamics and regulators of cell fate decisions are
revealed by pseudotemporal ordering of single cells. Nat Biotechnol. 32(4), 381-386.

Tsutsumi, H., Kurimoto, R., Nakamichi, R., Chiba, T., Matsushima, T., Fujii, Y., Sanada, R.,

Kato, T., Shishido, K., Sakamaki, Y., Kimura, T., Kishida, A., Asahara, H. (2022). Generation

of a tendon-like tissue from human iPS cells. J Tissue Eng. 13, 20417314221074018.

Walia,　B., Huang, AH. (2019). Tendon stem progenitor cells: Understanding the biology to

inform therapeutic strategies for tendon repair. J Orthop Res . 37(6), 1270-1280.

Wang, B., Liu, W., Zhang, Y., Jiang, Y., Zhang, WJ., Zhou, G., Cui, L., Cao, Y. (2008).

Engineering of extensor tendon complex by an ex vivo approach. Biomaterials. 29(20): 2954-

61.

Yin, Z., Hu, JJ., Yang, L., Zheng, ZF., An, CR., Wu, BB., Zhang, C., Shen, WL., Liu, HH.,

Chen, JL., Heng, BC., Guo, GJ., Chen, X., Ouyang, HW. (2016). Single-cell analysis reveals

a nestin + tendon stem/progenitor cell population with strong tenogenic potentiality. Sci Adv.

2(11), e1600874.

Yu, G., Wang, LG., Han, Y., He, QY. (2012). clusterProfiler: an R package for comparing

biological themes among gene clusters. OMICS. 16(5), 284-7.

Yu, G., Wang, LG., He, QY. (2015). ChIPseeker: an R/Bioconductor package for ChIP peak

annotation, comparison and visualization. Bioinformatics. 31(14), 2382-3.

Zhang, C., Zhu, J., Zhou, Y., Thampatty, BP., Wang, JH. (2019). Tendon Stem/Progenitor

Cells and Their Interactions with Extracellular Matrix and Mechanical Loading. Stem Cells

Int .2019, 3674647.

Acknowledgements

We thank all the members of the Department of Systems BioMedicine at Institute of Science

Tokyo for their support. We also thank the Research Core at Institute of Science Tokyo for

supporting cell sorting.

Author contributions

Conceptualization, H.T., T.C., and H.A.; Methodology, H.T., T.C., and H.A.; Formal analysis,

H.T.; Investigation, H.T., F. Y., T. M., T. K., A. K., K. A., Y. S.; Writing – Original Draft, H.T.;

Writing – Review & Editing, T.C., and H.A.; Funding Acquisition, H.A.; Supervision, H.A.

Data availability

FASTQ data of RNA-Seq and ATAC-seq are deposited in DDBJ under accession number PRJDB18857

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

This work was supported by JSPS KAKENHI (Grant Numbers JP15H02560, JP20H05696, 16H06279 (PAGS)), AMED (Grant Numbers JP21gm0810008, JP23ym0126805, JP24gm0010009), and NIH (Grant Number R01AR080127) to H.A.

Figure legends

Figure 1. scRNA-seq of tendon cells from 2-week-old and 6-week-old mice and the identification of surface markers of TSPC.

(A) Integrated uniform manifold approximation and projection (UMAP) scRNA-seq clustering of cells harvested from 2-week-old and 6-week-old mouse Achilles tendons.

(B) Dot plot of average gene expression levels of the indicated genes in each scRNA-seq cluster. The size of the dot reflects the percentage of cells in the cluster that express each gene. TC, tenocyte; SP1, tendon stem/progenitor cell_1; CA, cartilage; RB, ribosomal RNA; SP2, tendon stem/progenitor cell_2; LC1, lymphocyte_1; EC, endothelial cell; RBC, red blood cell; SM, smooth muscle cell; PC, proliferating cell; LC2, lymohocytes_2; MTJ, myotendinous junction cell; SW, Schwann cell; VEC, vascular endothelial cell; MC, macrophage.

(C) Proportions of cells in clusters identified from scRNA-seq. Clusters are colored according to cluster type.

(D) Volcano plot of gene expression in the SP2 cluster and the identification of candidate TSPC marker genes (red under line).

(E) Feature plot of Cd55 and Cd248 expression.

(F) Correlation of gene expression of TSPC candidate genes in 2-week data.

(G) Violin plots presenting the gene expression changes for a selection of differentially expressed genes.

Figure 2. snATAC-seq of tendon cells from a 2-week-old mouse and the validation of Cd55 and Cd248 as candidate markers of TSPC.

(A) UMAP snATAC-seq clustering of cells derived from the Achilles tendon of a 2-week-old mouse. Annotation was based on each gene activity (ground-truth annotation). SP1, tendon stem/progenitor cell_1; TC1, tenocyte_1; MTJ, myotendinous junction cell; SM, smooth muscle cell; MC, macrophage; CA, cartilage; TC2, tenocyte_2; SK, skeletal muscle cell; SW, Schwann cell; LC, lymphocyte; NC, neutrophil; EC, endothelial cell; RBC, red blood cell; UC, unspecified cell; SP2, tendon stem/progenitor cell_2; SP3, tendon stem/progenitor cell_3; VEC, vascular endothelial cell.

(B) Dot plot of average gene activity of the indicated genes in each snATAC-seq cluster. The size of the dot reflects the percentage of cells in the cluster that express each gene.

(C) UMAP visualization and predicted annotation of 2-week snATAC-seq after integration and label transfer of 2-week snRNA-seq data.

(D) Identification of matching cell clusters between the 2-week snRNA- and 2-week snATAC-seq data from visualized as heatmap. The heatmap shows the proportions of cells from each snATAC-seq cluster across all sample conditions assigned to each snRNA-seq cluster as part of the label-transfer process.

(E) Violin plot of tenocytes and TSPC-related gene expression in each cluster.

Figure 3. Trajectory analysis and peak visualization of snRNA-seq and snATAC-seq data for the tendon and tendon stem/progenitor cell-related cluster.

(A) UMAP representation of snRNA-seq differentiation trajectory of tenocytes and TSPC lineage and pseudotime-dependent gene expression changes of *Tppp3*, *Cd55*, *Cd248*, *Mkx,* and *Scx,* as inferred using Monocle3.

(B) UMAP representation of snATAC-seq differentiation trajectory of tenocytes and the TSPC lineage and pseudotime-dependent gene expression changes, as inferred using Cicero.

(C) Coverage plots of *Mkx*, *Scx*, *Cd55,* and *Cd248*. Selected peaks that differ across each cluster are highlighted.

Figure 4. Transcription factor landscapes of 2-week mouse Achilles tendons.

SCENIC analysis of transcription factor activity based on 2-week snRNA-seq data for tenocytes and the TSPC lineage (Left). Validation was performed based on the gene activity and motif activity of 2-week snATAC-seq and gene expression of 2-week snRNA-seq (right).

Figure 5. *In vitro* analysis of CD55+/CD248+ TSPCs.

(A) Immunohistochemical image of 2-week mouse Achilles tendons. CD55 and CD248, green; DAPI, blue.

(B) Schema of the *in vitro* assessment of the capacity of CD55+/CD248+ TSPCs as the

differentiation toward tenocytes.

(C) Colony-forming efficiency of CD55+/CD248+ and CD55-/CD248- (negative) TSPCs.

Colonies were stained with crystal violet (n = 6). CD55+/CD248+ TSPC exhibited higher

clonogenic capacity. Data are presented as means ± SEM. **$p < 0.01$.

(D) Morphological changes of CD55+/CD248+ and negative TSPCs after tenogenic induction.

(E) Quantitative PCR of tendon-related gene expression in CD55+/CD248+ and negative

TSPCs after tenogenic induction (n = 3). Data are presented as means ± SD. **$p < 0.01$, *p

< 0.05.

(F) SEM and TEM imaging of artificial tendons derived from CD55+/CD248+ and negative

TSPCs. Data are presented as means ± SD. **$p < 0.01$, *$p < 0.05$.

(G) Proportions of fiber alignment for each artificial tendon (n = 4). Data are presented as

means ± SEM. ***$p < 0.005$, **$p < 0.01$, *$p < 0.05$.

(H) Diameter of collagen fiber in each artificial tendon based on TEM imaging (n = 4). Data

are presented as means ± SD. ***$p < 0.005$.

(I) Tensile strength (MPa) of each artificial tendon (n = 5). Data are presented as means ±

SEM. ***$p < 0.005$.

Table1: Estimated TF activity in each gene

Supplemental Figure 1. Comparison of 2w- and 6w-scRNA seq results.

(A) Violin plot of gene expression enriched in the limb bud (Development (2014) 141, 3683-

3696) in each cluster.

(B) Gene Ontology (GO) terms associated with genes with upregulated expression in the

SP2 cluster.

(C) Number of inferred interactions and interaction strength of genes expressed within 2w-

and 6w-mouse scRNA-seq data.

Supplemental Figure 2. Analysis of 2-week snRNA-seq data.

(A) UMAP snRNA-seq clustering of cells harvested from 2-week mouse Achilles tendons.

(B) Dot plot of average gene expression levels of the indicated genes in 2-week snRNA-seq

clusters. RB, ribosomal RNA; TC1, tenocyte_1; SP1, stem/progenitor cell_1; SP2,

stem/progenitor cell_2; SM, smooth muscle cell; MTJ, myotendinous junction cell; CA,

cartilage; SP3, stem/progenitor cell_3; LC, lymphocyte; MC, macrophage; RBC, red blood

cell; SW, Schwann cell; EC, endothelial cell; TC2, tenocyte_2; PC, proliferating cell; NC,

neutrophil; UC, unspecified cell.

(C) Violin plot of tenocytes and TSPC-related gene expression in each cluster.

(D) Feature plot of TSPC-related gene expression.

1006

Supplemental Figure 3. Analysis of 6-week snATAC-seq.

(A) UMAP snATAC-seq clustering of cells harvested from 6-week mouse Achilles tendons.

Annotation was based on gene activity (ground-truth annotation, left) and the predicted

annotation inferred from 2-week snATAC-seq (right).

(B) Identification of matching cell clusters between the 2-week and 6-week snATAC-seq data,

visualized as a heatmap.

(C) Violin plot of tenocytes and TSPC-related gene activity in each cluster.

1014

Supplemental Figure 4. Analysis of 6-week snRNA-seq.

(A) UMAP snRNA-seq clustering of cells harvested from 6-week mouse Achilles tendons.

Annotation was based on each gene expression (ground-truth annotation, left) and predicted

annotation inferred from 2-week snRNA-seq (right).

(B) Identification of matching cell clusters between the 2-week and 6-week snRNA-seq data

visualized as a heatmap.

(C) Violin plot of tenocytes and TSPC-related gene expression in each cluster.

1022

1023   **Supplemental Figure 5. Comparison of 2-week and 6-week snATAC-seq.**

1024   (A) Circle plot of annotated differentially accessible regions for each data.

1025   (B) Integrated UMAP snATAC-seq clustering of cells harvested from 2-week and 6-week

1026   mouse Achilles tendons.

1027   (C) Feature plot of TSPC-related gene activity in each dataset.

1028

1029   **Supplemental Figure 6. Gene expression changes in CD55+/CD248+ and negative**

1030   **TSPCs.**

1031   Quantitative PCR of gene expression in CD55+/CD248+ and negative TSPCs (n = 4). Data

1032   are presented as means ± SEM. *$p < 0.05$, ***$p < 0.005$.

1033

1034   **Supplemental Figure 7. Chondrogenic and osteogenic induction of CD55+/CD248+ and**

1035   **negative TSPCs.**

1036   Quantitative PCR of tendon-related genes in CD55+/CD248+ and negative TSPCs after

1037   chondrogenic and osteogenic induction (n = 4). Data are presented as means ± SEM. **$p <$

1038 0.01, *$p < 0.05$.

Figure 1

Figure 2

Figure 3

Figure 4

Estimated TF activity
(stem/progenitor cell and tenocytes lineage)

Gene activity, motif activity and gene expression

Figure 5

Table1

Estimated TF activity : Cd55	
TF	**spearman correlation**
Klf4	0.264195541
Klf3	0.262413411
Zeb1	0.219593135
Pbx1	0.204085218
Hic1	0.202481054
Klf2	0.196232716
Nfe2l2	0.196076187

Estimated TF activity : Cd248	
TF	**spearman correlation**
Hic1	0.375565336
Klf3	0.277768834
Zmiz1	0.272991765
Klf4	0.260526766
Zeb1	0.250583629
Sp3	0.2502976
Kdm5b	0.193229132

Estimated TF activity : Mkx	
TF	**spearman correlation**
Creb3l1	0.320638653
Zfhx3	0.169341424
Mxi1	0.149048856
Bhlhe40	0.144301452
Ets2	0.137445497
Max	0.116134957
Elk3	0.098325126

Estimated TF activity : Scx	
TF	**spearman correlation**
Creb3l1	0.316522613
Npdc1	0.163452737
Ets2	0.145463784
Mix1	0.121248221
Bhlhe41	0.115930755
Elk3	0.10803805
Bmyc	0.104968457

Supplemental figure 1

Supplemental figure 2

Supplemental figure 3

A

6w: snATAC-seq (ground-truth annotation)

6w: snATAC-seq (predicted annotation)

B

Predicted annotation

Ground-truth annotation

Proportion of 6w_snATAC-seq cells
assigned to 2w_snATAC-seq cluster

C

Gene activity in 6w_snATAC-seq

Mkx · *Scx* · *Tnmd* · *Ly6a* · *Tppp3* · *Pdgfra* · *Cd55* · *Cd248* · *Cd34*

Supplemental figure 4

A

6w: snRNA-seq (ground-truth annotation)

6w: snRNA-seq (predicted annotation)

B

Predicted annotation

C Gene expression in 6w_snRNA-seq

Proportion of 6w_snRNA-seq cells
assigned to 2w_snRNA-seq cluster

Supplemental figure 5

Supplemental figure 6

Supplemental figure 7

chondrogenic

osteogenic

relative expression(/β2MG)

: CD55/CD248, induction+

: CD55/CD248, induction-

: negative, induction+

: negative, induction-