Title

Single-nucleus transcriptional and chromatin accessibility analyses of maturing mouse Achilles tendon uncover the molecular landscape of tendon stem/progenitor cells

Authors

Hiroki Tsutsumi[1#], Tomoki Chiba[1#], Yuta Fujii[1,2], Takahide Matsushima[1], Tsuyoshi Kimura[3,4], Akinori Kanai[5], Akio Kishida[3], Yutaka Suzuki[5] and Hiroshi Asahara[1,2*]

Affiliations

1 Department of Systems BioMedicine, Graduate School of Medical and Dental Sciences, Institute of Science Tokyo, Tokyo, Japan

2 Department of Molecular and Cellular Biology, The Scripps Research Institute, La Jolla, USA

3 Laboratory for Biomaterials and Bioengineering, Institute of Integrated Research, Institute of Science Tokyo, Tokyo, Japan

4 Department of Biomedical Engineering, Faculty of Life Science, Toyo University, Saitama, Japan

5 Laboratory of Systems Genomics, Department of Computational Biology and Medical Sciences, Graduate School of Frontier Sciences, the University of Tokyo, Chiba, Japan

#: These authors contributed equally to this work.

Contact info

*Correspondence: asahara@scripps.edu

Abstract

Tendons and ligaments are crucial connective tissues linking bones and muscles, yet achieving full functional recovery after injury remains challenging. We investigated the characteristics of tendon stem/progenitor cells (TSPCs) by focusing on the declining tendon repair capacity with growth. Using single-cell RNA sequencing on Achilles tendon cells from 2- and 6-week-old mice, we identified *Cd55* and *Cd248* as novel surface antigen markers for TSPCs. Combining single-cell RNA sequencing with single-nucleus RNA and ATAC sequencing analyses revealed that *Cd55* and *Cd248* positive fractions in tendon tissue represent TSPCs, as confirmed by their expression of established TSPC markers, with this population decreasing at 6 weeks. We also identified candidate upstream transcription

factors regulating these fractions. Functional analyses of isolated CD55/CD248 positive cells

demonstrated high clonogenic potential and tendon differentiation capacity, forming

functional tendon-like tissue *in vitro*. This study establishes CD55 and CD248 as novel TSPC

surface antigens, potentially advancing tendon regenerative medicine and contributing to the

development of new treatment strategies for tendon and ligament injuries.

Introduction

Tendons are connective tissues involved in the contractile movement of muscle to bone.

Tendons are rich in extracellular matrix components, such as type 1 collagen and

proteoglycans, which both have elastic and viscous properties to withstand overload

(Andarawis-Puri et al., 2015; Asahara et al., 2017; Connizzo et al., 2013).

Tendons have a hierarchical structure. Collagen fibrils aggregate to form fibers, which then

bundle into fascicles. Each fascicle is surrounded by a thin connective tissue layer called the

endotenon. Fascicles are further enclosed by the epitenon, and in some cases, an additional

outer layer called the paratenon. The number of fascicles and the presence of a paratenon

can vary among different tendons and species (Jozsa et al., 1991; Walia et al., 2019).

Tendon and ligament injuries potentially account for the majority of musculoskeletal

disorders and can lead to arthritis and spondylitis. (Gracey et al., 2020). The most popular

treatment approaches are surgery and conservation (Steinmann et al., 2020). However, full

functional recovery is often not achieved due to scarring and fibrosis in the injured area

(Guilak et al., 2014; Thomopoulos S et al., 2015). Furthermore, the risk of postoperative

rupture is high (Leong et al., 2020; Loiacono et al., 2019). In this context, tendon regeneration

focused on tendon stem/progenitor cells (TSPCs) is attracting attention (Leong et al., 2016).

Tendons have few cellular components, including tendon cells (tenocytes) and TSPCs (Tang

et al., 2016; Bi et al., 2007). Tenocytes are responsible for tendon homeostasis, while TSPCs

self-renew and differentiate into tenocytes. Thus, investigating TSPCs biology is important

for understanding tendon regeneration and homeostasis.

Murine TSPCs were first reported in 2007 by Bi Y et al. (Bi et al., 2007). They

demonstrated that TSPCs possess self-renewal capacity, colony-forming ability, and multi-

differentiation potential *in vitro*. Furthermore, murine TSPCs were reported to

simultaneously express stem cell markers such as *Cd44* and *Stem cells antigen-1* (*Ly6a*),

as well as tendon-related genes like *Scleraxis* (*Scx*) and Collagen type I alpha 1 chain

(*Col1a1*), and were identified as a subset of tendon cells within the tendon fascicle.

Subsequent research on murine TSPC localization has yielded diverse perspectives.

Harvey et al. reported that *Tppp3*/*Pdgfra*-positive cells in the epitenon are induced upon

tendon injury and contribute to repair (Harvey et al., 2019). Additionally, Yin et al. and

Tempfer et al. demonstrated the presence of TSPCs around blood vessels (Yin et al., 2016;
Tempfer et al., 2009).
Markers used to characterize murine TSPCs include CD73, CD105, and CD90, which are
known criteria for mesenchymal stem cells (MSCs) (Lui et al., 2015). Additionally, TSPCs
have been reported to express *POU class 5 homeobox 1* (*Oct-4*), *Nanog*, *Nucleostemin*,
stage-specific embryonic anitigen-4 *(SSEA-4)*, *c-Myc*, *SRY-box transcription factor (Sox*2),
Fucosyltransferase 4 (*Fut-4*), and other genes (Zhang et al., 2016). Expression of CD146
and CD44 has been confirmed as well (Ruzzini et al., 2014). The *Tppp3*/*Pdgfra*-positive
TSPCs reported by Harvey et al. highly express *Cd34*, which is generally considered to
have low expression in MSCs (Harvey et al., 2019; Tachibana, et al., 2022). *Cd34* is known
to be highly expressed in mouse embryonic limb buds at E14.5 compared to E11.5 (Havis
et al., 2014), suggesting that *Cd34* positive cells might reflect progenitor cells that
constitute the limb bud, including tendons. However, these markers are not specific to
TSPCs, thus, a definitive method to distinguish TSPCs from mature tendons *in vivo* has not
been established yet (Li et al., 2019; Wang et al., 2008; Chen et al., 2012; Cho et al., 2018;
Fang et al., 2022).
Regarding tendon regenerative capacity, Howell et al. reported an interesting observation.
Tendons in juvenile mice can regenerate functional tissue after injury, but this ability is lost
in mature mice, resulting in scar tissue formation (Howell et al., 2017). This finding
suggests the possibility of abundant TSPCs in juvenile mouse tendons.
Given this background, we hypothesized that evaluating tendon tissue heterogeneity at
the single-cell level using juvenile mouse tendons would enable a more detailed
characterization of TSPCs. This approach is expected to advance the identification and
characterization of TSPCs, which has been challenging with conventional methods.
Therefore, we performed scRNA-seq using cells collected from 2- and 6-week-old mouse
Achilles tendons and investigated clusters that co-express known TSPCs markers, such as
Tppp3, *Pdgfra*, and *Ly6a*. Then, we analyzed the expression of surface antigens in these
clusters and identified *Cd55* and *Cd248* as novel candidate surface antigens for murine
TSPCs.
Furthermore, snATAC-seq and snRNA-seq were performed simultaneously using cells
collected from Achilles tendons to evaluate the validity of *Cd55* and *Cd248* as surface
antigens for TSPCs and to identify the landscape of transcription factors (TFs) involved in
tendon maturation. We sorted mouse Achilles tendon cells based on CD55 and CD248 and
confirmed their phenotypes, demonstrating high clonogenicity and highly efficient induction
into tendon cells. These results suggested that CD55 and CD248 are novel surface antigens
of murine TSPCs and may be useful for understanding the process of tendon maturation and

for applications in cell therapy.

Material and Methods

Single cell isolation

Mice were euthanized by exposure to CO_2 followed by cervical dislocation. To remove Achilles tendons, a longitudinal incision through the skin was made down the midline of the posterior aspect of the lower limb. A sharp transverse incision was made just distal to the myotendinous junction and again just proximal to the enthesis, and the Achilles tendon was carefully removed. The procedure was performed bilaterally. All animals were purchased from Sankyo Lab Service (Tokyo, Japan). Achilles tendons were digested to obtain a single-cell suspension. In total, 40 or 60 (2-week and 6-week, respectively) mouse Achilles tendons were processed together as a single sample. After washing with 1× phosphate-buffered saline (PBS) several times and cutting 1-mm-wide sections using scissors, tendons were digested for 1 h at 37°C with continuous shaking at 1,200 rpm and 37°C in a dissociation solution consisting of 30 mg/mL collagenase (Wako, Osaka, Japan). After digestion, the single-cell suspension was filtered for debris using a 40 µm cell strainer and washed twice with 1× PBS. The samples were centrifuged for 15 min at 1,500 rpm and 4°C. Dead cells were removed using a Dead Cell Removal Kit (Miltenyi Biotec, Bergisch Gladbach, Germany). Cell viability was 76% (2-week) and 73% (6-week), respectively. The experiment involving mice was approved by Animal Experimental Committee of the Institute of Science Tokyo.

scRNA-seq library construction and sequencing

Cells isolated from Achilles tendons of both 2-week-old and 6-week-old mice were processed separately. Cells were resuspended in PBS with 1% fetal bovine serum (FBS) at a concentration of 10,000 cells per µL. Then, 10,000 cells per sample were loaded on a Chromium Controller (10x Genomics, Pleasanton, CA, USA) for single cell capture. Libraries were prepared using Single Cell 3′ Library & Gel Bead Kit v3 (10x Genomics) following the manufacturer's instructions. A single-cell emulsion (Gel Bead-In-EMulsions, GEMs) was created by making barcoded cDNA unique to each individual emulsion. A recovery agent was added to break GEM and cDNA was then amplified. A library was produced via end repair, dA-tailing, adapter ligation, post-ligation cleanup with SPRIselect, and sample index PCR. The quality and concentration of the amplified cDNA were evaluated using the Bioanalyzer (Agilent 2100) on a High Sensitivity DNA chip (Agilent, #5065-4401; Santa Clara, CA, USA). Sequencing was performed using the HiSeq X system (Illumina, San Diego, CA, USA) to generate 28/90 bp paired-end reads.

Cell dissociation, nuclei isolation, and snRNA-seq and snATAC-seq library

construction and sequencing

After obtaining cells from the mouse Achilles tendons of 2-week-old and 6-week-old mice (processed separately for each age group), nuclei were isolated following the Chromium Next GEM Single Cell Multiome ATAC + Gene Expression Reagent Bundle (10XGenomics) and all of the buffers were made according to the manufacturer's instructions. Briefly, after centrifugation, the supernatant was discarded, and the cells were re-suspended in 1 mL of 1% FBS in PBS. Approximately 500,000 cells were transferred to a new tube for further lysis. To remove the supernatant, cells were centrifuged again for 5 min at 300 rcf and 4°C and resuspended in 100 mL of chilled lysis buffer. Cells were lysed for 9 min on ice and 1 mL of wash buffer was added to stop the reaction. Then, the suspension was filtered through a 40 μm cell strainer, cells were centrifuged and resuspended in 66.2 mL of chilled Diluted Nuclei Buffer aiming to target 7000 nuclei. The final concentration of nuclei was determined, followed by transposition, GEM generation, barcoding, and library construction according to the Chromium Next GEM Single Cell Multiome ATAC + Gene Expression (10XGenomics). Libraries were sequenced with the parameters recommended by the manufacturer, using the NovaSeq 6000 (Illumina) to generate 28/90 bp paired-end reads for gene expression and 50/49 bp paired-end reads for ATAC sequencing.

scRNA-seq and snRNA-seq analyses

Sequencing reads were processed with the Cellranger_arc (10X Genomics, v2.0.0) using the mouse reference mm10. We performed separate analyses for scRNA-seq data (collected from both 2-week and 6-week samples) and snRNA-seq data (collected from both 2-week and 6-week samples) to enable comprehensive characterization of transcriptional landscapes. From the gene expression matrix, downstream analyses were carried out using R. Quality control, filtering, data clustering and visualization, and a differential expression analysis were carried out using Seurat (Butler et al., 2018). For each dataset, cells with unique feature counts over 2,500 or less than 200 and >5% mitochondrial counts were filtered. Then, heterotypic doublets (assuming 5% of barcodes represent doublets) were removed using DoubletFinder (McGinnis et al., 2019).

Unsupervised shared nearest neighbor (SNN) clustering was performed with varying resolution and the results were visualized using uniform manifold approximation and projection (UMAP) (Becht et al., 2019). Differentially expressed genes (DEGs) among each cell cluster were determined using the FindAllMarkers function in Seurat. The criteria of DEGs as a marker gene for the cluster was logFC > 0.25, adjusted $p < 0.05$, expression in

Pseudotime analysis

187 Monocle3 (Trapnell et al., 2014) was used to convert the snRNA-seq dataset into a cell

188 dataset object (CDS), preprocess data, correct for batch effects, embed with dimensional

189 reduction, and perform pseudotemporal ordering. Cicero (Pliner et al., 2018) was used to

190 generate pseudo-temporal trajectories for the snATAC-seq dataset.

Cell-cell communication analysis

193 Intercellular communication networks were quantitatively inferred and analyzed using

194 scRNA-seq data. The R package CellChat (Jin et al., 2021) was used to visualize the

195 interactions among different cell groups. Two hundred twenty-nine signaling pathway

196 families were grouped as a library to analyze cell–cell communication.

GO analysis

199 The R package ClusterProfiler (Yu et al., 2012) was used to perform a gene enrichment

200 analysis. The p-values were corrected by the Benjamini & Hochberg method.

Motif enrichment analysis (ChiP seeker)

203 Genomic regions containing snATAC-seq peaks were annotated using ChIPSeeker (Yu et

204 al., 2015) and the UCSC database on mouse (mm10).

snATAC-seq analysis

The Cell Ranger ATAC pipeline (1.2.0) (Satpathy et al., 2019) was used to preprocess the

data resulting from sequencing. First, Tn5 sites were mapped to the mouse reference

transcriptome mm10, and duplicate reads and background cells were removed. This

returned barcoded fragment files, which were loaded into Signac (Stuart et al., 2021) for

downstream analyses using the standard Signac/Seurat pipeline. Macs2 (Zhang et al.,

2008) was run on the fragment files to call peaks using the Signac 'CallPeaks' function.

Fragments were mapped to the Macs2 called peaks and assigned to cells using the Signac

'FeatureMatrix' function. Nucleosome signal strength and transcription start site (TSS)

enrichment for each cell were calculated using the Signac 'NucleosomeSignal' and

'TSSEnrichment' functions, respectively. Outliers in the QC metric categories were removed

as per Signac's standard processing guidelines. Latent semantic indexing (LSI), a form of

dimensional reduction, was performed using the Signac 'RunTFIDF' and 'RunSVD'

functions. The UMAP hyperparameters were varied to produce consistent object shapes

(using R). Once hyperparameters were chosen, the Signac/Seurat's 'RunUMAP' function

was run on the LSI dimensions chosen earlier for UMAP embedding. The Signac/Seurat

'FindNeighbors' function was run using the same LSI dimensions used for UMAP to

compute the nearest neighbor graph. Signac/Seurat 'FindClusters' was then run at varying

resolutions. A gene activity matrix was constructed by counting ATAC peaks within the gene

body and 2 kb upstream of the transcriptional start site for protein-coding genes annotated

in the Ensembl database. The gene activity matrix was log-normalized prior to label transfer

with the aggregated snRNA-seq Seurat object using a canonical correlation analysis. The

aggregated snATAC-seq object was filtered using a 97% confidence threshold for cell-type

assignment following label transfer to remove heterotypic doublets. The filtered snATAC-

seq object was reprocessed with TFIDF, SVD, and batch effect correction followed by

clustering and annotation based on lineage- specific gene activity. Differential chromatin

accessibility between cell types was assessed with the Signac FindMarkers function for

peaks detected in at least 20% of cells using a likelihood ratio test and a log-fold-change

threshold of 0.25. Bonferroni-adjusted p-values were used to determine significance at an

FDR < 0.05.

Integration of snRNA-seq and snATAC-seq data

snRNA-seq and snATAC-seq data were integrated using the cluster-label transfer

procedure as implemented in Signac and Seurat. Each snRNA-seq sample. as clustered

individually and its cluster labels were projected onto the matching, individually clustered

snATAC-seq sample or vice versa. Anchors were identified for condition-matched snRNA-

and snATAC-seq samples using the FindTransferAnchors function and a canonical

correlation analysis (CCA) was performed using the snRNA expression values and the

snATAC-imputed gene expression values. The anchors were used to transfer cluster-label

identifiers between the two data types using the TransferData function. Each cell in the

query was assigned the cluster label with the highest prediction score, and label transfer

was considered successful for query cells with prediction scores above 0.3.

SCENIC analysis

To identify TFs and characterize cell states, a cis-regulatory analysis was performed using

the R package SCENIC (Lake et al., 2018), which infers gene regulatory networks based

on co-expression and DNA motif analyses. The network activity was then analyzed for each

cell to identify recurrent cellular states. In short, TFs were identified using GENIE3 and

compiled into modules (regulons), which were subsequently subjected to cis-regulatory

motif analysis using RcisTarget with two gene-motif rankings: 10 kb around the TSS and

500 bp upstream. Regulon activity in every cell was then scored using AUCell.

Immunohistochemistry

Immunohistochemistry was performed as previously described (Tsutsumi et al., 2022).

Briefly, tissue samples were fixed in 4% paraformaldehyde overnight at 4°C and embedded

in paraffin. Sections of 10 μm in thickness were deparaffinized and activated with citric acid

buffer (10 mM sodium citrate, 1 mM EDTA; pH 6.0) at 80 °C for 60 min in a decloaking

chamber NxGen (BIOCARE Medical, CA, USA) or with 0.1% trypsin/PBS at 37 °C for

30 min. After they were blocked with Blocking One solution (Nacalai tesque, Kyoto, Japan)

for 60 min, they were incubated with rabbit anti-CD55 antibody (1:100; A13918, ABclonal,

Woburn, MA, USA) or rat anti-CD248 (1:100; MAB7535, R&D Systems, Minneapolis, MN,

USA) overnight at 4 °C. Following this step, they were incubated with Alexa Fluor plus 488

goat anti-rabbit antibody (1:1000; A32731, Thermo Fisher Scientific , Waltham, MA, USA)

or Alexa Fluor plus 488 donkey anti-rat antibody (1:1000; A21208, Thermo Fisher Scientific)

for 60 min. Hoechst 33342 (100ng/ml, Thermo Fisher Scientific) was added during this

incubation. The sections were then mounted with ProLong Glass Antifade Mountant

(P36980, Thermo Fisher Scientific).

Fluorescence-activated cell sorting (FACS)

Cells harvested from the Achilles tendon were incubated for 60 min on ice with APC-CD55

(#131812; BioLegend, San Diego, CA, USA) and a primary antibody against CD248 (#LS-

B2712, LSBio) using a FACS buffer (PBS containing 1 [v/v%] FBS). After washing, cells

were reacted with fluorescent-conjugated rabbit secondary antibody. Cell sorting was

performed using MoFlo XDP FACS (Beckman Coulter, Brea, CA, USA). DAPI was used as

the live/dead discriminator to the gate. CD55/CD248 dual positive and negative cell cluster

were sorted.

Primary cell culture

Primary TSPCs were isolated from the Achilles tendons of 2-week-old mice with

collagenase digestion described above. Single-cell suspensions were cultured in the

culture medium (MEMα + 20% FBS + 1% penicillin-streptomycin + 1 [v/v%] 100× non-

essential amino acid solution [NEAA; Gibco], 1 [v/v%] 100× GlutaMAX [Gibco]). At 80%–

90% confluence, cells were trypsinized, centrifuged, resuspended in culture medium as

passage 1 cells, and incubated in 5% CO_2 at 37°C, with fresh medium every 2–3 days.

Colony formation assay

For the colony formation assay, single-cell suspensions of TSPCs (1000 cells/well) were

seeded and incubated in six-well plates for 12–14 days in the growth medium and fixed

with 4% paraformaldehyde (PFA) (Sigma-Aldrich, St. Louis, MO, USA). Then, 0.1% crystal

violet solution (Wako) was used to stain the cells. Colonies of >30–50 cells were defined as

a single colony unit, and the number of clusters was counted using the ImageJ package

ColonyArea (Guzmán et al., 2014)

RT-PCR

Total RNA was purified using TRIzol reagent (Invitrogen, Grand Island, NY, USA). Reverse

transcription of mRNA was carried out using the PrimeScript RT Reagent Kit (Takara,

Shiga, Japan). Quantitative PCR (qRT-PCR) was performed on cDNA with the Thunderbird

SYBR mix (Toyobo, Osaka, Japan). B2m was used as a reference gene, and relative gene

expression levels were calculated through the ΔCT method.

Tenocytes, cartilage, and osteocyte differentiation

For the differentiation experiment, TSPCs were cultured in 6-well plates (50,000

cells/well). Osteogenic, chondrogenic, and tenogenic differentiation were induced using a

corresponding differentiation medium. The osteogenic differentiation medium contained a

culture medium supplemented with 10 nM dexamethasone (Sigma-Aldrich), 5 mM β-

glycerophosphate (APEXBIO), and 0.05 mM L-ascorbic acid 2-phosphate (Sigma-Aldrich).

The chondrogenic differentiation medium contained a culture medium supplemented with

100 nM dexamethasone and 10 ng/mL BMP2 (Sigma-Aldrich). The tenogenic differentiation

medium contained a culture medium supplemented with 10 ng/mL TGF-β1 (Peprotech,

Rocky Hill, NJ, USA), 10 ng/mL GDF-5 (R&D Systems, Minneapolis, MN, USA), and 0.05

mM L-ascorbic acid 2-phosphate. After a 2 week induction period, cells were harvested for

RT-PCR.

Bio-cultured tendon (Bio-tendon)

Generation of Bio-tendon has reported previously (Kataoka et al., 2020; Tsutsumi et al.,

2022) Briefly, sorted cells were embedded in a 3D-culture cocktail, which is consistent with

2 mg/mL Cellmatrix (Type I-A, Nitta Gelatin Inc., Osaka, Japan), pro-survival cocktail (final

concentrations: 100 nM Bcl-Xl BH4 4-23 (Merck Millipore, Burlington, MA, USA), 100 μM

carbobenzoxy-valyl-alanyl-aspartyl-[*O*-methyl]-fluoromethylketone (Z-VAD-FMK) (Promega,

Madison, WI, USA)) in culture medium. The 3D chamber was coated with 1% gelatin and

incubated at 37°C and 5% CO_2 for 30 min. After washing with 1× PBS 3 times, 1.0×10^6

cells were transferred to a 3D-culture cocktail mixture and incubated at 37°C and 5%

CO_2 for 60 min for gelation. Following gelation, tenogenic differentiation medium was added

to the chamber. Following 24 h of incubation, the chamber was placed within a cell

stretching system (Shellpa Pro, Menicon Life Science, Aichi, Japan). Cyclic mechanical

stretch was applied for 1 week, with gradual increase in stretch load: 1% on day 1, 2% on

day 2, 3% on day 3, 4% on day 4, and 5% from days 5 to 7. The stretching cycle was

programed at 0.25 Hz for 18 h/day, followed by resting for 6 h/day at 37°C and 5% CO_2.

Daily medium changes were also required.

Decellularization by HHP and chemical treatment

The cultured bio-tendon was placed into a plastic pack filled with saline and sealed to

prevent implosion and leakage during the procedure. The pack was then pressurized at

1,000 MPa at 30°C for 10 min using an HHP machine (Dr. CHEF; Kobe Steel, Hyogo,

Japan). After pressurization, the bio-tendon was washed thrice with 30% ethanol (EtOH) by

continuous shaking for 5 min at each step. Finally, 1-ethyl-3-(3-dimethylaminopropyl)

carbodiimide (EDC)/*N*-hydroxysuccinimide (NHS)-based cross-linking was performed by

adding 70 mM EDC (Wako) and 70 mM NHS (Wako) with 30% EtOH for 24 h at 4°C. The

processed bio-tendon was incubated in 1× PBS at 4°C until the experiment.

Electron microscopy

Bio-tendon tissues were dissected and fixed overnight in 2.5% glutaraldehyde in 0.1 M

phosphate buffer (PB). For transmission electron microscopy (TEM), the specimens ($n = 3$)

were rinsed with 0.1 M PB, post-fixed in 1% osmium buffered with 0.1 M PB for 2 h, and

dehydrated in a graded ethanol series. The specimens were then embedded in Epon 812,

sectioned into ultrathin sections (70 nm), mounted on copper grids, and double-stained with

uranyl acetate and lead citrate. TEM observation was conducted JEM-1400Flash (JEOL,

Tokyo, Japan). For scanning electron microscopy (SEM), the specimens ($n = 2$) were dried

in a critical-point dryer (HCP-2; Hitachi, Tokyo, Japan) with liquid CO_2 and coated with

platinum. The specimens were subsequently observed under SEM (JSM-7900F/JED-2300;

JEOL). Fiber orientation within the microscopic sections was analyze using the OrientationJ

image processing tool, a plugin for ImageJ.

Stretch test

The mechanical properties were measured using a creep meter (RE-3305S; Yamaden,

Tokyo, Japan). After measuring the initial length (mm), diameter (mm), and thickness (mm)

using a micrometer, the samples were fixed with two grips, which were pulled at a constant

speed of 0.05 mm/s until failure, and the tensile strength (N) and failure strain (mm) were

measured. The cross-sectional area (mm^2) was calculated using the initial diameter and

thickness. The stiffness was manually determined from the slope in the linear region of the

failure-stress curve.

The tensile strength (MPa), failure strain (%), stiffness, and Young's modulus were

calculated using the following formulas:

Tensile strength (MPa) = tensile strength (N)/cross-sectional area (mm^2).

Failure strain (%) = failure strain (mm)/initial length (mm).

Stiffness = Stress (N)/Strain (mm)

Young's modulus = stiffness × initial length (mm)/cross-sectional area (mm^2)

Statistical analysis

All values are presented as means ± SEM. Statistically significant differences were

assessed using unpaired two-tailed Student's *t*-tests and one-way analysis of variance

(ANOVA) with Tukey's post hoc tests. Statistical significance was set at $p < 0.05$.

Results

Identification of Novel Surface Antigens for Tendon Stem/Progenitor Cells (TSPCs)

In mice, the healing capacity for tendon injuries is high up to 2 weeks of age, but

decreases as the musculoskeletal system matures at 6 weeks (Howell et al., 2017). To

investigate whether this phenomenon is due to changes in the cellular population of tendon

tissue, including fluctuations in progenitor cells, we performed single-cell RNA sequencing

on mouse Achilles tendons at these two time points.

Achilles tendons were harvested from mice at 2 and 6 weeks of age. Following

collagenase digestion and dead cell removal, we employed massively parallel, droplet-

enabled scRNA-seq analysis (10X Genomics Chromium). Data processing was conducted

using the Cell Ranger pipeline (10X Genomics). We analyzed 10,314 cells from 2-week-old

mice (median 3,167 genes/cell and 41,611 mean reads/cell) and 6,513 cells from 6-week-

old mice (median 732 genes/cell and 60,386 mean reads/cell). After doublet removal using

Doubletfinder, we merged the datasets and performed unbiased clustering using Seurat,

identifying a total of 15 clusters (Fig. 1A). The top differentially expressed genes (DEGs) for

each cluster, based on log2 fold change and statistical significance, are summarized in Fig. 1B.

We identified two clusters (0 and 11) with high expression of *Mkx* and *Scx*, transcription factors involved in tendon development and growth. Cluster 0 showed higher expression of ECM-related genes such as *Col1a1* and *Fmod* compared to cluster 11, suggesting these were tenocytes (TC). Cluster 11, with high *Periostin* (*Postn*) expression (Jacobson, et al., 2020), was classified as myotendinous junction (MTJ). Other clusters were characterized as follows: cartilage (CA, high *SRY-box transcription factor 9* (*Sox9*) expression), lymphocytes (LC1, LC2, high *Lysozyme 2* (*Lyz2*) and *Complement component 1, q subcomponent, alpha polypeptide* (*C1qa*)), endothelial cells (EC, high *Platelet and endothelial cell adhesion molecule 1* (*Pecam1*)), red blood cells (RBC, high *Hemoglobin alpha, adult chain 1* (*Hba-a1*)), smooth muscle cells (SM, high *Myosin light chain 9* (*Myl9*)), proliferating cells (PC, high *Mki67* and *Stathmin 1* (*Stmn1*)), Schwann cells (SW, high *Myelin basic protein* (*Mbp*)), vascular endothelial cells (VEC, high *Multimerin 1* (*Mmrn1*)), and macrophages (MC, high *Protein tyrosine phosphatase receptor type C* (*Ptprc*)). In cluster3, *Col1a1* and *Fmod* were expressed; however, *Gm42418* and *AY036118* were highly expressed. These long non-coding RNAs are related to RN45s and therefore cluster3 represents ribosomal contamination (Isola et al., 2024). It is also speculated that the cartilage cluster (CA) reflects enthesis, with expression of *Scx* as well as *Sox9* (Zhan et al., 2023). DEG s of each cluster was summarized in supplementary table1-15.

To identify progenitor populations within these clusters, we analyzed expression patterns of previously reported markers *Tppp3* and *Pdgfra* (Harvey et al., 2019; Tachibana, et al., 2022), along with the known stem/progenitor cell marker *Ly6a* (Holmes et al., 2007; Sung et al., 2008; Hittinger et al., 2013; Sidney et al., 2014; Fang et al., 2022) (Supplemental Figure 1B). We identified subclusters within clusters 1 and 4 showing high expression of these genes, which we defined as SP1 and SP2. SP2 exhibited the highest expression of these genes, suggesting it had the strongest progenitor characteristics. It has been reported (Harvey et al., 2019) that the *Tppp3*-positive population is localized to peritenon, and SP clusters might reflect peritenon as well.

SP2 also showed strong expression of genes associated with early tendon development in mouse embryos, as identified by RNA-seq of mouse limb tendon cells at E14.5 (Havis et al., 2014) and the Eurexpress mouse embryo transcriptome atlas database (Diez-Roux et al., 2011) (Supplemental Figure 2A). Gene Ontology analysis revealed upregulation of tendon development-related pathways in SP2, including TGFβ production and collagen-containing extracellular matrix (Supplemental Figure 2B). Evaluation of cell numbers in

each cluster showed a decrease in SP2, TC, and MTJ at 6 weeks (Fig. 1C, Supplemental
Figure 1A).

We then examined DEGs in SP2 compared to SP1 (Fig. 1D). Whole list of DEGs
comparing SP2 and SP1 was summarized in supplementary table16. Among these, we
identified *Cd34*, *Cd55*, and *Cd248* as genes encoding surface antigens. *Cd34* is known to
be highly expressed in mouse embryonic limb bud E14.5 compared to E11.5 (Havis et al.,
2014), suggesting that *Cd34* positive cluster might reflect a progenitor cell that consist of
the limb bud, including tendons. However, *Cd55* and *Cd248* have not been discussed with
this context. Expression of *Cd55* and *Cd248* was localized to areas corresponding to SP1
and SP2 (Fig. 1E). We evaluated the correlation coefficients between the expression of
these genes and previously suggested TSPC markers (*Cd73*, *Cd90*, *Cd105*, *Cd44*, *Cd146*)
as well as *Tppp3*, *Pdgfra*, and *Ly6a* at 2 weeks of age. *Cd34*, *Cd55*, and *Cd248* showed
high correlation, suggesting these genes as new candidate surface antigens for TSPCs
(Fig. 1F).

Comparing gene expression between 2 and 6 weeks, we observed a decrease in *Mkx* and
Scx expression, consistent with previous reports (Grinstein et al., 2019). Furthermore, the
expression of our candidate tendon progenitor cell markers *Cd34*, *Cd55*, and *Cd248* all
decreased at 6 weeks. Inflammatory cells, including macrophages, increased at 6 weeks
(Fig. 1G). Gene expression analysis revealed a shift from M2 macrophage-related genes
(*Mannose receptor C-type 1* (*Mrc1*) and *Cathepsin C* (*Ctsc*)) at 2 weeks to M1
macrophage-related genes (*Chemokine (C-C motif) ligand 6* (*Ccl6*) and *Chemokine (C-C
motif) ligand 9* (*Ccl9*)) at 6 weeks. Comparative analysis between 2-week and 6-week
tendons was summarized as a heatmap in supplemental Figure 3.

Analysis of cell-cell communication changes in postnatal tendon and surrounding tissues
(Lui et al., 2019) using CellChat (Jin et al., 2021) showed a significant decrease in overall
interaction at 6 weeks compared to 2 weeks. These results demonstrate that the cellular
profile of tendon tissue changes dramatically at 6 weeks, with decreased expression of
stem progenitor markers and reduced interaction with surrounding tissues.

Single Nucleus RNA + ATAC Analysis of 2-Week-Old Mouse Achilles Tendon Cells
Gene regulation is mediated by the binding of transcription factors to cis-regulatory
elements proximal to the gene. Consequently, epigenetic changes such as chromatin
accessibility play a crucial role in gene expression (Miller et al., 2013). Moreover,
transcription factors are typically expressed at low levels, potentially leading to false
negatives in scRNA-seq due to detection limits. Chromatin accessibility changes often
precede gene expression changes, potentially allowing us to predict transcriptional

changes (Ranzoni et al., 2020). Therefore, we performed simultaneous ATAC-seq and

RNA-seq at the single-cell level to identify gene expression changes and their associated

epigenetic alterations, aiming to elucidate the post-natal growth mechanisms of tendon

tissue and evaluate the validity of *Cd55* and *Cd248* as markers.

We extracted nuclei from cells isolated from 2-week-old mouse Achilles tendons and

conducted droplet-enabled multi-omics analysis (10X Genomics Chromium), performing

simultaneous snATAC-seq and snRNA-seq as a reference. We first analyzed the snATAC-

seq data. After mapping reads to the genome and calling peaks, we annotated the location

of peaks in terms of genomic features. The peaks were associated with promoters, introns,

exons, and intergenic regions.

We evaluated 6,571 cells (median high-quality fragments per cell: 17,408) and clustered

them using latent semantic indexing and Uniform Manifold Approximation and Projection

(UMAP) with the R package Signac (Stuart et al., 2021) (Figure 2A). Given the limited

knowledge of cell-specific chromatin accessibility, we assessed gene activity by computing

counts per cell within the gene body and promoter of protein-coding genes. Using this gene

activity data, we performed unsupervised clustering and identified 17 clusters (ground-truth

annotation). Clusters expressing tendon-related genes such as *Mkx* and *Scx* were

identified as clusters 1 and 6, defined as A1: TC1 and A6: TC2, respectively. Clusters 0, 14,

and 15 were identified as expressing stem progenitor markers including *Cd34*, *Ly6a*,

Pdgfra, *Cd55*, and *Cd248*, and were defined as A2-0: SP1, A2-14: SP2, and A2-15: SP3,

respectively. The characteristic gene activities and annotations for each cluster are

summarized in Fig. 2B.

Next, we analyzed the snRNA-seq data. We evaluated 10,314 cells (median 3,167

genes/cell and 41,611 mean reads/cell). After doublet removal, we performed unbiased

clustering using Seurat and identified 17 clusters (Supplemental Figure 4A). Clusters

expressing tendon-related genes *Mkx* and *Scx* were clusters 1 and 13, defined as R2-1:

TC_1 and R2-13: TC_2, respectively. Clusters 2, 3, and 7 were identified as expressing

stem/progenitor-related genes and defined as R2-2: SP_1, R2-3: SP_2, and R2-7: SP_3,

respectively. To validate the consistency of our approaches, we conducted a

comprehensive comparison between the cell clusters identified in our scRNA-seq analysis

(which included both 2-week and 6-week samples) and our snRNA-seq analysis. We

confirmed that all clusters identified in the scRNA-seq data could be similarly annotated in

the snRNA-seq data, demonstrating strong concordance between these complementary

approaches (Supplemental Figure 5). This validation supports the reliability of our cell type

identification across different single-cell methodologies. Interestingly, R2-5: The MTJ cluster

expressed not only *Tnc* but also *Sox9*, consistent with previous reports (Nagakura et al.,

2020). Using the Signac package's "FindTransferAnchors" function, we calculated

predicted IDs for snATAC-seq clusters based on snRNA-seq annotations (predicted

annotation) (Fig. 2C). Evaluation of the two annotation methods for snATAC-seq revealed

that both methods allowed annotation of major cell types, and correlation between the two

was maintained. Thus, the validity of predicted annotations based on snRNA-seq was

confirmed (Figure 2D).

R2-7: SP3, considered the most undifferentiated stem/progenitor fraction in snRNA-seq,

could be divided into A2-0: SP1 and A2-14: SP2 in snATAC-seq.

Comparing gene activity of tendon-related and stem/progenitor genes in each cluster of the

ground truth annotation, A2-14: SP2 showed high expression of *Ly6a*, *Pdgfra*, and *Tppp3*,

and inverse correlation with *Mkx*, *Scx*, and *Tnmd*. Furthermore, the newly identified surface

antigens *Cd55* and *Cd248* also showed high expression in cluster 14. In contrast, *Cd34* did

not show a high expression pattern in A2-14: SP2. These results suggest that A2-14: SP2

is the most immature cluster, with *Cd55* and *Cd248* showing characteristic expression (Fig.

2E). Subcluster analysis of R7, considered the most immature fraction in snRNA-seq,

revealed two clusters based on *Cd55*/*Cd248* and *Cd34* expression, similar to the snATAC-

seq analysis. The former (R7-1) showed high expression of *Ly6a*, *Tppp3*, and *Pdgfra* (Fig.

2F). Trajectory analysis using R2-7 and A2-14 as roots in snRNA-seq and snATAC-seq,

respectively, showed increased expression of tendon-related genes *Mkx* and *Scx* along the

trajectory, while expression of *Tppp3*, *Cd55*, and *Cd248* decreased (Figure 3A). Evaluation

of peaks in each cluster based on gene expression levels in snRNA-seq revealed multiple

peaks with heights correlated to gene expression for *Mkx* and *Scx*. Similar evaluation of

Cd55 and *Cd248* showed multiple peaks upstream of the TSS for *Cd248* (Fig. 3B). These

results indicate correlation between snRNA-seq and snATAC-seq, and consistent with

scRNA-seq results, suggest that A2-14 represents the most undifferentiated TSPCs based

on *Cd55* and *Cd248* expression.

Comparison of Single Nucleus RNA+ATAC Analysis between 2-week and 6-week

Mouse Achilles Tendon Cells

To confirm the presence of the identified *Cd55*/*Cd248* fraction at 6 weeks, we performed

snRNA-seq and snATAC-seq on 6-week-old mouse Achilles tendon cells. Using the 2-week

snATAC-seq data as a reference, we annotated the 6-week snATAC-seq clusters. Clusters

A2-14 and A2-0 corresponded to clusters A6-12 and A6-0, respectively (Supplemental

Figure 6). We also annotated the 6-week snATAC-seq clusters using 2-week snRNA-seq

with the same results (Supplemental Figure 7). Gene activity evaluation revealed that,

similar to the 2-week data, the 6-week A6-12 and A6-0 clusters showed *Cd55*/*Cd248* high

and *Cd34* dim, and Cd55/*Cd248* dim and *Cd34* high patterns, respectively. Furthermore, annotation of the 6-week snRNA-seq data using the 2-week snRNA-seq as a reference revealed clusters with expression patterns similar to those observed in snATAC-seq. In both cases, clusters with high *Cd55* and *Cd248* expression showed high gene activity (snATAC-seq) and gene expression (snRNA-seq) of *Tppp3*, *Pdgfra*, and *Ly6a*. These results confirm that the *Cd55* and *Cd248* high clusters identified in the 2-week snRNA-seq+snATAC-seq analysis are similarly detected at 6 weeks.

We also compared the snATAC-seq data between 2 and 6 weeks. No significant differences were observed in genomic annotations between the two time points (Supplemental Figure 8A). After merging the datasets (Supplemental Figure 8B), we compared gene activity of tendon/stem-related genes to evaluate changes in gene activity. We found that the activity of *Tppp3*, *Pdgfra*, *Ly6a*, *Cd55*, *Cd248*, and *Cd34* all decreased at 6 weeks (Supplemental Fig 8C). This was consistent with the decreased gene expression observed when comparing 2-week and 6-week data.

Estimation of Transcription Factor Activity in 2-week Mouse Achilles Tendon Cells

To estimate transcription factor activity in each cluster, we used the Single-Cell Regulatory Network Inference and Clustering (SCENIC) package (Lake et al., 2018) to calculate gene regulatory network activity from scRNA-seq gene expression data. SCENIC constructs gene expression networks centered on transcription factors and infers transcription factor activity in each cluster. We summarized the predicted transcription factor activities in tendon and stem/progenitor-related clusters R2-7, R2-2, R2-3, R2-1, and R2-13 identified by snRNA-seq (Figure 4).

Next, we evaluated the gene activity of these identified transcription factors in each snATAC-seq cluster, along with motif activity calculated by chromVAR (Schep et al., 2017). We also assessed the expression levels of transcription factors in each snRNA-seq cluster. In the most immature fractions, represented by cluster A2-14 in snATAC-seq and cluster R2-7 in snRNA-seq, *KLF transcription factor 3* (*Klf3*), *KLF transcription factor 4* (*Klf4*), and *cAMP responsive element binding protein 5* (*Creb5*) showed consistent behavior in gene activity, motif activity, and gene expression. Additionally, *Signal transducer and activator of transcription 2* (*Stat2*) and cAMP responsive element binding protein 3 like 1 (*Creb3l1*) showed high values in A2-0/A2-15 and A2-1, respectively. *Stat2* gene expression was not observed, likely due to false positives resulting from low expression levels. The gene activity and motif activity of each transcription factor estimated by snATAC-seq correlated with the transcription factor activity calculated by SCENIC. Therefore, simultaneous

analysis of snRNA-seq and snATAC-seq could be used to more accurately evaluate the

function of transcription factors in each cluster.

Furthermore, SCENIC can predict candidate transcription factors regulating each gene

(Table 1). *Cd55* and *Cd248* were predicted to be regulated by *Klf3* and *Klf4*, while *Mkx* and

Scx were predicted to be under the control of *Creb3l1*. These predictions were consistent

with the data calculated from gene activity and motif activity in snATAC-seq.

In vitro Evaluation of CD55 and CD248

Immunostaining evaluation of CD55 and CD248, the identified candidate stem/progenitor

markers, revealed expression in the peritenon (Figure 5C, 5A). This was consistent with

previous reports of localized expression of *Tppp3*/*Pdgfra*-positive cells in the peritenon and

our analysis showing co-expression of *Tppp3*/*Pdgfra* and *Cd55*/*Cd248*. The results support

the possibility that SP clusters reflect peritenon. Given that *Cd55* and *Cd248* expression

appeared to reflect *Tppp3*, *Pdgfra*, and *Ly6a* expression more sensitively than *Cd34*, we

extracted tendon cells from 2-week-old mice and sorted them by FACS to determine the

biological phenotype of Cd55 and Cd248 positive cells (Figure 5B).

Gene expression analysis of sorted cells showed that CD55/CD248 positive cells,

compared to negative cells, had lower expression of *Mkx*, *Scx*, *Col1a1*, and *Creb3l1*, but

higher expression of *Ly6a*, *Tppp3*, *Pdgfra*, *Creb5*, and *Klf3*, consistent with snRNA-seq

analysis (Supplemental Figure 9). To assess stem/progenitor capacity, we performed

colony formation assays. CD55/CD248 sorted cells showed significantly increased colony

formation compared to CD55/CD248 negative cells (control) (Figure 5C).

We then evaluated the differentiation tendencies of these cells towards tendon, cartilage,

and bone. Tendon differentiation resulted in clusters of spindle-shaped cells from

CD55/CD248 positive cells, suggesting tenogenic differentiation (Figure 5D). Gene

expression analysis revealed that while CD55/CD248 positive cells initially had lower *Mkx*

and *Scx* expression than negative cells, this pattern reversed after differentiation (Figure

5E). Increased expression of other tendon-related genes such as *Col1a1* and *Bgn* was also

observed. In contrast, no significant increases in *Sox9* and *Collagen type II alpha 1 chain*

(*Col2a1*) (chondrogenic) or *RUNX family transcription factor 2* (*Runx2*), and *Alkaline*

phosphatase, biomineralization associated (*Alpl*) (osteogenic) expression were observed

compared to negative cells during cartilage and bone differentiation assays (Supplemental

Figure 10). Cells negative for CD55/CD248 could be mixed cell populations, including

hematopoietic lineages, cells from tendon mid substance, immune cells, and/or endothelial

cells. However, given that CD55/CD248 negative cells have increased expression of genes

characteristic of tendons such as *Mkx*, *Scx* and *Col1a1*, we could speculate that this

popularity might reflect terminally differentiated tenocytes and CD55/CD248 positive cells

possess tenogenic differentiation capacity.

To further evaluate the tenogenic potential of CD55/CD248 positive cells, we created

tendon-like tissue (bio-tendon) using our previously reported 3D stretch stimulation culture

system. SEM and TEM analysis to assess collagen fiber density and thickness showed that

bio-tendons derived from CD55/CD248 positive cells had an increased proportion of

collagen fibers parallel to the stretch direction and increased collagen fiber diameter. TEM

also revealed characteristic banding patterns and triple helix structures in these bio-

tendons, indicating mature collagen organization (Wieczorek et al., 2015) (Figure 5F-H).

Stretch tests to evaluate the mechanical capacity of the bio-tendons showed high tensile

strength (Figure 5I).

These results demonstrate that CD55/CD248 positive cells have a tendency to

differentiate into tendon cells.

Discussion

Tendons are known for their low cellular content, making complete functional recovery

after injury challenging and increasing the risk of re-rupture. As a result, cell therapy has

garnered attention as a novel treatment strategy, distinct from current conservative and

surgical approaches. However, this requires a thorough understanding of TSPCs. Research

on TSPCs has been limited due to the lack of identified characteristic surface antigens.

To our knowledge, no multi-omics analysis comparing juvenile and mature mouse tendon

cells has been conducted. In this study, we performed single-cell RNA-seq along with

snRNA-seq+snATAC-seq on nuclei isolated from 2-week and 6-week mouse Achilles

tendons. While comparisons using older mice (e.g., 12 weeks or more) were initially

considered, they were excluded due to extremely low cell yield and viability, making 2- and

6-week-old mice more suitable for this analysis. This approach allowed us to identify

candidate surface antigens for TSPCs and elucidate the dynamism of transcription factors

involved in post-developmental tendon growth. As a result, we discovered a novel

combination of *Cd55* and *Cd248* as surface antigens showing characteristic expression in

the TSPC fraction.

While scRNA-seq analysis also showed *Cd34* expression characteristic of the TSPC

fraction, consistent with previous reports, snRNA-seq + snATAC-seq analysis revealed that

the *Cd34* high-expression cluster could be further classified into two clusters based on

Cd55 and *Cd248* expression patterns. The cluster showing high expression of both *Cd55*

and *Cd248* also exhibited high expression of *Tppp3*, *Pdgfra*, and *Ly6a*, suggesting that

isolating *Cd55* and *Cd248* positive fractions may more sensitively capture immature

populations compared to *Cd34*.

snRNA-seq+snATAC-seq analysis provided simultaneous information on gene expression

levels and open chromatin regions for each cluster. Notably, *Cd248* showed several peaks

upstream of the TSS in high-expression clusters, likely reflecting its expression regulation

mechanism. We also identified several peaks that increased proportionally with expression

for *Mkx* and *Scx*. Given the many unknowns in *Mkx* and *Scx* expression regulation

mechanisms (Guerquin et al., 2013; Otabe et al., 2015), we used the SCENIC package to

predict upstream transcription factors based on expressed genes in each snRNA-seq

cluster. Combined with gene activity and motif activity data from snATAC-seq, Creb3l1 was

identified as a transcription factor regulating *Mkx* and *Scx* expression. *Creb3l1* has been

reported to increase in expression throughout development (Liu et al., 2015).

Klf3, *Klf4*, and *Creb5* were identified as characteristic transcription factors in the TSPC

fraction. Recent reports have highlighted the role of the *Klf* family in limb development (Kult

et al., 2021). Given that tendons are components of the developing limb, these transcription

factors may have roles in tendon biology. *Creb5* has been reported to show increased

expression from E11 to E13 (Liu et al., 2015), suggesting a role in early developmental

stages. Furthermore, *Creb5* has been reported to regulate Proteoglycan 4 (*Prg4*)

expression (Zhang et al., 2021). Further investigation into the functions of these

transcription factors in TSPCs is necessary.

CD55 was identified by Hoffmann et al. in 1969 as a surface antigen functioning as a

complement inhibitory factor on erythrocytes. CD55 is known to be expressed from early

developmental stages and characterizes the initial differentiation stage of hematopoietic

stem cells (Guo et al., 2013). It is also expressed in MSCs and has been reported as an

early progenitor marker for mouse mammary epithelial cells (Pal et al., 2017). CD248 was

identified in 1992 as an antigen for the FB5 antibody reacting with vascular wall cells

(Rettig et al., 1992). CD248 is a transmembrane glycoprotein expressed in pericytes and

fibroblasts during developmental stages. Regarding their relationship, CD55 and CD248

have been reported to be expressed in stromal cells during the early stages of arthritis

(Choi et al., 2017), but their detailed mechanisms in tendons remain unclear.

Previously suggested TSPC surface antigen candidates such as *Cd73*, *Cd90*, and *Cd105*

showed poor correlation with the expression of genes like *Tppp3*. Moreover, the initial

report on TSPCs (Bi et al., 2007) described TSPCs as *Scx+Cd34-*. The high expression of

CD55 and CD248 in the peritenon, similar to *Tppp3* (Harvey et al., 2019; Staverosky et al.,

2019), suggests the possibility of cells with tenogenic differentiation potential exhibiting

Scx+Cd34- expression patterns within the tendon. Spatial information is important to

investigate further. Future studies using lineage tracing experiments with mice labeled for

CD55 and CD248 or mouse model of selective ablation of CD55 and CD248 are necessary

to analyze the developmental functions of CD55 and CD248 positive cells and their roles in

injury healing in more detail.

Clinically, to our knowledge, few studies have sorted TSPCs based on surface antigens

and examined their *in vitro* tendon tissue-generating ability. In this study, we found that cells

sorted for CD55/CD248 showed higher clonogenicity compared to CD55-/CD248- cells and

demonstrated superior tendon tissue-generating ability in an artificial tendon model. In the

future, CD55/CD248 double-positive TSPC cells may prove useful in clinical applications

such as *in vitro* artificial tendon creation (Tsutsumi et al., 2022) and cell therapy for tendon

injuries (Huang et al., 2021).

Reference

Andarawis-Puri, N., Flatow, EL., Soslowsky, LJ. (2015). Tendon basic science: Development,

repair, regeneration, and healing. J Orthop Res. 33(6), 780-4.

Asahara, H., Inui, M., Lotz, MK. (2017). Tendons and Ligaments: Connecting Developmental

Biology to Musculoskeletal Disease Pathogenesis. J Bone Miner Res. 32(9), 1773-1782.

Becht, E., McInnes, L., Healy, J., Dutertre, CA., Kwok, IWH., Ng, LG., Ginhoux, F., Newell.

EW. (2019). Dimensionality reduction for visualizing single-cell data using UMAP. Nat

Biotechnol. 37, 38-44.

Bi Y, Ehirchiou D, Kilts TM, Inkson CA, Embree, MC., Sonoyama, W., Li, L., Leet, AI., Seo,

BM., Zhang, L., Shi, S., Young, MF. (2007). Identification of tendon stem/progenitor cells and

the role of the extracellular matrix in their niche. Nat Med. 13(10), 1219-27.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., Satija, R. (2018). Integrating single-cell

transcriptomic data across different conditions, technologies, and species. Nat Biotechnol.

36(5), 411-420.

Cao, Y., Liu, Y., Liu, W., Shan, Q., Buonocore, SD., Cui, L. (2002). Bridging tendon defects

using autologous tenocyte engineered tendon in a hen model. Plast Reconstr Surg. 110(5),

1280-9.

Chen, B., Wang, B., Zhang, WJ., Zhou, G., Cao, Y., Liu, W. (2012). In vivo tendon engineering

with skeletal muscle derived cells in a mouse model. Biomaterials. 33(26), 6086-97.

Chen, L., Liu, JP., Tang, KL., Wang, Q., Wang, GD., Cai, XH., Liu, XM. (2014). Tendon derived
stem cells promote platelet-rich plasma healing in collagenase-induced rat achilles
tendinopathy. Cell Physiol Biochem. 34(6), 2153-68.

Cho, YT., Hsu, WY., Lin, LF., Lin, YN. (2018). Kinesio taping reduces elbow pain during
resisted wrist extension in patients with chronic lateral epicondylitis: a randomized, double-
blinded, cross-over study. BMC Musculoskelet Disord. 19(1), 193.

Connizzo, BK., Yannascoli, SM., Soslowsky, LJ. (2013). Structure-function relationships of
postnatal tendon development: a parallel to healing. Matrix Biol. 32(2), 106-16.

Craft, A. M., Ahmed, N., Rockel, J. S., Baht, G. S., Alman, B. A., Kandel, R. A., Grigoriadis,
A. E. and Keller, G. M. (2013). Specification of chondrocytes and cartilage tissues from
embryonic stem cells. Development 140, 2597-2610.

Diez-Roux, G., Banfi, S., Sultan, M., Geffers, L., Anand, S., Rozado, D., Magen, A., Canidio,
E., Pagani, M., Peluso, I., Lin-Marq, N., Koch, M., Bilio, M., Cantiello, I., Verde, R., De, Masi,
C., Bianchi, SA., Cicchini, J., Perroud, E., Mehmeti, S., Dagand, E., Schrinner, S., Nürnberger,
A., Schmidt, K., Metz, K., Zwingmann, C., Brieske, N., Springer, C., Hernandez, AM., Herzog,
S., Grabbe, F., Sieverding, C., Fischer, B., Schrader, K., Brockmeyer, M., Dettmer, S., Helbig,
C., Alunni, V., Battaini, MA., Mura, C., Henrichsen, CN., Garcia-Lopez, R., Echevarria, D.,
Puelles, E., Garcia-Calero, E., Kruse, S., Uhr, M., Kauck, C., Feng, G., Milyaev, N., Ong, CK.,
Kumar, L., Lam, M., Semple, CA., Gyenesei, A., Mundlos, S., Radelof, U., Lehrach, H.,
Sarmientos, P., Reymond, A., Davidson, DR., Dollé, P., Antonarakis, SE., Yaspo, ML.,
Martinez, S., Baldock, RA., Eichele, G., Ballabio, A. (2011). A High-Resolution Anatomical
Atlas of the Transcriptome in the Mouse Embryo. PLoS Biol. 9(1), e1000582.

Gracey, E., Burssens, A., Cambré, I., Schett, G., Lories, R., McInnes, IB., Asahara, H.,
Elewaut, D. (2020). Tendon and ligament mechanical loading in the pathogenesis of
inflammatory arthritis. Nat Rev Rheumatol. 16(4):193–207.

Fang, F., Xiao, Y., Zelzer, E., Leong, KW., Thomopoulos, S. (2022). A mineralizing pool of
Gli1-expressing progenitors builds the tendon enthesis and demonstrates therapeutic
potential. Cell Stem Cell. 29(12), 1669-1684.

Franken, NA., Rodermond, HM., Stap, J., Haveman, J., van Bree C. (2006). Clonogenic
assay of cells in vitro. Nat Protoc. 2006. 1(5), 2315-9

Grinstein, M., Dingwall, HL., O'Connor, LD., Zou, K., Capellini, TD., Galloway, JL. (2019). A
distinct transition from cell growth to physiological homeostasis in the tendon. Elife. 8, e48689.

Harvey, T., Flamenco, S., Fan, CM. (2019). A Tppp3 + Pdgfra + tendon stem cell population
contributes to regeneration and reveals a shared role for PDGF signalling in regeneration
and fibrosis. Nat Cell Biol. 21(12), 1490-1503.

Guilak, F., Butler, DL., Goldstein, SA., Baaijens, FP. (2014). Biomechanics and
mechanobiology in functional tissue engineering. J Biomech. 47(9), 1933-40

Gumucio, JP., Sugg, KB., Mendias, CL. (2015). TGF-β superfamily signaling in muscle and
tendon adaptation to resistance exercise. Exerc Sport Sci Rev. 43(2), 93-9.

Guzmán, C., Bagga, M., Kaur, A., Westermarck, J., Abankwa, D. (2014). ColonyArea: an
ImageJ plugin to automatically quantify colony formation in clonogenic assays. PLoS One.
9(3), e92444.

Havis, E., Bonnin, MA., Olivera-Martinez, I., Nazaret, N., Ruggiu, M., Weibel, J., Durand, C.,
Guerquin, MJ., Bonod-Bidaud, C., Ruggiero, F., Schweitzer, R., Duprez, D. (2014).
Transcriptomic analysis of mouse limb tendon cells during development. Development.
141(19), 3683-96.

Hittinger, M., Czyz, ZT., Huesemann, Y., Maneck, M., Botteron, C., Kaeufl, S., Klein, CA.,
Polzer, B. (2013). Molecular profiling of single Sca-1+/CD34+,- cells--the putative murine lung
stem cells. PLoS One. 8(12), e83917.

Hoffmann EM. (1969). Inhibition of complement by a substance isolated from human
erythrocytes. II. Studies on the site and mechanism of action. Immunochemistry. 6(3), 405-
19.

Holmes, C., Stanford, WL. (2007). Concise review: stem cell antigen-1: expression, function,
and enigma. Stem Cells. 25(6), 1339-47.

Howell, K., Chien, C., Bell, R., Laudier, D., Tufa, SF., Keene, DR., Andarawis-Puri, N., Huang,
AH. (2017). Novel Model of Tendon Regeneration Reveals Distinct Cell Mechanisms
Underlying Regenerative and Fibrotic Tendon Healing. Sci Rep. 7, 45238.

Isola, JVV., Ocañas, SR., Hubbart, CR., Ko, S., Mondal, SA., Hense, JD., Carter, HNC.,
Schneider, A., Kovats, S., Alberola-Ila, J., Freeman, WM., Stout, MB. (2024). A single-cell
atlas of the aging mouse ovary. Nat Aging. 4(1), 145-162.

Jacobson, KR., Lipp, S., Acuna, A., Leng, Y., Bu, Y., Calve, S. (2020). Comparative Analysis
of the Extracellular Matrix Proteome across the Myotendinous Junction. 19(10), 3955-3967.

Jozsa, L., Kannus, P., Balint, JB., Reffy, A. (1991). Three-dimensional ultrastructure of human
tendons. Acta Anat (Basel). 142(4), 306-12.

Kataoka, K., Kurimoto, R., Tsutsumi, H., Chiba, T., Kato, T., Shishido,. K., Kato, M., Ito, Y.,
Cho, Y., Hoshi, O., Mimata, A., Sakamaki, Y., Nakamichi, R., Lotz, MK., Naruse, K., Asahara,
H. (2020). In vitro Neo-Genesis of Tendon/Ligament-Like Tissue by Combination of Mohawk
and a Three-Dimensional Cyclic Mechanical Stretch Culture System. Front Cell Dev Biol.
2020;8:307.

Kendal, AR., Layton, T., Al-Mossawi, H., Appleton, L., Dakin, S., Brown, R., Loizou, C.,
Rogers, M., Sharp, R., Carr, A. (2020). Multi-omic single cell analysis resolves novel stromal
cell populations in healthy and diseased human tendon. Sci Rep. 10(1): 13939.

Leong, DJ., Sun, HB. (2016). Mesenchymal stem cells in tendon repair and regeneration:
basic understanding and translational challenges. Ann. N. Y. Acad. Sci. 1383, 88–96.

Leong, NL., Kator, JL., Clemens, TL., James, A., Enamoto-Iwamoto, M., Jiang, J. (2020).
Tendon and Ligament Healing and Current Approaches to Tendon and Ligament
Regeneration. J Orthop Res. 38(1), 7-12.

Li, Y., Wu, T., Liu, S. (2021). Identification and Distinction of Tenocytes and Tendon-Derived
Stem Cells. Front Cell Dev Biol. 9, 629515.

Liu, H., Xu, J., Liu, CF., Lan, Y., Wylie, C., Jiang, R. (2015). Whole transcriptome expression

profiling of mouse limb tendon development by using RNA-seq. J Orthop Res. 33(6), 840–
848.

Loiacono, C., Palermi, S., Massa, B., Belviso, I., Romano, V., Gregorio, AD., Sirico, F., Sacco,
AM. (2019). Tendinopathy: Pathophysiology, Therapeutic Options, and Role of Nutraceutics.
A Narrative Literature Review. Medicina (Kaunas). 55(8), 447.

Lui, PP. (2015). Markers for the identification of tendon-derived stem cells in vitro and tendon
stem cells in situ - update and future development. Stem Cell Res Ther. 6(1), 106.

Lui, PPY., Wong, CM. (2019). Biology of Tendon Stem Cells and Tendon in Aging. Front
Genet. 10, 1338.

McGinnis, CS., Murrow, LM., Gartner, ZJ. (2019). DoubletFinder: Doublet Detection in Single-
Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst. 8(4), 329-337.e4.

Mendias, CL., Gumucio, JP., Bakhurin, KI., Lynch, EB., Brooks, SV. (2012). Physiological
loading of tendons induces scleraxis expression in epitenon fibroblasts. J Orthop Res. 30(4),
606-12.

Morikawa, S., Mabuchi, Y., Kubota, Y., Nagai, Y., Niibe, K., Hiratsu, E., Suzuki, S., Miyauchi-
Hara, C., Nagoshi, N., Sunabori, T., Shimmura, S., Miyawaki, A., Nakagawa, T., Suda, T.,
Okano, H., and Matsuzaki, Y. (2009). Prospective identification, isolation, and systemic
transplantation of multipotent mesenchymal stem cells in murine bone marrow. J. Exp.
Med. 206, 2483-2496.

Nagakura, R., Yamamoto, M., Jeong J, Hinata N, Katori Y, Chang WJ, Abe S. (2020).
Switching of Sox9 expression during musculoskeletal system development. Sci Rep. 10(1),
8425.

Nusspaumer, G., Jaiswal, S., Barbero, A., Reinhardt, R., Ishay Ronen, D., Haumer, A., Lufkin,
T., Martin, I. and Zeller, R. (2017). Ontogenic identification and analysis of mesenchymal
stromal cell populations during mouse limb and long bone development. Stem Cell Rep. 9,
1124-1138.

Pliner, HA., Packer, JS., McFaline-Figueroa, JL., Cusanovich, DA., Daza, RM., Aghamirzaie,

D., Srivatsan, S., Qiu, X., Jackson, D., Minkina, A., Adey, AC., Steemers, FJ., Shendure, J.,
Trapnell, C. (2018). Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell
Chromatin Accessibility Data. Mol. Cell. 71, 858–871. e8.

Ruzzini, L., Abbruzzese, F., Rainer, A., Longo, UG., Trombetta, M., Maffulli, N., Denaro, V.
(2014). Characterization of age-related changes of tendon stem cells from adult human
tendons. Knee Surg Sports Traumatol Arthrosc. 22(11), 2856-66.

Satpathy, AT., Granja, JM., Yost, KE., Qi, Y., Meschi, F., McDermott, GP., Olsen, BN.,
Mumbach, MR., Pierce, SE., Corces, MR., Shah, P., Bell, JC., Jhutty, D., Nemec, CM., Wang,
J., Wang, L., Yin, Y., Giresi, PG., Chang, ALS., Zheng, GXY., Greenleaf, WJ., Chang, HY.
(2019). Massively parallel single-cell chromatin landscapes of human immune cell
development and intratumoral T cell exhaustion. Nat Biotechnol. 37(8), 925-936.

Sidney, LE., Branch, MJ., Dunphy, SE., Dua, HS., Hopkinson, A. (2014). Concise review:
evidence for CD34 as a common marker for diverse progenitors. Stem Cells. 32(6), 1380-9.
876

Somerville, JM., Aspden, RM., Armour, KE., Armour, KJ., Reid, DM. (2004). Growth of
C57BL/6 mice and the material and mechanical properties of cortical bone from the tibia.
Calcif Tissue Int. 74(5), 469-75.

Staverosky, JA., Pryce, BA., Watson, SS., and Schweitzer, R. (2019). Tubulin polymerization-
promoting protein family member 3, Tppp3, is a specific marker of the differentiating tendon
sheath and synovial joints. Dev Dyn. 238(3), 685-92.

Steinmann, S., Pfeifer, CG., Brochhausen, C., Docheva, D. (2020). Spectrum of Tendon
Pathologies: Triggers, Trails and End-State. Int J Mol Sci. 21(3), 844.

Still, C, 2nd., Chang, WT., Sherman, SL., Sochacki, KR., Dragoo, JL., Qi, LS. (2021). Single-
cell transcriptomic profiling reveals distinct mechanical responses between normal and
diseased tendon progenitor cells. Cell Rep Med. 2(7), 100343.

Stuart, T., Hanna, J., Gutruf, P. (2021). Single-cell chromatin state analysis with Signac. Nat
Methods. 18(11), 1333-1341.

Sung, JH., Yang, HM., Park, JB., Choi, GS., Joh, JW., Kwon, CH., Chun, JM., Lee, SK., Kim,

SJ. (2008). Isolation and characterization of mouse mesenchymal stem cells. Transplant Proc. 40(8), 2649-54.

Tachibana, N., Chijimatsu, R., Okada, H., Oichi, T., Taniguchi, Y., Maenohara, Y., Miyahara, J., Ishikura, H., Iwanaga, Y., Arino, Y., Nagata, K., Nakamoto, H., Kato, S., Doi, T., Matsubayashi, Y., Oshima, Y., Terashima, A., Omata, Y., Yano, F., Maeda, S., Ikegawa, S., Seki, M., Suzuki, Y., Tanaka, S., Saito, T. (2022). RSPO2 defines a distinct undifferentiated progenitor in the tendon/ligament and suppresses ectopic ossification. Sci Adv. 8(33), eabn2138.

Takakura, N., Yoshida, H., Ogura, Y., Kataoka, H., Nishikawa, S., and Nishikawa, S. (1997). PDGFR alpha expression during mouse embryogenesis: immunolocalization analyzed by whole-mount immunohistostaining using the monoclonal anti-mouse PDGFR alpha antibody APA5. J. Histochem. Cytochem. 45, 883-893.

Tang, JB., Wu, YF., Cao, Y., Chen, CH., Zhou, YL., Avanessian, B., Shimada, M., Wang, XT., Liu, PY. (2016). Basic FGF or VEGF gene therapy corrects insufficiency in the intrinsic healing capacity of tendons. Sci Rep. 6, 20643.

Tempfer, H., Wagner, A., Gehwolf, R., Lehner, C., Tauber, M., Resch, H., Bauer, HC. (2009). Perivascular cells of the supraspinatus tendon express both tendon- and stem cell-related markers. Histochem Cell Biol. 131(6), 733-41.

Thomopoulos, S., Parks, WC., Rifkin, DB., Derwin KA. (2015). Mechanisms of tendon injury and repair. J Orthop Res. 33(6), 832-9.

Trapnell, C., Cacchiarelli. D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, NJ., Livak, KJ., Mikkelsen, TS., Rinn, JL. (2014). The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. Nat Biotechnol. 32(4), 381-386.

Tsutsumi, H., Kurimoto, R., Nakamichi, R., Chiba, T., Matsushima, T., Fujii, Y., Sanada, R., Kato, T., Shishido, K., Sakamaki, Y., Kimura, T., Kishida, A., Asahara, H. (2022). Generation of a tendon-like tissue from human iPS cells. J Tissue Eng. 13, 20417314221074018.

Walia, B., Huang, AH. (2019). Tendon stem progenitor cells: Understanding the biology to inform therapeutic strategies for tendon repair. J Orthop Res . 37(6), 1270-1280.

Wang, B., Liu, W., Zhang, Y., Jiang, Y., Zhang, WJ., Zhou, G., Cui, L., Cao, Y. (2008).
Engineering of extensor tendon complex by an ex vivo approach. Biomaterials. 29(20): 2954-
61.

Yin, Z., Hu, JJ., Yang, L., Zheng, ZF., An, CR., Wu, BB., Zhang, C., Shen, WL., Liu, HH.,
Chen, JL., Heng, BC., Guo, GJ., Chen, X., Ouyang, HW. (2016). Single-cell analysis reveals
a nestin + tendon stem/progenitor cell population with strong tenogenic potentiality. Sci Adv.
2(11), e1600874.

Yu, G., Wang, LG., Han, Y., He, QY. (2012). clusterProfiler: an R package for comparing
biological themes among gene clusters. OMICS. 16(5), 284-7.

Yu, G., Wang, LG., He, QY. (2015). ChIPseeker: an R/Bioconductor package for ChIP peak
annotation, comparison and visualization. Bioinformatics. 31(14), 2382-3.

Zhang, C., Zhu, J., Zhou, Y., Thampatty, BP., Wang, JH. (2019). Tendon Stem/Progenitor
Cells and Their Interactions with Extracellular Matrix and Mechanical Loading. Stem Cells
Int .2019, 3674647.

Zhang, T., Wan, L., Xiao, H., Wang, L., Hu, J., Lu, H.(2023) Single-cell RNA sequencing
reveals cellular and molecular heterogeneity in fibrocartilaginous enthesis formation. Elife.
12: e85873.

Acknowledgements
We thank all the members of the Department of Systems BioMedicine at Institute of Science
Tokyo for their support. We also thank the Research Core at Institute of Science Tokyo for
supporting cell sorting.

Author contributions
Conceptualization, H.T., T.C., and H.A.; Methodology, H.T., T.C., and H.A.; Formal analysis,
H.T.; Investigation, H.T., F. Y., T. M., T. K., A. K., K. A., Y. S.; Writing – Original Draft, H.T.;
Writing – Review & Editing, T.C., and H.A.; Funding Acquisition, H.A.; Supervision, H.A.

Data availability
FASTQ data of RNA-Seq and ATAC-seq are deposited in DDBJ under accession number

PRJDB18857

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research,

authorship, and/or publication of this article.

Funding

This work was supported by JSPS KAKENHI (Grant Numbers JP15H02560, JP20H05696,

16H06279 (PAGS)), AMED (Grant Numbers JP21gm0810008, JP23ym0126805,

JP24gm0010009), and NIH (Grant Number R01AR080127) to H.A.

Figure legends

Figure 1. scRNA-seq of tendon cells from 2-week-old and 6-week-old mice and the

identification of surface markers of TSPC.

(A) Integrated uniform manifold approximation and projection (UMAP) scRNA-seq clustering

of cells harvested from 2-week-old and 6-week-old mouse Achilles tendons.

(B) Dot plot of average gene expression levels of the indicated genes in each scRNA-seq

cluster. The size of the dot reflects the percentage of cells in the cluster that express each

gene. TC, tenocyte; SP1, tendon stem/progenitor cell_1; CA, cartilage; RB, ribosomal RNA;

SP2, tendon stem/progenitor cell_2; LC1, lymphocyte_1; EC, endothelial cell; RBC, red blood

cell; SM, smooth muscle cell; PC, proliferating cell; LC2, lymohocytes_2; MTJ, myotendinous

junction cell; SW, Schwann cell; VEC, vascular endothelial cell; MC, macrophage.

(C) Proportions of cells in clusters identified from scRNA-seq. Clusters are colored according

to cluster type.

(D) Volcano plot of gene expression in the SP2 cluster and the identification of candidate

TSPC marker genes (red under line).

(E) Feature plot of *Cd55* and *Cd248* expression.

(F) Correlation of gene expression of TSPC candidate genes in 2-week data.

(G) Violin plots presenting the gene expression changes for a selection of differentially

expressed genes.

Figure 2. snATAC-seq of tendon cells from a 2-week-old mouse and the validation of

Cd55 and Cd248 as candidate markers of TSPC.

(A) UMAP snATAC-seq clustering of cells derived from the Achilles tendon of a 2-week-old

mouse. Annotation was based on each gene activity (ground-truth annotation). SP1, tendon

stem/progenitor cell_1; TC1, tenocyte_1; MTJ, myotendinous junction cell; SM, smooth

muscle cell; MC, macrophage; CA, cartilage; TC2, tenocyte_2; SK, skeletal muscle cell; SW, Schwann cell; LC, lymphocyte; NC, neutrophil; EC, endothelial cell; RBC, red blood cell; UC, unspecified cell; SP2, tendon stem/progenitor cell_2; SP3, tendon stem/progenitor cell_3; VEC, vascular endothelial cell.

(B) Dot plot of average gene activity of the indicated genes in each snATAC-seq cluster. The size of the dot reflects the percentage of cells in the cluster that express each gene.

(C) UMAP visualization and predicted annotation of 2-week snATAC-seq after integration and label transfer of 2-week snRNA-seq data.

(D) Identification of matching cell clusters between the 2-week snRNA- and 2-week snATAC-seq data from visualized as heatmap. The heatmap shows the proportions of cells from each snATAC-seq cluster across all sample conditions assigned to each snRNA-seq cluster as part of the label-transfer process.

(E) Violin plot of tenocytes and TSPC-related gene expression in each cluster.

Figure 3. Trajectory analysis and peak visualization of snRNA-seq and snATAC-seq data for the tendon and tendon stem/progenitor cell-related cluster.

(A) UMAP representation of snRNA-seq differentiation trajectory of tenocytes and TSPC lineage and pseudotime-dependent gene expression changes of *Tppp3*, *Cd55*, *Cd248*, *Mkx,* and *Scx,* as inferred using Monocle3.

(B) UMAP representation of snATAC-seq differentiation trajectory of tenocytes and the TSPC lineage and pseudotime-dependent gene expression changes, as inferred using Cicero.

(C) Coverage plots of *Mkx*, *Scx*, *Cd55,* and *Cd248*. Selected peaks that differ across each cluster are highlighted.

Figure 4. Transcription factor landscapes of 2-week mouse Achilles tendons.

SCENIC analysis of transcription factor activity based on 2-week snRNA-seq data for tenocytes and the TSPC lineage (Left). Validation was performed based on the gene activity and motif activity of 2-week snATAC-seq and gene expression of 2-week snRNA-seq (right).

Figure 5. *In vitro* analysis of CD55+/CD248+ TSPCs.

(A) Immunohistochemical image of 10-week mouse Achilles tendons. Scale bars show 100μm. CD55 and CD248, green; Hoechst 33342, blue.

(B) Schema of the *in vitro* assessment of the capacity of CD55+/CD248+ TSPCs as the differentiation toward tenocytes.

(C) Colony-forming efficiency of CD55+/CD248+ and CD55-/CD248- (negative) TSPCs. Colonies were stained with crystal violet (n = 6). CD55+/CD248+ TSPC exhibited higher

clonogenic capacity. Data are presented as means ± SEM. **$p < 0.01$.

(D) Morphological changes of CD55+/CD248+ and negative TSPCs after tenogenic induction.

(E) Quantitative PCR of tendon-related gene expression in CD55+/CD248+ and negative

TSPCs after tenogenic induction (n = 3). Data are presented as means ± SD. **$p < 0.01$, *p

< 0.05.

(F) SEM and TEM imaging of artificial tendons derived from CD55+/CD248+ and negative

TSPCs. Data are presented as means ± SD. **$p < 0.01$, *$p < 0.05$.

(G) Proportions of fiber alignment for each artificial tendon (n = 4). Data are presented as

means ± SEM. ***$p < 0.005$, **$p < 0.01$, *$p < 0.05$.

(H) Diameter of collagen fiber in each artificial tendon based on TEM imaging (n = 4). Data

are presented as means ± SD. ***$p < 0.005$.

(I) Tensile strength (MPa) of each artificial tendon (n = 5). Data are presented as means ±

SEM. ***$p < 0.005$.

1053

Table1: Estimated TF activity in each gene

Supplemental Figure 1. 2w- and 6w-scRNA seq results.

(A) Integrated uniform manifold approximation and projection (UMAP) scRNA-seq clustering

of cells harvested from 2-week-old and 6-week-old mouse Achilles tendons. Clusters are

colored according to cluster type.

(B) Feature plot of *Tppp3*, *Pdgfra* and *Ly6a* expression.

1060

Supplemental Figure 2. Comparison of 2w- and 6w-scRNA seq results.

(A) Violin plot of gene expression enriched in the limb bud (Development (2014) 141, 3683-

3696) in each cluster.

(B) Gene Ontology (GO) terms associated with genes with upregulated expression in the

SP2 cluster.

(C) Number of inferred interactions and interaction strength of genes expressed within 2w-

and 6w-mouse scRNA-seq data.

1068

1069   **Supplemental Figure 3. Comparative analysis of 2w- and 6w-scRNA seq results.**

1070   Heatmap showing differential gene expression between red (2w) and green (6w).

1071

Supplemental Figure 4. Analysis of 2-week snRNA-seq data.

(A) UMAP snRNA-seq clustering of cells harvested from 2-week mouse Achilles tendons.

(B) Dot plot of average gene expression levels of the indicated genes in 2-week snRNA-seq

clusters. RB, ribosomal RNA; TC1, tenocyte_1; SP1, stem/progenitor cell_1; SP2,

stem/progenitor cell_2; SM, smooth muscle cell; MTJ, myotendinous junction cell; CA,

cartilage; SP3, stem/progenitor cell_3; LC, lymphocyte; MC, macrophage; RBC, red blood

cell; SW, Schwann cell; EC, endothelial cell; TC2, tenocyte_2; PC, proliferating cell; NC,

neutrophil; UC, unspecified cell.

(C) Violin plot of tenocytes and TSPC-related gene expression in each cluster.

(D) Feature plot of TSPC-related gene expression.

1082

Supplemental Figure 5. Analysis of 6-week snATAC-seq.

(A) UMAP snATAC-seq clustering of cells harvested from 6-week mouse Achilles tendons.

Annotation was based on gene activity (ground-truth annotation, left) and the predicted

annotation inferred from 2-week snATAC-seq (right).

(B) Identification of matching cell clusters between the 2-week and 6-week snATAC-seq data,

visualized as a heatmap.

(C) Violin plot of tenocytes and TSPC-related gene activity in each cluster.

1090

1091   **Supplemental Figure 6. Comparison of snRNA-seq and scRNA-seq.**

1092   (A) Dot plot of representative DEGs in snRNA-seq (2w) and snRNA-seq (2w+6w).

1093   (B) Relevance of annotation in each data.

1094

Supplemental Figure 7. Analysis of 6-week snRNA-seq.

(A) UMAP snRNA-seq clustering of cells harvested from 6-week mouse Achilles tendons.

Annotation was based on each gene expression (ground-truth annotation, left) and predicted

annotation inferred from 2-week snRNA-seq (right).

(B) Identification of matching cell clusters between the 2-week and 6-week snRNA-seq data

visualized as a heatmap.

(C) Violin plot of tenocytes and TSPC-related gene expression in each cluster.

1102

1103   **Supplemental Figure 8. Comparison of 2-week and 6-week snATAC-seq.**

1104   (A) Circle plot of annotated differentially accessible regions for each data.

1105   (B) Integrated UMAP snATAC-seq clustering of cells harvested from 2-week and 6-week

1106   mouse Achilles tendons.

1107   (C) Feature plot of TSPC-related gene activity in each dataset.

1108

1109   **Supplemental Figure 9. Gene expression changes in CD55+/CD248+ and negative**

1110   **TSPCs.**

1111   Quantitative PCR of gene expression in CD55+/CD248+ and negative TSPCs (n = 4). Data

1112   are presented as means ± SEM. *$p < 0.05$, ***$p < 0.005$.

1113

1114   **Supplemental Figure 10. Chondrogenic and osteogenic induction of CD55+/CD248+**

1115   **and negative TSPCs.**

1116   Quantitative PCR of tendon-related genes in CD55+/CD248+ and negative TSPCs after

1117   chondrogenic and osteogenic induction (n = 4). Data are presented as means ± SEM. **$p <$

1118 0.01, *$p < 0.05$.

Figure 1

Figure 2

Figure 3

Figure 4

Estimated TF activity
(stem/progenitor cell and tenocytes lineage)

Gene activity, motif activity and gene expression

Figure 5

Table1

Estimated TF activity : Cd55	
TF	**spearman correlation**
Klf4	0.264195541
Klf3	0.262413411
Zeb1	0.219593135
Pbx1	0.204085218
Hic1	0.202481054
Klf2	0.196232716
Nfe2l2	0.196076187

Estimated TF activity : Cd248	
TF	**spearman correlation**
Hic1	0.375565336
Klf3	0.277768834
Zmiz1	0.272991765
Klf4	0.260526766
Zeb1	0.250583629
Sp3	0.2502976
Kdm5b	0.193229132

Estimated TF activity : Mkx	
TF	**spearman correlation**
Creb3l1	0.320638653
Zfhx3	0.169341424
Mxi1	0.149048856
Bhlhe40	0.144301452
Ets2	0.137445497
Max	0.116134957
Elk3	0.098325126

Estimated TF activity : Scx	
TF	**spearman correlation**
Creb3l1	0.316522613
Npdc1	0.163452737
Ets2	0.145463784
Mix1	0.121248221
Bhlhe41	0.115930755
Elk3	0.10803805
Bmyc	0.104968457

Supplemental figure 1

A

B

Supplemental Figure 2

Supplemental Figure 3

Supplemental Figure 4

Supplemental Figure 5

A

Supplemental Figure 6

A

6w: snATAC-seq (ground-truth annotation)   6w: snATAC-seq (predicted annotation)

B

Predicted annotation

Proportion of 6w_snATAC-seq cells
assigned to 2w_snATAC-seq cluster

C

Gene activity in 6w_snATAC-seq

Supplemental Figure 7

Supplemental Figure 8

Supplemental figure 9

Supplemental Figure 10

: CD55/CD248, induction+

\- \- \- \- \- \- \- \- \-

: CD55/CD248, induction-

: negative, induction+

\- \- \- \- \- \- \- \- \-

: negative, induction-