TITLE

Conflicts between the DNA replication and repair machineries promote cell death in Gram-positive bacteria

AUTHORS

Hannah Gaimster, Daniel Stevens, Julia Hubbard, Katarzyna Mickiewicz, Heath Murray and Charles Winterhalter*

AFFILIATION

Centre for Bacterial Cell Biology, Biosciences Institute, Newcastle University, Newcastle-upon-Tyne, NE2 4AX, UK

AUTHOR LIST FOOTNOTES

* Correspondence: Charles.winterhalter@newcastle.ac.uk

ABSTRACT

Cellular proliferation relies on successful coordination and completion of genome replication and segregation. To help achieve this, many bacteria utilise regulatory pathways that ensure DNA replication initiation only occurs once per cell cycle. When dysregulated, loss of DNA replication control can have severe consequences. In *Escherichia coli* it has been observed that hyper-initiation of DNA synthesis leads to pleiotropic genome instability and cell death. Therefore, targeting DNA replication initiation proteins to promote hyper-initiation may be an approach to generate novel antimicrobials. However, it is not clear whether DNA replication hyper-initiation generally inhibits bacterial proliferation, and if so, via which pathways. To address this question, we devised genetic systems to artificially induce hyper-initiation in the model organism *Bacillus subtilis* and the pathogen *Staphylococcus aureus*. In both species, hyper-initiation elicited cellular degeneration culminating in growth inhibition by cell death. During this process in *B. subtilis*, temporal analyses revealed the early onset of the DNA damage response, followed by membrane depolarisation and cell lysis. This phenotype could be supressed by removing pathways that repair damaged DNA, suggesting that cell death is a consequence of conflicts between DNA replication and repair. In *S. aureus,* cells quickly accumulated striking morphological changes associated with rapid loss of chromosomal DNA and death via a lysis-independent pathway. Moreover, inducing hyper-initiation in *S. aureus* was observed to decrease bacterial survival during infection of murine macrophages. Taken together, these results suggest that stimulating the initiation of bacterial DNA replication could be an alternative approach to inhibiting bacterial growth, particularly in combination with compounds that inhibit or poison DNA repair, akin to cancer therapies.

INTRODUCTION

A successful cell cycle requires replication and segregation of the genome. In most cells, DNA synthesis begins at specific chromosomal loci called origins and requires accurate coordination with chromosome segregation and cell division to produce viable daughter cells [1, 2]. Importantly, while replication initiation culminates in loading of ring-shaped helicases enabling double-strand DNA unwinding prior to synthesis, the molecular mechanisms required to achieve helicase loading in prokaryotes and eukaryotes are distinct [3]. Therefore, the unique and essential characteristics of the bacterial DNA replication initiation pathway make it an attractive target for drug development [4].

In bacteria, chromosome replication generally initiates from a single origin (*oriC*), where the universal prokaryotic initiator protein DnaA binds to sequence specific elements (DnaA-boxes and DnaA-trios) and unwinds the DNA duplex [5-7], followed by recruitment and loading of helicases around single-stranded DNA [8-10]. During this process, DnaA activity is tightly regulated to ensure genome duplication occurs only once per cell cycle [11, 12].

Abnormal initiation of genome duplication can result in either replication inhibition or excessive DNA synthesis, both of which can inhibit bacterial proliferation. Intuitively, under-initiation of DNA replication produces cells that lack a copy of the genome [13]. A more complex situation is observed during hyper-initiation of DNA replication. In the Gram-negative model organism *Escherichia coli* cultured aerobically, hyper-initiation causes DNA double-strand breaks (DSBs) as a result of the DNA replication machinery (replisome) encountering repair systems acting on the genome [14, 15]. However, the lethal phenotype associated with hyper-initiation of DNA replication remains enigmatic, as suppression of cell death did not correlate with a significant decrease in DSBs [14]. Moreover, it is not yet clear whether hyper-initiation of DNA replication generally inhibits bacterial proliferation, and if so, via which pathways.

In this study, we investigated the consequence of DNA replication hyper-initiation in the Gram-positive model organism *Bacillus subtilis* and the pathogen *Staphylococcus*

aureus. In both cases hyper-initiation resulted in loss of cell viability, however the lethal phenotypes were not identical. The *B. subtilis* strain displayed cell elongation, DNA damage, membrane depolarisation, and eventually cell lysis. The *S. aureus* strain also displayed altered cell morphology with clear defects in chromosome inheritance, but here loss of cell viability occurred through a lysis-independent mechanism. Interestingly, induction of *S. aureus* hyper-initiation inhibited bacterial growth in a murine macrophage model, highlighting the potential of stimulating DNA replication initiation as an approach to reduce bacterial load during infection.

MATERIALS AND METHODS

Reagents and growth conditions

Nutrient agar (NA; Oxoid) was used for routine selection and maintenance of *E. coli* and *B. subtilis* strains. Tryptic soy agar (TSA; Oxoid) was used for routine selection and maintenance of *S. aureus* strains. Unless otherwise stated, *B. subtilis* and *S. aureus* strains were grown at 30°C and *E. coli* was grown at 37°C. Supplements were added as required: ampicillin (100 µg/ml), erythromycin (1 µg/ml) in conjunction with lincomycin (25 µg/ml), chloramphenicol (5 µg/ml for *B. subtilis*, 10 µg/ml for *S. aureus*), kanamycin (5 µg/ml), spectinomycin (50 µg/ml), tetracycline (10 µg/ml), xylose (1% w/v) and anydrotetracycline (aTc, 20 ng/ml). All chemicals and reagents were obtained from Sigma-Aldrich. RAW-Blue cells were maintained in Dulbecco's Modified Eagle's medium (DMEM, Sigma Cat#D6429) supplemented with 5% fetal bovine serum (FBS) at 37°C and 5% CO_2.

Biological resources: *E. coli*, *B. subtilis,* and *S. aureus* strains

All strains and cell lines used in this study are listed in Supplementary Table 1.

B. subtilis strain construction

Transformation of competent *B. subtilis* cells was performed using an optimized two-step starvation procedure as previously described [16]. Briefly, recipient strains were grown overnight at 30°C in transformation medium (Spizizen salts supplemented with 1 µg/ml Fe-NH_4-citrate, 6 mM $MgSO_4$, 0.5% w/v glucose, 0.02 mg/ml tryptophan and 0.02% w/v casein hydrolysate) supplemented with xylose where required. Overnight cultures were diluted 1:20 into fresh transformation medium supplemented with xylose where required and grown at 30°C for 3 hours with continual shaking. An equal volume of prewarmed starvation medium (Spizizen salts supplemented with 6 mM $MgSO_4$ and 0.5% w/v glucose) was added and the culture was incubated at 30°C for 2 hours with continuous shaking. DNA was added to 350 µl cells and the mixture was incubated at 30°C for 1 hour with continual shaking. 20-200 µl of

each transformation was plated onto selective media supplemented with xylose where required and incubated at 30°C for 24-48 hours.

S. aureus strain construction

Transformation of competent *S. aureus* cells was performed using plasmid electroporation [17]. Briefly, recipient cells were grown overnight in tryptic soy broth (TSB; Oxoid). Overnight cultures were diluted 1:100 in fresh TSB and grown to an absorbance at 600 nanometers (A_{600}) of 0.6. Cells were rapidly cooled down in an ice water bath and washed three times using an equal volume of ice-cold sterile deionised water followed by two washes in an equal volume of ice-cold 10% w/v glycerol solution. After the final centrifugation, cells were resuspended in 1:100 of the initial culture volume in ice-cold 10% w/v glycerol, snap frozen using liquid nitrogen, and stored at -80°C for up to six months. Before electroporation, electrocompetent cells were thawed at room temperature for 5 min, centrifuged at 10,000 g for 1 min, and resuspended in 100 µl of electrocompetent buffer (10% w/v glycerol supplemented with 500 mM sucrose). DNA was added to the cell mixture and electroporation was performed at 2.3 kV, 100 Ω, 25 µF (Bio-rad Gene Pulser II) using a 1 mm gap cuvette (VWR #732-1135). Cells were immediately resuspended in 1 ml of TSB supplemented with 500 mM sucrose, incubated for 1.5 hours at the permissive temperature, plated onto selective medium and incubated at permissive temperature for 48 hours.

E. coli plasmid construction and propagation

E. coli transformation was performed in DH5α via heat-shock following the Hanahan method [18]. Descriptions of plasmid construction, where necessary, are provided below.

pDS150 was generated using Gibson assembly to introduce *SaDnaA* into pRAB11 [19]. pDS157 was created using Quickchange mutagenesis on pDS150 to introduce the *SaDnaA*^{G159S} point mutation.

Spot titre assays

B. subtilis cells were grown in penassay broth medium (PAB) overnight at 30°C in the presence of xylose. Overnight cultures were diluted 1:100 into fresh PAB with or without xylose (time = 0 h). Samples were harvested at time points indicated, 10-fold serial dilutions were made into PAB with xylose, and 5 μL aliquots were spotted onto NA plates supplemented with xylose. For *S. aureus* spot titre assays, cells were grown in TSB supplemented with chloramphenicol overnight at 30°C and 5 μL of serial dilutions were spotted onto TSA supplemented with chloramphenicol in the presence or absence of aTc. All plates were incubated at 30°C for 24 hours. Experiments were performed independently at least three times and representative data are shown.

Automated plate reader analyses

Strains were grown overnight at 30°C in PAB with xylose (*B. subtilis*) or TSB supplemented with chloramphenicol (*S. aureus*). Overnight cultures were diluted 1:100 into PAB with or without xylose (*B. subtilis*), or 1:1000 into TSB with or without aTc (*S. aureus*) and automated absorbance measurements were captured over time at 30°C using a Tecan Sunrise plate reader (high shaking parameters with 2.8 mm shake width at 12.3 Hz, absorbance measurement every six minutes). All experiments were independently performed at least three times and representative data are shown.

Microscopy and image analysis

Microscopy was performed on an inverted epifluorescence microscope (Nikon Ti) fitted with a Plan Apochromat Objective (Nikon DM 100x/1.40 Oil Ph3). Light was transmitted from a CoolLED pE-300 lamp through a liquid light guide (Sutter Instruments), and images were collected using a Prime CMOS camera (Photometrics). Fluorescence filter sets were from Chroma: GFP (49002, 537 EX470/40 (EM), DM495lpxr (BS), EM525/50 (EM)) and mCherry (49008, EX560/40 (EM), 538 DM585lprx (BS), EM630/75 (EM)). Digital images were acquired using Metamorph software (version 7.7). Cells were mounted on ~1.2% agar pads

and a 0.13-0.17 mm glass coverslip (VWR) was placed on top. Wavelengths used: Brightfield and mCherry (150 ms), GFP/UV (50 ms), all at 100% exposure.

For timelapse microscopy experiments, a GeneFrame was used to create a flat agarose pad using 0.7% agarose dissolved into PAB with or without xylose (*B. subtilis*) or TSB with or without aTc (*S. aureus*). An aliquot of cells (1.5 μL) was spotted onto the pad and a 0.13-0.17 mm glass coverslip (VWR) was placed on top. Digital images were acquired using Metamorph software (version 7.7) every 15 minutes for 8-12 hours.

For cultures stained with $DiBAC_4(3)$ (Invitrogen, Thermofisher), 0.5 μl dye was added to 100 μl of cells and the mixture was incubated at 30°C with shaking at 800 rpm for 5 minutes, using a Thermomixer C (Eppendorf), then spotted onto an agarose slide. $DiBAC_4(3)$ preferentially binds cellular membranes that are depolarized, emitting a green fluorescence signal to allow for detection (i.e. cells appear fluorescent if the cell envelope is compromised).

For cultures stained with 4',6-diamidino-2-phenylindole (DAPI, Thermofisher), the dye was added to 100 μl of cells at a final concentration of 1 μg/ml and the mixture was incubated at 30°C with shaking at 800 rpm for 5 minutes, using a Thermomixer C (Eppendorf), then spotted onto an agarose slide.

The Fiji imaging software along with the ObjectJ and Chain tracer was used for image analysis [20]. Brightness and contrast used for each time point and condition remained consistent and all images are scaled to 0.065 microns.

Sample preparation for marker frequency analysis

B. subtilis cells were grown overnight in PAB at 30°C with xylose. Overnight cultures were diluted 1:100 into PAB with or without xylose (time = 0 h) and incubated at 30°C. Samples (500 μl) were harvested at times indicated and immediately mixed with sodium azide (1% w/v final concentration) to arrest growth and genome replication. For *S. aureus*, cells were grown overnight in TSB in the presence of chloramphenicol, followed by 1:100 dilution into TSB with or without aTc for two hours before harvesting samples. Cultures were collected by

centrifugation, the supernatant discarded, and pellets were flash frozen in liquid nitrogen before genomic DNA (gDNA) extraction using the DNeasy blood and tissue kit (Qiagen).

Quantitative PCR (qPCR)

qPCR was performed using the Luna qPCR mix (NEB) to measure the relative amount of origin DNA compared to the terminus. All PCR reactions were run in a Rotor-Gene Q instrument (Qiagen) using 20 µl reaction volumes in a Rotor-Disc 100 (Qiagen). Standard curves were obtained using the Rotor-Gene Q Software v2.0.2 (Qiagen) to calculate the efficiency of each primer pair, which varied ~5% between sets. Oligonucleotide primers designed to amplify *incC* (qSF19/qSF20 for *B. subtilis* or oDS263/oDS264 for *S. aureus*) and the terminus (qPCR57/qPCR58 for *B. subtilis* or oDS269/oDS270 for *S. aureus*) were typically 20-25 bases in length and amplified a ~100 bp PCR product (Supplementary Table 2). Individual *ori:ter* ratios were obtained in three steps: first, every Ct value was converted to $1/2^{Ct}$ and technical triplicates were averaged to generate a single enrichment value; second, origin enrichment was normalised by corresponding terminus values; third, relative *ori:ter* values were normalised by the enrichment obtained in control conditions. For *B. subtilis*, data was normalised to spore DNA as non-replicating control (*ori:ter* ratio = 1). For *S. aureus*, data was normalised to cells grown with the pRAB11 plasmid (empty vector used for the construction of *SaDnaA* variants) harvested at the same time point in exponential phase. Error bars indicate the standard error of the mean for 2-4 biological replicates.

Macrophage infection with *S. aureus* and gentamycin protection assay

RAW-Blue cells were seeded the day before bacterial infection experiments at 0.5×10^6 cells per ml in DMEM 5% FBS medium in 25 cm angled neck flasks (NUNC). Macrophages were challenged at a multiplicity of infection (MOI) of 1:5 in duplicates for *S. aureus* infection with CW1150 or CW1151 bacterial strains. The challenge was carried out in 10 mL of DMEM 5% FBS and cells were incubated for 1 hour at 37°C and 5% CO_2 to allow bacterial uptake by the macrophages. Following this incubation, gentamycin was added to all flasks (100 ug/ml

final concentration) to kill extracellular bacteria and aTc was supplemented to one of each of the duplicates (400 ng/ml final concentration) to induce DnaA variants overexpression. Flasks were then incubated for 20 hours at 37°C and 5% CO_2. Adherent cells were dislodged with a scrapper and 1 ml was transferred into an Eppendorf tube and spun down at 13,000 rpm. The supernatant was aspirated and the pellet was resuspended by pipetting up and down in 1 ml of Triton X100 (0.5% v/v) to lyse macrophages. Serial dilutions were performed and plated on TSA supplemented with 10 ug/ml chloramphenicol for selection of *S. aureus* carrying the inducible plasmid. Bacterial cells were incubated overnight at 37°C and used to quantify CFUs for survival plots. Macrophage experiments were performed as biological duplicates.

RESULTS

DNA replication hyper-initiation causes cell lysis in *B. subtilis*

A genetic system was employed to control the rate of DNA replication initiation in *B. subtilis.* This system exploits a hyperactive variant of DnaA (DnaA^{G154S}) that causes cells to over-initiate DNA replication [21]. To control hyper-initiation, a variant of the regulator ParA (ParA^{G12V}) was employed to inhibit DnaA activity [22]. The experimental strain encodes *dnaA*^{G154S} at the endogenous locus (under the control of its native expression system) and *parA*^{G12V} at an ectopic locus (under the control of a xylose-inducible promoter). Hereafter, we will refer to this strain (*dnaA*^{G154S} P_{xyl}-*parA*^{G12V}) as *Bs*^{HI} (for *B. subtilis* hyper-initiation). The *Bs*^{HI} strain was cultured in the presence of xylose to repress hyper-initiation, followed by growth in the absence of xylose to repress and dilute ParA^{G12V}, thereby enabling full activity of the hypermorphic DnaA^{G154S} variant (Fig. 1A).

Using this system, strains were streaked onto solid growth medium with or without xylose and incubated overnight. While both wild-type and *Bs*^{HI} strains showed similar growth in the presence of inducer, the *Bs*^{HI} strain displayed a severe growth defect in the absence of xylose (Fig. 1B). To further characterise this phenotype, cell growth in liquid media was followed over time. The results confirmed that the *Bs*^{HI} strain has a similar growth rate to wild-type in the presence of xylose, whereas *Bs*^{HI} cells could not reach the same density and appeared to lyse after ~5 hours in the absence of the inducer (Fig. 1C).

The impact of hyper-initiation on cell viability was quantified by growing cells in liquid medium with or without xylose, followed by aliquoting a serial dilution of each sample onto growth medium containing xylose (i.e. to repress hyper-initiation and allow growth of any viable cells). This spot titre analysis showed a similar number of colony forming units (CFU) between wild-type and *Bs*^{HI} strains after four hours of growth. However, by six and eight hours of growth without xylose, the *Bs*^{HI} strain displayed a >100-fold decrease in CFU compared to both wild-type and *Bs*^{HI} cultured in the presence of xylose (Fig. 1D).

Marker frequency analysis (MFA) was used to determine the frequency of DNA replication initiation under these experimental conditions. MFA showed similar *ori:ter* ratios

for all cells in lag phase regardless of the presence of absence of inducer (two-hour timepoint, Fig. 1E). Strikingly, Bs^{HI} cells displayed a significantly increased $ori{:}ter$ ratio following four hours growth without xylose ($ori{:}ter$ = 26 in exponential phase), whereas the same strain grown in the presence of inducer only mildly over-initiated ($ori{:}ter$ = 6 compared to wild-type $ori{:}ter$ = 3, Fig. 1E). Taken together, the results suggest that under this growth regime of depleting ParA^{G12V} to allow full activity of DnaA^{G154S}, hyper-initiation of DNA replication precedes significant cell death.

Phase contrast microscopy was used to investigate the fate of single *B. subtilis* cells experiencing DNA replication hyper-initiation. Even in the presence of xylose, Bs^{HI} cells show an elongated phenotype compared to wild-type (Fig. 1F), indicating that the normal cell cycle is perturbed in this strain [23]. Incubation in the absence of xylose resulted in dramatic changes to the cell envelope ("bulging" phenotype), with many cells appearing phase light, a hallmark of cell lysis (Fig. 1F). These phenotypes can be seen in greater detail in time lapse videos of Bs^{HI} cells grown in the presence or absence of xylose (Supplementary Videos 1 and 2, respectively). Together, these data indicate that in *B. subtilis,* DNA replication hyper-initiation causes a loss of viability via cell lysis.

Hyper-initiation activates the DNA damage response and leads to membrane depolarisation

In bacteria it is established that severe DNA damage elicits a stress response (the SOS response), allowing time to repair the damage and segregate replicating chromosomes [23-25]. To induce the SOS response, the critical recombination protein RecA binds to single-stranded DNA and stimulates the autocleavage of LexA, the transcriptional repressor of the SOS regulon [26]. Upon cleavage, LexA can no longer bind to DNA to repress transcription, resulting in induction of the SOS response.

One of the SOS responsive genes most strongly upregulated in *B. subtilis* encodes the cell division inhibitor YneA [27]. We hypothesised that the cell elongation observed during hyper-initiation (Fig. 1F) is caused by induction of the DNA damage response. To test

this model, a fluorescent transcriptional reporter under control of the LexA-dependent *yneA* promoter (P_{yneA}-*gfp*) was transformed into *Bs*[HI] (Fig. S1A). As a control, it was confirmed using spot titre assays that addition of the *yneA* reporter did not alter the phenotype associated with hyper-initiation (Fig. S1B). Following growth in liquid media with or without xylose, expression of P_{yneA}-*gfp* was observed using fluorescence microscopy. While a minority of *Bs*[HI] cells grown in the presence of xylose appeared to express the GFP reporter, in sharp contrast the majority of cells displayed a GFP signal following depletion of ParA[G12V] for five hours (Fig. 2A). Induction of the SOS response suggests that DNA replication hyper-initiation leads to the production of DNA damage intermediates (i.e. single-stranded DNA) recognised by RecA.

Timecourse and timelapse microscopy, following the growth of hyper-initiating cells, indicated that morphological abnormalities and lysis occur after DNA damage is generated (Fig. 1F and Supplementary Video 2). To further explore the observed compromise to cell envelope integrity, strains were incubated with the fluorescent dye $DiBAC_4$(3), which preferentially enters cells with depolarised phospholipid membranes [28]. Under the conditions tested to investigate hyper-initiation in liquid medium, fluorescence microscopy indicated that membrane depolarisation is a rare event in wild-type *B. subtilis* (Fig. S2). However, a significant number of *Bs*[HI] cells displayed fluorescent signals following depletion of ParA[G12V] for six hours (Fig. 2B). Using an *hbs-gfp* cassette to visualise DNA [29], after eight hours of growth in the absence of xylose the *Bs*[HI] strain showed abnormal chromosome content, whereas only a mild over-replication phenotype was observed in the presence of xylose (Fig. 2C). Together with cell growth analyses (Fig. 1C-D), the results indicate that membrane depolarisation occurs after DNA damage is detected, at a similar time as the onset of cell lysis.

The lethal phenotype caused by DnaA[G154S] is independent of prophage induction

The bacterial DNA damage response is known to activate lysogenic bacteriophages that can contribute to cell lysis. To determine whether the lysis phenotype observed in the *Bs*[HI] strain

was due to prophage activation, the hyper-initiation system was transformed into a *B. subtilis* strain lacking the six prophage elements harboured in the parental lab strain (Δ6^{prophage}) [30]. Spot titre analyses showed that after eight hours growth in the absence of inducer, the Δ6^{prophage} *Bs^{HI}* strain displayed a ~100-fold decrease in CFU (Fig. 3A). MFA validated that Δ6^{prophage} *Bs^{HI}* cells experienced significant hyper-initiation (Fig. 3B). These results indicate that prophage induction cannot explain the growth defects observed in cells hyper-initiating DNA replication. Nonetheless, to avoid issues of prophage induction, all subsequent experiments were performed using the Δ6^{prophage} genetic background.

The lethal phenotype caused by DnaA^{G154S} requires DNA replication hyper-initiation from *oriC*

In addition to being the master initiator of DNA replication in bacteria, DnaA also plays roles in diverse cellular processes such transcription regulation, chromosome organisation, cell division, cell differentiation, and metabolism [31, 32]. To ascertain whether the lethal phenotype observed with DnaA^{G154S} was dependent upon hyper-initiation from the endogenous replication origin *oriC*, a strain that replicates from a DnaA-independent origin *oriN* (*Bs^{HI} oriN*) was constructed, thus allowing mutagenesis of an essential DnaA-box within *oriC* (*Bs^{HI} oriN incC^{ΔBox6}*) [33]. Grown in the absence of xylose for eight hours, the *Bs^{HI} oriN* strain displayed a similar drop in CFU to the parental strain, indicating that the presence of *oriN* does not alleviate the hyper-initiation phenotype (Fig. 3C). In contrast, the *Bs^{HI} oriN incC^{ΔBox6}* strain showed no decrease in CFU following growth in the absence of xylose (Fig. 3C). MFA confirmed that the *Bs^{HI} oriN* strain hyper-initiates DNA replication, whereas the addition of the *incC^{ΔBox6}* mutation alleviated the ability of DnaA^{G154S} to over activate *oriC* (Fig. 3D). These results indicate that hyper-initiation of DNA replication is necessary for DnaA^{G154S} to elicit cell death.

Suppression of lethal DNA replication hyper-initiation by inactivation of base excision repair systems that target oxidised nucleobases

Studies in *E. coli* have suggested that DNA replication hyper-initiation causes DNA damage when the replisome encounters DNA repair events occurring on the chromosome [14, 15]. As the DNA repair enzymes must excise damaged bases before replacing them, the resulting single-stranded DNA gaps can promote replisome collapse and the emergence of DNA double-strand breaks (DSB) [34, 35]. Under aerobic growth conditions, nucleobase oxidation has been proposed to be the major source of DNA damage requiring repair [36, 37]. To explore this model, genes encoding factors required for recognising and initiating excision repair of oxidised purine nucleobases (*mutM* or *mutY* encoding the glycosylases MutM/MutY, respectively) were deleted in the Bs^{HI} strain. Spot titre analyses revealed that removing either of these DNA repair genes supressed the lethal phenotype observed in the parental Bs^{HI} strain (Fig. 3E). Critically, MFA showed that the strains harbouring either *mutM* or *mutY* knockouts remained capable of hyper-initiating DNA replication (Fig. 3F). Taken together, the data suggest that excessive replication from *oriC* can lead to conflicts between DNA replication and base excision repair, resulting in production of DNA damage, induction of the SOS response, and ultimately cell death.

Hyper-initiation induced cell death is conserved in *S. aureus* but independent of lysis

The replication initiator protein DnaA is broadly conserved in prokaryotes and we wondered whether DNA hyper-initiation could be exploited to inhibit the growth of bacterial pathogens. To test this idea, the consequence of DNA replication hyper-initiation was determined in *S. aureus*. An ectopic copy of *S. aureus* wild-type *dnaA* (encoding SaDnaA) or *dnaA*^{G159S} (encoding SaDnaA^{G159S}, analogous to *B. subtilis* DnaA^{G154S}) was placed under the control of an anhydrotetracycline-inducible (aTc) promoter on a self-replicating plasmid (Fig. 4A) [19, 38].

To characterise the effect of the SaDnaA^{G159S} variant, strains were grown in liquid media in the absence or presence of aTc (e.g. repression or induction of *SaDnaA*/*SaDnaA*^{G159S}). Expression of SaDnaA^{G159S} significantly inhibited growth of the culture, while expression of wild-type SaDnaA led to a longer lag phase but comparable

doubling-time during exponential growth (Fig. 4B). These results are consistent with a previous report that overexpression of SaDnaA in *S. aureus* is toxic [38].

To quantify the effect of SaDnaA^{G159S} on cell viability, cultures were grown overnight in the absence of aTc, then serially diluted and aliquoted onto solid media with or without inducer. Spot titre analyses showed that in the absence of inducer to express the *S. aureus dnaA* alleles, each strain produced a similar number of CFU, indicating that these plasmids are not cytotoxic under the conditions tested (Fig. 4C). In the presence of aTc, both the strain with the empty vector and the strain expressing wild-type SaDnaA showed a small colony phenotype, but nonetheless produced similar CFU compared to cultures grown without inducer. In contrast, expression of SaDnaA^{G159S} was associated with a ~100-fold loss in CFU (Fig. 4C). MFA demonstrated that induction of the hyper-active allele *SaDnaA^{G159S}* led to a significant increase in the rate of DNA replication initiation compared to wild-type (Fig. 4D). Together, these data indicate that DNA replication hyper-initiation inhibits bacterial growth in *S. aureus*.

To follow the consequence of hyper-initiation at a single cell level, timelapse microscopy was used to observe the growth of a strain harbouring the plasmid expressing SaDnaA^{G159S}. Cultures were grown to early exponential phase, then spotted onto agarose pads in the presence or absence of inducer aTc. Over eight hours, uninduced *S. aureus* cells were able to grow and fill in the field of view without notable morphological abnormalities (Supplementary Video 3). However, in the presence of aTc, strong growth inhibition and a significant increase in cell size was observed (Supplementary Video 4). Note that in this context cell lysis was not detected, likely reflecting inherent differences between the cell envelopes of *S. aureus* and *B. subtilis* (i.e. thick staphylococcal cell wall).

To further understand the growth inhibition observed during hyper-initiation in *S. aureus*, fluorescence microscopy was employed to visualise DNA content and the cell membrane, using the dyes DAPI and Nile red, respectively. Wild-type and hyper-initiation SaDnaA proteins were induced with aTc in early exponential phase and cells were imaged after 90 minutes. Under these conditions, cells expressing wild-type SaDnaA showed

uniform DAPI staining (i.e. normal chromosome content, Fig. 4E). In contrast, cells expressing the SaDnaA^{G159S} variant appeared significantly larger (up to four times the volume of the wild-type control) with heterogeneous DNA content including many cells devoid of DNA (Fig. 4E). Thus, DNA replication hyper-initiation in *S. aureus* leads to chromosome loss and cell death.

Hyper-initiation decreases *S. aureus* survival in murine macrophages

S. aureus is an opportunistic pathogen able to survive within immune cells and can cause respiratory, gut, blood and skin infections [39]. During the immune response, macrophages engulf *S. aureus* and are thought to release reactive oxygen species, potentially creating DNA damage. We hypothesised that DNA replication hyper-initiation in *S. aureus* might provide synergy with DNA damage generated in macrophages to inhibit bacterial survival during infection.

To test this model, RAW 264.7 murine macrophage-like cells were infected with *S. aureus* harbouring plasmids encoding either wild-type or *SaDnaA^{G159S}* alleles. Following a one-hour incubation to enable phagocytosis, extracellular bacteria were killed using gentamycin treatment and cultures were incubated for 20 hours in the presence or absence of aTc. Macrophages were then harvested and lysed to assess the number of surviving intracellular *S. aureus* cells. In uninduced conditions, strains expressing SaDnaA and SaDnaA^{G159S} yielded similar *S. aureus* survival (Fig. 4F). In the presence of aTc, macrophages infected with wild-type *SaDnaA* showed a small reduction in bacterial load compared to uninduced cells (1.4x less CFUs). In contrast, expression of SaDnaA^{G159S} produced a 25-fold reduction in *S. aureus* survival compared to the control. Together, these results suggest that *S. aureus* experiences oxidative DNA damage within macrophages which is exacerbated by DNA replication hyper-initiation, thereby inhibiting bacterial proliferation.

DISCUSSION

Here we report that hyper-initiation of DNA replication, achieved via expressing variants of the master initiator DnaA, inhibits proliferation of both *B. subtilis* and *S. aureus*. In *B. subtilis*, hyper-initiation induces the DNA damage response, membrane depolarisation, morphological deformations, and cell lysis (Figs. 1-2). Importantly, these mechanisms cannot be attributed to the activation of prophage and are *oriC*-dependent, indicating that DNA replication hyper-initiation is causative of the induced lethality (Fig. 3A-D). Consistent with observations made in previous studies [14, 15, 40], limiting repair of oxidised nucleobases suppressed this lethality, suggesting that the source of stress during hyper-initiation likely results from conflicts between the replication and repair machineries, leading to the emergence of DSBs (Fig. 3E-F).

Inspired by these findings, we further found that hyper-initiation can be exploited to limit proliferation of the human pathogen *S. aureus* (Fig. 4). The ability of *S. aureus* to survive within immune cells is thought to contribute to bacterial persistence [39]. Therefore, exploring approaches to reduce the intracellular pool of bacteria could favour clearance of *S. aureus* associated infections. Under standard laboratory growth conditions, we note that hyper-initiation of DNA replication generates significant morphological defects associated with cell death. During infection, experiments showed that hyper-initiation can also decrease *S. aureus* survival in macrophages. We speculate that hyper-initiation and associated replication/repair conflicts provides synergy with reactive oxygen species produced by macrophages during infection, thereby resulting in increased DNA damage and strong bacterial growth inhibition.

Our findings open new avenues for the development of alternative antimicrobial strategies. In this context, while several adverse approaches have been proposed to inhibit bacterial DNA replication [41], hyper-activation of DNA replication remains relatively unexplored. Interestingly, clinical isolates of *Mycobacterium tuberculosis* with mutations in *dnaA* were found to modulate resistance upon exposure to different classes of antimicrobials [42]. Therefore, the development of novel compounds targeting DnaA to exploit hyper-

initiation as an antibiotic adjuvant may be an attractive alternative to potentiate antimicrobials targeting DNA synthesis.

AUTHOR CONTRIBUTIONS

HG, DS, JH, KM, HM and CW contributed to the conception/design of the work. HG, DS, KM, HM and CW generated results presented in the manuscript. HG and CW created figures. HG, HM and CW wrote the manuscript. KM, HM and CW edited the manuscript.

FUNDING

Research support was provided to JH by an MRC Newcastle Impact Acceleration Award (MR/X50290X/1). KM was supported by a UKRI Future Leader Fellowship (MR/W009587/1). HM was supported by a Wellcome Trust Senior Research Fellowship (204985/Z/16/Z) and a Wellcome Trust Discovery Award (225811/Z/22/Z). CW was supported by a Wellcome Trust Early-Career Award (226338/Z/22/Z).

Conflict of interest: none declared.

SUPPLEMENTARY TABLES

Supplementary Table 1. Strains used in this study.

Strain	Genotype / description	Reference
Δ6	*trpC2 ΔSPβ Δskin ΔPBSX Δprophage1 Δpks Δprophage3*	[8]
168CA (HM715)	*trpC2*	[43]
CW1095	*ΔrbsU Δagr (cat bla tetR P_{tet})*	This work
CW1150	*ΔrbsU Δagr (cat bla tetR P_{tet}-SaDnaA)*	This work
CW1151	*ΔrbsU Δagr (cat bla tetR P_{tet}-SaDnaA^{G159S})*	This work
DH5α	*F– Φ80lacZΔM15 Δ(lacZYA-argF) U169 recA1 endA1 hsdR17 (rK–, mK+) phoA supE44 λ– thi-1 gyrA96 relA1*	CGSC
HG53	*trpC2 amyE::cat(P_{vneA}-gfp)*	This work (P*yneA-gfp* [26])
HG135	*trpC2 ΔmutM::erm*	BGSC
HG356	*trpC2 ΔmutY::erm*	BGSC
HG431	*trpC2 Δ6 amyE::spc(P_{xyl}-parA^{G12V}) dnaA^{G154S}::kan ΔmutM::erm*	This work
HG433	*trpC2 Δ6 amyE::spc(P_{xyl}-parA^{G12V}) dnaA^{G154S}::kan ΔmutY::erm*	This work
HM946	*trpC2 amyE::spc(P_{xyl}-parA^{G12V}) dnaA^{G154S}::kan*	[21]
HM1970	*trpC2 Δ6 amyE::spc(P_{xyl}-parA^{G12V}) dnaA^{G154S}::cat*	This work
HM1971	*trpC2 Δ6 amyE::spc(P_{xyl}-parA^{G12V}) dnaA^{G154S}::kan*	This work
HM1974	*trpC2 Δ6 amyE::spc(P_{xyl}-parA^{G12V}) dnaA^{G154S}::kan sacA::cat(hbs-mgfp)*	This work
HM1981	*trpC2 Δ6 aprE::spc(P_{xyl}-parA^{G12V}) dnaA^{G154S}::kan amyE::cat(P_{vneA}-gfp)*	This work
HM1990	*trpC2 Δ6 aprE::spc(P_{xyl}-parA^{G12V}) dnaA^{G154S}::cat amyE::kan(P_{spac}-gam-gfp)*	This work
HM2014	*trpC2 Δ6 amyE::spc(P_{xyl}-parA^{G12V}) dnaA^{G154S}::cat spoIIIJ::(oriN kan tet)*	This work
HM2015	*trpC2 Δ6 amyE::spc(P_{xyl}-parA^{G12V}) dnaA^{G154S}-incC^{ΔBox6}-dnaN::cat spoIIIJ::(oriN kan tet)*	This work
RAW 264.7	Murine adult male macrophages	InvivoGen
RN4220	*ΔrbsU Δagr*	[44]

Supplementary Table 2. Primers used for qPCR.

Species	Name	Sequence
B. subtilis	qPCR57	5□-TTTGCATGAACTGGGCAATA-3□
B. subtilis	qPCR58	5□-TCCGAACATGTCCAATGAGA-3□
B. subtilis	qSF19	5□-GATCAATCGGGGAAAGTGTG-3□
B. subtilis	qSF020	5□-CATTTTTATCCTCCTAACGGATAATG-3□
S. aureus	oDS263	5□-ACATTTCTATATGCTAATGTGGCAAG-3□
S. aureus	oDS264	5□-AGTAGGTGCTGTGGATTGGTG-3□
S. aureus	oDS269	5□-TGACCACCATGTTGTGCTTC-3□
S. aureus	oDS270	5□-AAAAATGCAAGAAGTTGGTGTAAC-3□

REFERENCES

1. Haeusser, D.P. and P.A. Levin, *The great divide: coordinating cell cycle events during bacterial growth and division.* Curr Opin Microbiol, 2008. **11**(2): p. 94-9.
2. Cooper, G., *The Cell: A Molecular Approach*, ed. S.M.S. Associates. 2000.
3. Bell, S.P. and J.M. Kaguni, *Helicase loading at chromosomal origins of replication.* Cold Spring Harb Perspect Biol, 2013. **5**(6).
4. Yi, L. and X. Lu, *New Strategy on Antimicrobial-resistance: Inhibitors of DNA Replication Enzymes.* Curr Med Chem, 2019. **26**(10): p. 1761-1787.
5. Fujikawa, N., et al., *Structural basis of replication origin recognition by the DnaA protein.* Nucleic Acids Res., 2003. **31**(8): p. 2077-86.
6. Richardson, T.T., O. Harran, and H. Murray, *The bacterial DnaA-trio replication origin element specifies single-stranded DNA initiator binding.* Nature, 2016. **534**(7607): p. 412-6.
7. Pelliciari, S., et al., *Evidence for a chromosome origin unwinding system broadly conserved in bacteria.* Nucleic Acids Res, 2021. **49**(13): p. 7525-7536.
8. Winterhalter, C., et al., *The DNA replication initiation protein DnaD recognises a specific strand of the Bacillus subtilis chromosome origin.* Nucleic Acids Res, 2023. **51**(9): p. 4322-4340.
9. Jameson, K.H. and A.J. Wilkinson, *Control of Initiation of DNA Replication in Bacillus subtilis and Escherichia coli.* Genes (Basel), 2017. **8**(1).
10. Kaguni, J.M., *Replication initiation at the Escherichia coli chromosomal origin.* Curr Opin Chem Biol, 2011. **15**(5): p. 606-13.
11. O'Donnell, M., L. Langston, and B. Stillman, *Principles and concepts of DNA replication in bacteria, archaea, and eukarya.* Cold Spring Harb Perspect Biol, 2013. **5**(7).
12. Skarstad, K. and T. Katayama, *Regulating DNA replication in bacteria.* Cold Spring Harb Perspect Biol, 2013. **5**(4): p. a012922.
13. Wagner, J.K., K.A. Marquis, and D.Z. Rudner, *SirA enforces diploidy by inhibiting the replication initiator DnaA during spore formation in Bacillus subtilis.* Mol. Microbiol., 2009. **73**(5): p. 963-74.
14. Charbon, G., et al., *Oxidative DNA damage is instrumental in hyperreplication stress-induced inviability of Escherichia coli.* Nucleic Acids Res, 2014. **42**(21): p. 13228-41.
15. Simmons, L.A., et al., *Hyperinitiation of DNA replication in Escherichia coli leads to replication fork collapse and inviability.* Molecular Microbiology, 2004. **51**(2): p. 349-358.
16. Anagnostopoulos, C. and J. Spizizen, *Requirements for transformation in Bacillus subtilis.* J. Bacteriol., 1961. **81**(5): p. 741-6.
17. Grosser, M.R. and A.R. Richardson, *Method for Preparation and Electroporation of S. aureus and S. epidermidis.* Methods Mol Biol, 2016. **1373**: p. 51-7.
18. Hanahan, D., J. Jessee, and F.R. Bloom, *Plasmid transformation of Escherichia coli and other bacteria.* Methods Enzymol, 1991. **204**: p. 63-113.
19. Helle, L., et al., *Vectors for improved Tet repressor-dependent gradual gene induction or silencing in Staphylococcus aureus.* Microbiology (Reading), 2011. **157**(Pt 12): p. 3314-3323.
20. Schindelin, J., et al., *Fiji: an open-source platform for biological-image analysis.* Nat. Methods, 2012. **9**(7): p. 676-82.
21. Murray, H. and J. Errington, *Dynamic control of the DNA replication initiation protein DnaA by Soj/ParA.* Cell, 2008. **135**(1): p. 74-84.
22. Scholefield, G., J. Errington, and H. Murray, *Soj/ParA stalls DNA replication by inhibiting helix formation of the initiator protein DnaA.* EMBO J., 2012. **31**(6): p. 1542-55.
23. Au, N., et al., *Genetic composition of the Bacillus subtilis SOS system.* J Bacteriol, 2005. **187**(22): p. 7655-66.

24. Hill, T.M., et al., *sfi-independent filamentation in Escherichia coli Is lexA dependent and requires DNA damage for induction.* J Bacteriol, 1997. **179**(6): p. 1931-9.

25. Trusca, D., et al., *Bacterial SOS checkpoint protein SulA inhibits polymerization of purified FtsZ cell division protein.* J Bacteriol, 1998. **180**(15): p. 3946-3953.

26. Gozzi, K., et al., *Bacillus subtilis utilizes the DNA damage response to manage multicellular development.* NPJ Biofilms Microbiomes, 2017. **3**: p. 8.

27. Kawai, Y., S. Moriya, and N. Ogasawara, *Identification of a protein, YneA, responsible for cell division suppression during the SOS response in Bacillus subtilis.* Mol Microbiol, 2003. **47**(4): p. 1113-1122.

28. Te Winkel, J.D., et al., *Analysis of Antimicrobial-Triggered Membrane Depolarization Using Voltage Sensitive Dyes.* Front Cell Dev Biol, 2016. **4**: p. 29.

29. Kohler, P. and M.A. Marahiel, *Association of the histone-like protein HBsu with the nucleoid of Bacillus subtilis.* J Bacteriol, 1997. **179**(6): p. 2060-2064.

30. Westers, H., et al., *Genome engineering reveals large dispensable regions in Bacillus subtilis.* Mol Biol Evol, 2003. **20**(12): p. 2076-90.

31. Goranov, A.I., et al., *A transcriptional response to replication status mediated by the conserved bacterial replication protein DnaA.* Proc Natl Acad Sci U S A, 2005. **102**(36): p. 12932-7.

32. Scholefield, G., J.W. Veening, and H. Murray, *DnaA and ORC: more than DNA replication initiators.* Trends Cell Biol, 2011. **21**(3): p. 188-94.

33. Richardson, T.T., et al., *Identification of a basal system for unwinding a bacterial chromosome origin.* EMBO J, 2019. **38**(15): p. e101649.

34. Kuzminov, A., *Single-strand interruptions in replicating chromosomes cause double-strand breaks.* Proc Natl Acad Sci U S A, 2001. **98**(15): p. 8241-6.

35. Caldecott, K.W., *Causes and consequences of DNA single-strand breaks.* Trends Biochem Sci, 2024. **49**(1): p. 68-78.

36. Imlay, J.A., *The molecular mechanisms and physiological consequences of oxidative stress: lessons from a model bacterium.* Nat Rev Microbiol, 2013. **11**(7): p. 443-54.

37. Foster, P.L., et al., *Determinants of spontaneous mutation in the bacterium Escherichia coli as revealed by whole-genome sequencing.* Proc Natl Acad Sci U S A, 2015. **112**(44): p. E5990-9.

38. Kurokawa, K., et al., *Rapid exchange of bound ADP on the Staphylococcus aureus replication initiation protein DnaA.* J Biol Chem, 2009. **284**(49): p. 34201-10.

39. Pidwill, G.R., et al., *The Role of Macrophages in Staphylococcus aureus Infection.* Front Immunol, 2020. **11**: p. 620339.

40. Fan, X.Y., et al., *Oxidation of dCTP contributes to antibiotic lethality in stationary-phase mycobacteria.* Proc Natl Acad Sci U S A, 2018. **115**(9): p. 2210-2215.

41. Grimwade, J.E. and A.C. Leonard, *Blocking the Trigger: Inhibition of the Initiation of Bacterial Chromosome Replication as an Antimicrobial Strategy.* Antibiotics (Basel), 2019. **8**(3).

42. Hicks, N.D., et al., *Mutations in dnaA and a cryptic interaction site increase drug resistance in Mycobacterium tuberculosis.* PLoS Pathog, 2020. **16**(11): p. e1009063.

43. Kunst, F. and e. al., *The complete genome sequence of the Gram positive bacterium Bacillus subtilis.* Nature, 1997. **390**: p. 249-256.

44. Kreiswirth, B.N., et al., *The toxic shock syndrome exotoxin structural gene is not detectably transmitted by a prophage.* Nature, 1983. **305**(5936): p. 709-12.

FIGURE 1: Hyper-initiation of DNA replication causes lysis and cell death in *B. subtilis*. (A) Genetic system used to artificially modulate DNA replication hyper-initiation. Xylose induction controls expression of *parA^G12V^*, which enables ParA^G12V^ to downregulate the activity of the hyperactive DnaA^G154S^ variant resulting in repression of hyper-initiation. In the absence of xylose, *parA^G12V^* is no longer expressed and DnaA^G154S^ allows *B. subtilis* to hyper-initiate. **(B)** Restreaks showing the phenotype associated with *dnaA^G154S^ P_xyl_-parA^G12V^* (*Bs^HI^*) cells in the absence of xylose after 24 h growth. **(C)** Growth curves showing the arrest of cell growth in *Bs^HI^* cells after 6 h incubation in the absence of xylose. **(D)** Spot titre analyses showing the loss of viability observed in *Bs^HI^* cells after 6 h growth in the absence of xylose. **(E)** Marker frequency analyses demonstrate that *Bs^HI^* cells start to hyper-initiate DNA replication after 4 h growth in the absence of xylose. **(F)** Phase-contrast microscopy showing the lysis phenotype observed in *Bs^HI^* cells after 8 h growth in the absence of xylose. Scale bar indicates 10 μm. Red arrows highlight lysed cells. (B-F) Strains: Wild-type (168CA), *Bs^HI^* (HM946).

FIGURE 2: Hyper-initiation causes DNA damage followed by membrane depolarisation and cell lysis via an apoptotic-like mechanism. (A) Microscopy experiments showing activation of the DNA damage response in Bs^{HI} cells after 5 h growth in the absence of xylose. Phase contrast microscopy images are overlayed with fluorescence signal corresponding to expression of the P_{yneA}-*gfp* SOS regulon reporter. Strain: HG287. **(B)** Microscopy experiments showing membrane depolarisation in Bs^{HI} cells following 6 h of growth in the absence of xylose. Phase contrast microscopy images are overlayed with fluorescence signal corresponding to DiBAC$_4$(3) entering depolarised cells. Strain: HM946. **(C)** Microscopy experiments detect chromosome loss in Bs^{HI} cells in the absence of xylose. Microscopy images are merged between green fluorescence for visualisation of DNA content (Hbs-GFP signal) and red fluorescence for membrane stain (Nile red dye). Strain: HM1974. (A-C) Scale bar indicates 10 μm.

FIGURE 3: Cell lysis caused by hyper-initiation is not due to prophage induction, is dependent on the replication origin *oriC* and can be rescued by limiting DNA repair. (A, C, E) Spot titre analyses showing that the hyper-initiation induced lethal phenotype **(A)** is not due to the activity of major prophages, **(C)** is dependent on the presence of the functional replication origin *oriC* recognised by DnaA, **(E)** can be rescued by a knockout of DNA repair genes *mutM* or *mutY*. **(B, D, F)** Marker frequency analyses showing that **(B)** a strain lacking major prophages remains capable of hyper-initiating DNA replication, **(D)** the presence of the ectopic origin *oriN* does not affect the ability of Bs^{HI} cells to hyper-initiate, whereas combination with an inactive endogenous origin (*oriN oriC$^{\Delta Box6}$*) abolishes hyper-initiation, **(F)** deletion of mutM or mutY does not affect the ability of $dnaA^{G154S}$ P_{xyl}-$parA^{G12V}$ cells to hyper-initiate. (A-B) Strains: *Δ6* (CW483), Bs^{HI} *Δ6* (HM1971). (C-D) Strains: Bs^{HI} (HM1970), Bs^{HI} *oriN* (HM2014), Bs^{HI} *oriN oriC$^{\Delta box6}$* (HM2015). (E-F) Strains: Bs^{HI} (HM1971), Bs^{HI} *ΔmutM* (HG431), Bs^{HI} *ΔmutY* (HG433).

FIGURE 4: Hyper-initiation in *S. aureus* causes cell death and reduces survival in murine macrophages. (A) Genetic system used to artificially modulate DNA replication hyper-initiation in *S. aureus*. Anhydrotetracycline (aTc) induction controls expression of *SaDnaA* variants (wild-type or hyper-initiator allele *SaDnaA*^{G159S}). **(B)** Plate reader analyses showing the absence of cell growth in cells harbouring the hyper-initiator allele *SaDnaA*^{G159S} in the presence of aTc. **(C)** Spot titre assay showing that a hyper-initiation allele (*SaDnaA*^{G159S}) inhibits the growth of *S. aureus*. **(D)** Marker frequency analyses showing that *SaDnaA*^{G159S} cells hyper-initiate DNA replication in the presence of aTc. Fold change *ori:ter* ratios are normalised to the empty vector control (*P*$_{tet}$). **(E)** Microscopy experiments showing that hyper-initiation (*P*$_{tet}$-*SaDnaA*^{G159S} cells) leads to the accumulation of drastic morphological changes. Cells harbouring wild-type *SaDnaA* show normal chromosome content whereas the *SaDnaA*^{G159S} variant is associated with loss of chromosome. Microscopy images are merged between cyan fluorescence for visualisation of DNA content (DAPI dye) and red fluorescence for membrane stain (Nile red dye). Scale bar indicates 2 μm. **(F)** Survival plot showing that *S. aureus* infection is attenuated when inducing the hyper-initiator allele *SaDnaA*^{G159S}. (B-F) Strains: Control *P*$_{tet}$ (CW1095), *P*$_{tet}$-*SaDnaA* (CW1150), *P*$_{tet}$-*SaDnaA*^{G159S} (CW1151).

A

B

SUPPLEMENTARY FIGURE 1: DNA damage response reporter in *B. subtilis*. (A) Genetic system used to artificially modulate hyper-initiation and to visualise the onset of the DNA damage response simultaneously via the P_{yneA}-*gfp* reporter cassette. **(B)** Spot titre assay showing that the P_{yneA}-*gfp* reporter cassette does not affect the lethal phenotype observed upon induction of hyper-initiation. Strains: Wild-type P_{yneA}-*gfp* (HG53), Bs^{HI} P_{yneA}-*gfp* (HM1981).

Membrane depolarisation: DiBAC$_4$(3)

SUPPLEMENTARY FIGURE 2: Membrane polarisation in wild-type _B. subtilis_. Microscopy experiments showing that wild-type _B. subtilis_ does not undergo membrane depolarisation in the presence or absence of xylose. Phase contrast microscopy images are overlayed with a green fluorescence channel (DiBAC$_4$(3) dye). Scale bar indicates 10 μm. Strain: HM715.