THE GEOMETRY OF GAMETIC DISPERSAL IN A FLYING MAMMAL, *RHINOLOPHUS HIPPOSIDEROS*

Thomas Brazier[1,2*] , Diane Zarzoso-Lacoste[3] , Lisa Lehnen[4,5] , Pierre-Loup Jan[1,6],
Sebastien J. Puechmaille[4,7,8] , Eric J. Petit[1]

[1] DECOD (Ecosystem Dynamics and Sustainability), INRAE, Ifremer, Institut Agro, Rennes, France
[2] ECOBIO UMR 6553 (Ecosystems, Biodiversity, Evolution), University of Rennes, CNRS, Rennes, France
[3] EDYSAN UMR 7058, Ecologie et Dynamique des Systèmes Anthropisés, CNRS, Université de Picardie
Jules Verne, Amiens, France
[4] Applied Zoology and Nature Conservation, Zoological Institute and Museum, University of Greifswald,
Greifswald, Germany
[5] Senckenberg Biodiversity and Climate Research Center, Frankfurt am Main, Germany
[6] CEBC, Centre for Biological Studies of Chizé, CNRS, Villiers-en-bois, France
[7] ISEM, University of Montpellier, CNRS, EPHE, IRD, Montpellier, France
[8] Institut Universitaire de France, Paris, France

Corresponding author: thomas.brazier@univ-rennes.fr

October 24, 2024

Abstract

Dispersal influences population and evolutionary dynamics, in a manner that depends on
by which dispersal strategies gene flow occurs. In some species, mating partners move exclu-
sively for mating, dispersing genes but not individuals. This is the case in many bat species,
of which the lesser horseshoe bat (*Rhinolophus hipposideros*) shows a genetic structure at
a fine spatial scale suggesting restricted dispersal. We investigated how natal and mating
dispersal shape gene flow in this species in two meta-populations using paternity and popu-
lation assignments. Half of the inferred paternities were intra-colonial and gave an estimate
of the mean mating dispersal distance of around 11 km, explaining the observed genetic
structure. Complete gametic dispersal distances were further estimated by combining natal
with mating dispersal distances. The resulting gametic dispersal kernels showed a mean
distance of around 20 km and a fat-tailed distribution typical of an excess of long-distance
dispersal movements. It is the first time that natal and mating dispersal distances have been
separately estimated and then combined in animals, documenting quantitatively how mating
dispersal decorrelates gene and individual flows. It is important to consider this mechanism
to explain dispersal evolution.

Keywords: gametic dispersal, mating behaviour, gene flow, sex-biased dispersal, parentage assignment,
dispersal kernel

Introduction

Individuals or propagules may move between their birthplace and any other place, habitat, social group, patch, etc. These movements, called dispersal, change the composition of the community they leave and the community they settle in, modify the demography and trait distribution of both the population they leave and the population they settle in, and exchange genes between populations. As individuals move outside their natal site, dispersal is involved in range expansion and colonizations, the rescue of small populations from extinction, invasiveness of exotic species but also the spread of pathogens or diseases, and it affects source-sink dynamics (Brown and Kodric-Brown, 1977; Gadgil, 1971; Hastings et al., 2004; Perkins et al., 2013; Phillips et al., 2008). In addition, dispersing individuals contribute to gene flow, affecting local adaptation or the spread of beneficial mutations, for example (Peischl and Gilbert, 2020; Tigano and Friesen, 2016). Dispersal is thus of crucial importance for both ecology and evolution, since population and evolutionary dynamics are intertwined through, among others, this particular trait.

Dispersal strategies are largely diversified among species (Bowler and Benton, 2005), mostly depending on proportions of dispersing individuals, mobility, and life-stage involved (Broquet and Petit, 2009). For example, natal dispersal, the definitive movement of offspring outside their natal location, generally occurs before the first reproduction event. On the contrary, mating dispersal is a later temporary movement relative to the pairing of opposite-sex individuals from different resident sites aimed for reproduction (Hazlitt et al., 2006; Sugg et al., 1996). Dispersal can be gametic and/or zygotic, depending on which propagules are involved in movements (gametes vs. individuals). This is most easily illustrated in plants, in which seed dispersal is zygotic while pollen dispersal is gametic. Natal dispersal can be either gametic or zygotic (see the example with plants) whereas mating dispersal does not involve a permanent movement of parents and leads to gametic dispersal only. Thus, natal and mating dispersal contribute to different degrees to the dispersal of individuals and gene flow (Waser and Elliott, 1991).

Importantly, male gametic dispersal may result from the addition of different dispersal behaviours: in mobile species, the addition of natal dispersal movements to mating dispersal movements results in male gametes moving further than the male home site. Natal dispersal is assessable by multiple methods (e.g. Capture-Mark-Recapture, direct observation or tagging). Mating dispersal, in contrast, involves home site fidelity and only short (sometimes a few hours), temporary movement to another site and is thus difficult to assess. Consequently, the contribution of mating dispersal to gene flow may be underestimated, because movements for mating often remain cryptic. Yet, mating dispersal has been documented in invertebrates, fishes, lizards, birds and mammals (Archie et al., 2008; Chapple and Keogh, 2005; Johannesen and Lubin, 1999; Kesler et al., 2010; Portnoy et al., 2015).

We aim to characterize the geometry of male gametic dispersal by following the spatial displacement of sperm through both natal and mating dispersal movements, adopting a dispersal distance kernel perspective (Nathan et al., 2012).

The lesser horseshoe bat, *Rhinolophus hipposideros* (André, 1797), is an interesting species to study natal and mating dispersal separately. Like most temperate bat species, it is sedentary and females are philopatric. Dispersal is thus supported by males (Bontadina et al., 2002; Gaisler and Chytil, 2002). As in many other bat species, little is known about the mating system and male dispersal (McCracken and Wilkinson, 2000). Because male-biased dispersal is pervasive among mammals (Dobson, 1982; Greenwood, 1980; Lawson Handley and Perrin, 2007), including bats (Kerth et al., 2002; Petit et al., 2001), it is assumed that males actively disperse and participate to gene flow through natal dispersal and mating with females that are not from their natal colony. In such species, natal and mating dispersal movements need to be studied simultaneously to reveal their relative contribution to gametic dispersal and ultimately gene flow (Archie et al., 2008; Chapple and Keogh, 2005; Kerth and Morf, 2004).

Lehnen et al. (2021) recently demonstrated that the lesser horseshoe bat's genetic structure follows an isolation-by-distance pattern at a scale of a few kilometres in two different meta-populations. This pattern is unusual among bats, suggesting very limited gene flow in this species (Aguillon et al., 2017; Rousset, 1997). We thus tested the hypotheses that (1) some lesser horseshoe bat males adopt a philopatric behaviour (Entwistle et al., 2000) and (2) males mate preferably with females of their home colony, which is either their natal colony or the colony they settle in after their natal dispersal movement, if any. We used molecular methods, assigning fathers to their colony of birth for estimating natal dispersal, and assigning offspring to their father to estimate mating dispersal (Broquet and Petit, 2009; Dussex et al., 2016; Pemberton, 2008). Natal and mating dispersal distances were then combined in a continuous-space model to estimate the complete male gametic dispersal distance kernel.

Materials and methods

Study meta-populations

We analysed data previously gathered on two lesser horseshoe meta-populations with different population dynamics (Jan et al., 2019; Lehnen et al., 2021; Zarzoso-Lacoste et al., 2018). The French meta-population (Picardy) was stable while the German one (Thuringia) was expanding. Seventeen colonies were sampled over four years (2013-2016) in Picardy (France), and 19 colonies were sampled over three years (2015-2017) in Thuringia (Germany). Distances between colonies ranged from 0.7 to 56 km in Picardy and from 1.8 to 99 km in Thuringia.

Genotype dataset

The dataset comprised all unique genotypes obtained from bats faeces sampled in the colonies during the respective study periods. Each unique genotype of six to eight microsatellite markers was considered as one individual (see Zarzoso-Lacoste et al. 2018, 2020 for the complete procedure). There were two sampling sessions per year, during summer, before and after parturition. All individuals sampled for the first time after parturition were considered as putative offspring. Females and males were separated by the sexual marker DDX3X/Y-Mam with the method described by Zarzoso-Lacoste et al. (2018, 2020).

Parentage assignment

Maternities and paternities were inferred with COLONY 2 (Jones and Wang, 2010) version 2.0.6.4 for Linux. COLONY was run with a full-pedigree likelihood method. Following Foley et al. (2020), parental links were inferred from 30 independent COLONY runs of the same dataset with different random seeds to achieve convergence on the best putative father. The eight microsatellite markers were used in the parentage assignment procedure. As COLONY is able to accommodate for genotyping errors, allelic dropout rates and other typing errors at each locus were set with values estimated by Zarzoso-Lacoste et al. (2018). Females were considered as polygamous with only one offspring per year (the father could change every year) and males as polygamous with no offspring limitation. Detection probabilities for mothers and fathers were set to 0.9 for mothers and 0.56 for fathers (details in Supplementary Information). Cohorts of the different years (four in Picardy, three in Thuringia) were analysed together.

In this species, both sexes are sexually mature at the end of their first year (Gaisler, 1966). Thus, putative offspring were all individuals sampled for the first time after parturition in any given year and these potential offspring were assumed to become candidate parents the year after their birth. Candidate parents were assumed to breed every year. As a consequence, for a given offspring, candidate mothers (respectively fathers) were all females (respectively males) born before the year of sampling of the offspring.

Then, we built lists of excluded paternities and maternities (Jones and Wang, 2010). Excluded paternities were all males born the same or later years for each given offspring. Excluded maternities were also all females born the same or later years for each given offspring, with an additional exclusion of all females from a different colony than the offspring colony as offsprings stay in their mother's colony for at least several weeks after birth. Females give birth to at most one offspring per year (Gaisler, 1966), so all offspring born the same year as the focal one were excluded from maternal sibship.

As in Foley et al. (2020), the comparison of simulated parent/offspring genotypes and the parentage inferred by COLONY was used both to find a statistic discriminant that allowed to exclude wrong paternities and to perform a power analysis of the categorical assignment method (i.e. type I and type II error rates). We simulated genotypes sampled in a theoretical population with known parent-offspring relationships and used it as an input in COLONY to assess type I (the wrong parent was inferred) and type II (no parent was found among the sampled ones) errors. We simulated the genotype datasets with the COLONY simulation module for Windows version 2.0.6.4 (Wang, 2013) to mimic demography, genetic diversity and genotyping errors observed in each sampled meta-population (details in supplementary materials and methods). Each simulated dataset was replicated 50 times with different random seeds then analysed with COLONY to assess for a convergence on true fathers over multiple replicated runs.

Male mating dispersal

The empirical distribution of mating dispersal distances was directly assessed by pairwise distances between the home colony of offspring and fathers inferred with categorical assignment (i.e. COLONY). We compared

these estimates to expected values obtained by a random association between true juveniles (candidate juveniles for which a true parent was found) and candidate fathers. A 95% confidence envelope for this null expectation was obtained by using a bootstrap procedure. The expected and observed distributions of offspring-father distances were compared with a Kolmogorov-Smirnov test. Besides, the exponential parameter β of the mating dispersal distance kernel was estimated by a Bayesian approach provided in the MasterBayes package (Hadfield et al., 2006) for cross-validation.

Male natal dispersal

The natal colony of male individuals was assessed by genetic assignment of fathers to colonies in the dataset with STRUCTURE (Pritchard et al., 2000). Male adults were assigned to colonies based on the genetic structure inferred from all female individuals with the USEPOPINFO model, as described in Dussex et al. (2016). Natal dispersal was then estimated by calculating the distance between the source colony inferred by STRUCTURE and the colony where the male was sampled. Only migrants of the actual generation were searched (GENSBACK = 0). The colony with the highest assignment probability was retained as the source colony for migrants. Ten replicated STRUCTURE analyses were run with a total of 2,000,000 iterations including a burnin of 1,000,000 (thinning = 100). Within-chain convergence was assessed with trace plots and diagnostics implemented in the CODA R package (Plummer et al., 2006). Replicates were compared and the replicate minimizing the Geweke statistics for both Alpha and Log-Likelihood sampling chains was retained for categorical assignments.

Male gametic dispersal

Parameters of the gametic dispersal distance kernel were estimated based on the distribution of empirical distances. Gametic dispersal distances were the result of both natal and mating dispersal; in other words, the distance between the natal colony of the male and the natal colony of its offspring. Competing probabilistic models were fitted to the data with the FitDistRplus package to estimate the parameters of the dispersal kernel with maximum likelihood estimation. Goodness-of-fit were compared based on AICc (Mazerolle, 2020) and BIC criterion (Delignette-Muller and Dutang, 2015). The most classical dispersal distributions (Gaussian, Exponential, Exponential Power, Weibull and Gamma) were compared (Nathan et al., 2012).

Additionally, we investigated hurdle models to deal with the excess of zeroes in dispersal distances (Lachenbruch, 2002). Hurdle models were constructed as a mixture of a binomial distribution for the probability of mating inside/outside of the colony and a continuous distribution excluding null dispersal distances. The parameter of the binomial distribution was estimated with a Generalized Linear Model in R and parameters of the positive continuous distribution were estimated with FitDistRplus. Parameters' confidence intervals were computed by bootstrapping. The log-likelihood of hurdle models was the sum of the log-likelihoods of the binomial model and the conditional model separately estimated (McDowell, 2003).

Statistical analysis

All statistical analyses were conducted with R version 4.1.2 (R Core Team, 2022). From the distributions of observed mating and gametic offspring-father distances, we estimated the mean and median dispersal distances with a bootstrap method.

Results

Dataset

Of the 3,706 unique genotypes in Picardy (France), 1,602 individuals were labelled as offspring (43%). 72% of individuals considered as adults were females (58% to 92% females depending on colony). 21% of individuals had an incomplete genotype (1 or 2 missing loci). The mean sample size per colony was 218±36 (s.e.m.) individuals for cumulated years 2013-2016, ranging from a minimal sample size of 53 bats to a maximal sample size of 746 bats. In Thuringia (Germany), there were 3,913 unique genotypes with 2,128 potential offspring (54%). 62% of individuals considered as adults were females. 19% of individuals had an incomplete genotype (1 or 2 missing loci). The mean sample size per colony was 126±22 (s.e.m.) for years 2015-2017 (range: 10 to 385).

Simulations

The proportion of successful father assignments was low with a high prevalence of type II errors (no father found for a given offspring, Table S1). In simulations, a conservative frequency value of 0.3 in Picardy and 0.25 in Thuringia accurately excluded wrong fathers when at least 20 runs of COLONY were used, even for the least powerful analysis with 6/8 complete loci (Fig. 1).

Figure 1: Assignment frequency of the inferred father according to the number of runs in simulated datasets. Black dots indicate cases where a true parent was found; grey dots indicate cases where the wrong parent was found for 3892 simulated offspring. Assignment probabilities of simulated father/offspring dyads with allelic frequencies of Picardy (A) or Thuringia (B). For each number of runs R, 100 assignment frequencies were estimated by randomly sampling R runs. The dashed line indicates the exclusion threshold of 0.3 (Picardy) or 0.25 (Thuringia) determined in the simulations.

Parentage assignment

We pooled inferences from genotypes with no more than 2 missing loci ($\geq 6/8$ loci) with the inferences from the complete genotypes (8/8 loci). Ambiguous assignments (different fathers for the same offspring) were resolved in favour of the most powerful dataset (8/8 complete loci). COLONY found 530 unique paternities (33% of putative offspring) in the French dataset. Based on simulation results (Fig. 1), we removed inferred paternities found in less than 30% of 30 runs, resulting in 82 paternities being considered as true paternities. There were 17 fathers responsible for these 82 paternities over four years. From the German dataset with the same procedure, 62 paternities were retained with only 18 fathers over three years.

Mating dispersal: where do the fathers come from?

Based on the empirical distribution of mating dispersal distances, the Kolmogorov-Smirnov test between the two distributions of offspring-father distances (expected and observed) showed a significant deviation (D = 0.40 in France and 0.53 in Germany, p < 0.001 in both cases, n = 82 and 62) with more zero distances than expected by chance (Fig. 2). The mean mating dispersal distance was 11.3 km (IC95 = [7.4; 15.7], 10,000 bootstraps, n = 82) in France and 11.6 km (IC95 = [7.3; 16.6], 10,000 bootstraps, n = 62) in Germany. However, the median distance were much lower in these two populations (7.7 km, IC95 = [5.5, 8.6] and 4.7 km, IC95 = [0, 7.8], respectively). Both empirical distributions were globally congruent (Fig. 2) and the MasterBayes Bayesian approach yielded similar results (Supplementary Information).

Natal dispersal: where were the fathers born?

Assignment to natal colonies with STRUCTURE indicated a high proportion of natal dispersers. Of 17 fathers in Picardy, eight were inferred as natal dispersers, yet their assignment probabilities were low with a mean probability of 0.13. There were 11 dispersing fathers in Thuringia (of a total of 18 fathers) with a mean assignment probability of 0.52. All sampling chains of STRUCTURE globally converged to stationarity and Geweke Z-scores of the selected replicate for Picardy were -0.45 (α) and 0.44 (Log-Likelihood). Geweke Z-scores for Thuringia were 0.06 (α) and 0.84 (Log-Likelihood).

Figure 2: Distribution of offspring-father mating dispersal distances in Picardy (A) and Thuringia (B). Black dots represent observed distances for accepted paternities (n = 82 offspring and 17 fathers in A and n = 62 offspring and 17 fathers in B). The grey ribbon marks the 95% confidence interval for expected distances between candidate fathers and true offspring under a random mating hypothesis, estimated with resampling methods. The black curve shows the theoretical Weibull dispersal kernel (i.e. cumulative density function of distances) with parameters fitted with fitditrplus on individual mating distances (dashed curves correspond to 95% confidence intervals estimated with 1,000 bootstraps). The grey curve shows the negative exponential cumulative density function of distances with the β parameter estimated by Masterbayes (dashed curves correspond to 95% confidence intervals estimated with 1,000 bootstraps).

Gametic dispersal: the whole sperm journey

The mean gametic dispersal distance (i.e. pairwise distance between the natal colony and the colony where the offspring was born resulting from both natal and mating dispersal) was slightly higher than for mating dispersal alone: 17.6 km (IC95 = [14.4; 20.7], 10,000 bootstraps, n = 82) in France and 22.2 km (IC95 = [15.2; 29.6], 10,000 bootstraps, n = 62) in Germany. However, as for mating dispersal, the median gametic dispersal

distances were lower than the mean (16.2 km, IC95 = [10.7, 18.9] and 6.9 km, IC95 = [4.1, 8.7], respectively), revealing a strong skewness of the distribution of distances. Upon the individual distances of gametic dispersal, model comparison based on AICc/BIC showed that hurdle models consistently outperformed all single continuous distributions (Table 1). When the conditional probability of mating inside/outside the colony and dispersal distances outside the colony were modelled separately in the hurdle model, the best fitting distribution in Picardy was the binomial + Weibull (Fig. S1). The proportion of gametic dispersal outside the natal colony, estimated by the binomial parameter, was 0.78 (IC95 = [0.68; 0.87], 1,000 bootstraps). The shape and scale parameters of the fitted Weibull were 1.90 (IC95 = [1.54; 2.37], 1,000 bootstraps) and 25.3 (IC95 = [20.9; 29.4], 1,000 bootstraps). In Thuringia, hurdle models outperformed all other ones and Log-Normal was the best fit, yet with very similar AICc/BIC values for different fat-tailed families ($\Delta_{AICc/BIC} \leq 3$; Table 1; Fig.S1). We selected the Weibull model for comparison between Picardy and Thuringia (Fig. 3) and compared it to the negative exponential dispersal kernel inferred by Lehnen et al. (2021) on a large European meta-population (mean distance = 16.77 km). The binomial parameter in Thuringia was slightly lower than Picardy, at 0.70 (IC95 = [0.56; 0.82], 1,000 bootstraps). The fitted Weibull had a shape of 0.99 (IC95 = [0.87; 1.16], 1,000 bootstraps) and a scale of 31.8 (IC95 = [22.4; 42.5], 1,000 bootstraps).

Table 1: Model comparison between competing probabilistic kernels of dispersal distances fitted to the empirical gametic dispersal distribution (i.e. paternity distances inferred with COLONY + natal dispersal inferred with STRUCTURE), based on the number of parameters of the model, AICc and BIC. The best model based on AICc/BIC and all competing models with $\Delta_{AICc/BIC} \leq 3$ are indicated by values in bold.

Model	N parameters	Thuringia		Picardy	
		AICc	BIC	AICc	BIC
Single continuous distribution					
Gaussian	2	600	604	675	679
Negative Exponential	1	**511**	**513**	**636**	**638**
Exponential power	3	549	555	747	754
Weibull	2	**513**	517	**638**	643
Gamma	2	2758	2762	2884	2889
Hurdle models					
Binomial + Gaussian	3	499	497	597	596
Binomial + Negative Exponential	2	**465**	**465**	617	617
Binomial + Exponential power	4	495	491	594	**590**
Binomial + Weibull	3	**467**	**465**	**590**	**588**
Binomial + Gamma	3	**467**	**465**	**592**	**590**
Binomial + Log-normal	3	**464**	**462**	599	598

Figure 3: Best model (AICc and BIC criterion) of gametic dispersal distance kernels fitted on strictly positive dispersal distances in Picardy (A) and Thuringia (B). Histograms show densities of empirical gametic dispersal distances. The best model is a Weibull function. In each case, our dispersal kernel (solid line) is compared to the negative exponential dispersal kernel inferred by Lehnen et al. (2021) (dashed line) based on the IBD slope on a large European meta-population (effective dispersal distance = 16.77 km, shape = 1/16.77).

Discussion

Both natal and mating dispersal mechanisms contribute to dispersal, and to its impacts on gene, population and community dynamics. These impacts depend on the distribution of dispersal distances, especially when long-distance dispersal is important (Nathan et al., 2012). To the best of our knowledge, our study is the first to consider both mating and natal dispersal to infer the complete gametic dispersal distance kernel in animals (Nathan, 2007; Rogers et al., 2019).

While mating dispersal has already been explicitly assessed on its own, the contribution of mating dispersal events to gametic dispersal is still scarcely investigated (Archie et al., 2008; Buchalski et al., 2014; Chapple and Keogh, 2005; Randall et al., 2007; Ruch et al., 2009). A previous study separately estimated mating and natal dispersal in a philopatric mammal with a similar indirect approach as ours, revealing that mating dispersal contributes most to gene flow in this species (Winters and Waser, 2003). The combination of pollen and seed dispersal recently received attention in plants (Browne et al., 2018). However, to our knowledge, the combination of mating and natal dispersal to estimate the total gametic dispersal kernel has never been reported in metazoans (Nathan, 2007; Nathan et al., 2012; Rogers et al., 2019).

Natal and mating dispersal both shape the geometry of the gametic dispersal distance kernel

Dispersal kernels are typically fitted to fat-tailed distributions (i.e. leptokurtic distributions) and most often clearly outperform the Gaussian and Exponential models (Nathan et al., 2012). The gametic dispersal kernel we inferred for the lesser horseshoe bat belongs to this family of models and shows a distribution already observed in many phylogenetically distant species (Browne et al., 2018; Catalano et al., 2020; Collin et al., 2013; Kesler et al., 2010; Paradis et al., 2002). For instance, fat-tailed dispersal kernels are common amongst plant studies (Austerlitz et al., 2004; Bullock et al., 2016; Robledo-Arnuncio and García, 2007). Our results are thus consistent with a large panel of dispersal studies, and fat-tailed dispersal kernels seem to be a common property applicable to both our meta-populations (Klein et al., 2006; Petrovskii and Morozov, 2009). The comparison of our dispersal kernels with the one inferred by Lehnen et al. (2021) across a wide European meta-population shows that our method is able to detect fine differences in kernel shape and leptokurtosis between two different meta-populations (see Fig. 3).

Although long-distance dispersal events seem relatively rare, they remain important for gene flow and shape the leptokurtic dispersal kernel. Most strikingly, some of the actively dispersing males are involved in extreme long-distance mating events. Long-distance dispersal is often dependent on individual variation, with few individuals expressing extreme behaviours (Hawkes, 2009; Kesler et al., 2010), as was already observed for this species (Heymer, 1964). Interestingly, we observed that mating dispersal introduces variance in individual behaviours that define the leptokurtic dispersal kernel (Petrovskii and Morozov, 2009). Indeed we reported low median dispersal distances and the mean effective dispersal distances were strongly impacted by extreme values. Thus, individual variation in mating behaviours should be considered as a strong component of the dispersal kernel. Niggemann et al. (2012) reported that uncertainty in the dispersal kernel increases for long distances. Parentage assignment permits to assess this individual variation, but underestimation is expected. Some long-distance dispersal may remain cryptic due to the low number of fathers inferred (n = 34) and the finite size of the study area. Yet, our study species usually flies less than 40 km (Lehnen et al., 2021) and the sampling spatial scale was twice this distance.

When sperm go further than individuals: mating dispersal decorrelates the flows of genes and individuals

At least 50% of males were moving temporarily outside their residence colony exclusively for mating, sometimes at very long distances, which suggests that mating dispersal is indeed a major component of gene flow in the lesser horseshoe bat.

Amongst mammals, longer mating than natal dispersal distances have been documented in kangaroo rats (*Dipodomys spectabilis*) and Spix's disc-winged bat (*Thyroptera tricolor*), two species characterized by strong philopatry of both sexes (Buchalski et al., 2014; Winters and Waser, 2003).

Mating dispersal plays the same role in metazoans as gamete dispersal in aquatic (mainly marine) organisms and pollen dispersal in terrestrial plants: it allows gametes to move further than the individuals producing gametes, ultimately decoupling meta-population dynamics and gene flow. This is an interesting context for enhancing our understanding of dispersal evolution. Available models and experiments highlight the role played by kin competition in favouring long-distance dispersal and thus fat-tailed dispersal distance

distributions (Bitume et al., 2013; Ronce, 2007; Rousset and Gandon, 2002). This makes sense when considering mating dispersal in species characterized by isolation-by-distance, such as the lesser horseshoe bat (Fig. S2). The relatedness between mating partners decreases when mating dispersal movements increase in distance, so that their offspring are less likely to live in a kin environment. By contrast, mating dispersal has no influence on local density or the probability to reach empty habitat patches, two factors that are also thought to drive the evolution of dispersal distances (Bitume et al., 2013; Ronce, 2007). If kin competition is a potential key driver of mating dispersal, it may interact with other selective forces such as dispersal costs and inbreeding load. An attempt by Ravigné et al. (2006) to model the interactions between these different factors on the joint evolution of seed and pollen dispersal at different distances provides an example for the kind of information that could be gained from modelling dispersal in species that rely both on zygotic and gametic dispersal with different life-stages and behaviours involved. In this model, pollen dispersal distance depends on seed dispersal, with the outcome being conditional on the relative dispersal costs associated with short- versus long-distance and seed versus pollen dispersal. Finally, our indirect approach decomposing the total gametic dispersal into natal and mating dispersal components should be a useful addition to study complex dispersal patterns in animal species in which gene flow and movements of individuals are decorrelated, such as phylopatric species.

Methodological limitations for inferences of mating and gametic dispersal

Only a few paternities were considered as reliable after a stringent selection based on simulation results. This contrasts with the high number of mothers recovered for the same colonies (Zarzoso-Lacoste et al., 2018), indicating that the reduced number of inferred paternities is not linked to a lack of informativeness of the set of markers. An alternative and more plausible explanation is that many potential fathers were not sampled. In particular, only males haunting maternity colonies were sampled, i.e. males searching a priori the proximity of females. Some of the undetected males may avoid maternity colonies to form male colonies or to live solitarily, yet this behaviour is known to diminish the mating success of fathers excluded from maternity colonies (Senior et al., 2005). In bat species, males seeking to access females generally reside in maternity colonies (Burland et al., 1999; Heckel and Von Helversen, 2003). Potential fathers are thus more likely to be among colonial fathers. Moreover, subsequent analyses will only be biased by this lack of potential fathers if unsampled fathers adopt a mating behaviour different from colonial fathers, which remains to be investigated. In contrast to other bat species (Burland et al., 1999; Petri et al., 1997; Rossiter et al., 2000), we found considerable reproductive skew amongst our colonial males, as only 17 and 18 males were responsible for all inferred paternities in Picardy and Thuringia, respectively.

The two methods we employed for estimating mating dispersal distances (a categorical approach with COLONY and a Bayesian probabilistic approach with Masterbayes) yielded similar results, suggesting that COLONY estimates are robust. We also estimated natal dispersal distances by categorical genetic assignment to colonies, based on the selection of the best replicate among independent STRUCTURE runs. The reliability of this approach has already been assessed in a previous study, and genetic assignment with STRUCTURE remains robust to identify individual dispersers, compared to other methods (Dussex et al., 2016). Finally, we combined mating and natal dispersal to estimate the gametic dispersal kernel based on a probabilistic modelling approach (Diefenbach et al., 2008; Williamson, 2002). Our results are in agreement with the dispersal kernel inferred by Lehnen et al. 2021.

Male philopatry and its implications

Mating dispersal distance is generally higher in other, swarming bat species (Kerth and Morf, 2004; Rivers et al., 2005), compared to what we observed in the lesser horseshoe bat. This species has a sedentary lifestyle and low dispersal ability, and swarming has not yet been reported for this species (Schofield et al., 2022). This results in a strong genetic structure (Fig. S2) compared to other sedentary and forest specialist species such as the Bechstein's bat (*Myotis bechsteinii*) (Kerth and Petit, 2005) or the brown long-eared bat (*Plecotus auritus*) (Burland et al., 1999). It is striking that genetic structure is stronger in the lesser horseshoe bat than the brown long-eared bat given that both are sedentary and that males and females share roosts in the latter one. However, in *Plecotus auritus*, males seek mating partners outside their colony (Burland and Worthington Wilmer, 2001).

Unexpectedly, we found 41 and 48% intra-colony paternities in two different meta-populations, which is a high percentage for a temperate bat (Kerth and Morf, 2004; Metheny et al., 2008) and also contrasts with swarming species (Burland et al., 1999; Kerth et al., 2003; Rivers et al., 2005). In the lesser horseshoe bat, pairs are formed between individuals of the same colony more often than by chance. If this trait is shared

among rhinolophids, it could explain intra-lineage polygyny and more generally, mating among relatives (Rossiter et al., 2005). Our results should be relevant to a finer comprehension of why species with apparent sex-biased dispersal show nevertheless genetic structure at a fine scale.

Conclusions

The population genetics approaches used here decomposed natal-dispersal and mating-dispersal components of the global gametic dispersal. This allowed to directly assess dispersal that would otherwise have remained cryptic if short-term movements for mating stayed undetected. Gene flow is intrinsically limited between lesser horseshoe colonies even when considering both natal and mating dispersal, despite long-distance dispersal events being more frequent than expected. We observed that variations in individuals mating behaviour shape a skewed dispersal kernel while promoting fine-scale genetic structure, with potential implications for genetic differentiation between populations. Hence, despite their limited number, the most actively dispersing males are important for the maintenance of genetic diversity in small breeding units.

Acknowledgments

We thank Gerald Kerth for sharing the Thuringia dataset ahead of publication. We thank every member of the INRAE DECOD lab (former ESE team) for their insighful discussions and support.

Statement of authorship

E.J.P. conceived the study; D.Z.-L., L.L. and P.-L.J. produced the data, T.B. analysed the data; T.B. and E.J.P. wrote the manuscript. All authors commented on the findings and reviewed the manuscript.

Data and code availability

The data reported in this article and the code to analyse the data are freely available in the INRAE data repository - DECOD collection (https://doi.org/10.57745/GBV4EA).

References

Aguillon, S. M., Fitzpatrick, J. W., Bowman, R., Schoech, S. J., Clark, A. G., Coop, G., & Chen, N. (2017). Deconstructing isolation-by-distance: The genomic consequences of limited dispersal (G. S. Barsh, Ed.). *PLOS Genetics*, *13*(8), e1006911. https://doi.org/10.1371/journal.pgen.1006911

Archie, E. A., Maldonado, J. E., Hollister-Smith, J. A., Poole, J. H., Moss, C. J., Fleischer, R. C., & Alberts, S. C. (2008). Fine-scale population genetic structure in a fission-fusion society: Fine-scale genetic structure in elephants. *Molecular Ecology*, *17*(11), 2666–2679. https://doi.org/10.1111/j.1365-294X.2008.03797.x

Austerlitz, F., Dick, C. W., Dutech, C., Klein, E. K., Oddou-Muratorio, S., Smouse, P. E., & Sork, V. L. (2004). Using genetic markers to estimate the pollen dispersal curve. *Molecular Ecology*, *13*(4), 937–954. https://doi.org/10.1111/j.1365-294X.2004.02100.x

Bitume, E. V., Bonte, D., Ronce, O., Bach, F., Flaven, E., Olivieri, I., & Nieberding, C. M. (2013). Density and genetic relatedness increase dispersal distance in a subsocial organism (J. Fryxell, Ed.). *Ecology Letters*, *16*(4), 430–437. https://doi.org/10.1111/ele.12057

Bontadina, F., Schofield, H., & Naef-Daenzer, B. (2002). Radio-tracking reveals that lesser horseshoe bats (*Rhinolophus hipposideros*) forage in woodland. *Journal of Zoology*, *258*(3), 281–290. https://doi.org/10.1017/S0952836902001401

Bowler, D. E., & Benton, T. G. (2005). Causes and consequences of animal dispersal strategies: Relating individual behaviour to spatial dynamics. *Biological Reviews*, *80*(2), 205–225. https://doi.org/10.1017/S1464793104006645

Broquet, T., & Petit, E. J. (2009). Molecular Estimation of Dispersal for Ecology and Population Genetics. *Annual Review of Ecology, Evolution, and Systematics*, *40*(1), 193–216. https://doi.org/10.1146/annurev.ecolsys.110308.120324

Brown, J. H., & Kodric-Brown, A. (1977). Turnover rates in insular biogeography: Effect of immigration on extinction. *Ecology*, *58*(2), 445–449.

Browne, L., Ottewell, K., Sork, V. L., & Karubian, J. (2018). The relative contributions of seed and pollen dispersal to gene flow and genetic diversity in seedlings of a tropical palm. *Molecular Ecology*, *27*(15), 3159–3173. https://doi.org/10.1111/mec.14768

Buchalski, M. R., Chaverri, G., & Vonhof, M. J. (2014). When genes move farther than offspring: Gene flow by male gamete dispersal in the highly philopatric bat species *Thyroptera tricolor. Molecular Ecology*, *23*(2), 464–480. https://doi.org/10.1111/mec.12593

Bullock, J. M., Gonza, L. M., White, S. M., & Pa, M. (2016). A synthesis of empirical plant dispersal kernels. *Journal of Ecology*, 14.

Burland, T. M., Barratt, E. M., Beaumont, M. A., & Racey, P. A. (1999). Population genetic structure and gene flow in a gleaning bat, *Plecotus auritus*, 6.

Burland, T. M., & Worthington Wilmer, J. (2001). Seeing in the dark: Molecular approaches to the study of bat populations. *Biological Reviews of the Cambridge Philosophical Society*, *76*(3), 389–409. https://doi.org/10.1017/S1464793101005747

Catalano, K. A., Dedrick, A. G., Stuart, M. R., Puritz, J. B., Montes, H. R., & Pinsky, M. L. (2020). Quantifying dispersal variability among nearshore marine populations. *Molecular Ecology*, mec.15732. https://doi.org/10.1111/mec.15732

Chapple, D. G., & Keogh, J. S. (2005). Complex mating system and dispersal patterns in a social lizard, *Egernia whitii. Molecular Ecology*, *14*(4), 1215–1227. https://doi.org/10.1111/j.1365-294X.2005.02486.x

Collin, S. B., Edwards, P. K., Leung, B., & Johnson, L. E. (2013). Optimizing early detection of non-indigenous species: Estimating the scale of dispersal of a nascent population of the invasive tunicate Ciona intestinalis (L.) *Marine Pollution Bulletin*, *73*(1), 64–69. https://doi.org/10.1016/j.marpolbul.2013.05.040

Delignette-Muller, M. L., & Dutang, C. (2015). **Fitdistrplus** : An *R* Package for Fitting Distributions. *Journal of Statistical Software*, *64*(4). https://doi.org/10.18637/jss.v064.i04

Diefenbach, D. R., Long, E. S., Rosenberry, C. S., Wallingford, B. D., & Smith, D. R. (2008). Modeling Distribution of Dispersal Distances in Male White Tailed Deer. *The Journal of Wildlife Management*, 8.

Dobson, F. S. (1982). Competition for mates and predominant juvenile male dispersal in mammals. *Animal behaviour*, *30*(4), 1183–1192.

Dussex, N., Broquet, T., & Yearsley, J. M. (2016). Contrasting dispersal inference methods for the greater white-toothed shrew. *The Journal of Wildlife Management*, *80*(5), 812–823. https://doi.org/10.1002/jwmg.21075

Entwistle, A. C., Racey, P. A., & Speakman, J. R. (2000). Social and population structure of a gleaning bat, *Plecotus auritus. Journal of Zoology*, *252*(1), 11–17. https://doi.org/10.1111/j.1469-7998.2000.tb00815.x

Foley, N. M., Petit, E. J., Brazier, T., Finarelli, J. A., Hughes, G. M., Touzalin, F., Puechmaille, S. J., & Teeling, E. C. (2020). Drivers of longitudinal telomere dynamics in the long-lived bat species, *Myotis myotis. Molecular Ecology*, mec.15395. https://doi.org/10.1111/mec.15395

Gadgil, M. (1971). Dispersal: Population Consequences and Evolution. *Ecology*, *52*(2), 253–261. https://doi.org/10.2307/1934583

Gaisler, J., & Chytil, J. (2002). Mark-recapture results and changes in bat abundance at the cave of Na Turoldu, Czech Republic, 174.

Gaisler, J. (1966). Reproduction in the lesser horseshoe bat (*Rhinolophus hipposideros*, Bechstein, 1800). *Bijdragen tot de Dierkunde*, *36*(1), 45–62.

Greenwood, P. J. (1980). Mating systems, philopatry and dispersal in birds and mammals. *Animal Behaviour*, *28*(4), 1140–1162. https://doi.org/10.1016/S0003-3472(80)80103-5

Hadfield, J. D., Richardson, D. S., & Burke, T. (2006). Towards unbiased parentage assignment: Combining genetic, behavioural and spatial data in a Bayesian framework. *Molecular Ecology*, *15*(12), 3715–3730. https://doi.org/10.1111/j.1365-294X.2006.03050.x

Hastings, A., Cuddington, K., Davies, K. F., Dugaw, C. J., Elmendorf, S., Freestone, A., Harrison, S., Holland, M., Lambrinos, J., Malvadkar, U., Melbourne, B. A., Moore, K., Taylor, C., & Thomson, D. (2004). The spatial spread of invasions: New developments in theory and evidence: Spatial spread of invasions. *Ecology Letters*, *8*(1), 91–101. https://doi.org/10.1111/j.1461-0248.2004.00687.x

Hawkes, C. (2009). Linking movement behaviour, dispersal and population processes: Is individual variation a key?: Movement behaviour, dispersal and population. *Journal of Animal Ecology*, *78*(5), 894–906. https://doi.org/10.1111/j.1365-2656.2009.01534.x

Hazlitt, S. L., Sigg, D. P., Eldridge, M., & Goldizen, A. W. (2006). Restricted mating dispersal and strong breeding group structure in a mid-sized marsupial mammal (*Petrogale penicillata*). *Molecular Ecology*, *15*(10), 2997–3007. https://doi.org/10.1111/j.1365-294X.2006.02985.x

Heckel, G., & Von Helversen, O. (2003). Genetic mating system and the significance of harem associations in the bat *Saccopteryx bilineata. Molecular Ecology*, *12*(1), 219–227.

Heymer, A. (1964). Résultats du baguage de chauve-souris dans les Pyrénées-Orientales de 1945 à 1959. *Vie et Milieu*, 765–800.

Jan, P.-L., Lehnen, L., Besnard, A.-L., Kerth, G., Biedermann, M., Schorcht, W., Petit, E. J., Le Gouar, P., & Puechmaille, S. J. (2019). Range expansion is associated with increased survival and fecundity in a long-lived bat species. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1906), 20190384. https://doi.org/10.1098/rspb.2019.0384

Johannesen, J., & Lubin, Y. (1999). Group founding and breeding structure in the subsocial spider *Stegodyphus lineatus* (*Eresidae*). *Heredity*, *82*(6), 677–686. https://doi.org/10.1046/j.1365-2540.1999.00528.x

Jones, O. R., & Wang, J. (2010). COLONY: A program for parentage and sibship inference from multilocus genotype data. *Molecular Ecology Resources*, *10*(3), 551–555. https://doi.org/10.1111/j.1755-0998.2009.02787.x

Kerth, G., Mayer, F., & Petit, E. (2002). Extreme sex-biased dispersal in the communally breeding, nonmigratory Bechstein's bat (*Myotis bechsteinii*). *Molecular Ecology*, *11*(8), 1491–1498. https://doi.org/10.1046/j.1365-294X.2002.01528.x

Kerth, G., Kiefer, A., Trappmann, C., & Weishaar, M. (2003). High gene diversity at swarming sites suggest hot spots for gene flow in the endangered Bechstein's bat. *Conservation Genetics*, *4*(4), 491–499.

Kerth, G., & Morf, L. (2004). Behavioural and Genetic Data Suggest that Bechstein's Bats Predominantly Mate Outside the Breeding Habitat. *Ethology*, *110*(12), 987–999. https://doi.org/10.1111/j.1439-0310.2004.01040.x

Kerth, G., & Petit, E. (2005). Colonization and dispersal in a social species, the Bechstein's bat (*Myotis bechsteinii*). *Molecular Ecology*, *14*(13), 3943–3950. https://doi.org/10.1111/j.1365-294X.2005.02719.x

Kesler, D. C., Walters, J. R., & Kappes, J. J. (2010). Social influences on dispersal and the fat-tailed dispersal distribution in red-cockaded woodpeckers. *Behavioral Ecology*, *21*(6), 1337–1343. https://doi.org/10.1093/beheco/arq158

Klein, E. K., Lavigne, C., & Gouyon, P.-H. (2006). Mixing of propagules from discrete sources at long distance: Comparing a dispersal tail to an exponential. *BMC Ecology*, *6*(1), 3. https://doi.org/10.1186/1472-6785-6-3

Lachenbruch, P. A. (2002). Analysis of data with excess zeros. *Statistical Methods in Medical Research*, *11*(4), 297–302. https://doi.org/10.1191/0962280202sm289ra

Lawson Handley, L. J., & Perrin, N. (2007). Advances in our understanding of mammalian sex-biased dispersal: Mammalian sex-biased dispersal. *Molecular Ecology*, *16*(8), 1559–1578. https://doi.org/10.1111/j.1365-294X.2006.03152.x

Lehnen, L., Jan, P.-L., Besnard, A.-L., Fourcy, D., Kerth, G., Biedermann, M., Nyssen, P., Schorcht, W., Petit, E. J., & Puechmaille, S. J. (2021). Genetic diversity in a long-lived mammal is explained by the past's demographic shadow and current connectivity. *Molecular Ecology*, *30*(20), 5048–5063. https://doi.org/10.1111/mec.16123

Mazerolle, M. J. (2020). *AICcmodavg: Model selection and multimodel inference based on (Q)AIC(c)*. Manual.

McCracken, G. F., & Wilkinson, G. S. (2000). Bat Mating Systems. In *Reproductive Biology of Bats* (pp. 321–362). Elsevier. https://doi.org/10.1016/B978-012195670-7/50009-6

McDowell, A. (2003). From the Help Desk: Hurdle Models. *The Stata Journal: Promoting communications on statistics and Stata*, *3*(2), 178–184. https://doi.org/10.1177/1536867X0300300207

Metheny, J. D., Kalcounis-Rueppell, M. C., Willis, C. K. R., Kolar, K. A., & Brigham, R. M. (2008). Genetic relationships between roost-mates in a fission–fusion society of tree-roosting big brown bats (*Eptesicus fuscus*). *Behavioral Ecology and Sociobiology*, *62*(7), 1043–1051. https://doi.org/10.1007/s00265-007-0531-y

Nathan, R. (2007). Total dispersal kernels and the evaluation of diversity and similarity in complex dispersal systems. In A. J. Dennis, E. W. Schupp, R. J. Green, & D. A. Westcott (Eds.), *Seed dispersal: Theory and its application in a changing world* (1st ed., pp. 252–276). CABI. https://doi.org/10.1079/9781845931650.0252

Nathan, R., Klein, E., Robledo-Arnuncio, J. J., & Revilla, E. (2012). Dispersal kernels: Review. In J. Clobert, M. Baguette, T. G. Benton, & J. M. Bullock (Eds.), *Dispersal Ecology and Evolution* (pp. 186–210). Oxford University Press. https://doi.org/10.1093/acprof:oso/9780199608898.003.0015

Niggemann, M., Wiegand, T., Robledo-Arnuncio, J. J., & Bialozyt, R. (2012). Marked point pattern analysis on genetic paternity data for uncertainty assessment of pollen dispersal kernels: Point pattern analysis on paternity data. *Journal of Ecology*, *100*(1), 264–276. https://doi.org/10.1111/j.1365-2745.2011.01854.x

Paradis, E., Baillie, S. R., & Sutherland, W. J. (2002). Modeling large-scale dispersal distances. *Ecological Modelling*, *151*(2-3), 279–292. https://doi.org/10.1016/S0304-3800(01)00487-2

Peischl, S., & Gilbert, K. J. (2020). Evolution of Dispersal Can Rescue Populations from Expansion Load. *The American Naturalist*, *195*(2), 349–360. https://doi.org/10.1086/705993

Pemberton, J. (2008). Wild pedigrees: The way forward. *Proceedings of the Royal Society B: Biological Sciences*, *275*(1635), 613–621. https://doi.org/10.1098/rspb.2007.1531

Perkins, A. T., Phillips, B. L., Baskett, M. L., & Hastings, A. (2013). Evolution of dispersal and life history interact to drive accelerating spread of an invasive species (J. Fryxell, Ed.). *Ecology Letters*, *16*(8), 1079–1087. https://doi.org/10.1111/ele.12136

Petit, E., Balloux, F., & Goudet, J. (2001). Sex-biased dispersal in a migratory bat: A characterization using sex-specific demographic parameters. *Evolution*, *55*(3), 635–640. https://doi.org/10.1111/j.0014-3820.2001.tb00797.x

Petri, B., Paabo, S., Von Haeseler, A., & Tautz, D. (1997). Paternity assessment and population subdivision in a natural population of the larger mouse-eared bat *Myotis myotis*. *Molecular Ecology*, *6*(3), 235–242. https://doi.org/10.1046/j.1365-294X.1997.00176.x

Petrovskii, S., & Morozov, A. (2009). Dispersal in a Statistically Structured Population: Fat Tails Revisited. *The American Naturalist*, *173*(2), 278–289. https://doi.org/10.1086/595755

Phillips, B. L., Brown, G. P., Travis, J. M. J., & Shine, R. (2008). Reid's Paradox Revisited: The Evolution of Dispersal Kernels during Range Expansion. *The American Naturalist*, *172*(S1), S34–S48. https://doi.org/10.1086/588255

Plummer, M., Best, N., Cowles, K., & Vines, K. (2006). CODA: Convergence Diagnosis and Output Analysis for MCMC. *R News*, *6*(1), 7–11.

Portnoy, D. S., Puritz, J. B., Hollenbeck, C. M., Gelsleichter, J., Chapman, D., & Gold, J. R. (2015). Selection and sex-biased dispersal in a coastal shark: The influence of philopatry on adaptive variation. *Molecular Ecology*, *24*(23), 5877–5885. https://doi.org/10.1111/mec.13441

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, *155*(2), 945–959.

R Core Team. (2022). *R: A language and environment for statistical computing*. Manual. R Foundation for Statistical Computing. Vienna, Austria.

Randall, D. A., Pollinger, J. P., Wayne, R. K., Tallents, L. A., Johnson, P. J., & Macdonald, D. W. (2007). Inbreeding is reduced by female-biased dispersal and mating behavior in Ethiopian wolves. *Behavioral Ecology*, *18*(3), 579–589. https://doi.org/10.1093/beheco/arm010

Ravigné, V., Olivieri, I., Martinez, S. G., & Rousset, F. (2006). Selective interactions between short-distance pollen and seed dispersal in self-compatible species. *Evolution*, *60*(11), 2257–2271. https://doi.org/10.1111/j.0014-3820.2006.tb01863.x

Rivers, N. M., Butlin, R. K., & Altringham, J. D. (2005). Genetic population structure of Natterer's bats explained by mating at swarming sites and philopatry. *Molecular Ecology*, *14*(14), 4299–4312. https://doi.org/10.1111/j.1365-294X.2005.02748.x

Robledo-Arnuncio, J. J., & García, C. (2007). Estimation of the seed dispersal kernel from exact identification of source plants. *Molecular Ecology*, *16*(23), 5098–5109. https://doi.org/10.1111/j.1365-294X.2007.03427.x

Rogers, H. S., Beckman, N. G., Hartig, F., Johnson, J. S., Pufal, G., Shea, K., Zurell, D., Bullock, J. M., Cantrell, R. S., Loiselle, B., Pejchar, L., Razafindratsima, O. H., Sandor, M. E., Schupp, E. W., Strickland, W. C., & Zambrano, J. (2019). The total dispersal kernel: A review and future directions (C. Aslan, Ed.). *AoB PLANTS*, *11*(5), plz042. https://doi.org/10.1093/aobpla/plz042

Ronce, O. (2007). How Does It Feel to Be Like a Rolling Stone? Ten Questions About Dispersal Evolution. *Annual Review of Ecology, Evolution, and Systematics*, *38*(1), 231–253. https://doi.org/10.1146/annurev.ecolsys.38.091206.095611

Rossiter, S. J., Jones, G., Ransome, R. D., & Barratt, E. M. (2000). Parentage, reproductive success and breeding behaviour in the greater horseshoe bat (*Rhinolophus ferrumequinum*). *Proceedings of the Royal Society B: Biological Sciences*, *267*(1443), 545–551. https://doi.org/10.1098/rspb.2000.1035

Rossiter, S. J., Ransome, R. D., Faulkes, C. G., Le Comber, S. C., & Jones, G. (2005). Mate fidelity and intra-lineage polygyny in greater horseshoe bats. *Nature*, *437*(7057), 408–411. https://doi.org/10.1038/nature03965

Rousset, F., & Gandon, S. (2002). Evolution of the distribution of dispersal distance under distance-dependent cost of dispersal: Evolution of dispersal distributions. *Journal of Evolutionary Biology*, *15*(4), 515–523. https://doi.org/10.1046/j.1420-9101.2002.00430.x

Rousset, F. (1997). Genetic Differentiation and Estimation of Gene Flow from *F* -Statistics Under Isolation by Distance. *Genetics*, *145*(4), 1219–1228. https://doi.org/10.1093/genetics/145.4.1219

Ruch, J., Heinrich, L., Bilde, T., & Schneider, J. M. (2009). The evolution of social inbreeding mating systems in spiders: Limited male mating dispersal and lack of pre-copulatory inbreeding avoidance in a subsocial predecessor. *Biological Journal of the Linnean Society*, *98*(4), 851–859. https://doi.org/10.1111/j.1095-8312.2009.01322.x

Schofield, H., Reiter, G., & Dool, S. E. (2022). Lesser Horseshoe Bat *Rhinolophus hipposideros* (André, 1797). In K. Hackländer & F. E. Zachos (Eds.), *Handbook of the Mammals of Europe* (pp. 1–34). Springer International Publishing. https://doi.org/10.1007/978-3-319-65038-8_39-1

Senior, P., Butlin, R. K., & Altringham, J. D. (2005). Sex and segregation in temperate bats. *Proceedings of the Royal Society B: Biological Sciences*, *272*(1580), 2467–2473. https://doi.org/10.1098/rspb.2005.3237

Sugg, D. W., Chesser, R. K., Stephen Dobson, F., & Hoogland, J. L. (1996). Population genetics meets behavioral ecology. *Trends in Ecology & Evolution*, *11*(8), 338–342. https://doi.org/10.1016/0169-5347(96)20050-3

Tigano, A., & Friesen, V. L. (2016). Genomics of local adaptation with gene flow. *Molecular Ecology*, *25*(10), 2144–2164. https://doi.org/10.1111/mec.13606

Wang, J. (2013). A simulation module in the computer program COLONY for sibship and parentage analysis. *Molecular Ecology Resources*, *13*(4), 734–739. https://doi.org/10.1111/1755-0998.12106

Waser, P. M., & Elliott, L. F. (1991). Dispersal and genetic structure in kangaroo rats. *Evolution*, *45*(4), 935–943. https://doi.org/10.1111/j.1558-5646.1991.tb04361.x

Williamson, M. (2002). Overview and synthesis: The tale of the tail. *Dispersal Ecology: The 42nd Symposium of the British Ecological Society Held at the University of Reading 2-5 April 2001*, 431–443.

Winters, J. B., & Waser, P. M. (2003). Gene dispersal and outbreeding in a philopatric mammal. *Molecular Ecology*, *12*(8), 2251–2259. https://doi.org/10.1046/j.1365-294X.2003.01896.x

Zarzoso-Lacoste, D., Jan, P.-L., Lehnen, L., Girard, T., Besnard, A.-L., Puechmaille, S. J., & Petit, E. J. (2018). Combining noninvasive genetics and a new mammalian sex-linked marker provides new tools to investigate population size, structure and individual behaviour: An application to bats. *Molecular Ecology Resources*, *18*(2), 217–228. https://doi.org/10.1111/1755-0998.12727

Zarzoso-Lacoste, D., Jan, P.-L., Lehnen, L., Girard, T., Besnard, A.-L., Puechmaille, S. J., & Petit, E. J. (2020). Corrigendum. *Molecular Ecology Resources*, *20*(6), 1787–1787. https://doi.org/10.1111/1755-0998.13254