Improving the monitoring of the invasive Blue Crab (*Callinectes sapidus*): combining environmental DNA and citizen observations

Erwan Delrieu-Trottin[1]*, Mathieu Bianic[1], Claude Miaud[1],
Manon Garcia[1], Véronique Arnal[1], Régis Hocdé[2], Christophe Cochet[3],
Nathalie Barre[4],Laetitia Cornil[5] and Stéphanie Manel[1,6]

[1]CEFE, Univ Montpellier, CNRS, EPHE-PSL University, IRD, Montpellier, France
[2]MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Montpellier, France
[3]ISEM, Univ Montpellier, CNRS, IRD, EPHE, CIRAD, INRAP, Montpellier, France
[4]Conservatoire d'Espaces Naturels (CEN) Occitanie, Montpellier, France
[5]Direction Régionale de l'Environnement, de l'Aménagement et du Logement (DREAL) d'Occitanie, Montpellier, France
[6]Institut Universitaire de France, IUF, Paris

Abstract

1. Early detection is a crucial tool for identifying and managing the spread of invasive species.

2. In this study, we validated a probe-based quantitative polymerase chain reaction (qPCR) assay for the detection of the invasive Blue Crab, *Callinectes sapidus*, in the Mediterranean Sea, using 22 initial eDNA environmental samples (eDNA) collected from three coastal lagoons.

3. A subsequent large-scale eDNA sampling campaign (61 samples in 31 sites), conducted in collaboration with local stakeholders, was carried out to map the distribution of *Callinectes sapidus* along the Occitanie coastline (Western Mediterranean, France).

4. Using eDNA probe-based qPCR, *Callinectes sapidus* was detected in 32 out the 61 samples (52%), confirming its presence in 24 out of 31 sites surveyed, including the 13 lagoons where its occurrence had already been reported, as well as two additional lagoons and at sea where no prior records existed.

5. Our results demonstrate the utility of eDNA probe-based qPCR for effective monitoring the invasive Blue Crab. The integration of eDNA analysis with citizen science observations enhances the monitoring framework, facilitating early detection and contributing to improved management strategies from a prevention perspective.

INTRODUCTION

Species that have been introduced to areas outside their natural ranges, known as non-native species, pose a major threat to marine biodiversity with notable socio-economic impacts once they become invasive (Alidoost Salimi et al., 2021; Giakoumi et al., 2019). The introduction of non-native species, and specifically invertebrates, has become a recurring phenomenon over the last century, largely attributed to globalization, international trade (Bailey et al., 2020; Seebens et al., 2021; Zenetos et al., 2022), and has been facilitated by climate change and maritime traffic (Pearman et al., 2021; Seebens et al., 2016). The arrival of these invasive non-native species can significantly jeopardize local biodiversity through phenomena such as competitive exclusion, predation, introgression or niche modification (Mooney & Cleland, 2001). Prevention is likely the most effective approach to managing such biological pressures and can be done by establishing monitoring programs to detect invasive species (Giakoumi et al., 2019).

The Blue Crab, *Callinectes sapidus* Rathbun, 1896, is indigenous to the estuaries and coastal waters of the western Atlantic coast (USA) where it is actively targeted by fisheries. Its presence in the Mediterranean Sea has been documented as early as 1930 (Galil et al. 2002, Lee et al., 2020). Recent scientific publications confirmed sightings of *Callinectes sapidus* in various locations across the Mediterranean Sea, including Spain, Algeria, Italy and France (Labrune et al. 2019, Mancinelli et al., 2021; Sabelli, 2023). Consequently, the French government has initiated a regional action plan (http://www.occitanie.developpement-durable.gouv.fr/presentation-a25426.html) to provide insights pertinent to the inquiries outlined in European directives concerning lagoon, coastal, and marine environments. As of 2022, Europe has strictly regulated 88 species, including 10 species of fish, 1 species of mollusk and 8 species of aquatic invertebrates including the Chinese mitten crab (*Eriocheir sinensis*, https://circabc.europa.eu/ui/group/4cd6cb36-b0f1-4db4-915e-65cd29067f49/library/79885406-e439-4961-ab2e-717191190f34/details) while *Callinectes sapidus* is in the process of being regulated.

*Corresponding author: erwan.delrieu-trottin@ephe.psl.eu

One first preventive action is the early detection of this species. Indeed, most reports of the presence of the species in French West Mediterranean Sea are made by fishermen whose nets are often damaged by *C. sapidus* in spring and summer - yet not all lagoons are surveyed, concentration are not the same in the different surveyed lagoons, there are no data for a large part of the potential habitat of *C. sapidus* while the question of its presence in lagoons in winter remains open.

Environmental DNA (eDNA) emerges as a promising solution, enabling early detection of invasive species in low-impacted sites or monitor their persistence in managed sites (Dejean et al., 2012; Kress et al., 2015; Taberlet et al., 2018). Specifically, quantitative polymerase chain reaction (qPCR)-based eDNA detection assays aim to identify a target species through the analysis of DNA present in water samples collected from target sites. Additionally, it can potentially allow for the estimation of a minimum density threshold for species detection (Klymus et al., 2020a, 2020b). The use of qPCR-based approach instead of the eDNA metabarcoding approach has been shown to enhance detection capabilities (McColl-Gausden et al., 2023; Tsuji et al., 2024; Yu et al., 2022). Inventories based on eDNA are particularly interesting in challenging environments such as turbid wetlands colonized by *Callinectes sapidus* (Saenz-Agudelo et al., 2022). Here, we first tested the specificity of a mitochondrial genetic marker to detect Blue Crab from environmental samples using qPCR-based eDNA detection assays. We then applied qPCR-based eDNA detection assays to monitor specifically the potential presence of *C. sapidus* across 31 sites comprising 17 lagoons, one channel between lagoons and the sea, three river mouths, one channel between lagoons, and one coastal site along the Occitanie coast (French West Mediterranean Sea). Comparing our eDNA-based data with citizen science data and the literature, we discuss how combining such data can improve the monitoring of the invasive Blue Crab.

Material and Methods

Study area, eDNA sampling and extraction

Sampling was done along the French Mediterranean coasts following the protocol of (Boulanger et al., 2021). Basically, each water eDNA sample consisted of 30 liters of seawater filtered using a 0.20μM filtration capsule over a 30-minute transect from a boat, using a 12 V peristaltic pump with a flow rate of one liter per minute. Immediately after filtration, the capsule was drained of any remaining water, filled with 80 mL of CL1 conservation buffer, and stored at room temperature until extraction. Three sampling campaigns were conducted in collaboration with regional stakeholders with different objectives:

1. To test the ability of our genetic marker to detect *Callinectes sapidus* through water filtration and our protocol of qPCR based eDNA assay, a total of 13 water eDNA samples were collected in situ and in an aquarium (Table S1). Two to three replicates were performed per site. Sampling in situ were conducted in two sites where a high concentration of *C. sapidus* had been reported by professional fishermen (Canet and Thau (Marseillan) lagoons) and at two periods: early in the year in March 2021 at the end of winter when *C. sapidus* is potentially dormant and later during summing (June-July 2021) when the species is active and more abundant. Two additional samples were collected from a 1500L tank hosting around forty *C. sapidus* individuals at the Biodiversarium (Banyuls-sur-Mer) in July 2021 (Table S1).

2. To assess the influence of Blue Crab density on detection, a total of 9 water eDNA samples were collected in July 2022 in three sites with different densities of *Callinectes sapidus*. In each site, three replicates were collected (Table S1). Sites were chosen in concertation with fishermen and stakeholder based, with two sites in Canet and one site in Méjean-Perols. In Canet, the three replicates collected nearby the channel between the lagoon and the sea were considered as the high density site as Blue Crab was often caught in fishing nets with high densities. A second site within the same lagoon (Roselière) corresponded to a lower density site of Blue Crab (lower capture rate). Finally, the three replicates collected in Méjean-Perols lagoon also corresponded to a low density sample site with only episodic captures.

3. With a spatial monitoring perspective, the presence of *Callinectes sapidus* along the French West Mediterranean Sea coast was evaluated sampling 31 sites from May to June 2023. Two replicates except for one site were collected each time, representing a total of 61 eDNA samples (Figure 1, Table S2). Overall, we sampled a diversity of coastal habitats, including (i)17 lagoons (Ayrolles, Bages-Sigean, Canet, Grazel, Gruissan, Ingril (Sud), La Palme, Médard, Méjean-Pérols, Or, Pierre Blanche, Ponant, Salses-Leucate, Salonique, Thau, Vendres, and Vic) (ii) one channel between lagoons and the sea (Grau du Prévost), (iii) three river mouths (Grau d'Agde (river mouth of the Hérault river), Grau de Vendres (river mouth of the Aude river) and Grau

D'Orgon (river mouth of the Petit Rhône river), (iv) one channel between lagoons (Canal du Rhône à Sète) and (v) one coastal site (Canet Mer). Given the size of some lagoons, we sampled up to four sites within a single lagoon (e.g. Thau). Overall, our study includes not only sites that are surveyed by professional sentinel fishermen where *C. sapidus* had already been reported such as Canet but also sites where no data were available at sampling time: Salonique lagoon, the channel between lagoons, the three river mouths and the channel between lagoon and the sea. *Callinectes sapidus* is suspected to be able to use these accesses to reach the sea or move from one lagoon to another.

qPCR-based eDNA detection assays

The DNA extraction of eDNA samples was performed at SPYGEN (Le Bourget du Lac, France), following the protocol published by Pont et al. (2018) for the first and the third campaign and in house in a dedicated room for water DNA sample extraction at CEFE for the second campaign (Faure et al., 2023). Some samples of the first campaign extracted at SPYGEN (Le Bourget du Lac, France) have also been re-extracted at CEFE to compare and validate the in-house extraction protocol. Negative extraction controls were carried out in parallel with each batch of eDNA extraction to monitor for potential contamination. Amplifications of the 83 eDNA samples were performed using qPCR following Faure et al., (2023). We used primers and PCR-probe designed by Andersen et al (2018), developed from *Callinectes sapidus* tissue samples in the North Sea to amplify a portion of the mitochondrial cytochrome oxidase 1 genome (see Table S3).

We first verified the specificity of this barcode performing in silico PCR using the online tool Primer-BLAST, with a maximum of 3 mismatches (https://www.ncbi.nlm.nih.gov/tools/primer-blast/index.cgi). We then tested in vivo this barcode on Blue Crab tissue and on six related species found in Canet Lagoon (*Gammarus aequicauda*, *Palaemon elegans*, *Eriphia verrucosa*, *Palaemon adspersus*, *Idotea chelipes*, *Palaemon longirostris*), which could potentially confound our qPCR-based eDNA detection assays.

The qPCR-based eDNA detection assays were designed to selectively amplify the DNA of *Callinectes sapidus* and quantify the DNA concentration. An initial calibration step was performed to associate the number of cycles required to reach the exponential phase of DNA amplification, ie cycle threshold (CT) values with a DNA concentration. The qPCR was then applied to each eDNA sample to generate CT values, which were compared to the initial calibration curve to determine the initial DNA quantity. The qPCR runs were performed on the CeMEB labex high-throughput qPCR platform (University of Montpellier), a facility physically separated from the preqPCR laboratory (GEMEX platform, CEFE). We followed Faure et al. (2023) protocol for the qPCR amplification: the final reaction volume of 25 μl contained 3 μl of template DNA (eDNA extract), 12.5 μl of TaqMan™ Environmental Master Mix 2.0 (Life Technologies, now ThermoFisher Scientific), 6.5 μl of double-distilled water (ddH2O), 1 μl of each primer (10 μM) and 1 μl of probe (2.5 μM). Samples were run on a LightCycler® 480 qPCR system (Roche, Basel, Switzerland) under the following thermal cycling conditions: 5 min at 50°C and 10 min at 95°C, followed by 55 cycles of 30 s at 95°C and 1 min at 62°C. The first campaign (2021) was performed in sites with high density of Blue crab and 12 to 24 replicates were performed. Following the first results with detections as low as only 5 of 24 replicates positives, we opted for 24 qPCR replicates for all other samples to optimize the early detection of *C. sapidus*. Finally, we sequenced two eDNA qPCR products that were positive under our detection assay to confirm that only Blue Crab had been amplified.

Citizen science data

The citizen data primarily consists of information available from professional fishermen, which was then relayed by the Regional Committee of Maritime Fisheries and Marine Farming of Occitanie (CRP-MEMO) into the Information System established by Conservatoire d'Espaces Naturels d'Occitanie (SICEN). This information system is a set of resources and devices for collecting, storing, processing and disseminating naturalist data, useful to standardize the data, whether it comes from citizen science or naturalist studies conducted on natural sites. The data is then transferred to the Occitanie SINP (Information System for the Inventory of Natural Heritage). For any new observer participating in citizen science, their entered data is first validated by DREAL and CEN Occitanie before being integrated into the database, provided that the presence of the species is confirmed with photographic evidence. The capture data reported from professional fishermen are considered as already verified observations of *Callinectes sapidus*.

Figure 1: Sampling sites for the eDNA assay targeting the Blue Crab *Callinectes sapidus*. The species is recorded as present when its DNA has been detected in at least one of the two samples collected in each site.

Results

Sensitivity of the qPCR assay

Two standard curves were produced along the qPCR assay for this study. The first standard curve (y = 3.40x +23,45, r2 = 0.98, efficiency = 97%) has been used for analyzing eDNA samples from the first and second campaign (2023). DNA extracted from Blue Crab tissues was successfully amplified for a range of dilutions with concentrations as low as 6x10-5 ng.μL^{-1}, representing the limit of quantification (LOQ). As stock of dilution had been totally used, a second standard curve was produced for the third sampling campaign samples qPCR tests (y = 5.03 x +25.60, r2 = 0.90, efficiency = 58.1%). DNA extracted from Blue Crab tissues was amplified for a range of dilutions with concentrations as low as 4x10-4 ng.μL-1 (LOQ), an order of magnitude higher than for the first standard curve representing the limit of quantification.

Specificity of the qPCR assay

Positive controls performed on DNA extracted from *Callinectes sapidus* tissues samples displayed the expected DNA concentration values, thereby validating the qPCR assays. None of the six species (tissue) present in the Canet lagoons with Blue Crab were

amplified when using the qPCR conditions selected for our test, indicating selectivity of our marker. All eDNA samples from the first campaign, collected in March and July in two lagoons with high densities of Blue Crab, and in aquariums, revealed positive detection of *C. sapidus* (Table S1). We retrieved between 20% (5 positive wells out of 24 for Canet) and 75% of positive wells (9 out of 12 wells for Grau de Canet) in natural habitat and 100% of positive wells for aquariums (24 positive wells out of 24). Sanger sequencing of the qPCR products from two eDNA samples collected in the Biodiversarium (positive control, SPY 212708, SPY 212702) and from two eDNA samples collected in Canet lagoon (SPY212706, SPY212707) (Table S1) matched at 100% with existing Blue Crab sequences already available in NCBI database and confirmed the specificity of our qPCR assay.

qPCR and effect of Blue Crab density

Callinectes sapidus has been detected in all three lagoons of the second field campaign with seven out of nine samples positive. Only two out of three samples detected *C. sapidus* in the high and intermediate concentration sites while the three samples were positive in the low concentration site. In the high concentration site, *C. sapidus* had been detected in two out of the three filters, with 0%, 16% and 94% positive

wells, with CTs when positive ranging from 36.457 +/- 0.57 to 39.05 +/- 1,033 corresponding to concentrations from 3.5 E-4 +/- 1.5 E-4 to 6.65 E-5 +/- 3.34 E-5 ng/µL.

For the site with the intermediate concentration of Blue Crab, *Callinectes sapidus* had been detected in two out of the three filters, with 0%, 8 and 25% of positive wells, with CTs when positive ranging from 39.99 +/- 0.55 to 40.72 +/- 0.165 and corresponding concentrations ranging from 1.86 E-5 +/- 2.07 E-6 to 3.18 E-5 +/- 1.05 E-5 to ng/µL. Finally, *C. sapidus* has been detected in the three filters of the low concentration site, with 4% to 16% of positive wells. CTs ranged from 25.77 +/- 0.28 to 40.35 +/- 0.89, corresponding to concentrations from 2.72 E-5 +/- 1.78 E-5 to 0.47 +/- 0.09. Overall, the lowest Blue Crab eDNA concentration has been retrieved in the intermediate concentration site 1.86 E-5 +/- 2.07 E-6 and the highest Blue Crab eDNA concentration has been retrieved in the site where *C. sapidus* was supposed to be at the lowest concentration. Overall, our qPCR quantifications seem to not be correlated with those known densities (Table S1). It is worth noting that when samples are pooled by sites, 31% of wells were positive for the high concentration site (Canet) while only 11% of the wells were positive for the medium and the low concentration site. During the first sampling campaign (2021), 56% of wells were positive in March for this site, and 44% in July. However, only 6% of wells were positive for this site during the last campaign (June 2023).

Comprehensive mapping of Blue Crab distribution in 2023

A total of 32 eDNA samples (52%) exhibited positive genetic detection in 2023 (Figure 1, Table S2). During each qPCR run, all technical blank controls yielded negative results. Across the 31 sites, *Callinectes sapidus* was detected in 24 sites. Among those 24 sites, detection was effective in the two samples for 8 sites, and in one sample for the other 16 sites. Finally, *C. sapidus* was not detected in 7 sites (0/2 samples). We detected *C. sapidus* in most kinds of habitats we sampled. We found that *Callinectes sapidus* was present in 14 out of the 17 lagoons sampled. *Callinectes sapidus* was not detected during this 2023 survey solely in Or and Gruissan lagoons. It is worth noting that in Thau and Salses-Leucate lagoons, it has not been detected in all sites covering each lagoon. In the same manner, if *C. sapidus* was detected in 2022 in both sites in Or lagoon, it has not been detected in 2023. It is worth noting that *C. sapidus* has been detected in Canet three years in a row (2021, 2022, 2023). No Blue Crab has been detected in the channel between lagoons and the sea (Grau du Prévost).

We detected *C. sapidus* in two out of the three river mouths surveyed (Grau d'Agde and Grau d'Orgon) while it has not been detected in the third one (Grau de Vendres). *Callinectes sapidus* has also been detected in Canal du Rhône à Sète which connects several lagoons where *C. sapidus* has been detected and in the coastal site that were in front of Canet lagoon (Canet Mer).

Comparison of eDNA-based detection with previous records and citizen data

We detected *Callinectes sapidus* in the 10 lagoons where it had previously been reported in the literature (Labrune et al 2019) (Figure 2). Compared to the literature, current citizen science data reported Blue Crab in two of the river mouths prospected (Grau d'Orgon and Agde) and in three other lagoons (Médard, Vic, Ingril (Sud)) where we have also detected it using eDNA probe-based qPCR (Figure 2). Finally, our eDNA probe-based qPCR assay detected for the first time *C. sapidus* in three sites; two lagoons (Salonique and Pierre Blanche) and at sea in front of Canet (Figure 2).

DISCUSSION

Our study demonstrates the effectiveness of eDNA probe-based qPCR for monitoring the invasive Blue Crab *Callinectes sapidus* in lagoons and coastal site in the Mediterranean Sea. The protocol successfully detected the species in 32 samples out 61 tested, confirming its presence in 24 out of the 31 sampled sites. Notably, our eDNA probe-based qPCR tests revealed the presence of *C. sapidus* in the 13 lagoons where the species had been previously documented through independent citizen observations, as well as in three additional sites where no prior records existed at the time of sampling. This study highlights the significant potential of eDNA-based methods for Blue Crab and more generally invasive species detection and showcases the value of integrating eDNA monitoring with existing databases derived from citizen observations for comprehensive conservation management.

Previous studies (Labrune et al. 2019) and citizen observations have already indicated the expansion of *Callinectes sapidus* in the study area (Occitanie), highlighting the necessity for early detection and the urgency of management strategies to mitigate the impact of this invasive species. Our findings not only confirm the presence of *C. sapidus* at sites previously identified by stakeholders and citizen observers but also document new occurrences, demonstrating the high efficacy of eDNA based methods for monitoring invasions in such a context. The utility of eDNA

Figure 2: Comparison of the Blue Crab *Callinectes sapidus* detection for all sites sampled and tested with qPCR-based eDNA detection assays (2022; 2 sites and 2023; 31 sites) and comparison with current citizen science data (December 2023) (Stakeholders) and previous records up to 2018 (Labrune et al. 2019). The presence of *Callinectes sapidus* (in black) is reported for eDNA results when its DNA has been detected in at least one of the two samples of a site. Numbers in brackets indicate the number of sites considered for a given location.

has been established for other invasive invertebrate species in lagoons (Klymus et al., 2017; Roux et al., 2020; Tréguier et al., 2014) and in marine ecosystems (Gargan et al., 2022; Simpson et al., 2023), where it has showed significant advantages over conventional survey methods. For example, the detection rate (per site) of an elusive freshwater turtle species using eDNA qPCR barcoding was 5.55 times higher than conventional trapping methods and the approach proved to be 18.7% more cost effective (Sternhagen et al., 2024). Given this recognized utility, a study developed specific eDNA qPCR assays to detect 60 invasive, threatened, and exploited freshwater vertebrates and invertebrates in Eastern Canada (Hernandez et al., 2020).

In our study, eDNA did not enable the quantification of *Callinectes sapidus* populations, as no correlation was found between eDNA concentration and the known density of *C. sapidus* . For fishes, such relationships have been established for some fish species (Karlsson et al., 2022; Rourke et al., 2022) but is not universally (Rourke et al., 2023). For invertebrates, the relation appear less consistent, since they have typically lower shedding rates compared to fish (Allan et al., 2021). Additional factors, such as reproductive behaviors, periods of dormancy (Crane et al., 2021) or density (Uthicke et al., 2022), could further explain the lack of correlation between eDNA concentration and crab density (Marques et al., 2024). Finally, there is a great variability between replicates, with most detection retrieved by only one of the two replicates, which might explicate the lack of correlation and advocating for the use of several replicates per site.

Combining both eDNA and citizen/stakeholder observations has proven to be an effective way to enhance avian ecological research (Padró, 2024). Here, we show that it offers a dual advantage. First it provides an opportunity to validate the eDNA as an adequate tool for monitoring *Callinectes sapidus* populations. Second, the two approaches are complementary. Citizen and stakeholders-based monitoring may be readily accessible in some sites while it may prove to be more challenging in others where eDNA provides an effective alternative. While observations provide a direct result, eDNA probe-based qPCR approach is faster, cheaper and more sensitive than eDNA metabarcoding approach. Yet, they are still not as fast as direct observations. One key benefit of eDNA is that it enables more systematic and less opportunistic monitoring compared to citizen observations. This facilitates standardized, large-scale spatial monitoring (e.g. Mathon et al., 2023) and/or seasonal or temporal assessments (Bálint et al., 2018). By enabling broad spatial and systematic sampling, eDNA

qPCR enhances our understanding of the Blue Crab's habitat and helps identify environmental factors such as salinity, temperature, turbidity, and the presence of seagrass beds or bivalves, which may influence its distribution (e.g., Marques et al., 2024). While citizen observations primarily detect *C. sapidus* from May to October, little is known about their distribution during winter. Additionally, citizen observations are unable to detect larval or juvenile stages (2-5 cm), whereas the development of eRNA is promising, with the potential to identify early life stages and even distinguish between juvenile and adult individuals (Parsley & Goldberg, 2024). eDNA can thus be a powerful tool to address these knowledge gaps in future studies. Regarding *C. sapidus* monitoring, channels connecting the sea and lagoons, serve as strategic passageways throughout the year. We recommend prioritizing monitoring in such critical sites to determine their role in the invasion process, either as vectors of dispersal or as refugia during the winter months.

Integrating environmental DNA analysis with targeted barcode-based detection offers a straightforward and reproducible approach for monitoring population fluctuations over time. This methodology has significant potential to facilitate stakeholder engagement in tracking population dynamics, providing an accessible and reliable tool for long-term monitoring and conservation efforts.

Data Availability Statement

The data that support the findings of this study are available in Supp mat. Detection of the Callinectes sapidus have also been uploaded on the Natural Héritage Inventory Information System ((SINP).

Ethics and Permit Approval Statement

No ethical nor permit were required for the research should be stated here.

Funding Statement

This study received public funding from DREAL Occitanie (Ministry of Ecology) as part of a public policy on invasive alien species and specifically of the regional action plan to combat the invasive species blue crab.

Conflicts of Interests

The authors declare that they have no conflict of interest.

Acknowledgements

The authors would like thank Stéphane Hourdez (LECOB, Banyuls-sur-Mer) for sharing tissues of invertebrates. The authors would like to address a special thank to all sites' managers for their help with this study: Canet (Rolland Mivière & Thierry Auga-Bascou ; Syndicat mixte du bassin versant du Réart (SMBVR), Office Français de la Biodiversité – Parc Natural Marin du Golfe du Lion (PN-MGL)); Salses-Leucate (Jean-Francois Laffon ; Syndicat mixte RIVAGE, Conseil Départemental des Pyrénées-Orientales) ; La Palme and Bages-Sigean (Parc naturel régional de la Narbonnaise, OFB SD 11) ; Vendres (Rémi Belleza ; Communauté de communes La Domitienne et sa Capitainerie) ; Thau (Michela Patrissi ; Centre d'étude pour la Promotion des Activités Lagunaires et Maritimes (Cépralmar), Syndicat Mixte du Bassin de Thau (SMBT)); Ingril Sud (Romain Nathan Danielou ; Sète Agglopole Méditerranée, Conservatoire du Littoral), Vic and Pierre Blanche (Romain Nathan Danielou CEN Occitanie); Méjean-Pérols et Or (Max Lecouturier ; Ville de Lattes, Syndicat Mixte du Bassin de l'étang de l'Or (SYMBO)); Grau d'Agde (Sylvain Blouet, Florent Keller, Patrick Ramy ; Aire Marine de la côte Agathoise); Grau du Prévost (Mondy Laigle ; Association de gestion de la Réserve Marine de la côte palavasienne); Ponant, Médard et Salonique (Christophe Rosso ; Mairie du Grau du Roi, Syndicat Mixte pour la protection et la gestion de la Camargue Gardoise (SMCG)); Grau d'Orgon (Delphine Marobin; Parc naturel régional de Camargue). We also thank the numerous students from CEFE and MARBEC labs wo helped in the field, including Dimitiri Medetian, Maurine Vilcot, Margot Dentan, Andrea Reyes Camargo, Simon Bettinger, Marie Orblin, and Loïc Sanchez.

Author contributions

EDT, CM, NB, LC and SM contributed to the study conception and design. All authors were involved in the Material preparation, data collection and analysis. The first draft of the manuscript was written by EDT and SM, and all authors commented on previous versions of the manuscript. All authors have read and approved the final manuscript.

References

Alidoost Salimi, P., Creed, J. C., Esch, M. M., Fenner, D., Jaafar, Z., Levesque, J. C., ..., & Sweet, M. (**2021**). A review of the diversity and impact of invasive non-native species in tropical marine ecosystems. *Marine Biodiversity Records*, 14, 11.

Allan, E. A., DiBenedetto, M. H., Lavery, A. C., Govindarajan, A. F., & Zhang, W. G. (**2021**). Modeling characterization of the vertical and temporal variability of environmental DNA in the mesopelagic ocean. *Scientific Reports*, 11, 21273.

Bailey, S. A., Brown, L., Campbell, M. L., Canning-Clode, J., Carlton, J. T., Castro, N., ..., & Zhan, A. (**2020**). Trends in the detection of aquatic non-indigenous species across global marine, estuarine and freshwater ecosystems: A 50-year perspective. *Diversity and Distributions*, 26, 1780–1797.

Bálint, M., Pfenninger, M., Grossart, H.-P., Taberlet, P., Vellend, M., Leibold, M. A., ..., & Bowler, D. (**2018**). Environmental DNA Time Series in Ecology. *Trends in Ecology & Evolution*, 33, 945–957.

Boulanger, E., Loiseau, N., Valentini, A., Arnal, V., Boissery, P., Dejean, T., ..., & Mouillot, D. (**2021**). Environmental DNA metabarcoding reveals and unpacks a biodiversity conservation paradox in Mediterranean marine reserves. *Proceedings of the Royal Society B: Biological Sciences*, 288, 20210112.

Crane, L. C., Goldstein, J. S., Thomas, D. W., Rexroth, K. S., & Watts, A. W. (**2021**). Effects of life stage on eDNA detection of the invasive European green crab (*Carcinus maenas*) in estuarine systems. *Ecological Indicators*, 124, 107412.

Dalongeville, A., Boulanger, E., Marques, V., Charbonnel, E., Hartmann, V., Santoni, M. C., ... Mouillot, D. (**2022**). Benchmarking eleven biodiversity indicators based on environmental DNA surveys: More diverse functional traits and evolutionary lineages inside marine reserves. *Journal of Applied Ecology*, 59, 2803–2813.

Dejean, T., Valentini, A., Miquel, C., Taberlet, P., Bellemain, E., & Miaud, C. (**2012**). Improved detection of an alien invasive species through environmental DNA barcoding: the example of the American bullfrog *Lithobates catesbeianus*. *Journal of Applied Ecology*, 49, 953–959.

Faure, N., Manel, S., Macé, B., Arnal, V., Guellati, N., Holon, F., ..., & Deter, J. (**2023**). An environmental DNA assay for the detection of Critically Endangered angel sharks (*Squatina* spp.). *Aquatic Conservation: Marine and Freshwater Ecosystems*, 33, 1088–1097.

Galil BS, Froglia C, Noël P (**2002**) CIESM atlas of exotic species in the Mediterranean. Volume 2: Crustaceans: decapods and stomatopods. CIESM publishers (Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée), Monaco, 192 pp, https://www.ciesm.org/atlas/Callinectessapidus.php

Gargan, L. M., Brooks, P. R., Vye, S. R., Ironside, J. E., Jenkins, S. R., Crowe, T. P., & Carlsson, J. (**2022**). The use of environmental DNA metabarcoding and quantitative PCR for molecular detection of marine invasive non-native species associated with artificial structures. *Biological Invasions*, 24, 635–648.

Giakoumi, S., Katsanevakis, S., Albano, P. G., Azzurro, E., Cardoso, A. C., Cebrian, E., ... Sghaier, Y. R. (**2019**).

Management priorities for marine invasive species. *Science of The Total Environment*, 688, 976–982.

Hernandez, C., Bougas, B., Perreault-Payette, A., Simard, A., Côté, G., & Bernatchez, L. (**2020**). 60 specific eDNA qPCR assays to detect invasive, threatened, and exploited freshwater vertebrates and invertebrates in Eastern Canada. *Environmental DNA*, 2, 373–386.

Karlsson, E., Ogonowski, M., Sundblad, G., Sundin, J., Svensson, O., Nousiainen, I., & Vasemägi, A. (**2022**). Strong positive relationships between eDNA concentrations and biomass in juvenile and adult pike (*Esox lucius*) under controlled conditions: Implications for monitoring. *Environmental DNA*, 4, 881–893.

Klymus, K. E., Ramos, D. V. R., Thompson, N. L., & Richter, C. A. (**2020a**). Development and Testing of Species-specific Quantitative PCR Assays for Environmental DNA Applications. *JoVE*, e61825.

Klymus, K. E., Merkes, C. M., Allison, M. J., Goldberg, C. S., Helbing, C. C., Hunter, M. E., . . . & Richter, C. A. (**2020b**). Reporting the limits of detection and quantification for environmental DNA assays. *Environmental DNA*, 2, 271–282.

Klymus, K. E., Marshall, N. T., & Stepien, C. A. (**2017**). Environmental DNA (eDNA) metabarcoding assays to detect invasive invertebrate species in the Great Lakes. *PLOS ONE*, 12, e0177643.

Andersen, J. H., Kallenbach, E., Thaulow, J., Hesselsøe, M., Bekkevold, D., Hansen, B. K., ..., & Knudsen, S. W. (**2018**). Development of species-specific eDNA-based test systems for monitoring of non-indigenous Decapoda in Danish marine waters. NIVA-rapport. https://niva.brage.unit.no/niva-xmlui/bitstream/handle/11250/2689446/7544-2020_high.pdf?sequence=2

Kress, W. J., García-Robledo, C., Uriarte, M., & Erickson, D. L. (**2015**). DNA barcodes for ecology, evolution, and conservation. Trends in Ecology & Evolution, 30, 25–35.

Labrune, C., Amilhat, E., Amouroux, J.-M., Jabouin, C., Gigou, A., & Noël, P. (**2019**). The arrival of the American blue crab, *Callinectes sapidus* Rathbun, 1896 (Decapoda: Brachyura: Portunidae), in the Gulf of Lions (Mediterranean Sea). *BioInvasions Records*, 8, 876–881.

Mancinelli, G., Bardelli, R., & Zenetos, A. (**2021**). A global occurrence database of the Atlantic blue crab *Callinectes sapidus*. *Scientific Data*, 8, 111.

Marques, V., Loot, G., Blanchet, S., Miaud, C., Planes, S., Peyran, C., . . . Manel, S. (**2024**). Optimizing detectability of the endangered fan mussel using eDNA and ddPCR. *Ecology and Evolution*, 14, e10807.

Mathon, L., Marques, V., Manel, S., Albouy, C., Andrello, M., Boulanger, E., . . . ,& Mouillot, D. (**2023**). The distribution of coastal fish eDNA sequences in the Anthropocene. *Global Ecology and Biogeography*, 32, 1336–1352.

McColl-Gausden, E. F., Weeks, A. R., Coleman, R., Song, S., & Tingley, R. (**2023**). Using hierarchical models to compare the sensitivity of metabarcoding and qPCR for eDNA detection. *Ecological Informatics*, 75, 102072.

Mooney, H. A., & Cleland, E. E. (**2001**). The evolutionary impact of invasive species. Proceedings of the National Academy of Sciences, 98, 5446–5451.

Padró, J. (**2024**). Integrating eDNA metabarcoding and citizen science enhances avian ecological research. *Journal of Animal Ecology*, 93, 1192–1196.

Parsley, M. B., & Goldberg, C. S. (**2024**). Environmental RNA can distinguish life stages in amphibian populations. *Molecular Ecology Resources*, 24, e13857.

Pearman, J. K., von Ammon, U., Laroche, O., Zaiko, A., Wood, S. A., Zubia, M., . . . , & Pochon, X. (**2021**). Metabarcoding as a tool to enhance marine surveillance of nonindigenous species in tropical harbors: A case study in Tahiti. *Environmental DNA*, 3, 173–189.

Rourke, M. L., Fowler, A. M., Hughes, J. M., Broadhurst, M. K., DiBattista, J. D., Fielder, S., . . . , & Furlan, E. M. (**2022**). Environmental DNA (eDNA) as a tool for assessing fish biomass: A review of approaches and future considerations for resource surveys. *Environmental DNA*, 4, 9–33.

Rourke, M. L., Walburn, J. W., Broadhurst, M. K., Fowler, A. M., Hughes, J. M., Fielder, D. S., . . . , & Furlan, E. M. (**2023**). Poor utility of environmental DNA for estimating the biomass of a threatened freshwater teleost; but clear direction for future candidate assessments. *Fisheries Research*, 258, 106545.

Roux, L.-M. D., Giblot-Ducray, D., Bott, N. J., Wiltshire, K. H., Deveney, M. R., Westfall, K. M., & Abbott, C. L. (**2020**). Analytical validation and field testing of a specific qPCR assay for environmental DNA detection of invasive European green crab (*Carcinus maenas*). *Environmental DNA*, 2, 309–320.

Sabelli, C. (**2023**). What scientists know about the blue crab invasion. Nature Italy.

Saenz-Agudelo, P., Delrieu-Trottin, E., DiBattista, J. D., Martínez-Rincon, D., Morales-González, S., Pontigo, F., . . . & Correa, C. (**2022**). Monitoring vertebrate biodiversity of a protected coastal wetland using eDNA metabarcoding. *Environmental DNA*, 4, 77–92.

Seebens, H., Schwartz, N., Schupp, P. J., & Blasius, B. (**2016**). Predicting the spread of marine species introduced by global shipping. *Proceedings of the National Academy of Sciences*, 113, 5646–5651.

Seebens, H., Bacher, S., Blackburn, T. M., Capinha, C., Dawson, W., Dullinger, S., . . . &Essl, F. (**2021**). Projecting the continental accumulation of alien species through to 2050. *Global Change Biology*, 27, 970–982.

Simpson, T. J., Wellington, C. M., Lukehurst, S. S., Huerlimann, R., Veilleux, H., Snow, M., . . . & McDonald, J. I. (**2023**). Development of a real-time PCR (qPCR) method for the identification of the invasive paddle crab *Charybdis japonica* (Crustacea, Portunidae). PeerJ, 11, e15522.

Sternhagen, E. C., Davis, M. A., Larson, E. R., Pearce, S.

E., Ecrement, S. M., Katz, A. D., & Sperry, J. H. (**2024**). Comparing cost, effort, and performance of environmental DNA sampling and trapping for detecting an elusive freshwater turtle. Environmental DNA, 6, e525.

Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (**2018**). Environmental DNA: For Biodiversity Research and Monitoring. Oxford University Press. *Journal of Applied Ecology*, 51, 871–879.

Tsuji, S., Doi, H., Hibino, Y., Shibata, N., & Watanabe, K. (**2024**). Rapid assessment of invasion front and biological impact of the invasive fish *Coreoperca herzi* using quantitative eDNA metabarcoding. *Biological Invasions*, 26, 3107–3123.

Uthicke, S., Robson, B., Doyle, J. R., Logan, M., Pratchett, M. S., & Lamare, M. (**2022**). Developing an effective marine eDNA monitoring: eDNA detection at pre-outbreak densities of corallivorous seastar (*Acanthaster cf. solaris*). *Science of The Total Environment*, 851, 158143.

Yu, Z., Ito, S., Wong, M. K.-S., Yoshizawa, S., Inoue, J., Itoh, S., . . . Hyodo, S. (**2022**). Comparison of species-specific qPCR and metabarcoding methods to detect small pelagic fish distribution from open ocean environmental DNA. *PLOS ONE*, 17, e0273670.

Zenetos, A., Albano, P. G., Garcia, E. L., Stern, N., Tsiamis, K., & Galanidi, M. (**2022**). Established non-indigenous species increased by 40% in 11 years in the Mediterranean Sea. *Mediterranean Marine Science*, 23.