An IFN-STAT1-CYBB Axis Defines Protective Plasmacytoid DC to

Neutrophil Crosstalk During *Aspergillus fumigatus* Infection

Yahui Guo[1, 2, 3], Mariano A. Aufiero[1, 2, 3, 4], Kathleen A.M. Mills[1, 2, 3, 5], Simon A. Grassmann[3],

Hyunu Kim[3], Paul Zumbo[6], Mergim Gjonbalaj[1, 2, 3], Audrey Billips[1, 2, 3], Katrina B. Mar[1, 2, 3], Yao

Yu[1, 2, 3], Doron Betel[7], Joseph C. Sun[3], Tobias M. Hohl[1, 2, 3, 4, 5, #]

[1]Infectious Disease Service, Department of Medicine, and [2]Human Oncology and Pathogenesis

Program, and [3]Immunology Program, [4]Louis V. Gerstner Jr. Graduate School of Biomedical

Sciences, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York,

NY10065, USA

[5]Immunology and Microbial Pathogenesis Graduate Program, Weill Cornell Graduate School,

New York, NY 10065, USA

[6]Applied Bioinformatics Core, Department of Physiology and Biophysics, Weill Cornell

Medicine, New York, NY 10065, USA.

[7]Applied Bioinformatics Core, Division of Hematology and Medical Oncology, Department of

Medicine, Institute for Computational Biomedicine, Weill Cornell Medicine, New York, NY

10065, USA.

[#], Corresponding author and lead contact. E-mail: hohlt@mskcc.org (T.M.H.)

Footnote

[#]Address correspondence to:

Tobias M. Hohl, MD, PhD

Memorial Sloan Kettering Cancer Center

1275 York Avenue, Box 9

New York, NY 10065

Phone: 646-888-3596

Fax: 646-422-0502

hohlt@mskcc.org

Abstract

Aspergillus fumigatus is the most common cause of invasive aspergillosis (IA), a devastating

infection in immunocompromised patients. Plasmacytoid dendritic cells (pDCs) regulate host

defense against IA by enhancing neutrophil antifungal properties in the lung. Here, we define the

pDC activation trajectory during *A. fumigatus* infection and the molecular events that underlie

the protective pDC - neutrophil crosstalk. Fungus-induced pDC activation begins after bone

marrow egress and results in pDC-dependent regulation of lung type I and type III IFN levels.

These pDC-derived products act on type I and type III IFN receptor-expressing neutrophils and

control neutrophil fungicidal activity and reactive oxygen species production via STAT1

signaling in a cell-intrinsic manner. Mechanistically, neutrophil STAT1 signaling regulates the

transcription and expression of *Cybb*, which encodes one of five NADPH oxidase subunits. Thus,

pDCs regulate neutrophil-dependent immunity against inhaled molds by controlling the local

expression of a subunit required for NADPH oxidase assembly and activity in the lung.

Keywords: *Aspergillus fumigatus*, plasmacytoid DC, neutrophil, interferons, STAT1, NADPH

oxidase, *Cybb,* lung, fungus, crosstalk

Introduction

Invasive pulmonary aspergillosis, a life-threatening mold infection, occurs when the respiratory immune system fails to eradicate ubiquitous inhaled *Aspergillus* spores (i.e., conidia) prior to their germination into tissue-invasive hyphae (1). At-risk populations include patients with acute leukemia and other bone marrow disorders, recipients of hematopoietic cell and lung transplants, individuals with immune-related or neoplastic diseases treated with prolonged corticosteroid therapy or with novel targeted biologics (e.g., ibrutinib) that blunt fungal immune surveillance pathways (2-4) and patients with severe respiratory virus infections, including influenza and COVID-19 (5, 6). As a result of these medical advances and global pandemic viruses, *A. fumigatus* has become the most common agent of mold pneumonia worldwide (6-12).

Host defense against airborne mold conidia depends on intact myeloid cell numbers and function at the respiratory mucosa. Lung-infiltrating neutrophils and monocyte-derived dendritic cells (Mo-DCs) play essential roles in killing phagocytosed conidia, with a central role for products of NADPH oxidase in this process (3, 13-15). Patients with chronic granulomatous disease (CGD) and defective NADPH oxidase function are uniquely vulnerable to invasive aspergillosis (IA, with a lifetime prevalence of 40-55% (16). Exposure to products of NADPH oxidase induces a regulated cell death process in conidia trapped within neutrophil phagosomes, resulting in sterilizing immunity at respiratory mucosal barrier (17).

An emerging theme of host defense against *Aspergillus* is the essential role of intercellular crosstalk to license neutrophil effector properties *in situ*. On one hand, recruited monocytes and Mo-DCs promote neutrophil ROS production through type I and III interferon (IFN) release, though the link between IFN production and neutrophil ROS activity remains undefined (15). On the other hand, we discovered that fungus-engaged neutrophils and Mo-DCs release CXCL9 and

CXCL10 which results in the recruitment of CXCR3[+] pDCs by promoting their influx from the

circulation into the lung. In the lung, pDCs enhance neutrophil fungicidal properties and are

essential for host defense, even in the presence of lung-infiltrating monocytes and Mo-DCs (18).

However, the underlying molecular mechanisms that regulate pDC to neutrophil crosstalk remain

undefined. Thus, both Mo-DCs and pDCs independently enhance neutrophil antifungal activity,

but it is unknown whether pDCs employ either similar or distinct mechanisms of intercellular

crosstalk with neutrophils compared to monocytes and Mo-DCs. Critical open questions relate to

the identity of pDC-derived molecules that are required for protective crosstalk with neutrophils

and to the ensuing molecular changes in neutrophils that regulate antifungal activity *in situ*.

In this study, we demonstrate that pDCs represent a key and indispensable source of type

I and type III interferons (IFNs) in the lung during respiratory *A. fumigatus* infection. In turn,

pDC-dependent and IFN-induced STAT1 signaling controls neutrophil *Cybb* expression, which

encodes an essential subunit of the NADPH oxidase complex. Thus, protective pDC to

neutrophil crosstalk primarily harnesses intercellular IFN-STAT1 signaling to calibrate the

synthesis of a critical component of the NADPH oxidase complex, thereby licensing neutrophils

to achieve optimal antifungal activity and promoting sterilizing responses against inhaled mold

spores.

Results

Aspergillus fumigatus **induces pDC activation in the lung.**

In response to respiratory fungal infection, pDCs exit the bone marrow (BM), enter the

circulation, and traffic to the lung. To gain an understanding of the pDC activation trajectory

during this infection-induced trafficking event, we conducted an unbiased transcriptome analysis

of BM pDCs isolated from uninfected mice and of BM and lung pDCs isolated from mice at 72 h

post-infection (pi), since this time point represents the peak of lung pDC influx (Fig. 1A, 1B,

Supplemental Fig. S1A). For each pDC RNA-seq sample, we pooled sorted pDCs from 10 mice

from each tissue examined and analyzed 4 biological replicates. The bulk pDC transcriptome

was remarkably similar between naïve and infected BM pDCs, with only 51 differentially

expressed genes, supporting a model in which pDC activation occurs en route to the site of

infection. In these two groups of BM pDCs, we did not observe notable differences in the

transcription of genes that encode cytokines or interferons (Fig. 1C, Supplemental Fig. S1B). In

contrast, RNA-seq analysis revealed significant changes in the pDC transcriptome when lung

pDCs were isolated from *Aspergillus*-infected mice and compared to BM pDCs from the same

animals (Fig. 1B). Overall, pDCs from infected lungs had 4475 and 4474 differentially regulated

genes compared to pDCs isolated from the BM of naïve and infected mice, respectively. (Fig.1C,

Supplemental Fig. S1C and S1D). Using Kyoto Encyclopedia of Genes and Genomes (KEGG)

pathway enrichment analysis, we found that pathways involved in the cytokine-cytokine receptor

interaction, Toll-like receptor, RIG-I, and JAK-STAT signaling were among the most

upregulated pathways in pDCs isolated from infected lungs compared to BM pDCs isolated from

the same mice (Fig. 1D and Supplemental Fig. S1E).

Lung-infiltrating pDCs upregulated genes implicated in fungal recognition, including the

C-type lectin receptors *Clec7a*, *Clec4n*, *Clec4d*, *Clec4e*, *Clec9a*, *Cd69* and *Cd209e* and

downstream signaling molecules (*Card9* and *Syk*). pDCs upregulated Toll-like (*Tlr2*, *Tlr7*, *Tlr9*,

and *Myd88)*, and growth factor (*Csf2ra*, *Csf2rb*) signaling pathways as well. Notably, we found

that expression of type I interferon (IFN), type III IFN, and the type I IFN receptor (*Ifnar1* and

Ifnar2), were markedly increased in lung-infiltrating pDCs isolated from infected mice (Fig. 1E),

consistent with prior reports that *A. fumigatus* infection induces type I and type III IFN release in

the lung (15). In addition, lung-infiltrating pDCs upregulated cytokine and chemokine (*Cxcl9*,

Cxcl10 and *Il12b)*, integrin receptor (i.e., *Icam1*, Intercellular Adhesion Molecule) and *Hif1a*

mRNAs (Fig. 1E).

To examine the impact of pDC transcriptional changes on protein expression, we infected

IFNβ reporter mice with *A. fumigatus* and found that *ifnb* promoter-driven fluorescent protein

expression increased when BM pDCs trafficked to the lung (Fig. 1F and 1H). Similarly, we

observed pDC trafficking-dependent increases in ICAM1 surface expression (Fig. 1G and 1I).

Thus, *A. fumigatus* infection substantially alters the pDC transcriptome at the portal of infection.

pDCs regulate type I and type III IFN in the lung during *A. fumigatus* infection.

To examine the contribution of pDCs to the lung inflammatory milieu, we examined type

I (*Ifna1/2/5/6*) and type III (*Ifnl2/3*) IFN induction and found that induction peaked at 48 to 72

hpi (Fig. 1E and Fig. 2A-2C), temporally coincident with the peak lung pDC influx observed in a

prior study (18). To determine whether pDCs directly control type I and type III induction, we

infected BDCA2-DTR (pDC Depleter) mice, in which diphtheria toxin (DT) administration

specifically ablates pDCs at rest and under inflammatory conditions (19), but does not ablate -

CC- chemokine receptor 2 (CCR2)-expressing monocytes and Mo-DCs that have been

implicated in type I and type III IFN release following *A. fumigatus* infection (15, 18). pDC

depleter mice exhibited a 60-80% reduction in lung *Ifna1/2/5/6* and *Ifnl2/3* mRNA levels at 72

hpi, demonstrating that pDCs directly control the induction of type I and III IFN mRNA,

consistent with the bulk RNA-seq data and with their activation trajectory in the *Aspergillus-*

infected lung (Fig. 1E).

To measure the importance of pDCs for lung cytokine levels, we next compared lung

cytokine profiles by ELISA from pDC-depleted mice and from non-transgenic, co-housed

littermate controls. pDC ablation resulted in a partial depletion (40-60%) of lung type I

(IFNα 2/4) and type III IFN levels (IFNλ 2/3). In contrast, other pro-inflammatory cytokines

implicated in pulmonary antifungal defense (GMCSF, TNF, IL1β, IL6, and IL12) were not

impacted by pDC ablation, due to their production by other cellular sources (12) in the lung (Fig.

2E and 2F). These data establish that pDCs play a critical role in regulating lung type I and type

III IFN levels during *A. fumigatus* infection.

STAT1 signaling in neutrophils controls the intracellular killing of *Aspergillus* conidia.

Type I and type III IFNs both activate STAT1 signaling in target cells and are essential

for host defense against *A. fumigatus* (15). Under baseline conditions and during *Aspergillus*

infection, lung neutrophils expressed both type I IFN receptor and type III IFN receptor mRNA,

as judged by RNAscope analysis (Supplemental Fig. S2A and S2B), and both signal through

STAT1. Targeted ablation of *Stat1* in neutrophils renders mice susceptible to invasive

aspergillosis, yet it is unknown how STAT1 signaling is coupled to myeloid cell antifungal

activity and whether STAT1 signaling is required for killing in a myeloid cell-intrinsic or -

extrinsic manner.

To distinguish these possibilities, we generated mixed BM chimeric mice (1:1 mixture of

CD45.2$^+$ *Stat1$^{-/-}$* and CD45.1$^+$ *Stat1$^{+/+}$* donor BM cells → lethally irradiated CD45.1$^+$ CD45.2$^+$

Stat1$^{+/+}$ recipients) and compared the fungicidal activity of *Stat1$^{-/-}$* and *Stat1$^{+/+}$* leukocytes within

the same lung inflammatory context (Fig. 3A). To accomplish this, we utilized fluorescent

Aspergillus reporter (FLARE) conidia that encode a red fluorescent protein (DsRed; viability

fluorophore) and are labeled with an Alexa Fluor 633 (AF633; tracer fluorophore) (20). FLARE

conidia enable us to distinguish live ($DsRed^+AF633^+$) and dead ($DsRed^-AF633^+$) conidia during

leukocyte interactions with single encounter resolution (Fig. 3B).

Following infection with FLARE conidia, we measured the frequency of neutrophil

fungal uptake (Fig. 3C, frequency of neutrophil uptake = R1 + R2) and the proportion of fungus-

engaged neutrophils that contain live conidia (Fig. 3C, proportion of fungus-engaged neutrophils

with live conidia = R1/ (R1 + R2)). $Stat1^{-/-}$ neutrophils engulfed conidia at a slightly lower rate

compared to $Stat1^{+/+}$ neutrophils in the same lung (Fig. 3D). However, the frequency of fungus-

engaged neutrophils that contained live conidia was higher in $Stat1^{-/-}$ neutrophils compared to

$Stat1^{+/+}$ neutrophils (Fig. 3E). Thus, neutrophil-engulfed conidia were more likely to be viable in

$Stat1^{-/-}$ neutrophils than in $Stat1^{+/+}$ neutrophils in the same lung tissue environment (Fig. 3E),

indicating that STAT1 signaling enhances neutrophil fungicidal activity in a cell-intrinsic

manner. Consistent with these findings and in line with published studies (15) we found that

mice that lacked $Stat1$ in radiosensitive hematopoietic cells were more susceptible to A.

$fumigatus$ challenge than mice with $Stat1$ sufficiency in the same compartment (Fig. 3F and 3G).

$Stat1$ expression in radiosensitive hematopoietic cells was dispensable for lung IFNα2/4, IFNλ

2/3, IL1β, IL6, IL12p70, IL23, and TNF levels (Supplemental Fig. S3A and 3H).

Our previous work found that lung pDCs regulate neutrophil ROS generation during

respiratory $A. fumigatus$ challenge (18). Here we measured neutrophil ROS production in $Stat1^{+/+}$

and $Stat1^{-/-}$ neutrophils and found that the ROS median fluorescence intensity (MFI) in $Stat1^{-/-}$

ROS^+ lung neutrophils were significantly reduced compared $Stat1^{+/+}$ ROS^+ lung neutrophils at 72

hpi (Fig. 3I and 3J). We enumerated myeloid cells infiltration in $Stat1^{+/+}$ and $Stat1^{-/-}$ lungs at 72

hpi, and found no difference in the recruitment of neutrophils, monocyte, and pDCs in $Stat1^{-/-}$

mice. There was a slight decrease in Mo-DC numbers in $Stat1^{-/-}$ mice, which suggests that

monocytes may exhibit a limited differentiation into Mo-DCs, resulting in reduced lung Mo-DC

numbers in $Stat1^{-/-}$ mice compared to control mice (Fig. S3B-S3E).

pDCs regulate neutrophil STAT1-dependent antifungal activity.

206   To explore whether pDCs regulate neutrophil STAT1-dependent antifungal activity, we

bred CD45.2^{+} BDCA2-DTR$^{Tg/+}$ mice with CD45.2^{+} $Stat1^{-/-}$ mice to generate CD45.2^{+} BDCA2-

DTR$^{Tg/+}$ $Stat1^{-/-}$ mice and bred BDCA2-DTR$^{Tg/+}$ $Stat1^{+/+}$ mice to the CD45.1^{+} (C57BL6.SJL)

background. BM cells from both strains were mixed in a 1:1 ratio and utilized as donor cells to

generate mixed BM chimeric mice (CD45.1^{+} BDCA2-DTR$^{Tg/+}$ $Stat1^{+/+}$ and CD45.2^{+} BDCA2-

DTR$^{Tg/+}$ $Stat1^{-/-}$→ CD45.1^{+}CD45.2^{+} recipient mice). This experimental design enabled us to

compare the fungicidal activity of $Stat1^{+/+}$ and $Stat1^{-/-}$ neutrophils in the same lung, either in the

absence or in the presence of pDCs through the administration or omission of DT to mixed BM

chimeric mice (Fig. 4A).

215   This experimental approach yielded 4 groups of neutrophils that were analyzed 72 hpi

with FLARE conidia. Group 1 (G1) neutrophils were $Stat1^{+/+}$ neutrophils isolated from pDC-

sufficient mice; G2 neutrophils were $Stat1^{+/+}$ neutrophils isolated from pDC-ablated mice; G3

neutrophils were $Stat1^{-/-}$ neutrophils isolated from pDC-sufficient mice; and G4 neutrophils were

$Stat1^{-/-}$ neutrophils isolated from pDC-ablated mice (Fig. 4A). There was no difference in

conidial uptake among these four groups of neutrophils (Fig. 4B and 4C), indicating that pDCs

do not control neutrophil conidial uptake and that neutrophil-intrinsic STAT1 signaling is

dispensable for this process.

The frequency of neutrophils that contained live conidia was markedly increased in G3

(pDC$^+$ lungs; *Stat1$^{-/-}$*) neutrophils compared to G1 (pDC$^+$ lungs; *Stat1$^{+/+}$*) neutrophils (Fig. 4B

and 4D), consistent with prior experimental results (Fig. 3E). Critically, pDC ablation increased

the frequency of *Stat1$^{+/+}$* neutrophils that contained live conidia (comparison of G2 versus G1

neutrophils; P = 0.027, Fig. 4B and 4D). This result indicates that pDC-derived products

contribute to STAT1-dependent neutrophil conidiacidal activity. In contrast, pDC ablation did

not significantly increase the frequency of *Stat1$^{-/-}$* neutrophils that contained live conidia

(comparison of G4 versus G3 neutrophils; P = 0.09, Fig. 4B and 4D).

To obtain a complementary measurement of pDC to neutrophil STAT1 crosstalk, we

measured ROS production in the 4 neutrophil groups in parallel. Consistent with the direct

measurements of fungicidal activity in Fig. 4D and with the ROS measurements in Fig. 3I and 3J,

Stat1$^{+/+}$ neutrophils displayed a higher ROS mean fluorescence intensity (MFI) than *Stat1$^{-/-}$*

neutrophils (G1 versus G3). pDC ablation markedly reduced ROS production by *Stat1$^{+/+}$*

neutrophils (G2 versus G1), in line with the reduction in neutrophil fungicidal activity (Fig. 4E).

In fact, *Stat1$^{+/+}$* neutrophils isolated from pDC-depleted lungs (G2) had a similar ROS MFI as

Stat1$^{-/-}$ neutrophils isolated from pDC-sufficient lungs (G3). ROS levels observed in *Stat1$^{-/-}$*

neutrophils isolated from pDC-depleted lungs (G3) was lower than the ROS levels observed in

Stat1$^{-/-}$ neutrophils isolated from pDC-sufficient lungs (Fig. 4E), consistent with the idea that

pDCs can further modulate neutrophil ROS production in a *Stat1*-independent fashion.

Collectively, these findings indicate that pDCs regulate STAT1-dependent neutrophil fungicidal

activity and ROS production.

STAT1-dependent guanylate-binding proteins are dispensable for the neutrophil anti-fungal activity.

245
246

To gain further insight into how STAT1 regulates cell-intrinsic neutrophil fungicidal

activity and ROS generation during respiratory *A. fumigatus* challenge, we generated mixed bone

marrow chimeric mice (1:1 mix of CD45.2$^+$ *Stat1*$^{-/-}$ and CD45.1$^+$ *Stat1*$^{+/+}$ donor bone marrow

cells → lethally irradiated CD45.1$^+$CD45.2$^+$ *Stat1*$^{+/+}$ recipient mice) and performed bulk RNA-

seq on *Stat1*$^{-/-}$ and *Stat1*$^{+/+}$ neutrophils sorted from *A. fumigatus*-infected recipient mice. There

were marked differences in the transcriptome of *Stat1*$^{-/-}$ and *Stat1*$^{+/+}$ lung neutrophils (Fig. 5A),

with 2586 genes showing differential expression in *Stat1*$^{-/-}$ neutrophils compared to *Stat1*$^{+/+}$

neutrophils. KEGG pathway enrichment analysis showed downregulation of many pathways in

Stat1$^{-/-}$ neutrophils, including the cytosolic DNA sensing pathway, proteasome, RIG-I like

receptor signaling pathway, Toll-like receptor signaling pathway, cytokine-cytokine receptor

pathway (Fig. 5B). We identified commonly downregulated genes in *Stat1*$^{-/-}$ neutrophils, many

of which were known interferon-regulated genes (ISGs; Fig. 5C).

Several GTPases guanylate-binding proteins (GBPs), including *Gbp2*, *Gbp3*, *Gbp5* and

Gbp7 were significantly downregulated in *Stat1*$^{-/-}$ neutrophils (Fig. 5C). To address the function

of these genes in neutrophil antifungal activity in otherwise immune competent mice, we utilized

Gbp$^{chr3-/-}$ mice, which lack the entire chromosome 3 cluster that contains *Gbp1*, *Gbp2*, *Gbp3*,

and *Gbp5*. We generated single chimeric mice (CD45.2$^+$ Gbp$^{chr3-/-}$ or CD45.2$^+$ Gbp$^{chr3+/+}$ →

lethally irradiated CD45.1$^+$ Gbp$^{chr3+/+}$ recipients) (Fig. 5D) and compared the mortality and

fungal burden of Gbp$^{chr3-/-}$ and Gbp$^{chr3+/+}$ chimeric mice. As expected from a prior study in

corticosteroid and cyclophosphamide-treated mice (21), there was no difference in mortality or

fungal colony forming unit (CFU) between Gbp$^{chr3-/-}$ and Gbp$^{chr3+/+}$ chimeric mice (Fig. 5E and

5F). Using mixed chimeric mice (CD45.2$^+$ Gbp$^{chr3-/-}$ and CD45.1$^+$ Gbp$^{chr3+/+}$ →

CD45.1$^+$CD45.2$^+$ Gbp$^{chr3+/+}$ recipients) (Fig. 5G) and FLARE conidia, we quantified the cell-

intrinsic antifungal activity of Gbp$^{chr3-/-}$ and Gbp$^{chr3+/+}$ leukocytes and found no difference in

neutrophil conidial uptake and killing (Fig. 5H and 5I). Moreover, Gbp$^{chr3-/-}$ and Gbp$^{chr3+/+}$

neutrophils isolated from the same lung exhibited no difference in ROS MFI (Fig. 5J).

The pDC-IFN-STAT1 axis regulates neutrophil *Cybb* expression during A. *fumigatus*

infection.

Since GBPs did not contribute to STAT1-regulated neutrophil defense against *A.*

fumigatus, we investigated other candidate genes that were downregulated in *Stat1*$^{-/-}$ neutrophils

(Fig. 5C) and focused next on *Cybb,* which encodes CYBB, the p91 subunit of the NADPH

oxidase complex (NOX2). The four genes (*Cyba*, *Ncf1*, *Ncf2*, and *Ncf4*) that encode other

subunits of the neutrophil NADPH oxidase complex (CYBA/p22 subunit, NCF1/p47 subunit,

NCF2/p67 subunit and NCF4/p40 subunit) were not downregulated in *Stat1*$^{-/-}$ neutrophils (Fig.

6A). During *A. fumigatus* infection lung *Cybb* expression peaked at 48 and 72 hpi (Fig. S4A),

temporally coincident with the observed peaks in lung pDC influx and type I and type III IFN

expression (Fig. 2A-C). The production of reactive oxygen species (ROS) by neutrophils is a

vital mechanism for effectively eradicating fungal infections. Consistent with a prior study, we

confirmed that *Cybb*-deficient mice (*gp91$^{phox-/-}$*) were highly susceptible to *A. fumigatus* infection

(Fig. S4B) (22). To investigate the hypothesis that STAT1 signaling in neutrophils regulates

Cybb expression, we isolated neutrophils from infected *Stat1*$^{-/-}$ and *Stat1*$^{+/+}$ mice and found that

Cybb mRNA level were decreased in *Stat1*$^{-/-}$ neutrophils compare to *Stat1*$^{+/+}$ neutrophils (Fig.

6B), consistent with STAT1-dependent regulation of *Cybb* transcription.

To determine chromatin accessibility at the *Cybb* locus, we performed assay for

transposase-accessible chromatin (ATAC) sequencing of sorted *Stat1*$^{-/-}$ and *Stat1*$^{+/+}$ lung

neutrophils from *A. fumigatus*-infected mice. Within the *Cybb* locus, we found three regions

(blue squares) that were less accessible in *Stat1*$^{-/-}$ neutrophils than in *Stat1*$^{+/+}$ neutrophils (Fig.

6C), consistent with a modest STAT1-dependent regulation of *Cybb* expression (Fig. 6A and 6B).

As a positive control, within Gbp2 locus, we found one region (red square) that was less

accessible in *Stat1*$^{-/-}$ neutrophils compared to *Stat1*$^{+/+}$ neutrophils (Fig. 6C), consistent with

STAT1-dependent regulation of *Gbp* expression (Fig. 5C).

To probe STAT1 binding to the Cybb gene locus, we performed <u>C</u>leavage <u>U</u>nder <u>T</u>argets

<u>&</u> <u>R</u>elease <u>U</u>sing <u>N</u>uclease (CUT&RUN) experiments of sorted *Stat1*$^{-/-}$ and *Stat1*$^{+/+}$ lung

neutrophils from *A. fumigatus*-infected mice. We did not observe direct STAT1 binding to *Cybb*

locus and promoter region (Fig. 6D). In CUT&RUN experiments, STAT1 bound to the *Gbp2*

locus and promoter region (Fig. 6D). Collectively, these results suggest that STAT1 regulates

Cybb transcription via an indirect mechanism.

Because STAT1 did not appear to regulate *Cybb* transcription through direct effects on

chromosomal accessibility or binding to the *Cybb* locus, we next explored whether lung

neutrophil CYBB protein levels were regulated by STAT1 signaling. *Stat1*$^{-/-}$ and *Stat1*$^{+/+}$

neutrophils were isolated from *A. fumigatus* infected mice lungs at 72 hpi and analyzed for

CYBB expression by Western blotting. CYBB protein levels, but not other subunits (p22/CYBA

and p40/NCF4) of NADPH oxidase protein levels were lower in *Stat1*$^{-/-}$ neutrophils compared to

Stat1$^{+/+}$ neutrophils (Fig. 6E – 6H), linking neutrophil STAT1 activation to CYBB expression,

and to the oxidative burst.

Discussion

Our data introduce a model of protective pDC to neutrophil crosstalk in which pDCs

undergo a defined activation trajectory in transit to the *Aspergillus*-infected lung. pDC activation

provides a critical source of type I and type III IFN at the portal of infection, and licenses

neutrophil antifungal properties in the lung in a direct manner. The latter step occurs through

neutrophil-intrinsic STAT1-dependent control of CYBB protein levels. Local pDC-dependent

control of NADPH oxidase assembly regulates the strength of neutrophil oxidative burst, as

judged by reactive oxygen species production, boosts neutrophil intracellular conidial killing,

and confers sterilizing immunity against inhaled spores.

pDCs originate in the bone marrow (BM), travel through the bloodstream, and migrate to

lymphoid and nonlymphoid tissues during both normal and inflammatory states (18, 23-25). In a

recent study we demonstrated that *Aspergillus*-infected Mo-DCs and neutrophils release CXCR3

ligands (i.e., CXCL9 and CXCL10) into the inflamed lung, coupling fungal recognition and

fungus-induced inflammation to CXCR3 signaling-dependent pDC influx from the circulation

into the lung. In this study, *A. fumigatus* infection induces a pDC activation trajectory that

substantially alters the lung rather than the BM pDC transcriptome, implying that pDC

trafficking to peripheral tissue precedes activation occurs at the portal of infection. While not a

focus of this work, the precise mechanism of pDC activation in the fungus-infected lung remains

an open question, in part because conditional gene deletion strategies do not exist for pDCs,

precluding facile comparison of gene-deficient and gene-sufficient pDCs in infected tissues in

mice with no other genetic perturbations.

Our data indicate that lung-infiltrating pDCs upregulate genes implicated in fungal

recognition, including the C-type lectin receptors (CLRs) and downstream signaling molecules.

pDCs express the C-type lectin receptors Dectin1, Dectin2, and Dectin3 and can secrete

IFNα and TNF when co-incubated with *Aspergillus* hyphae via Dectin2 signaling (26-28),

though the interaction of pDCs with conidia, the infectious propagules, were not examined.

pDCs do not internalize *Aspergillus* conidia readily in the test tube or in the infected lung (4, 26),

supporting the notion that activation occurs via fungal cell contact or via the presence of

activating cytokines or other inflammatory mediators. Support for the latter scenario comes from

an experiment in which curdlan (i.e., a particulate Dectin1 agonist)-induced upregulation of pDC

CD40 and CD83 expression could be increased by simultaneous co-incubation with an acellular

curdlan-stimulated peripheral blood mononuclear cell supernatant (26). Beyond *Aspergillus*,

pDC-enriched human cell fractions can release TNF in a Dectin2 and Dectin3 signaling-

dependent fashion in response to *Paracoccidioides brasiliensis* co-incubation (29). Beyond CLR

signaling, we found that *Aspergillus* infection triggered upregulation of Toll-like receptors in

lung-infiltrating pDCs. Prior work demonstrated that *Aspergillus*-derived unmethylated CpG

sequences can activate TLR9 signaling *in vitro* (30, 31), and this may represent an additional

mechanism by which pDCs become activated *during Aspergillus* infection.

During Dengue, Zika, and Hepatitis C viral infections, physical contact with virally

infected cells stimulates pDC-mediated antiviral responses (32, 33). $\alpha_L\beta_2$ integrin (lymphocyte

function-associated antigen-1; LFA1) expressed by the pDC can bind to ICAM1 on infected cells

to promote a sustained interaction, termed an interferonogenic synapse, during which viral RNA

is transferred to pDCs, leading to IFN production via the nucleic acid sensor TLR7. This process

activates type I IFN-dependent antiviral programs in infected tissues. pDC commitment to type I

IFN production is further regulated by antecedent cell-intrinsic TNF receptor and leukemia

inhibitory factor signaling during murine cytomegalovirus infection, underscoring the

contribution of local cytokine signaling to pDC activation by contact-dependent and -

independent mechanisms (34). The lung pDC transcriptomic data detected increased expressed

ICAM1 in the lung-infiltrating pDCs isolated from infected mice. This observation raises the

possibility that ICAM1 expression may facilitate lung pDC contact with β_2-integrin-expressing

myeloid cells to facilitate reciprocal interactions. In sum, the upregulation of multiple receptors

from various classes suggests that pDCs likely employ a blend of receptors to identify and react

to fungal pathogens, leading to full pDC activation in the fungus-infected lung. Notably, we were

unable to isolate sufficient pDCs to compare the bulk lung pDC transcriptome in naïve mice with

the lung pDC transcriptome in *Aspergillus*-infected mice.

Recent studies have advanced the concept that circulating neutrophils exhibit

transcriptional heterogeneity in the steady state and during microbial infection (35). In addition,

neutrophils can acquire transcription-dependent non-canonical functions upon entry into

peripheral tissues, exemplified by a regulatory role in angiogenesis in the lung (36). Single-cell

analysis has revealed the presence of three separate circulating murine and human populations

that differ with respect to interferon-stimulated gene (ISG) and CXCR4 expression. Interestingly,

all three populations expressed similar levels of *Cybb* mRNA and exhibited similar NADPH

oxidase scores by gene ontogeny analysis during homeostasis and systemic bacterial (i.e.,

Escherichia coli) infection (35). During tuberculosis, malaria, and hematopoietic cell

transplantation, circulating human neutrophils exhibit an IFN transcriptional signature (37-39),

which in the case of malaria, is reduced by receipt of anti-malarial therapy. The finding that

pDC-derived type I and type III IFN licenses neutrophil antifungal activity through STAT1-

dependent control of *Cybb* transcription and CYBB protein levels raises the important question

whether the ISG[hi] neutrophil subset is enriched in the lung and particularly effective at killing

Aspergillus conidia compared to the other two ISG^{lo} circulating subsets. The ability to track and

quantify fungal uptake killing with FLARE conidia will facilitate future studies to link lung-

infiltrating neutrophil subsets, defined by distinct transcriptional profiles, to the quality of

effector functions at single-encounter resolution. These studies support idea that dynamic

transcriptional plasticity represents a cardinal feature of the circulating neutrophil response to

microbial infection and other external stressors. It remains unclear whether the observed

transcriptional plasticity represents dynamic changes in the production of transcriptionally

heterogeneous circulating neutrophil subsets or represents an interconversion between circulating

subsets.

The effects of type III IFN on neutrophils are context-dependent. In a DSS model of

sterile colitis, IFNλ has been linked to neutrophil ROS suppression and a reduction in neutrophil

degranulation (40). In human neutrophils, recombinant IFNλ can inhibit platelet-induced

NETosis (41). IFNλ has various roles in bacterial infections (42, 43), with a previous study

demonstrated that *Bordetella pertussis* infection induces IFNλ and IFNλ receptor (IFNLR1)

expression as well as inflammation in the lung (42). IFNλ signaling in neutrophils can suppress

B. pertussis killing and neutrophil production of ROS, MMP9, NETs, MPO, and IFNγ (44, 45).

In contrast, conditional deletion of the IFNλ receptor in neutrophils is linked to a reduction in

neutrophil ROS during pulmonary *A. fumigatus* infection (15). During HSV corneal infection,

application of recombinant IFNλ suppressed neutrophil recruitment but did not impact virucidal

activity or ROS production (46). These results indicate that IFNλ may have different biological

effects based on cellular targets and responsiveness at sites of inflammation and on the type of

inflammatory stimulus. In this study, the STAT1-dependent action of pDC-dependent IFNs (both

type I and III) on neutrophils is critical for their local licensing of their cell-intrinsic cytotoxic

activity against *Aspergillus* conidia. The individual contribution of type I versus type III IFN-

dependent activation on STAT1-dependent CYBB expression remains unclear. An open question

remains how STAT1 signaling precisely regulates *Cybb* transcription and CYBB translation,

since we could not detect clear evidence of STAT1 binding to the *Cybb* promoter or clear

STAT1-dependent changes in chromatin accessibility.

Overall, these findings provide important insights into the pDC-STAT1-CYBB axis as a

key regulator of NADPH oxidase expression and highlight the critical role of this pathway in

promoting antifungal immunity in neutrophils. The discovery that pDCs can regulate ROS

induction by neutrophils by controlling the STAT1-dependent expression of a single NADPH

oxidase subunit adds to our understanding of the complex and protective interplay between

innate immune cells during fungal infection.

ACKNOWLEDGMENTS

We thank Dr. Thirumala-Devi Kanneganti (St. Jude Children's Research Hospital)

providing the *GBP$^{chr3-/-}$* bone marrows. We thank the MSKCC IGO core for performing RNA

sequencing, and Chen Liao (Geisel School of Medicine at Dartmouth), Paul Zumbo (Weill

Cornell Medicine) for assistance with the RNA sequencing data analysis. We thank Robert

Cramer (Geisel School of Medicine at Dartmouth), Iliyan Iliev (Weill Cornell Medicine), and

members of the Hohl laboratory for numerous conversations and insights into this study. In this

study, T. M. H. was supported by NIH grants P30 CA 008748 (to MSKCC), R37 AI093808, and

R01 AI139632; Y. G. was supported by the Ludwig Center Post-Doctoral Award, MSKCC;

M.A.A. was supported by NIH F31 AI161996; K.A.M.M. was supported by NIH F31 AI167511;

K. B. M. was supported by Experimental Immuno-oncology Scholars' Program, MSKCC and

NIH 5T32CA009149; J. C. S. was supported by the Ludwig Center for Cancer Immunotherapy,

the American Cancer Society, the Burroughs Wellcome Fund, and the NIH grants R01 AI100874,

R01 AI130043, R01 AI155558, and P30CA008748. S. G. was supported by CRI / Donald J.

Gogel Postdoctoral Fellowship CRI Award CRI3934 and NIH K99AI180360. The funders had

no role in study design, data collection and analysis, decision to publish or preparation of

manuscript. We acknowledge the use of the Integrated Genomics Operation Core, funded by the

NCI Cancer Center Support Grant (CCSG, P30 CA08748), Cycle for Survival, and the Marie-

Josée and Henry R. Kravis Center for Molecular Oncology.

AUTHOR CONTRIBUTIONS

Conceptualization, T.M.H., and Y.G.; Methodology, T.M.H., and Y.G.; Investigation, Y.G.,

M.A.A., K.A.M.M, S. A. G., H. K., P. Z., M.G., A.B., K.M., Y. Y.; Writing – Original Draft,

Y.G. and T.M.H.; Writing – Review & Editing, Y.G. and T.M.H.; Funding Acquisition, T.M.H.

and Y.G.; Resources, J. S., and D. B.

DECLARATION OF INTERESTS

The authors declare no competing interests.

MATERIALS AND METHODS

Chemicals and reagents

Unless otherwise noted, chemicals were purchased from Sigma-Aldrich or Fisher

Scientific, cell culture reagents from Thermo Fisher Scientific, and microbiological culture

media from BD Biosciences. Antibodies for flow cytometry were acquired from BD Biosciences,

Thermo Fisher Scientific and Tonbo.

Mice

Our study examined male and female animals, and similar findings are reported for both

sexes. C57BL/6J (JAX: 00664), BDCA2-DTR (JAX: 014176) mice were from Jackson

Laboratories. $GBP^{chr3-/-}$ bone marrows were provided by Dr. Thirumala-Devi Kanneganti.

C57BL/6.SJL (Stock: 4007) were purchased from Taconic. C57BL/6 and C57BL/6.SJL mice

were crossed to generate CD45.1$^+$CD45.2$^+$ recipient mice for mixed BM chimeras. CD45.2$^+$

BDCA2-DTR$^{Tg/+}$ were backcrossed to C57BL/6.SJL mice to obtain CD45.1$^+$ BDCA2- DTR$^{Tg/+}$

mice. BDCA2- DTR$^{Tg/+}$ were crossed with CD45. 2$^+$ $Stat1^{-/-}$ mice to obtain CD45. 2$^+$ BDCA2-

DTR$^{Tg/+}$ $Stat1^{-/-}$ mice. For experiments in which the breeding strategy did not yield littermate

controls, gene-knockout mice were co-housed with C57BL/6 mice for 14 days prior to infection,

whenever possible.

Generation of Bone Marrow Chimeric Mice

For single BM chimeras, CD45.1$^+$ C57BL/6.SJL recipients were lethally irradiated

(900cGy), reconstituted with either 2-5 $\times$ 10^6 CD45.2$^+$ $GBP^{chr3-/-}$, CD45.2$^+$ C57BL/6J or CD45.2$^+$

$Stat1^{-/-}$ BM cells. For mixed BM chimeras, CD45.1$^+$CD45.2$^+$ recipients were irradiated and

reconstituted with a 1:1 mixture of CD45.1$^+$ C57BL/6.SJL and CD45.2$^+$ $Stat1^{-/-}$ or CD45.2$^+$

$GBP^{chr3-/-}$ BM cells, CD45.1$^+$ BDCA2- DTR$^{Tg/+}$ and CD45.2$^+$ BDCA2- DTR$^{Tg/+}$ $Stat1^{-/-}$. After

BM transplantation, recipient mice received 400 µg/ml enrofloxacin in the drinking water for 21

days to prevent bacterial infections and rested for 6-8 weeks prior to experimental use.

Aspergillus fumigatus **Infection Model**

A. fumigatus Af293, Af293-dsRed (20), and CEA10 (47) strains were cultured on glucose

minimal medium slants at 37°C for 4-7 days prior to harvesting conidia for experimental use. To

generate FLARE conidia, briefly, 7×10^8 Af293-dsRed conidia were rotated in 10 µg/ml Biotin

XX, SSE in 1 ml of 50 mM carbonate buffer (pH 8.3) for 2 hr at 4 °C, incubated with 20 µg/ml

Alexa Fluor 633 succinimidyl ester at 37 °C for 1 h, resuspended in PBS and 0.025% Tween 20

for use within 24 hrs. (20, 48). To generate morphologically uniform heat-killed swollen conidia,

5×10^6/ml conidia were incubated at 37° C for 14 hours in RPMI-1640 and 0.5 µg/ml

voriconazole and heat killed at 100 °C for 30 minutes (49). To infect mice with 30-60 million

live or heat-killed *A. fumigatus* cells, conidia were resuspended in PBS, 0.025% Tween-20 at a

concentration of $0.6\text{-}1.2 \times 10^9$ cells and 50 µl of cell suspension was administered via the

intranasal route to mice anesthetized by isoflurane inhalation.

Analysis of *in vivo* and *in vitro* conidial uptake and killing

To analyze of conidia uptake and killing, FLARE conidia were used to infect mice. In

data analyses for a given leukocyte subset, conidial uptake refers to the frequency of fungus-

engaged leukocytes, i.e., the sum of dsRed$^+$ AF633$^+$ and dsRed$^-$ AF633$^+$ leukocytes. Conidial

viability within a specific leukocyte subset refers to the frequency of leukocytes that contain live

conidia (dsRed$^+$ AF633$^+$) divided by the frequency of all fungus-engaged leukocytes (dsRed$^+$

AF633$^+$ and dsRed$^-$ AF633$^+$).

In vivo **Cell Depletion**

To ablate specific cells, BDCA2- DTR$^{Tg/+}$, BDCA2- DTR$^{Tg/+}$ *Stat1$^{-/-}$*, and non-transgenic

littermate controls were injected i.p. with 10 ng/g body weight DT on Day -1, Day 0, and Day +2

pi (19), unless noted otherwise.

Analysis of Infected Mice

To prepare single cell lung suspensions for flow cytometry, we followed the method

outlined in (Hohl et al., 2009) with minor modifications. After perfusing murine lungs, they were

placed in a gentle MACS™ C tube and mechanically homogenized in 5 ml RPMI-1640, 10%

FBS, and 0.1 mg/ml DNAse using a gentle MACS™ Octo Dissociator (Miltenyi Biotech)

without the use of collagenase. The lung cell suspensions were then lysed of RBCs, enumerated,

and stained with fluorophore-conjugated antibodies prior to flow cytometric analysis. We used

either a Cytoflex or a BD Aria for flow cytometric sorting and performed flow plot analysis

using FlowJo v.9.6.6 software.

Neutrophils were identified as CD45$^+$ CD11b$^+$ Ly6C^{lo} Ly6G$^+$ cells, monocytes as

CD45$^+$ CD11b$^+$ CD11c$^-$ Ly6G$^-$ Ly6C^{hi} cells, Mo-DCs as CD45$^+$ CD11b$^+$ CD11c$^+$ Ly6G$^-$

Ly6C^{hi} MHC class II$^+$ cells, and pDCs as CD45$^+$ CD11c^{int} SiglecF$^-$ CD19$^-$ NK1.1$^-$ CD11b$^-$

B220$^+$ SiglecH$^+$ cells.

To assess the lung fungal burden, perfused murine lungs were homogenized using a

PowerGen 125 homogenizer (Fisher) in 2 mL of PBS containing 0.025% Tween-20 and plated

onto Sabouraud dextrose agar. For analysis of cytokine levels by ELISA, whole lungs were

weighed and mechanically homogenized in 2 mL of PBS containing protease inhibitors. To

analyze cytokine levels by qRT-PCR, we extracted total RNA from cells using TRIzol

(Invitrogen). One to two micrograms of total RNA were reverse-transcribed using the High-

Capacity cDNA Reverse Transcription Kit (Applied Biosystems). We used TaqMan Fast

Universal Master Mix (2×) and TaqMan probes (Applied Biosystems) for each gene and

normalized to glyceraldehyde-3-phosphate dehydrogenase. Gene expression was calculated

using the Ct method relative to the naïve sample.

Intracellular ROS levels were measured in cells using CM-H2DCFDA [5-(and 6-)

chloromethyl-2,7-dichlorodihydrofluorescein diacetate, acetyl ester] as described in (50). Briefly,

single cell lung suspensions were incubated with 1µM CM-H2DCFDA in Hanks' balanced salt

solution at 37º C for 45 min according to manufacturer's instruction and analyzed by flow

cytometry.

Immunoblotting assay

Cells were lysed in lysis buffer (150 mM NaCl, 50 mM HEPES (pH 7.4), 1 mM EDTA, 1%

Nonidet P-40, protease inhibitors). Total cell lysates were subjected to SDS-PAGE and then

blotted using indicated antibodies, Cybb polyclonal antibody (Invitrogen, cat. no. PA5-76034),

p22phox mAb (CST, cat. no. 37570), p40phox mAb (cat. no. AB76158), GAPDH mAb (CST,

cat. no. 5174s).

RNA sequencing

RNA extraction. Phase separation in cells lysed in 1 mL TRIzol Reagent (ThermoFisher cat. no.

15596018) was induced with 200 µL chloroform. RNA was extracted from 350 µL of the

aqueous phase using the miRNeasy Micro or Mini Kit (Qiagen cat. no. 217084/217004) on the

QIAcube Connect (Qiagen) according to the manufacturer's protocol. Samples were eluted in

13-15 μL RNase-free water.

Transcriptome sequencing. After RiboGreen quantification and quality control by Agilent

BioAnalyzer, 1.9-2.0 ng total RNA with RNA integrity numbers ranging from 7.8 to 10

underwent amplification using the SMART-Seq v4 Ultra Low Input RNA Kit (Clonetech cat. no.

63488), with 12 cycles of amplification. Subsequently, 7.4-10 ng of amplified cDNA was used to

prepare libraries with the KAPA Hyper Prep Kit (Kapa Biosystems cat. no. KK8504) using 8

cycles of PCR. Samples were barcoded and run on a HiSeq 4000 or NovaSeq 6000 in a PE50

(HiSeq) or PE100 (NovaSeq) run, using the HiSeq 3000/4000 SBS Kit or NovaSeq 6000 S4

Reagent Kit (200 Cycles) (Illumina). An average of 40 million paired reads were generated per

sample and the percent of mRNA bases per sample averaged 84%.

RNA-Sequencing data analysis. Raw reads were quality checked with FastQC v0.11.7

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/), and adapters were trimmed using

Trim Galore v0.6.7 (http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/). Reads

were aligned to the mouse reference genome (GRCm38.p6) using STAR v2.6.0c (51) with

default parameters. Gene abundances were calculated with featureCounts v1.6.2 (52) using

composite gene models from Gencode release vM17 . Principle component analysis was

performed using the plotPCA function from DESeq2 v1.32.0 (53). Differentially expressed genes

were determined with DESeq2 v1.32.0 with a two-factor model incorporating batch as a

covariate, with significance determined by Wald tests ($q < 0.05$). Gene set enrichment analysis

was performed using fgsea v1.18.0 (54) with gene sets from the Broad Institute's MSigDB (55,

56) collections; genes were ranked by the DESeq2 Wald statistic. Only pathways with an

adjusted P value < 0.05 were considered enriched. Expression heatmaps were generated using

variance-stabilized data, with the values centered and scaled by row. Code has been deposited to

GitHub (https://github.com/abcwcm/Guo2024).

RNAscope microscopy

Formaldehyde-fixed, paraffin embedded (FFPE) surgical tissue Sects. FFPE tissue

sections (5 µm) were processed exactly as described in the manufacturer's instructions (ACD

Bio, cat. no. 323100). Probes targeting the following genes were used: Ly6G (ACD Bio, cat. no.

455701-C3), IFNλR (ACD Bio, cat. no. 512981-C1), IFNαR1(ACD Bio, cat. no. 512971-C2).

Slides were mounted with Prolong Diamond mounting media (ThermoFisher Scientific, cat. no.

P36965). Slides were scanned using a Pannoramic Digital Slide 517 Scanner (3DHISTECH,

Budapest, Hungary) using a 20×/0.8NA objective.

Assay for transposase-accessible chromatin (ATAC) sequencing and epigenome analyses

Freshly harvested WT and KO mouse neutrophils were directly sent to MSKCC's

Epigenetics Research Innovation Lab. ATAC was performed as previously described (Corces et

al. Nature Methods 2017) using 50,000 cells per replicate and the Tagment DNA TDE1 Enzyme

(Illumina, 20034198). Sequencing libraries were prepared using the ThruPLEX DNA-Seq Kit

(Takarabio, R400676) and sent to the MSKCC Integrated Genomics Operation core facility for

sequencing on a NovaSeq 6000. Raw sequencing reads were trimmed and filtered for quality

(Q>15) and adapter content using version 0.4.5 of TrimGalore

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore) and running version 1.15 of

cutadapt and version 0.11.5 of FastQC. Version 2.3.4.1 of bowtie2 (http://bowtie-

bio.sourceforge.net/bowtie2/index.shtml) was employed to align reads to mouse assembly mm10

and alignments were deduplicated using Mark Duplicates in Picard Tools v2.16.0. Enriched

regions were discovered using MACS2 (https://github.com/taoliu/MACS) with a p-value setting

of 0.001, filtered for blacklisted regions

(http://mitra.stanford.edu/kundaje/akundaje/release/blacklists/ mm10-

mouse/mm10.blacklist.bed.gz), and a peak atlas was created using +/- 250 bp around peak

summits. The BEDTools suite (http://bedtools.readthedocs.io) was used to create normalized

bigwig files. Version 1.6.1 of featureCounts (http://subread.sourceforge.net) was used to build a

raw counts matrix and DESeq2 was employed to calculate differential enrichment for all

pairwise contrasts. Peak-gene associations were created by assigning all intragenic peaks to that

gene, while intergenic peaks were assigned using linear genomic distance to transcription start

sites (TSS).

Histone and Transcription Factor CUT&RUN

For CUT&RUN, 400,000 sorted neutrophils were used for H3K4me3 and STAT1

analysis. Anti-H3K4me3 (Epicypher, cat. no. 13-0028) or polyclonal anti-STAT1 antibodies

(Proteintech, cat. no. 10144-2-AP) were employed for each target. Sorted cells were washed with

PBS and resuspended in Antibody Buffer (1 × eBioscience Perm/Wash Buffer, 1× Roche

cOmplete EDTA-free Protease Inhibitor, 0.5 µM Spermidine, and 2 µM EDTA in H2O). They

were incubated overnight at 4°C with control IgG, H3K4me3, or STAT1 antibodies diluted

1:100 in Antibody Buffer in a 96-well V-bottom plate. Following antibody incubation, cells were

washed twice with Buffer 1 (1× eBioscience Perm/Wash Buffer, 1× Roche cOmplete EDTA-free

Protease Inhibitor, 0.5 µM Spermidine in H2O) and resuspended in 50 µL of Buffer 1 plus 1×

pA/G-MNase (Cell Signaling, cat. no. 57813). This mixture was incubated on ice for 1 hour, then

washed twice with Buffer 2 (0.05% w/v Saponin, 1× Roche cOmplete EDTA-free Protease

Inhibitor, 0.5 µM Spermidine in 1X PBS) three times. Cells were resuspended in Calcium Buffer

(Buffer 2 plus 2 µM CaCl2) and incubated on ice for 30 minutes to activate the pA/G-MNase

reaction. An equal volume of 2× STOP Buffer (Buffer 2 plus 20 µM EDTA plus 4 µM EGTA)

and 1 pg of *Saccharomyces cerevisiae* spike-in DNA (Cell Signaling, cat. no. 29987) were added.

Samples were incubated at 37 °C for 15 minutes, followed by DNA isolation and purification

using the Qiagen MinElute Kit per the manufacturer's instructions.

Immunoprecipitated DNA was quantified by PicoGreen, and the size was evaluated using

an Agilent BioAnalyzer: fragments between 100 and 600 bp were size selected using aMPure XP

beads (Beckman Coulter, cat. no. A63882). KAPA HTP Library Preparation Kit (Kapa

Biosystems, cat. no. KK8234) was used to prepare Illumina sequencing libraries according to the

manufacturer's instructions with 0.001–0.5 ng input DNA and 14 cycles of PCR. Barcoded

libraries were run on the NovaSeq 6000 in a PE100 run using S4 kit version 1.5 with XP mode to

generate approximately 23 million paired reads per sample.

CUT&RUN Data Processing

CUT&RUN datasets were processed by trimming paired reads for adaptors and low-

quality sequences using Trimmomatic (v0.39) and aligning to the mm10 reference genome with

Bowtie 2 (v2.4.1). Peaks were identified using MACS2 (v2.2.7.1) with input samples as control,

employing narrow peak parameters and cutoff - analysis - ple-5 - keep - dup all -B - SPMR.

Irreproducible discovery rate (IDR) calculations were performed using ENCODE project scripts

(IDR v2.0.4.2). Reproducible peaks with an IDR value of 0.05 or less in each condition were

retained, aggregated, and merged to create the final atlas, which was annotated with the UCSC

Known Gene model. Reads were mapped to this atlas and counted using the summarize Overlaps

function from the Genomic Alignment package (v1.34.1).

Visualization. Genomic tracks were visualized using GViz (v1.42.1) or IGV (v2.9.4).

Quantitation and Statistical Analysis

All data presented are representative of at least two independent experiments, as

indicated. Unless stated otherwise, all results are expressed as mean ($\pm$ SEM). We used the

Mann-Whitney test for comparisons of two groups, and the Kruskal-Wallis test for multi-group

comparisons, unless noted otherwise. Survival data were analyzed using the long-rank test. All

statistical analyses were performed using GraphPad Prism software (v9.2.0).

Study approval

All mouse strains were bred and housed in the MSKCC or St. Jude Children's Research

Hospital Animal Resource Center under specific pathogen-free conditions. All animal

experiments were conducted with sex- and age-matched mice and performed with MSKCC

Institutional Animal Care and Use Committee approval. Animal studies complied with all

applicable provisions established by the Animal Welfare Act and the Public Health Services

Policy on the Humane Care and Use of Laboratory Animals.

Resource availability and Lead Contact

Further information and requests for resources or reagents should be directed to the Lead Contact,

Tobias M. Hohl (hohlt@mskcc.org).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

Values for all data points in graphs are reported in the Supporting Data Values file. RNA

sequencing data, ATAC sequencing data and CUT&RUN sequencing data were uploaded into

NCBI database. Raw RNA sequencing datasets generated in this study are available at

GSE280164. Raw ATAC sequencing data and CUT&RUN sequencing data generated in this

study are available at GSE280229.

Key Resources Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
APC Mouse anti-Mouse CD45.1 (clone A20)	BD Bioscience	Cat#558701; RRID: AB_1645214
PerCP-Cy™5.5 Mouse anti-Mouse CD45.2 (clone 104)	BD Bioscience	Cat#552950; RRID: AB_394528
APC-Cy™7 Rat anti-Mouse CD45 (clone 30-F11)	BD Bioscience	Cat#557659; RRID: AB_396774
PerCP-Cy™5.5 Rat anti-CD11b (clone M1/70)	BD Bioscience	Cat#550993; RRID: AB_394002
PE-Cy™7 Hamster anti-Mouse CD11c (clone HL3)	BD Bioscience	Cat#558079; RRID: AB_647251
BV 605™ anti-mouse CD11c Antibody (clone N418)	BioLegend	Cat#117334; RRID: AB_117334
PE-Cy™7 Rat Anti-Mouse Ly6C (clone AL-21)	BD Bioscience	Cat#560593; RRID: AB_1727557
FITC Rat anti-Mouse Ly6G (clone 1A8)	BD Bioscience	Cat#551460; RRID: AB_394207
FITC Rat anti-Mouse CD45R/B220 (clone RA3-6B2)	BD Bioscience	Cat#553087; RRID: AB_394617
PE anti-mouse CD183 Antibody (clone CXCR3-173)	BioLegend	Cat#126505; RRID: AB_1027656
BV421 Rat Anti-Mouse SiglecF (clone E50-2440)	BD Bioscience	Cat#562681; RRID: AB_2722581

BV605 Rat Anti-Mouse SiglecH (clone 440c)	BD Bioscience	Cat#747673; RRID: AB_2744234
Purified Rat Anti-Mouse CD16/CD32 (clone 2.4G2)	BD Bioscience	Cat#553142; RRID: AB_394657
eFluor 450 NK1.1 Monoclonal Antibody (clone PK136)	eBioscience	Cat#48-5941-80; RRID: AB_2043878
eFluor 450 CD19 Monoclonal Antibody (clone eBio1D3)	eBioscience	Cat#48-0193-82; RRID: AB_2734905
Alexa Fluor 700, I-A/I-E Monoclonal Antibody (clone M5/114.15.2)	eBioscience	Cat#56-53-21; RRID: AB_494009
FITC Mouse IgG1 kappa Isotype Control	eBioscience	Cat#11-4714-42; RRID: AB_10596964
PE Rat IgG2a kappa Isotype Control	eBioscience	Cat#12-4321-42; RRID: AB_1518773
PerCP-Cyanine5.5 IgG Isotype Control	eBioscience	Cat#45-4888-80; RRID: AB_906260
APC Rat IgG2b kappa Isotype Control	eBioscience	Cat#17-4031-82; RRID: AB_470176
APC-eFluor 780 Rat IgG2b kappa Isotype Control	eBioscience	Cat#47-4031-80; RRID: AB_1272021
Alexa Fluor 700 Rat IgG2b kappa Isotype Control	eBioscience	Cat#56-4031-80; RRID: AB_837123
PE-Cyanine7 Rat IgG2a kappa Isotype Control	eBioscience	Cat#25-4321-82; RRID: AB_470200
eFluor 450 Rat IgG1 kappa Isotype Control	eBioscience	Cat#48-4301-82; RRID: AB_1271984

Bacterial and Virus Strains		
AF293	Fungal genetics stock center	#A1100
AF293 ds-Red	(20)	N/A
CEA10 (also known as CBS144.89)	(47)	Received form Robert A. Cramer
Chemicals, Peptides, and Recombinant Proteins		
Alexa Fluor 633 succinimidyl ester	Invitrogen	Cat#S12375
Collagenase IV	ThermoFisher Scientific	Cat#LS004189
RPMI-1640	RPMI-1640	21870092
Enrofloxacin	Bayer	Baytril 100
Voriconazole	Pfizer	N/A
Tween-20	Sigma	P9416
Protease Inhibitor Cocktail	Roche	Cat#11697498001
DNase I, Grade II	Sigma	Cat#10104159001
isoflurane	Henry Schein Animal Health	Cat#29405
Diphtheria toxin (DT)	List Biological Laboratories	Cat#150
Paraformaldehyde, 32% Solution	ThermoFisher Scientific	Cat#50-980-495
CM-H$_2$DCFDA	ThermoFisher Scientific	Cat#C6827
10× HBSS	ThermoFisher Scientific	14065056
TRIzol-LS	Invitrogen	10296028
Mouse IFNa2/4 ELISA kit	Invitrogen	BMS6027
Mouse IL-28B/IFN-lambda 3 DuoSet ELISA	Invitrogen	DY1789B-05
Mouse GM-CSF DuoSet ELISA	Invitrogen	DY415-05
Mouse TNF alpha ELISA Ready-SET-Go!™ Kit	Invitrogen	50-173-31

Mouse IL-6 ELISA Ready-SET-Go!™ Kit	Invitrogen	50-172-19
Mouse IL-1 beta ELISA Kit	Invitrogen	BMS6002
Mouse IL-12 p70 ELISA Kit	Invitrogen	BMS6004
High-Capacity RNA-to-cDNA Kit	Appliedbiosystems	4387406
Ribosomal RNA Control Reagents	Appliedbiosystems	4308329
IFNa2 TaqMan Assay Mm00833961_s1	Thermo Fisher	4331182
IFNa1/a5/a6 TaqMan Assay Mm03030145_gH	Thermo Fisher	4331182
IFNL2/3 TaqMan Assay Mm04204155_gH	Thermo Fisher	4331182
Experimental Models: Organisms/Strains		
BDCA2-DTR	The Jackson Laboratory	JAX: 014176
C57BL/6	The Jackson Laboratory	JAX: 000664
C57BL/6.SJL	Taconic	Stock No. 4007
$Stat1^{-/-}$ mice	The Jackson Laboratory	JAX: 012606
$GBP^{chr3-/-}$ mice		Received from Dr. Thirumala-Devi Kanneganti.
$Cybb^{-/-}$ mice	The Jackson Laboratory	JAX: 002365
Software and Algorithms		
Prism 10	Prism 10	N/A
Flow Jo 10.10.0	Flow Jo 10.10.0	N/A

672   **References**

673   1.   Thompson GR, 3rd, and Young JH. Aspergillus Infections. *N Engl J Med.*

674   2021;385(16):1496-509.

675   2.   Lionakis MS, Dunleavy K, Roschewski M, Widemann BC, Butman JA, Schmitz R, et al.

676   Inhibition of B Cell Receptor Signaling by Ibrutinib in Primary CNS Lymphoma. *Cancer*

677   *Cell.* 2017;31(6):833-43 e5.

678   3.   Varughese T, Taur Y, Cohen N, Palomba ML, Seo SK, Hohl TM, et al. Serious

679   Infections in Patients Receiving Ibrutinib for Treatment of Lymphoid Cancer. *Clin Infect*

680   *Dis.* 2018;67(5):687-92.

681   4.   Desai JV, Zarakas MA, Wishart AL, Roschewski M, Aufiero MA, Donkó Á, et al. BTK

682   drives neutrophil activation for sterilizing antifungal immunity. *J Clin Invest.*

683   2024;134(12):0021-9738.

684   5.   Hoenigl M, Seidel D, Sprute R, Cunha C, Oliverio M, Goldman GH, et al. COVID-19-

685   associated fungal infections. *Nat Microbiol.* 2022;7(8):1127-40.

686   6.   Feys S, Carvalho A, Clancy CJ, Gangneux JP, Hoenigl M, Lagrou K, et al. Influenza-

687   associated and COVID-19-associated pulmonary aspergillosis in critically ill patients.

688   *Lancet Respir Med.* 2024;12(9):728-42.

689   7.   Tischler BY, and Hohl TM. Menacing Mold: Recent Advances in Aspergillus

690   Pathogenesis and Host Defense. *J Mol Biol.* 2019;431(21):4229-46.

691   8.   Brown GD, Denning DW, Gow NA, Levitz SM, Netea MG, and White TC. Hidden

692   killers: human fungal infections. *Sci Transl Med.* 2012;4(165):165rv13.

693   9.   Lionakis MS, and Levitz SM. Host Control of Fungal Infections: Lessons from Basic

694   Studies and Human Cohorts. *Annu Rev Immunol.* 2018;36:157-91.

695   10.   Lionakis MS, Drummond RA, and Hohl TM. Immune responses to human fungal

696   pathogens and therapeutic prospects. *Nat Rev Immunol.* 2023;23(7):433-52.

697   11.   Latgé JP, and Chamilos G. Aspergillus fumigatus and Aspergillosis in 2019. *Clin*

698   *Microbiol Rev.* 2019;33(1):0893-8512.

699   12.   Mills KAM, Aufiero MA, and Hohl TM. Epithelial responses to fungal pathogens. *Curr*

700   *Opin Microbiol.* 2024;80:102508.

701   13.   Gerson SL, Talbot GH, Hurwitz S, Strom BL, Lusk EJ, and Cassileth PA. Prolonged

702   granulocytopenia: the major risk factor for invasive pulmonary aspergillosis in patients

703   with acute leukemia. *Ann Intern Med.* 1984;100(3):345-51.

704   14.   Espinosa V, Jhingran A, Dutta O, Kasahara S, Donnelly R, Du P, et al. Inflammatory

705   monocytes orchestrate innate antifungal immunity in the lung. *PLoS Pathog.*

706   2014;10(2):e1003940.

707   15.   Espinosa V, Dutta O, McElrath C, Du P, Chang YJ, Cicciarelli B, et al. Type III

708   interferon is a critical regulator of innate antifungal immunity. *Sci Immunol.*

709   2017;2(16):2470-9468.

710   16.   Marciano BE, Spalding C, Fitzgerald A, Mann D, Brown T, Osgood S, et al. Common

711   severe infections in chronic granulomatous disease. *Clin Infect Dis.* 2015;60(8):1176-83.

712   17.   Shlezinger N, Irmer H, Dhingra S, Beattie SR, Cramer RA, Braus GH, et al. Sterilizing

713   immunity in the lung relies on targeting fungal apoptosis-like programmed cell death.

714   *Science.* 2017;357(6355):1037-41.

715   18.   Guo Y, Kasahara S, Jhingran A, Tosini NL, Zhai B, Aufiero MA, et al. During

716   Aspergillus Infection, Monocyte-Derived DCs, Neutrophils, and Plasmacytoid DCs

Enhance Innate Immune Defense through CXCR3-Dependent Crosstalk. *Cell host &*

microbe. 2020;28(1):104-16 e4.

19. Swiecki M, Gilfillan S, Vermi W, Wang Y, and Colonna M. Plasmacytoid dendritic cell

ablation impacts early interferon responses and antiviral NK and CD8(+) T cell accrual.

Immunity. 2010;33(6):955-66.

20. Jhingran A, Kasahara S, and Hohl TM. Flow Cytometry of Lung and Bronchoalveolar

Lavage Fluid Cells from Mice Challenged with Fluorescent Aspergillus Reporter

(FLARE) Conidia. *Bio Protoc.* 2016;6(18):2331-8325.

21. Briard B, Karki R, Malireddi RKS, Bhattacharya A, Place DE, Mavuluri J, et al. Fungal

ligands released by innate immune effectors promote inflammasome activation during

Aspergillus fumigatus infection. *Nat Microbiol.* 2019;4(2):316-27.

22. Pollock JD, Williams DA, Gifford MA, Li LL, Du X, Fisherman J, et al. Mouse model of

X-linked chronic granulomatous disease, an inherited defect in phagocyte superoxide

production. *Nat Genet.* 1995;9(2):202-9.

23. Fujimura N, Xu B, Dalman J, Deng H, Aoyama K, and Dalman RL. CCR2 inhibition

sequesters multiple subsets of leukocytes in the bone marrow. *Sci Rep.* 2015;5:11664.

24. Sozzani S, Vermi W, Del Prete A, and Facchetti F. Trafficking properties of

plasmacytoid dendritic cells in health and disease. *Trends Immunol.* 2010;31(7):270-7.

25. Swiecki M, Miller HL, Sesti-Costa R, Cella M, Gilfillan S, and Colonna M. Microbiota

induces tonic CCL2 systemic levels that control pDC trafficking in steady state. *Mucosal*

Immunol. 2017;10(4):936-45.

26.   Maldonado SD, Dai J, Dutta O, Hurley HJ, Singh S, Gittens-Williams L, et al. Human

739   Plasmacytoid Dendritic Cells Express C-Type Lectin Receptors and Attach and Respond

740   to Aspergillus fumigatus. *J Immunol.* 2022;209(4):675-83.

741   27.   Seeds RE, Gordon S, and Miller JL. Characterisation of myeloid receptor expression and

742   interferon alpha/beta production in murine plasmacytoid dendritic cells by flow

743   cytomtery. *J Immunol Methods.* 2009;350(1-2):106-17.

744   28.   Taylor PR, Brown GD, Reid DM, Willment JA, Martinez-Pomares L, Gordon S, et al.

745   The beta-glucan receptor, dectin-1, is predominantly expressed on the surface of cells of

746   the monocyte/macrophage and neutrophil lineages. *J Immunol.* 2002;169(7):3876-82.

747   29.   Preite NW, Feriotti C, Souza de Lima D, da Silva BB, Condino-Neto A, Pontillo A, et al.

748   The Syk-Coupled C-Type Lectin Receptors Dectin-2 and Dectin-3 Are Involved in

749   Paracoccidioides brasiliensis Recognition by Human Plasmacytoid Dendritic Cells. *Front

750   Immunol.* 2018;9:464.

751   30.   Ramirez-Ortiz ZG, Specht CA, Wang JP, Lee CK, Bartholomeu DC, Gazzinelli RT, et al.

752   Toll-like receptor 9-dependent immune activation by unmethylated CpG motifs in

753   Aspergillus fumigatus DNA. *Infection and immunity.* 2008;76(5):2123-9.

754   31.   Herbst S, Shah A, Mazon Moya M, Marzola V, Jensen B, Reed A, et al. Phagocytosis-

755   dependent activation of a TLR9-BTK-calcineurin-NFAT pathway co-ordinates innate

756   immunity to Aspergillus fumigatus. *EMBO molecular medicine.* 2015;7(3):240-58.

757   32.   Assil S, Coleon S, Dong C, Decembre E, Sherry L, Allatif O, et al. Plasmacytoid

758   Dendritic Cells and Infected Cells Form an Interferogenic Synapse Required for Antiviral

759   Responses. *Cell Host Microbe.* 2019;25(5):730-45 e6.

33.   Webster B, Werneke SW, Zafirova B, This S, Coleon S, Decembre E, et al. Plasmacytoid

dendritic cells control dengue and Chikungunya virus infections via IRF7-regulated

interferon responses. *Elife.* 2018;7:2050-084X

34.   Abbas A, Vu Manh TP, Valente M, Collinet N, Attaf N, Dong C, et al. The activation

trajectory of plasmacytoid dendritic cells in vivo during a viral infection. *Nat Immunol.*

2020;21(9):983-97.

35.   Xie X, Shi Q, Wu P, Zhang X, Kambara H, Su J, et al. Single-cell transcriptome profiling

reveals neutrophil heterogeneity in homeostasis and infection. *Nat Immunol.*

2020;21(9):1119-33.

36.   Ballesteros I, Rubio-Ponce A, Genua M, Lusito E, Kwok I, Fernandez-Calvo G, et al. Co-

option of Neutrophil Fates by Tissue Environments. *Cell.* 2020;183(5):1282-97 e18.

37.   Berry MP, Graham CM, McNab FW, Xu Z, Bloch SA, Oni T, et al. An interferon-

inducible neutrophil-driven blood transcriptional signature in human tuberculosis. *Nature.*

2010;466(7309):973-7.

38.   Rocha BC, Marques PE, Leoratti FMS, Junqueira C, Pereira DB, Antonelli L, et al. Type

I Interferon Transcriptional Signature in Neutrophils and Low-Density Granulocytes Are

Associated with Tissue Damage in Malaria. *Cell Rep.* 2015;13(12):2829-41.

39.   Montaldo E, Lusito E, Bianchessi V, Caronni N, Scala S, Basso-Ricci L, et al. Cellular

and transcriptional dynamics of human neutrophils at steady state and upon stress. *Nat*

Immunol. 2022;23(10):1470-83.

40.   Broggi A, Tan Y, Granucci F, and Zanoni I. IFN-lambda suppresses intestinal

inflammation by non-translational regulation of neutrophil function. *Nat Immunol.*

2017;18(10):1084-93.

41.   Chrysanthopoulou A, Kambas K, Stakos D, Mitroulis I, Mitsios A, Vidali V, et al.

Interferon lambda1/IL-29 and inorganic polyphosphate are novel regulators of

neutrophil-driven thromboinflammation. *J Pathol.* 2017;243(1):111-22.

42.   Ardanuy J, Scanlon K, Skerry C, Fuchs SY, and Carbonetti NH. Age-Dependent Effects

of Type I and Type III IFNs in the Pathogenesis of

Infection and Disease. *Journal of Immunology.* 2020;204(8):2192-202.

43.   Lazear HM, Nice TJ, and Diamond MS. Interferon-lambda: Immune Functions at Barrier

Surfaces and Beyond. *Immunity.* 2015;43(1):15-28.

44.   Antos D, Parks OB, Duray AM, Abraham N, Michel JJ, Kupul S, et al. Cell-intrinsic

regulation of phagocyte function by interferon lambda during pulmonary viral, bacterial

super-infection. *PLoS Pathog.* 2024;20(8):e1012498.

45.   Kumar A, Johnson D, Bukowski A, Noto MJ, and Carbonetti NH. IFNlambda signaling

in neutrophils enhances the pathogenesis of Bordetella pertussis infection. *J Leukoc Biol.*

2024:0741-5400.

46.   Antony F, Pundkar C, Sandey M, Jaiswal AK, Mishra A, Kumar A, et al. IFN-lambda

Regulates Neutrophil Biology to Suppress Inflammation in Herpes Simplex Virus-1-

Induced Corneal Immunopathology. *J Immunol.* 2021;206(8):1866-77.

47.   Girardin H, Latge JP, Srikantha T, Morrow B, and Soll DR. Development of DNA probes

for fingerprinting Aspergillus fumigatus. *J Clin Microbiol.* 1993;31(6):1547-54.

48.   Heung LJ, Jhingran A, and Hohl TM. Deploying FLAREs to Visualize Functional

Outcomes of Host-Pathogen Encounters. *PLoS pathogens.* 2015;11(7):e1004912.

49.   Hohl TM, Van Epps HL, Rivera A, Morgan LA, Chen PL, Feldmesser M, et al.

805   Aspergillus fumigatus triggers inflammatory responses by stage-specific beta-glucan

806   display. *PLoS pathogens.* 2005;1(3):e30.

807   50.   Hackstein H, Wachtendorf A, Kranz S, Lohmeyer J, Bein G, and Baal N. Heterogeneity

808   of respiratory dendritic cell subsets and lymphocyte populations in inbred mouse strains.

809   *Respir Res.* 2012;13:94.

810   51.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

811   universal RNA-seq aligner. *Bioinformatics.* 2013;29(1):15-21.

812   52.   Liao Y, Smyth GK, and Shi W. featureCounts: an efficient general purpose program for

813   assigning sequence reads to genomic features. *Bioinformatics.* 2014;30(7):923-30.

814   53.   Love MI, Huber W, and Anders S. Moderated estimation of fold change and dispersion

815   for RNA-seq data with DESeq2. *Genome Biol.* 2014;15(12):550.

816   54.   Korotkevich G, Sukhov V, and Sergushichev A. Fast gene set enrichment analysis.

817   *bioRxiv.* 2019:060012.

818   55.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al.

819   Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide

820   expression profiles. *Proc Natl Acad Sci U S A.* 2005;102(43):15545-50.

821   56.   Loures FV, Rohm M, Lee CK, Santos E, Wang JP, Specht CA, et al. Recognition of

822   Aspergillus fumigatus hyphae by human plasmacytoid dendritic cells is mediated by

823   dectin-2 and results in formation of extracellular traps. *PLoS Pathog.*

824   2015;11(2):e1004643.

Figure 1. pDC transcriptome following *A. fumigatus* infection.

(A) Experimental scheme for bulk RNA-seq of BM pDCs sorted from naïve mice (red symbols)

and BM (green symbols) and lung (blue symbols) pDCs sorted from *A. fumigatus*-infected mice.

(B) Principal component analysis of gene expression in sorted BM pDCs from naïve (triangles)

and infected mice (dots) and of sorted lung pDCs from infected mice (squares). Each symbol

represents a biological replicate. pDCs from 10 mice were pooled for each replicate, and 2

replicates were included in each of 2 experiments, denoted as Expt1 (filled symbol) and Expt 2

(open symbols). (C) Number of differentially expressed genes for three comparisons of 2 pDC

subsets. (D) Differentially enriched KEGG pathways ($q < 0.05$) in pDCs isolated from infected

lungs vs infected BM. Black bars indicate pathways enriched in lung pDCs from infected mice,

white bars indicate pathways enriched in BM pDCs from infected mice. (E) Heatmap for 35

selected genes with a >2 -fold difference in expression and a false discovery rate (FDR) $p < 0.05$.

Each lane represents 1 replicate from 2 expts. (F - I) Representative flow cytometry plots (F and

G) and quantified mean fluorescence intensity MFI (H and I) of IFNβ YFP and ICAM1

expression in pDCs isolated from (top) naïve BM, (middle), the BM of *A. fumigatus*-infected

mice or (bottom) the lungs of *A. fumigatus*-infected mice. (A-I) Infection dose: 3×10^7 CEA10

conidia via intratracheal route, analysis 72 hpi. (B - E) Data were pooled from 2 independent

experiments. (H and I) Statistical analysis: Kruskal-Wallis test. See also Figure S1.

Figure 2. Cytokine profiles in pDC-depleted mice during *A. fumigatus* infection

(A - C) *Ifn* gene expression, measured by qRT-PCR using TaqMan probes, in the lung of

C57BL6/J mice at indicated times, n = 5-6 per group. (D) Lung *Ifn* gene expression and (E and

F) lung cytokine levels measured by ELISA in DT-treated pDC depleter mice (BDCA2-DTR[Tg/+];

red symbols) and DT-treated non-Tg littermate controls (black dashed line), n = 8 per group. (A -

E) Infection dose: 3×10^7 CEA10 conidia via intratracheal route, analysis 72 hpi. Data were

pooled from 2 independent experiments and presented as mean ± SEM. Dots represent individual

mice. Statistical analysis: Kruskal-Wallis test.

Figure 3. STAT1 signaling regulates neutrophil-intrinsic *Aspergillus* killing.

(A) Scheme to generate *Stat1*$^{-/-}$ and *Stat*$^{+/+}$ mixed bone chimeric mice and (B) illustration of the

two-component fluorescent *Aspergillus* reporter (FLARE) system used to measure conidial

uptake and killing by *Stat1*$^{-/-}$ and *Stat*$^{+/+}$ lung leukocytes in mixed bone marrow chimeric mice.

(C) Representative plot that display DsRed and AF633 fluorescence intensity of lung neutrophils.

The R1 gate denotes neutrophils that contain live conidia, the R2 gate denotes neutrophils that

contain killed conidia. (D and E) The plots show (D) normalized lung neutrophil conidial uptake

(R1 + R2) ± SEM and (E) conidial viability (R1/ (R1 + R2) ± SEM in indicated lung neutrophils

isolated from *Stat1*$^{-/-}$ (purple symbols) and *Stat1*$^{+/+}$(black symbols) mixed BM chimeric mice

(n=8 per group). (F) Scheme to generate *Stat1*$^{-/-}$ and WT (*Stat*$^{+/+}$) chimeric mice. (G) Kaplan

Meier survival of *Stat1*$^{-/-}$ and *Stat*$^{+/+}$ chimeric mice (n = 7-8 per group) infected with 3-6 $\times 10^7$

CEA10 conidia. (H) Lung cytokine levels in *Stat1*$^{-/-} \rightarrow$ WT (purple symbols) and *Stat1*$^{+/+} \rightarrow$ WT

(black symbols) single chimeric mice. (I) Representative plot and (J) mean ± SEM neutrophil

ROS production in cells isolated from *Stat1*$^{-/-} \rightarrow$ WT (purple symbols) and *Stat1*$^{+/+} \rightarrow$ WT (black

symbols) single chimeric mice (n=5 per group). (C - E) Infection dose: 3×10^7 Af293 FLARE

conidia via intratracheal route, analysis 72 hpi. (F - I) Infection dose: 3×10^7 CEA10 conidia via

intratracheal route, analysis 72 hpi. (D and E) Data were pooled from 2 independent experiments.

Dots represent individual mice. Statistical analysis: paired t test. (H - J) Data are representative

of 2 experiments. Dots represent individual mice. Statistical analysis: Mann-Whitney test. See

also Figure S2.

Figure 4. pDCs regulate neutrophil STAT1-dependent antifungal activity.

(A) Scheme to generate and to deplete pDCs or not in CD45.1$^+$ BDCA2- DTR$^{Tg/+}$ $Stat1^{+/+}$ and

CD45.2$^+$ BDCA2- DTR$^{Tg/+}$ $Stat1^{-/-}$ mixed bone marrow chimeric mice, resulting in 4

experimental groups (G1-G4). (B-D) (B) Representative plots that display DsRed and Af633

fluorescence intensity in lung neutrophils from 4 experimental groups: G1: pDC$^+$ $Stat1^{+/+}$

($Stat1^{+/+}$ neutrophils from pDC-sufficient mice), G2: pDC$^-$ $Stat1^{+/+}$ ($Stat1^{+/+}$ neutrophils from

pDC-depleted mice), G3: pDC$^+$ $Stat1^{-/-}$ ($Stat1^{-/-}$ neutrophils from pDC-sufficient mice), G4: pDC$^-$

$Stat1^{-/-}$ ($Stat1^{-/-}$ neutrophils from pDC-depleted mice). The R1 gate indicates the frequency of

neutrophils that contain live conidia, the gate R2 indicates the frequency of neutrophils that

contain killed conidia. (C and D) The plots show mean neutrophil (C) conidial uptake (R1 + R2)

± SEM and (D) conidial viability (R1/ (R1 + R2) ± SEM in indicated lung neutrophils isolated

from the 4 groups, n = 12 per group, data pooled from 2 experiments. (E) Mean ± SEM ROS

production in indicated lung neutrophils, n = 6 per group. (B - D) Infection dose: 3×10^7 Af293

FLARE conidia via intratracheal route, analysis 72 hpi. (C and D) Data were pooled from 2

independent experiments and presented as mean ± SEM, (E) Data are representative of 2

experiments. Dots represent individual mice. Statistical analysis: Kruskal-Wallis test. See also

Figure S3.

Figure 5. STAT1-dependent guanylate-binding proteins are dispensable for the neutrophil

antifungal response.

(A) Principal component analysis of global gene expression, (B) differentially enriched KEGG

pathways, and (C) Volcano plot of differentially expressed genes in $Stat1^{-/-}$ (purple symbols) and

$Stat1^{+/+}$(black symbols) neutrophil sorted from mixed bone marrow chimeric mice (n = 6) at 72

hpi with 3×10^7 CEA10 conidia. Selected downregulated genes in $Stat1^{-/-}$ neutrophils are

highlight in the Volcano plot. (D) Experimental scheme to generate $GBP^{chr3-/-}$ and $GBP^{chr3+/+}$

single chimeric mice. (E) Kaplan Meier survival (n = 7-8 per group), and (F) mean ± SEM lung

CFU (n = 5 per group) in $GBP^{chr3-/-}$ (red symbols) and $GBP^{chr3+/+}$ (black symbols) single

chimeric mice infected with $3\text{-}6 \times 10^7$ CEA10 conidia. (G) Experimental scheme to generate

$GBP^{chr3-/-}$ and $GBP^{chr3+/+}$ mixed chimeric mice. (H and I) The plots show normalized neutrophil

(H) conidial uptake (R1 + R2) ± SEM and (I) conidial viability (R1/ (R1 + R2) ± SEM in lung

neutrophils isolated from $GBP^{chr3-/-}$ (red symbols) and $GBP^{chr3+/+}$ (black symbols) mixed bone

marrow chimeric mice (n = 6 per group). (J) Mean ± SEM neutrophil ROS production in

neutrophils isolated from $GBP^{chr3-/-}$ (red symbols) and $GBP^{chr3+/+}$ (black symbols) mixed bone

marrow chimeric mice (n=6 per group). (H - J) Infection dose: 3×10^7 Af293 FLARE conidia

via intratracheal route, analysis 72 hpi. (F, H - J) Data are representative of 2 experiments and

presented as mean ± SEM. Dots represent individual mice. Statistical analysis: Mann-Whitney

test.

Figure 6. STAT1-dependent control of *Cybb* expression and CYBB protein levels in

neutrophils.

(**A**) Heatmap of *Cybb*, *Cyba*, *Ncf1*, *Ncf3*, *Ncf4*, *Nox1*, *Nox3*, and *Nox4* expression in neutrophils

sorted from *Stat1*$^{-/-}$ (purple symbols) and *Stat1*$^{+/+}$(black symbols) mixed bone marrow chimeric

mice (n = 6). Genes with a STAR* are DEG. Each lane is an independent biological replicate. (B)

qRT-PCR of *Cybb* mRNA expression in neutrophils sorted from *Stat1*$^{-/-}$ (purple symbols) and

Stat1$^{+/+}$ (black symbols) mice. (C) ATAC-seq analysis of neutrophils were sorted from *Stat1*$^{-/-}$

(purple peaks) and *Stat*$^{+/+}$ (black peaks) mice. Gene tracks show chromatin accessibility at the

Cybb and *Gbp2* locus. (D) Bone marrow and lung neutrophils were sorted from *Stat1*$^{-/-}$ (purple

peaks) and *Stat1*$^{+/+}$ (black peaks) mice and processed for CUT&RUN. Gene tracks show STAT1

signal as normalized fragment pileup (y-axis) plotted by genome position (x-axis). The shaded

box highlights a putative STAT1 binding site at the Gbp2 promoter region. Gene tracks show

H3K4me3 ChIP-seq signal as normalized fragment pileup (top rows; green). (E - H) Western

blot and quantitation of (F) CYBB, (G) CYBA, (H) NCF4 vs GAPDH protein levels in lung

neutrophils sorted from *Stat1*$^{-/-}$ (purple symbols) and *Stat1*$^{+/+}$ (black symbols) mice, neutrophils

from 4-5 mice were pooled to obtain sufficient protein for Western blot analysis in each

biological replicate. (A - H) Infection dose: 3×10^7 CEA10 conidia via intratracheal route,

analysis 72 h pi. (C) Data were calculated by Kruskal-Wallis nonparametric test for multiple

comparisons for each group compared with control group. (G – I) Data are representative of 2

experiments. Dots represent individual mice. Statistical analysis: Mann-Whitney test. See also

Figure S4.

Figure S1. Related to Fig. 1. pDC transcriptome analysis following *A*. *fumigatus* infction.

(A) The plots indicate the FACS sorting strategy for BM and lung pDCs (top row) and the

typical (>95%) pDC purity after FACS sorting (bottom row). (B) Volcano plot of the

differentially expressed genes in pDCs sorted from the bone marrow of infected vs. naïve mice.

(C) Volcano plot of the differentially expressed genes in pDCs sorted from infected lungs vs

naïve bone marrow. (D) Volcano plot of the differentially expressed genes in pDCs sorted from

infected lungs vs infected bone marrow. (E) The plot shows differentially exriched KEGG

pathways (q$\square$<$\square$0.05) observed in pDCs isolated from infected lungs vs infected BM. Black

bars indicate pathways enriched in lung pDCs from infected mice, white bars indicate pathways

enriched in BM pDCs from infected mice.

Figure S2. Related to Fig. 3. Type I and III IFN receptor expression on Ly6G^{+} lung

neutrophils.

(A) Lung sections from naïve and *A. fumigatus*-infected mice were analyzed by RNAscope

using probes to Ly6G (first column) and IFNαR1 (second column), and DAPI (third column),

and merged images are shown in fourth column, and yellow arrows indicate examples of co-

localization of the Ly6G and IFNαR1 probes within the same nuclei. (B) Lung sections from

naïve and *A. fumigatus*-infected mice were analyzed by RNAscope using probes to Ly6G (first

column) and IFNλR1 (second column), and DAPI (third column), and merged images are shown

in fourth column, and yellow arrows indicate examples of co-localization of the Ly6G and

IFNλR1 probes within the same nuclei. (A and B) Infection dose: 3×10^{7} CEA10 conidia via

intratracheal route, analysis at 72 hpi. Scale bar = 20 μm.

Figure S3. Related to Fig. 4. STAT1 modulates the neutrophil antifungal response.

(A) Lung cytokine levels measured by ELISA of $Stat1^{-/-}$ (purple symbols) and $Statt1^{+/+}$ (black

symbols) mice. (B - E) Lung (B) neutrophil, (C) monocyte, (D) Mo-DC and (E) pDC numbers in

$Stat1^{-/-}$ (purple symbols) and $Stat1^{+/+}$ (black symbols). (A - E) Infection dose: 3×10^7 CEA10

conidia via intratracheal route, analysis 72 hpi. Data are representative of 2 experiments. Dots

represent individual mice. Statistical analysis: Mann-Whitney test.

Figure S4. Related to Fig. 6. $Cybb^{-/-}$ mice are susceptible to *A. fumigatus* infection.

(A) *Cybb* gene expression in neutrophils isolated from $Stat1^{-/-}$ (purple symbols) and $Stat1^{+/+}$

(black symbols) mice. Gene expression as determined by qRT-PCR (n = 3 - 5 per group) (B)

Kaplan Meier survival (n = 7 - 8 per group) of $Cybb^{-/-}$ (blue symbols) and WT control (black

symbols) mice infected with $3 - 6 \times 10^7$ CEA10 conidia. (C) Genome browser tracks of ATAC-

seq data at the *Cybb* locus in lung neutrophils isolated from $Stat1^{-/-}$ (purple symbols) and

$Stat^{+/+}$ (black symbols) mice. (D) Statistics of peaks at indicated *Cybb* loci in lung neutrophils

isolated from $Stat1^{-/-}$ (purple symbols) and $Stat^{+/+}$ (black dash line) mice. (A - D) Infection dose:

3×10^7 CEA10 conidia via intratracheal route, analysis 72 hpi. *P < 0.05; **P < 0.01; ***P <

0.001. Data presented as mean ± SEM. Dots represent individual mice. Statistical analysis:

Kruskal-Wallis test for multiple comparisons.

A *Ifnα-1/5/6*

B *Ifnα-2*

C *Ifnλ-2/3*

D

E

F

A

WT (CD45.1⁺)

Stat1⁻/⁻ (CD45.2⁺)

BM cells 1:1 → 900 cGy → WT (CD45.1/.2⁺) → 6 wks → i.n. *A. fumigatus*

B

Leukocyte → Uptake → DsRed⁺AF633⁺ Leukocyte + live *A. fumigatus* → Killing → DsRed⁻AF633⁺ Leukocyte + killed *A. fumigatus*

C

WT

NØ Gate

R1 4.3%

R2 32.2%

Stat1⁻/⁻

NØ Gate

R1 5.5%

R2 23.9%

Aspergillus Viability (dsRed)

Aspergillus Tracer (Alexa Fluor 633)

D

0.0010

% Uptake R1+R2 (Normalized to WT)

WT   *Stat1⁻/⁻*

E

0.0002

% Aspergillus Viability R1/(R1+R2) (Normalized to WT)

WT   *Stat1⁻/⁻*

F

Stat1⁻/⁻ or WT (CD45.2⁺)

BM cells → 900 cGy → WT (CD45.1⁺) → 6 wks → i.n. *A. fumigatus*

G

% Survival

P = 0.0009

WT → WT

Stat1⁻/⁻ → WT

Days

H

Log10 pg / ml

○ WT → WT
○ *Stat1⁻/⁻* → WT

IFNα-2/4 IFNλ-2/3 IL-1β IL-6 IL-12p70 IL-23 TNF-α

I

NØ Gate

ROS⁺

Count

ROS

J

0.0317

ROS (MFIX1000)

WT   *Stat1⁻/⁻*

A Principal component analysis of neutrophils

● *Stat1⁻/⁻* □ WT

PC2: 6% variance

PC1: 90% variance

B KEGG pathways enriched: WT vs *Stat1⁻/⁻*

Cytosolic DNA sensing pathway
Proteasome
RIG-I like receptor signaling pathway
Toll like receptor signaling pathway
Cytokine-cytokine receptor interaction
Type I diabetes mellitus
Antigen processing and presentation
Phosphatidylinositol signaling system
Systemic lupus erythematosus

-log10 (p Value)

C

-log10 (padj)

Gbp2
Gbp3
Gbp7
Gbp5
Cybb

log2 Fold Change

D

Gbp^chr3-/- or WT (CD45.2⁺) → BM cells → 900 cGy → WT (CD45.1⁺) → 6 wks → i.n. *A. fumigatus*

G

WT (CD45.1⁺)
Gbp^chr3-/- (CD45.2⁺) → BM cells 1:1 → 900 cGy → WT (CD45.1/.2⁺) → 6 wks → i.n. *A. fumigatus*

E

% Survival

WT → WT
Gbp^chr3-/- → WT

Days

F

CFU/g lung x 10⁴

0.5204

○ WT → WT
○ *Gbp^chr3-/-* → WT

H

% Uptake R1+R2 (Normalized to Control)

n.s.

I

% Aspergillus Viability R1/(R1+R2) (Normalized to Control)

n.s.

J

ROS (MFI X 1000)

n.s.